Circadian photoreceptor CRYPTOCHROME promotes wakefulness under short winter-like days via a GABAergic circuitry

Lixia Chen[1,2], Danya Tian[1], Chang Su[1] and Luoying Zhang[1,3]*

[1]Key Laboratory of Molecular Biophysics of Ministry of Education, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China

[2]National "111" Center for Cellular Regulation and Molecular Pharmaceutics, Key Laboratory of Fermentation Engineering (Ministry of Education), Hubei University of Technology, Wuhan, Hubei 430068, China

[3]Hubei Province Key Laboratory of Oral and Maxillofacial Development and Regeneration, Wuhan, Hubei 430022, China

*Correspondence: Luoying Zhang, E-mail: zhangluoying@hust.edu.cn

Abstract

A cardinal symptom of seasonal affective disorder (SAD, also known as winter depression) is hypersomnolence, while the cause of this "winter sleepiness" is not known. Here we found that lack of the circadian photoreceptor *cryptochrome* (*cry*) leads to increased sleep under short winter-like days in fruit flies, reminiscent of the hypersomnolence in SAD. CRY functions in neurons that synthesize the major inhibitory neurotransmitter GABA, including the small ventral lateral neurons which are known to be circadian pacemakers, and down-regulates the GABAergic tone. This in turn leads to increased neural activity of the wake-promoting large ventral lateral neurons, a subset of circadian neurons that are inhibited by GABA-A receptor. CRY protein is known to be degraded by light, thus rendering CRY to be functional within this GABAergic circuitry to enhance wakefulness only under short day length. Taken together, we demonstrate a mechanism that specifically regulates wakefulness under short winter-like days, which may provide insights regarding the winter sleepiness in SAD.

Keywords: wakefulness, circadian pacemaker, CRYPTOCHROME, GABA, photoperiod, photoreceptor

Introduction

Seasonal affective disorder (SAD), also known as winter depression, is characterized by the onset of depression in fall/winter months and spontaneous remission in the spring/summer (Rosenthal et al., 1984). It is generally agreed that SAD is caused by lack of day light in the fall/winter months due to the short day length (or photoperiod), as bright light therapy is effective and most commonly used for treating the depression associated with SAD (Golden et al., 2005). The prevalence of SAD is ~1-10% world-wide, with symptoms lasting for approximately 40% of the year (Kurlansik and Ibay, 2012). About 64-80% of SAD patients report a winter increase in sleep, ranging from 30 min to 2 h longer in duration compared to controls, which is considered to be a distinguishing symptom in the characterization and diagnosis of SAD (Wescott et al., 2020). Currently almost nothing is known regarding the underlying mechanism of this winter hypersomnolence.

This phenomenon of winter hypersomnolence in SAD patients implicates distinct mechanisms that regulate sleep under short winter-like photoperiods vs. longer non-winter-like photoperiods, as the sleep of these individuals appear to be selectively perturbed under short photoperiods. However, the mechanisms by which sleep duration is determined under different photoperiods have not been characterized. Since the circadian clock is believed to be important for adaptations to seasonal changes in the environment and in particular, seasonal changes of photoperiod, we hypothesize that the circadian clock may also participate in regulating sleep duration under different photoperiods (Wood and Loudon, 2014).

To address our hypothesis, we tested fruit flies mutant for different circadian

clock genes under a range of photoperiods. We found that flies lacking the circadian

photoreceptor CRYPTOCHROME (CRY) display increased sleep duration

specifically under short photoperiods, similar to the winter hypersomnolence in SAD.

Genetic and pharmacological analysis identified that CRY is functioning in

GABAergic neurons and acts upon GABA-A receptor to promote wakefulness. We

further narrowed down the neural circuitry mediating the influences of CRY on sleep

by demonstrating that *cry* deficiency increases the GABAergic tone and reduces

calcium concentration in the wake-promoting large ventral lateral neurons (l-LNvs)

which are known to be GABA-A+ (Hamasaka et al., 2005; Parisky et al., 2008).

Consistently, inhibiting these neurons increases sleep while activating them blocks the

effects of *cry* deficiency on sleep. CRY likely functions in part in the GABAergic

small ventral lateral neurons (s-LNvs), and lack of *cry* increases GABA level and the

activity of these cells while impairing their GABA transmission suppresses the sleep

phenotype of *cry* mutants (Allada and Chung, 2010). In summary, here we identify a

potential role for CRY in down-regulating the GABAergic signaling from the s-LNvs

to the l-LNvs specifically under short photoperiod. This in turn enhances the neural

activity of the wake-promoting l-LNvs, resulting in increased wakefulness during

short winter-like days. These findings reveal a mechanism underlying how sleep

duration is determined under winter-like photoperiod, while disruptions of this

regulatory system may be related to the winter hypersomnolence associated with

SAD.

Results

cry mutation increases sleep duration specifically under short photoperiods

We assessed the sleep of flies mutant for circadian clock gene *period* (per^0), *timeless* (tim^0), *clock* (clk^{jrk}), *cycle* (cyc^0) and *cry* (cry^b) under a range of photoperiods (Figure 1A and B)(Allada, 1998; Konopka and Benzer, 1971; Sehgal et al., 1994; Stanewsky, 1998). We found that cry^b mutation, which is known to be a loss of function or severe hypomorphic allele, leads to increased sleep duration under 4 h light: 20 h dark condition (4L20D) and 8L16D but not under longer photoperiods (Stanewsky, 1998). Since this phenotype recapitulates the winter hypersomnolence of SAD patients, we further characterized the effects of *cry* deficiency on sleep. We focused on sleep under 4L20D, as the extent of sleep increase is slightly larger than that of 8L16D. We also demonstrated that *cry* mutation lengthens sleep duration in both male and female flies, while waking activity is not significantly reduced in the mutants (Figure 1C and D). This means the increased sleep in the mutants is not caused by defects of locomotor ability. We next examined the sleep architecture of these flies and found that *cry* mutation enhances sleep by extending the duration of average sleep bout rather than increasing sleep bout number, indicating that *cry* deficiency promotes sleep consolidation under short photoperiod (Figure 1E and F). Because CRY exerts influences on sleep/wakefulness in a gender-independent manner, we used male flies for the remainder of the study. In addition, we tested the effects of a *cry* knock-out allele (cry^0) on sleep under 4L20D and also observed significantly prolonged sleep

duration, similar to cry^b mutation (Figure 1—figure supplement 1) (Dolezelova et al.,

2007).

CRY functions in GABAergic neurons and promotes wakefulness via

GABA/GABA-A

Upon light activation, CRY binds to the clock protein TIM and results in its

degradation, which is followed subsequently by degradation of CRY itself (Busza et

al., 2004; Ceriani et al., 1999; Lin et al., 2001). Therefore, we first tested whether the

influences of CRY on sleep/wakefulness requires TIM. We monitored sleep in flies

mutant for both *cry* and *tim*, and found that the sleep duration of these double mutants

are comparable to that of *cry* mutants (Figure 2—figure supplement 1). This indicates

that CRY promotes wakefulness in a TIM-independent manner.

To identify anatomical substrates that mediate the effects of CRY on wakefulness,

we employed the UAS/GAL4 system to knock down *cry* in different brain structures

and cell types and verified that *cry* is indeed knocked down by assessing its mRNA

level (Figure 2—figure supplement 2). We found that knocking down *cry* in

GABAergic neurons using *vesicular GABA transporter* (*VGAT*)GAL4 or two

independent *glutamic acid decarboxylase 1* (*Gad1*)GAL4 lines results in increased

sleep duration under 4L20D but not 12L12D, similar to the *cry* mutant phenotype

(Figure 2A-D; Figure 2—figure supplement 3A-3L)(Deng et al., 2019). Given that

*Gad1*GAL4-2 generated a more prominent sleep phenotype, we used this driver for

the remainder of this study.

Since CRY appears to act in GABAergic neurons to promote wakefulness, we

tested whether GABA is involved in this regulatory process. We first took a

pharmacological approach and fed flies with drugs to inhibit GABA transaminase

(ethanolamine-O-sulphate, EOS) or GABA transporter (nipecotic acid, NipA). Both of

these treatments are known to increase GABA level, and indeed lengthens the sleep

duration in WT flies (Ki and Lim, 2019; Leal and Neckameyer, 2002). *cry* mutation

fails to enhance sleep when treated with NipA, while the extent of sleep increase

caused by *cry* mutation is smaller when treated with EOS compared to the control

(Figure 2E; Figure 2—figure supplement 4A and B). Similarly, knocking down *cry* in

GABAergic neurons no longer lengthens sleep duration after NipA treatment (Figure

2F; Figure 2—figure supplement 4C). For further validation, we adopted a genetic

approach. We found that knocking down *Gad1*, the synthetic enzyme for GABA,

suppresses the long sleep phenotype of *cry* RNAi flies (Figure 2G; Figure 2—figure

supplement 4D) (Jackson et al., 1990). In addition, we knocked down *VGAT* which

encodes an essential transporter responsible for packing GABA into synaptic vesicles

(Enell et al., 2007). *cry* deficiency fails to lengthen sleep duration when *VGAT* is

knocked down (Figure 2H and I; Figure 2—figure supplement 4E and F). We verified

that *Gad1* and *VGAT* are indeed knocked down by measuring their mRNA levels

(Figure 2—figure supplement 4G and H). As a control for the genetic interaction

experiments, we co-expressed a GFP RNAi and found this does not significantly alter

the sleep duration of *cry* RNAi flies (Figure 2—figure supplement 4I). These series of

results indicate that CRY acts via GABA to promote wakefulness.

We next sought to identify GABA receptor that mediates the effects of CRY on

sleep/wakefulness. We fed flies with agonists of GABA-A (THIP) and GABA-B

receptor (SKF-97541) (Ki and Lim, 2019; Matsuda et al., 1996; Mezler et al., 2001).

Both drugs enhance sleep in WT, while *cry* mutation can increase sleep in flies fed

with SKF-97541 but not THIP, implicating that CRY acts through GABA-A to

promote wake (Figure 3A-C). Consistently, GABA-A receptor antagonist

carbamezapine (CBZ) reduces sleep in WT flies while *cry* mutation fails to lengthen

sleep duration after CBZ treatment (Figure 3D and E) (Agosto et al., 2008). We also

treated *cry* RNAi flies with THIP or CBZ and found this treatment abolished the long

sleep phenotype as well (Figure 3F-I). To validate that CRY modulates

sleep/wakefulness by acting upon GABA-A receptor, we tested for genetic interaction

between *cry* and *Resistant to dieldrin* (*Rdl*), a gene that encodes GABA-A receptor in

flies and has previously been shown to be involved in sleep regulation (Chung et al.,

2009; ffrench-Constant et al., 1993; Parisky et al., 2008). We found that *Rdl* mutation

(*Rdl^{MDRR}*) blocks the sleep-enhancing effect of *cry* RNAi (Figure 3J-L)(Agosto et al.,

2008). These findings demonstrate that GABA-A receptor mediates the

wake-promoting function of CRY.

In short, pharmacological and genetic approaches reveal that CRY acts in

GABAergic neurons and impinges on GABA/GABA-A signaling to promote

wakefulness.

CRY acts upon l-LNvs to promote wakefulness

Previous studies reported that GABA enhances sleep at least in part by activating

RDL on the l-LNvs, thus inhibiting the activities of these cells and reducing the

release of the arousal-promoting neuropeptide pigment dispersing factor (PDF)

(Chung et al., 2009; Parisky et al., 2008). Therefore, we tested whether the l-LNvs

mediate the effects of CRY on sleep/wakefulness. We first tested whether the l-LNvs

indeed receive projections from GABAergic neurons. We expressed a GFP-tagged

synaptotagmin (*syt-GFP*) which labels axon terminals using *Gad1*GAL4, and GFP

signal was observed at the l-LNv soma (Figure 4—figure supplement 1A) (Zhang et

al., 2002). We employed the GFP Across Synaptic Partners (GRASP) method with

enhanced specificity (t-GRASP) to check for synaptic connections between

GABAergic neurons and l-LNvs (Shearin et al., 2018). Pre-mGRASP was expressed

in the axon terminal of GABAergic neurons (using *Gad1*GAL4), whereas its

t-GRASP partner post-mGRASP was expressed in the l-LNvs (using the driver

*Pdf*LexA) (Shang et al., 2008). GFP signal can be detected at the l-LNv soma, which

means GABAergic neurons may send synaptic projections to the l-LNvs (Figure 4

—figure supplement 1B). We further validated this connection using the *trans*-Tango

technique which labels down-stream synaptic targets of GABAergic neurons with the

HA tag, and we were able to observe HA expression in the l-LNvs (Figure 4—figure

supplement 1C) (Talay et al., 2017). All in all, these results strongly suggest that

GABAergic neurons project to the l-LNvs and form synaptic connections.

Consistently, knocking down *Rdl* in the l-LNvs rescues the long sleep phenotype of

cry mutants (Figure 4—figure supplement 1D).

We next examined the effects of *cry* mutation on GABA level at the l-LNv soma

by immunostaining. As expected, GABA signal is significantly increased in *cry*

mutants under 4L20D but not 12L12D (Figure 4A-D). Consistent with elevated

GABA, GCaMP6m signal, an indicator of intracellular calcium concentration, is

significantly reduced in the l-LNvs under 4L20D (Figure 4E and F) (Chen et al.,

2013). This implies that the neural activity of these cells is down-regulated, possibly

due to increased GABA signaling.

To test whether the long sleep phenotype in *cry* mutants is due to reduced activity

of the l-LNvs, we electrically silenced these neurons by expressing an inward

rectifying potassium channel Kir2.1 (Baines et al., 2001). This results in lengthened

sleep duration under 4L20D, mimicking the sleep phenotype of *cry* mutants (Figure

4G; Figure 4—figure supplement 2A). On the other hand, activating the l-LNvs by

expressing the temperature-gated depolarizing cation channel TrpA1 or the bacterial

depolarization-activated sodium channel NachBac blocks the sleep enhancing effect

of *cry* mutation (Figure 4H-K; Figure 4—figure supplement 2B-E) (Hamada et al.,

2008; Nitabach et al., 2006). These findings support the notion that *cry* mutation

lengthens sleep duration by down-regulating the neural activity of the l-LNvs. Since it

has been shown that the l-LNvs promote arousal by releasing PDF, we assessed

whether PDF or its receptor PDFR is required by CRY to exert influence on

sleep/wakefulness (Chung et al., 2009; Parisky et al., 2008). *cry* mutation fails to

lengthen sleep duration on *Pdf* or *Pdfr* mutant background, indicating the necessity of

PDF signaling in mediating the arousal-promoting function of CRY (Figure 4L and M;

Figure 4—figure supplement 2F and G) (Hyun et al., 2005; Lear et al., 2005; Mertens

et al., 2005; Renn et al., 1999).

To summarize, these results strongly suggest that CRY promotes wakefulness by

reducing GABA signaling and thus increasing the activity of the l-LNvs, which may

in turn lead to increased release of PDF.

CRY acts in the s-LNvs to promote wakefulness

To identify the subset of GABAergic neurons in which CRY functions to regulate

wakefulness, we first examined the expression pattern of *Gad1*GAL4 by labeling the

GAL4+ cells with a nuclear GFP (nls-GFP) (Shiga et al., 1996). While the l-LNvs are

not GFP+, we noticed that the s-LNvs which are close to the l-LNvs and also express

PDF appear to be GFP+ (Figure 5A and B; Figure 5-figure supplement 1A). In

addition, nls-GFP was expressed in GABAergic neurons (using *Gad1*GAL4), whereas

RFP was expressed in the l- and s-LNvs (using the driver *Pdf*LexA). GFP signal can

be detected in the s-LNvs but not the l-LNvs (Figure 5-figure supplement 1B).

Because the s-LNvs are known to express CRY, we suspected that CRY may be acting

in these s-LNvs to regulate the activity of the l-LNvs via GABA signaling (Benito et

al., 2008; Yoshii et al., 2008). To test this idea, we knocked down *cry* in the s-LNvs

using R6GAL4 while over-expressing *dicer2* (*dcr2*) to enhance RNAi efficiency and

observed a modest but significant lengthening of sleep duration (Figure 5C; Figure

5—figure supplement 2A) (Helfrich-Forster et al., 2007). However, when we adopted

a *Pdf*GAL80 to block the actions of GAL4 in the PDF neurons in

*Gad1*GAL4/UAS*cry*RNAi, this does not alter the long-sleep phenotype (Figure

5—figure supplement 2B) (Stoleru et al., 2004). These series of results suggest that

cry expression in the s-LNvs are necessary but not sufficient to maintain normal

sleep/wakefulness. We were indeed able to detect GABA signal at the s-LNvs, while

GABA intensity is enhanced in *cry* mutants under 4L20D but not 12L12D (Figure

5D-G). To further validate that the s-LNvs are GABAergic, we knocked down *VGAT*

in these cells and observed a decrease of GABA intensity (Figure 5—figure

supplement 3A and B).

Next, we examined the effects of *cry* deficiency on the activity level of the

s-LNvs. We found that *cry* mutation increases calcium concentration in these cells

under 4L20D, in stark contrast to that of the l-LNvs (Figure 5H and I). This strongly

suggests that *cry* deficiency leads to elevated neural activity in the s-LNvs.

Consistently, we observed increased sleep when we activated these cells using TrpA1,

similar to the long-sleep phenotype of *cry* mutants (Figure 5J; Figure 5—figure

supplement 4A). Because R6GAL4 is also expressed in several other cells in the brain,

we combined *Pdf*GAL80 to verify that the sleep-enhancing effect is caused by

over-activation of the s-LNvs (Helfrich-Forster et al., 2007). Indeed, we no longer

observed the long-sleep phenotype when GAL4 expression is inhibited in the PDF

neurons by GAL80 (Figure 5J; Figure 5—figure supplement 4A). Moreover, we found

that *cry* mutation can no longer exert effects on sleep when the s-LNvs are activated,

further validating that the s-LNvs mediate the influences of CRY on sleep (Figure 5K;

Figure 5—figure supplement 4B). Because CRY has been shown to regulate

membrane depolarization and input resistance together with the redox sensor of the

voltage-gated potassium channel β-subunit HYPERKINETIC (HK), we tested

whether CRY and HK also function together in the s-LNvs to regulate arousal

(Agrawal et al., 2017; Fogle et al., 2015). We found that while over-expressing *hk* in

the s-LNvs of WT flies does not alter sleep duration under 4L20D, over-expressing it

in *cry* mutants significantly increases sleep (Figure 5L; Figure 5—figure supplement

4C). This genetic interaction indicates that CRY and HK cooperate to regulate

wakefulness, possibly by modulating the electric activity of the s-LNvs.

Next, we expressed syt-GFP in the s-LNvs and observed GFP signal at the soma

of the l-LNvs, indicating that the s-LNvs send axonal terminals to the l-LNvs (Figure

6A). *trans*-Tango technique further demonstrated that the s-LNvs project to form

synaptic connections with the l-LNvs (Figure 6B). Moreover, when we disrupted

GABA transmission in the s-LNvs by knocking down *VGAT* or *Gad1*, this reduced

GABA intensity in the l-LNvs, implicating that the s-LNvs release GABA onto the

l-LNvs (Figure 6C-E). At the behavioral level, these manipulations supress the

long-sleep phenotype of *cry* mutation (Figure 6F-I).

Taken together, these results suggest that *cry* deficiency in the s-LNvs results in

increased neuronal activity and thus enhanced GABA release at the l-LNvs via direct

synaptic projections, ultimately leading to decreased wakefulness and increased sleep.

Short photoperiod reduces GABA level at the l-LNvs

Our findings thus far point to an inhibitory role of CRY on the GABAergic tone, and

this in turn removes the inhibition on the neural activities of the l-LNvs. Since CRY is

degraded by light, we hypothesized that CRY should exert a stronger effect on GABA

under short photoperiod (Emery et al., 1998)**.** Consistently, GABA intensity does

appear to be down-regulated at the l-LNvs under 4L20D vs. 12L12D (Figure 7A and

B). On the other hand, GABA intensity does not exhibit photoperiod-dependent

alteration in the s-LNvs (Figure 7C and D). Presumably, this reduced GABA level at

the l-LNvs under 4L20D can result in elevated activation of these cells and increased

wakefulness. In line with this, WT flies display shortened sleep duration under 4L20D

compared to 12L12D (Figure 7E and F). This sleep reduction is a result of decreased

sleep bout length but not bout number, while wake activity is not altered by

photoperiod (Figure 7G-I). These observations indicate that short photoperiod

hampers sleep maintenance, similar to the effects of CRY.

Discussion

Previous studies have shown that CRY mediates light-induced electrical activity of the

l-LNvs and acute arousal, which indicates that CRY can immediately act to promote

wake and terminate sleep in response to light pulse (Fogle et al., 2015; Sheeba et al.,

2008). At the molecular level, this is believed to be accomplished by a direct coupling

of light-activated CRY with HK in the l-LNvs (Fogle et al., 2015). Here we found that

CRY promotes extended wakefulness under short photoperiod by functioning as an

inhibitor of GABAergic tone. In contrast to the previously characterized roles of CRY

which are activated by light, we believe this novel function we identify here reflects a

role for CRY in the dark. Under short photoperiods, CRY acts in the dark to inhibit

GABA signaling and thus promote wakefulness, leading to reduced sleep during the

dark phase compared to longer photoperiods. Indeed, CRY is an ideal signal for

conveying photoperiodic information, for it is degraded by light and only accumulates

during darkness (Emery et al., 1998). It can measure the length of the day/night and

thus instruct downstream signaling components to modulate photoperiod-dependent

processes. Consistent with this idea, the largest increases of sleep in *cry* mutants occur

immediately after lights-off and prior to lights-on (Fig 1B), indicating that CRY

functions to promote wakefulness during these time windows which are exactly the

time windows that would be affected by photoperiod changes (i.e. there will be light

during these time windows under longer photoperiods). In other words, CRY appears

to act at the time of the day that is most sensitive to photoperiodic changes, which fits

perfectly with the idea that it serves as an instructive signal of day/night length.

While EOS and NipA treatment lengthen sleep duration as previously reported,

we are somewhat surprised by the observation that knocking down *Gad1* or *VGAT* in

GABAergic neurons also extends sleep duration (Ki and Lim, 2019; Leal and

Neckameyer, 2002). We reason this may be due to some sort of over-compensation

induced by chronic GABA deficiency to maintain excitation/inhibition balance, as

previous studies have reported that the amplitude of glutamatergic current is

substantially down-regulated in *Gad1* mutant flies and increased in *Gad1*

over-expression flies (Featherstone et al., 2000), (Featherstone et al., 2002).

Consistent with this idea, we noticed when *cry* is knocked down in *Gad1/VGAT* RNAi

flies, sleep duration is shortened and comparable to that of the controls. This is

probably because *cry* depletion enhances the GABAergic tone, thus normalizing

GABA signaling in these flies which results in normal sleep duration.

We acknowledge that THIP treatment leads to prominent lengthening of sleep

duration, and thus it is possible in this case *cry* deficiency no longer increases sleep

duration due to a ceiling effect rather than epistatic interaction. Nonetheless,

considering that CBZ and *rdl* mutation also block the effect of *cry* deficiency on sleep

duration, it is highly likely that GABA-A receptor mediates the influences of CRY on

sleep. One caveat is that the $rdl^{MD\text{-}RR}$ mutation has been shown to diminish the

desensitization of GABA-A receptor and is thus believed to be a gain-of-function

allele, but here we found that it eliminates the effect of *cry* RNAi on sleep (Zhang et

al., 1994). Since treatment with GABA-A agonist THIP leads to substantially

increased sleep while under the same condition $rdl^{MD\text{-}RR}$ mutation does not appear to

significantly alter sleep duration, we suspect that similar to *Gad1/VGAT* RNAi,

chronic enhancement of GABA-A function associated with $rdl^{MD\text{-}RR}$ mutation may

also trigger some kind of compensatory mechanism that counteracts this increased

GABA-A activity. Consequently, the influences of *cry* RNAi on sleep are blocked. Of

course, a lot more in depth characterizations are required to elucidate these issues.

The s-LNvs have been reported to receive GABAergic inputs possibly via

GABA-B receptors, but have not been shown to be able to synthesize GABA (Dahdal

et al., 2010; Hamasaka et al., 2005). One study using cell-type specific gene

expression profiling demonstrates *Gad1* and *VGAT* expression in both s-LNvs and

l-LNvs, although with relatively low signal (Nagoshi et al., 2010). Here we observed

that *Gad1*GAL4 is expressed in the s-LNvs, and their GABA intensity is reduced

when we use R6GAL4 to knock down *VGAT* in these cells. R6GAL4 drives

prominent expression in the s-LNvs with very little if any expression in the l-LNvs,

and weaker (and not very consistent) expression in several other neurons in the

protocerebrum, pars intercerebralis and subesophageal area which are all believed to

lie outside of the circadian neuron network (Helfrich-Forster et al., 2007). Therefore,

we reason that the alteration of GABA signal associated with knocking down *VGAT*

should arise from VGAT deficiency within the s-LNvs. Taken together, these results

suggest that the s-LNvs are GABAergic. Although knocking down *VGAT* or *Gad1* in

these cells can suppress the long sleep phenotype of *cry* mutants, knocking down *cry*

in these cells only lead to a modest lengthening of sleep duration. Moreover,

inhibiting *cry* RNAi expression in PDF neurons does not eliminate the long-sleep

phenotype of *Gad1*GAL4/UAS*cry*RNAi flies. Therefore, we suspect that *cry*

deficiency in other GABAergic neurons are also required for the long-sleep

phenotype. Given that the s-LNvs are known to express CRY and appear to be

GABAergic based on our findings here, we believe that CRY acts at least in part in

the s-LNvs to promote wakefulness under short photoperiod.

The molecular mechanism by which CRY down-regulates the GABAergic tone

remains unclear. Besides conducting light-driven depolarization via HK, CRY has

also been shown to act in synergy with HK to prevent membrane input resistance

from falling to a low level in larval salivary glands (Agrawal et al., 2017; Fogle et al.,

2015). In contrast to previous studies, here we found that lack of CRY increases the

activity of the s-LNvs. Instead of functioning in synergy with HK, CRY appears to act

in the opposite direction of HK as over-expressing *hk* enhances the long-sleep

phenotype caused by *cry* mutation. We reason that the coupling between CRY and HK

as well as their influences on the electric activity in the s-LNvs may be different from

that of the l-LNvs and larval salivary glands. Nonetheless, our results also support an

interaction between CRY and HK to promote arousal. We suspect that CRY acts via

HK to inhibit the activity of the s-LNvs, which results in decreased GABA release and

dis-inhibition of the l-LNvs. Extensive further investigations will be needed to

elucidate the mechanism by which CRY regulates the activity of the s-LNvs.

In conclusion, here we describe a CRY-controlled GABAergic circuitry involving

the l-LNvs and the s-LNvs that adjusts sleep duration in adaptation to changes in day

length and propose a mechanistic explanation regarding how this circuitry functions

(Fig 7J). Under short photoperiods, more CRY accumulates and inhibits the activity of

the GABAergic s-LNvs, leading to a dis-inhibition of the l-LNvs which can release

more PDF and promote arousal. Under longer photoperiods, on the other hand, less

CRY can accumulate and thus the s-LNvs will exert more inhibitory influences on the

l-LNvs, leading to decreased release of PDF and wakefulness. Notably, almost all

neurons in the mammalian pacemaker, the suprachiasmatic nucleus, are GABAergic,

and GABA/GABA-A signaling have been shown to mediate neuronal coupling in

response to photoperiod changes (Ono et al., 2021). We believe a similar GABAergic

circuitry may exist in the mammalian system that adjusts sleep/wakefulness to

photoperiodic changes.

Materials and Methods

Fly strains

All strains were obtained from Bloomington *Drosophila* Stock Center, Vienna

Drosophila Resource Center and TsingHua Fly Center or as gifts from colleagues.

Except for *Gad1*GAL4-2, all neurotransmitter related GAL4 lines were generated in

Dr. Yi Rao's laboratory (Deng et al., 2019). The following fly strains were used in this

study: cry^b (BDSC:80921), per^0 (BDSC:80917), tim^0 (BDSC:80922), clk^{jrk}

(BDSC:80927), cyc^0 (BDSC:80929), Pdf^{01} (BDSC:26654), $Pdfr^{han5304}$ (BDSC:33068),

UAS-*cry*RNAi-1 (THU4761), UAS-*cry*RNAi-2 (THU5148), cry^{03} (Dolezelova et al.,

2007), UAS-*gad1*RNAi-1 (THU2920), Rdl^{MD-RR} (BDSC:1675), UAS-*VGAT*RNAi-1

(THU4303), UAS-*VGAT*RNAi-2 (V45916), UAS-*hk* (BDSC:86270), isogenic w^{1118}

(BDSC:5905), UAS-*NachBac* (BDSC:9469), *Pdf*GAL4 (BDSC:6899), *VGAT*GAL4

(BDSC:58980), UAS-*TrpA1* (BDSC:26263), UAS-*Kir2.1* (BDSC:6596)*,* UAS-*dcr2*

(V6009), UAS-syt*GFP* (BDSC:6925), UAS-nls*GFP* (BDSC:4775),

UAS-*myr*GFP,QUAS-mtdtomato(3XHA);*Trans*-tango (BDSC:77480) ,

13XLexAop2-*post-t*-GRASP,20XUAS-*pre*-mGRASP (BDSC:79040),

LexAop-rCD2-RFP (Perisse et al., 2013), *Gad1*GAL4-1, *Gad1*GAL4-2

(BDSC:51630), *Dop2R*GAL4, *TH*GAL4, *Trh*GAL4, *AdoR*GAL4, *DAT*GAL4,

*GABAR1*GAL4, *GABAR3*GAL4, *SerT*GAL4, *GluRIA*GAL4, R6GAL4

(Helfrich-Forster et al., 2007), c929GAL4 (Park et al., 2008), *R10H10*GAL4

(Sekiguchi et al., 2020)*, R78G01*GAL4 (Yoshii et al., 2015), *Pdf*GAL80 (Stoleru et al.,

2004), UAS-GCaMP6m-Tdtomato (Chen et al., 2013) and *Pdf*-LexA (Shang et al.,

2008). All flies used for sleep monitoring were backcrossed with the isogenic w^{1118}

strain for at least 5 times except for cry^{03} which was backcrossed three times. All

experiments were conducted in male flies unless otherwise specified.

Fly sleep monitoring and analysis

Flies were raised on standard cornmeal-yeast-sucrose medium and kept in 12L12D at

25°C until behavior monitoring. ~3 to 4-day-old flies were entrained under different

photoperiods at 25℃ for 4 days, and then their activities in the next 3 days were

analyzed. Sleep is defined as 5 min consecutive inactivity. Sleep was analyzed with

Counting Macro written in Excel (Microsoft) following previously published protocol

(Pfeiffenberger et al., 2010). Flies were fed with agar-sucrose food (2% agar, 5%

sucrose) during entire sleep monitoring. TrpA1 flies were raised at 21°C and baseline

sleep was monitored at 21°C. Temperature was then raised at lights on to 29°C for

further sleep monitoring.

Drug treatment

For pharmacological experiments, drugs were mixed in the fly food at the following

concentrations. For nipecotic acid (10 mg/ml, Sigma) and EOS (10 mM, Sigma),

drugs were fed during the entire sleep monitoring. For THIP (10 ug/ml, Sigma),

SKF-97541 (10 ug/ml, Tocris) and CBZ (0.15 mg/ml, Sinopharm Chemical Reagent),

drugs were fed for 1 day after baseline sleep monitoring. The same amount of solvent

was added into the fly food as vehicle control.

RNA extraction and quantitative real-time PCR (qRT-PCR)

Approximately 50 5-day-old flies were collected and frozen immediately on dry ice.

Fly heads were isolated and homogenized in Trizol reagent (Life Technologies). Total

RNA was extracted and qRT-PCR conducted following our previously published

procedures (Bu et al., 2019). The following primers were used to quantify mRNA

level, *cry* F: TGCAGGTACCAAGAATGTGG, R: GTCCACGTCCATCAGTTGC.

Gad1 F: TGCCACCACATTGAAGTACC, R: GGTGAACATGTTGGTGTTCG.

VGAT F: ACGGCTTTAGGCAAGGTAGC, R: TTGGAATTCGTCGATTTTGC.

Immunostaining

Male flies were entrained for 3 days under indicated photoperiod and collected on

Day 4. flies were anesthetized with CO_2 and dissected and fixed with 4%

paraformaldehyde diluted in PBS at the indicated time points. The brains were then

fixed with 4% paraformaldehyde for 30 minutes. Samples were washed with PBT

(PBS with 0.3% TritonX-100) for 3×10min. Then samples were incubated in PBS

with 1% TritonX-100 for 20 min at room temperature. Brain samples were then

blocked in PBT with 5% fetal bovine serum (Hyclone) for 30 minutes and

subsequently incubated with mouse anti-PDF (1:100, DSHB), rabbit anti-GABA

(1:200, Sigma), and anti-HA (1:100, DSHB) for 2-4 days at 4°C (4 days for

anti-GABA and 2 days for the other antibodies). After PBT rinses for 3 times, the

brains were incubated with donkey anti-mouse Alexa Fluor-594 (1:1000, Life

Technologies), donkey anti-rabbit Alexa Fluor-488 (1:1000, Abcam) and donkey

anti-rabbit Alexa Fluor-647 (1:1000, Abcam) overnight at 4°C. Then the brains were

rinsed 3 times in PBS and mounted and imaged using Olympus FV3000 confocal

microscope with a 60× objective lens. The intensity of GABA signal was quantified

by ImageJ software. For each cell, the image slice with the strongest signal in the Z

stack was selected and average intensity was quantified. A region on the same image

slice was then selected as background and its intensity value was subtracted from the

average intensity value of the cell. This subtracted value was subsequently normalized

to the average intensity of the control group as indicated in figure legends. Sample

sizes in the legends indicate number of cells examined per genotype. For *trans*-Tango

experiment, offsprings were raised at 18°C for 3-4 weeks before HA and PDF

immunostaining.

Calcium imaging

For live imaging experiments (GCaMP and tdTomato), flies 2-3 days old were

collected into tubes with standard food and entrained under LD for 3 days. Flies were

anesthetized with CO_2 and brains were dissected at the indicated time points in

Drosophila adult hemolymph-like saline solution. The dissected brain samples were

put on glass slide and sealed with cover slide. The duration for dissection and

microscopy should be completed within 0.5 hours for each time point. Images were

captured with Olympus FV3000 confocal microscopy with a 20× objective lens. The

intensity of GCaMP and tdTomato signals were quantified by ImageJ software.

Sample sizes in the legends indicate number of cells examined per genotype.

Statistical Analysis

For data that fit normal distribution, two-tailed Student's *t*-test (Microsoft Excel) was

used to compare the difference between two genotypes. For data that do not fit normal

distribution, Mann-Whitney test (Prism Graphpad) was used to compare the

difference between two genotypes. For multiple comparisons, one-way ANOVA with

Bonferroni multiple comparison test (Prism Graphpad) was used. Sample size,

statistical test, and significance values are indicated in figure legends. Sample size is

determined based on previous studies with similar experimental assays.

Acknowledgements

We would like to thank Drs. Chang Liu, Yi Rao, Liming Wang, Junhai Han, Zhihua

Liu and Yi Zhong for kindly providing flies used in this study. We would also like to

thank Dr. Chang Liu for helpful advice on GABA immunostaining. This work was

supported by grants from the Ministry of Science and Technology of China STI

2030-Major Projects (2021ZD0203202 and 2024YFA1803201), Natural Science

Foundation of China (82530044 and 32341021) and Science and Technology

Department of Hubei Province (2022CFA049) to Luoying Zhang and grant from the

Natural Science Foundation of China (32300984) to Lixia Chen.

Author contributions

488   L.C. and L.Z. designed the experiments. L.C. and D.T. conducted the experiments.

489   L.C., C.S. and L.Z. wrote the paper.

Disclosure and competing interests statement

492   The authors declare no competing interests.

Data Availability

495   This study includes no data deposited in external repositories.

References

Agosto, J., Choi, J.C., Parisky, K.M., Stilwell, G., Rosbash, M., and Griffith, L.C. (2008). Modulation of GABA(A) receptor desensitization uncouples sleep onset and maintenance in *Drosophila*. Nat Neurosci *11*, 354-359.

Agrawal, P., Houl, J.H., Gunawardhana, K.L., Liu, T., Zhou, J., Zoran, M.J., and Hardin, P.E. (2017). *Drosophila* CRY Entrains Clocks in Body Tissues to Light and Maintains Passive Membrane Properties in a Non-clock Body Tissue Independent of Light. Curr Biol *27*, 2431-2441 e2433.

Allada, R., and Chung, B.Y. (2010). Circadian organization of behavior and physiology in *Drosophila*. Annu Rev Physiol *72*, 605-624.

Allada, R., White, N.E., So, W.V., Hall, J.C., and Rosbash, M. (1998). A mutant *Drosophila* homolog of mammalian *Clock* disrupts circadian rhythms and transcription of *period* and *timeless*. Cell *93*, 791-804.

Baines, R.A., Uhler, J.P., Thompson, A., Sweeney, S.T., and Bate, M. (2001). Altered electrical properties in *Drosophila* neurons developing without synaptic transmission. J Neurosci *21*, 1523-1531.

Benito, J., Houl, J.H., Roman, G.W., and Hardin, P.E. (2008). The blue-light photoreceptor CRYPTOCHROME is expressed in a subset of circadian oscillator neurons in the *Drosophila* CNS. J Biol Rhythms *23*, 296-307.

Bu, B., He, W., Song, L., and Zhang, L. (2019). Nuclear Envelope Protein MAN1 Regulates the *Drosophila* Circadian Clock via *Period*. Neurosci Bull *35*, 969-978.

Busza, A., Emery-Le, M., Rosbash, M., and Emery, P. (2004). Roles of the two *Drosophila* CRYPTOCHROME structural domains in circadian photoreception. Science *304*, 1503-1506.

Ceriani, M.F., Darlington, T.K., Staknis, D., Mas, P., Petti, A.A., Weitz, C.J., and Kay, S.A. (1999). Light-dependent sequestration of TIMELESS by CRYPTOCHROME. Science *285*, 553-556.

Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V.*, et al.* (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature *499*, 295-300.

Chung, B.Y., Kilman, V.L., Keath, J.R., Pitman, J.L., and Allada, R. (2009). The GABA(A) receptor RDL acts in peptidergic PDF neurons to promote sleep in *Drosophila*. Curr Biol *19*, 386-390.

Dahdal, D., Reeves, D.C., Ruben, M., Akabas, M.H., and Blau, J. (2010). *Drosophila* pacemaker neurons require G protein signaling and GABAergic inputs to generate twenty-four hour behavioral rhythms. Neuron *68*, 964-977.

Deng, B., Li, Q., Liu, X., Cao, Y., Li, B., Qian, Y., Xu, R., Mao, R., Zhou, E., Zhang, W.*, et al.* (2019). Chemoconnectomics: Mapping Chemical Transmission in *Drosophila*. Neuron *101*, 876-893 e874.

Dolezelova, E., Dolezel, D., and Hall, J.C. (2007). Rhythm defects caused by newly engineered null mutations in *Drosophila*'s *cryptochrome* gene. Genetics *177*, 329-345.

Emery, P., So, W.V., Kaneko, M., Hall, J.C., and Rosbash, M. (1998). CRY, a *Drosophila* clock and light-regulated cryptochrome, is a major contributor to circadian rhythm resetting and photosensitivity. Cell *95*, 669-679.

Enell, L., Hamasaka, Y., Kolodziejczyk, A., and Nassel, D.R. (2007). gamma-Aminobutyric acid (GABA) signaling components in *Drosophila*: immunocytochemical localization of GABA(B) receptors in relation to the GABA(A) receptor subunit RDL and a vesicular GABA transporter. J Comp

Neurol *505*, 18-31.

Featherstone, D.E., Rushton, E., and Broadie, K. (2002). Developmental regulation of glutamate receptor field size by nonvesicular glutamate release. Nat Neurosci *5*, 141-146.

Featherstone, D.E., Rushton, E.M., Hilderbrand-Chae, M., Phillips, A.M., Jackson, F.R., and Broadie, K. (2000). Presynaptic glutamic acid decarboxylase is required for induction of the postsynaptic receptor field at a glutamatergic synapse. Neuron *27*, 71-84.

ffrench-Constant, R.H., Steichen, J.C., Rocheleau, T.A., Aronstein, K., and Roush, R.T. (1993). A single-amino acid substitution in a gamma-aminobutyric acid subtype A receptor locus is associated with cyclodiene insecticide resistance in *Drosophila* populations. Proc Natl Acad Sci U S A *90*, 1957-1961.

Fogle, K.J., Baik, L.S., Houl, J.H., Tran, T.T., Roberts, L., Dahm, N.A., Cao, Y., Zhou, M., and Holmes, T.C. (2015). CRYPTOCHROME-mediated phototransduction by modulation of the potassium ion channel beta-subunit redox sensor. Proc Natl Acad Sci U S A *112*, 2245-2250.

Golden, R.N., Gaynes, B.N., Ekstrom, R.D., Hamer, R.M., Jacobsen, F.M., Suppes, T., Wisner, K.L., and Nemeroff, C.B. (2005). The efficacy of light therapy in the treatment of mood disorders: a review and meta-analysis of the evidence. Am J Psychiatry *162*, 656-662.

Hamada, F.N., Rosenzweig, M., Kang, K., Pulver, S.R., Ghezzi, A., Jegla, T.J., and Garrity, P.A. (2008). An internal thermal sensor controlling temperature preference in *Drosophila*. Nature *454*, 217-220.

Hamasaka, Y., Wegener, C., and Nassel, D.R. (2005). GABA modulates *Drosophila* circadian clock neurons via GABAB receptors and decreases in calcium. J Neurobiol *65*, 225-240.

Helfrich-Forster, C., Shafer, O.T., Wulbeck, C., Grieshaber, E., Rieger, D., and Taghert, P. (2007). Development and morphology of the clock-gene-expressing lateral neurons of *Drosophila melanogaster*. J Comp Neurol *500*, 47-70.

Hyun, S., Lee, Y., Hong, S.T., Bang, S., Paik, D., Kang, J., Shin, J., Lee, J., Jeon, K., Hwang, S., *et al.* (2005). *Drosophila* GPCR *Han* is a receptor for the circadian clock neuropeptide PDF. Neuron *48*, 267-278.

Jackson, F.R., Newby, L.M., and Kulkarni, S.J. (1990). *Drosophila* GABAergic systems: sequence and expression of glutamic acid decarboxylase. J Neurochem *54*, 1068-1078.

Ki, Y., and Lim, C. (2019). Sleep-promoting effects of threonine link amino acid metabolism in *Drosophila* neuron to GABAergic control of sleep drive. eLife *8*.

Konopka, R.J., and Benzer, S. (1971). Clock mutants of *Drosophila melanogaster*. Proc Natl Acad Sci U S A *68*, 2112-2116.

Kurlansik, S.L., and Ibay, A.D. (2012). Seasonal affective disorder. Am Fam Physician *86*, 1037-1041.

Leal, S.M., and Neckameyer, W.S. (2002). Pharmacological evidence for GABAergic regulation of specific behaviors in *Drosophila melanogaster*. J Neurobiol *50*, 245-261.

Lear, B.C., Merrill, C.E., Lin, J.M., Schroeder, A., Zhang, L., and Allada, R. (2005). A G protein-coupled receptor, groom-of-PDF, is required for PDF neuron action in circadian behavior. Neuron *48*, 221-227.

Lin, F.J., Song, W., Meyer-Bernstein, E., Naidoo, N., and Sehgal, A. (2001). Photic signaling by *cryptochrome* in the *Drosophila* circadian system. Mol Cell Biol *21*, 7287-7294.

Matsuda, K., Hosie, A.M., Buckingham, S.D., Squire, M.D., Baylis, H.A., and Satella, D.B. (1996). pH-Dependent actions of THIP and ZAPA on an ionotropic *Drosophila melanogaster* GABA receptor. Brain Res *739*, 335-338.

Mertens, I., Vandingenen, A., Johnson, E.C., Shafer, O.T., Li, W., Trigg, J.S., De Loof, A., Schoofs, L.,

and Taghert, P.H. (2005). PDF receptor signaling in *Drosophila* contributes to both circadian and

geotactic behaviors. Neuron *48*, 213-219.

Mezler, M., Müller, T., and Raming, K. (2001). Cloning and functional expression of GABA receptors

from *Drosophila*. European Journal of Neuroscience *13*, 477-486.

Nagoshi, E., Sugino, K., Kula, E., Okazaki, E., Tachibana, T., Nelson, S., and Rosbash, M. (2010).

Dissecting differential gene expression within the circadian neuronal circuit of. Nat Neurosci *13*,

60-U220.

Nitabach, M.N., Wu, Y., Sheeba, V., Lemon, W.C., Strumbos, J., Zelensky, P.K., White, B.H., and

Holmes, T.C. (2006). Electrical hyperexcitation of lateral ventral pacemaker neurons desynchronizes

downstream circadian oscillators in the fly circadian circuit and induces multiple behavioral periods. J

Neurosci *26*, 479-489.

Ono, D., Honma, K.I., and Honma, S. (2021). GABAergic mechanisms in the suprachiasmatic nucleus

that influence circadian rhythm. J Neurochem *157*, 31-41.

Parisky, K.M., Agosto, J., Pulver, S.R., Shang, Y., Kuklin, E., Hodge, J.J., Kang, K., Liu, X., Garrity,

P.A., Rosbash, M.*, et al.* (2008). PDF cells are a GABA-responsive wake-promoting component of the

Drosophila sleep circuit. Neuron *60*, 672-682.

Park, D., Veenstra, J.A., Park, J.H., and Taghert, P.H. (2008). Mapping peptidergic cells in *Drosophila*:

where DIMM fits in. PLoS One *3*, e1896.

Perisse, E., Yin, Y., Lin, A.C., Lin, S.W., Huetteroth, W., and Waddell, S. (2013). Different Kenyon Cell

Populations Drive Learned Approach and Avoidance in. Neuron *79*, 945-956.

Pfeiffenberger, C., Lear, B.C., Keegan, K.P., and Allada, R. (2010). Processing sleep data created with

the *Drosophila* Activity Monitoring (DAM) System. Cold Spring Harb Protoc *2010*, pdb prot5520.

Renn, S.C., Park, J.H., Rosbash, M., Hall, J.C., and Taghert, P.H. (1999). A *pdf* neuropeptide gene

mutation and ablation of PDF neurons each cause severe abnormalities of behavioral circadian rhythms

in *Drosophila*. Cell *99*, 791-802.

Rosenthal, N.E., Sack, D.A., Gillin, J.C., Lewy, A.J., Goodwin, F.K., Davenport, Y., Mueller, P.S.,

Newsome, D.A., and Wehr, T.A. (1984). Seasonal affective disorder. A description of the syndrome and

preliminary findings with light therapy. Arch Gen Psychiatry *41*, 72-80.

Sehgal, A., Price, J.L., Man, B., and Young, M.W. (1994). Loss of circadian behavioral rhythms and

per RNA oscillations in the *Drosophila* mutant *timeless*. Science *263*, 1603-1606.

Sekiguchi, M., Inoue, K., Yang, T., Luo, D.G., and Yoshii, T. (2020). A Catalog of GAL4 Drivers for

Labeling and Manipulating Circadian Clock Neurons in. J Biol Rhythm *35*, 207-213.

Shang, Y., Griffith, L.C., and Rosbash, M. (2008). From the Cover: Feature Article: Light-arousal and

circadian photoreception circuits intersect at the large PDF cells of the *Drosophila* brain. Proc Natl

Acad Sci U S A *105*, 19587-19594.

Shearin, H.K., Quinn, C.D., Mackin, R.D., Macdonald, I.S., and Stowers, R.S. (2018). t-GRASP, a

targeted GRASP for assessing neuronal connectivity. J Neurosci Methods *306*, 94-102.

Sheeba, V., Gu, H., Sharma, V.K., O'Dowd, D.K., and Holmes, T.C. (2008). Circadian- and

light-dependent regulation of resting membrane potential and spontaneous action potential firing of

Drosophila circadian pacemaker neurons. J Neurophysiol *99*, 976-988.

Shiga, Y., Tanaka-Matakatsu, M., and Hayashi, S. (1996). A nuclear GFP/ beta-galactosidase fusion

protein as a marker for morphogenesis in living *Drosophila*. Dev Growth Differ *38*, 99-106.

Stanewsky, R., Kaneko, M., Emery, P., Beretta, B., Wagner-Smith, K., Kay, S.A., Rosbash, M., and

Hall, J.C. (1998). The *cry^b* mutation identifies cryptochrome as a circadian photoreceptor in *Drosophila*.

Cell *95*, 681-692.

Stoleru, D., Peng, Y., Agosto, J., and Rosbash, M. (2004). Coupled oscillators control morning and

evening locomotor behaviour of *Drosophila*. Nature *431*, 862-868.

Talay, M., Richman, E.B., Snell, N.J., Hartmann, G.G., Fisher, J.D., Sorkac, A., Santoyo, J.F.,

Chou-Freed, C., Nair, N., Johnson, M.*, et al.* (2017). Transsynaptic Mapping of Second-Order Taste

Neurons in Flies by trans-Tango. Neuron *96*, 783-795 e784.

Wescott, D.L., Soehner, A.M., and Roecklein, K.A. (2020). Sleep in seasonal affective disorder. Curr

Opin Psychol *34*, 7-11.

Wood, S., and Loudon, A. (2014). Clocks for all seasons: unwinding the roles and mechanisms of

circadian and interval timers in the hypothalamus and pituitary. J Endocrinol *222*, R39-59.

Yoshii, T., Hermann-Luibl, C., Kistenpfennig, C., Schmid, B., Tomioka, K., and Helfrich-Förster, C.

(2015). Cryptochrome-Dependent and -Independent Circadian Entrainment Circuits in. Journal of

Neuroscience *35*, 6131-6141.

Yoshii, T., Todo, T., Wulbeck, C., Stanewsky, R., and Helfrich-Forster, C. (2008). Cryptochrome is

present in the compound eyes and a subset of *Drosophila*'s clock neurons. J Comp Neurol *508*,

952-966.

Zhang, H.G., ffrench-Constant, R.H., and Jackson, M.B. (1994). A unique amino acid of the

Drosophila GABA receptor with influence on drug sensitivity by two mechanisms. J Physiol *479 (Pt*

1), 65-75.

Zhang, Y.Q., Rodesch, C.K., and Broadie, K. (2002). Living synaptic vesicle marker:

synaptotagmin-GFP. Genesis *34*, 142-145.

Figure Legend

Figure 1. *cry* mutation increases sleep duration selectively under short photoperiods. (A) Daily sleep duration in male circadian clock gene mutants per^0, tim^0, clk^{jrk}, cyc^0, and cry^b compared to wild-type (WT) males under 4L20D, 8L16D, 12L12D, 16L8D, 20L4D (12L12D, n = 86, 63, 91, 14, 91, 53 flies; 8L16D, n = 21, 31, 31, 10, 32, 31 flies; 12L12D, n = 31, 14, 32, 14, 27, 32 flies; 16L8D, n = 36, 30, 33, 17, 36, 30 flies; 20L4D, n = 42, 55, 83, 26, 91, 89 flies). (B) Average sleep traces of cry^b and WT male flies under 4L20D. (C-F) The daily sleep duration (C), waking activity (D), sleep bout number (E), average sleep bout length (F) for cry^b and WT female and male flies under 4L20D (n = 56, 88, 58, 91 flies). For (B), statistical differences between cry^b and WT is determined by paired two-tailed *t*-test, *$P < 0.001$. For A and C-E, statistical differences is determined by two-tailed Student's *t*-test, ***$P < 0.001$. Error bars represent standard error of the mean (SEM). ZT, Zeitgeber Time (ZT0 is the time of lights on).

Figure 1—figure supplement 1. *cry* knock-out mutation lengthens sleep duration under 4L20D. Daily sleep duration of male WT flies and *cry* knock-out flies, monitored under 4L20D (n = 53, 35, 47). Student's *t*-test: *$P < 0.05$. Error bars represent SEM.

Figure 2. CRY functions in GABAergic neurons and promotes wakefulness via GABA signaling. (A, B, C) Daily sleep duration of male flies with *cry* knocked down in GABAergic neurons by *VGAT*GAL4 (A) (n = 72, 74, 96, 49, 54 flies), *Gad1*GAL4-1 (B) (n = 29, 26, 29, 23, 32 flies) or *Gad1*GAL4-2 (C) (n = 34, 31, 31, 25, 23 flies) monitored under 4L20D. (D) Daily sleep duration of *cry* knocked down in GABAergic neurons by *Gad1*GAL4 monitored under 12L12D (n = 20, 19, 30, 26, 32, 20 flies). For A-D, One-way ANOVA with Bonferroni multiple comparison test: compared to GAL4 control, ***$P < 0.001$; compared to UAS control, #$P < 0.05$, ###$P < 0.001$. (E) Daily sleep duration of male WT and cry^b flies fed with EOS or NipA under 4L20D (n = 31, 32, 30, 30, 25, 32 flies). Two-tailed Student's *t*-test: compared to WT, ###$P < 0.001$; compared to vehicle control, *$P < 0.05$, ***$P < 0.001$. (F) Daily sleep duration of *cry* RNAi and control flies fed with NipA under 4L20D (n = 72, 74, 95, 49, 54, 23, 24, 26, 15, 24 flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, ***$P < 0.001$; compared to UAS control, ###$P < 0.001$. For comparing to vehicle control, two-tailed Student's *t*-test was used, $$$$P < 0.001$. (G) Daily sleep duration of male flies with *cry* and *gad1* knocked down in GABAergic neurons monitored under 4L20D (n = 72, 73, 24, 33, 45, 33, 27 flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, **$P < 0.01$, ***$P < 0.001$; compared to UAS control, ###$P < 0.001$. For comparing to RNAi control, two-tailed Student's *t*-test was used: compared to UAS*cry*RNAi/GAL4 control, $$$$P < 0.001$; compared to UAS*Gad1*RNAi/GAL4 control, &&&$P < 0.001$. (H) Daily sleep duration of male flies with *cry* and *VGAT* knocked down in GABAergic neurons monitored under 4L20D (n = 72, 73, 46, 41, 45, 66, 34 flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, ***$P < 0.001$; compared to UAS control, #$P < 0.05$, ### $P < 0.001$. For comparing to RNAi control, two-tailed Student's *t*-test was used: compared to UAS*cry*RNAi/GAL4 control, $$$$P < 0.001$; compared to UAS*VGAT*RNAi/GAL4 background, not significant. (I) Daily sleep duration of *cry* mutant flies

with *VGAT* knocked down in GABAergic neurons monitored under 4L20D (n = 72, 59, 20, 23, 69,

22 flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with

Bonferroni multiple comparison test was used: compared to GAL4 control, ***P < 0.01, ***P <

0.001; compared to UAS control, #P < 0.05, ###P < 0.001. For comparison between mutant vs.

control, two-tailed Student's *t*-test was used: compared to WT background, &&&P < 0.001;

compared to U*VGAT*RNAi/GAL4, not significant. Error bars represent SEM; G4, GAL4; U, UAS;

NS, not significant.

Figure 2—figure supplement 1. CRY does not regulate sleep duration via TIM. Daily sleep

duration of male WT flies and flies mutant for *cry* and/or *tim*, monitored under 4L20D (n = 58, 63,

91, 33 flies). Two-tailed Student's *t*-test: compared to WT background, ***P < 0.001; compared to

tim^0, ###P < 0.001. Error bars represent SEM.

Figure 2—figure supplement 2. Screening for anatomical substrates that mediate the effects

of CRY on sleep/wakefulness. (A) Difference in sleep duration between male *cry* RNAi flies

compared to GAL4 and UAS controls under 4L20D (n = 20-92). (B) Plots of relative mRNA

abundance for cry determined by qRT-PCR in whole-head extracts of *cry*RNAi and control flies

under 4L20D (n = 4). The value of the GAL4 control group was set to 1. Mann-Whitney test:

compared to GAL4 control, *P < 0.05; compared to UAS control, #P < 0.05. G4, GAL4; U, UAS.

Figure 2—figure supplement 3. CRY promotes wakefulness in GABAergic neurons. (A-D)

Sleep profile (A), daily waking activity (B), sleep bout number (C), average sleep bout length (D)

of male flies with *cry* knocked down in GABAergic neurons by *VGAT*GAL4 monitored under

4L20D (n = 72, 74, 96, 49, 54 flies). (E-H) Sleep profile (E), daily waking activity (F), sleep bout

number (G), average sleep bout length (H) of male flies with *cry* knocked down in GABAergic

neurons by *Gad1*GAL4-1 monitored under 4L20D (n = 29, 26, 29, 23, 32 flies). (I-L) Sleep profile

(I), daily waking activity (J), sleep bout number (K), average sleep bout length (L) of male flies

with *cry* knocked down in GABAergic neurons by *Gad1*GAL4-2 monitored under 4L20D (n = 34,

31, 31, 25, 23 flies). One-way ANOVA with Bonferroni multiple comparison test: compared to

GAL4 control, ***P < 0.001; compared to UAS control, #P < 0.05, ###P < 0.001.

Figure 2—figure supplement 4. CRY promotes wakefulness via GABA signaling. (A, B) Sleep

profile of male WT and *cry* mutant flies fed with NipA (A) or EOS (B) under 4L20D in Fig 2E.

White box indicates light period while black box indicates dark period. (C) Sleep profile of *cry*

RNAi and control flies fed with NipA under 4L20D in Fig 2F. (D) Sleep profile of male flies with

cry and *gad1* knocked down in GABAergic neurons monitored under 4L20D in Fig 2G. (E) Sleep

profile of male flies with *cry* and *VGAT* knocked down in GABAergic neurons monitored under

4L20D in Fig 2H. (F) Sleep profile of *cry* mutant flies with *VGAT* knocked down in GABAergic

neurons monitored under 4L20D in Fig 2I. (G, H) Plots of relative mRNA abundance for *Gad1* (G)

and *VGAT* (H) determined by qRT-PCR in whole-head extracts of RNAi and control flies under

4L20D (n = 6). The value of the GAL4 control group was set to 1. Mann-Whitney test: compared

to GAL4 control, **P < 0.01; compared to UAS control, ##P < 0.01. (I) Daily sleep duration of

male flies with *cry* and *gfp* knocked down in GABAergic neurons by *Gad1*GAL4 monitored under

4L20D (n = 72, 26, 29, 20, 30, 31, 68, 32, 31, 29, 29). One-way ANOVA with Bonferroni multiple

comparison test: compared to GAL4 control, ***P < 0.001; compared to UAS control, #P < 0.05,

###P < 0.001. Error bars represent SEM; G4, GAL4; U, UAS; NS, not significant.

Figure 3. GABA-A receptor mediates the effects of CRY on sleep/wakefulness. (A, C, D)
Daily sleep duration of male WT and *cry* mutant flies fed with THIP (A) (n = 28, 29, 16, 19 flies),
SKF-97541 (C) (n = 25, 21, 23, 18 flies) or CBZ (D) (n = 28, 25, 31, 29 flies) under 4L20D.
Two-tailed Student's *t*-test: compared to WT, ###$P < 0.001$; compared to vehicle control, ***$P <$
0.001.(B, E) Sleep profile of male WT and *cry* mutant flies fed with THIP (B) or CBZ (E) under
4L20D in A and D, respectively. White box indicates light period while black box indicates dark
period. (F, H) Daily sleep duration of male *cry* RNAi and control flies fed with THIP (F) (n = 72,
74, 95, 49, 54, 23, 30, 27,27, 20 flies) or CBZ (H) (n = 72, 74, 95, 49, 54, 18, 22, 26, 30, 24 flies)
under 4L20D. For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA
with Bonferroni multiple comparison test was used: compared to GAL4 control, ***$P < 0.001$;
compared to UAS control, ##$P < 0.01$, ###$P < 0.001$. For comparing to vehicle control, two-tailed
Student's *t*-test was used, $$$$P < 0.001$. (G, I) Sleep profile of male *cry* RNAi and control flies
fed with THIP (G) or CBZ (I) under 4L20D in F and H, respectively. White box indicates light
period while black box indicates dark period. (J) Daily sleep duration of male rdl^{MD-RR} flies with
cry knocked down in GABAergic neurons under 4L20D, along with relevant controls (n = 72, 73,
66, 44, 45, 33, 45, 54, 38, 33 flies). For comparison between RNAi flies vs. UAS/GAL4 controls,
one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control,
***$P < 0.001$; compared to UAS control, ###$P < 0.001$. For comparing to WT or
UAS*cry*RNAi/GAL4 control, two-tailed Student's *t*-test was used: compared to WT, &$P < 0.05$;
compared to UAS*cry*RNAi/GAL4 control, $$$$P < 0.001$. (K, L) Sleep profile of male rdl^{MD-RR}
flies with *cry* knocked down in GABAergic neurons using *cry*RNAi-1 (K) or *cry*RNAi-2 (L)
under 4L20D in J. Error bars represent SEM; G4, GAL4; U, UAS.

Figure 4. CRY increases the neural activity of the l-LNvs to promote wakefulness. (A-D)
Brains from male WT and *cry* mutant flies dissected at ZT1, 7, 13, 19 under 4L20D (A) or
12L12D (C) are immunostained with GABA (cyan) and PDF (magenta) antisera, and
representative l-LNvs are displayed. Merged signal is shown as yellow. Bar graphs represent
normalized GABA intensity in the l-LNvs under 4L20D (B) (n = 26-56 cells) and 12L12D (D) (n
= 32-128 cells). The average value of the control group at ZT1 is set to 1. (E) Representative live
image of the l-LNvs expressing GCaMP6m and tdTomato using *Pdf*GAL4.Brain samples are
dissected at the indicated time points under 4L20D. (F) Quantification of GCaMP6m signal
intensity normalized to that of tdTomato (n = 25-54 cells). Student's *t*-test: *$P < 0.05$, **$P < 0.01$,
***$P < 0.001$. (G) Daily sleep duration of male flies expressing Kir2.1 in PDF neuron using
*Pdf*GAL4 and controls, monitored under 4L20D (n = 32, 42, 30 flies). One-way ANOVA with
Bonferroni multiple comparison test: compared to GAL4 control, *$P < 0.05$; compared to UAS
control, ###$P < 0.001$. (H, I) Daily sleep duration of male *cry* mutant flies expressing TrpA1 in the
l-LNvs using c929GAL4 (H) (n = 42, 42, 38, 35, 45, 37 flies) or R10H10GAL4 (I) (n = 62, 62, 17,
30, 29, 30 flies) and relevant controls, monitored under 4L20D and 29℃ to activate TrpA1. For
comparison with UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison
test was used: compared to GAL4 control, ***$P < 0.001$; compared to UAS control, ###$P < 0.001$.
For comparison between mutant vs. control, two-tailed Student's *t*-test was used: &$P < 0.05$,
&&&$P < 0.001$. (J, K) Daily sleep duration of *cry* mutant flies expressing NachBac in the l-LNvs
using c929GAL4 (J) (n = 45, 30, 32, 48, 55, 48 flies) or R10H10GAL4 (K) (n = 25, 59, 29, 74, 25,
32 flies) and relevant controls, monitored under 4L20D. For comparison with UAS/GAL4 controls,

one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, *$P < 0.05$, ***$P < 0.001$; compared to UAS control, ###$P < 0.001$. For comparison between mutant vs. control, two-tailed Student's *t*-test was used: &$P < 0.05$, &&&$P < 0.001$. (L, M) Daily sleep duration of male Pdf^{01}-cry^{b} (L) (n = 58, 91, 89, 31 flies) and $Pdfr^{han5304}$;cry^{b} (M) (n = 58, 91, 31, 31 flies) mutants along with relevant controls, monitored under 4L20D. Two-tailed Student's *t*-test: compared to WT background, ***$P < 0.001$; compared to Pdf^{01}or $Pdfr^{han5304}$, ###$P < 0.001$; compared to cry^{b}, &&$P < 0.01$, &&&$P < 0.001$. The scale bar represents 15 µm. Error bars represent SEM. G4, GAL4; U, UAS.

Figure 4—figure supplement 1. GABAergic neurons project to and form synapses with the l-LNvs. (A) l-LNvs of flies expressing syt-GFP in GABAergic neurons using *Gad1*Gal4 maintained under 4L20D and immunostained with PDF antisera. (B) l-LNvs of flies expressing Pre-mGRASP in GABAergic neurons using *Gad1*Gal4 and t-GRASP in the l-LNvs using *Pdf*LexA maintained under 4L20D. Brains are immunostained with PDF antisera. (C) l-LNvs of flies expressing trans-Tango in GABAergic neurons using *Gad1*Gal4 maintained under 4L20D and immunostained with HA and PDF antisera. (D) Daily sleep duration of *cry* mutant flies with *rdl* knocked down in the l-LNvs using R78G01GAL4 and relevant controls, monitored under 4L20D (n = 31, 31, 64, 32, 31, 32 flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, *$P < 0.05$; compared to UAS control, ###$P < 0.001$. For comparison between mutant vs. control, two-tailed Student's *t*-test was used: compared to WT background, &$P < 0.05$, &&&$P < 0.001$; compared to U*rdl*RNAi/GAL4, not significant. The scale bar represents 15 µm. Error bars represent SEM. G4, GAL4; U, UAS; LexAop, LexA operator.

Figure 4—figure supplement 2. CRY promote wakefulness by impinging on the excitability of l-LNv and PDF signaling. (A) Sleep profile of male flies expressing Kir2.1 in PDF neuron using *Pdf*GAL4 and controls, monitored under 4L20D in Fig 4G. White box indicates light period while black box indicates dark period. (B, C) Sleep profile of male cry mutant flies expressing TrpA1 in the l-LNvs using c929GAL4 (B) or R10H10GAL4 (C) and relevant controls in Fig 4H and I, monitored under 4L20D and 29°C to activate TrpA1. (D, E) Sleep profile of cry mutant flies expressing NachBac in the l-LNvs using c929GAL4 (D) or R10H10GAL4 (E) and relevant controls in Fig 4J and K, monitored under 4L20D. (F, G) Daily sleep duration of male Pdf^{01}-cry^{b} (D) and $Pdfr^{han5304}$;cry^{b} (E) mutants along with relevant controls in Fig 4L and M, respectively, monitored under 4L20D.

Figure 5. CRY acts in the s-LNvs to inhibit their neural activity and promote wakefulness. (A) Brains of male flies expressing nls-GFP driven by *Gad1*GAL4 maintained under 4L20D and immunostained with PDF antisera. Representative l-LNv and s-LNv are displayed. (B) Quantification of GFP signal intensity of PDF neuron (n = 20, 49, 43 cells). Two-tailed Student's *t* test: ***$P < 0.001$. (C) Daily sleep duration of flies with *cry* knocked down in the s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D (n = 45, 60, 63, 34, 39 flies). (D-G) Brains from male WT and *cry* mutant flies dissected at ZT1, 7, 13, 19 under 4L20D (D) or 12L12D (F) are immunostained with GABA (cyan) and PDF (magenta) antisera, and representative s-LNvs are displayed. Merged signal is shown as yellow. Bar graphs represent normalized GABA intensity in the s-LNvs under 4L20D (E) (n = 20-53 cells) and 12L12D (G) (n = 29-67 cells). The average value of the control group at ZT1 is set to 1. (H) Representative live

image of the s-LNvs expressing GCaMP6m and tdTomato using *Pdf*GAL4. Brain samples are dissected at the indicated time points under 4L20D. (I) Quantification of GCaMP6m signal intensity normalized to that of tdTomato (n = 29-47 cells). Two-tailed Student's *t*-test: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. (J) Daily sleep duration of male flies expressing TrpA1 in the s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D and 29℃ to activate TrpA1 (n = 45, 54, 29, 22, 56, 22 flies). (K) Daily sleep duration of male *cry* mutant flies expressing TrpA1 in the s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D and 29℃ to activate TrpA1 (n = 33, 42, 23, 35, 36, 20 flies). (L) Daily sleep duration of male *cry* mutant flies over-expressing HK in the s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D (n = 24, 29, 23, 30, 37, 22 flies). For comparison with UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, *$P < 0.05$, ***$P < 0.001$; compared to UAS control, ##$P < 0.01$, ###$P < 0.001$. For comparison between mutant vs. control, two-tailed Student's *t*-test was used: compared to WT background, &&$P < 0.01$, &&&$P < 0.001$. The scale bar represents 15 μm. Error bars represent SEM. G4, GAL4; U, UAS; NS, not significant.

Figure 5—figure supplement 1. *Gad1*GAL4 drives expression in the s-LNvs but not l-LNvs. (A) Brains of male flies expressing nls-GFP driven by *Gad1*GAL4 maintained under 4L20D and immunostained with PDF antisera. (B) Brains of male flies expressing nls-GFP in GABAergic neurons using *Gad1*Gal4 and rCD2-RFP in the PDF neurons using *Pdf*LexA maintained under 4L20D. The scale bar represents 15 μm.

Figure 5—figure supplement 2. *cry* expression in the s-LNvs is necessary but not sufficient for maintaining normal sleep/wakefulness. (A) Sleep profile of *cry* knocked down in s-LNvs by R6GAL4 monitored under 4L20D in Fig 5C. White box indicates light period while black box indicates dark period. (B) Daily sleep duration of *cry* knocked down in PDF- GABAergic neurons using *Gad1*GAL4 and *Pdf*GAL80, monitored under 4L20D (n =101, 26, 29, 27, 31, 23, 32, 73, 63). One-way ANOVA with Bonferroni multiple comparison test: compared to GAL4 control, ***$P < 0.001$; compared to UAS control, #$P < 0.05$, ###$P < 0.001$. Error bars represent SEM. G4, GAL4; G80, GAL80; U, UAS.

Figure 5—figure supplement 3. s-LNvs appear to be GABAergic neurons. (A) Brains from male flies with *VGAT* knocked down in the s-LNvs using R6GAL4 and control maintained under 4L20D and immunostained with GABA (cyan) and PDF (magenta) antisera. Representative s-LNvs are displayed. Merged signal is shown as yellow. (B) Bar graphs represent normalized GABA intensity in the s-LNvs with *VGAT* knocked down using R6GAL4 (n = 25, 32 cells). The average value of the control group is set to 1. Two-tailed Student's *t*-test, ***$P < 0.001$. The scale bar represents 15 μm. Error bars represent SEM. G4, GAL4; U, UAS.

Figure 5—figure supplement 4. CRY promotes wakefulness by impinging on the excitability of the s-LNvs. (A) Sleep profile of male flies expressing TrpA1 in the s-LNvs using R6GAL4 and relevant controls in Fig 5J, monitored under 4L20D and 29°C to activate TrpA1. White box indicates light period while black box indicates dark period. (B) Sleep profile of male *cry* mutant flies expressing TrpA1 in the s-LNvs using R6GAL4 and relevant controls in Fig 5K, monitored under 4L20D and 29°C to activate TrpA1. (C) Sleep profile of male *cry* mutant flies over-expressing HK in the s-LNvs using R6GAL4 and relevant controls in Fig 5L, monitored under 4L20D. Error bars represent SEM. G4, GAL4; G80, GAL80; U, UAS.

Figure 6. s-LNvs release GABA onto the l-LNvs. (A) l-LNvs of male flies expressing syt-GFP in
the s-LNvs using R6GAL4 maintained under 4L20D and immunostained with PDF antisera. (B)
l-LNvs of male flies expressing trans-Tango in the s-LNvs using R6GAL4 maintained under
4L20D and immunostained with HA and PDF antisera. (C) Brains from male flies with *VGAT* (top)
or *Gad1* (bottom) knocked down in the s-LNvs using R6GAL4 and controls maintained under
4L20D and immunostained with GABA (cyan) and PDF (magenta) antisera. Representative
l-LNvs are displayed. Merged signal is shown as yellow. (D, E) Bar graphs represent normalized
GABA intensity in the l-LNvs of flies with *VGAT* (D) (n = 29, 37 cells) or *Gad1* (E) (n = 37, 51
cells) knocked down in the s-LNvs using R6GAL4, monitored under 4L20D. The average value of
the control group is set to 1. Two-tailed Student's *t*-test: $**P < 0.01$, $***P < 0.001$. (F) Daily sleep
duration of male *cry* mutant flies with *VGAT* knocked down in the s-LNvs using R6GAL4 and
relevant controls, monitored under 4L20D (n = 31, 31, 32, 29, 43, 19 flies). For comparison
between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple
comparison test was used: compared to GAL4 control, $*P < 0.05$, $**P < 0.01$; compared to UAS
control, $###P < 0.01$. For comparison between mutant vs. control, two-tailed Student's *t*-test was
used: compared to WT background, $&&&P < 0.001$. (G) Sleep profile of male *cry* mutant flies
with *VGAT* knocked down in the s-LNvs using R6GAL4 and relevant controls in F, monitored
under 4L20D. (H) Daily sleep duration of male *cry* mutant flies with *Gad1* knocked down in the
s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D (n = 59, 59, 53, 60, 62, 45
flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with
Bonferroni multiple comparison test was used: compared to GAL4 control, $*P < 0.05$, $**P < 0.01$,
$***P < 0.001$; compared to UAS control, $#P < 0.05$, $##P < 0.01$. For comparison between mutant
vs. control, two-tailed Student's *t*-test was used: compared to WT background, $&&&P < 0.001$. (I)
Sleep profile of male *cry* mutant flies with *Gad1* knocked down in the s-LNvs using R6GAL4 and
relevant controls in H, monitored under 4L20D.The scale bar represents 15 μm. Error bars
represent SEM. G4, GAL4; U, UAS; NS, not significant.

Figure 7. Short photoperiod reduces GABA level at the l-LNvs and increases sleep duration.
(A-D) Brains from male WT flies dissected at ZT1, 7, 13, 19 under 4L20D or 12L12D are
immunostained with GABA (cyan) and PDF (magenta) antisera. Representative l-LNvs (A) and
s-LNvs (C) are displayed. Merged signal is shown as yellow. Bar graphs represent normalized
GABA intensity in the l-LNvs (B) (n = 28-55 cells) and s-LNvs (D) (n = 21-59 cells). The average
value of the control group at ZT1 is set to 1. Student's *t*-test: $***P < 0.001$. (E-I) The daily sleep
duration (E), sleep bout number (F), average sleep bout length (G), waking activity (H) and sleep
profile (I) of male WT flies under 4L20D and 12L12D (n = 91, 32 flies). Two-tailed Student's
t-test: $***P < 0.001$. The scale bar represents 15 μm. Error bars represent SEM. (J) A model
demonstrating how CRY regulates the GABAergic s-LNv/l-LNv circuitry under short vs. longer
photoperiods to promote arousal.

Source Data

Figure 1—source data 1

Numerical data with associated statistical analyses underlying Figure 1.

Figure 2—source data 1

Numerical data with associated statistical analyses underlying Figure 2.

Figure 3—source data 1

Numerical data with associated statistical analyses underlying Figure 3.

Figure 4—source data 1

Numerical data with associated statistical analyses underlying Figure 4.

Figure 5—source data 1

Numerical data with associated statistical analyses underlying Figure 5.

Figure 6—source data 1

Numerical data with associated statistical analyses underlying Figure 6.

Figure 7—source data 1

Numerical data with associated statistical analyses underlying Figure 7.

Figure 1

Figure 1—figure supplement 1

Figure 2

Figure 2—figure supplement 1

Figure 2—figure supplement 2

Figure 2—figure supplement 3

Figure 2—figure supplement 4

Figure 3

Figure 4

Figure 4—figure supplement 1

Figure 4—figure supplement 2

Figure 5

Figure 5—figure supplement 1

A

*Gad1*G4-1/UnIs-GFP

GFP   PDF   GFP/PDF

l-LNvs

s-LNvs

B

LexAop-rCD2-RFP;*Gad1*G4-2/*Pdf*LexA;UnIs-GFP/+

GFP   RFP   GFP/RFP

l-LNvs

s-LNvs

Figure 5—figure supplement 2

Figure 5—figure supplement 3

A

GABA GABA/PDF

R6G4/+;U*dcr2*/+

U*VGATRNAi*-2/R6G4;
U*dcr2*/+

B

s-LNvs

Relative GABA intensity

1.5

1.0

0.5

0.0

***

R6G4/+;U*dcr2*/+

U*VGATRNAi*-2/R6G4;U*dcr2*/+

Figure 5—figure supplement 4

Figure 6

A R6G4/Usyt-GFP

l-LNvs · GFP · PDF · merge

B R6G4/U*trans*-Tango

l-LNvs · *trans*-Tango(HA) · PDF · merge

C

GABA · GABA/PDF
R6G4/+;U*dcr2*/+

GABA · GABA/PDF
U*VGATRNAi*-2/R6G4;U*dcr2*/+

l-LNvs

R6G4/+;U*dcr2*/+

R6G4/+;U*Gad1RNAi*/U*dcr2*

l-LNvs

D l-LNvs

Relative GABA intensity

R6G4/+;U*dcr2*/+
U*VGATRNAi*-2/R6G4;U*dcr2*/+ ***

E l-LNvs

Relative GABA intensity

R6G4/+;U*dcr2*/+
R6G4/+;U*gad1RNAi*/U*dcr2* **

F

Sleep duration (min)

R6G4/+
U*VGATRNAi*-2/+
R6G4/+;*cry^b* &&&
U*VGATRNAi*-2/+;*cry^b* &&&
U*VGATRNAi*-2/R6G4 *
U*VGATRNAi*-2/R6G4;*cry^b* ### **
NS

G

Sleep duration (min/0.5h)

— R6G4/+
— U*VGAT*RNAi-2/+
— R6G4/+;*cry^b*
— U*VGAT*RNAi-2/+;*cry^b*
— U*VGAT*RNAi-2/R6G4
— U*VGAT*RNAi-2/R6G4;*cry^b*

ZT0 · ZT24

H

Sleep duration (min)

R6G4/+
U*VGad1RNAi*/+
R6G4/+;*cry^b* &&
U*Gad1RNAi*/+;*cry^b* &&&
U*Gad1RNAi*/R6G4 ##
U*Gad1RNAi*/R6G4;*cry^b* # **
NS

I

Sleep duration (min/0.5h)

— R6G4/+
— U*Gad1*RNAi/+
— R6G4/+;*cry^b*
— U*Gad1*RNAi/+;*cry^b*
— U*Gad1*RNAi/R6G4
— U*Gad1*RNAi/R6G4;*cry^b*

ZT0 · ZT24

Figure 7