Streptococcus agalactiae and _Escherichia coli_ Induce Distinct Effector $\gamma\delta$ T Cell

Responses During Neonatal Sepsis and Neuroinflammation

Lila T. Witt[1,2], Kara G. Greenfield[1], Kathryn A. Knoop[1,3,*]

[1] Department of Immunology, Mayo Clinic, Rochester MN, USA 55901

[2] Mayo Graduate School of Biomedical Sciences, Mayo Clinic

[3] Department of Pediatrics, Mayo Clinic

Send correspondence to: Knoop.Kathryn@mayo.edu

Abstract

The neonatal phase of life is a time during which susceptibility to infection is particularly high, with

prematurely born neonates being especially vulnerable to life-threatening conditions such as

bacterial sepsis. While *Streptococcus agalactiae*, also known as group B *Streptococcus* (GBS)

and *Escherichia coli* are frequent causative pathogens of neonatal sepsis, it is still unclear how

the neonatal adaptive immune system responds to these pathogens. In the present study, we find

that γδ T cells in neonatal mice rapidly respond to single-organism sepsis infections of GBS and

E. coli, and that these infections induce distinct activation and effector functions from IFN-γ and

IL-17 producing γδ T cells, respectively. We also report differential reliance on γδTCR signaling

to elicit effector cytokine responses during neonatal sepsis, with IL-17 production during *E. coli*

sepsis being associated with TCR signaling, whereas IFN-γ production during GBS sepsis is

TCR-independent. Furthermore, we report that the divergent effector responses of γδ during GBS

and *E. coli* sepsis impart distinctive neuroinflammatory phenotypes on the neonatal brain. The

present study sheds light on how the neonatal adaptive immune response responds differentially

to bacterial stimuli and how these responses impact neonatal sepsis-associated

neuroinflammation.

Introduction

Late-onset neonatal sepsis remains a leading cause of neonatal morbidity and mortality

worldwide, particularly amongst preterm infants (Bergin et al., 2015; Stoll et al., 2011). The

neonatal period, defined as the first 28 days of life in humans, is a time during which risk of

infection is especially high, due in part to the relative immaturity of the neonatal immune system

(Bergin et al., 2015; Knoop et al., 2020; Segura-Cervantes et al., 2016). Premature neonates are

highly vulnerable to life-threatening conditions such as sepsis (Bizzarro et al., 2005; Dong and

Speer, 2015; Wynn et al., 2015). Gram-negative bacilli such as *Klebsiella*, *Pseudomonas* and *E.*

coli are prevalent in the gastrointestinal tract of premature neonates and are capable of

translocating from the gut and causing sepsis (Basu, 2015; Carl et al., 2014; Knoop et al., 2020).

Increased bacterial translocation from the neonatal gut is facilitated in part by selective

deficiencies in gut barrier defense mechanisms, including decreased production of protective

factors such as mucous, anti-microbial peptides, and IgA (Basu, 2015).

Conversely, Gram-positive neonatal sepsis is frequently caused by *Staphylococcus aureus* and

*Streptococcous agalactia*e, or group B *Streptococcus* (GBS). GBS colonizes the neonate via

vertical transmission during birth, often as the result of the neonate aspirating GBS-infected

amniotic fluid (Heath and Jardine, 2014; Stoll et al., 2011). Although rates of GBS sepsis are

declining due to improved early detection methods and prophylactic maternal antibiotic

administration, the mortality rate of GBS sepsis can be as high as 10%, with 30-50% of survivors

going on to experience neurological comorbidities in early childhood (Mynarek et al., 2021).

Although both GBS and *E. coli* can be found as components of a healthy microbiota, they have

the potential to cause severe disease in vulnerable populations, such as neonates (Remington et

al., 2010; Tavares et al., 2022). The increased risk of sepsis development amongst preterm

neonates is further compounded by deficiencies in several innate immunological defense

mechanisms, including decreased levels of circulating complement proteins, impaired neutrophil

function, and reduced secretion of pro-inflammatory cytokines by dendritic cells compared to

adults (Tsafaras et al., 2020). Similar to the innate immune system, the adaptive immune system

in neonates bears several striking differences to that of adults (Tsafaras et al., 2020). Compared

to adults, neonates have impaired memory T cell formation and are largely skewed toward Th2

over Th1 differentiation, thereby impairing their ability to mount a proper immune response to

microbial infections (Barrios et al., 1996; Basha et al., 2014; Li et al., 2004). Neonates also have

deficiencies in humoral immunity, such as delayed germinal center formation and impaired

antibody responses to both T cell dependent and independent antigens (Semmes et al., 2021)

Despite their relative impairments in the conventional T and B cell compartments, neonates have

a significant population of γδ T cells (Basha et al., 2014). γδ T cells are innate-like lymphocytes

that are abundant in barrier sites and act as early immune sentinels during infection (Chien et al.,

2014; Vantourout and Hayday, 2013). In contrast to conventional CD4+ and CD8+ T cells, γδ T

cells are exported from the thymus as functionally mature cells and are poised to rapidly deploy

their effector functions upon the detection of microbial ligands or pro-inflammatory cytokines

(Ribot et al., 2021). As the first T cells to develop in the embryonic thymus, γδ T cells are critical

players in the neonatal immune response during a time when CD4+ and CD8+ T cells, and B cells

are still developing and maturing (Dimova et al., 2015; Gibbons et al., 2009; Vantourout and

Hayday, 2013). Indeed, γδ T cells have been found to play a critical role in host protection during

neonatal influenza (Guo et al., 2018) and *Clostridium difficile* infection (Chen et al., 2020),

underscoring their importance during early life.

In the present study, we characterize the immune responses to two major neonatal sepsis

pathogens, *Streptococcus agalactiae* (Group B *Streptococcus*) and *Escherichia coli.* We report

that these two pathogens induce distinct effector cytokine responses from gamma delta (γδ) T

cells in postnatal day 7 (P7) pups and that these responses differentially impact mortality. We

also report that these two pathogens drive distinct neuroinflammatory phenotypes in neonatal

mice. This study sheds light on how distinct sepsis pathogens drive differential adaptive immune

responses in neonatal mice, impacting sepsis mortality and neuroinflammation.

Results

γδ T cells Respond to E. coli *and GBS Neonatal Sepsis and Differentially Drive Mortality*

While group B *Streptococcus* (GBS) and *Escherichia coli* are frequent causative pathogens of

neonatal sepsis, it is still unclear how features of these bacteria differentially drive the neonatal

adaptive immune response (Tsai et al., 2014). Therefore, we infected C57BL/6 P7 mice with a

single-organism infection of either 10^6 CFU *Streptococcus agalactiae* (GBS) or $2x10^4$ CFU *E. coli.*

Both *E. coli* and GBS septicemia induced robust activation from γδ T cells in the spleen 18 hours

post-infection, as measured by an increase the proportion of CD69+ CD62L- γδ T cells compared

to uninfected controls (Fig. 1a). A modest increase in the activation status of conventional CD4+

and CD8+ T cells, and B cells was also observed (Fig. S1a-c). Paralleling clinical findings, P7

mice infected with Gram-negative *E. coli* experienced decreased survival compared to Gram-

positive GBS (Tsai et al., 2014), despite the higher bacterial burden in GBS-infected pups at 18

hours post-infection (Fig.1b-d). Similarly, greater γδ T cell activation was also observed in *E. coli*

compared to GBS-infected pups (Fig. 1a). Interestingly, antibody-mediated depletion of γδ T

cells in *E. coli* infected mice was sufficient to completely rescue mortality (Fig. 1c), however no

differences in survival were observed in GBS- infected pups upon depletion of γδ T cells (Fig. 1b).

Of note, depleting γδ T cells did not impact either GBS or *E. coli* bacterial burden in neonatal pups

(Fig. 1d), suggesting that γδ T cells do not impact anti-microbial immune defenses during

neonatal sepsis. These data indicate that although γδ T cells undergo activation during both GBS

and *E. coli* neonatal sepsis infection, those responses differentially impact mortality independent

of systemic bacterial burden.

E. coli and GBS Neonatal Sepsis Drive Distinct Effector Cytokine Responses from γδ T cells

Due to the differential impact of γδ T cells on mortality during GBS vs. *E. coli* neonatal sepsis, we

next sought to further characterize the responses of γδ T cells during these two single-organism

sepsis infections. γδ T cells undergo developmental programming in the thymus and exist in the

periphery as either those that produce either IL-17 or IFN-γ (Haas et al., 2009; Muñoz-Ruiz et al.,

2017; Ribot et al., 2009). These distinct effector programs can also be discerned based on the

expression of certain surface markers, such as CCR6, restricted to IL-17 producing γδ T cells,

and CD27, restricted to IFN-γ producing γδ T cells (Ribot et al., 2009). We therefore asked if

different subsets of γδ T cells were being activated during GBS and *E. coli* infection in neonates.

Cytokine staining of γδ T cells revealed a robust increase in IL-17 expression during *E. coli*, but

not GBS infection whereas GBS induced increased IFN-γ, but not IL-17, expression from γδ T

cells (Fig. 2a-c). These findings were validated with serum ELISA, showing systemic increases in

these cytokines during GBS and *E. coli* sepsis infections (Fig. 2d,e). Accordingly, γδ T cell

activation during GBS infection was restricted to those expressing CD27, whereas during *E. coli*

infection, γδ T cell activation was restricted to the CCR6+ expressing fraction (Fig. 2f,g),

suggesting GBS and E. coli infections induce discrete activation of γδ T cell subsets. Therefore,

single-organism GBS and *E. coli* infections in neonates induces specific activation of different γδ

T cell populations, resulting in distinct effector cytokine profiles.

Neuroinflammation is a Feature of E. coli and GBS Neonatal Sepsis

Adverse neurologic outcomes are associated with inflammatory events in early life (Mynarek et

al., 2021; Stoll et al., 2004) however, how the adaptive immune response contributes to these

outcomes remains poorly understood. Both *E. coli* and GBS were present in the brains of P7 pups

18 hours post-infection, with increased CFUs of GBS present compared to *E. coli* (Fig. 3a). Flow

cytometry analysis of CD45^{hi} immune cells in the perfused brains of P7 pups revealed a significant

increase in monocytes and neutrophils in the brains of *E. coli* infected pups compared to

uninfected and GBS-infected pups (Fig. 3b,c). Interestingly, no significant increase in brain-

infiltrating monocytes or neutrophils was noted in GBS-infected pups compared to control mice,

despite the presence of GBS in the brain (Fig. 3b,c). To further investigate the neuroinflammatory

phenotype associated with GBS and *E. coli* sepsis infection in neonates, we measured mRNA of

immunological genes from bulk brain tissue on the Nanostring nCounter Gene Expression

platform. In the brains of both GBS and *E. coli* infected P7 pups, there was a significant increase

in the expression of monocyte and neutrophil chemotactic factors, such as *Ccl2* and *Cxcl1*,

respectively (Fig. 3d,e). Compared to uninfected pups, the brains of GBS-infected pups had

increased expression of genes involved in the innate immune response to Gram-positive bacteria,

such as *Tlr1* and *Tlr2* (Fig. 3d). *E. coli* brains had increased expression of genes associated with

TLR4 activation, such as *Cd14*, and the complement pathway, such as *C3* and *Marco* (Fig. 3e).

There were twenty-four differentially expressed genes between the brains of GBS and *E. coli*

infected mice including increased expression of *casp-3* during GBS infection, and increased *Il23*

expression during *E. coli* infection (Fig. 3e). These findings indicate that although both GBS and

E. coli can cause neuroinflammation during single-organism neonatal sepsis infection, they

induce distinct inflammatory phenotypes.

TCR-Specific Activation of γδ T cells Occurs During E. coli, but not GBS, Septicemia and Sepsis-

Associated Neuroinflammation

γδ T cells are capable of undergoing activation via multiple pathways, including MHC-independent

TCR activation by pathogen-derived non-peptide antigens (Constant et al., 1994). Nur77 is

transcription factor that is rapidly and specifically expressed during antigen-receptor mediated

signaling and activation in T and B cells (Ashouri and Weiss, 2017; Moran et al., 2011). Therefore,

to determine if γδTCR signaling was occurring in GBS and *E. coli* sepsis infection in mice, we

utilized P7 Nur77-GFP reporter pups. In the spleens of *E. coli* infected P7 pups, there was an

increase in the proportion of Nur77+ CD69+ γδ T cells (Fig. 4a), indicating TCR-mediated

activation of γδ T cells was occurring during *E. coli* neonatal sepsis infection. Importantly, nearly

all of these Nur77+ CD69+ cells in the spleen are CCR6+ (Fig. 4c), suggesting that the IL-17

signature during *E. coli* neonatal sepsis is associated with TCR signaling. Interestingly, there was

no change in the proportion of Nur77+ CD69+ γδ T cells in the spleens of GBS infected pups (Fig.

4a). These data suggest that during GBS infection, CD27+ γδ T cells undergo TCR-independent

activation. These data suggest that the CCR6+ γδ T cells responding to *E. coli* and CD27+ γδ T

cells responding to GBS neonatal sepsis undergo distinct pathways of activation to elicit their

effector cytokine responses.

γδ T cells play an important role in CNS homeostasis (Park et al., 2022; Ribeiro et al., 2019a) and

have been shown to have both a protective (Gentles et al., 2015) and detrimental (Gelderblom et

al., 2014; Welte et al., 2008) effect on the CNS under inflammatory conditions. We therefore

hypothesized that distinct γδ T cell responses in the periphery during GBS and *E. coli* sepsis

would impart a differential role for γδ T cells during sepsis-associated neuroinflammation. γδ T

cells within the CNS are highly skewed toward IL-17 production and CCR6 expression (Ribeiro et

al., 2019b; Wo et al., 2020). Similar to the spleen, we observed an increase in the proportion of

Nur77+ CD69+ γδ T cells in the brains of *E. coli*, but not GBS infected mice (Fig. 4b), with the

majority of Nur77+ CD69+ γδ T cells expressing CCR6 (Fig. 4c), suggesting that in the brain,

there is γδTCR specific signaling of IL-17 producing γδ T cells.

γδ T cell Responses During E. coli and GBS Septicemia Differentially Impact Sepsis-Associated

Neuroinflammation

In order to determine the contribution of γδ T cells to GBS and *E. coli* sepsis-associated

neuroinflammation, we compared immunological gene expression between BL/6 and TCRδ-/-

infected pups. TCRδ-/- were used in this instance due to the lack of depletion of brain-resident γδ

T cells during anti-TCRγδ antibody administration (Fig. S3a, b). Analysis of the brain by

Nanostring revealed sixty differentially expressed genes between BL/6 and TCRδ-/- *E. coli*

infected pups, and twenty-four differentially expressed genes between BL/6 and TCRδ-/- GBS

infected pups. Compared to BL/6 *E. coli* infected pups, TCRδ-/- pups had increased expression

of genes associated with apoptosis, such as *Casp-3,* and *Mapk1* (Fig. 4d). These findings

suggesting that during *E. coli* neonatal sepsis, γδ T cells may protect the neonatal brain from

inflammation-induced cell death. Compared to BL/6 GBS-infected mice, TCRδ-/- GBS infected

pups had decreased expression of genes involved in antigen presentation, such as *H2-Dmb2* and

genes involved in TGF-β signaling, such as *Tgfbr1* and *Smad5* (Fig. 4e). Genes involved in

apoptosis, such as *Casp-3*, were not significantly differentially expressed between TCRδ-/- and

BL/6 GBS infected pups, though it was increased in BL/6 GBS infected pups compared to BL/6

E. coli infected pups (Fig. 4e). Taken together, these data suggest that γδ T cells play a context-

specific role during sepsis-associated neuroinflammation.

Discussion

The present study reveals that γδ T cells undergo rapid activation and cytokine production during

a murine model of single-organism *Streptococcus agalactiae* (GBS) and *Escherichia coli* neonatal

sepsis infection. We also report that GBS and *E. coli* septicemia induce specific activation of IFN-

γ and IL-17 producing γδ T cells, respectively (Fig. 2). Although the γδ T cell compartment had

the highest proportion of cells undergoing activation in response to both *E. coli* and GBS sepsis

infections, a modest increase in activation status was observed for CD4+ and CD8+ T cells (Fig.

S1). This small proportion of activated CD4 and CD8 T cells may represent "virtual memory" (T$_{VM}$)

T cells, which are abundant in neonates (Akue et al., 2012; Schüler et al., 2004). Increased

activation status of CD4+ and CD8+ T cells at the early timepoint of 18 hours post-infection may

further implicate T$_{VM}$ cells, as they are capable of responding to infection faster naïve T cells due

to their memory-like capacity (Haluszczak et al., 2009; Lee et al., 2013).

In the context of *E. coli* infection, the presence of γδ T cells was found to be detrimental to the

survival of neonatal mice, whereas the presence of γδ T cells had no effect on survival of neonatal

pups during GBS infection (Fig. 1b, c). Furthermore, improved survival of pups in the absence of

γδ T cells during *E. coli* sepsis infection may be directly due to IL-17, the neutralization of which

has been shown to improve sepsis survival outcomes in other models of neonatal sepsis (Wynn

et al., 2016). This contrasts with other infections, such as neonatal murine influenza, in which

neonatal mice lacking γδ T cells rapidly succumbed to infection (Guo et al., 2018). The apparent

contradictory role of γδ T cells as pathogenic during sepsis vs. protective during other neonatal

infections could be due to the state of immune dysregulation and hyperinflammation that occurs
during sepsis.

Despite greater bacterial burden in the GBS sepsis infection, γδ T cells were more activated in
the context of *E. coli* sepsis infection (Fig. 1a,d), suggesting that specific bacterial factors, not just
bacterial burden, are more important in dictating the magnitude of a the neonatal γδ T cell
response. An additional outstanding question raised from this study is the extent to which γδ
cytokine responses depend directly on bacterial factors vs. the upstream immune response to
those bacterial factors, and whether γδ T cells are more likely to produce a certain cytokine in
response to specific virulence factors expressed by the bacteria. The clinical isolate of *E. coli*
used herein is an extraintestinal pathogenic *E. coli* (ExPEC) that expresses a number of virulence
genes, including the capsular polysaccharide K1 (Carl et al., 2014; Knoop et al., 2020). The K1
antigen has been implicated in various neonatal infections, including meningitis and sepsis
(Kaczmarek et al., 2014) and plays a critical role in the ability of *E. coli* to resist phagocytosis and
invade the central nervous system (Hoffman et al., 1999). Similarly, GBS COH-1 (ATCC) is a
highly virulent, encapsulated serotype III clinical isolate that expresses several virulence genes
involved in immune resistance and neuroinvasion. Included amongst these virulence factors is
the serine protease *CspA*, which has been implicated in the ability of GBS COH-1 to evade
opsonophagocytosis (Harris et al., 2003). GBS is also highly neuroinvasive and represents a
major causative pathogen of neonatal meningitis (Tavares et al., 2022). GBS COH-1 expresses
invasion associated gene (*IagA*), which has been implicated in the ability of GBS to infect human
brain microvascular endothelial cells (Doran et al., 2005). Furthermore, the extent to which
individual virulence factors expressed by *E. coli* and GBS induce specific γδ T cell activation
requires further investigation.

We also report that during *E. coli* and GBS sepsis infection, γδ T cells undergo distinct pathways
of activation in order to elicit their effector cytokine responses. While TCR-mediated activation

was associated with the production of IL-17 by γδ T cells during *E. coli* infection, we observed a

lack of Nur77 induction that during GBS infection, suggesting that γδ T cells likely use a TCR-

independent mechanism to undergo activation and produce IFN-γ (Fig. 4a). Similar to NK cells,

γδ T cells can produce IFN-γ in response to cytokines, such as IL-12, and IL-18 (Silva-Santos et

al., 2019), and in response to NKG2D ligands (Sedlak et al., 2014). Furthermore, the exact

mechanism by which CD27+ γδ T cells become activated and make IFN-γ during GBS infection

will be the subject of future studies.

In addition to the danger that sepsis poses to infants in the short-term, there are also several long-

term consequences that are common in survivors of neonatal sepsis. Poor neurological

outcomes, such as impaired neurologic growth, and the development of cerebral palsy, have been

associated with early life inflammatory events, including sepsis (Stoll et al., 2004). Although

mechanisms by which early life inflammation contributes to this phenomenon remain under

investigation, it is thought that inflammatory factors disrupt neuronal connectivity in the developing

brain (Cardoso et al., 2015). While both GBS and *E. coli* can cause severe neuroinflammation in

neonates (Kim, 2006; Tavares et al., 2022) how the neonatal adaptive immune response impacts

sepsis-associated neuroinflammation remains poorly understood. The present study therefore

sought to characterize the unique neuroinflammatory signatures associated with GBS and *E. coli*

systemic infection, and the extent to which these signatures are dependent upon γδ T cells. Both

GBS and *E. coli* were found in the brains of infected P7 pups, with slightly higher CFUs of GBS

compared to *E. coli*, similar to the periphery (Fig. 4a compared to 1d). *E. coli* infection also induced

greater infiltration of monocytes and neutrophils in the brain (Fig. 3b,c) despite both GBS and *E.

coli* showing increased gene expression of *Cxcl1* and *Cxcl10* (Fig. 3d,e). Therefore, this

recruitment may instead be due to the expression of bacterial factors, specific to *E. coli*, that act

as a direct chemotactic signal for these cells. Increased *casp-3* expression was observed in the

brains of *E. coli* infected TCRδ-/- pups compared to WT infected pups, suggesting that the γδ T

cell response in the brain during *E. coli* infection helps to prevent inflammation-induced cell death

in the developing brain (Fig. 4d). Interestingly, no significant increase in *casp-3* expression was

observed in GBS infected TCRδ-/- pups compared to WT infected pups (Fig. 3f). These data may

suggest that the specific γδ T cell responses during *E. coli* infection, possibly by IL-17 production,

promote cell survival in the developing brain. IL-17 has been shown to have an antiapoptotic effect

on tumor cells (Nam et al., 2008), and to help promote the survival of B cells and their

differentiation into plasma cells (Xu and Cao, 2010), but its specific role in this context remains

unknown. Furthermore, whether the induction of *Casp-3* and cell death in the brain is protective

or pathogenic requires further study.

During neonatal GBS infection, there was decreased expression of the *H2-Dmb2* gene in the

absence of γδ T cells. H2-Dmb2 is involved in the removal of CLIP from MHC class II molecules

and is critical for proper antigen presentation (Doebele et al., 2000; Santambrogio et al., 2019).

Although it is well-known that IFN-γ induces MHC II presentation (Kambayashi and Laufer, 2014;

Wijdeven et al., 2018), whether the decreased expression of molecules associated with MHC II

expression is dependent upon γδ T cell-derived IFN-γ during GBS infection also requires further

study. Similarly, *Il6ra*, *Tgfbr1*, and *Smad3* were also increased in the BL/6 compared to TCRδ-/-

GBS infected pups, suggesting that during GBS infection, γδ T cells may impact TGF-β signaling.

Overall, these findings present evidence of context-specific γδ T cell responses during neonatal

sepsis, with these responses differentially impacting survival and neuroinflammation. Taken

together, we report that γδ T cell responses during neonatal sepsis rely heavily on the initiating

pathogen. This work will help elucidate the contributions of neonatal γδ T cells to sepsis induced

mortality and neuroinflammation.

Methods

Bacteria

Clinical *E. coli* isolates were prepared as described previously (Knoop et al., 2020). GBS COH-1

was obtained from ATCC. Single bacterial colonies of GBS and *E. coli* were taken from a streak

plate and placed in a 15 mL conical (Fisher Healthcare) containing 5 mL of LB broth (Fisher

Healthcare) and placed into a 37°C incubator overnight. The following day, 10 mL of LB broth was

measured into a 50 mL conical tube (Fisher Healthcare) and a sterile dropper was used to place

2-3 drops of overnight bacterial stock into fresh LB broth. Bacteria was shaken at 150 rpm at 37°C

until an OD of 0.3 was reached. The bacterial culture was spun down at 12000 rpm for 10 minutes

(*E. coli*) or 30 minutes (GBS) and the LB supernatant was discarded. The bacterial pellet was

resuspended in 10mL of sterile PBS (Life Technologies) and a dose of $2x10^4$ CFU (*E. coli*) or 10^6

CFU (GBS) was intraperitoneally injected into the neonatal pups with an insulin syringe (Cardinal

Healthcare).

Mice

C57BL/6, TCRδ-/- and Nur77-GFP mice were purchased from Jackson Laboratory. All animals

were bred in accordance with the Institutional Animal Care and Use Committee at Mayo Clinic.

Anti-TCRγδ antibody (UC7-13D5, Biolegend) was given intraperitoneally for *in vivo* depletion of

γδ T cells (15 µg/g body weight).

Infection Model

Neonatal mice were infected on postnatal day 7 (P7). Spleens were harvested 18 hours post-

infection and blood was collected and allowed to clot for 45 minutes at room temperature and

spun down at 10,000xg for 2 minutes. Serum was collected and was stored at -20°C until use.

The liver was digested in 1mL of sterile PBS (Life Technologies) and 0.5 g of zirconium beads

(Fisher Healthcare) in a safe-lock snap cap tube (Fisher Healthcare) and placed into a tissue

homogenizer for 5 minutes. The liver homogenate was then serially diluted in PBS to achieve a

1:10^5 dilution (*E. coli*) and 1:10^6 dilution (GBS). Bacterial homogenate from *E. coli* and GBS

infected pups was plated on either MacConkey agar plates (Fisher Healthcare) or Tryptic Soy

Agar (DIFCO) respectively and CFUs were counted the following day.

ELISA

The following mouse ELISA kits were used: IL-17 DuoSet (Fisher Healthcare), IFN-γ ELISAmax

(Biolegend), ELISAs were performed according to manufacturer's instructions. Serum samples

were diluted 1:5 in ELISA diluent buffer (Biolegend).

Flow Cytometry

Spleens were harvested from mice and were placed in 1 mL of RPMI 1640 (VWR International

LLC), spleens were then mechanically homogenized using the frosted end of two glass slides

(Fisher Healthcare). Spleens were counted using a hemocytometer. Cells were then spun down

and resuspended in 500 µL FACS buffer (PBS containing 5% human serum, 0.5% BSA, 0.1%

sodium azide) and allowed to block for 20 minutes at 4°C. Surface master mix was made in FACS

buffer and staining was performed for 30 minutes in the dark at 4°C. Following this staining,

samples were washed twice with FACS buffer and samples were acquired on the CyTek Northern

Lights Spectral Flow Cytometer (Cytek Biosciences).

Intracellular cytokine staining

Cells were placed in 250 µL RPMI supplemented with 10% FBS, 2mM Glutamine (Gibco), 2mM

Pyruvate (BioWhittaker), 50 µg/mL Pen/Strep (BioWhittaker), and 0.55 mM 2-ME (Gibco). 1X

protein transport inhibitor (Fisher Healthcare) and 1:1000 PMA/Ionomycin (Biolegend) were

added, and samples were placed in an incubator at 37°C for four hours. Samples were spun down

and resuspended in FACS buffer to block for 15 minutes. Samples were then stained for surface

markers for 30 minutes at 4°C before 100 µL of fixative (Life Technologies) was added to each

sample for 30 minutes at room temperature. Samples were then washed once with FACS buffer

and once with 1X perm buffer (Biolegend) and spun down at 1500 rpm for 5 minutes. Samples

were then resuspended in 100 µL 1X perm buffer and intracellular antibodies and were placed at

4°C overnight. The following day, samples were washed twice with FACS buffer and acquired on

the CyTek Northern Lights Spectral Flow Cytometer.

The following antibodies were used for flow cytometry analysis:

Target	Fluorophore	Clone	Vendor	Titration
CD45	PE/Cy7, BV605	30-F11	Biolegend	1:500
CD4	BV421	GK1.5	Biolegend	1:500
CD8	BV510	53-6.7	Biolegend	1:250
CD11b	BV510	M1/70	Biolegend	1:250
TCRgd	BV421, PE	GL3	Biolegend	1:250
Viability	Zombie NIR, Violet	N/A	Biolegend	1:1000
Ly6G	APC	1A8	Biolegend	1:250
CD69	FITC, BV650	H1.2F3	Biolegend	1:250
CD62L	PE/Cy5	MEL-14	Biolegend	1:250
CCR6	BV785, APC	29-2L17	Biolegend	1:250
CD27	BV421	LG.3A10	Biolegend	1:250
Ly6C	PerCP	HK1.4	Biolegend	1:250
IL-17A	APC, PE	TC-11-18H10.1	Biolegend	1:100
IFN-g	PE, FITC	XMG1.2	BD Biosciences	1:100
TCRβ	BV711	H57-597	Biolegend	1:250
CD19	APC/Cy7	6D5	Biolegend	1:250
CD3	APC/Cy7	17A2	Biolegend	1:250

Brain Isolation for Analysis by Flow Cytometry

Mice were anesthetized with 10 ug/g of Ketamine/Xyalizine mixture and transcardially perfused

with 10 mL of cold, sterile PBS (Life Technologies). Brains were digested as previously described

(Cumba Garcia et al., 2016). In brief, brains were isolated and placed into a 50 mL conical tube

(Fisher Healthcare) containing 5 mL of RPMI (VWR International LLC). The RPMI containing the

brains of the mice were then transferred to a 7 mL glass Tenbroeck dounce homogenizer (Pyrex)

and homogenized until the brain was visibly digested (about 10 plunges). The homogenate was

then poured through a 70 μm filter into a new 50 mL conical tube, and 10 more mL of RPMI 1640

was added, along with 1 mL of 10X PBS and 9 mL of Percoll (Sigma-Aldrich INC). The 50 mL

conical tubes were then placed into a centrifuge and pelleted at 7840xg for 30 minutes at 4°C.

Following the spin, the supernatant was fully aspirated off and samples were washed with 50 mL

of fresh RPMI 1640 and spun again at 1500 rpm for 10 minutes. Samples were then blocked in

FACS buffer for 15 minutes before surface staining was performed at 4°C for 30 minutes.

mRNA Isolation from Brains Brains were homogenized in 1 mL of sterile PBS (Life Technologies)

using a dounce homogenizer (Pyrex). 100 μL of brain homogenate was used for mRNA isolation

with the Quiagen RNeasy Mini Kit according to manufacturer's instructions. RNA samples were

stored at -80°C until ready for use.

Nanostring

Nanostring nCounter Mouse Immunology Max Kit was used following mRNA isolation from the

brain. RNA hybridization was performed according to Nanostring manufacturer's instructions.

Samples were incubated for 24 hours at 65°C and then were run on the nCounter Prep Station

5s before being placed on the nCounter Digital Analyzer. The ROSALIND platform was used for

data analysis.

Statistical Analysis

Student's unpaired t-test, One-way ANOVA, and Kaplan-Meier tests were conducted using

GraphPad Prism (GraphPad Software, Inc.,)

Author Contributions: LTW and KAK conceived the studies and wrote the manuscript. LTW

performed animal experiments, ELISAs, flow cytometry, RNA analysis, and data analysis. KGG

assisted with animal husbandry and experiments. All authors reviewed the data and manuscript.

Work was funded by T32 AI170478 (LTW), and R01 DK134366 (KAK).

Akue, A.D., Lee, J.Y., and Jameson, S.C. (2012). Derivation and maintenance of virtual memory CD8 T cells.
J Immunol *188*, 2516-2523.
Ashouri, J.F., and Weiss, A. (2017). Endogenous Nur77 Is a Specific Indicator of Antigen Receptor Signaling
in Human T and B Cells. J Immunol *198*, 657-668.
Barrios, C., Brawand, P., Berney, M., Brandt, C., Lambert, P.H., and Siegrist, C.A. (1996). Neonatal and early
life immune responses to various forms of vaccine antigens qualitatively differ from adult responses:
predominance of a Th2-biased pattern which persists after adult boosting. Eur J Immunol *26*, 1489-1496.
Basha, S., Surendran, N., and Pichichero, M. (2014). Immune responses in neonates. Expert Rev Clin
Immunol *10*, 1171-1184.
Basu, S. (2015). Neonatal sepsis: the gut connection. European Journal of Clinical Microbiology &
Infectious Diseases *34*, 215-222.
Bergin, S.P., Thaden, J.T., Ericson, J.E., Cross, H., Messina, J., Clark, R.H., Fowler, V.G., Jr., Benjamin, D.K.,
Jr., Hornik, C.P., and Smith, P.B. (2015). Neonatal Escherichia coli Bloodstream Infections: Clinical
Outcomes and Impact of Initial Antibiotic Therapy. Pediatr Infect Dis J *34*, 933-936.
Bizzarro, M.J., Raskind, C., Baltimore, R.S., and Gallagher, P.G. (2005). Seventy-five years of neonatal sepsis
at Yale: 1928-2003. Pediatrics *116*, 595-602.
Cardoso, F.L., Herz, J., Fernandes, A., Rocha, J., Sepodes, B., Brito, M.A., McGavern, D.B., and Brites, D.
(2015). Systemic inflammation in early neonatal mice induces transient and lasting neurodegenerative
effects. J Neuroinflammation *12*, 82.
Carl, M.A., Ndao, I.M., Springman, A.C., Manning, S.D., Johnson, J.R., Johnston, B.D., Burnham, C.A.,
Weinstock, E.S., Weinstock, G.M., Wylie, T.N.*, et al.* (2014). Sepsis from the gut: the enteric habitat of
bacteria that cause late-onset neonatal bloodstream infections. Clin Infect Dis *58*, 1211-1218.
Chen, Y.S., Chen, I.B., Pham, G., Shao, T.Y., Bangar, H., Way, S.S., and Haslam, D.B. (2020). IL-17-producing
γδ T cells protect against Clostridium difficile infection. J Clin Invest *130*, 2377-2390.
Chien, Y.-h., Meyer, C., and Bonneville, M. (2014). γδ T Cells: First Line of Defense and Beyond. Annual
Review of Immunology *32*, 121-155.
Constant, P., Davodeau, F., Peyrat, M.-A., Poquet, Y., Puzo, G., Bonneville, M., and Fournié, J.-J. (1994).
Stimulation of Human γδ T Cells by Nonpeptidic Mycobacterial Ligands. Science *264*, 267-270.
Cumba Garcia, L.M., Huseby Kelcher, A.M., Malo, C.S., and Johnson, A.J. (2016). Superior isolation of
antigen-specific brain infiltrating T cells using manual homogenization technique. J Immunol Methods
439, 23-28.
Dimova, T., Brouwer, M., Gosselin, F., Tassignon, J., Leo, O., Donner, C., Marchant, A., and Vermijlen, D.
(2015). Effector Vγ9Vδ2 T cells dominate the human fetal γδ T-cell repertoire. Proc Natl Acad Sci U S A
112, E556-565.
Doebele, R.C., Busch, R., Scott, H.M., Pashine, A., and Mellins, E.D. (2000). Determination of the HLA-DM
interaction site on HLA-DR molecules. Immunity *13*, 517-527.
Dong, Y., and Speer, C.P. (2015). Late-onset neonatal sepsis: recent developments. Archives of disease in
childhood Fetal and neonatal edition *100*, F257-F263.
Doran, K.S., Engelson, E.J., Khosravi, A., Maisey, H.C., Fedtke, I., Equils, O., Michelsen, K.S., Arditi, M.,
Peschel, A., and Nizet, V. (2005). Blood-brain barrier invasion by group B Streptococcus depends upon
proper cell-surface anchoring of lipoteichoic acid. The Journal of Clinical Investigation *115*, 2499-2507.
Gelderblom, M., Arunachalam, P., and Magnus, T. (2014). γδ T cells as early sensors of tissue damage and
mediators of secondary neurodegeneration. Front Cell Neurosci *8*, 368.
Gentles, A.J., Newman, A.M., Liu, C.L., Bratman, S.V., Feng, W., Kim, D., Nair, V.S., Xu, Y., Khuong, A.,
Hoang, C.D.*, et al.* (2015). The prognostic landscape of genes and infiltrating immune cells across human
cancers. Nat Med *21*, 938-945.

Gibbons, D.L., Haque, S.F., Silberzahn, T., Hamilton, K., Langford, C., Ellis, P., Carr, R., and Hayday, A.C. (2009). Neonates harbour highly active gammadelta T cells with selective impairments in preterm infants. Eur J Immunol *39*, 1794-1806.

Guo, X.J., Dash, P., Crawford, J.C., Allen, E.K., Zamora, A.E., Boyd, D.F., Duan, S., Bajracharya, R., Awad, W.A., Apiwattanakul, N*., et al.* (2018). Lung γδ T Cells Mediate Protective Responses during Neonatal Influenza Infection that Are Associated with Type 2 Immunity. Immunity *49*, 531-544.e536.

Haas, J.D., González, F.H.M., Schmitz, S., Chennupati, V., Föhse, L., Kremmer, E., Förster, R., and Prinz, I. (2009). CCR6 and NK1.1 distinguish between IL-17A and IFN-γ-producing γδ effector T cells. European Journal of Immunology *39*, 3488-3497.

Haluszczak, C., Akue, A.D., Hamilton, S.E., Johnson, L.D., Pujanauski, L., Teodorovic, L., Jameson, S.C., and Kedl, R.M. (2009). The antigen-specific CD8+ T cell repertoire in unimmunized mice includes memory phenotype cells bearing markers of homeostatic expansion. J Exp Med *206*, 435-448.

Harris, T.O., Shelver, D.W., Bohnsack, J.F., and Rubens, C.E. (2003). A novel streptococcal surface protease promotes virulence, resistance to opsonophagocytosis, and cleavage of human fibrinogen. The Journal of Clinical Investigation *111*, 61-70.

Heath, P.T., and Jardine, L.A. (2014). Neonatal infections: group B streptococcus. BMJ Clin Evid *2014*.

Hoffman, J.A., Wass, C., Stins, M.F., and Kim, K.S. (1999). The capsule supports survival but not traversal of Escherichia coli K1 across the blood-brain barrier. Infect Immun *67*, 3566-3570.

Kaczmarek, A., Budzyńska, A., and Gospodarek, E. (2014). Detection of K1 antigen of Escherichia coli rods isolated from pregnant women and neonates. Folia Microbiol (Praha) *59*, 419-422.

Kambayashi, T., and Laufer, T.M. (2014). Atypical MHC class II-expressing antigen-presenting cells: can anything replace a dendritic cell? Nat Rev Immunol *14*, 719-730.

Kim, K.S. (2006). Meningitis-Associated <i>Escherichia coli</i>. EcoSal Plus *2*, 10.1128/ecosalplus.1128.1126.1121.1122.

Knoop, K.A., Coughlin, P.E., Floyd, A.N., Ndao, I.M., Hall-Moore, C., Shaikh, N., Gasparrini, A.J., Rusconi, B., Escobedo, M., Good, M*., et al.* (2020). Maternal activation of the EGFR prevents translocation of gut-residing pathogenic Escherichia coli in a model of late-onset neonatal sepsis. Proceedings of the National Academy of Sciences of the United States of America *117*, 7941-7949.

Lee, J.Y., Hamilton, S.E., Akue, A.D., Hogquist, K.A., and Jameson, S.C. (2013). Virtual memory CD8 T cells display unique functional properties. Proc Natl Acad Sci U S A *110*, 13498-13503.

Li, L., Lee, H.H., Bell, J.J., Gregg, R.K., Ellis, J.S., Gessner, A., and Zaghouani, H. (2004). IL-4 utilizes an alternative receptor to drive apoptosis of Th1 cells and skews neonatal immunity toward Th2. Immunity *20*, 429-440.

Moran, A.E., Holzapfel, K.L., Xing, Y., Cunningham, N.R., Maltzman, J.S., Punt, J., and Hogquist, K.A. (2011). T cell receptor signal strength in Treg and iNKT cell development demonstrated by a novel fluorescent reporter mouse. J Exp Med *208*, 1279-1289.

Muñoz-Ruiz, M., Sumaria, N., Pennington, D.J., and Silva-Santos, B. (2017). Thymic Determinants of γδ T Cell Differentiation. Trends Immunol *38*, 336-344.

Mynarek, M., Bjellmo, S., Lydersen, S., Afset, J.E., Andersen, G.L., and Vik, T. (2021). Incidence of invasive Group B Streptococcal infection and the risk of infant death and cerebral palsy: a Norwegian Cohort Study. Pediatr Res *89*, 1541-1548.

Nam, J.S., Terabe, M., Kang, M.J., Chae, H., Voong, N., Yang, Y.A., Laurence, A., Michalowska, A., Mamura, M., Lonning, S*., et al.* (2008). Transforming growth factor beta subverts the immune system into directly promoting tumor growth through interleukin-17. Cancer Res *68*, 3915-3923.

Park, J.H., Kang, I., and Lee, H.K. (2022). γδ T Cells in Brain Homeostasis and Diseases. Front Immunol *13*, 886397.

Remington, J.S., Wilson, C.B., Nizet, V., Klein, J.O., and Maldonado, Y. (2010). Infectious diseases of the fetus and newborn E-book (Elsevier Health Sciences).

Ribeiro, M., Brigas, H.C., Temido-Ferreira, M., Pousinha, P.A., Regen, T., Santa, C., Coelho, J.E., Marques-
Morgado, I., Valente, C.A., Omenetti, S*., et al.* (2019a). Meningeal γδ T cell-derived IL-17 controls synaptic
plasticity and short-term memory. Sci Immunol *4*.
Ribeiro, M., Brigas, H.C., Temido-Ferreira, M., Pousinha, P.A., Regen, T., Santa, C., Coelho, J.E., Marques-
Morgado, I., Valente, C.A., Omenetti, S*., et al.* (2019b). Meningeal γδ T cell–derived IL-17 controls synaptic
plasticity and short-term memory. Science Immunology *4*, eaay5199.
Ribot, J.C., deBarros, A., Pang, D.J., Neves, J.F., Peperzak, V., Roberts, S.J., Girardi, M., Borst, J., Hayday,
A.C., Pennington, D.J*., et al.* (2009). CD27 is a thymic determinant of the balance between interferon-
gamma- and interleukin 17-producing gammadelta T cell subsets. Nat Immunol *10*, 427-436.
Ribot, J.C., Lopes, N., and Silva-Santos, B. (2021). γδ T cells in tissue physiology and surveillance. Nature
Reviews Immunology *21*, 221-232.
Santambrogio, L., Berendam, S.J., and Engelhard, V.H. (2019). The Antigen Processing and Presentation
Machinery in Lymphatic Endothelial Cells. Front Immunol *10*, 1033.
Schüler, T., Hämmerling, G.n.J., and Arnold, B. (2004). Cutting Edge: IL-7-Dependent Homeostatic
Proliferation of CD8+ T Cells in Neonatal Mice Allows the Generation of Long-Lived Natural Memory T Cells
1. The Journal of Immunology *172*, 15-19.
Sedlak, C., Patzl, M., Saalmüller, A., and Gerner, W. (2014). IL-12 and IL-18 induce interferon-γ production
and de novo CD2 expression in porcine γδ T cells. Dev Comp Immunol *47*, 115-122.
Segura-Cervantes, E., Mancilla-Ramírez, J., González-Canudas, J., Alba, E., Santillán-Ballesteros, R.,
Morales-Barquet, D., Sandoval-Plata, G., and Galindo-Sevilla, N. (2016). Inflammatory Response in
Preterm and Very Preterm Newborns with Sepsis. Mediators Inflamm *2016*, 6740827.
Semmes, E.C., Chen, J.-L., Goswami, R., Burt, T.D., Permar, S.R., and Fouda, G.G. (2021). Understanding
Early-Life Adaptive Immunity to Guide Interventions for Pediatric Health. Frontiers in Immunology *11*.
Silva-Santos, B., Mensurado, S., and Coffelt, S.B. (2019). γδ T cells: pleiotropic immune effectors with
therapeutic potential in cancer. Nat Rev Cancer *19*, 392-404.
Stoll, B.J., Hansen, N.I., Adams-Chapman, I., Fanaroff, A.A., Hintz, S.R., Vohr, B., Higgins, R.D., National
Institute of Child, H., and Human Development Neonatal Research Network, f.t. (2004).
Neurodevelopmental and Growth Impairment Among Extremely Low-Birth-Weight Infants With Neonatal
Infection. JAMA *292*, 2357-2365.
Stoll, B.J., Hansen, N.I., Sánchez, P.J., Faix, R.G., Poindexter, B.B., Van Meurs, K.P., Bizzarro, M.J., Goldberg,
R.N., Frantz, I.D., 3rd, Hale, E.C*., et al.* (2011). Early onset neonatal sepsis: the burden of group B
Streptococcal and E. coli disease continues. Pediatrics *127*, 817-826.
Tavares, T., Pinho, L., and Bonifácio Andrade, E. (2022). Group B Streptococcal Neonatal Meningitis. Clin
Microbiol Rev *35*, e0007921.
Tsafaras, G.P., Ntontsi, P., and Xanthou, G. (2020). Advantages and Limitations of the Neonatal Immune
System. Front Pediatr *8*.
Tsai, M.H., Hsu, J.F., Chu, S.M., Lien, R., Huang, H.R., Chiang, M.C., Fu, R.H., Lee, C.W., and Huang, Y.C.
(2014). Incidence, clinical characteristics and risk factors for adverse outcome in neonates with late-onset
sepsis. Pediatr Infect Dis J *33*, e7-e13.
Vantourout, P., and Hayday, A. (2013). Six-of-the-best: unique contributions of γδ T cells to immunology.
Nature Reviews Immunology *13*, 88-100.
Welte, T., Lamb, J., Anderson, J.F., Born, W.K., O'Brien, R.L., and Wang, T. (2008). Role of two distinct
gammadelta T cell subsets during West Nile virus infection. FEMS Immunol Med Microbiol *53*, 275-283.
Wijdeven, R.H., van Luijn, M.M., Wierenga-Wolf, A.F., Akkermans, J.J., van den Elsen, P.J., Hintzen, R.Q.,
and Neefjes, J. (2018). Chemical and genetic control of IFNγ-induced MHCII expression. EMBO Rep *19*.
Wo, J., Zhang, F., Li, Z., Sun, C., Zhang, W., and Sun, G. (2020). The Role of Gamma-Delta T Cells in Diseases
of the Central Nervous System. Frontiers in Immunology *11*.

Wynn, J.L., Guthrie, S.O., Wong, H.R., Lahni, P., Ungaro, R., Lopez, M.C., Baker, H.V., and Moldawer, L.L.
(2015). Postnatal Age Is a Critical Determinant of the Neonatal Host Response to Sepsis. Mol Med *21*, 496-
504.
Wynn, J.L., Wilson, C.S., Hawiger, J., Scumpia, P.O., Marshall, A.F., Liu, J.H., Zharkikh, I., Wong, H.R., Lahni,
P., Benjamin, J.T.*, et al.* (2016). Targeting IL-17A attenuates neonatal sepsis mortality induced by IL-18.
Proc Natl Acad Sci U S A *113*, E2627-2635.
Xu, S., and Cao, X. (2010). Interleukin-17 and its expanding biological functions. Cell Mol Immunol *7*, 164-
174.

Figure 1: γδ T cells Respond to *E. coli* and GBS Neonatal Sepsis and Differentially Drive Mortality
A) Gating scheme (L) and quantification (R) of CD69+, CD62L- splenic γδ+ T cells in postnatal day 7 (P7) 18 hours post-infection with either 2×10^4 CFU *E. coli* or 10^6 CFU GBS COH-1 B) Survival curves of BL/6 P7 pups infected with *E. coli* or C) GBS treated with either isotype IgG or 15 μg/g anti-TCRγδ antibody D) Bacterial CFUs from the livers of GBS and *E. coli* infected pups treated with anti-TCRγδ antibody. Data shown is from four independent experiments, n>3 mice per group. Statistical tests used include one-way ANOVA (A, D), and Kaplan-Meier (B, C). with ns p>0.5

Figure 2: *E. coli* and GBS Neonatal Sepsis Drive Distinct Effector Cytokine Responses from γδ T cells
A) Flow cytometry gating scheme of IFN-y+ and IL-17+ splenic γδ+ T cells from uninfected, GBS, and *E. coli* infected BL/6 P7 pups. B) Mean fluorescent intensity (MFI) of IFN-y and C) IL-17 from splenic γδ+ T cells 18 hours post-infection. D) IFN-y and E) IL-17 serum ELISA. F) Proportion of activated CD27+ and G) CCR6+ γδ+ T cells from uninfected, GBS, and *E. coli* infected BL/6 P7 pups. Data shown is from three independent experiments, n>3 mice per group, statistical tests used include one-way ANOVA with ns p>0.5.

Figure 3: Neuroinflammation is a Feature of *E. coli* and GBS Neonatal Sepsis
A) GBS and *E. coli* CFUs from the brains of BL/6 P7 pups B) Absolute number of monocytes (CD45hi, CD11b+, Ly6C+, Ly6G-) and C) Neutrophils (CD45hi, CD11b+, Ly6C-, Ly6G+) in the perfused brains of *E. coli* and GBS infected BL/6 P7 pups 18 hours post-infection. D) Volcano plot of differentially expressed genes between BL/6 uninfected control vs. *E. coli* infected P7 pups and E) GBS infected P7 pups. Data shown is from four independent experiments, n>4 mice per group. Statistical tests used include Student's unpaired t-test (A), one-way ANOVA (B,C) with ns p>0.5.

Figure 4: TCR-Specific Activation of γδ+ T cells Occurs During E. coli, but not GBS, Septicemia and Differentially Drives Neuroinflammation
Proportion of Nur77+ CD69+ γδ+ T cells in the A) Spleen and B) Brain of GBS and *E. coli* infected Nur77-GFP P7 pups. C) CCR6 expression of Nur77+ CD69+ γδ+ T cells in the spleen and brain of *E. coli*-infected pups. D) Volcano plots of differentially expressed genes in the brains of *E. coli* infected, or E) GBS-infected BL/6 vs. TCRδ-/- P7 pups. Data shown is from three independent experiments, n>3 mice per group. Statistics used include one-way ANOVA (A,B) with ns p>0.5.

Fig. 1: γδ T cells Respond to *E. coli* and GBS Neonatal Sepsis and Differentially Contribute to Mortality

Fig. 2: *E. coli* and GBS Neonatal Sepsis Drive Distinct Effector Cytokine Responses from γδ T cells

Fig. 3: Neuroinflammation is a Feature of *E. coli* and GBS Neonatal Sepsis

Fig. 4: TCR-Specific Activation of γδ T cells Occurs During *E. coli*, but not GBS, Septicemia and Differentially Drives Neuroinflammation

Figure S1: Responses of CD4+ and CD8+ T Cells, and B cells to GBS and *E. coli* Neonatal Sepsis

A) Flow cytometric staining of CD69+, CD62L- splenic CD4, B) CD8 T cells and C) B cells 18 hours-post *E. coli* or GBS infection in P7 BL/6 pups. Data shown is from two independent experiments, n>3 mice per group, statistical tests used include one-way ANOVA, ns p>0.5.

Figure S2: Flow Cytometry Gating Scheme of Neonatal Brain Immune Cell Populations

A) Live, single cells in the brain were gated on CD45, with CD45mid population representing microglia (CD11b+, CX3CR1+). The CD45hi compartment was gated into TCRγδ+ or CD11b+. CD11b+ cells were then gated into Ly6C+, Ly6G- (Monocytes), or Ly6C-, Ly6G+ (Neutrophils).

Figure S3: Lack of Depletion of Brain-Resident T Cells Upon Systemic anti-TCRγδ Antibody Administration

A) Absolute number of γδ+ T cells in the spleen and B) brain of BL/6 P7 pups intraperitoneally injected with 15 µg/g anti-TCRγδ antibody (clone: UC7-13D5). Data shown is from two independent experiments, n>4 mice per group. Statistical tests used include Student's unpaired t-test (A,B), with ns p>0.5.