SARS-CoV-2 neurotropism-induced anxiety/depression-like behaviors require Microglia activation

Qian Ge [1#], Shan Zhou [2#], Jose Porras [2], Panfeng Fu[4], Ting Wang[4], Jianyang Du[1,3] *,
Kun Li[2] *

Author affiliations:
[1] Department of Anatomy and Neurobiology, University of Tennessee Health Science Center, Memphis, TN 38163, USA.
[2] Florida Research and Innovation Center, Cleveland Clinic, Port St. Lucie, FL 34987, USA
[3] Neuroscience Institute, University of Tennessee Health Science Center, Memphis, TN, United States.
[4] Center for Translational Science, Florida International University, Port St. Lucie, FL, 34987, USA

* Correspondence to:

Kun Li
Florida Research and Innovation Center,
Cleveland Clinic,
Port St. Lucie, FL 34987, USA
Tel:772-419-2239
E-mail:lik11@ccf.org

Jianyang Du
Department of Anatomy and Neurobiology,
University of Tennessee Health Science Center,
Memphis, TN 38163, USA
Tel: 901-448-3463
E-mail: jdu15@uthsc.edu

Authorship note: [#] Q.G. and S.Z. contributed equally to this work.

KEYWORDS COVID-19, SARS-CoV-2, long COVID, post-acute sequelae of SARS-CoV-2 infection, spike protein, Microglia, Amygdala, neurotropism, anxiety- and depression-like behaviors, synaptic transmission and plasticity.

ABSTRACT -196 words

The coronavirus disease 2019 (COVID-19) pandemic, caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has been associated with a wide range of "long COVID" neurological symptoms. However, the mechanisms governing SARS-CoV-2 neurotropism and its effects on long-term behavioral changes remain poorly understood. Using a highly virulent mouse-adapted SARS-CoV-2 strain, denoted as SARS2-N501Y$_{MA30}$, we demonstrated that intranasal inoculation of SARS2-N501Y$_{MA30}$ results in viral dissemination to multiple brain regions, including the

amygdala and hippocampus. Behavioral assays indicated a marked elevation in

anxiety- and depression-like behaviors post infection. A comparative analysis of RNA

expression profiles disclosed alterations in the post-infected brains. Additionally, we

observed dendritic spine remodeling on neurons within the amygdala after infection.

Infection with SARS2-N501Y$_{MA30}$ was associated with microglial activation and a

subsequent increase in microglia-dependent neuronal activity in the amygdala.

Pharmacological inhibition of microglial activity subsequent to viral spike inoculation

mitigates microglia-dependent neuronal hyperactivity. Transcriptomic analysis of

infected brains revealed the upregulation of inflammatory and cytokine-related

pathways, implicating microglia-driven neuroinflammation in the pathogenesis of

neuronal hyperactivity and behavioral abnormality. Overall, these data provide critical

insights into the neurological consequences of SARS-CoV-2 infection and underscore

microglia as a potential therapeutic target for ameliorating virus-induced

neurobehavioral abnormalities.

INTRODUCTION:

As of April, 3rd, 2024, the global pandemic of COVID-19 has resulted in 774,954,393

reported cases and 7,040264 confirmed deaths (WHO COVID-19 Dashboard

https://data.who.int/dashboards/covid19/deaths?n=c). The death rate and hospital

admissions related to COVID-19 have been dramatically reduced as a result of

extensive vaccination rollouts and improved treatments. However, a substantial number

of patients (10-20%) are experiencing a persistent or newly developed set of symptoms

following the acute phase of the illness. This condition is commonly referred to as post-

acute sequelae of SARS-CoV-2 infection (1) also known as "long COVID," (1). While

PASC initially attracted attention for its severe impact on older adults and those with

underlying health concerns, it has since been clear that it may also occur in otherwise

healthy young people, and it can develop after even a modest initial infection (2, 3).

These long-lasting symptoms may persist for weeks or even months, posing challenges

for healthcare providers and necessitating further research and support to address the

long-term health consequences of the disease.

Among the enduring symptoms of PASC, neurological manifestations stand out

prominently. Symptoms include cognitive difficulties, autonomic dysfunction, extreme

fatigue, sleep disturbances, and mental health complications such as anxiety and

depression (2, 3). The exact cause of these syndromes remains uncertain and is

constantly being studied. Possible factors contributing to these symptoms include viral

infection of brain cells, immune-mediated phenomena, coincidental events, or a

combination of these factors. In vitro studies have provided clear evidence of SARS-

CoV-2 infection in human brain organoids or cell cultures (4, 5) Additionally, several

autopsy studies have reported the presence of viral RNA or proteins in the brains of

patients who died from COVID-19, suggesting the possibility of SARS-CoV-2

neurotropism in the central nervous system (CNS) (5, 6). The neuromechanism of

SARS-CoV-2 is not clearly defined, both a systematic mechanism and direct

neurotropism have been proposed (7). Recent studies suggest that SARS-CoV-2

triggered the COVID-19 pandemic, and possesses brain neurotropism primarily through

binding to the ACE2 on neurons (5), or through tunneling nanotubes (8). In addition to

direct invasion of neuronal cells, the activation of microglia and/or astrocytes might

contribute to the onset and progression of neurological disorders through abnormal

maintenance of homeostasis, leading to altered neuronal activities, and thus is

considered critical in defining the neurological damage and neurological outcome of

COVID-19 (9).

As the principal innate immune cells of the brain, microglia form the primary focus of

research in the field of neuroimmune disease. Microglia are the most dynamic neural

cells found to date (10-12). Their profoundly dynamic processes constantly survey the

local microenvironment, monitor neuronal activity (13, 14), and respond to infection and

injury by releasing pro-inflammatory molecules and phagocytic clearance of apoptotic

cells (15). During a homeostatic situation, secretion of molecules and phagocytosis by

microglia likewise maintain synaptic transmission and plasticity (16, 17). In contrast,

hyperactive microglia can be pathogenic and are associated with symptoms of

psychiatric disorders, including anxiety disorders and major depression (18), which have

commonly appeared during the COVID-19 pandemic. The activity of microglia can be

modulated by neuronal activity (19-21), suggesting the existence of microglia-neuron

crosstalk. Despite the high potential relationship between SARS-CoV-2 infection and

the neuroimmune system, the mechanisms by which SARS-CoV-2 infection activates

pro-inflammatory microglia regulates microglia-neuron interaction and alters neuronal

activity have not been adequately studied.

Further research is essential to comprehensively unravel the mechanisms underlying

PASC and its neurological manifestations, as this knowledge is critical for developing

effective management and treatment strategies for affected patients. Mouse models

serve as invaluable tools in conducting such studies. Unfortunately, mice are naturally

resistant to ancestral SARS-CoV-2 infection due to the low affinity of the viral spike (S)

glycoprotein to mouse ACE2 receptors (mACE2) (22, 23). To overcome this limitation,

several strategies have been employed to overexpress human ACE2 (hACE2) in mice.

These approaches include the delivery of exogenous hACE2 using a replication-

deficient adenovirus (Ad5-hACE2) or Adeno-associated Virus (AAV-hACE2), and the

generation of K18-hACE2 transgenic mice (24-26). These models have been used to

investigate the acute infection of SARS-CoV-2 as well as the neurological

consequences (5). However, it should be noted that these models have drawbacks that

limit their application in studying long-term PASC after SARS-CoV-2 infection. The

former may be incapable of getting a brain infection, whereas the latter may acquire

severe artificial multiorgan infections with a high mortality rate, rendering them poor

candidates for long-term post-infection behavioral testing.

A highly virulent mouse-adapted SARS-CoV-2 strain, SARS2-N501Y$_{MA30}$, was

successfully isolated through serial passage of a recombinant SARS-CoV-2/spike

N501Y virus in BALB/c mice (27). After SARS2-N501Y$_{MA30}$ infection, we observed viral

titers, viral genomic RNA (gRNA), and subgenomic RNA (sgRNA) in both the lungs and

the brains of C57BL/6 mice, indicating the occurrence of virus infection. Fourteen days

after the SARS2-N501Y$_{MA30}$ infection, the mice displayed abnormal anxiety- and

depression-like behaviors in multiple behavioral paradigms including the open-field test,

elevated plus maze test, tail suspension test, and forced swimming test. Due to the

critical role of the amygdala in anxiety and depression-like behaviors in rodents and

humans (28), we first examined this region and discovered microglia activation following

SARS2-N501Y$_{MA30}$ infection. Furthermore, heightened microglia-dependent neuronal

activity in the amygdala was evident upon exposure to the spike protein. Our findings

uncover the neurotropic potential of SARS-CoV-2 and its direct link to anxiety- and

depression-like behaviors through the activation of microglia-mediated

neuroinflammatory pathways. This study sheds valuable insights into the neurological

repercussions of SARS-CoV-2 infection and suggests microglia as prospective

therapeutic targets for mitigating virus-induced neurobehavioral deficits. Understanding

the neurological basis of COVID-19-related neuropsychiatric symptoms is critical for

developing effective treatments to reduce the long-term impact of PASC on mental

health.

MATERIALS AND METHODS

Mice, virus, and cells

Specific pathogen-free 6-9 weeks male and female Balb/c and C57BL/6 mice were

purchased from the Jackson Laboratory. All protocols were approved by the Institutional

Animal Care and Use Committees of Cleveland Clinic-Florida Research and Innovation

Center (CC-FRIC) and the University of Tennessee Health Science Center (UTHSC).

The mouse-adapted SARS-CoV-2-N501Y$_{MA30}$ was provided by Drs. Stanley Perlman

and Paul McCray at the University of Iowa, USA (27). The virus was propagated in

Calu-3 cells and tittered by plaque assay in VeroE6 cells. Calu-3 cells were maintained

in minimal essential medium (MEM) supplemented with 20% fetal bovine serum (FBS),

0.1 mM nonessential amino acids (NEAA), 1 mM sodium pyruvate, 2 mM l-glutamine,

1% penicillin and streptomycin, and 0.15% NaHCO$_3$ at 37°C in 5% CO$_2$. Vero E6 cells

were grown in Dulbecco's modified Eagle medium (DMEM) supplemented with 10%

FBS, 0.1 mM NEAA, 1 mM sodium pyruvate, 2 mM l-glutamine, 1% penicillin and

streptomycin, and 0.15% NaHCO$_3$ at 37°C in 5% CO$_2$.

Virus infection and titration.

The research involving SARS-CoV-2 was conducted within the biosafety level 3 (BSL3)

Laboratory at CC-FRIC. Balb/c or C57BL/6 mice were gently anesthetized using

isoflurane and subsequently intranasally infected with 10^4 FPU of SARS-CoV-2-

N501Y$_{MA30}$. Post-infection, daily monitoring and weight measurements of the mice were

conducted. Tissues were aseptically collected and dissociated in PBS using disposable

homogenizers. The viral preparation or supernatants from lung or brain tissue

homogenates were subject to sequential dilution in DMEM. These diluted samples were

then introduced to VeroE6 cells in 12-well plates to conduct plaque assays (29). After

one hour of incubation, the viral inoculums were removed, and the cells were overlaid

with a 1.2% agarose solution supplemented with 4% FBS. After 3-day incubation, the

cells were fixed with formaldehyde, and the overlays were meticulously eliminated,

facilitating visualization of the resulting plaques through the application of a 0.1% crystal

violet stain.

Behavioral experiments

C57BL/6 mice aged 6-9 weeks were utilized to investigate behavioral responses using a

battery of tests as described in **Fig. 2**. We conducted a well-established behavioral test

battery in mice fourteen days after intranasal administration of SARS2-N501Y$_{MA30}$ (10^4

PFU/ mouse). See detailed descriptions below.

Anxiety-like behaviors

Open Field Test Assessed exploratory and anxiety behaviors in an open-field box for 6

minutes. The Open Field task was used to determine the general activity levels, gross

locomotor activity, exploration habits of the mice, and anxiety-like behaviors. The

evaluation occurred within a cubic, white plastic enclosure measuring 30 cm × 30 cm ×

30 cm. Mice were introduced to a corner of the arena and given 6 minutes to acclimate

to the unfamiliar surroundings. Their activities were recorded via video. Subsequently,

the recorded footage, encompassing metrics such as distance traversed, velocity,

resting intervals, movement duration, and time spent in both the center and corners of

the arena, was analyzed utilizing Anymaze software. This examination serves as an

initial assessment of locomotor activity and anxiety-related behaviors, particularly

focusing on the duration spent in the central area.

Elevated Plus Maze Test Evaluated anxiety-like behavior by recording entries into and

time spent in the open arms of a maze for 6 minutes. It was conducted as previously

described (30). The EPM apparatus, 40 cm high from the floor, consists of two open

arms (35 × 10 cm) and two closed arms (35 × 10 cm), those two parts stretch

perpendicular to each other and connect in a center platform (5 cm). Mice were placed

in the center zone facing an open arm and allowed to explore the maze freely for 6

minutes. The time spent in open arms, closed arms, and behaviors of head dipping,

sniffing, and grooming were analyzed.

Anxiety-like behaviors

Tail Suspension Test Conduct the tail suspension test to assess depression-like

behaviors. Suspend the mice by their tails for 6 minutes using adhesive tape in a

controlled environment. Record and analyze the duration of immobility during the

predetermined time period (6 minutes). The state of immobility is defined by the

absence of movement. The detection of immobility time in these tests serves as a

behavioral measure to evaluate the impact of various interventions, such as drug

treatments or genetic manipulations, on the animal's response to stress or despair.

Forced Swimming Test Conduct the forced swimming test to assess depression-like

behavior. Place the mice individually in a water-filled container for 6 minutes and record

their swimming and immobility behaviors. Manually analyze the behavioral results by

determining the immobility time for each mouse. Compile the data and calculate the

mean immobility time for each experimental group.

Bulk RNA sequencing

C57BL/6J mice were intranasally (i.n.) infected with 10^4 FPU of SARS-CoV-2-

N501Y$_{MA30}$, as described previously. At specified time points post-infection, whole

brains were harvested for RNA extraction using the RNeasy Lipid Tissue Mini Kit

(74804, Qiagen, Hilden, Germany). RNA concentration and purity were determined by

NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA).

Library preparation was performed with the Illumina Stranded mRNA Prep Kit

(20040532, Illumina, San Diego, CA, USA), adhering to the manufacturer's instructions.

Sequencing was executed on the DNBSEQ-G400 platform (Innomics Inc. Cambridge,

MA, USA). Data analysis and visualization was conducted on the Dr. Tom online

analysis platform, utilizing the following pipeline: 1. Data Cleaning: Sequencing data

was filtered with SOAPnuke, yielding clean reads stored in FASTQ format. 2. Alignment:

Clean reads were aligned to the reference genome using HISAT2. 3. Fusion Gene

Detection: Ericscript (0.5.5-5) was employed to identify fusion genes. 4. Differential

Splicing Gene (DSG) Analysis: rMATS (v4.1.1) was utilized for detecting DSGs. 5. Gene

Set Alignment: Bowtie2 aligned the clean reads to the gene set. 6. Expression

Quantification: RSEM (v1.2.28) calculated gene expression levels, providing read count,

FPKM, and TPM metrics.7. Differential Expression Gene (DEG) Analysis: DESeq2

conducted DEG analysis, with a significance threshold set at Q value $\leqslant$ 0.05. 8.

Visualization: A heatmap of DEGs was generated using Pheatmap, based on the DEG

analysis results. 9. Phenotypic Insight: GO (Gene Ontology, http://www.geneontology

.org/) and KEGG (Kyoto Encyclopedia of Genes and Genomes, https://www.kegg.jp/)

enrichment analyses of annotated DEGs were performed using Phyper, based on the

Hypergeometric test. Significance levels of terms and pathways were adjusted by Q

value, with a stringent cutoff (Q value $\leqslant$ 0.05).

Quantitative real-time PCR analysis of viral RNA.

Total cellular RNA was isolated using the Direct-zol RNA miniprep kit (Zymo Research,

Irvine, CA) following the manufacturer's protocol including a DNase treatment step.

Total RNA (200 ng) was used as the template for first-strand cDNA synthesis. The

resulting cDNA was used to quantify the SARS-CoV-2 RNA levels by real-time

quantitative PCR using Power SYBR green PCR master mix (Applied Biosystems,

Waltham, MA). Average values from duplicates of each sample were used to calculate

the viral RNA level relative to the GAPDH gene and presented as 2−ΔCT, as indicated

(where CT is the threshold cycle). CT values of gRNA and sgRNA from uninfected mice

(0 dpi) are constantly > 35. The sequences of the primers used are listed in the **S1**

Table.

Brain slice preparation, S1 protein perfusion, and fEPSP recordings.

Mice were euthanized with overdosed isoflurane and whole brains were dissected into

pre-oxygenated (5% CO_2 and 95% O_2) ice-cold high sucrose dissection solution

containing (in mM): 205 sucrose, 5 KCl, 1.25 NaH_2PO_4, 5 $MgSO_4$, 26 $NaHCO_3$, 1 $CaCl_2$,

and 25 Glucose and sliced into 300 μm on a Leica VT1000S vibratome (31). The

transverse amygdala slices were then transferred into the normal artificial cerebrospinal

fluid (ACSF) containing (in mM): 115 NaCl, 2.5 KCl, 2 $CaCl_2$, 1 $MgCl_2$, 1.25 NaH_2PO_4,

11 glucose, 25 $NaHCO_3$, bubbled with 95% O_2/5% CO_2, pH 7.35 at 20°C-22°C. Slices

were incubated in the ACSF at least 1 hour before recording. Individual slices were

transferred to a submersion-recording chamber and were continuously perfused with

the 5% CO_2/95% O_2 solution (~3.0 ml/min) at room temperature (20°C - 22°C). The

spike protein (BPSbIoscience, #510333) was diluted to 167 ng/ml and perfused to the

brain slices in the ACSF. For the field EPSP (fEPSP) experiments, neurons were held in

current-clamp mode with a pipette solution containing (in mM): 2 KCl (mOsm = 290,

adjusted to pH 7.25 with KOH). A concentric bipolar stimulating electrode (FHC Inc.,

Bowdoin, ME) was positioned in the cortical inputs in the amygdala slices. Away from

the stimulating electrode around 400 μm is a glass recording electrode. EPSPs were

recorded in current-clamp mode every 20 seconds and continuously recorded the

EPSPs for at least 1 hour. Data were acquired at 10 kHz using Multiclamp 700B and

pClamp 10.

ATP measurement

Brains were removed from the skull and dissected. Brain slices were then placed in 200

μl ice-cold PBS, in the presence of vehicle or spike protein (167 ng/ml, BPS BIoscience,

#510333) for the indicated time (10, 30, 60 minutes). To quantify ATP released from the

brain slice, we employed an ATP Determination kit (Invitrogen, A22066). Briefly, a 100

μl reaction mixture was added to the 96-well cell culture plate, which contains a 10 μl

sample or standard solution. After 15 minutes of incubation in the dark, the plate was

read using a Synergy Neo2 hybrid multimode microplate reader (BioTek, Winooski, VT,

USA). ATP concentrations were determined by reference to a standard curve.

Histology.

Tissues (lungs, brain) were collected and fixed in zinc formalin. Following fixation, the

lungs were processed for paraffin embedding and sliced into a 4 µm section and the

brain was sectioned into 30 µm by a vibratome for subsequent hematoxylin and eosin

(H&E) staining by Immunohistochemistry Core of Cleveland Clinic Lerner Research

Institute and Immunohistochemistry Core at the University of Tennessee Health Science

Center. We have used two serial lung sections (six fields/section) from each animal and

a total of 4 to 5 animals per group. Acute lung injury severity was evaluated with ATS

guidelines (32) for neutrophil infiltration in alveolar and interstitial space, hyaline

membranes, alveolar wall thickening, and proteinaceous debris deposition. Briefly, a

scoring system (0-2) was employed for each of the criteria mentioned. An average

score of 0 indicated absence of injury, 1 indicated mild to moderate injury, and 2

indicated severe injury.

Immunohistochemistry

Following the behavioral procedures indicated in the text and figures, the mice were

euthanized with overdosed isoflurane and were fixed in Zinc Formalin. Following

fixation, we used a vibratome (Leica VT-1000S) to dissect 30 µm amygdala coronal

slices, which were collected in ice-cold PBS. To complete immunofluorescence staining,

slices were placed in Superblock solution (Thermo Fisher Scientific) plus 0.2% Triton X-

100 for 1 hour and incubated with primary antibodies (1:1000 dilution) at 4°C for 24

hours (31). Primary antibodies we used include mouse monoclonal antibody to dsRNA

(Millipore Cat# MABE1134); Rabbit polyclonal to Iba1 (Abcam Cat# ab108539); Rabbit

polyclonal to GFAP (Abcam Cat# ab7260) and mouse anti-NeuN (Cell Signaling Cat#

24307). We then washed and incubated slices for one hour with secondary antibodies

diluted at a ratio of 1:200 (Alexa Fluor 488 goat anti-rabbit IgG (H+L) (Thermo Fisher

Scientific Cat# A-11008); Alexa Fluor 594 goat anti-mouse IgG (H+L) (Thermo Fisher

Scientific Cat# A-11032); Alexa Fluor 647 goat anti-mouse IgG (H+L) (Thermo Fisher

Scientific Cat# A-21235). VectaShield H-1500 (Vector Laboratories Cat# H-1500) was

used to mount slices, while regular fluorescent DIC microscopy and confocal

microscopy were used to image the slices.

For SARS-CoV-2 antigen detection, tissue slides underwent a series of incubation

steps. Initially, slides were treated with a blocking reagent (10% normal goat serum) for

30 minutes to reduce non-specific binding. Subsequently, a rabbit polyclonal antibody

against the SARS-CoV-2 nucleocapsid protein (dilution: 1:4,000, catalog number:

40143-T62, Sino Biological) was applied to the slides for 15 minutes. Following the

primary antibody incubation, slides were further incubated with Rabbit Envision (Dako)

and diaminobenzidine (Dako) as a chromogen to visualize the antigen-antibody

complexes.

Dendrites and spine morphologic analyses.

After their behavioral testing, we measured the density of dendritic spines on amygdala

neurons by using Golgi staining (n = 5 per group). Mice brains were collected, fixed, and

processed for the Golgi staining according to the protocol provided by the FD Rapid

Golgi Stain Kit (PK401, FD NeuroTechnologies). All images were deconvolved within

the Zeiss Application Suite software. The number of dendritic spines was analyzed

using plug-in SpineJ as described (33), with modifications. Spines were examined on

dendrites of amygdala neurons that satisfied the following criteria: (1) presence of

untruncated dendrites, (2) dark and consistent Golgi staining throughout all dendrites,

and (3) visual separability from neighboring neurons. We counted the number of

dendritic spines along the second dendritic branch at distances from 50 to 100 μm from

the soma, in images obtained at 630× magnification. For each neuron, three to five

dendritic segments 10 μm in length were analyzed. For each group, 6-10 neurons/mice

were analyzed. We used ImageJ software to analyze spines. The analysis of dendritic

spines includes the number of spines and spine density, which are critical indicators of

synaptic function (34, 35). Spines were classified as thin if they had a long, slender neck

and small, Spine length > 0.5 μm; Head perimeter = 2 μm to 3 μm; mushroom, if they

had a well-defined, thick neck and spine length > 0.5 μm; Head perimeter ≥ 3 μm; or

339 stubby, if they were short and thick, without a distinguishable neck, spine length ≤ 0.5

340 µm; or filopodia if they were long and curved, spine length > 0.5 µm, head perimeter ≤ 2

341 µm.

343 **Statistical analysis.**

344 One-way ANOVA and Tukey's post-hoc multiple comparison tests were used for

345 statistical comparison of groups. An unpaired Student's t-test was used to compare

346 results between the two groups. $p < 0.05$ was considered statistically significant, and we

347 did not exclude potential outliers from our data except the ones that did not receive

348 successful aversive conditioning. The graphing and statistical analysis software

349 GraphPad Prism 8 was used to analyze statistical data, which was presented as means

350 ± SEM. Sample sizes (n) are indicated in the figure legends, and data are reported as

351 biological replicates (data from different mice, and different brain slices). Each group

352 contained tissues pooled from 4-5 mice. Due to variable behavior within groups, we

353 used sample sizes of 10-16 mice per experimental group as we previously described in

354 earlier experiments (31). In behavioral studies, we typically studied groups with five

355 randomly assigned animals per group. The experiments were repeated with another set

356 of four animals until we reached the target number of experimental mice per group.

358 **RESULTS**

359 **Neuroinvasion following Respiratory Infection of SARS-CoV-2**

360 The inherent low affinity of the ancestral viral spike (S) glycoprotein for mouse ACE2

361 (mACE2) renders mice naturally resistant to SARS-CoV-2 infection (22, 23). However,

362 specific mutations in the viral spike protein, such as Q498Y, P499T, and N501Y, can

363 enhance binding to mACE2, resulting in asymptomatic infection in mice (36). Through

364 serial passages of a recombinant SARS-CoV-2/spike N501Y virus in BALB/c mice, we

365 successfully isolated a highly virulent mouse-adapted SARS-CoV-2 strain, designated

366 as SARS2-N501Y$_{MA30}$ (27). Interestingly, young C57BL/6 mice exhibited reduced

367 sensitivity to symptomatic infection by SARS2-N501Y$_{MA30}$ **(Fig. 1A)**, making them an

368 optimal model for studying the long-term repercussions of COVID-19. Thus, we

369 intranasally infected mice with 10^4 PFU/mouse of SARS2-N501Y$_{MA30}$ (27). Comparable

to our previous findings, young Balb/c mice developed lethal disease, whereas young

C57BL/6 mice only displayed minimal weight loss of less than 20% and swift recovery

within approximately one week **(Fig. 1B, C)**. In C57BL/6 mice, viral titers peaked at 2

days post infection (dpi) in the lungs, followed by the presence of viral titers in the brain

peaking at 4 dpi **(Fig. 1D, E)**. Consistent with the observation, both viral genomic RNA

(gRNA) and subgenomic RNA (sgRNA) were identified in the lung **(Supplemental**

Fig.1) and brain **(Fig. 1F, G)**. Immunofluorescence targeting double-stranded RNA

(dsRNA) further validated virus RNA replication presence in amygdala brain slices at 4

dpi, which subsequently disappeared by 14 dpi. **(Fig. 1H-J)**, with similar observations in

the prefrontal cortex (PFC), albeit with slight distinctions **(Fig. 1K)**. These results

collectively suggest direct viral infiltration into the brain, alongside respiratory infection.

Remarkably, the production of infectious SARS-CoV-2 appeared to be controlled in the

brain, as the titer is extremely low **(Fig 1E)** and it did not induce major pathological

changes within brain tissues **(Supplemental Fig. 2)**. Further investigation revealed that

SARS-CoV-2 replication was predominant in neuron cells in the amygdala, rather than

in microglia or astrocytes **(Fig. 1L-N).** These findings indicated a SARS2-N501Y$_{MA30}$

neuroinvasion with a peak at 4 dpi followed by a quick clearance.

Histological analysis of lung samples revealed acute lung injury using American

Thoracic Society (ATS) guidelines for experimental acute lung injury (ALI) based on

neutrophil infiltration in alveolar and interstitial space, hyaline membranes, alveolar wall

thickening, and proteinaceous debris deposition (32). SARS-CoV-2 infection induced

mild lung injury which peaked on day 6, accompanied by viral antigen presence at 2 dpi

and disappeared after 6 dpi **(Supplemental Fig 3)**. However, these changes, including

infiltration and viral presence, resolved after a 14-week period. To explore host defense

responses in SARS-CoV-2-infected mice, we assessed cytokine/chemokine gene

expression in the lungs **(Supplemental Fig 1)**. Notably, these genes exhibited

significant upregulation at 2 dpi or 4 dpi, gradually subsiding to baseline levels after 6

dpi. These findings collectively illustrate a mild respiratory disease in the infected mice,

which subsequently resolved within 14 days post-infection.

SARS2-N501Y$_{MA30}$ infection induces significant changes in anxiety- and depression-like behaviors.

Our investigation successfully detected viral dsRNA within the amygdala **(Fig. 1K)**, an area pivotal in coordinating defensive responses in mice (37) and emotional behaviors in humans (28). This finding prompted us to postulate that the infection may still exert influence over the functionality of this region and consequently modify the defensive behaviors of infected mice. To evaluate the impact of SARS-CoV-2 infection on anxiety- and depression-like behaviors, we conducted a well-established behavioral test battery in mice fourteen days after intranasal administration of SARS2-N501Y$_{MA30}$ (10^4 PFU/ mouse) **(Fig. 2A)**. We chose the 14-day time point for behavioral testing since viral infections were observed to have cleared by then. C57BL/6 mice typically recovered from symptomatic infection at 10^4 PFU/mouse within about a week **(Fig. 1B)**. Additional indicators, such as body temperature, fur condition, daily food, and water intake over time (daily records), and mouse locomotion behavior, were similar to those of mock-infected mice at 14 dpi.

In the open-field test, the infected mice exhibited a decreased duration spent in the center area compared to the mock infection group, indicative of heightened anxiety-like behaviors **(Fig. 2B, left)**. Remarkably, both the SARS2-N501Y$_{MA30}$ and mock infection groups demonstrated similar locomotor activity, implying a complete recovery of locomotion post-infection **(Fig. 2B, right)**. In the elevated plus-maze test, mice in the infected group significantly reduced their time spent in the open arms **(Fig. 2C, left)**, while the number of entries into the open arms remained unaltered **(Fig. 2C, right)**, further substantiating an augmentation in anxiety-like behaviors. Consistent outcomes were observed in the tail suspension and forced swimming tests, both are widely used for evaluating depression-like behaviors, where mice in the virus-infected groups displayed increased immobile time compared to the vehicle groups **(Fig. 2D, E)**. These findings substantiate our conclusion that SARS2-N501Y$_{MA30}$ infection leads to an escalation in anxiety- and depression-like behaviors in mice, indicating potential repercussions on brain function that warrant further exploration.

Temporal Gene Signatures in the Brain after SARS-CoV-2 Infection

To further elucidate the molecular mechanisms associated with the observed neuroinvasion and subsequent behavioral alterations, we investigated the temporal dynamics of host transcriptional responses in the brain. This was achieved through bulk RNA sequencing (RNA-seq) on total brain RNA from both mock-infected animal cohorts (0 dpi) and animals infected with SARS2-N501Y$_{MA30}$ at 4 dpi, and 14 dpi, respectively **(Fig. 3 and Supplemental Fig. 4)**. Principal component analysis (PCA) unveiled distinct differences among mock and infected groups with samples from both 4 and 14 dpi **(Supplemental Fig. 4A).** The volcano map of differentially expressed genes (DEGs) revealed significant changes in host genes compared to mock-infected animals or between 4 and 14 dpi based on |log2fold| $\geqslant$ 1.5, Q value $\leqslant$ 0.05 **(Supplemental Fig. 4B)**. In the 4 dpi vs 0 dpi group, 124 genes were upregulated and 6 were downregulated, while in the 14 dpi vs 0 dpi group, 29 genes were upregulated and 2 were downregulated **(Fig. 3A)**. There were only 8 genes upregulated and 5 downregulated between 4 and 14 dpi. Notably, 29 upregulated genes in the 14 dpi group also appeared in the 4 dpi group, many of which were associated with synaptic transmission and neuronal development **(Fig. 3B and Supplemental Fig. 4).** Overall, there were 140 genes significantly changed **(Supplemental Fig. 4).**

We performed pathway analysis based on the Kyoto Encyclopedia of Genes and Genomes (KEGG; http://www.genome.jp/kegg/) and Gene Ontology (GO) based enrichment analysis on the 140 genes **(Supplementary Fig. 4C, E).** Among them, 36 genes were implicated to be involved in different neuronal functions in the brain **(Fig. 3C),** such as synaptic transmission, including Car8, Gabra6, Pcp2, Slc5a1, Tfap2a, or neuronal development, such as Barhl2, Hoxa5, Ppp1r17, Skor1, Fat2, and Nrk. We conducted a KEGG network analysis to delineate the interconnections among the 36 identified genes and identified genes related to synaptic vesicle cycle (slc6a2, slc6a4, and slc6a5), long-term depression (Ppp1r17 and Grid2), and neuroactive ligand-receptor interaction (Grid2, Glra1, and Gabra6) **(Fig. 3D).** In conclusion, our comprehensive RNA-seq analysis has unveiled significant changes in host gene expression following viral infection. These alterations suggest a reprogramming of

neuronal transcriptional states, corroborating the observed behavioral modifications.

Moreover, the data indicate a profound influence on synaptic function and

neurodevelopmental processes, underscoring the far-reaching effects of SARS-CoV-2

on the central nervous system.

SARS2-N501Y$_{MA30}$ infection induces spine remodeling in the amygdala.

To complement the gene expression changes revealed by our RNA-seq analysis, we

next examined the morphological alterations in neurons within the amygdala, a region

implicated in depression-related behaviors. This investigation aimed to confirm the

impact of SARS2-N501Y$_{MA30}$ infection on synaptic plasticity through the observation of

dendritic spine remodeling. Dendritic morphology has been widely implicated in the

mechanisms governing synaptic plasticity (35, 38, 39). Dendritic spines constitute the

principal target of neurotransmission input within the CNS (40), and their density and

structure form the foundation for physiological alterations in synaptic efficacy that

underlie learning and memory processes (41). We hypothesized that dendritic structure

and plasticity undergo alterations following SARS2-N501Y$_{MA30}$ infection. Fourteen days

after SARS2-N501Y$_{MA30}$ infection, brain slices containing the amygdala were

meticulously dissected, fixed, and subjected to Golgi staining to label the spines **(Fig.**

4A). Neurons were randomly selected from within the basolateral amygdala, with a

focus on secondary branches located 50-100 μm away from the soma **(Fig. 4B, C)**.

Subsequently, spine morphology was manually visualized and analyzed **(Fig. 4D)**.

To characterize spine morphology, we categorized dendritic processes into two distinct

morphological classes: mature and immature spines (42). Mature spines, predominantly

exhibiting a "mushroom-like" morphology, exhibit more stable postsynaptic structures

enriched in α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors

(AMPARs) and are considered functional spines. Conversely, immature spines,

characterized by their thin, stubby, and filopodial features, represent unstable

postsynaptic structures with transitional properties. Immature dendritic spines are

believed to hold the potential for future synaptic plasticity, either maturing into functional

spines or disappearing from the dendrite (43). Spine categories were identified based

on parameters described in previous studies and Methods (35, 44).In the brains of mice

infected with SARS2-N501Y$_{MA30}$ (14 dpi), we found variations in spine subtypes, notably

an elevated proportion of mature spines compared to the mock infection group **(Fig.**

4E). Combined with the gene expression profiles from **Fig. 3** and observations on

dendritic morphology, our findings indicate enhanced synaptic plasticity in amygdala

neurons fourteen days after SARS2-N501Y$_{MA30}$ infection.

Spike protein increases microglia-dependent fEPSPs in amygdala slices

Exposure to the SARS-CoV-2 spike protein has emerged as a topic of interest in

neurobiology, particularly regarding its impact on neuronal function. Previous research

has provided initial insights into how spike protein can influence neuronal morphology

and function (45-47). To further explore the mechanistic aspects of this phenomenon,

we focused on the amygdala, a region critically involved in emotional processing and

defensive behaviors. We perfused amygdala slices from mice with the recombinant

B1.351 variant spike protein (167 ng/ml), which has a high affinity to mouse ACE, and

monitored changes in fEPSPs for 2 hours **(Fig. 5A)**. We observed a significant increase

in fEPSPs following exposure to the B1.351 variant spike protein, indicating heightened

synaptic transmission and neuronal excitability **(Fig. 5B, C)**. Despite the fact that

whether SARS-CoV-2 spike protein increases or decreases neuronal activity is

controversial (45-48), this observation is noteworthy, as it suggests that the spike

protein, when directly interacting with neurons, has the potential to amplify their

responsiveness to incoming signals. This heightened neuronal response could, in turn,

render individuals more sensitive to various stimuli, potentially leading to alterations in

behaviors or physiological reactions. Interestingly, the administration of 50 µM

Resveratrol, a known inhibitor of microglial activation (49), effectively counteracted the

spike protein-induced enhancement of fEPSPs **(Fig. 5C, D)**. This finding suggests that

microglia, the intrinsic resident immune sentinels of the brain, may play a pivotal role in

mediating the effects of the spike protein on neuronal activity.

Microglia activation after SARS-CoV-2 infection or exposure to spike protein

Following the discovery in **Fig. 5**, we conducted further investigations into microglial

reactivity during SARS2-N501Y$_{MA30}$ infection. Through immunofluorescence staining for

Iba1, we documented a significant activation of microglia, evidenced by their

morphological shift from ramified to amoeboid shapes (50). This shift was notable at 4

days post-infection, by 14 days post-infection, microglia activity had not fully reverted to

the baseline quiescent state observed at 0 dpi, although it was substantially reduced

compared to 4 dpi, as illustrated in **Fig. 6A**. This finding provides further support for the

hypothesis that microglial activation may contribute to the observed alterations in

neuronal activities within the amygdala following SARS-CoV-2 infection.

Microglia, the brain's resident immune cells, are highly dynamic, engaging with neurons

and other brain cells upon activation. Their processes reach out to neighboring cells,

forming a complex network of interactions crucial for understanding the neuroimmune

impact of SARS-CoV-2 infection (51). To further elucidate the mechanism by which the

spike protein promoted microglia activation, we visualized microglia by

immunofluorescence staining for Iba1 in mouse amygdala slices. Our observations

confirmed the activation of microglia following SARS-CoV-2 B1.351 spike protein

perfusion **(Fig. 6B, C)**. Interestingly, the activation was attenuated by the purinoceptor

P2Y$_{12}$ (P2Y12R) antagonist ticlopidine **(Fig. 6C, D)**, indicating that the P2Y12R

signaling pathway mediates this response. P2Y$_{12}$, selectively expressed in microglia

(52), serves as the main adenosine triphosphate (ATP) receptor, initialing microglial

activation and mobilization in response to ATP, a potent neuronal chemoattractant

released from neurons (19, 53). Additionally, it is worth noting that SARS-CoV-2

infection has been shown to induce ATP release from host cells, including neural cells,

via specific channels (54, 55), a phenomenon we corroborated by detecting elevated

ATP levels in brain slices perfused with spike protein **(Fig. 6E, F)**.

Brain Immune Inflammation and Anti-Viral Responses during Viral Infection

Cytokines and chemokines serve as crucial indicators of anti-viral response and

immune inflammation. To delineate the immune responses within the brain during

SARS-CoV-2 infection, we conducted quantitative reverse-transcription PCR (RT-

qPCR) to profile cytokines and chemokines expression across various time points in the

brain **(Fig. 7)**. Contrary to expectations, the expression levels of interferon genes,

including IFN-β, γ, and λ, as well as the tumor necrosis factor gene, remained unaltered

in the infected brain tissues. However, it is noteworthy that transcripts encoding pro-

inflammatory cytokines, such as interleukin-1α (IL-1α), interleukin-6 (IL-6), interleukin-8

(IL8/CXCL1), and especially monocyte chemoattractant protein-1 (MCP1/CCL2),

exhibited significant upregulation at 2- or 4-days post-infection. These temporal

increases coincided with the presence of viral RNA in the brain. Importantly, it is worth

highlighting that the alterations observed in immune gene expression within the brain

were modest in comparison to those observed in the lungs and returned to baseline

levels after 6 days post-infection. This discovery underscores the presence of restrained

immune inflammation and anti-viral responses within the brain. The observed changes

in immune gene expression warrant further comprehensive investigations to unravel the

underlying mechanisms and identify potential targets for therapeutic intervention.

DISCUSSION

Neurotropism of SARS-CoV-2 and Behavioral Alterations

Our findings revealed the presence of SARS-CoV-2 in the CNS following intranasal

inoculation in a SARS2-N501Y$_{MA30}$ infection mouse model **(Fig.1)**. These results are

consistent with previous reports indicating that SARS-CoV-2 can invade the human

brain (5, 6), supporting the notion of neurotropism. SARS-CoV-2 probably enters the

central nervous system (CNS) through two potential pathways (56). The first involves

accessing the CNS via the neural-mucosal interface in the olfactory mucosa, enabling

the virus to spread from the periphery olfactory neurons into the neurons of the olfactory

bulb. The second pathway is through entry into the brain via blood circulation,

potentially breaching the blood-brain barrier (BBB). In this scenario, the integrity of the

BBB could be disrupted by inflammatory responses triggered by the infection.

Notably, the observed viral invasion of the CNS was accompanied by significant

alterations in anxiety- and depression-like behaviors, as evidenced by behavioral

assays **(Fig. 2)**. Anxiety and depression are prevalent neuropsychiatric disorders

associated with significant morbidity and mortality worldwide. The link between other

viral infections and mental health disturbances has been reported in previous studies,

with evidence of viral neuroinvasion contributing to behavioral changes (57, 58). Our

study provides valuable evidence supporting a direct association between SARS-CoV-2

neurotropism and anxiety- and depression-like behaviors. These findings underscore

the importance of considering the potential neuropsychiatric consequences of COVID-

19 and highlight the need for comprehensive mental health assessments and

interventions during and after the acute phase of infection.

Microglial Activation, Neuroinflammation, and Neuronal Activity

Microglia, as the primary immune cells of the CNS, play a critical role in maintaining

brain homeostasis and protecting against infections (59). Upon encountering pathogens

or damage signals, microglia undergo rapid activation, transforming from a surveillant to

an immune-responsive phenotype. In our study, we observed robust microglial

activation in brain regions where viral particles were detected, suggesting a potential

role for microglia in mediating SARS-CoV-2-induced neurobehavioral alterations. While

microglia itself may not be a direct target of SARS-CoV-2 infection (4, 60), the initial

invasion of SARS-CoV-2 in the neuron cells of the brain likely triggers their activation.

Microglial activation can lead to neuroinflammation, characterized by the release of pro-

inflammatory cytokines, chemokines, and reactive oxygen species (ROS). Chronic

neuroinflammation has been implicated in the pathogenesis of various neuropsychiatric

disorders, including anxiety and depression (61). Notably, our transcriptomic analysis of

SARS-CoV-2-infected brains revealed upregulation of inflammatory and cytokine-

related pathways, supporting the hypothesis of microglia-driven neuroinflammation as a

mechanism underlying behavioral changes. While our whole-brain qPCR and bulk RNA

seq revealed minimal changes in immune responses, it is conceivable that the immune

responses induced by SARS-CoV-2 are relatively subdued and localized to specific

brain regions. Therefore, the comprehensive assessment of whole-brain studies may

underestimate the extent of regional immune responses.

Purinergic receptor $P2Y_{12}$, a G protein-coupled receptor exclusively found in microglia

(52), played a pivotal role in the brain's responses to extracellular nucleotides like ATP

and ADP. Upon activation by ATP, microglia not only accumulate but also modulate

neuronal activity (16, 17), implicating ATP, known for its swift release from activated

neurons (62), as a swift mediator in the neuroimmune response to SARS2-N501Y$_{MA30}$

infection. This activation leads to the release of pro-inflammatory cytokines, which are

known to influence neuronal excitability and synaptic transmission (63), potentially

exacerbating anxiety- and depression-like behaviors (64). Indeed SARS-CoV-2 infection

has been shown to induce ATP release from brain cells, including neural cells, through

ATP-release channels (54, 55), reinforcing the link between infection and rapid ATP-

mediated microglia response. Clinical and laboratory studies support the idea that

elevated cytokines like IL-1β, IL-6, CCL2 and TNF-α associated with the development of

anxiety and depression-like behaviors in mice (65-68), underscoring the contribution of

neuroinflammation to the behavioral manifestations observed following SARS-CoV-2

infection. Our hypotheses detailing the interactions are visually summarized in **Fig. 8**,

providing a clear representation of the underlying mechanisms.

Microglial Inhibition and Neuronal Activity Reversal

To elucidate the specific role of microglia in mediating neuronal activity following SARS-

CoV-2 infection, we selectively inhibit microglia using a pharmacological approach.

Intriguingly, this inhibition markedly reduced the heightened neuronal activity induced by

viral spike protein. This observation indicates a direct involvement of microglia in the

development of abnormal neuronal activity in response to SARS-CoV-2 infection. The

observed neuronal activity rescue upon microglial inhibition aligns with previous studies

showing that microglial activation is a key driver of neuronal activity changes in various

CNS disorders (69). Understanding the interactions between microglia and neurons and

how SARS-CoV-2 infection influences these processes in the amygdala is crucial to

comprehending the neurological consequences of the virus. Such insights could offer

valuable information to guide potential interventions or treatments for neurological

manifestations observed in long COVID. Further research is needed to fully elucidate

the intricate mechanisms underlying microglial activation and its impact on neuronal

functions in response to SARS-CoV-2 infection in the brain.

Inflammatory Pathways and Therapeutic Interventions

Our hypothesis suggests that microglia-driven neuroinflammation contributes to the behavioral changes seen in long COVID-19. This highlights the therapeutic potential of targeting neuroinflammatory pathways to ameliorate anxiety- and depression-like symptoms associated with the disease. The transcriptomic analysis of brains affected by the virus has identified upregulated inflammatory pathways, indicating potential targets for therapeutic interventions. Anti-inflammatory agents and immunomodulatory drugs that selectively dampen microglial activation may offer a promising approach for treating the neuropsychiatric consequences of SARS-CoV-2 infection. However, further research is needed to identify specific inflammatory mediators and downstream signaling pathways responsible for the observed behavioral changes.

Limitations and Future Directions

While our study provides valuable insights into the link between SARS-CoV-2 neurotropism, microglial activation, neuronal activity, and anxiety- and depression-like behaviors, several limitations should be acknowledged.

First, in our pursuit to unravel the etiology of neurological disorders post-COVID-19, we identified behavioral phenotypes emerging within weeks post-infection. The onset of neurological symptoms post-COIVD-19 can vary greatly among individuals, manifesting during the acute phase, post-recovery, or even several months later. The persistence of PASC symptoms in humans can range from weeks to several months, and in some cases, may extend even further. We acknowledge the differences in lifespan and sensitivity to SARS-CoV-2 between our murine model and human subjects. We chose the 14-day time point for behavioral testing as viral infections were observed to have cleared, and the conditions of infected mice had fully recovered. This rapid recovery in the mouse model confers a significant advantage for our mechanistic studies, eliminating the need for a year-long wait. To improve upon our research model and better mimic human conditions, we plan to extend the testing duration by incorporating additional time points up to at least one year in future studies. Furthermore, the prolonged effects of SARS-CoV-2 on mental health necessitate further investigation.

680   Second, we used a murine model, and findings in rodents may not fully translate to
681   human conditions. Future research should explore the temporal dynamics of microglial
682   activation and neuroinflammation throughout different stages of SARS-CoV-2 infection.
683
684   Third, we identified that brain infection in mice developed mild respiratory diseases. We
685   did not further evaluate the case of asymptomatic infection because a clinical study
686   suggested that the risk of developing long-term symptoms in asymptomatic SARS-CoV-
687   2 infected persons was significantly lower than those in symptomatic SARS-CoV-2
688   infection cases (70).
689
Fourth, we utilized a mouse-adapted SARS-CoV-2 strain that shares some key spike
mutations with the Beta variant. However, the specific potential, impact, and severity of
neuroinvasion resulting from different SARS-CoV-2 variants, such as the Omicron
variant, and the potential effects of breakthrough infections in vaccinated individuals,
remain areas that require further investigation to provide a comprehensive
understanding.
696
Fifth, we did not systematically investigate the potential impact of elevated cytokines
and chemokines induced by respiratory COVID-19 on microglial activation and their
potential effects on impairing CNS neuron function. It is well-established that exposure
to chemotherapy drugs, brain radiation, or systemic inflammation can lead to persistent
activation of certain microglia (1, 71-74). Interestingly, a recent study of SARS-CoV-2
infection in AAV-hACE2 sensitized mice or H1N1 influenza infection in mice revealed
that mild respiratory disease-activated white-matter-selective microglia, leading to
oligodendrocyte loss, impaired neurogenesis, and elevated CCL11 levels (75). These
findings highlight the potential impact of systemic inflammation on microglia activation
and brain function. Longitudinal studies in human cohorts are necessary to establish
causality between viral neurotropism, microglial activation, and neuropsychiatric
outcomes. Moreover, investigating potential sex-dependent differences in behavioral
responses and microglial activation can provide a more comprehensive understanding
of COVID-19-associated neuropsychiatric sequelae. Addressing these limitations

through further research is essential to advance our understanding of the complex neurological effects of SARS-CoV-2 infection and its variants. By exploring these areas further, we can gain valuable insights to develop more effective interventions and treatment strategies for individuals affected by COVID-19 and its neurological consequences. The evaluation of the consequences of N501Y$_{MA30}$ infection on the CNS has provided valuable insights, positioning this mouse-adapted strain as an auspicious model for studying the neurological manifestations of PASC. Continuation of research utilizing this mouse model, alongside other relevant models, is pivotal in advancing our understanding of the neurological effects of SARS-CoV-2 and facilitating the development of effective interventions for PASC.

Finally, we utilized bulk RNA-sequencing to analyze gene changes following viral infection. However, this approach may not fully capture gene changes in specific cell types, such as microglia, which could potentially confound the observed neuron-microglia interactions. To precisely elucidate the immune response and subsequent changes in neuronal activity genes, future studies should consider supplementing with single-cell RNA-seq focusing on specific brain regions.

Conclusion

Our study demonstrates that SARS-CoV-2 neurotropism induces anxiety- and depression-like behaviors through the activation of microglia and subsequent neuroinflammation. These findings underscore the importance of considering mental health disturbances in COVID-19 patients and emphasize the need for integrated approaches to address both the physical and mental health aspects of the pandemic. Targeting microglial activation and neuroinflammatory pathways may offer promising therapeutic avenues to mitigate the neurobehavioral consequences of SARS-CoV-2 infection. Further investigations in human cohorts are warranted to validate and extend our preclinical findings for potential translational applications.

AUTHOR CONTRIBUTIONS

J.D. and K.L. conceived the project. J.D., K.L., and Q.G. designed the experiments.

Q.G. and J.D. performed the patch-clamp and immunostaining experiments and data

analysis. S.Z. and K.L. performed the behavior and molecular experiments and data

analysis. K.L., J.D., and Q.G. wrote the manuscript. All authors reviewed and edited the

manuscript.

ACKNOWLEDGMENTS

J.D. is supported by the National Institutes of Mental Health (1R01MH113986), the

Cystic Fibrosis Foundation (002544I221), and the University of Tennessee Health

Science Center start-up fund. K.L. is supported by the Cystic Fibrosis Foundation and

the Cleveland Clinic Florida Research and Innovation Center start-up fund.

CONFLICT-OF-INTEREST

The authors have declared that no conflict of interest exists.

REFERENCES

1.   A. C. Geraghty *et al.*, Loss of Adaptive Myelination Contributes to Methotrexate
Chemotherapy-Related Cognitive Impairment. *Neuron* **103**, 250-265 e258
(2019).

2.   Z. A. Sherif *et al.*, Pathogenic mechanisms of post-acute sequelae of SARS-
CoV-2 infection (PASC). *Elife* **12** (2023).

3.   B. Bowe, Y. Xie, Z. Al-Aly, Postacute sequelae of COVID-19 at 2 years. *Nat Med*
29, 2347-2357 (2023).

4.   M. G. Andrews *et al.*, Tropism of SARS-CoV-2 for human cortical astrocytes.
Proc Natl Acad Sci U S A **119**, e2122236119 (2022).

5.   E. Song *et al.*, Neuroinvasion of SARS-CoV-2 in human and mouse brain. *J Exp*
Med **218** (2021).

6.   S. R. Stein *et al.*, SARS-CoV-2 infection and persistence in the human body and
brain at autopsy. *Nature* **612**, 758-763 (2022).

7.   M. F. DosSantos *et al.*, Neuromechanisms of SARS-CoV-2: A Review. *Front*
Neuroanat **14**, 37 (2020).

8.   A. Pepe, S. Pietropaoli, M. Vos, G. Barba-Spaeth, C. Zurzolo, Tunneling
nanotubes provide a route for SARS-CoV-2 spreading. *Sci Adv* **8**, eabo0171
(2022).

9.   M. G. Savelieff, E. L. Feldman, A. M. Stino, Neurological sequela and disruption
of neuron-glia homeostasis in SARS-CoV-2 infection. *Neurobiol Dis* **168**, 105715
(2022).

10.   D. Davalos *et al.*, ATP mediates rapid microglial response to local brain injury in
vivo. *Nat Neurosci* **8**, 752-758 (2005).

11.   A. Nimmerjahn, F. Kirchhoff, F. Helmchen, Resting microglial cells are highly
dynamic surveillants of brain parenchyma in vivo. *Science* **308**, 1314-1318
(2005).
12.   R. M. Ransohoff, V. H. Perry, Microglial physiology: unique stimuli, specialized
responses. *Annu Rev Immunol* **27**, 119-145 (2009).
13.   H. Wake, A. J. Moorhouse, S. Jinno, S. Kohsaka, J. Nabekura, Resting microglia
directly monitor the functional state of synapses in vivo and determine the fate of
ischemic terminals. *J Neurosci* **29**, 3974-3980 (2009).
14.   M. E. Tremblay, R. L. Lowery, A. K. Majewska, Microglial interactions with
synapses are modulated by visual experience. *PLoS Biol* **8**, e1000527 (2010).
15.   K. M. Lucin, T. Wyss-Coray, Immune activation in brain aging and
neurodegeneration: too much or too little? *Neuron* **64**, 110-122 (2009).
16.   Q. Li, B. A. Barres, Microglia and macrophages in brain homeostasis and
disease. *Nat Rev Immunol* **18**, 225-242 (2018).
17.   D. P. Schafer, E. K. Lehrman, B. Stevens, The "quad-partite" synapse: microglia-
synapse interactions in the developing and mature CNS. *Glia* **61**, 24-36 (2013).
18.   D. J. Stein, M. F. Vasconcelos, L. Albrechet-Souza, K. M. M. Cereser, R. M. M.
de Almeida, Microglial Over-Activation by Social Defeat Stress Contributes to
Anxiety- and Depressive-Like Behaviors. *Front Behav Neurosci* **11**, 207 (2017).
19.   U. B. Eyo *et al.*, Neuronal hyperactivity recruits microglial processes via neuronal
NMDA receptors and microglial P2Y12 receptors after status epilepticus. *J
Neurosci* **34**, 10528-10540 (2014).
20.   A. M. Fontainhas *et al.*, Microglial morphology and dynamic behavior is regulated
by ionotropic glutamatergic and GABAergic neurotransmission. *PLoS One* **6**,
e15973 (2011).
21.   Y. Li, X. F. Du, C. S. Liu, Z. L. Wen, J. L. Du, Reciprocal regulation between
resting microglial dynamics and neuronal activity in vivo. *Dev Cell* **23**, 1189-1202
(2012).
22.   J. Damas *et al.*, Broad host range of SARS-CoV-2 predicted by comparative and
structural analysis of ACE2 in vertebrates. *Proc Natl Acad Sci U S A* **117**, 22311-
22322 (2020).
23.   M. A. A. Mahdy, W. Younis, Z. Ewaida, An Overview of SARS-CoV-2 and Animal
Infection. *Front Vet Sci* **7**, 596391 (2020).
24.   D. V. Glazkova *et al.*, Generation of SARS-CoV-2 Mouse Model by Transient
Expression of the Human ACE2 Gene Mediated by Intranasal Administration of
AAV-hACE2. *Mol Biol* **56**, 705-712 (2022).
25.   J. Sun *et al.*, Generation of a Broadly Useful Model for COVID-19 Pathogenesis,
Vaccination, and Treatment. *Cell* **182**, 734-743 e735 (2020).
26.   J. Zheng *et al.*, COVID-19 treatments and pathogenesis including anosmia in
K18-hACE2 mice. *Nature* **589**, 603-607 (2021).
27.   L. R. Wong *et al.*, Eicosanoid signalling blockade protects middle-aged mice from
severe COVID-19. *Nature* **605**, 146-151 (2022).
28.   P. H. Janak, K. M. Tye, From circuits to behaviour in the amygdala. *Nature* **517**,
284-292 (2015).

29.   L. R. Wong *et al.*, Sensitization of Non-permissive Laboratory Mice to SARS-
825   CoV-2 with a Replication-Deficient Adenovirus Expressing Human ACE2. *STAR
826   Protoc* **1**, 100169 (2020).
827   30.   A. A. Walf, C. A. Frye, The use of the elevated plus maze as an assay of anxiety-
828   related behavior in rodents. *Nat Protoc* **2**, 322-328 (2007).
829   31.   J. Du *et al.*, Transient acidosis while retrieving a fear-related memory enhances
830   its lability. *Elife* **6** (2017).
831   32.   G. Matute-Bello *et al.*, An official American Thoracic Society workshop report:
832   features and measurements of experimental acute lung injury in animals. *Am J
833   Respir Cell Mol Biol* **44**, 725-738 (2011).
834   33.   F. Levet, J. Tonnesen, U. V. Nagerl, J. B. Sibarita, SpineJ: A software tool for
835   quantitative analysis of nanoscale spine morphology. *Methods* **174**, 49-55
836   (2020).
837   34.   C. J. Kreple *et al.*, Acid-sensing ion channels contribute to synaptic transmission
838   and inhibit cocaine-evoked plasticity. *Nat Neurosci* **17**, 1083-1091 (2014).
839   35.   W. J. Wright *et al.*, Silent synapses dictate cocaine memory destabilization and
840   reconsolidation. *Nat Neurosci* **23**, 32-46 (2020).
841   36.   H. Shuai *et al.*, Emerging SARS-CoV-2 variants expand species tropism to
842   murines. *EBioMedicine* **73**, 103643 (2021).
843   37.   F. Rigoli, M. Ewbank, T. Dalgleish, A. Calder, Threat visibility modulates the
844   defensive brain circuit underlying fear and anxiety. *Neurosci Lett* **612**, 7-13
845   (2016).
846   38.   K. M. Woolfrey, D. P. Srivastava, Control of Dendritic Spine Morphological and
847   Functional Plasticity by Small GTPases. *Neural Plast* **2016**, 3025948 (2016).
848   39.   G. Yang, F. Pan, W. B. Gan, Stably maintained dendritic spines are associated
849   with lifelong memories. *Nature* **462**, 920-924 (2009).
850   40.   J. N. Bourne, K. M. Harris, Balancing structure and function at hippocampal
851   dendritic spines. *Annu Rev Neurosci* **31**, 47-67 (2008).
852   41.   C. H. Bailey, E. R. Kandel, K. M. Harris, Structural Components of Synaptic
853   Plasticity and Memory Consolidation. *Cold Spring Harb Perspect Biol* **7**, a021758
854   (2015).
855   42.   A. Peters, I. R. Kaiserman-Abramof, The small pyramidal neuron of the rat
856   cerebral cortex. The perikaryon, dendrites and spines. *Am J Anat* **127**, 321-355
857   (1970).
858   43.   K. P. Berry, E. Nedivi, Spine Dynamics: Are They All the Same? *Neuron* **96**, 43-
859   55 (2017).
860   44.   E. E. Koffman *et al.*, Acid-sensing ion channel 1a regulates the specificity of
861   reconsolidation of conditioned threat responses. *JCI Insight*
862   10.1172/jci.insight.155341 (2022).
863   45.   J. Oh *et al.*, SARS-CoV-2 spike protein induces cognitive deficit and anxiety-like
864   behavior in mouse via non-cell autonomous hippocampal neuronal death. *Sci
865   Rep* **12**, 5496 (2022).
866   46.   C.-Y. Chen, Y.-C. Chou, Y.-P. Hsueh, SARS-CoV-2 D614 and G614 spike
867   variants impair neuronal synapses and exhibit differential fusion ability. *BioRxiv*
868   10.1101/2020.12.03.409763 (2020).

47. M. G. Frank *et al.*, SARS-CoV-2 spike S1 subunit induces neuroinflammatory,
microglial and behavioral sickness responses: Evidence of PAMP-like properties.
Brain Behav Immun **100**, 267-277 (2022).

48. G. Datta *et al.*, SARS-CoV-2 S1 Protein Induces Endolysosome Dysfunction and
Neuritic Dystrophy. *Front Cell Neurosci* **15**, 777738 (2021).

49. S. Zhang *et al.*, Resveratrol Attenuates Microglial Activation via SIRT1-SOCS1
Pathway. *Evid Based Complement Alternat Med* **2017**, 8791832 (2017).

50. A. Vidal-Itriago *et al.*, Microglia morphophysiological diversity and its implications
for the CNS. *Front Immunol* **13**, 997786 (2022).

51. M. Colonna, O. Butovsky, Microglia Function in the Central Nervous System
During Health and Neurodegeneration. *Annu Rev Immunol* **35**, 441-468 (2017).

52. S.-S. Lin, Y. Tang, P. Illes, A. Verkhratsky, The Safeguarding Microglia: Central
Role for P2Y12 Receptors. *Frontiers in Pharmacology* **11** (2021).

53. L. J. Wu, K. I. Vadakkan, M. Zhuo, ATP-induced chemotaxis of microglial
processes requires P2Y receptor-activated initiation of outward potassium
currents. *Glia* **55**, 810-821 (2007).

54. R. Luu *et al.*, Pannexin-1 channel opening is critical for COVID-19 pathogenesis.
iScience **24**, 103478 (2021).

55. D. E. Ribeiro *et al.*, Hyperactivation of P2X7 receptors as a culprit of COVID-19
neuropathology. *Mol Psychiatry* **26**, 1044-1059 (2021).

56. S. M. Burks, H. Rosas-Hernandez, M. Alejandro Ramirez-Lee, E. Cuevas, J. C.
Talpos, Can SARS-CoV-2 infect the central nervous system via the olfactory bulb
or the blood-brain barrier? *Brain Behav Immun* **95**, 7-14 (2021).

57. K. Tomonaga, Virus-induced neurobehavioral disorders: mechanisms and
implications. *Trends Mol Med* **10**, 71-77 (2004).

58. I. Kotsiri *et al.*, Viral Infections and Schizophrenia: A Comprehensive Review.
Viruses **15** (2023).

59. M. E. Tremblay, C. Madore, M. Bordeleau, L. Tian, A. Verkhratsky,
Neuropathobiology of COVID-19: The Role for Glia. *Front Cell Neurosci* **14**,
592214 (2020).

60. D. Beckman *et al.*, SARS-CoV-2 infects neurons and induces neuroinflammation
in a non-human primate model of COVID-19. *Cell Rep* **41**, 111573 (2022).

61. S. J. Rhie, E. Y. Jung, I. Shim, The role of neuroinflammation on pathogenesis of
affective disorders. *J Exerc Rehabil* **16**, 2-9 (2020).

62. R. D. Fields, Nonsynaptic and nonvesicular ATP release from neurons and
relevance to neuron-glia signaling. *Semin Cell Dev Biol* **22**, 214-219 (2011).

63. M. Schafers, L. Sorkin, Effect of cytokines on neuronal excitability. *Neurosci Lett*
437, 188-193 (2008).

64. N. Vogelzangs, P. de Jonge, J. H. Smit, S. Bahn, B. W. Penninx, Cytokine
production capacity in depression and anxiety. *Transl Psychiatry* **6**, e825 (2016).

65. S. Rossi *et al.*, Interleukin-1beta causes anxiety by interacting with the
endocannabinoid system. *J Neurosci* **32**, 13896-13905 (2012).

66. R. K. Farooq, K. Asghar, S. Kanwal, A. Zulqernain, Role of inflammatory
cytokines in depression: Focus on interleukin-1beta. *Biomed Rep* **6**, 15-20
(2017).

67.   M. A. Alshammari *et al.*, Systemic TNF-alpha blockade attenuates anxiety and
depressive-like behaviors in db/db mice through downregulation of inflammatory
signaling in peripheral immune cells. *Saudi Pharm J* **28**, 621-629 (2020).
68.   J. Wojcieszak, K. Kuczynska, J. B. Zawilska, Role of Chemokines in the
Development and Progression of Alzheimer's Disease. *J Mol Neurosci* **72**, 1929-
1951 (2022).
69.   J. Qin, Z. Ma, X. Chen, S. Shu, Microglia activation in central nervous system
disorders: A review of recent mechanistic investigations and development efforts.
Front Neurol **14**, 1103416 (2023).
70.   Y. Ma *et al.*, Long-Term Consequences of Asymptomatic SARS-CoV-2 Infection:
A Systematic Review and Meta-Analysis. *Int J Environ Res Public Health* **20**
(2023).
71.   E. M. Gibson *et al.*, Methotrexate Chemotherapy Induces Persistent Tri-glial
Dysregulation that Underlies Chemotherapy-Related Cognitive Impairment. *Cell*
176, 43-55 e13 (2019).
72.   M. L. Monje, S. Mizumatsu, J. R. Fike, T. D. Palmer, Irradiation induces neural
precursor-cell dysfunction. *Nat Med* **8**, 955-962 (2002).
73.   M. L. Monje, H. Toda, T. D. Palmer, Inflammatory blockade restores adult
hippocampal neurogenesis. *Science* **302**, 1760-1765 (2003).
74.   M. L. Monje *et al.*, Impaired human hippocampal neurogenesis after treatment for
central nervous system malignancies. *Ann Neurol* **62**, 515-520 (2007).
75.   A. Fernandez-Castaneda *et al.*, Mild respiratory COVID can cause multi-lineage
neural cell and myelin dysregulation. *Cell* **185**, 2452-2468 e2416 (2022).

Figure. 1

Figure. 1. Outcomes of Intranasal Infection with SARS2-N501Y$_{MA30}$. (A) Schematic depicting the outcomes of infection in young BALB/c and C57BL/6 mice following administration of 10^4 PFU of SARS2-N501Y$_{MA30}$. **(B-C)** Daily monitoring of body weight (B) and survival (C) in young BALB/c and C57BL/6 mice post-infection. **(D-E)** Virus titers in the lungs (D) and brains (E) of C57BL/6 mice infected with SARS2-N501Y$_{MA30}$ at the indicated days post-infection. **(F-G)** Viral genomic RNA (gRNA) (F) and subgenomic RNA (sgRNA) (G) levels in brain tissues from SARS2-N501Y$_{MA30}$ infected C57BL/6 mice. The levels of viral gRNA and sgRNA were normalized to GAPDH and presented as $2^{\wedge}$-ΔCT (n = 4 or 5 mice per group). CT values for viral genomic RNA (gRNA) or subgenomic RNA (sgRNA) from mock-infected tissues were consistently greater than 35. Statistical significance: ***p = 0.001, **p = 0.0017, determined by ordinary one-way ANOVA. **(H-J)** Immunofluorescence staining targeting dsRNA (red), neuronal nuclear protein (NeuN, green), and nuclei (DAPI, blue) in amygdala brain slices collected from mice infected with SARS2-N501Y$_{MA30}$ infection at 4 or 14 days post (I-J), compared to mock infection (H). Arrows indicate dsRNA and NeuN-positive neurons. Inserts show enlarged single-cell images. **(K)** Percentage of dsRNA-positive cells (dsRNA$^+$) in slices from the amygdala and the prefrontal cortex (PFC). The peak of dsRNA$^+$ cells is observed at 4 dpi. Statistical significance: **p = 0.0012, determined by a two-tailed unpaired Student's t-test. **(L-M)** Immunofluorescence targeting dsRNA (red), astrocyte (glial fibrillary acidic protein, GFAP, green) or microglia (Ionized calcium-binding adaptor molecule 1, Iba1, green), and nuclei (DAPI, blue) in amygdala slices from infected mice. **(N)** Comparison of dsRNA$^+$ cells with NeuN$^+$, GFAP$^+$, and Iba1$^+$ at 4 dpi. . Data are presented as mean $\pm$ SEM.

Figure. 2

Figure. 2. Induction of anxiety- and depression-like Behaviors in Mice Following SARS2-N501Y$_{MA30}$ infection. (A) Experimental design and timeline illustrating the administration of SARS2-N501Y$_{MA30}$ and the behavioral test battery. **(B)** Open field test. Upper: Representative heat map tracking of activity in the mock and SARS2-N501Y$_{MA30}$ infected mice. Lower: Results depicting the total travel distance and time spent in the center area. Statistical significance: **p = 0.0014 (total travel distance); ns, non-significant, p = 0.2927 (center area). **(C)** Elevated plus maze test. Upper: Representative heat map tracking of activity in mock and SARS2-N501Y$_{MA30}$ infected mice in closed arms (C) and open arms (O). Lower: Results indicating the time spent in open arms and the number of entries into open arms. Statistical significance: *p = 0.0299 (time spent in open arms); ns, non-significant, p = 0.6897 (number of entries). **(D-E)** Schematics of the tail suspension (D) and forced swimming (E) tests and results showing immobile time. Statistical significance: *p = 0.0118 (tail suspension test); **p = 0.0029 (forced swimming test). All statistical analyses were performed using a two-tailed unpaired Student's t-test. The analysis includes data from 10 mice in each group. Data are presented as mean $\pm$ SEM.

Figure. 3

Figure. 3. RNA sequencing reveals distinct alterations in the brains of SARS2-N501Y$_{MA30}$ Infected Mice. (A) Histogram represents the corresponding numbers of differentially expressed genes (DEGs) between groups. (|log2fold| ≥ 1.5, Q value ≤ 0.05) **(B)** Venn diagram illustrating the differential gene expression analysis between groups of gene sets. The intersections show shared gene expression patterns, while non-overlapping regions indicate uniquely expressed genes. The numbers within each section denote the count of genes. **(C)** Heatmap displaying hierarchical clustering of 36 selected genes associated with neuron functions. **(D)** KEGG Network analysis of key driver genes and enriched KEGG pathways among the 36 selected genes. Dots symbolize genes, squares represent KEGG pathways, and lines connecting genes to squares signify gene enrichment within the respective pathway.

Figure. 4

Figure. 4. Golgi-Cox Staining of Neurons in the Amygdala of SARS2-N501Y$_{MA30}$ Infected Mice. (A) Experimental design and timeline depicting the administration of SARS2-N501Y$_{MA30}$, dissection of brain slices, and Golgi staining process. **(B-C)** Left: Representative Golgi-Cox staining images of amygdala slices from mock-infected (B) and mice infected with SARS2-N501Y$_{MA30}$ at 14 days post-infection (dpi) (C). Scale bar = 200 µm. Right: Enlarged images from the red boxes in the left images for each respective group. Scale bar = 50 µm. **(D)** Representative dendrite images from the red boxes in (B) and (C). Scale bar = 2 µm. Green arrows point out the mushroom spines. **(E)** Comparison of the percentage of mature spines in the mock and SARS2-N501Y$_{MA30}$ infection groups. Statistical significance: **$p = 0.0038$. The analysis includes data from 11 neurons sourced from 4 mice in each group. Statistical analysis was performed using a two-tailed unpaired Student's t-test. Data are presented as mean $\pm$ SEM.

Figure. 5

Figure. 5. Enhancement of Microglia-Dependent fEPSPs in the Amygdala by SARS-CoV-2 Variant B.1.351 Spike Protein. (A) Schematic representation of field excitatory postsynaptic potential (fEPSP) recordings in the Amygadala brain slices. **(B)** Representative fEPSP traces were recorded at the beginning (0 hours) and the end of the 2-hour recordings, during perfusion with either vehicle or spike protein (S1+S2, B.1.351, β variant, BPS Bioscience #510333, 167 ng/ml). **(C)** Average fEPSP data. The spike protein was applied to the slice (167 ng/ml) at the indicated time point (red trace). In the vehicle group, a mock aqueous buffer solution without the spike protein was used (gray trace). The enhancement of fEPSPs by the spike protein was attenuated by pre-treating brain slices with 50 µM Resveratrol (green trace). **(D)** Summarized amplitudes of the last five fEPSPs as shown in (C). Statistical significance: ***p = 0.0003, **p = 0.0082, determined by one-way ANOVA with Tukey's post hoc multiple comparisons. The analysis includes data from 5 slices in each group. Data are presented as mean $\pm$ SEM.

Figure. 6

Figure. 6. Microglial Activation in Response to SARS-CoV-2 infection and B1.351 Spike Protein. (A) Dynamics of Microglial Activation Post-Infection in the Amygdala of Mice. Immunofluorescence staining for Iba1 highlights microglia in amygdala slices from mice at various time points following infection. An enlarged view of individual microglia provided to detail the changes in shape and morphology indicative of activation states. **(B-D)** Microglial response to perfusion with the spike protein. Immunostaining with Iba-1 illustrates microglial morphology in brain slices after a one-hour perfusion with the (B) vehicle or (C) spike protein (S1+S2, B.1.351, β variant, 167 ng/ml), and (D) spike protein following a 30-minute pretreatment with 50 μM Ticlopidine, a P2Y12R antagonist. The series showcases the microglial response to the spike protein, with additional magnified images highlighting the individual microglia and their morphological analysis using Sholl analysis, presented to the right of each panel. **(E)** Comparison of microglial activation assessed by Sholl analysis, which measures the total branch length and radius of cell area. Statistical significance: ****p < 0.0001, ***p = 0.0006, determined by one-way ANOVA with Tukey's post hoc multiple comparisons. The analysis includes data from 30 cells across 4 slices in each group. **(F)** Quantification of ATP release. Brain slices were incubated in cold PBS with either vehicle or spike protein (167 ng/ml). Supernatants were collected at 10, 30, and 60 minutes post-incubation. The amount of ATP released into the supernatant was quantified. Each dot is an independent brain slice and represents the mean of 2 technical replicates. Statistical significance: *p =0.0374, ****p < 0.0001, determined by two-way ANOVA. Data are presented as mean ± SEM.

Figure. 7

Figure. 7. Cytokines and Chemokines Induced in the Brain of SARS-CoV-2-Infected Mice.
Brains of C57BL/6 mice intranasally infected with 10^4 PFU of SARS2-N501Y$_{MA30}$ were
harvested at indicated days post-infection. Cytokine and chemokine transcripts were measured
by quantitative real-time PCR (qRT-PCR) analyzing total RNA extracted from mock-infected (0
dpi) and infected young C57BL/6 mice. Each brain was collected from one individual mouse.
Mock (0 dpi), 2, 4, 8, 14 dpi: n=4; 6 dpi: n=5. The levels of transcripts were normalized to
GAPDH and presented as 2^-ΔCT. Statistical significance: *p < 0.05, **p < 0.01, ****p < 0.0001,
determined by ordinary one-way ANOVA. Data are presented as mean $\pm$ SEM.

Figure. 8

Figure. 8. Modeling Neurotropism-induced Behavioral Changes by SARS2-N501Y$_{MA30}$.
This schematic illustrates the potential mechanism of SARS-CoV-2 targeting neurons, prompting ATP release, which in turn recruits and activates microglia. The ensuing cascade of microglial response and pro-inflammatory cytokine production is hypothesized to amplify neuronal activity, potentially altering defensive behaviors associated with anxiety and depression.

Supplemental Figure 1. SARS2-N501Y$_{MA30}$ Virus RNA Replication and Host Immune Responses in the Lungs.
Total RNA was extracted from lungs collected from C57BL/6 mice (n=3 to 5), intranasally infected with 10^4 PFU of SARS2-N501Y$_{MA30}$ at designated days post-infection. Viral RNA, as well as host cytokine and chemokine transcripts, were quantified using quantitative real-time polymerase chain reaction (qRT-PCR). Viral gRNA or sgRNA CT values from mock-infected tissues (0 dpi) exceeded 35. Transcript levels were normalized to GAPDH and presented as 2^-ΔCT. Statistical significance was assessed using ordinary one-way ANOVA, with significance levels indicated as follows: *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. Data are presented as mean ± SEM.

Supplementary Figure. 2

| Mock | 6 dpi | 14 dpi |

50 μm

Supplemental Figure 2. Brain Histopathology after Mock and SARS2-N501Y$_{MA30}$ Infection.
The representative images of brain tissue sections were stained with hematoxylin and eosin (H&E) at indicated days following either mock or SARS-CoV-2 infection. Scale bar: 50 μm

Supplemental Figure 3. Lung Pathology and Immunohistochemistry of Viral Protein. (A) Lung sections from mice infected with SARS2-N501Y$_{MA30}$ (n=5 for mock infection and for 6 dpi; n=4 for 14 dpi) underwent H&E staining. **(B)** Acute lung injury severity was assessed blindly following ATS guidelines (32), with 0 for no injury, 1 for mild to moderate injury, and 2 for severe injury. ****p < 0.0001. Scale bars: 100 μm. Data are shown as mean $\pm$ SEM. **(C)** Lung Tissue Immunohistochemistry revealing SARS-CoV-2 N protein (black arrows). Representative images show the presence of SARS-CoV-2 N protein (indicated by black arrows) at specified days post-infection. Scale bar: 100 μm.

Supplemental Figure 4. Transcriptomic Profile Overview in SARS2-N501Y$_{MA30}$ infected mouse brains. (A) Principal Component Analysis (PCA) of transcriptome profiles comparing mock-infected brains (0dpi) with those at 4 and 14 days post-infection. **(B)** Volcano Plot Analysis: The x-axis denotes log2-transformed fold changes, and y-axis indicates -log10-transformed significant values. (Thresholds set at |log2fold| ≥ 1.5, Q value ≤ 0.05) **(C)** Gene Ontology (GO) Enrichment Analysis of DEGs across groups. **(D)** Heatmap visualization of 140 DEGs identified post-infection. **(E)** Kyoto Encyclopedia of Genes and Genomes (KEGG) Pathway Analysis comparing groups at 4dpi vs 0dpi, 14dpi vs 0dpi.

S1 Table

Gene	Forward Primer	Reverse Primer
2019-nCoV_N1	GACCCCAAAATCAGCGAAAT	TCTGGT TACTGCCAGTTGAATCTG
SARS-CoV-2 sgRNA for E	CGATCTCTTGTAGATCTGTTCTC	ATATTGCAGCAGTACGCACACA
mGAPDH	AACAGCAACTCCCACTCTTC	CCTGTTGCTGTAGCCGTATT
mIFNβ	TCAGAATGAGTGGTGGTTGC	GACCTTTCAAATGCAGTAGATTCA
mIFNγ	CGGCACAGTCATTGAAAGCCTA	GTTGCTGATGGCCTGATTGTC
mIFNλ	AGCTGCAGGTCCAAGAGCG	GGTGGTCAGGGCTGAGTCATT
mTNFα	GAACTGGCAGAAGAGGCACT	AGGGTCTGGGCCATAGAACT
mIL1a	CGCTTGAGTCGGCAAAGAAAT	ACAAACTGATCTGTGCAAGTCTC
mIL-1b	ACTGTTTCTAATGCCTTCCC	ATGGTTTCTTGTGACCCTGA
mIL-6	GAGGATACCACTCCCAACAGACC	AAGTGCATCATCGTTGTTCATACA
mIL-8	CACCTCAAGAACATCCAGAGCT	CAAGCAGAACTGAACTACCATCG
mCCL2	CTTCTGGGCCTGCTGTTCA	CCAGCCTACTCATTGGGATCA
mCCL5	AGATCTCTGCAGCTGCCCTCA	GGAGCACTTGCTGCTGGTGTAG
mCXCL9	GCCATGAAGTCCGCTGTTCT	GGGTTCCTCGAACTCCACACT
mCXCL10	GCCGTCATTTTCTGCCTCAT	GCTTCCCTATGGCCCTCATT