Gene expression comparisons between captive and wild shrew brains reveal captivity effects

Maria Alejandra Bedoya Duque[1,2], William R. Thomas[2*], Dina K. N. Dechmann[3,4], John Nieland[5], Cecilia Baldoni[3,4], Dominik von Elverfeldt[6], Marion Muturi[3,4], Angelique P. Corthals[7] Liliana M. Dávalos [2,8*]

[1]Departamento de Ciencias Biológicas Bioprocesos y Biotecnología, Universidad Icesi, Cali, Valle del Cauca, Colombia
[2]Department of Ecology and Evolution, Stony Brook University, Stony Brook, New York, United States
[3]Max-Planck Institute for Animal Behavior, Radolfzell, Germany
[4]University of Konstanz, Konstanz, Germany
[5]Health Science and Technology, Aalborg University, Aalborg, Denmark
[6]Department of Radiology, University Medical Center Freiburg, Freiburg, Germany
[7]John Jay College of Criminal Justice; New York, United States
[8]Consortium for Inter-Disciplinary Environmental Research, Stony Brook University, Stony Brook, New York, United States

*Correspondence: *William R. Thomas (william.thomas@stonybrook.edu), Liliana M. Dávalos (liliana.davalos@stonybrook.edu).*

ORCIDs
MABD - 0000-0002-6879-9803
WRT - 0009-0002-3858-2440
DKN - 0000-0003-0043-8267
JN - 0000-0001-7423-0122
APC - 0000-0002-5610-2992
LMD - 0000-0002-4327-7697

ABSTRACT

Compared to their free-ranging counterparts, wild animals in captivity experience different conditions with lasting physiological and behavioral effects. Although shifts in gene expression are expected to occur upstream of these phenotypes, we found no previous gene expression comparisons of captive vs. free-ranging mammals. We assessed gene expression profiles of three brain regions (cortex, olfactory bulb, and hippocampus) of wild shrews (*Sorex araneus*) compared to shrews kept in captivity for two months and undertook sample drop-out to examine robustness given limited sample sizes. Consistent with captivity effects, we found hundreds of differentially expressed genes in all three brain regions, 104 overlapping across all three, that enriched pathways associated with neurodegenerative disease, oxidative phosphorylation, and genes encoding ribosomal proteins. In the shrew, transcriptomic changes detected under captivity resemble responses in several human pathologies, including major depressive disorder and neurodegeneration. While interpretations of individual genes are tempered by small sample sizes, we propose captivity influences brain gene expression and function and can confound analyses of natural processes in wild individuals under captive conditions.

Keywords: Captivity, brain, shrew, transcriptomics.

INTRODUCTION

Wild animals placed in captivity are subject to altered environmental conditions in comparison to their free-ranging counterparts. Shelter, movement space, diet, climate, lighting, social factors, and human presence vary drastically between the wild and captivity [1–4]. Organisms respond to these changes by altering behavior, physiology, biochemistry and gene expression, but these responses occur at different time scales [5]. While long-term habituation to captivity can activate multiple stress responses, even short-term habituation to captive conditions affects brain morphology and function [1,3,6–8]. Symptoms of chronic stress appear because of frequent exposure to perceived threats [2] that with reduced environmental stimulation can lead to depressive-like states in captive animals [9]. Thus, despite the perception of a comfortable, predictable, and controlled environment, captive conditions can generate detrimental responses in captive individuals.

Short-term and long-term environmental changes also lead to lasting changes in gene expression, with a central role in cellular adaptation through extensive transcriptional and post-transcriptional regulation [10–13]. Although studies in mammals have addressed the effects of captivity in many phenotypes downstream of transcription, including morphology [14,15], longevity [16] and stress [17,18], to our knowledge, there are no experiments comparing brain gene expression in non-domesticated wild and captive mammals. This is of particular interest to modern research, as many experiments are performed on mammals in laboratory settings and may yield different results from those obtained in the wild [6,19], potentially confounding the effects of captivity with phenotypes of interest.

To explore potential changes in the brain from captivity, we assessed the brain gene expression profiles of wild common shrews (*Sorex araneus*) compared to same-age shrews held captive for two months (>10% of their lifespan). The common shrew is ideal for revealing potential captivity effects on gene expression, as many research resources are available: a complete genome assembly and a brain MRI-atlas with region-specific gene expression profiles [20]. *S. araneus* is also relevant to neuroscience for its seasonal change in brain size [21]. We focused on three brain regions that may shift functional importance in an environment that does not require exploring complex surroundings: 1) the cortex,

whose evolution in mammals has improved information processing and decision making, 2) the
hippocampus, involved in spatial memory and learning, and 3) the olfactory bulb, which receives
olfactory information from the environment. We outline differences in brain gene expression between
wild and captive shrews detected using differential gene expression analysis of mRNA. Our results
support a captivity effect in the shrew brain, and we highlight similar responses found in studies of
metabolism, neurodegenerative diseases, stress, and depression.

MATERIALS AND METHODS

Shrew Collection

Shrews were caught in Liggeringen, Germany, as previously described [22] to reduce stress. Our study
had small sample sizes, as only ten individuals from the population could be sampled in autumn and
logistical barriers from the COVID-19 pandemic restricted sampling over multiple years. Captive juvenile
shrews were caught in September of 2020 and housed individually for two months at the Max Planck
Institute of Animal Behavior before samples were taken (Table 1). Animals were allowed to move freely
between two connected cages, one containing a running wheel, bowls of food and water, a nesting pot,
and the other consisting of several layers of soil and hay (detailed housing in [23]). Captive shrews were
kept at ambient outdoor temperatures, ranging from -5°C to 22°C, and provided a daily diet of 5g of raw
meat and 2g of mealworm. Only four autumn wild juvenile shrews (two female and two male) were
caught in November, and were sampled the same day of capture. Although sex is used as a covariate in
subsequent analyses, there were two female wild individuals, but no captive females. This confounded
sex and captivity in experimental design. Season was used as a proxy for age, as *S. araneus* have an
unusually short lifespan of ~13 months, with one reproductive season followed by senescence. Overlap
between generations occurs in summer and early fall, when adults have worn teeth, disheveled fur, and
developed gonads. Adults were excluded from the experiment.

Tissue Sampling, RNA Extraction, and Sequencing

Shrews were weighed, then anesthetized and euthanized under deep anesthesia by perfusion of the
vascular system with PAXgene Tissue Fixative (perfusion described in Lázaro et al., 2018 [22]). The brain
was extracted immediately, weighed, dissected into individual regions, and submerged in cold fixative.
Three brain regions were compared between captive and wild individuals: cortex, olfactory bulb, and
hippocampus. Brain regions were held in fixative, and within 24 hours were switched to PAXgene Tissue
Stabilizer and frozen in liquid nitrogen. Skull toothrow length was measured to assess the size of
individuals. RNA was extracted from each brain region using a modified Qiagen Micro RNAeasy protocol
designed to retrieve RNA from small sensory tissues [24]. Quality control, library preparation, and
sequencing were conducted by Genewiz. RNA integrity numbers (RIN) were measured with RNA
ScreenTape and samples below a RIN of 6 were removed from the experiment. mRNAs were selected by
polyA enrichment followed by paired-end sequencing with Illumina HiSeq.

Bioinformatics

Samples with low sequencing depths (<20 million reads) were removed from the experiment. Adapters
were trimmed and low-quality reads removed with fastp 0.20.1 [25]. Reads were mapped by pseudo-
alignment to the *S. araneus* reference transcriptome (GCF_000181275.1_SorAra2.0_rna) using Kallisto
0.46.1 [26]. Samples with mapping percentage under 45% were removed. Read counts were then

normalized for library size and content using the median of ratios from DESeq2 1.36.0 [27]. To visualize
the overall effect, we ran a principal component analysis (PCA) of the 500 most varying genes for each
brain region using variance stabilized read counts [28].

Differential gene expression between captive and wild individuals for each brain region was analyzed in
R (version 4.2.0) with DESeq2 [27]. Sex and body size were used as additional covariates for the analysis
(~sex+toothrow+condition). Maxillary tooth row length was used as a proxy for body size, as it does not
change seasonally within individuals [29,30]. Statistical significance for each region comparison was
determined by conducting a modified Wald test, using an alternative null of 20% change, and then
adjusting p-values for multiple testing using Benjamini and Hochberg correction [31] ($p_{adj} < 0.05$). The
proportion of true nulls was analyzed with limma [32]. Given a statistical significance threshold of 0.05,
the expected proportion of true nulls approaches 0.95. Higher values indicate false negatives, and lower
values indicate false positives. Genes identified as differentially expressed for each brain region were
separated by direction (upregulated or downregulated). KEGG pathway enrichment analysis (Modified
Fisher's Exact test) was carried out for both directional gene lists of each brain region using DAVID
[33,34]. To examine result robustness, we conducted a drop-out analysis for each brain region
(Supplemental Figures 1-3). We removed one individual per analysis, identified overlapping differentially
expressed genes across all analyses, reran DAVID pathway enrichments, and identified overlapping
pathways.

RESULTS

We identified divergence in the expression profiles of all three brain regions associated with a potential
captivity effect. For each brain region, PCA produced two principal component axes that accounted for
most of the variation from these genes (Figure 1). The first principal component separated wild from
captive individuals, accounting for 42% of the variation in gene expression in the hippocampus and 47%
in the olfactory bulb. Percent variation expressed was even greater in the cortex, where the first
principal component accounted for 63% of the variation. As for PC2, each region highlighted one sample
within each analysis with variation not found in other samples. We determined this was not a single
outlier sample, as we found each case corresponded to different individuals for each brain region
(Supplemental Figure 4), with no discernible pattern across regions.

Differential expression analysis confirmed these patterns (Figure 1). Of the 19,296 genes analyzed, 603
genes were differentially expressed (DEGs; $p_{adj}<0.05$) in at least one brain region (Figure 2). In the
cortex, we found 434 DEGs, 162 upregulated and 272 downregulated in captive shrews. The olfactory
bulb had 298 DEGs, with 185 upregulated and 112 downregulated in captive animals. In the
hippocampus, we identified 180 DEGs, 48 upregulated and 132 downregulated in captive animals. The
proportion of true nulls was high, 0.97 in the cortex, 1.00 in the hippocampus, and 0.99 in the olfactory
bulb. This indicates lower statistical power to detect many true positives in the hippocampus and
olfactory bulb because of small samples sizes and high variability [35,36]. We compared lists of DEGs
between brain regions and found 104 genes that were differentially expressed in the same direction
across all regions, with another 101 overlapping in two of three brain regions (Figure 2).

To partially overcome small sample sizes, we undertook pathway enrichment analyses, which revealed
captive gene expression differences are linked to ribosomal and mitochondrial function, with
associations to neurodegenerative processes (Figure 3). We identified 21 pathways significantly

enriched (p-value < 0.05, modified Fisher's Exact Test) for the cortex, 21 for the olfactory bulb and 21 for
the hippocampus. Out of these, 14 pathways were significantly enriched (p<0.05) across the three brain
regions (Figure 3), with the largest enrichment in the ribosome pathway (p_{cor}: 7.4×10^{-32}, n= 35genes; p_{olf}:
4.0×10^{-11}, n=13 genes; p_{hip}: 5.0×10^{-15}, n=17 genes). Within enriched pathways, we identified expression
changes that likely affect mitochondrial function in the brain. Nineteen genes encoding mitochondrial
complex I NADH:ubiquinone oxidoreductase subunit proteins (largest enzyme of the respiratory chain),
of which 8 overlap all three regions and were found across 12 of the 14 overlapping enriched pathways
(Figure 3). Six of the pathways containing complex I proteins were associated with neurodegenerative
diseases, including Parkinson's disease, amyotrophic lateral sclerosis, pathways of neurodegeneration-
multiple diseases, prion disease, Huntington's disease, and Alzheimer's disease. Out of the eight
enriched pathways remaining, six of them were associated with changes in metabolism, such as
oxidative phosphorylation, retrograde endocannabinoid signaling, non-alcoholic fatty liver disease,
diabetic cardiomyopathy, chemical carcinogenesis - ROS, and thermogenesis, highlighting the influence
of mitochondrial genes on both metabolically related and neurodegenerative mechanisms.

While principal component analyses show little impact from sex, drop-out analyses indicate robust gene
expression results in the cortex (Supplemental Figures 5 and 1-3). First, we examined the expression of
each individual mitochondrial gene of interest for each brain region (Supplemental Figures 6A-B).
Normalized counts did show some dependance on sex. But instead of driving differences in gene
expression, sex differences appear to dampen overall signal, and differences are greater between wild
and captive samples than between sexes. Pathway enrichment for drop-out analyses was robust for the
cortex, where 109 significant genes from the full experimental analysis remained differentially
expressed across all dropouts (p_{adj}<0.05). This subset of robust genes enriches most of the proposed
pathways, including oxidative phosphorylation, neurodegeneration, Parkinson's disease, Alzheimer's
disease, prion disease, Huntington's disease, and ALS (p<0.05). The olfactory bulb and hippocampus
were not robust to sample drop-out, as only 27 genes for the hippocampus and 15 genes for the
olfactory bulb remained across all drop-out analyses and did not enrich highlighted pathways.

DISCUSSION

Both genomic [37,38] and environmental variation [39–41] impact neural structures and function. Past
studies have focused on fixed domestication-associated neural changes in subsets of mammals [42–44],
and identified differences in gene expression and transcription factors involved in domestication-
associated changes in both brain and craniofacial phenotypes [37,38]. While brain gene expression
responses to captive environment change have been documented in diverse organisms [45–48], this is
the first study documenting potential captivity effects specifically on mammalian brain transcripts in the
absence of artificial selection. Although captivity is routine in many experiments, effects on expression
may parallel or counter responses to domestication or other forms of induced or natural environmental
change. Potentially large expression changes in the brain after a brief stint in captivity highlight the
importance of working with both wild and captive individuals to avoid confounding gene expression
changes from phenotypes of interest with those associated with captivity.

We found potential region-specific responses to captivity, but small sample sizes mean these results are
limited by low statistical power and should be interpreted with caution. Principal components separate
samples by captivity first (Figure 1) so that overall expression supports a captivity effect. Individual
DEGs, however, are not robust to sample drop-out for the underpowered hippocampus and olfactory

bulb (Supplemental Figure 2 and 3). For example, in the hippocampus we found apparent upregulation
of four myosin genes (*MYH6*, *MYL2*, *MYL3*, *MYBPC3*) (Figure 1). This may relate to changes in synaptic
plasticity, as myosins regulate actin cytoskeleton dynamics in dendrites essential for long-term
potentiation and memory formation in rodents [49–52]. But based on power analyses from vertebrate
brain expression [36], when sampling four individuals, there is ~70% power to accurately detect
differential expression in genes with more than 40 reads. While most observed DEGs are above this
threshold (Supplemental Figure 7), nearly half of the total genes sequenced had fewer than 40 reads,
reducing the power to detect differential expression (~5% to 45%), and therefore the reliability for each
individual differentially expressed gene.

Instead, we focus biological interpretations on pathway changes found across all three brain regions
(Figure 3), as gene sets reduce noise and are more stable compared to individual gene expression
changes [53]. We found downregulation of ribosomal protein genes in captive shrews. Ribosomes
compose the cell's protein synthesis machinery, and their own synthesis, ribosomal biogenesis, is a
highly conserved process [54,55]. In the captive shrew brain, downregulation of 12 ribosomal proteins
(RPs) suggests impaired regulation of ribosomal biogenesis and subsequent functions. By determining
translational capacity of proliferating cells [56], ribosomal biogenesis constrains animal growth and
development [57,58]. Dysregulation of ribosomal biogenesis is therefore implicated in numerous
phenotypes including neurite survival and synaptic function associated with human neurodegeneration
[54]. These effects seem to be mirrored in captive shrews, as we found downregulated gene enrichment
for pathways associated with neurodegeneration. Neurodegeneration has been linked to changes in
stress [59,60], but these transcriptomic profiles and potential shifts in processes such as reduced
synaptic efficiency might be occurring because of captivity and not disease.

Differential expression of ribosomal protein-coding genes in the brains of captive shrews could influence
behavior, as many such genes have been previously highlighted for their regulatory changes under
domestication and social defeat stress [44,61]. For example, comparisons between domestic rabbits and
wild counterparts identified 612 DEGs across four brain regions that also enriched ribosomal pathways
[44]. Additionally, changes in expression of ribosomal genes have also been associated with decreased
proliferation in the hippocampus of mice under chronic social defeat stress, which consists of repeated
negative social experiences (defeats) produced by daily agonistic interactions with an aggressive
partner, leading to states of anxiety and depression [61]. Captivity-associated differential expression of
ribosomal proteins in the shrew brain may thus indicate a depressed neural state, especially as past
studies have argued that animals in captivity often experience repeated stressors used to induce
depressive-like states [9].

Captivity conditions such as altered diet and physical activity may also influence shrew energy
metabolism in the brain. In captive shrews, we found downregulation of mitochondrial complex I
proteins, essential for energy production, as they comprise the first and largest complex of the electron
transport chain [62,63]. Impairments to complex I function are linked to pathophysiological states,
genetic diseases [62,64–66], and several neurodegenerative disorders, leading to decline of energy
production and increased production of reactive oxygen species [67–70]. Downregulation of 8 genes
encoding for mitochondrial complex I subunit proteins in three shrew brain regions suggests a
dysfunction of the oxidative phosphorylation (OXPHOS) system and dysregulation of energy metabolism
in captivity. These genes are also downregulated in the brains of patients with autism [71], and
neurodegenerative diseases such as Huntington's [72] and Alzheimer's [73]. Indeed, the effects of

certain conditions found in captivity, such as reduced stimulation, could also modify OXPHOS function in
association with neurodegeneration, as normal brains exposed to chronic sensory deprivation exhibit
downregulation of OXPHOS subunits as found in the brain of Alzheimer's patients [74]. However, studies
have found differences in mitochondrial function/gene expression between sexes [75–77], and sex-
dependent susceptibility to neurodegenerative disease [76]. Therefore, including only two female
samples is a limitation of our study to be improved in the future.

As with differential ribosomal protein expression, mitochondrial dysfunction and changes in expression
of complex I subunits have also been linked to depressed activity and stress-related behaviors. For
example, female mice bred for low activity showed upregulation of complex I subunit genes [78],
including three genes downregulated in the three brain regions tested in captive shrews (*NDUFA2*,
NDUFA4, and *NDUFS5*). These genes exhibit opposite expression patterns between low activity mice and
captive shrews. But other studies have found the direction of changes in mitochondrial gene expression
related to stress and anxiety may depend on stressor duration, genetic background, and strain of the
model organism studied [79]. Nevertheless, published and newly detected shifts still point to
mitochondrial pathway dysregulation as a major brain stress response [79]. Differential expression of
complex I subunit genes in captive shrews is consistent with a transcriptional phenotype associated with
depression and stress in model organisms. While the validation of specific mechanisms for such a
transcriptional phenotype (e.g., by western blot validation) is outside the scope of this paper, we have
identified key potential targets for future analyses.

Conclusion

Despite attempts to limit differential stressors and optimize holding conditions, these results indicate
transcriptomic changes in the brain of shrews after being placed in a captive environment. Such changes
could influence phenotype through shifts in energy metabolism, synaptic function, cell survival, and
behavior. Across pathways, captivity-associated changes in the shrew resemble responses in several
human diseases, notably, neurodegeneration. These are unlikely to be shrew-specific, as similar
expression shifts have been found in other organisms with disturbance in their natural environments.
When investigating the molecular mechanisms of disease using model organisms, captivity can confound
analyses of natural physiological processes. There are benefits to using captive organisms in disease
research, but our results indicate future studies should use natural environments to illuminate brain
function and responses to environmental change. These experiments could involve collaborative efforts
with ecologists and evolutionary biologists [80] to study natural models of brain change [21,81] that can
be validated in laboratory settings using cell lines. Alternatively, findings from laboratory-based disease
models can be paired with experimental groups in exclosures as close to wild as possible to assess the
impact of environmental factors on experimental outcomes.

ACKNOWLEDGMENTS
279 Images were created with a licensed version of BioRender.com.

GRANTS
281 Human Frontiers of Science Program, award: RGP0013/2019 (to Dina K. N. Dechmann, John Nieland,
282 Liliana M. Dávalos).

The Hearst Foundation, APS Hearst Summer Undergraduate Research Fellowship (to Maria Alejandra Bedoya Duque).

Stony Brook University, Presidential Innovation and Excellence award (to Liliana M. Dávalos, William R. Thomas was supported in part by award).

REFERENCES

[1]   Fischer CP, Wright-Lichter J, Romero LM. Chronic stress and the introduction to captivity: How wild house sparrows (*Passer domesticus*) adjust to laboratory conditions. Gen Comp Endocrinol 2018;259:85–92. https://doi.org/10.1016/j.ygcen.2017.11.007.

[2]   Fischer CP, Romero LM. Chronic captivity stress in wild animals is highly species-specific. Conserv Physiol 2018;7. https://doi.org/10.1093/conphys/coz093.

[3]   Love AC, Lovern MB, DuRant SE. Captivity influences immune responses, stress endocrinology, and organ size in house sparrows (*Passer domesticus*). Gen Comp Endocrinol 2017;252:18–26. https://doi.org/10.1016/j.ygcen.2017.07.014.

[4]   Seeber PA, Morrison T, Ortega A, East ML, Greenwood AD, Czirják GÁ. Immune differences in captive and free-ranging zebras (*Equus zebra* and *E. quagga*). Mamm Biol 2020;100:155–64. https://doi.org/10.1007/s42991-020-00006-0.

[5]   Schulte PM. Changes in gene expression as biochemical adaptations to environmental change: A tribute to Peter Hochachka. Comp Biochem Physiol 2004;139:519–29. https://doi.org/10.1016/j.cbpc.2004.06.001.

[6]   Calisi RM, Bentley GE. Lab and field experiments: Are they the same animal? Horm Behav 2009;56:1–10. https://doi.org/10.1016/j.yhbeh.2009.02.010.

[7]   Dickens MJ, Earle KA, Romero LM. Initial transference of wild birds to captivity alters stress physiology. Gen Comp Endocrinol 2009;160:76–83. https://doi.org/10.1016/j.ygcen.2008.10.023.

[8]   Cucchi T, Neaux D, Féral L, Goussard F, Adriensen H, Elleboudt F, et al. How domestication, feralization and experience-dependent plasticity affect brain size variation in *Sus scrofa*. R Soc Open Sci 2024;11. https://doi.org/10.1098/RSOS.240951.

[9]   Lecorps B, Weary DM, von Keyserlingk MAG. Captivity-Induced Depression in Animals. Trends Cogn Sci 2021;25:539–41. https://doi.org/10.1016/J.TICS.2021.03.010.

[10]   López-Maury L, Marguerat S, Bähler J. Tuning gene expression to changing environments: From rapid responses to evolutionary adaptation. Nat Rev Genet 2008;9:583–93. https://doi.org/10.1038/nrg2398.

[11]   Morin M, Jönsson M, Wang CK, Craik DJ, Degnan SM, Degnan BM. Captivity induces a sweeping and sustained genomic response in a starfish. Mol Ecol 2023;32:3541–56. https://doi.org/10.1111/MEC.16947.

317   [12]   Zhang P, Chen JS, Li QY, Sheng LX, Gao YX, Lu BZ, et al. Neuroprotectants attenuate hypobaric
318   hypoxia-induced brain injuries in cynomolgus monkeys. Zool Res 2020;41:3.
319   https://doi.org/10.24272/J.ISSN.2095-8137.2020.012.

320   [13]   Yang M, Huang Y, Wu H, Li C, Ling S, Sun J, et al. Blood transcriptome analysis revealed the
321   immune changes and immunological adaptation of wildness training giant pandas. Molecular
322   Genetics and Genomics 2022;297:227–39. https://doi.org/10.1007/S00438-021-01841-
323   7/METRICS.

324   [14]   Hartstone-Rose A, Selvey H, Villari JR, Atwell M, Schmidt T. The Three-Dimensional Morphological
325   Effects of Captivity. PLoS One 2014;9:e113437. https://doi.org/10.1371/journal.pone.0113437.

326   [15]   Zack EH, Smith SM, Angielczyk KD. Effect of captivity on the vertebral bone microstructure of
327   xenarthran mammals. Anat Rec 2022;305:1611–28. https://doi.org/10.1002/ar.24817.

328   [16]   Tidière M, Gaillard JM, Berger V, Müller DWH, Lackey LB, Gimenez O, et al. Comparative analyses
329   of longevity and senescence reveal variable survival benefits of living in zoos across mammals. Sci
330   Rep 2016;6:1–7. https://doi.org/10.1038/srep36361.

331   [17]   Marino L, Rose NA, Visser IN, Rally H, Ferdowsian H, Slootsky V. The harmful effects of captivity
332   and chronic stress on the well-being of orcas (Orcinus orca). J Vet Behav 2020;35:69–82.
333   https://doi.org/10.1016/j.jveb.2019.05.005.

334   [18]   Quispe R, Villavicencio CP, Addis E, Wingfield JC, Vasquez RA. Seasonal variations of basal cortisol
335   and high stress response to captivity in *Octodon degus*, a mammalian model species. Gen Comp
336   Endocrinol 2014;197:65–72. https://doi.org/10.1016/j.ygcen.2013.12.007.

337   [19]   Stryjek R, Parsons MH, Fendt M, Święcicki J, Bębas P. Let's get wild: A review of free-ranging rat
338   assays as context-enriched supplements to traditional laboratory models. J Neurosci Methods
339   2021;362:109303. https://doi.org/10.1016/j.jneumeth.2021.109303.

340   [20]   Baldoni C, Thomas WR, von Elverfeldt D, Reisert M, Làzaro J, Muturi M, et al. Histological and MRI
341   brain atlas of the common shrew, *Sorex araneus*, with brain region-specific gene expression
342   profiles. Front Neuroanat 2023;17. https://doi.org/10.3389/fnana.2023.1168523.

343   [21]   Lázaro J, Hertel M, Sherwood CC, Muturi M, Dechmann DKN. Profound seasonal changes in brain
344   size and architecture in the common shrew. Brain Struct Funct 2018;223:2823–40.
345   https://doi.org/10.1007/s00429-018-1666-5.

346   [22]   Lázaro J, Hertel M, LaPoint S, Wikelski M, Stiehler M, Dechmann DKN. Cognitive skills of common
347   shrews (*Sorex araneus*) vary with seasonal changes in skull size and brain mass. J Exp Biol
348   2018;221. https://doi.org/10.1242/jeb.166595.

349   [23]   Lázaro J, Hertel M, Muturi M, Dechmann DKN. Seasonal reversible size changes in the braincase
350   and mass of common shrews are flexibly modified by environmental conditions. Sci Rep 2019;9.
351   https://doi.org/10.1038/s41598-019-38884-1.

352   [24]   Yohe LR, Davies KTJ, Rossiter SJ, Dávalos LM. Expressed Vomeronasal Type-1 Receptors (V1rs) in
353   Bats Uncover Conserved Sequences Underlying Social Chemical Signaling. Genome Biol Evol
354   2019;11:2741. https://doi.org/10.1093/gbe/evz179.

[25]   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics
2018;34:i884. https://doi.org/10.1093/bioinformatics/bty560.

[26]   Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. Nat
Biotechnol 2016;34:525–7. https://doi.org/10.1038/nbt.3519.

[27]   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. Genome Biol 2014;15:1–21. https://doi.org/10.1186/s13059-014-0550-8.

[28]   Love MI, Anders S, Huber W. Analyzing RNA-seq data with DESeq2 2023.
https://bioconductor.org/packages/devel/bioc/vignettes/DESeq2/inst/doc/DESeq2.html#varianc
e-stabilizing-transformation (accessed February 5, 2023).

[29]   Keicher L, Teague O'mara M, Voigt CC, Dechmann DKN. Stable carbon isotopes in breath reveal
fast metabolic incorporation rates & seasonally variable but rapid fat turnover in the common
shrew (Sorex araneus). J Exp Biol 2017;220:2834–41. https://doi.org/10.1242/jeb.159947.

[30]   Lázaro J, Dechmann DKN, LaPoint S, Wikelski M, Hertel M. Profound reversible seasonal changes
of individual skull size in a mammal. Curr Biol 2017;27:R1106–7.
https://doi.org/10.1016/j.cub.2017.08.055.

[31]   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach
to Multiple Testing. J R Stat Soc Series B Stat Methodol 1995;57:289–300.
https://doi.org/10.1111/J.2517-6161.1995.TB02031.X.

[32]   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. Limma powers differential expression
analyses for RNA-sequencing and microarray studies. Nucleic Acids Res 2015;43:e47.
https://doi.org/10.1093/nar/gkv007.

[33]   Sherman BT, Hao M, Qiu J, Jiao X, Baseler MW, Lane HC, et al. DAVID: a web server for functional
enrichment analysis and functional annotation of gene lists (2021 update). Nucleic Acids Res
2022;50:W216–21. https://doi.org/10.1093/nar/gkac194.

[34]   Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using
DAVID bioinformatics resources. Nat Protoc 2009;4:44–57.
https://doi.org/10.1038/nprot.2008.211.

[35]   Jiang H, Doerge RW. Estimating the Proportion of True Null Hypotheses for Multiple
Comparisons. Cancer Inform 2008:25–32.

[36]   Todd E V., Black MA, Gemmell NJ. The power and promise of RNA-seq in ecology and evolution.
Mol Ecol 2016;25:1224–41. https://doi.org/10.1111/mec.13526.

[37]   Benítez-Burraco A, Uriagereka J, Nataf S. The genomic landscape of mammal domestication
might be orchestrated by selected transcription factors regulating brain and craniofacial
development. Development Genes and Evolution 2023 233:2 2023;233:123–35.
https://doi.org/10.1007/S00427-023-00709-7.

[38] Lu C, Li M, Sun X, Li N, Wang W, Tong P, et al. Comparing the hippocampal miRNA expression
profiles of wild and domesticated Chinese tree shrews (*Tupaia belangeri chinensis*). BMC Ecol
Evol 2021;21. https://doi.org/10.1186/S12862-020-01740-2.

[39] O' Regan HJ, Kitchener AC. The effects of captivity on the morphology of captive, domesticated
and feral mammals. vol. 35. 2005. https://doi.org/10.1111/j.1365-2907.2005.00070.x.

[40] Reyes AS, Bittar A, Ávila LC, Botia C, Esmeral NP, Bloch NI. Divergence in brain size and brain
region volumes across wild guppy populations. Proceedings of the Royal Society B 2022;289.
https://doi.org/10.1098/RSPB.2021.2784.

[41] Frahm H, Stephan H. Vergleichende Volumenmessungen an Hirnen von Wild- und
Gefangenschaftstieren des Spitzhornchens (*Tupaia*). Journal fuer Hirnforschung. 1976;17:449–
462.

[42] Albert FW, Somel M, Carneiro M, Aximu-Petri A, Halbwax M, Thalmann O, et al. A Comparison of
Brain Gene Expression Levels in Domesticated and Wild Animals. PLoS Genet 2012;8.
https://doi.org/10.1371/journal.pgen.1002962.

[43] Chadaeva I, Ponomarenko P, Kozhemyakina R, Suslov V, Bogomolov A, Klimova N, et al.
Domestication Explains Two-Thirds of Differential-Gene-Expression Variance between Domestic
and Wild Animals; The Remaining One-Third Reflects Intraspecific and Interspecific Variation.
Animals 2021;11:2667. https://doi.org/10.3390/ani11092667.

[44] Sato DX, Rafati N, Ring H, Younis S, Feng C, Blanco-Aguiar JA, et al. Brain Transcriptomics of Wild
and Domestic Rabbits Suggests That Changes in Dopamine Signaling and Ciliary Function
Contributed to Evolution of Tameness. Genome Biol Evol 2020;12:1918–28.
https://doi.org/10.1093/gbe/evaa158.

[45] DuRant S, Love AC, Belin B, Tamayo-Sanchez D, Santos Pacheco M, Dickens MJ, et al. Captivity
alters neuroendocrine regulators of stress and reproduction in the hypothalamus in response to
acute stress. Gen Comp Endocrinol 2020;295. https://doi.org/10.1016/j.ygcen.2020.113519.

[46] Yagound B, West AJ, Richardson MF, Gruber J, Reid JG, Whiting MJ, et al. Captivity induces large
and population-dependent brain transcriptomic changes in wild-caught cane toads (*Rhinella*
marina). Mol Ecol 2022:4949–61. https://doi.org/10.1111/mec.16633.

[47] Roznere I, Sinn BT, Daly M, Watters GT. Freshwater mussels (Unionidae) brought into captivity
exhibit up-regulation of genes involved in stress and energy metabolism. Sci Rep 2021;11:1–10.
https://doi.org/10.1038/s41598-021-81856-7.

[48] Maas AE, Lawson GL, Bergan AJ, Tarrant AM. Exposure to CO_2 influences metabolism,
calcification and gene expression of the thecosome pteropod *Limacina retroversa*. J Exp Biol
2018;221. https://doi.org/10.1242/jeb.164400.

[49] Kneussel M, Wagner W. Myosin motors at neuronal synapses: Drivers of membrane transport
and actin dynamics. Nat Rev Neurosci 2013;14:233–47. https://doi.org/10.1038/nrn3445.

[50] Ma X, Kawamoto S, Uribe J, Adelstein RS. Function of the Neuron-specific Alternatively Spliced
Isoforms of Nonmuscle Myosin II-B during Mouse Brain Development. Mol Biol Cell
2006;17:2138–49. https://doi.org/10.1091/mbc.E05-10.

[51] Miyazaki T, Watanabe M, Yamagishi A, Takahashi M. B2 exon splicing of nonmuscle myosin heavy
chain IIB is differently regulated in developing and adult rat brain. Neurosci Res 2000;37.
https://doi.org/https://doi.org/10.1016/S0168-0102(00)00130-9.

[52] Rex CS, Gavin CF, Rubio MD, Kramar EA, Chen LY, Jia Y, et al. Myosin IIb Regulates actin dynamics
during synaptic plasticity and memory formation. Neuron 2010;67:603–17.
https://doi.org/10.1016/j.neuron.2010.07.016.

[53] Abraham G, Kowalczyk A, Loi S, Haviv I, Zobel J. Prediction of breast cancer prognosis using gene
set statistics provides signature stability and biological context. BMC Bioinformatics 2010;11:277.
https://doi.org/https://doi.org/10.1186/1471-2105-11-277.

[54] Hetman M, Slomnicki LP. Ribosomal biogenesis as an emerging target of neurodevelopmental
pathologies. J Neurochem 2019;148:325. https://doi.org/10.1111/jnc.14576.

[55] Kumar V. Introduction to ribosome factory, origin, and evolution of translation. In: Emerging
Concepts in Ribosome Structure, Biogenesis, and Function. Cambridge: Academic Press; 2021. p.
1–13.

[56] Chaillou T, Kirby TJ, Mccarthy JJ. Ribosome biogenesis: emerging evidence for a central role in the
regulation of skeletal muscle mass. J Cell Physiol 2014;229:1584.
https://doi.org/10.1002/jcp.24604.

[57] Donati G, Montanaro L, Derenzini M. Ribosome biogenesis and control of cell proliferation: p53 is
not alone. Cancer Res 2012;72:1602–7. https://doi.org/10.1158/0008-5472.can-11-3992.

[58] 58. Kumar V. rDNA gene structure, transcription, and its coregulation. In: Emerging Concepts in
Ribosome Structure, Biogenesis, and Function. Cambridge: Academic Press; 2021. p. 33–45.

[59] Picard M, McEwen BS. Psychological Stress and Mitochondria: A Systematic Review. Psychosom
Med 2018;80:141. https://doi.org/10.1097/psy.0000000000000545.

[60] Trabjerg MS, Mørkholt AS, Lichota J, Oklinski MKE, Andersen DC, Jønsson K, et al. Dysregulation
of metabolic pathways by carnitine palmitoyl-transferase 1 plays a key role in central nervous
system disorders: experimental evidence based on animal models. Sci Rep 2020;10:1–19.
https://doi.org/10.1038/S41598-020-72638-8.

[61] Smagin DA, Kovalenko IL, Galyamina AG, Bragin AO, Orlov YL, Kudryavtseva NN. Dysfunction in
Ribosomal Gene Expression in the Hypothalamus and Hippocampus following Chronic Social
Defeat Stress in Male Mice as Revealed by RNA-Seq. Neural Plast 2016;2016.
https://doi.org/10.1155/2016/3289187.

[62] Ansari F, Yoval-Sánchez B, Niatsetskaya Z, Sosunov S, Stepanova A, Garcia C, et al. Quantification
of NADH:ubiquinone oxidoreductase (complex I) content in biological samples. J Biol Chem
2021;297:297–301. https://doi.org/10.1016/j.jbc.2021.101204.

[63] Stroud DA, Surgenor EE, Formosa LE, Reljic B, Frazier AE, Dibley MG, et al. Accessory subunits are
integral for assembly and function of human mitochondrial complex I. Nature 2016 538:7623
2016;538:123–6. https://doi.org/10.1038/nature19754.

[64] Koopman WJH, Verkaart S, Visch HJ, Van Emst-De Vries S, Nijtmans LGJ, Smeitink JAM, et al.
Human NADH:ubiquinone oxidoreductase deficiency: Radical changes in mitochondrial
morphology? Am J Physiol Cell Physiol 2007;293:22–9.
https://doi.org/10.1152/ajpcell.00194.2006.

[65] Raffa S, Chin XLD, Stanzione R, Forte M, Bianchi F, Cotugno M, et al. The reduction of NDUFC2
expression is associated with mitochondrial impairment in circulating mononuclear cells of
patients with acute coronary syndrome. Int J Cardiol 2019;286:127–33.
https://doi.org/10.1016/j.ijcard.2019.02.027.

[66] Torraco A, Bianchi M, Verrigni D, Gelmetti V, Riley L, Niceta M, et al. A novel mutation in
NDUFB11 unveils a new clinical phenotype associated with lactic acidosis and sideroblastic
anemia. Clin Genet 2017;91:441–7. https://doi.org/10.1111/cge.12790.

[67] Fato R, Bergamini C, Bortolus M, Maniero AL, Leoni S, Ohnishi T, et al. Differential effects of
mitochondrial Complex I inhibitors on production of reactive oxygen species. Biochimica et
Biophysica Acta (BBA) - Bioenergetics 2009;1787:384–92.
https://doi.org/10.1016/j.bbabio.2008.11.003.

[68] Giachin G, Bouverot R, Acajjaoui S, Pantalone S, Soler-López M. Dynamics of Human
Mitochondrial Complex I Assembly: Implications for Neurodegenerative Diseases. Front Mol
Biosci 2016;3:43. https://doi.org/10.3389/fmolb.2016.00043.

[69] Hoefs SJG, Dieteren CEJ, Distelmaier F, Janssen RJRJ, Epplen A, Swarts HGP, et al. NDUFA2
Complex I Mutation Leads to Leigh Disease. Am J Hum Genet 2008;82:1306–15.
https://doi.org/10.1016/j.ajhg.2008.05.007.

[70] McElroy GS, Chakrabarty RP, D'Alessandro KB, Vasan K, Tan J, Stoolman JS, et al. Reduced
expression of mitochondrial complex I subunit Ndufs2 does not impact healthspan in mice. Sci
Rep 2022;12. https://doi.org/10.1038/S41598-022-09074-3.

[71] Anitha A, Nakamura K, Thanseem I, Matsuzaki H, Miyachi T, Tsujii M, et al. Downregulation of the
Expression of Mitochondrial Electron Transport Complex Genes in Autism Brains. Brain Pathol
2013;23:294. https://doi.org/10.1111/bpa.12002.

[72] Weydt P, Pineda V V., Torrence AE, Libby RT, Satterfield TF, Lazarowski ERR, et al.
Thermoregulatory and metabolic defects in Huntington's disease transgenic mice implicate PGC-
1α in Huntington's disease neurodegeneration. Cell Metab 2006;4:349–62.
https://doi.org/10.1016/j.cmet.2006.10.004.

[73] Liang WS, Reiman EM, Valla J, Dunckley T, Beach TG, Grover A, et al. Alzheimer's disease is
associated with reduced expression of energy metabolism genes in posterior cingulate neurons.
PNAS 2008;18:4441–6. https://doi.org/10.1073pnas.0709259105.

[74] Chandrasekaran K, Hatanpää K, Brady DR, Rapoport SI. Evidence for Physiological Down-
regulation of Brain Oxidative Phosphorylation in Alzheimer's Disease. Exp Neurol 1996;142:80–8.
https://doi.org/10.1006/exnr.1996.0180.

[75] Ventura-Clapier R, Moulin M, Piquereau J, Lemaire C, Mericskay M, Veksler V, et al.
Mitochondria: a central target for sex differences in pathologies. Clin Sci 2017;131:803–22.
https://doi.org/10.1042/CS20160485.

[76] Gaignard P, Savouroux S, Liere P, Pianos A, Thérond P, Schumacher M, et al. Effect of Sex
Differences on Brain Mitochondrial Function and Its Suppression by Ovariectomy and in Aged
Mice. Endocrinology 2015;156:2893–904. https://doi.org/10.1210/EN.2014-1913.

[77] Junker A, Juster RP, Picard M. Integrating sex and gender in mitochondrial science. Curr Opin
Physiol 2022;26:100536. https://doi.org/10.1016/J.COPHYS.2022.100536.

[78] Booher WC, Vanderlinden LA, Hall LA, Thomas AL, Evans LM, Saba LM, et al. Hippocampal RNA
sequencing in mice selectively bred for high and low activity. Genes Brain Behav 2023;22:e12832.
https://doi.org/10.1111/gbb.12832.

[79] Misiewicz Z, Iurato S, Kulesskaya N, Salminen L, Rodrigues L, Maccarrone G, et al. Multi-omics
analysis identifies mitochondrial pathways associated with anxiety-related behavior. PLoS Genet
2019. https://doi.org/10.1371/journal.pgen.1008358.

[80] Bertile F, Matallana-Surget S, Tholey A, Cristobal S, Armengaud J. Diversifying the concept of
model organisms in the age of -omics. Commun Biol 2023;6. https://doi.org/10.1038/S42003-
023-05458-X.

[81] van Tussenbroek IA, Knörnschild M, Nagy M, Ten Cate CJ, Vernes SC. Morphological Diversity in
the Brains of 12 Neotropical Bat Species. Https://DoiOrg/103161/15081109ACC2023252011
2024;25:323–38. https://doi.org/10.3161/15081109ACC2023.25.2.011.

FIGURE LEGENDS

Figure 1.

Differential gene expression between wild (n=4) and captive (n=4) shrews in the cortex, olfactory bulb,
and hippocampus. *Left:* PCA of the top 500 varying genes for each brain region. *Right*: Volcano plots for
the cortex, olfactory bulb, and hippocampus gene expression. Significant ($p_{adj} < 0.05$) DEGs are shown in
red (downregulated) or blue (upregulated) relative to wild, DEGs of interest are labeled. Labeled
downregulated genes encode for ribosomal proteins and mitochondrial complex I subunit proteins,
which have also been differentially expressed in previous studies. Labeled upregulated genes
correspond to the top 6 genes from the most significant upregulated enriched pathways.

Figure 2.

Overlapping DEGs for the three brain regions. The side histogram corresponds to the number of DEGs
for each brain region (set size), and the top histogram corresponds to intersections between the sets of
DEGs across the three brain regions. All the 104 DEGs that overlap between the three brain regions
(highlighted in blue) have the same directional change in gene expression (79 downregulated, 25
upregulated).

Figure 3.

Left: Pathway analysis for captive and wild shrews for the cortex, olfactory bulb and hippocampus. These
pathways were significantly enriched across the three brain regions when analyzing downregulated
genes, and were also enriched when analyzing only the 104 overlapping significant genes. *Right:* Side
histogram represents percentage of complex I subunit genes in each of these pathways.

TABLES
546

547 **Table 1.** – Descriptions of individuals used in gene expression experiments.

Individual	Condition	Sex	Toothrow (mm)	Cortex	Hippocampus	Olfactory Bulb
1	Wild	F	7.15	X	X	X
2	Wild	M	7.05	X	X	X
3	Wild	F	7.1	X	X	X
4	Wild	M	7.15	X	X	X
5	Captive	M	7	X	X	X
6	Captive	M	7	X	X	NA
7	Captive	M	6.9	X	X	X
8	Captive	M	7.25	X	X	X
9	Captive	M	6.9	NA	NA	X

550 # SUPPLEMENTAL MATERIAL

551 Supplemental Figs. S1–S7, supplemental table 2: https://doi.org/10.6084/m9.figshare.27237300.v1

552 Supplemental Table 1: DOI: https://doi.org/10.5061/dryad.qz612jmng