Immediate early genes as a molecular switch for lasting vulnerability following

pubertal stress in mice

Karissa N. Gautier[1], Samantha L. Higley[1], John M. Mendoza[1], and Kathleen E.

Morrison[1,2]*

[1]Department of Psychology, West Virginia University, Morgantown, WV, USA

[2]Department of Neuroscience, West Virginia University, Morgantown, WV, USA

*Corresponding author
Kathleen E. Morrison
Assistant Professor
kathleen.morrison@mail.wvu.edu
304-293-1685

Word count: 4648
Running head: Molecular vulnerability following pubertal stress
Key words: stress, sex differences, puberty, transcriptome, hypothalamus,
allopregnanolone

Abstract

Why individuals have negative consequences following stress is a complex

phenomenon that is dictated by individual factors, the timing of stress within the

lifespan, and when the consequences are measured. Women who undergo adverse

childhood experiences are at risk for lasting biological consequences, including affective

and stress dysregulation. We have shown that pubertal adversity is associated with a

blunted glucocorticoid response within the hypothalamic-pituitary-adrenal axis in both

peripartum humans and mice. In mice, we examined puberty-stress reprogramming in

the paraventricular nucleus (PVN) of the hypothalamus, which initiates the HPA axis

response. We found that pubertal stress led to an upregulation of six immediate early

genes (IEGs) in the PVN of adult, pregnant mice. Separately, we showed that the

pregnancy-associated hormone allopregnanolone is necessary and sufficient to produce

the blunted stress response phenotype in pubertally stressed mice. Here, we examined

the response of the IEGs in the PVN to the primary disruption of pubertal stress in early

adolescence and to the secondary disruption of increased allopregnanolone in

pregnancy. We found that in adult female, but not male, mice previously stressed during

puberty, intra-PVN allopregnanolone was sufficient to recapitulate the pubertal stress

associated baseline IEG expression profile. We also examined baseline IEG expression

during adolescence, where we found that IEGs have sex-specific developmental

trajectories that were disrupted by pubertal stress. Altogether, these data establish that

IEGs can act as a key molecular switch that leads to increased vulnerability to negative

outcomes in adult, pubertally stressed animals. Understanding how the factors that

produce vulnerability combine throughout the lifespan will further our understanding of

50   the etiology of negative outcomes and will help guide both the nature and timing of

51   potential treatments.

Introduction

Risk for neuropsychiatric disorders in adulthood is increased by exposure to adverse childhood experiences (ACEs) (1,2). Women who are exposed to ACEs during puberty are at the greatest risk for neuropsychiatric disorders across the lifespan (3–5) and suffer unique physiological and biological symptoms that are not shared by those who experienced adversity in childhood or adulthood (6–8). We previously established that adversity during puberty in female mice mimicked pubertal ACEs in women, significantly blunting their stress reactivity only during the peripartum window (9). Risk of peripartum depression and anxiety are elevated in women with high adversity and are associated with long-term negative outcomes for both mother and baby (10). Although we know that multiple experiences such as pubertal stress and pregnancy compound over the lifespan to influence risk for negative outcomes, we have limited understanding of the developmental trajectory of the biological programming that underlies increasing risk.

We have shown lasting effects of stress during puberty on the physiology and behavior of female mice and humans (8,9,11,12). To understand the mechanisms underlying the lasting effects of pubertal stress, our focus has been on the paraventricular nucleus of the hypothalamus (PVN), both a key brain region that regulates the hypothalamic-pituitary-adrenal (HPA) stress axis response and a region we have previously identified to have disruption of both the epigenome and transcriptome following pubertal stress (9,11). It has been of great interest to scientists and clinicians to understand the molecular underpinnings of the lasting effects of exposure to stress or trauma early in life (13–15). Here, we leveraged our pubertal

stress-associated molecular signature, where six immediate early genes (IEGs) were

permissively expressed at baseline in the PVN of pregnant females, to provide insight

into whether early life stress-induced molecular changes are present as an immediate

consequence of pubertal stress or are apparent only in adulthood. In addition to the IEG

signature, we have shown that allopregnanolone may be the part of pregnancy that is

responsible for unmasking the latent molecular programming. However, whether

allopregnanolone and the IEGs are linked is unknown. Although IEGs are associated

with neural activation and plasticity after exposure to stimuli, IEG mRNA and proteins

are canonically very low to undetectable at baseline (16–18). So, their increased

expression at baseline in pregnant, pubertally stressed females suggests a potential

mechanistic role in influencing how the PVN is poised to respond to stimuli. Further,

pubertal stress occurs much earlier in the lifespan compared to when we examine

physiology and the transcriptional state of the brain (early adolescence vs adulthood).

We also examined whether IEGs are the primary programming of pubertal stress by

addressing whether baseline IEG expression is dynamic during puberty and whether

any potential change is influenced by pubertal stress.

Therefore, we examined the response of the IEGs in the PVN to the primary

disruption of pubertal stress in early adolescence and to the secondary disruption of

increased allopregnanolone in pregnancy. We hypothesized that if the IEGs are a

molecular trigger of the negative consequences we have observed in pubertally

stressed, pregnant mice and humans, we would observe increased responsiveness of

these genes in the PVN of pubertally stressed mice in both developmental phases.

Methods

Animals

All mice were virgin, in house mixed strain C57BL/6:129 (N = 143) (9,11,12). Some

mice were heterozygous *Crh-IRES-Cre;Ai14*, capable of expressing tdTomato in

corticotropin releasing factor cells, although the transgenic background was not utilized

in this study (19,20). All mice were kept on a 12-hour light-dark cycle and food and

water were available *ad libitum*. All procedures were approved by the West Virginia

University Institutional Animal Care and Use Committee and were conducted in

accordance with the National Institutes of Health Guide for the Care and Use of

Laboratory Animals.

Pubertal stress

Starting on postnatal day (PN) 21, mice underwent two weeks of chronic variable stress

(CVS) (9,11,12). The stressors began within the first hour of lights on (between 8:00 –

9:00 am) during the daily nadir of circadian corticosterone levels. CVS lasted two hours

each day, excluding the acute restraint, which was only 15 minutes in length. Two

stressors were randomly assigned each day and varied between three different sensory

modalities: tactile (wet bedding, wire mesh, no bedding, marbles, multiple cage

changes, 15-minute acute restraint stress), auditory (white noise, owl screech), and

olfactory (70% ethanol, puma odor [1:200 2-Phenethylamine, CAS 64-04-0, in mineral

oil]). Mice experienced rotating combinations of stressors (tactile and auditory, tactile

and olfactory, or auditory and olfactory), combinations of which also rotated between

specific stressors in each category. For Experiment 1, mice in the CVS group were

weaned into singly housed cages at the beginning of stress (PN21) and were pair-

housed with a same-sex, same-stress littermate at the end of the 14 days of stress

(PN34). Control mice were weaned on PN28 into same-sex, pair-housed cages. They

were left undisturbed until adulthood (10-18 weeks). For Experiment 2, mice in the CVS

group were weaned into singly housed cages at the beginning of stress (PN21). They

were left undisturbed for at least 24 hours after the last stressor and prior to brain

collection, which occurred at PN28 or PN35. Control mice collected at PN21 and PN28

were removed directly from the litter at the time of collection. Control mice used for

PN35 collations were weaned on PN28 into same-sex, pair-housed cages.

Experiment 1: pharmacological manipulation of allopregnanolone in adulthood

We previously observed pubertal stress-associated physiological, transcriptomic, and

chromatin outcomes in adulthood when outcomes were examined under conditions of

high allopregnanolone (pregnancy or peripheral allopregnanolone treatment, **Figure**

1A). Here we sought to mechanistically link allopregnanolone to the transcriptomic

outcomes, a phenotype where six immediate early genes (IEGs) are permissively

expressed in the PVN at baseline conditions. (**Figure 1B**). To test a direct role for

allopregnanolone in the IEG phenotype in the PVN, adult male and female mice were

treated with allopregnanolone (100 ng per side of PVN) or vehicle (20% weight/volume

2-Hydroxypropyl-β-cyclodextrin in water). Allopregnanolone was suspended in vehicle

at 100ng/200nL for intra-PVN administration. It was prepared in advance and stored at -

20°C.

In adulthood (age 10-18 weeks), female (N = 26) and male (N = 19) mice

underwent stereotaxic cannulation to selectively target the PVN. Mice were

anesthetized using 4% isoflurane in an induction chamber and immediately connected

to the surgery setup, where they received constant isoflurane (0.5%-2.5%). In addition,

local analgesic carprofen (SC, 5 mg/kg) and anesthetics bupivacaine (at incision site,

1.5 mg/kg) and lidocaine (at incision site, 0.5 mg/kg) were administered before

beginning the procedure. A double-barreled cannula (0.6mm center-to-center placing,

4mm cut below pedestal, P1 Technologies) was implanted in the brain at the PVN

relative to bregma (-0.3mm medial/lateral, -0.85mm anterior/posterior) and lowered

3.8mm below the skull. Dental cement was used to fill the incision and create a headcap

around the cannula, securing it to the skull. Mice were monitored for five consecutive

days and received peripheral carprofen (SC, 5 mg/kg) for the first three days post-

surgery. Following recovery (7-22 days), 200nL of allopregnanolone (100 ng/200 nl) or

200 nl vehicle was simultaneously infused into each side of the double-barreled cannula

using two Hamilton Microsyringes and Dual Syringe Nanoliter Pump over a 1 minute

period. The needle was left in place for an additional 1 minute to allow the infusion to

diffuse away from the needle. Mice were briefly restrained to accomplish the insertion

and removal of the injection needle. The dose and timing have been shown effective in

altering behavior in prior studies (21,22). Mice were euthanized two hours later via

cervical dislocation. Brains were frozen on dry ice and stored at -80°C until use.

Experiment 2: Influence of CVS on IEG expression during adolescence

Brains were collected from female and male mice at one of three time points: PN21,

PN28, or PN35. At PN28 and PN35, brains were either from Control, unstressed mice

or CVS mice 24 hours after the last stressor. Sample sizes started at N=10 per group,

although there was some attrition (**Table S1**). Mice were anesthetized with vaporized

Isoflurane and brains were collected under baseline (non-stimulated) conditions. Brains

were frozen on dry ice and stored at -80°C until use.

PVN collection and analysis

Brains were cryosectioned to collect precise samples of the PVN. Two, 300 um slices

were taken from the brain at the anatomic location of the PVN using Paxinos and

Franklin's stereotaxic coordinates (23) for all brains, except those from PN21 animals,

for which we utilized two, 250 um slices. Pre-chilled 1mm biopsy tissue punchers were

used to remove the PVN from each slice. PVN biopsies were ejected into a clean, pre-

chilled 1.5 mL tube and stored at -80°C until use.

Verification of accurate cannula placement for Experiment 1 was made by

inspecting the 300um slices of each sample for clear indications that the cannula and

infusion needles had passed through the tissue. Further confirmation of accurate PVN

location was completed by taking 10um slices before and after the region. The sections

were stained using neutral red to verify anterior and posterior sites around the PVN.

Samples were omitted from the study if the injection site fell outside of this region or

neutral red staining revealed incorrect cryosectioning of the PVN (1 male and 1 female,

Table S1).

Gene expression

We examined the expression of the six target IEGs (*Fos, Fosb, Arc, Egr1, Junb,* and

Jund). RNA was isolated from PVN tissue using the RNeasy Mini Kit (Qiagen) following

the manufacturer's instructions. The quality of isolated RNA was assessed using a

NanoDrop and Qubit RNA High Sensitivity Kit. The Applied Biosystems High-Capacity

cDNA Reverse Transcriptase Kit was used to generate cDNA, and the quality of cDNA

was assessed on a NanoDrop.

Gene expression was measured using the QuantStudio 5 quantitative real-time

PCR (qRT-PCR) system. All gene expression assays were prepared with TaqMan Fast

Advanced Master Mix and probes for the target genes and reference gene *Gapdh*

(**Table S2**). Each reaction was run in triplicate using an equal volume of cDNA and

Gapdh was run on each plate. Relative gene expression was calculated using the ΔΔCt

method.

Statistical analyses

For experiment 1, gene expression data were analyzed by a 3-way ANOVA with sex,

stress, and treatment (vehicle vs. allopregnanolone) as factors. Interactions were further

investigated by 2-way ANOVAs (stress x treatment) within sex. For experiment 2, gene

expression data were analyzed using several approaches: independent samples t-test

for baseline sex differences at PN21, linear regression for trajectory of gene expression

from PN21 to PN35, and 3-way ANOVA with sex, stress, and age (P28, PN35) as

factors. Fisher's least significant difference post-hoc tests were utilized if appropriate.

Data were considered outliers and excluded from analysis if they were ± two standard

deviations from the mean. Sample sizes for all groups are provided in the Supplement

(**Table S1**). All analyses were performed in Prism (GraphPad) with an alpha level of $p <$

0.05.

Results

Intra-PVN allopregnanolone is sufficient to recapitulate the pubertal stress associated

IEG expression profile in adult females.

We have previously shown that pubertal stress (CVS) resulted in a blunted

corticosterone response to acute restraint stress in adult mice during pregnancy or with

allopregnanolone treatment (**Figure 1A**). Further, we identified a distinct epigenetic

phenotype in the PVN in pregnant, pubertally stressed females, such that the chromatin

landscape was open and there was permissive expression of six IEGs at baseline.

Here, we tested whether it is the increased levels of allopregnanolone present in

pregnancy that interact with prior pubertal stress experience to increase baseline IEG

expression in the PVN (**Figure 2A**).

Adult female and male mice that had either been exposed to CVS or not received

an intra-PVN injection of allopregnanolone or vehicle. In five of the six IEGs (*Arc, Egr1,*

Fos, Junb, Jund), there was an effect of sex on baseline gene expression (**Table S3**).

There was no effect of sex, stress, or allopregnanolone on the expression of *Fosb*

(**Figure S1**). Our findings in vehicle-treated mice are consistent with our previous work,

such that there were no significant differences between Control and CVS vehicle-

treated adult mice (**Figure 2B,C**). The heatmap shows that males generally have

decreased expression relative to vehicle treated female Controls, and the expression in

males is less dynamic than in females in response to pubertal stress or intra-PVN

allopregnanolone. This is consistent with our previous findings showing that males are

less vulnerable to the effects of pubertal stress in the PVN and on the HPA axis than

females.

Because there was an effect of sex on IEG expression, we examined the

relationship between pubertal stress and intra-PVN allopregnanolone within sex. In

males, there were no significant effects of pubertal stress or intra-PVN allopregnanolone

on any of the IEGs ($p > 0.05$). However, in females, there was a significant interaction

between pubertal stress and intra-PVN allopregnanolone on baseline gene expression

for *Arc* (**Figure 2D**, $F(1, 18) = 8.68$, $p = 0.008$), *Egr1* (**Figure 2E**, $F 1, 18) = 7.54$, $p =$

0.01), and *Fos* (**Figure 2F**, $F(1, 18) = 4.84$, $p = 0.04$). For these genes, post-hoc testing

revealed the same thing. Under allopregnanolone treatment, the gene expression for

Control and CVS females diverged, such that Control females had significantly

decreased gene expression relative to Control Vehicle females (*Arc* $p = 0.01$, *Egr1* $p =$

0.03, *Fos* $p = 0.02$) and relative to CVS Allopregnanolone females (*Arc* $p = 0.001$, *Egr1*

$p = 0.002$, *Fos* $p = 0.03$). The same non-significant pattern was observed in the

expression of *Fosb, Junb,* and *Jund* (**Figure S1**). Thus, acute allopregnanolone

delivered into the PVN of adult females recapitulates the phenotype that we have

previously observed in the brain of late pregnant females, such that pubertally stressed

females have increased baseline IEG expression relative to control females. These

findings are specific to females, as adult males show no significant effect of pubertal

stress or intra-PVN allopregnanolone on baseline IEG expression.

IEGs have sex-specific developmental trajectories in adolescence that are disrupted by

chronic pubertal stress.

To further understand the potential mechanistic role for the baseline IEG

expression patterns in the lasting consequences of pubertal stress, we measured gene

expression in the PVN during the age range when CVS is applied (PN21-35, **Figure**

3A). We then examined whether each gene had any type of variability in baseline

expression between the three ages examined (PN21, PN28, PN35) in Control animals.

Linear regression analysis showed that there was a significant slope in the trajectory

from PN21 to PN35 for *Arc* ($F(1,47) = 6.58$, $p = 0.01$) and *Jund* ($F(1,50) = 6.04$, $p =$

0.02) in Controls (**Figure 3B, Figure S2**). For *Arc*, exposure to pubertal stress led to a

flattening of the gene expression trajectory, with CVS mice having no significant slope in

the trajectory from PN21 to PN35 ($p > 0.05$, **Table S4**). For *Egr1*, *Fos*, and *Fosb*, there

was no change in trajectory for expression from PN21 to PN35 in Controls ($p > 0.05$,

Table S4). However, exposure to pubertal stress led to a significant decrease in gene

expression from PN21 to PN35 in CVS mice for *Egr1* ($F(1,53) = 5.90$, $p = 0.02$), *Fos* (F

$(1,50) = 59.83$, $p < 0.0001$), *Fosb* ($F (1,51) = 18.57$, $p < 0.0001$), and *Jund* ($F (1,52) =$

7.83, $p = 0.01$). We performed further testing to confirm that the regression lines

significantly differed between Control and CVS mice. A test of differences in slope or

elevation showed that Control and CVS mice had significantly different slopes of

expression through the early adolescent window for *Egr1*, *Fos*, *Fosb*, *Junb*, and *Jund*.

Some of the effects on the trajectory of expression were more pronounced in one sex

(**Figure 3C**).

At PN21, there were no differences between females and males in baseline gene

expression (**Table S5**). Due to the unbalanced design that results from the inclusion of

PN21 pre-stress gene expression in an ANOVA, we analyzed gene expression in PN28

and PN35 mice. Gene expression was calculated relative to that of PN21 Controls. For

four of the six IEGs, there was a significant interaction between age and sex (*Fosb,*

Junb, Jund) or a main effect of age (*Egr1*) on gene expression in the PVN (**Table S6**).

This effect of age provides further support that, even in non-stimulated, baseline

conditions, IEG expression in the PVN has a dynamic developmental trajectory

throughout early adolescence.

IEG expression was altered by exposure to pubertal CVS. Gene expression was

examined either after 1 week (PN28) or the complete 2 weeks (PN35) of CVS. As

brains were collected 24 hours after the final stressor, these findings represent non-

stimulated, baseline levels of gene expression following chronic stress. For four of the

six genes, there was a significant effect of stress (*Fos, Fosb, Junb*) or interaction

between stress, sex, and age (*Egr1*) on gene expression in the PVN (**Table S6, Figure**

S3).

Overall, these analyses show that IEGs were influenced in an interactive way by

age, stress, and pubertal stress. For example, *Egr1* displayed a dynamic change

throughout the pubertal stress window in Control mice in which females and males

showed an opposite pattern of expression from PN21 to PN35 (**Figure 3D-E**). Some

IEGs, like *Fos*, were strongly influenced by pubertal stress (**Figure 3F-G**). Thus, for all

genes, pubertal stress altered the trajectory of normative baseline gene expression

during the two-week window during which CVS is applied (PN21 – PN35).

Discussion

Why individuals have negative consequences following stress is a complex phenomenon that is dictated by individual factors (e.g., genetics, sex, resilience), when in the lifespan the stress is experienced (e.g., childhood, adolescence), and when the consequences are measured, both proximity to the stressful event and timepoint within the lifespan (24–29). Understanding how these factors combine will further our understanding of the etiology of negative outcomes and will help guide both the nature and timing of potential treatments. For women, who are more likely than men to suffer from affective disorders, risk factors include experiencing stressful or traumatic events around the onset of puberty and experiencing later times of hormonal change such as pregnancy and aging (30,31). We have previously shown that in both humans and mice, adversity experienced around the onset of puberty led to a blunted hypothalamic-pituitary-adrenal (HPA) stress response during the peripartum window and increased reporting of postpartum depressive symptoms in humans (9). We have used this translationally-relevant mouse model to investigate the mechanisms underlying the lasting effect of pubertal stress on the HPA axis response during the peripartum window. Our previous work identified latent epigenetic changes in the paraventricular nucleus of the hypothalamus (PVN) that may be uncovered by the increased allopregnanolone levels during pregnancy (11). Here, we expand on those findings to directly link allopregnanolone action in the PVN to the molecular changes associated with pubertal stress.

Our previous transcriptomic analysis of the PVN revealed increases in the

baseline expression of six immediately early genes (IEGs) in pregnant females as a

result of pubertal adversity (9). Here, we utilized PVN-specific pharmacological

manipulations to address whether allopregnanolone alone and delivered acutely is

sufficient to induce an increase in baseline IEG expression in pubertally stressed mice.

In adult female mice, a single dose of allopregnanolone administered into the PVN

recapitulated the increased IEG expression only in pubertally stressed females.

Allopregnanolone is a neurosteroid metabolite of progesterone that is present at high

levels in the brain during pregnancy. It binds to several receptors, including

extrasynaptic GABA-A, pregnane X receptors, and progesterone receptors (32).

Notably, the function of allopregnanolone as an FDA approved treatment for postpartum

depression, under the names Brexanolone and Zuranolone, is attributed to the functions

of allopregnanolone at the GABA-A receptor (33–35). In our studies, allopregnanolone

does not act to reduce risk for negative outcomes in prenatal mice. One potentially

meaningful difference between the models that established allopregnanolone as an

effective postpartum depression treatment are that those do not involve early life stress

and our model does. Future work will need to resolve whether individuals who have a

history of preconception adversity are likely to fully benefit from neurosteroid-based

antidepressants. It is possible that the effect of allopregnanolone in altering the

transcriptome in response to pubertal stress is working via different mechanisms,

including the transcription-regulating pregnane X receptor (36).

These findings provide further evidence that increased baseline IEG expression

is a key molecular switch that may be 'wired' by pubertal stress and leads to the

vulnerability of an altered HPA axis in adulthood, during which allopregnanolone can

'flip the switch' to uncover the vulnerability. However, whether this molecular switch is

wired immediately after pubertal stress, at postnatal day 35, or whether it develops in

the weeks between PN35 and adulthood, was unknown. Understanding the first steps in

the molecular cascade that result in negative outcomes are typically identified in

adulthood will lead to early interventions and preventative approaches for

neuropsychiatric disorders (37). Therefore, we examined IEG expression in the PVN

during the window of pubertal stress in females and males. First, we examined

developmental changes in gene expression in female and male mice. Overall, baseline

IEG expression in the PVN was dynamic during the window when we apply pubertal

stress. Males and females did not differ in gene expression at PN21, but often showed

different trajectories of gene expression when examined 1 week (PN28) or 2 weeks

(PN35) later. These data show that IEG expression in the PVN is not static. We found

that pubertal stress altered IEG expression during this adolescent window. In general,

the effect of pubertal stress was to decrease baseline IEG expression in both females

and males.

Future studies will resolve these findings in a cell-type specific manner. In our

current and past studies, we have examined the complete PVN. While limited work has

addressed cell-type specific outcomes in the hypothalamus following adolescent stress,

there is work from early prenatal stress that informs future cell-type specific studies.

HPA axis activation is dictated by the signaling of corticotropin-releasing factor (CRF),

glutamatergic, and GABA cells. Limited bedding and nesting produced alterations to the

transcriptome of CRF cells in 1.5 week old mice. Single-cell sequencing of CRF cells in

the early postnatal period provides evidence for enhanced sensitivity of CRF cells to the

stress of limited bedding and nesting (38). This single-cell sequencing experiment

showed that, although GABA neurons were not analyzed, it was CRF cells that co-

expressed glutamate, and not GABA, that showed sensitivity to this early life adversity.

This is consistent with prior work showing increased synaptic glutamatergic drive onto

neonatal PVN CRF neurons following limited nesting (39). This study also showed that

mice exposed to limited nesting had altered sensitivity to allopregnanolone. It is possible

that what we and others are finding is a mechanism wherein CRF cells are being tuned

by both glutamatergic and GABAergic systems, and that early life and adolescent

stressors impinge upon both neurotransmitter systems to lead to vulnerability to

negative outcomes later in life.

Another future goal will be to merge this cell-type specificity with tracking of the

same cells throughout the lifespan, which will overcome our current limitation of

examining adolescent and adult outcomes in different animals. Recent work has shown

that a stress exposure early, in the form of maternal separation, led to the prolonged

sensitivity of a set of cells in the nucleus accumbens to stressors in adulthood (40).

Using a transgenic strategy in mice to permanently mark cells that had increased *Arc*

expression in response to early life stress, Balouek et al showed that neurons tagged by

maternal separation were the same cells reactivated by stress later in life and that

blocking their activation ameliorated the negative behavioral consequences.

Interestingly, it was males that were sensitive to the adult manipulations. This is in

contrast to our study and others conducted in adolescence, where females tend to be

more vulnerable to lasting negative outcomes.

While our model considers times in the lifespan when females are more

vulnerable than males, we included male mice in all experiments here. Consistent with

our previous work showing that adult females are more vulnerable to the lasting effects

of pubertal stress, we found that IEG expression in males is less sensitive in adulthood

than it is in females. This contrasts with our finding that adolescent males display

changes in IEG expression the PVN in response to pubertal stress during the stress

window. Future studies will resolve why the molecular responsiveness in the PVN of

males does not also result in negative outcomes in adulthood and will consider other

brain regions and behavioral outcomes that might provide evidence of negative

plasticity in males. Altogether, our study and others conducted recently continue to lend

support for to the literature on sex differences in vulnerability to early life stress (41–44).

Ongoing studies will continue to refine our knowledge on the molecular underpinnings of

how sex and early life stress interact to produce vulnerability to negative outcomes.

Few studies have examined the expression levels of immediate early genes in

non-stimulated conditions, as they are expressed at low levels at baseline and are

canonically associated with stimulus-induced expression (16–18). Studies where either

IEG mRNA or protein have been measured have been focused on the impact of

neonatal experiences. For instance, embryonic rearing conditions of chick eggs causes

lateralization of spontaneous (baseline) c-Fos expression in several brain regions (45).

Another study that examined the regulators of *Arc* expression in response to acute

immobilization stress showed that blocking brain-derived neurotrophic factor reduces

baseline *Arc* expression in the neocortex, which gives insight into molecules that may

participate in the lasting effects that we have observed in our model (46). In our model,

four of the six IEGs are from the Jun and Fos family of proteins, which dimerize to form

the AP-1 complex in order to regulate transcription (47). The AP-1 complex interacts

with the glucocorticoid receptor, such that AP-1 binding is required in the recruitment of

GR (48). It is possible that the lasting HPA axis physiological plasticity observed in

pregnant, pubertally stressed females is driven by alterations to the AP-1 complex that

subsequently impact glucocorticoid signaling.

There are competing hypotheses for the way in which multiple stressors

throughout the lifespan interact to influence risk for negative outcomes. Some findings

support a mismatch theory, where organisms that experience early life stress are well

suited to future stressful environments and appear resilient (49). Our prior results and

new findings support the idea of cumulative stress, such that a 'second hit' challenge

results in an increased risk for negative outcomes, also termed an increased allostatic

load (50). However, when humans and animals experience that 'second hit' of

challenge, whether another exposure to stress later in life or to physiological changes

such as pregnancy, it has not been clear what the molecular agents of that 'hit' are. We

have identified that allopregnanolone is a molecular agent of that 'second hit' of risk and

that it interacts directly with IEGs to influence their baseline expression (Figure 4). This

interaction at baseline could lead to differences in how the cells in the PVN respond to

stimuli and underlie the blunted HPA axis response and altered behavior we have

previously observed.

We have taken a developmental approach to the question of how stress during

puberty impacts an individual through the lifespan, wherein we are generating a

trajectory of molecular, physiological, and behavioral outcomes both during early life

stress and into adulthood. Indeed, our findings show that a key molecular switch for the

negative adult outcomes after pubertal stress – the permissive expression of a set of

IEGs - is plastic, displaying normative changes throughout development, and that stress

during puberty results in a disruption of baseline expression that is detectable in

adolescence. These studies provide novel insight into the mechanisms underlying

female-relevant risk for stress dysregulation, a central endophenotype of affective

disorders. Understanding the early signs and mechanisms of negative outcomes like

stress dysregulation and affective dysfunction that, both in humans and animal models,

are generally identified and treated in adulthood, will lead to early interventions,

preventative approaches, and better outcomes for mental health.

Author Contributions

KEM designed experiments. KNG, SLH, JMM, and KEM conducted experiments,

sorted, and analyzed data. KNG wrote an initial draft of the manuscript, which was

edited to a final version by KEM.

Acknowledgments

This work was supported by National Institutes of Health Grant R00 HD091376 (KEM)

and start-up funds to KEM. KNG and JMM were additionally funded by the West Virginia

University Summer Undergraduate Research Experience program.

Disclosure Statement

The authors report no conflicts of interest.

Data Availability

Data will be publicly available via the Open Science Framework repository upon

publication.

References

1. Merrick MT, Ports KA, Ford DC, Afifi TO, Gershoff ET, Grogan-Kaylor A (2017): Unpacking the impact of adverse childhood experiences on adult mental health. *Child Abuse Negl* 69: 10–19.

2. Rosenberg SD, Lu W, Mueser KT, Jankowski MK, Cournos F (2007): Correlates of Adverse Childhood Events Among Adults With Schizophrenia Spectrum Disorders. *Psychiatr Serv* 58: 245–253.

3. Chapman DP, Whitfield CL, Felitti VJ, Dube SR, Edwards VJ, Anda RF (2004): Adverse childhood experiences and the risk of depressive disorders in adulthood. *J Affect Disord* 82: 217–225.

4. Janssen I, Krabbendam L, Bak M, Hanssen M, Vollebergh W, de Graaf R, van Os J (2004): Childhood abuse as a risk factor for psychotic experiences. *Acta Psychiatr Scand* 109: 38–45.

5. Heim C, Shugart M, Craighead WE, Nemeroff CB (2010): Neurobiological and psychiatric consequences of child abuse and neglect. *Dev Psychobiol* 52: 671–690.

6. Leenarts LEW, Vermeiren RRJM, van de Ven PM, Lodewijks HPB, Doreleijers TAH, Lindauer RJL (2013): Relationships Between Interpersonal Trauma, Symptoms of Posttraumatic Stress Disorder, and Other Mental Health Problems in Girls in Compulsory Residential Care. *J Trauma Stress* 26: 526–529.

7. Jaffee SR (2017): Child Maltreatment and Risk for Psychopathology in Childhood and Adulthood. *Annu Rev Clin Psychol* 13: 525–551.

8. Morrison KE, Stenson AF, Marx-Rattner R, Carter S, Michopoulos V, Gillespie CF, *et al.* (2022): Developmental Timing of Trauma in Women Predicts Unique Extracellular Vesicle Proteome Signatures. *Biol Psychiatry* 91: 273–282.

9. Morrison KE, Epperson CN, Sammel MD, Ewing G, Podcasy JS, Hantsoo L, *et al.* (2017):

Preadolescent Adversity Programs a Disrupted Maternal Stress Reactivity in Humans

and Mice. *Biol Psychiatry* 81: 693–701.

10. Slomian J, Honvo G, Emonts P, Reginster J-Y, Bruyère O (2019): Consequences of

maternal postpartum depression: A systematic review of maternal and infant outcomes.

Womens Health Lond Engl 15. https://doi.org/10.1177/1745506519844044

11. Morrison KE, Cole AB, Kane PJ, Meadows VE, Thompson SM, Bale TL (2020): Pubertal

adversity alters chromatin dynamics and stress circuitry in the pregnant brain.

Neuropsychopharmacology 45: 1263–1271.

12. Morrison KE, Narasimhan S, Fein E, Bale TL (2016): Peripubertal Stress With Social

Support Promotes Resilience in the Face of Aging. *Endocrinology* 157: 2002–2014.

13. Davis EP, McCormack K, Arora H, Sharpe D, Short AK, Bachevalier J, *et al.* (2022): Early

life exposure to unpredictable parental sensory signals shapes cognitive development

across three species. *Front Behav Neurosci* 16: 960262.

14. Daskalakis NP, Bagot RC, Parker KJ, Vinkers CH, de Kloet ER (2013): The three-hit

concept of vulnerability and resilience: toward understanding adaptation to early-life

adversity outcome. *Psychoneuroendocrinology* 38: 1858–1873.

15. Warhaftig G, Almeida D, Turecki G (2023): Early life adversity across different cell- types in

the brain. *Neurosci Biobehav Rev* 148: 105113.

16. Gallo FT, Katche C, Morici JF, Medina JH, Weisstaub NV (2018): Immediate Early Genes,

Memory and Psychiatric Disorders: Focus on c-Fos, Egr1 and Arc. *Front Behav*

Neurosci 12. Retrieved September 19, 2023, from

https://www.frontiersin.org/articles/10.3389/fnbeh.2018.00079

17. Hughes P, Lawlor P, Druganow M (1992): Basal expression of Fos, Fos-related, Jun, and

Krox 24 proteins in rat hippocampus. *Mol Brain Res* 13: 355–357.

18. Pérez-Cadahía B, Drobic B, Davie JR (2011): Activation and function of immediate-early

516   genes in the nervous system. *Biochem Cell Biol Biochim Biol Cell* 89: 61–73.

517   19. Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, *et al.* (2010): A robust

518   and high-throughput Cre reporting and characterization system for the whole mouse

519   brain. *Nat Neurosci* 13: 133–140.

520   20. Taniguchi H, He M, Wu P, Kim S, Paik R, Sugino K, *et al.* (2011): A resource of Cre driver

521   lines for genetic targeting of GABAergic neurons in cerebral cortex. *Neuron* 71: 995–

522   1013.

523   21. Rabinowitz A, Cohen SJ, Finn DA, Stackman RW (2014): The neurosteroid

524   allopregnanolone impairs object memory and contextual fear memory in male C57BL/6J

525   mice. *Horm Behav* 66: 238–246.

526   22. Engin E, Treit D (2007): The anxiolytic-like effects of allopregnanolone vary as a function of

527   intracerebral microinfusion site: the amygdala, medial prefrontal cortex, or hippocampus.

528   *Behav Pharmacol* 18: 461.

529   23. Paxinos G, Franklin KBJ (2019): *Paxinos and Franklin's the Mouse Brain in Stereotaxic*

530   *Coordinates*. Academic Press.

531   24. Hodes GE, Pfau ML, Leboeuf M, Golden SA, Christoffel DJ, Bregman D, *et al.* (2014):

532   Individual differences in the peripheral immune system promote resilience versus

533   susceptibility to social stress. *Proc Natl Acad Sci* 111: 16136–16141.

534   25. Morrison KE, Epperson CN, Bale TL (2020): Chapter 6 - Sex differences in the programming

535   of stress resilience. In: Chen A, editor. *Stress Resilience*. Academic Press, pp 81–94.

536   26. Rincón-Cortés M, Herman JP, Lupien S, Maguire J, Shansky RM (2019): Stress: Influence

537   of sex, reproductive status and gender. *Neurobiol Stress* 10: 100155.

538   27. Peña CJ, Nestler EJ, Bagot RC (2019): Environmental Programming of Susceptibility and

539   Resilience to Stress in Adulthood in Male Mice. *Front Behav Neurosci* 13. Retrieved

540   September 22, 2023, from https://www.frontiersin.org/articles/10.3389/fnbeh.2019.00040

28. Feder A, Fred-Torres S, Southwick SM, Charney DS (2019): The Biology of Human

Resilience: Opportunities for Enhancing Resilience Across the Life Span. *Biol Psychiatry*

86: 443–453.

29. Boyce WT, Levitt P, Martinez FD, McEwen BS, Shonkoff JP (2021): Genes, Environments,

and Time: The Biology of Adversity and Resilience. *Pediatrics* 147: e20201651.

30. Morrison KE (2020): Animal models built for women's brain health: Progress and potential.

Front Neuroendocrinol 59: 100872.

31. Li H, Bowen A, Bowen R, Balbuena L, Feng C, Bally J, Muhajarine N (2019): Mood

instability during pregnancy and postpartum: a systematic review. *Arch Womens Ment*

Health. https://doi.org/10.1007/s00737-019-00956-6

32. Walton N, Maguire J (2019): Allopregnanolone-based treatments for postpartum depression:

Why/how do they work? *Neurobiol Stress* 11: 100198.

33. Morrison KE, Cole AB, Thompson SM, Bale TL (2019): Brexanolone for the treatment of

patients with postpartum depression. *Drugs Today* 55: 537.

34. Deligiannidis KM, Meltzer-Brody S, Maximos B, Peeper EQ, Freeman M, Lasser R, *et al.*

(2023): Zuranolone for the Treatment of Postpartum Depression. *Am J Psychiatry* 180:

668–675.

35. Meltzer-Brody S, Colquhoun H, Riesenberg R, Epperson CN, Deligiannidis KM, Rubinow

DR, *et al.* (2018): Brexanolone injection in post-partum depression: two multicentre,

double-blind, randomised, placebo-controlled, phase 3 trials. *Lancet Lond Engl* 392:

1058–1070.

36. Frye CA, Koonce CJ, Walf AA (2014): Novel receptor targets for production and action of

allopregnanolone in the central nervous system: a focus on pregnane xenobiotic

receptor. *Front Cell Neurosci* 8: 106.

37. Uhlhaas PJ, Davey CG, Mehta UM, Shah J, Torous J, Allen NB, *et al.* (2023): Towards a

youth mental health paradigm: a perspective and roadmap. *Mol Psychiatry* 1–11.

38. Short AK, Thai CW, Chen Y, Kamei N, Pham AL, Birnie MT, *et al.* (2023): Single-Cell

Transcriptional Changes in Hypothalamic Corticotropin-Releasing Factor–Expressing

Neurons After Early-Life Adversity Inform Enduring Alterations in Vulnerabilities to

Stress. *Biol Psychiatry Glob Open Sci* 3: 99–109.

39. Gunn BG, Cunningham L, Cooper MA, Corteen NL, Seifi M, Swinny JD, *et al.* (2013):

Dysfunctional Astrocytic and Synaptic Regulation of Hypothalamic Glutamatergic

Transmission in a Mouse Model of Early-Life Adversity: Relevance to Neurosteroids and

Programming of the Stress Response. *J Neurosci* 33: 19534–19554.

40. Balouek J-A, Mclain CA, Minerva AR, Rashford RL, Bennett SN, Rogers FD, Peña CJ

(2023): Reactivation of Early-Life Stress-Sensitive Neuronal Ensembles Contributes to

Lifelong Stress Hypersensitivity. *J Neurosci* 43: 5996–6009.

41. Lupien SJ, McEwen BS, Gunnar MR, Heim C (2009): Effects of stress throughout the

lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci* 10: 434–445.

42. Sandman CA, Glynn LM, Davis EP (2013): Is there a viability-vulnerability tradeoff? Sex

differences in fetal programming. *J Psychosom Res* 75: 327–335.

43. Kim DR, Bale TL, Epperson CN (2015): Prenatal programming of mental illness: current

understanding of relationship and mechanisms. *Curr Psychiatry Rep* 17: 5.

44. O'Connor TG, Heron J, Golding J, Beveridge M, Glover V (2002): Maternal antenatal anxiety

and children's behavioural/emotional problems at 4 years. Report from the Avon

Longitudinal Study of Parents and Children. *Br J Psychiatry J Ment Sci* 180: 502–508.

45. Lorenzi E, Mayer U, Rosa-Salva O, Morandi-Raikova A, Vallortigara G (2019): Spontaneous

and light-induced lateralization of immediate early genes expression in domestic chicks.

Behav Brain Res 368: 111905.

46. Benekareddy M, Nair AR, Dias BG, Suri D, Autry AE, Monteggia LM, Vaidya VA (2013):

Induction of the plasticity-associated immediate early gene Arc by stress and

hallucinogens: role of brain-derived neurotrophic factor. *Int J Neuropsychopharmacol* 16:

405–415.

47. Hai T, Curran T (1991): Cross-family dimerization of transcription factors Fos/Jun and

ATF/CREB alters DNA binding specificity. *Proc Natl Acad Sci U S A* 88: 3720–3724.

48. Biddie SC, John S, Sabo PJ, Thurman RE, Johnson TA, Schiltz RL, *et al.* (2011):

Transcription Factor AP1 Potentiates Chromatin Accessibility and Glucocorticoid

Receptor Binding. *Mol Cell* 43: 145–155.

49. Nederhof E, Schmidt MV (2012): Mismatch or cumulative stress: toward an integrated

hypothesis of programming effects. *Physiol Behav* 106: 691–700.

50. Kuhn M, Scharfenort R, Schümann D, Schiele MA, Münsterkötter AL, Deckert J, *et al.*

(2016): Mismatch or allostatic load? Timing of life adversity differentially shapes gray

matter volume and anxious temperament. *Soc Cogn Affect Neurosci* 11: 537–547.

Figures

Figure 1. Relevant prior findings and experimental timelines for the current study.

(A) We previously established that chronic variable stress (CVS) during puberty leads to

a blunted hypothalamic-pituitary-adrenal axis in adulthood only when allopregnanolone

is present at high levels in the brain, including during pregnancy and under

allopregnanolone treatment (9,11). We identified the paraventricular nucleus of the

hypothalamus (PVN) as a key tissue of the HPA axis that was reprogrammed by

pubertal stress. Pubertal stress and pregnancy combined to lead to an altered

chromatin landscape, including an increase in the number of open chromatin sites, and

permissive expression of six immediate early genes (IEGs) at baseline. These findings

suggest that IEGs play a mechanistic role in the puberty-stress induced phenotype that

is apparent during times of dynamic change in allopregnanolone, including pregnancy.

(B) Here, we directly tested the relationship between allopregnanolone within the PVN

and the puberty stress associated IEG expression. Female and male mice were

exposed to CVS from PN21-34. In adulthood, mice were implanted with a double-barrel

cannula aimed at the PVN. Following recovery, mice were injected with either vehicle or

allopregnanolone. Two hours later, brains were collected and processed for the

measurement of gene expression. **(C)** We further examined a potential mechanistic role

for IEG expression during the pubertal stress window. Brains were collected in baseline,

non-stimulated conditions from female and male mice at either at PN21 (Control only),

PN28 (Control and CVS), or PN35 (Control and CVS). Brains from CVS mice were

collected 24 hours after exposure to the last stressor.

Figure 2. Intra-PVN allopregnanolone interacted with pubertal stress and sex to alter baseline immediate early gene expression.

(A) Adult female and male mice that had been exposed to pubertal stress or not were cannulated so that vehicle or allopregnanolone could be microinjected into the PVN. Two hours after microinjection, brains were collected in non-stimulated conditions. The PVN was isolated, RNA was isolated, and qRT-PCR was performed for six puberty-stress associated IEGs (*Arc, Egr1, Fos, Fosb, Junb, Jund*). **(B)** Heat map showing expression relative to Control Vehicle females within each gene. Relative expression was calculated using the ΔΔCt method. Intra-PVN Allopregnanolone generally led to a decrease in IEG expression in Control females. In contrast, CVS females responded with an increase in IEG expression. **(C)** In contrast to females, males generally had less expression of all IEGs, except for *Jund*. IEG expression in males was less sensitive to

either pubertal stress or intra-PVN Allopregnanolone. **(D-F)** When IEGs were examined

within sex, there was a significant interaction between pubertal stress (CVS) and intra-

PVN allopregnanolone for *Arc*, *Egr1*, and *Fos.* This expression recapitulates previous

findings in pregnant females, where pubertally stressed females with high

allopregnanolone levels in the PVN had increased IEG expression at baseline. CVS =

chronic variable stress, IEG = immediate early gene, PVN = paraventricular nucleus of

the PVN. $*p < 0.05$ on Fisher's LSD post-hoc test.

Figure 3. Baseline immediate early genes expression was dynamic during early

adolescence and was altered by pubertal stress.

(A) Brains were collected from young female and male mice that were exposed to

pubertal stress or not at three time points. These time points represent prior to starting

CVS (PN21), after 1 week of CVS (PN28), and after two weeks of CVS (PN35). Brains

were collected 24 hours after the prior stressor and in non-stimulated conditions. The

PVN was dissected, RNA was isolated, and qRT-PCR was performed for six puberty-

stress associated immediate early genes (IEG; *Arc, Egr1, Fos, Fosb, Junb, Jund*). **(B)**

Heat map depicting the direction of the slope of the linear regression line calculated

from PN21 to PN35 for gene expression in the PVN of Control (female and male) and

CVS (female and male) mice. The heat map shows that CVS disrupts the trajectories of

each of the six IEGs, and always by decreasing gene expression. **(C)** There was a sex

difference in the impact of pubertal stress on the trajectory of baseline IEG expression

during adolescence for several genes. **(B,C)** Asterisks indicate a slope of the regression

line from PN21-PN35 is significantly non-zero, while pound signs indicate that the

slopes are significantly different between the groups within a gene. **(D,E)** Some IEGs

were influenced in an interactive way by age, stress, and pubertal stress. **(D)** *Egr1*

displayed a dynamic change throughout the pubertal stress window in Control mice in

which females and males showed an opposite pattern of expression from PN21 to

PN35. **(E)** The experience of pubertal stress interacted with age and sex to determine

Egr1 expression in the PVN. **(F,G)** Some IEGs were strongly influenced by pubertal

stress, like *Fos*. **(F)** In Control mice, females and males show little change during the

PN21-PN35 age window. **(F,G)** However, pubertal stress led to a significant decrease in

Fos expression at both PN28 and PN35 in female and male mice. CVS = chronic

variable stress, IEG = immediate early gene, PN = postnatal day, PVN = paraventricular

nucleus of the PVN. (B-C) *p < 0.05 for non-zero slope following linear regression; #p <

0.05 for significantly different slopes between groups but within gene (D-G) *p < 0.05 for

main effect of CVS or for interaction terms following 3-way ANOVA analysis.

Figure 4. Schematic exemplifying our findings on the influence of pubertal stress on the transcriptome of the developing and adult hypothalamus. Through prior and current findings, we have demonstrated that a set of immediate early genes in the paraventricular nucleus of the hypothalamus is sensitive to pubertal stress (chronic variable stress from postnatal days 21-34) both during the stress exposure and in adulthood. Of note, all IEG expression differences were measured in non-stimulated, baseline conditions, and so may represent how the PVN is poised to respond to stimuli. Given here is the example of *Fos*, which is decreased during the pubertal stress window relative to Controls. Whether adult mice are pregnant or are exposed to allopregnanolone, there is an increase in baseline IEG expression. This suggests that the PVN is poised to respond differently, which we have observed in the form of a blunted physiological response to restraint in adulthood. The new findings that baseline IEG expression is altered provide support for a role of IEGs in the mechanism underlying the lasting effect of pubertal stress on the PVN.

Figure Legends

Figure 1. Relevant prior findings and experimental timelines for the current study.

(A) We previously established that chronic variable stress (CVS) during puberty leads to a blunted hypothalamic-pituitary-adrenal axis in adulthood only when allopregnanolone is present at high levels in the brain, including during pregnancy and under allopregnanolone treatment (9,11). We identified the paraventricular nucleus of the hypothalamus (PVN) as a key tissue of the HPA axis that was reprogrammed by pubertal stress. Pubertal stress and pregnancy combined to lead to an altered chromatin landscape, including an increase in the number of open chromatin sites, and permissive expression of six immediate early genes (IEGs) at baseline. These findings suggest both that IEGs play a mechanistic role in the puberty-stress induced phenotype that is apparent during times of dynamic change in allopregnanolone, including pregnancy. (B) Here, we directly tested the relationship between allopregnanolone within the PVN and the puberty stress associated IEG expression. Female and male mice were exposed to CVS from PN21-34. In adulthood, mice were implanted with a double-barrel cannula aimed at the PVN. Following recovery, mice were injected with either vehicle or allopregnanolone. Two hours later, brains were collected and processed for the measurement of gene expression. (C) We further examined a potential mechanistic role for IEG expression during the pubertal stress window. Brains were collected in baseline, non-stimulated conditions from female and male mice at either at PN21 (Control only), PN28 (Control and CVS), or PN35 (Control and CVS). Brains from CVS mice were collected 24 hours after exposure to the last stressor.

Figure 2. Intra-PVN allopregnanolone interacts with pubertal stress and sex to

alter immediate early gene expression. (A) Adult female and male mice that had

been exposed to pubertal stress or not were cannulated so that vehicle or

allopregnanolone could be microinjected into the PVN. Two hours after microinjection,

brains were collected in non-stimulated conditions. The PVN was isolated, RNA was

isolated, and qRT-PCR was performed for six puberty-stress associated immediate

early genes (IEG; *Arc, Egr1, Fos, Fosb, Junb, Jund*). **(B)** Heat map showing expression

relative to Control Vehicle females within each gene. Relative expression was

calculated using the ∆∆Ct method. Intra-PVN Allopregnanolone generally led to a

decrease in IEG expression in Control females. In contrast, CVS females responded

with an increase in IEG expression. **(C)** In contrast to females, males generally had less

expression of all IEGs, except for *Jund*. IEG expression in males was less sensitive to

either pubertal stress or intra-PVN Allopregnanolone. **(D-F)** When IEGs were examined

within sex, there was a significant interaction between pubertal stress (CVS) and intra-

PVN allopregnanolone for *Arc*, *Egr1*, and *Fos*. This expression recapitulates previous

findings in pregnant females, where pubertally stressed females with high

allopregnanolone levels in the PVN had increased IEG expression at baseline. CVS =

chronic variable stress, PVN = paraventricular nucleus of the PVN. $*p < 0.05$ on Tukey

post-hoc test.

Figure 3. Baseline immediate early genes expression was dynamic during early

adolescence and was altered by pubertal stress. (A) Brains were collected from

young female and male mice that were exposed to pubertal stress or not at three time

points. These time points represent prior to starting CVS (PN21), after 1 week of CVS

(PN28), and after two weeks of CVS (PN35). Brains were collected 24 hours after the

prior stressor and in non-stimulated conditions. The PVN was dissected, RNA was

isolated, and qRT-PCR was performed for six puberty-stress associated immediate

early genes (IEG; *Arc, Egr1, Fos, Fosb, Junb, Jund*). **(B)** Heat map depicting the

direction of the slope of the linear regression line calculated from PN21 to PN35 for

gene expression in the PVN of Control (female and male) and CVS (female and male)

mice. The heat map shows that CVS disrupts the trajectories of each of the six IEGs,

and always by decreasing gene expression. **(C)** There was a sex difference in the

impact of pubertal stress on the trajectory of baseline IEG expression during

adolescence for several genes. **(B,C)** Asterisks indicate a slope that is significantly non-

zero, while pound signs indicate that the slopes are significantly different between the

groups within a gene. **(D,E)** Some IEGs were influenced in an interactive way by age,

stress, and pubertal stress. **(D)** *Egr1* displayed a dynamic change throughout the

pubertal stress window in Control mice in which females and males showed an opposite

pattern of expression from PN21 to PN35. **(E)** The experience of pubertal stress

interacted with age and sex to determine *Egr1* expression in the PVN. **(F,G)** Some IEGs

were strongly influenced by pubertal stress, like *Fos*. **(F)** In Control mice, females and

males show little change during the PN21-PN35 age window. **(F,G)** However, pubertal

stress led to a significant decrease in *Fos* expression at both PN28 and PN35 in female

and male mice. CVS = chronic variable stress, IEG = immediate early gene, PN =

postnatal day, PVN = paraventricular nucleus of the PVN. (B-C) *$p < 0.05$ for non-zero

slope following linear regression; #$p < 0.05$ for significantly different slopes between

groups but within gene (D-G) *p < 0.05 for main effect of CVS or for interaction terms

following 3-way ANOVA analysis.

Figure 4. Schematic exemplifying our findings on the influence of pubertal stress

on the transcriptome of the developing and adult hypothalamus. Through prior and

current findings, we have demonstrated that a set of immediate early genes in the

paraventricular nucleus of the hypothalamus is sensitive to pubertal stress (chronic

variable stress from postnatal days 21-34) both during the stress exposure and in

adulthood. Of note, all IEG expression differences were measured in non-stimulated,

baseline conditions, and so may represent how the PVN is poised to respond to stimuli.

Given here is the example of *Fos*, which is decreased during the pubertal stress window

relative to Controls. Whether adult mice are pregnant or are exposed to

allopregnanolone, there is an increase in baseline IEG expression. This suggests that

the PVN is poised to respond differently, which we have observed in the form of a

blunted physiological response to restraint in adulthood. The new findings that baseline

IEG expression is altered provide support for a role of IEGs in the mechanism

underlying the lasting effect of pubertal stress on the PVN.