Astrocyte aquaporin mediates a tonic water efflux maintaining brain

homeostasis

Cuong Pham[1], Yuji Komaki[2], Anna Deàs-Just[1], Benjamin Le Gac[1], Christine Mouffle[1], Clara

Franco[1], Agnès Chaperon[1], Vincent Vialou[1], Tomokazu Tsurugizawa[3,4,#], Bruno Cauli[1,#],

Dongdong Li[1,#]

[1]Sorbonne Université, Institute of Biology Paris Seine, Neuroscience Paris Seine, CNRS

UMR8246, INSERM U1130, Paris 75005, France

[2]Central Institute for Experimental Animals, Kawasaki, Japan

[3]Human Informatics and Interaction Research Institute, National Institute of Advanced Industrial

Science and Technology (AIST), Tsukuba, Japan

[4]Faculty of Engineering, University of Tsukuba, Tsukuba, Japan

[#]Correspondence: t-tsurugizawa@aist.go.jp, bruno.cauli@upmc.fr, dongdong.li@inserm.fr

Keywords: AQP4, imaging, swelling, diffusion

1 ABSTRACT

Brain water homeostasis not only provides a physical protection, but also determines the diffusion

of chemical molecules key for information processing and metabolic stability. As a major type of

glia in brain parenchyma, astrocytes are the dominant cell type expressing aquaporin water channel.

How astrocyte aquaporin contributes to brain water homeostasis in basal physiology remains to be

understood. We report that astrocyte aquaporin 4 (AQP4) mediates a tonic water efflux in basal

conditions. Acute inhibition of astrocyte AQP4 leads to intracellular water accumulation as

optically resolved by fluorescence-translated imaging in acute brain slices, and *in vivo* by fiber

photometry in mobile mice. We then show that aquaporin-mediated constant water efflux

maintains astrocyte volume and osmotic equilibrium, astrocyte and neuron Ca^{2+} signaling, and

extracellular space remodeling during optogenetically induced cortical spreading depression.

Using diffusion-weighted magnetic resonance imaging (DW-MRI), we observed that *in vivo*

inhibition of AQP4 water efflux heterogeneously disturbs brain water homeostasis in a region-

dependent manner. Our data suggest that astrocyte aquaporin, though bidirectional in nature,

mediates a tonic water outflow to sustain cellular and environmental equilibrium in brain

parenchyma.

Significance statement - Our brain is immersed, thus protected, in a water environment. It ensures

intra- and extracellular molecular diffusion, which is vital for brain function and health. Brain

water homeostasis is maintained by dynamic water transport between different cell types.

Astrocytes are a main type of glial cell widely distributed in brain parenchyma, expressing the

bidirectional aquaporin water channel. Here we show that in basal conditions, aquaporin channel

mediates a tonic water efflux from astrocytes. This mechanism maintains astrocyte volume

stability, activity-gated brain parenchyma remodeling and brain water homeostasis. Our finding

sheds light on how astrocytes regulate water states in the brain, and will help to understand brain

allostasis in specific life contexts.

1 Introduction

Every aspect of brain function relies on the delicately maintained water environment. It supports

brain structural stability and molecular diffusion, laying the ground for information processing,

metabolite shuttling and adaptation to living environments (Kimelberg, 2004). Water is the

fundamental constituent of the cerebrospinal fluid infiltrating into the central nervous system and

the interstitial fluid distributed in brain parenchyma (Agnati et al., 2017; Brinker et al., 2014).

Brain fluid transport is suggested to support the diffusion of energetic fuels like glucose and lactate

to warrant the metabolic circumstances in the parenchyma, and is also implicated in the clearance

of waste molecules from the brain as described in the glymphatic system (Abbott et al., 2018; Iliff

et al., 2013; Iliff et al., 2012; Mestre et al., 2018a; Rasmussen et al., 2022). In addition, the

restricted mobility of water molecules diffusion within intra- and extracellular space reflects the

brain microstructure. The diffusion-weighted magnetic resonance imaging (DW-MRI) detects the

mobility of water proton for structural and functional neuroimaging (Le Bihan et al., 2006;

Tsurugizawa et al., 2013), being adopted for clinical diagnostics in neurological diseases such as

ischemia, brain tumor and edema (Gaddamanugu et al., 2022; Le Bihan and Iima, 2015) and

recently leveraged for assessing brain glymphatic system (Giannetto et al., 2024; Gomolka et al.,

2023).

Water equilibrium in the brain is maintained by dynamic transport between different cell

entities. Aquaporin is a family of transmembrane channels facilitating bidirectional water flow,

with aquaporin 4 (AQP4) being the main subtype expressed in the central nervous system. In the

brain, AQP4 is expressed in astrocytes (Papadopoulos and Verkman, 2013; Xiao et al., 2023) that

are a major type of glial cell in the parenchyma (Barres, 2008). This feature enables astrocytes to

well balance osmotic oscillations imposed by the transmembrane transport of ions, metabolites,

and signaling molecules, all mediated by water. AQP4 has been proposed to maintain the

circulation of cerebrospinal fluid (CSF), underlying the clearance efficacy of glymphatic system

(Hablitz et al., 2020; Kress et al., 2014; Mestre et al., 2018a; Smith et al., 2017). Dysregulation in

astrocyte aquaporin water transport is implicated in pathological scenarios such as cerebral edema

and neuroinflammation (Verkman et al., 2011). Valuable insights have been obtained on the

function of astrocyte aquaporin in specific physiopathological conditions including osmotically

evoked volume responses, ischemia and neurodegenerative diseases (Verkman et al., 2017).

Meanwhile, how astrocyte aquaporin contributes to brain water homeostasis in basal physiology

has remained elusive. A better understanding will aid to discern astrocytic regulation of cerebral

water states, thereby the potential maladaptations in neurological disorders.

Here, combining *in vivo* chemical targeting, optical and MRI imaging of water dynamics,

we report that AQP4 sustains a tonic water efflux from brain astrocytes. This mechanism was

found to be critical in maintaining astrocyte volume and signaling stability, as well as the

extracellular space remodeling in mouse cortex during optogenetically induced cortical spreading

depression. Using DW-MRI, we observed that acute inhibition of astrocyte water efflux *in vivo*

heterogeneously alters water diffusion across brain regions. Our finding suggests that aquaporin

acts as a water export route in astrocytes in physiological conditions, contributing to maintain

cellular and parenchymal water equilibrium.

Results

Astrocyte aquaporin mediates a tonic water efflux

To optically follow astrocyte water transport *in situ*, we performed real-time imaging of

sulforhodamine B (SRB)-labeled astrocytes, whose fluorescence changes reflect real-time volume

oscillation due to transmembrane water flux. We chemically labeled mouse brain astrocytes *in vivo*

with a highly water-diffusible and astrocyte-specific red fluorescent dye, sulforhodamine B (SRB)

(Figure 1A, *top*) (Appaix et al., 2012; Pham et al., 2020). About one hour after its intraperitoneal

injection, astrocytic SRB labeling was widely distributed in the mouse cortex, as seen in living

acute brain slices (Fig. 1A, *middle*) and validated by colocalization with EGFP-identified

astrocytes in slices taken from GFAP-EGFP transgenic mice (Fig. 1A, *bottom*). Net water transport

across the astrocyte membrane alters cytoplasmic SRB concentration and cellular volume, which

can thus be followed by changes in fluorescence intensity when imaged in a fixed field of view.

To validate this fluorescence intensity-translated (FIT) imaging of astrocyte water transport, we

employed wide-field fluorescence microscopy for single-plane time-lapse imaging in acute brain

slices in the primary somatosensory (S1) cortex. This approach allowed us to collect fluorescence

from both the focal plane and along the axial extension, thereby imprinting volumetric

fluorescence into the single image plane. Indeed, water influx induced by the application of

hypoosmotic solution over different periods decreased astrocyte SRB fluorescence, concomitantly

reflecting the cell swelling (Fig. 1B, *top*); whereas water export and astrocyte shrinking upon

hyperosmotic manipulation increased astrocyte fluorescence (Fig. 1B, *bottom*). Hence, by

following the fluorescence change of SRB-labeled astrocytes, the FIT imaging strategy allows to

monitor the real-time changes in astrocyte water transport and volume change.

In basal conditions, flat fluorescence time course was recorded in brain slices suggesting

equilibrated water transport across the astrocyte membrane (Fig. 1B) and volume homeostasis in

the brain parenchyma. To inspect a tonic role of aquaporin in astrocyte water transport, we sought

to acutely block astrocytic aquaporin channel, aquaporin 4 (AQP4). We used the synthetic

compound 2-(nicotinamido)-1,3,4-thiadiazole (TGN-020) that is derived from the condensation of

nicotinamide and thiadiazole derivatives (Burnett et al., 2015; Huber et al., 2009; Igarashi et al.,

2011; Sun et al., 2022), whose specificity for AQP4 has been validated *in vitro* by ion channel

heterologous expression system (Toft-Bertelsen et al., 2021) and *in vivo* using the AQP4 knockout

(KO) mice (Harrison et al., 2020; Igarashi et al., 2013). This approach permitted the functional

pinpointing of astrocyte aquaporin under physiological conditions, while avoiding the chronic

compensations caused by genetic tools and mouse models that have been reported to alter brain

water content, volume and extracellular architecture potentially confounding the functional

readouts (Binder et al., 2004; Gomolka et al., 2023; Haj-Yasein et al., 2011; MacAulay, 2021; Yao

et al., 2015).

AQP4 is a bidirectional channel providing a path of least resistance for water transfer along

the osmotic gradient (Papadopoulos and Verkman, 2013). As an osmotic equilibrium is maintained

in the brain parenchyma in basal state, there might not be net water transport across AQP4.

However, we observed in cortical slices that while little impact was seen with vehicle (CTR) on

the tendency of fluorescence time courses of SRB-labeled astrocytes (Fig. 1C, *left and middle*),

acute inhibition of astrocyte AQP4 by TGN-020 caused a gradual decline, suggesting a progressive

intracellular water accumulation (Fig. 1C, *left and middle*). This observation implies that in basal

condition, astrocyte aquaporin mediates a constant water efflux; its blocking causes intracellular

water accumulation and swelling (Fig. 1C, *right*). To corroborate this observation *in vivo*, we

performed fiber photometry recording in moving mice (Fig. 1D, *left*). An optical fiber was

implanted into the mouse S1 cortex to follow fluorescence signals of local astrocyte population

post the intraperitoneal injection of SRB (Fig. 1D, *left*). After about one hour, when astrocytes

were labeled, either saline or TGN-020 was injected intraperitoneally. Although no significant

effect was observed with the saline control, TGN-020 induced a decrease in astrocyte SRB

fluorescence and a reduction in its oscillation range (Fig. 1D, *right*), mirroring an intracellular

dilution of fluorescent molecules. Together, these observations suggest that acute inhibition of

astrocyte aquaporin leads to intracellular water accumulation thereby cell swelling.

Aquaporin inhibition perturbs astrocyte and neuron signaling

Astrocyte volume equilibrium not only determines brain structural stability, but also associates

with dynamic cellular signals. Astrocyte swelling has been shown to alter intracellular Ca^{2+}

signaling (Benfenati et al., 2011; Eilert-Olsen et al., 2019). We performed Ca^{2+} imaging in acute

brain slices to confirm that acute aquaporin inhibition induces astrocyte swelling and then Ca^{2+}

oscillation. The genetically encoded Ca^{2+} sensor GCaMP6f was selectively expressed in astrocytes

by crossing the Glast-Cre^{ERT2} and GCaMP6^{floxP} mouse lines (Herrera Moro Chao et al., 2022;

Pham et al., 2020) (Fig. 2A, *top*). Because astrocyte Ca^{2+} signals often occur in local domains, we

adapted a light sheet microscope (Pham et al., 2020) for wide-field optical sectioning so as to

image them with high signal-to-noise ratio (Fig. S1). We first confirmed that water accumulation,

therefore astrocyte swelling, induced by hypotonic solution enhanced the intracellular Ca^{2+} signal

(Fig. 2A). Then we followed astrocyte Ca^{2+} in isotonic control condition while blocking astrocyte

AQP4 with TGN-020, and observed that Ca^{2+} signaling was augmented following the acute

aquaporin inhibition (Fig. 2B). The spontaneous Ca^{2+} signals in perivascular astrocyte end-feet,

which wrapped the blood vessels displaying as tube-like structures, showed relatively higher

sensitivity to the acute application of TGN-020, consistent with the enrichment of AQP4 in the

end feet (Fig. S2). This result confirms that inhibiting astrocyte tonic water efflux would have led

to intracellular water accumulation and swelling, thus altering the astrocytic Ca^{2+} signaling. In

addition, astrocyte swelling has been reported to induce the release of neuroactive molecules such

as glutamate thereby influencing nearby neuron activity (Fiacco et al., 2007; Yang et al., 2019).

Disrupting AQP4 tonic water outflow may not only cause astrocyte swelling but also influence the

activity of neighboring neurons. We then imaged Ca^{2+} signals as a surrogate for neuronal activity

in cortical somatostatin (SST) interneurons which exhibit high excitability (Karagiannis et al.,

2021) and express both ionotropic and metabotropic glutamate receptors (Cauli et al., 2000),

rendering them ideal to sense astrocyte-released signaling molecules. We expressed GCaMP6f in

SST interneurons by crossing homozygous SST-Cre (Taniguchi et al., 2011) and homozygous

GCaMP6^{floxP} mice. TGN-020 inhibition of astrocyte AQP4 led to a global Ca^{2+} elevation in SST

interneuron populations (Fig. 2C). These observations suggest that astrocyte aquaporin mediates a

tonic water efflux, contributing to maintain both the volume and signaling homeostasis.

Aquaporin water efflux regulates astrocyte volume response

We then examined the role of aquaporin water efflux in astrocyte volume response to osmotic

changes. The AQP4 blocker TGN-020 or equimolar vehicle was applied throughout the recording,

namely being continuously present before and after osmotic challenges. We first followed the

evoked water efflux and shrinking induced by hypertonic solution. As astrocyte AQP4 supports

preferentially water efflux, its inhibition would attenuate hypertonicity-imposed water extrusion

(Fig. 3A). Supporting this, application of TGN-020 slowed down the hypertonicity-induced water

efflux and shrinking, reflected by the longer time to peak in the SRB fluorescence time course as

compared to control (Fig. 3B, *left*), though the delay in the initial onset was not significantly

prolonged (Fig. 3B, *right*). The maximal increase in astrocyte SRB fluorescence was lower in the

presence of TGN-020, suggesting that AQP4 blocking reduced the overall amount of water efflux

(Fig. 3B, *right*).

We next evoked water influx, thereby astrocyte swelling, by hypotonic solution in the

presence or absence of TGN-020 (Fig. 3C). In contrast to the effects on hypertonicity-evoked water

efflux, AQP4 acute inhibition was observed to accelerate both the initial water accumulation and

the swelling rate in astrocytes, reflected by their earlier onset in SRB fluorescence decrease and

faster reaching to the plateau value relative to the control condition (Fig. 3D). This observation

cross-validates the role of astrocyte aquaporin in supporting water efflux, whereby its blockade

facilitates the initial accumulation of the evoked water influx (Fig. 3C). In contrast, the late-stage

maximum decrease in astrocyte SRB fluorescence was observed to be reduced in the presence of

TGN-020 (Fig. 3D, *right*), reflecting a reduction in the total amount of water accumulation and

swelling. As TGN-020 was present prior to hypotonic challenge, it would have slightly swelled

astrocytes due to the blockade of tonic water efflux, thereby constraining the range of further

swelling induced by subsequent hypotonicity. In addition, while astrocyte AQP4 sustains a water

outflow in basal physiology, in conditions when transmembrane water gradient is altered as

occurred in hypotonic solutions, the net water flow through AQP4 should be finally dictated by

the osmotic gradient due to its bidirectional nature. Therefore, along the imposed hypotonicity,

AQP4 started to instruct a water influx; its constant blocking by TGN-020 would have reduced the

total amount of water influx thereby the maximal extent of swelling.

Astrocytes are widely distributed throughout the brain parenchyma, orchestrating water

transport and volume homeostasis (Ochoa-de la Paz and Gulias-Canizo, 2022). Water outflow via

astrocyte AQP4 may play a role in the structural remodeling of parenchyma at the global level.

We then induced general cell swelling and changes in extracellular spaces by triggering cortical

spreading depression (CSD), a widespread depolarization implicated in brain physiopathology

(Chung et al., 2019; Holthoff and Witte, 1996; Zhao et al., 2019). To be orthogonal to the

pharmacological control of astrocyte aquaporin, the CSD depolarization wave was initiated by

optogenetically stimulating ChR2-expressing glutamatergic pyramidal cells in the cortical slices

of Emx1-Cre::Ai32ChR2 mice (Gorski et al., 2002; Madisen et al., 2012), termed Opto-CSD. As

cell swelling increases the transmittance of infrared light, we imaged Opto-CSD by the intrinsic

optical signal (IOS) derived from infrared illumination (Holthoff and Witte, 1996; MacVicar and

Hochman, 1991), which was spectrally distinct from ChR2 photoactivation (detailed in Methods).

The IOS displayed transient increase across cortical layers during the Opto-CSD (Fig. 4A-C),

exhibiting propagating kinetics characteristic of spreading depression (Chung et al., 2019; Holthoff

and Witte, 1996). The biphasic starting of IOS coincided with a sharp response in the extracellular

potential indicating the CSD initiation (Fig. S3A). IOS signal contains a first peak reflecting the

rapid CSD response (Fig. 4C, *b*) followed by a prolonged phase of general cellular swelling (Fig.

4C, *c*). By combining fluorescence imaging of SRB-labelled astrocytes, whose spectrum is

separated from the IOS infrared signal, we observed that astrocytes swelling (i.e., decrease in SRB

fluorescence) paralleled CSD swelling (Fig. S4). Consistent with our observation on astrocyte

volume response (Fig. 3D), when pre-incubating slices with TGN-020 to block AQP4 water

outflow, the initiation of both the CSD and general swelling was accelerated while their maximum

amplitude reduced (Fig. 4D-E; Fig. S3B). Indeed, TGN-020 also increased the speed and the

duration of the swelling while prolonging the recovery time from the swelling, confirming acute

inhibition of AQP4 water efflux facilitates astrocyte swelling while restrains shrinking (Fig. S5).

Blocking action potentials with tetrodotoxin (TTX) only reduced the amplitude of the initial CSD

response while the effect on general swelling is insignificant (Fig. 4D-E). We further followed the

swelling of individual SRB-labelled astrocytes during Opto-CSD. In consistence with the result

obtained from hypotonic challenge (Fig. 3D, *right*), the presence of TGN-020 reduced the peak

amplitude of astrocyte swelling (i.e., the maximal SRB fluorescence decrease; Fig. 4F). As a post

hoc adaptation to the transiently induced Opto-CSD, astrocyte swelling was followed by a

shrinking to recover cellular volume to the homeostatic level, which was recorded as a rebound in

SRB fluorescence intensity. We noted that TGN-020 presence significantly restrained the recovery

of SRB fluorescence back to the baseline (Fig. 4F, *bottom*), showing that AQP4 inhibition hindered

astrocyte water efflux therefore the shrinking.

Collectively, our data show that astrocyte AQP4 sustains a tonic water outflow regulating

the cellular volume response and the general cell swelling of the parenchyma.

Tonic astrocyte water transport underlies brain homeostasis

Water homeostasis sets the basis for molecular diffusion in the brain, which instructs

neurotransmitter availability, ion recycling and metabolite trafficking. We then examined *in vivo*

the role of astrocyte aquaporin outflow in brain water diffusion using DW-MRI (Le Bihan and

Iima, 2015). It uses the diffusion of water molecules to generate contrast that can be quantified by

the apparent diffusion coefficient (ADC) in the nerve tissue (Beaulieu, 2002; Le Bihan, 2014). We

used a 7T MRI to image global brain water diffusion in mice lightly anesthetized with

medetomidine that normalized the animals under the sedative state. Astrocyte AQP4 outflow was

acutely perturbed by TGN-020 applied via intraperitoneal injection. Brain water diffusion was

mapped every 5 min before and after TGN or saline (as control) administration (Fig. 5A). Each

acquisition sweep was performed with three b values (0, 250, 1800 s/mm²) that accounted for ADC,

so to fit the exponential curve (Fig. 5B). The representative images reveal the enough image quality

to calculate the ADC, which allowed us to examine the effect of TGN-020 on water diffusion rate

in multiple regions (Fig. 5C). The temporal features in water diffusion change appeared also

different: an early elevation followed by a reversible tendency was observed for cortical areas, a

rapid and long-lasting elevation for the striatum while a delayed and transient increase for the

hippocampus (Fig. 5D, *top*). The respiratory rate was not changed by the injection of TGN020 and

saline (Fig. S6). We assessed the time course of ADC change with three phantoms: distilled water,

n-undecane, and n-dodecane. Both n-undecane and n-dodecane have been used as phantom for

diffusion-weighted image (DWI) (Tofts et al., 2000; Wu and Alexander, 2007). The ADC was

fluctuated within 2% of the averaged ADC at basal period (Fig. S7), indicating that the ADC

changes induced by TGN-020 is over the noise level. The *in vivo* neuroimaging results confirm

that the tonic water efflux from astrocyte aquaporin contributes to maintain the homeostasis of

brain water diffusion, and also suggest spatiotemporal heterogeneities in brain water handling.

Discussion

Water equilibrium underlies brain function, plasticity and dynamics. Astrocytes are the principal

brain cell type expressing aquaporin water channels that are highly implicated in regulating local

environments in the parenchyma (Xiao et al., 2023). We show that albeit being a bidirectional

channel, astrocyte aquaporin sustains a tonic water outflow in basal states to ensure structural and

functional stability, as well as the water homeostasis at the brain level. We have used the chemical

dye SRB to label astrocytes *in vivo*, so as to follow the transmembrane water transport and volume

dynamics by real-time fluorescence changes. This sensitivity is ensured by its far lower molecular

weight (MW, 558.7 Da), thereby high solubility in the cytoplasm, as compared with fluorescence

proteins such as the GFP (MW, 26.9 kDa). In addition, the chemical labeling enables genetics-less

targeting of brain astrocytes, so to avoid protein overexpression that may impact astrocyte volume

and water transport responses.

The observation of a basal aquaporin water efflux implies there is a constitutive water

accumulation in brain astrocytes. As a ubiquitous vehicle for transporting ions, transmitters and

metabolites, water enters astrocytes via a wide range of ion channels, transporters and exchangers.

For instance, they express Na^+/K^+ and Na^+/HCO_3^- cotransporters to dynamically regulate intra-

and extracellular ion homeostasis (MacAulay, 2021). Standing as a major glial cell type controlling

the neuropil environment, astrocytes take up synaptically released K^+ through inwardly rectified

K^+ channels and neurotransmitters (e.g., glutamate, GABA) via high-affinity transporters to

safeguard synaptic transmission (Dallerac et al., 2018). The cotransport of water into the cells with

ions or metabolites would lead to water buildup in astrocytes. Moreover, astrocytes juxtapose the

cerebral vasculature, being a front relay station for brain metabolism. They express glucose

transporters and lactate-permeable monocarboxylate transporters facilitating energy substrate

uptake and transfer between blood vessels and neuron-glia networks (Cauli et al., 2023). As water

is constantly produced during metabolic processes (Bonvento and Bolanos, 2021), astrocyte

metabolism would also contribute to the constitutive water accumulation in the cytoplasm. An

efficient efflux pathway appears necessary to counterbalance the constitutive water buildup,

thereby maintaining astrocyte and brain homeostasis. Our data suggest that astrocyte AQP4 may

fulfill such a role by mediating a tonic water outflow (Fig. S8), providing functional evidence to

the previous suggestion (Amiry-Moghaddam et al., 2003). It has been observed that knock out of

AQP4 leads to water over accumulation in the brain, likely due to the reduction in water exit from

the pial membrane (Haj-Yasein et al., 2011; Vindedal et al., 2016). Our result also echoes that in the principal cells of the kidney collecting duct, AQP4 is suggested to mainly mediate water exit to balance the AQP2-sustained water entry (Noda et al., 2010). Moreover, we observed that inhibiting astrocyte basal water efflux alters Ca^{2+} activities in both astrocytes and neurons, reminiscent of the early finding that inhibition of AQP4 by TGN-020 enhances cerebral blood flow in mouse cortex (Igarashi et al., 2013), which might be due to the enhanced neuron-astrocyte activity tone. As a pharmacological compound, TGN-020 exerts a partial blocking effect on AQP4 (Huber et al., 2009), implying that the actual functional impact of AQP4 per se might be stronger than what we currently observed. Alternatively, the phenylbenzamide AER-270 and its water-soluble prodrug AER-271 have been reported as AQP4 blocker (Farr et al., 2019) and impairs brain water diffusion and glymphatic fluid (Giannetto et al., 2024). AER270(271) has been noted to also be the inhibitor for κB nuclear factor (NF-κB) whose inhibition could reduce CNS water content and influence brain fluid dynamics (e.g., the ADC of DW-MRI) in an AQP4-independent manner (Giannetto et al., 2024; Salman et al., 2022). The inhibition efficiency of AER-270 seems also lower than that of TGN-020 (Huber et al., 2009; Giannetto et al., 2024; Salman et al., 2022). Developing pharmacological compounds that can more specifically and efficiently target AQP4 will facilitate the functional and pre-clinical investigations.

This study also provides mechanistic hints to understand AQP4-relevant pathologies in specific contexts. For instance, astrocyte water accumulation and swelling are implicated in the development of brain edema. Different types of edema have been delineated depending on the pathological contexts, including such as cytotoxic, vasogenic and hydrocephalic subtypes. The presence of AQP4 has been found to ameliorate vasogenic (vascular leak) and hydrocephalic edema (Jeon et al., 2021; Tait et al., 2010), where the excessive water has been suggested to exit

the brain via AQP4-dependent route. This goes with our observation that astrocyte AQP4

contributes to basal water extrusion. Hence aquaporin facilitators may help to control vasogenic

and hydrocephalic edema. Differently, inhibiting AQP4 has been observed to ameliorate cytotoxic

edema encountered during cerebral ischemia (Verkman et al., 2011). Beneficial effects of TGN-

020 on edema have been reported in spinal cord injuries (Li et al., 2019), in ischemia (Sun et al.,

2022) and in retinal edema in diabetic animal models (Oosuka et al., 2020). These observations

suggest that the role of AQP4 in pathological conditions is context-dependent. Indeed, while our

results suggest astrocyte AQP4 mediates a water outflow in basal states, the direction of water

transport in specific contexts will be determined by the actual transmembrane water gradient. As

the origin of edema is variable and water is among the many other ions involved in edema

formation (Stokum et al., 2016), astrocyte aquaporin may either be the primary cause for cytosolic

water accumulation or secondarily sustain protective water extrusion. In neuromyelitis optica

spectrum disorders (NMOSD), an unusual nervous autoimmune disorder often featured with

circulating IgG autoantibody against AQP4, the impairment in astrocyte water homeostasis has

been observed (Hinson et al., 2012; Lucchinetti et al., 2014). The inhibition of astrocyte AQP4 by

autoantibody may initially affect the basal water efflux. Yet chronic pathological evolution could

convert astrocytes into reactive states, with either or both the aquaporin expression, intra- and

extracellular environments being altered, the role of AQP4 in water transport might be accordingly

shifted (Mireles-Ramirez et al., 2022). Additionally, pro-inflammatory cytokines and chemokines

during the immunological maladaptation could also contribute to shape astrocyte functions

including aquaporin-mediated water transport (Lucchinetti et al., 2014; Mireles-Ramirez et al.,

2022). Recent data show that TGN-020 targeting of AQP4 mitigates the inflammation during

ischemia-reperfusion injury, by enhancing glymphatic amyloid clearance and inhibiting the ERK

1/2 pathway (Li et al., 2023), suggesting that in pathological conditions, multiple factors could be recruited by astrocyte aquaporin targeting.

We observed that acutely blocking aquaporin caused basal water accumulation leading to swelling in astrocytes, and facilitated the initial occurrence of the evoked swelling by hypoosmoticity and during cortical spreading depression. This recalls early observation by electron microscopy that knocking out the anchoring protein alpha-syntrophin to disperse AQP4 clustering induces an enlargement of astrocyte end feet (Amiry-Moghaddam et al., 2003). It likely reflects astrocyte local swelling caused by the disruption of AQP4-mediated water efflux. Our observation also parallels the report that AQP4 knock out facilitates astrocytes swelling induced by hypoosmotic solution (Murphy et al., 2017). Nevertheless, genetic inactivation of AQP4 also reveals inhibitory effects on astrocyte swelling (Benfenati et al., 2011; Mola et al., 2016; Woo et al., 2018). Such discrepancy might be due to the variable compensations in brain water content and structure integrity during chronic genetic manipulations (Binder et al., 2004; Haj-Yasein et al., 2011; MacAulay, 2021; Yao et al., 2015). Using the calcein quenching assay and AQP4 KO (Solenov et al., 2004), it has been demonstrated in cultured astroctyes that AQP4 is a functional water channel. AQP4 deletion reduced both astrocyte water permeability and the absolute amplitude of swelling over comparable time, and also slowed down cell shrinking, which are overall confirmed by in our results from acute AQP4 blocking. Yet in the study (Solenov et al., 2004), the time to swelling plateau was prolonged in AQP4 KO astrocytes, which was observed to be accelerated in the present study. This difference may be due to compensatory mechanisms in chronic AQP4 KO, or reflect the different volume responses in cultured astrocytes from brain slices or *in vivo* contexts as noted previously (Risher et al., 2009). Nevertheless, in the present study, astrocyte volume and water transport dynamics have been inferred from the real-time

fluorescence change of SRB labeling. Rapid three-dimensional imaging of astrocyte volume

dynamics *in situ* will provide further quantitative information on the role of AQP4 in volume

regulations (MacAulay, 2021; MacAulay et al., 2004). In addition, though our data suggest that

under basal condition astrocyte AQP4 sustains a tonic water outflow, in conditions when massive

water influx persists as exemplified by the application of hypoosmotic solutions, the direction of

water transport of AQP4 would follow the imposed water gradient. Regulatory volume decrease

has been observed in cultured astrocyte volume responses behaving as a protective mechanism

(Mola et al., 2016). In current condition in acute brain slices, when we applied hypotonic solution

to induce astrocyte swelling, our protocol did not reveal rapid regulatory volume decrease. During

opto-CSD, we observed astrocyte volume decrease following the transient swelling (Fig. 4F),

where both the peak amplitude and the degree of recovery were reduced by inhibiting AQP4 with

TGN-020. These observations suggest astrocyte regulatory volume decrease likely to be condition-

dependent, while also recalling that the regulatory volume decrease is barely detectable in brain

slices or *in vivo* during hypo-osmotic challenge (Risher et al., 2009). Given the implication of CSD

in brain migraine aura, ischemia and seizure (Chung et al., 2019), the current observation suggests

a functional involvement of astrocyte AQP4 in the physiopathological adaptations.

Astrocyte aquaporin modulates brain water transport (MacAulay, 2021), though its role in

the glymphatic system is under deliberation (Mestre et al., 2018a; Rasmussen et al., 2022; Smith

and Verkman, 2019; Smith et al., 2017). Our data suggest that astrocyte aquaporin-mediated

outflow helps to maintain local water environment in the brain parenchyma, facilitating the

circulation of the interstitial and cerebrospinal fluid. Moreover, besides affecting water transport,

targeting AQP4 also impacts astrocyte volume therefore the size of extracellular space. These two

factors would need to be considered when evaluating AQP4 involvement in brain fluid transport.

We show by DW-MRI that impairing astrocyte water transport induces the changes of

ADC across the brain. When blocking AQP4 with TGN-020, therefore its basal water outflow, we

observed spatially heterogeneous elevations in brain water diffusion rate. It has been reported that

relative to wild-type mice, AQP4 KO mice show increased water diffusion along with the enlarged

interstitial space, brain volume yet reduced CSF spaces (Gomolka et al., 2023). Notably in AQP4

KO mice, the influx of gadolinium CSF tracer was reduced as followed by dynamic contrast-

enhanced MRI (Gomolka et al., 2023). As corroboration, the AQP4 blocker AER-271 also reduces

glymphatic water influx (Giannetto et al., 2024). Together, these observations suggest that AQP4

plays a critical role not only in modulating water diffusion in brain parenchyma but also in the

CSF transport. Our current data are derived from 5 min apart acquisitions, providing information

over the early phase of AQP4 inhibition therefore extending our early report (Debacker et al.,

2020). The regional heterogeneity likely reflects the various levels of AQP4 expression; its relative

enrichment in the cerebral cortex (Gomolka et al., 2023; Hubbard et al., 2015; Mestre et al., 2018a)

corresponds to the pronounced effect observed here by DW-MRI. An overall increase in brain

water diffusion rate was observed when blocking AQP4 water efflux. This treatment causes water

accumulation inside astrocytes and their swelling, which reduces the extracellular space but

increases the intracellular space. Water diffusion would be enhanced inside astrocytes and

decreased in extracellular space, respectively. DW-MRI maps global water diffusion of both intra-

and extracellular space. Likely, a net increase in brain water diffusion may reflect its intracellular

increase exceeding the extracellular decrease. In addition, convective brain fluid flow has been

suggested to be present in the perivascular space (Smith and Verkman, 2019), a scenario extended

by the glymphatic system to the extracellular space of the parenchyma (Mestre et al., 2018b). In

this sense, transiently squeezing extracellular space might increase the rate of the water flow.

Additionally, a transient upregulation of other water-permeable pathways would increase the water

diffusion in compensating the AQP4 blocking. It has been shown that TGN-020 increases regional

cerebral blood flow in wild-type mice but not in AQP4 KO mice (Igarashi et al., 2013). The current

MRI data yet lack sufficient spatial resolution to delineate the spatial compartmentalization of

brain fluid flow. We here used anisotropic resolution and minimum b-values to earn the temporal

resolution. The phantom study supports the quality of DW-MRI data to calculate the time course

of ADC. To further use isotropic resolution and more b-values will help to validate our current

results.

This study sheds light on the mechanisms by which astrocyte aquaporin contributes to the

water environment in brain parenchyma, and will help to understand the processes underlying

brain homeostasis and adaptation to life conditions.

Materials and methods

Animals and acute brain slice preparation

All procedures using animals were carried out in strict accordance with French regulations (Rural Code R214/87 to R214/130) and conformed to the ethical recommendations of the European Economic Community (Directive 86/609/EEC) and the National Charter French on ethics in animal experimentation. All protocols were approved by the Charles Darwin ethics committee and submitted to the French Ministry of Education and Research (Approval 2015 061011367540 APAFIS#573-2015061011367547 v1).

For water transport and volume imaging, coronal slices comprising somatosensory cortex were acutely prepared from C57BL/6 mice of both sexes at ages of 4-6 weeks, unless otherwise indicated. Mice were deeply anesthetized by isoflurane (ISOVET, Piramal) evaporation in a closed plexiglass box. The brain was taken out and placed in a modified artificial cerebrospinal fluid (aCSF) for slicing (in mM: 30 NaCl, 4.5 KCl, 1.2 NaH2PO4, 1 MgCl2, 26 NaHCO3, 10 D-Glucose, and 194 sucrose) maintained at 4 °C during sectioning, where depolarizing ions (Na$^+$, Ca^{2+}) were reduced in attempt to lower the potential excitotoxicity during the tissue dissection while sucrose used to balance the osmolarity (Jiang et al., 2016). The brain was cut into 300-μm thick slices with a vibratome (Leica VT1200S). Brain slices were recovered in standard aCSF (mM: 124 NaCl, 4.5 KCl, 1.2 NaH2PO4, 1 MgCl2, 2 CaCl2, 26 NaHCO3, and 10 D-Glucose) at 37 °C for about 1 h, and the same aCSF was used for brain slice imaging at room temperature. For optogenetic cortical spreading depression (opto-CSD) experiments, 18 to 21 day-old mice were deeply anesthetized with 150 μL of isoflurane. After the decapitation, the brain was quickly removed and placed into ice-colded oxygenated aCSF containing the following (in mM): NaCl 125; NaHCO3 26; Sucrose 15; Glucose 10; KCl 2,5; CaCl2 2; NaH2PO4 1,25; MgCl2 1; kynurenic acid 1 (nonspecific

glutamate receptor antagonist, Sigma-Aldrich). Coronal slices from mouse somatosensory cortex

(300 μm thick) were cut with a vibratome (VT1000, Leica) and allowed to recover at room

temperature for at least 45 min in aCSF saturated with O_2/CO_2 (95%/5%).

***In vivo* chemical and genetic targeting of astrocytes**

Mice of similar age were used also for *in vivo* labeling of astrocytes, where the astrocyte-specific

red fluorescent dye sulforhodamine B (10 mg/ml, Sigma) was intraperitoneally injected into awake

mice at a dose of 10 μL/g. The genetically encoded Ca^{2+} indicator GCaMP6 was expressed in

astrocytes *in vivo* by crossing a Cre-dependent GCaMP6f mouse line (Ai95, The Jackson

Laboratory) with an inducible conditional mouse line Glast-Cre^{ERT2}, which expresses Cre

recombinase selectively in astrocytes upon tamoxifen injection (Slezak et al., 2007). Tamoxifen

(T5648, Sigma) was dissolved in 100 % ethanol (10 mg/250μL). The corn oil (Sigma) was

subsequently added at a 9:1 proportion relative to the ethanol-tamoxifen solution. The resultant

solution was then heated at 37°c for 15 min, vortexed, sonicated for 15 min to reach transparency,

before being aliquoted at 250 μL each containing 1 mg tamoxifen. Aliquots were stored at -20 °C

until use. Glast-Cre^{ERT2}::Ai95$^{GCaMP6f//WT}$ mice were injected with tamoxifen at ~3–4 weeks of age,

once a day (1 mg) for two consecutive days. Genotyping of the Cre-dependent GCaMP6f mouse

line (Ai95, The Jackson Lab) used the standard primers and polymerase chain reaction protocols

provided by the supplier. Genotyping of GLAST-Cre^{ERT2} mouse line was performed using the

primers for Cre recombinase (TK139/141) as reported (Slezak et al., 2007). For Ca^{2+} imaging in

somatostatin interneurons, homozygous mice B6J.Cg-Sst$^{tm2.1(cre)Zjh}$/MwarJ (Jackson Laboratory)

(Taniguchi et al., 2011) have been crossed with homozygous Ai95$^{GCaMP6f/GCaMP6f}$ mice to obtain

heterozygous   SST-Cre$^{cre/WT}$::Ai95$^{GCaMP6f//WT}$ mice.   For   opto-CSD,   homozygous   mice

B6;129PEmx1tm1lr/J ((Gorski et al., 2002), Emx1-Cre[cre/cre], Jackson Laboratory) have been

crossed with homozygous mice B6;129SGt(ROSA)26Sortm32.1(CAG-

COP4*H134R/EYFP)Hze/J ((Madisen et al., 2012) Ai32[ChR2/ChR2], Jackson Laboratory) to obtain

heterozygous Emx1-cre[cre/WT]::Ai32[ChR2/WT] mice.

***Ex vivo* fluorescence imaging**

Light sheet and epifluorescence imaging were both performed using a wide-field upright

microscope (Zeiss Axioskop 50, Germany) equipped with water-immersion objectives.

Epifluorescence illumination was provided by a monochromator light source (Polychrome II,

TILL Photonics, Germany) directly coupled to the imaging objective via an optical fiber. We used

water immersion imaging objectives (ZEISS): ×10 NA0.3 used to have a global view, ×20 NA0.5

used to perform light sheet imaging, and ×63 NA1.0 to image individual astrocytes under

epifluorescence configuration so as to collect their whole-cell SRB fluorescence to follow

astrocyte water transport and volume dynamics. The green fluorescence of GCaMP6 was separated

from the red fluorescence of SRB, by spectrally exclusive double-band filters (Di03-R488/561-t3

and FF01- 523/610, Semrock). SRB-labeled astrocytes were excited at 550 nm. Fluorescence

signal was collected using a digital electron-multiplying charge-coupled device (EMCCD Cascade

512B, Photometrics).

A light sheet imaging system was adapted from our previous publication (Pham et al.,

2020), where a chamber with a slightly tilted surface was used so as to spare the agarose embedding

and allow the light sheet to access to any parts of the brain slice (Fig. S1). Briefly, the in-focus

light sheet was introduced laterally from an independent optical module (Alpha3 light sheet add-

on, Phaseview, France) equipped with an air objective (Zeiss EC EPIPlan ×10, 0.25NA). This

optical module was orthogonal to the upright imaging pathway and coupled via a wavelength combiner (Thorlabs) to two continuous wave lasers 473-nm and 561-nm (CNI, China), for the excitation of GCaMP6 and SRB, respectively. The imaging chamber ($20 \times 30 \times 46$ mm width, height and length) was custom-designed using 3D modeling software (Trimble SketchUP), produced by a Mojo 3D printer (Stratasys) with the acrylonitrile butadiene styrene material, where the horizontal slice-supporting surface in the chamber was tilted by 8° along the laser entry direction. In this way, the brain slice could be laid on the slightly tilted surface while being stabilized with a conventional Harp slice anchor grid (Warner Instruments), whereby the laterally generated laser light sheet could reach local regions throughout the brain slices. The chamber was mounted on a motorized stage (PI-Physik Instrumente GmbH, Germany) for axial micro-manipulation. Laser excitation and image acquisition were controlled by MetaMorph (Molecular Devices).

Based on the time-lapse image stacks (1 Hz), the active regions of interest (ROIs) of astrocytes displaying dynamic Ca^{2+} signals were identified using our established spatio-temporal screening method (Pham et al., 2020). The signal strength of astrocyte Ca^{2+} signals was estimated from the temporal integral of the normalized Ca^{2+} time courses (dF/F_0, F_0 is the average of the lowest 20-data point baseline) for the periods prior and post to TGN or vehicle application respectively, which was then normalized per minute. As for neuronal Ca^{2+} signals, we followed the global changes with low-frequency imaging (1 Hz) and calculate the mean fluorescence of single 200 µm × 200 µm field of view. SRB fluorescence dynamics of individual astrocytes was analyzed with ImageJ software (NIH). We sought to follow the temporal changes in SRB fluorescence signal. The acquired fluorescent images contained not only the SRB signal, but also the background signals consisting of, for instance, the biological tissue auto fluorescence, digital

camera background noise and the leak light sources from environments. The value of the background signal was estimated by the mean fluorescence of peripheral cell-free subregions (15 $\times$ 15 µm²) and removed from all frames of single time-lapse image stacks. F_0 was identified as the mean value of the ten data points immediately preceding the detected fluorescence changes, and the same timing used when comparing the paired conditions. Because single astrocytes typically occupy an area of ~50-100 µm in radius (Appaix et al., 2012), regions of interest of 50-µm diameter were delineated surrounding identified astrocyte footprint, to comprise the extended area of individual astrocytes while minimizing interference with neighbors. Time courses extracted from single regions of interest were normalized as dF/F_0.

Dynamic imaging of the extracellular space and astrocytes volume changes

Individual slices were transferred to a submerged recording chamber and perfused with oxygenated aCSF (~2 ml/min) lacking kynurenic acid. Slices were observed using a double port upright microscope (BX51WI, Olympus, Japan) equipped with Dodt gradient contrast optics (DGC; Luigs and Neumann), collimated light emitting device (LED; 780 nm, Thorlabs) as transmitted light sources, ×4 objective (PlanN, 0.10NA, Olympus), ×20 (UMPlanFL N, 0.50 W, Olympus) or ×40 (LUMPlanF/IR, 0.80 w, Olympus) objectives, and a digital camera (OrcaFlash 4.0, Hamamatsu) attached on the front port of the microscope. The observation with ×4 allows to delimit the cortical layers and to confirm the presence of "barrels"-like patterns in layer IV (Erzurumlu and Gaspar, 2020). The ×20 objective was used to observe infrared intrinsic optical signals (IOS) (Holthoff and Witte, 1996; MacVicar and Hochman, 1991) of the cortical layers I to IV. The ×40 objective was used to image SRB labelled astrocytes combined with IOS in layers I and II/III.

After one minute of baseline period, a 10-s continuous photostimulation at 470 nm of the

$ChR2_{H134R}$ (30% of the maximal power of the LED) was carried out via the epifluorescence port

of the microscope using a collimated LED system (CoolLED, PreciseExcite) and a set of multiband

filters consisting of an excitation filter (HC 392/474/554/635, Semrock), a dichroic mirror (BS

409/493/573/652, Semrock) and an emission filter (HC 432/515/595/730, Semrock) to observe the

induced changes of light transmittance. The infrared and epifluorescence images were acquired

each at 1 Hz (exposure 100 ms, binning 2×2) using the Imaging Workbench 6.1 software (Indec

Byosystems) for a total duration of 15 min.

In vivo **fiber photometry**

The optical fiber (200 µm core, NA = 0.37, Neurophotometrics) coupled to a stainless-steel ferrule

(1.25 mm) was stereotaxically implanted in mouse S1 cortex (relative to bregma, rostro-caudal 0

mm, lateral ± 2.5 mm, vertical -1.2mm), and fixed to the skull with dental cement (SuperBond,

Sun medical). Mice were familiarized with the setup for 30 minutes prior to fiber photometry

recording in mobile states. 5-min baseline was recorded before SRB injection, then SRB was

applied to label brain astrocytes via intraperitoneal injection and the fluorescence time course was

continuously recorded. To examine *in vivo* the effect on astrocyte water transport of acute AQP4

blocking, either TGN-020 (200 mg/kg; Tocris or MedChemExpress) or saline control was

intraperitoneally injected 1 h after SRB injection, a time point when astrocytes were well labeled

as confirmed by brain slice imaging.

Fluorescence signals were recorded using a Neurophotometrics fiber photometry system

(FP3002). A branched fiber optic patch cord (BFP_200/230/900-0.37_FC-MF1.25, Doric Lenses)

connected to the fiber photometry apparatus was attached to the implanted fiber optic cannula

using a cubic zirconia sleeve. To record fluorescence signals from SRB, excitation light from 560-

nm LED was band-pass filtered, collimated, reflected by a dichroic mirror and focused by a ×20

objective. The excitation light power at the tip of the patch cord was 50 µW. Emitted fluorescence

was band-pass filtered and focused on digital camera. Signals were collected at a rate of 1 Hz for

SRB and visualized using the open-source software Bonsai 2.4 (http://bonsai-rx.org) and analyzed

in Igor software (Wavemetrics).

***In vivo* diffusion-weighted MRI**

To test the effect of astrocyte AQP4 acute inhibition on brain water homeostasis, we used

diffusion-weighted magnetic resonance imaging (DW-MRI) to calculate the apparent diffusion

coefficient (ADC) of tissue water molecules in the brain. MRI experiment was performed with

medetomidine anesthesia (0.6 mg/kg/h, i.v.). Two groups of mice (seven for each) were injected

with either the water-soluble TGN-020 sodium salt (Key Organics, LR-0041) or saline. For the

crossover test, the order of injection of TGN-020 or saline was randomly determined. The

experiment was performed 2 days, which separated 1 month for the recover. The respiration was

monitored, and the body temperature was maintained at 36 °C throughout the measurement.

17   The MRI experiments were conducted on a 7.0 tesla MRI system equipped with actively

shielded gradients at a maximum strength of 700 mT/m (Biospec; 70/16 Bruker BioSpin, Ettlingen,

Germany) and with a cryogenic quadrature radio frequency surface probe (Cryoprobe). We chose

the cryoprobe because the signal-to-noise ratio appeared better than volume coil. We also

confirmed to obtain good image quality in whole brain (Fig. 5B-C). After 10 min from the start of

the scanning, either TGN-020 (200 mg/kg body weight) or Saline was injected. Then, scanning

was continued for 60 min. The respiration rate was monitored and was confirmed within the range

of 80-150 /min throughout the experiment. Mice were injected with 200 mg/kg of TGN-020. We

used a standard diffusion-weighted spin echo EPI sequence, with the following parameters: spatial

resolution = $175 \times 175 \times 500 \ \mu m^3$/voxel, 3 b-values (b = 0, 250 and 1800 s/mm^2), 6 directions, 1

segment, echo time = 37.1 ms / repetition time = 5769 ms, bandwidth = 300 kHz, δ = 3 ms, Δ =

24 ms, scan time = 5 min. The DWI scan was continued for 70 min (total 14 scans). We used

minimum direction of motion probing gradient and anisotropic resolution to image the time course

of ADC changes following the injection of the TGN-020 (Debacker et al., 2020). Previous study

has used long diffusion time (> 20 ms) and long echo time (40 ms) to follow the mean diffusivity

(Aggarwal et al., 2020), supporting the suitability of our protocol to investigate the ADC. The

stability of DW-MRI was evaluated through 70-minute scans using three different phantoms: water,

n-undecane (Merck, Darmstadt, Germany), and n-dodecane (Merck, Darmstadt, Germany).

The motion correction of DWI data was performed using statistical parametric mapping

software (SPM12, Welcome Trust Center for Neuroimaging, UK). The template image and regions

of interest (ROIs) were then co-registered to the DWI data using SPM12. The ROIs of the cortex,

the hippocampus, and the striatum were depicted with reference to Allen mouse brain atlas

(https://scalablebrainatlas.incf.org/mouse/ABA12). The ADC at each time point was calculated

from DWI data with all b-values (b = 0, 250, and 1800 s/mm²) using DSIstudio (https://dsi-

studio.labsolver.org/). The percentage change in ADC was then calculated according to the

equation: $\Delta ADC_i = (ADC_i / ADC_2 - 1) \times 100$ (%). The ADC_i is the ADC at the i-th time point. We

define the averaged ADC at 1st and 2nd time-point as the baseline because saline injection was

performed after the 2nd DWI scan. The averaged ADC changes within ROIs were calculated using

a homemade program.

Statistics

Experimental data are expressed as mean ± standard error unless otherwise mentioned. The *t*-test

was performed for two-group comparisons and significant difference was determined by p values

less than 0.05. Mann-Whitney U test was used when the data were non-parametric deviating from

normal distributions. Bonferroni-Holm correction was used for multiple comparisons. Statistical

tests were carried out with Matlab (The MathWorks) and Statistica 6 (Statsoft). The significance

levels are shown in figures by $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Acknowledgements

We thank the animal and imaging facilities of the IBPS (Sorbonne Université, Paris, France). We thank Nathalie Rouach for the GFAP-EGFP mice, Hervé Le Corronc and Thomas Panier for the discussion and the 3D printing of the light sheet chamber, respectively. This work was supported by the Agence Nationale de la Recherche (ANR-17-CE37-0010-03; ANR-20-CE14-0025-02), Japanese society for promotion of science (JSPS) 2022 summer grant, Grant-in-Aid for Challenging Research (Exploratory) in Japan (grant number 21K19464), the i-Bio initiative grant and the Emergence program of Sorbonne Université, and the grant from the Fondation de France. CP was supported by a fellowship from France Alzheimer and BLG by a fellowship from the Fondation pour la Recherche sur Alzheimer.

Supplemental file

There are eight supplementary figures (Fig. S1-S8).

Competing interest

The authors declare no competing financial interests.

Data availability

The authors commit to share data, documentation, and code used in analysis.

References:

Abbott, N.J., Pizzo, M.E., Preston, J.E., Janigro, D., and Thorne, R.G. (2018). The role of brain barriers in
fluid movement in the CNS: is there a 'glymphatic' system? Acta Neuropathol *135*, 387-407.

Aggarwal, M., Smith, M.D., and Calabresi, P.A. (2020). Diffusion-time dependence of diffusional kurtosis
in the mouse brain. Magn Reson Med *84*, 1564-1578.

Agnati, L.F., Marcoli, M., Leo, G., Maura, G., and Guidolin, D. (2017). Homeostasis and the concept of
'interstitial fluids hierarchy': Relevance of cerebrospinal fluid sodium concentrations and brain
temperature control (Review). Int J Mol Med *39*, 487-497.

Amiry-Moghaddam, M., Otsuka, T., Hurn, P.D., Traystman, R.J., Haug, F.M., Froehner, S.C., Adams, M.E.,
Neely, J.D., Agre, P., Ottersen, O.P.*, et al.* (2003). An alpha-syntrophin-dependent pool of AQP4 in
astroglial end-feet confers bidirectional water flow between blood and brain. Proceedings of the
National Academy of Sciences of the United States of America *100*, 2106-2111.

Appaix, F., Girod, S., Boisseau, S., Romer, J., Vial, J.C., Albrieux, M., Maurin, M., Depaulis, A., Guillemain,
I., and van der Sanden, B. (2012). Specific in vivo staining of astrocytes in the whole brain after
intravenous injection of sulforhodamine dyes. PLoS One *7*, e35169.

Barres, B.A. (2008). The mystery and magic of glia: a perspective on their roles in health and disease.
Neuron *60*, 430-440.

Beaulieu, C. (2002). The basis of anisotropic water diffusion in the nervous system - a technical review.
NMR Biomed *15*, 435-455.

Benfenati, V., Caprini, M., Dovizio, M., Mylonakou, M.N., Ferroni, S., Ottersen, O.P., and Amiry-
Moghaddam, M. (2011). An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex
is essential for cell-volume control in astrocytes. Proceedings of the National Academy of Sciences of the
United States of America *108*, 2563-2568.

Binder, D.K., Papadopoulos, M.C., Haggie, P.M., and Verkman, A.S. (2004). In vivo measurement of brain
extracellular space diffusion by cortical surface photobleaching. J Neurosci *24*, 8049-8056.

Bonvento, G., and Bolanos, J.P. (2021). Astrocyte-neuron metabolic cooperation shapes brain activity.
Cell metabolism *33*, 1546-1564.

Brinker, T., Stopa, E., Morrison, J., and Klinge, P. (2014). A new look at cerebrospinal fluid circulation.
Fluids Barriers CNS *11*, 10.

Burnett, M.E., Johnston, H.M., and Green, K.N. (2015). Structural characterization of the aquaporin
inhibitor 2-nicotinamido-1,3,4-thiadiazole. Acta Crystallogr C Struct Chem *71*, 1074-1079.

Cauli, B., Dusart, I., and Li, D. (2023). Lactate as a determinant of neuronal excitability, neuroenergetics
and beyond. Neurobiol Dis *184*, 106207.

Cauli, B., Porter, J.T., Tsuzuki, K., Lambolez, B., Rossier, J., Quenet, B., and Audinat, E. (2000).
Classification of fusiform neocortical interneurons based on unsupervised clustering. Proceedings of the
National Academy of Sciences of the United States of America *97*, 6144-6149.

Chung, D.Y., Sadeghian, H., Qin, T., Lule, S., Lee, H., Karakaya, F., Goins, S., Oka, F., Yaseen, M.A.,
Houben, T.*, et al.* (2019). Determinants of Optogenetic Cortical Spreading Depolarizations. Cereb Cortex
29, 1150-1161.

Dallerac, G., Zapata, J., and Rouach, N. (2018). Versatile control of synaptic circuits by astrocytes: where,
when and how? Nat Rev Neurosci *19*, 729-743.

Debacker, C., Djemai, B., Ciobanu, L., Tsurugizawa, T., and Le Bihan, D. (2020). Diffusion MRI reveals in
vivo and non-invasively changes in astrocyte function induced by an aquaporin-4 inhibitor. PLoS One *15*,
e0229702.

Eilert-Olsen, M., Hjukse, J.B., Thoren, A.E., Tang, W., Enger, R., Jensen, V., Pettersen, K.H., and Nagelhus,
E.A. (2019). Astroglial endfeet exhibit distinct Ca(2+) signals during hypoosmotic conditions. Glia *67*,
2399-2409.

Erzurumlu, R.S., and Gaspar, P. (2020). How the Barrel Cortex Became a Working Model for
Developmental Plasticity: A Historical Perspective. J Neurosci *40*, 6460-6473.

Farr, G.W., Hall, C.H., Farr, S.M., Wade, R., Detzel, J.M., Adams, A.G., Buch, J.M., Beahm, D.L., Flask, C.A.,
Xu, K.*, et al.* (2019). Functionalized Phenylbenzamides Inhibit Aquaporin-4 Reducing Cerebral Edema and
Improving Outcome in Two Models of CNS Injury. Neuroscience *404*, 484-498.

Fiacco, T.A., Agulhon, C., Taves, S.R., Petravicz, J., Casper, K.B., Dong, X., Chen, J., and McCarthy, K.D.
(2007). Selective stimulation of astrocyte calcium in situ does not affect neuronal excitatory synaptic
activity. Neuron *54*, 611-626.

Gaddamanugu, S., Shafaat, O., Sotoudeh, H., Sarrami, A.H., Rezaei, A., Saadatpour, Z., and Singhal, A.
(2022). Clinical applications of diffusion-weighted sequence in brain imaging: beyond stroke.
Neuroradiology *64*, 15-30.

Giannetto, M.J., Gomolka, R.S., Gahn-Martinez, D., Newbold, E.J., Bork, P.A.R., Chang, E., Gresser, M.,
Thompson, T., Mori, Y., and Nedergaard, M. (2024). Glymphatic fluid transport is suppressed by the
aquaporin-4 inhibitor AER-271. Glia.

Gomolka, R.S., Hablitz, L.M., Mestre, H., Giannetto, M., Du, T., Hauglund, N.L., Xie, L., Peng, W.,
Martinez, P.M., Nedergaard, M.*, et al.* (2023). Loss of aquaporin-4 results in glymphatic system
dysfunction via brain-wide interstitial fluid stagnation. eLife *12*.

Gorski, J.A., Talley, T., Qiu, M., Puelles, L., Rubenstein, J.L., and Jones, K.R. (2002). Cortical excitatory
neurons and glia, but not GABAergic neurons, are produced in the Emx1-expressing lineage. J Neurosci
22, 6309-6314.

Hablitz, L.M., Pla, V., Giannetto, M., Vinitsky, H.S., Staeger, F.F., Metcalfe, T., Nguyen, R., Benrais, A., and
Nedergaard, M. (2020). Circadian control of brain glymphatic and lymphatic fluid flow. Nature
communications *11*, 4411.

Haj-Yasein, N.N., Vindedal, G.F., Eilert-Olsen, M., Gundersen, G.A., Skare, Ã., Laake, P., Klungland, A.,
Thorén, A.E., Burkhardt, J.M., Ottersen, O.P.*, et al.* (2011). Glial-conditional deletion of aquaporin-4
(Aqp4) reduces blood-brain water uptake and confers barrier function on perivascular astrocyte endfeet.
Proceedings of the National Academy of Sciences of the United States of America *108*, 17815-17820.

Harrison, I.F., Ismail, O., Machhada, A., Colgan, N., Ohene, Y., Nahavandi, P., Ahmed, Z., Fisher, A.,
Meftah, S., Murray, T.K.*, et al.* (2020). Impaired glymphatic function and clearance of tau in an
Alzheimer's disease model. Brain *143*, 2576-2593.

Herrera Moro Chao, D., Kirchner, M.K., Pham, C., Foppen, E., Denis, R.G.P., Castel, J., Morel, C.,
Montalban, E., Hassouna, R., Bui, L.C.*, et al.* (2022). Hypothalamic astrocytes control systemic glucose
metabolism and energy balance. Cell metabolism *34*, 1532-1547 e1536.

Hinson, S.R., Romero, M.F., Popescu, B.F., Lucchinetti, C.F., Fryer, J.P., Wolburg, H., Fallier-Becker, P.,
Noell, S., and Lennon, V.A. (2012). Molecular outcomes of neuromyelitis optica (NMO)-IgG binding to
aquaporin-4 in astrocytes. Proceedings of the National Academy of Sciences of the United States of
America *109*, 1245-1250.

Holthoff, K., and Witte, O.W. (1996). Intrinsic optical signals in rat neocortical slices measured with near-
infrared dark-field microscopy reveal changes in extracellular space. J Neurosci *16*, 2740-2749.

Hubbard, J.A., Hsu, M.S., Seldin, M.M., and Binder, D.K. (2015). Expression of the Astrocyte Water
Channel Aquaporin-4 in the Mouse Brain. ASN Neuro *7*.

Huber, V.J., Tsujita, M., and Nakada, T. (2009). Identification of aquaporin 4 inhibitors using in vitro and
in silico methods. Bioorg Med Chem *17*, 411-417.

Igarashi, H., Huber, V.J., Tsujita, M., and Nakada, T. (2011). Pretreatment with a novel aquaporin 4
inhibitor, TGN-020, significantly reduces ischemic cerebral edema. Neurol Sci *32*, 113-116.

Igarashi, H., Tsujita, M., Suzuki, Y., Kwee, I.L., and Nakada, T. (2013). Inhibition of aquaporin-4
significantly increases regional cerebral blood flow. Neuroreport *24*, 324-328.

Iliff, J.J., Lee, H., Yu, M., Feng, T., Logan, J., Nedergaard, M., and Benveniste, H. (2013). Brain-wide
pathway for waste clearance captured by contrast-enhanced MRI. J Clin Invest *123*, 1299-1309.

Iliff, J.J., Wang, M., Liao, Y., Plogg, B.A., Peng, W., Gundersen, G.A., Benveniste, H., Vates, G.E., Deane, R.,
Goldman, S.A*., et al.* (2012). A paravascular pathway facilitates CSF flow through the brain parenchyma
and the clearance of interstitial solutes, including amyloid beta. Sci Transl Med *4*, 147ra111.

Jeon, H., Kim, M., Park, W., Lim, J.S., Lee, E., Cha, H., Ahn, J.S., Kim, J.H., Hong, S.H., Park, J.E*., et al.*
(2021). Upregulation of AQP4 Improves Blood-Brain Barrier Integrity and Perihematomal Edema
Following Intracerebral Hemorrhage. Neurotherapeutics *18*, 2692-2706.

Jiang, R., Diaz-Castro, B., Looger, L.L., and Khakh, B.S. (2016). Dysfunctional Calcium and Glutamate
Signaling in Striatal Astrocytes from Huntington's Disease Model Mice. J Neurosci *36*, 3453-3470.

Karagiannis, A., Gallopin, T., Lacroix, A., Plaisier, F., Piquet, J., Geoffroy, H., Hepp, R., Naude, J., Le Gac,
B., Egger, R*., et al.* (2021). Lactate is an energy substrate for rodent cortical neurons and enhances their
firing activity. eLife *10*, e71424.

Kimelberg, H.K. (2004). Water homeostasis in the brain: basic concepts. Neuroscience *129*, 851-860.

Kress, B.T., Iliff, J.J., Xia, M., Wang, M., Wei, H.S., Zeppenfeld, D., Xie, L., Kang, H., Xu, Q., Liew, J.A*., et al.*
(2014). Impairment of paravascular clearance pathways in the aging brain. Ann Neurol *76*, 845-861.

Le Bihan, D. (2014). Diffusion MRI: what water tells us about the brain. EMBO Mol Med *6*, 569-573.

Le Bihan, D., and Iima, M. (2015). Diffusion Magnetic Resonance Imaging: What Water Tells Us about
Biological Tissues. PLoS Biol *13*, e1002203.

Le Bihan, D., Urayama, S., Aso, T., Hanakawa, T., and Fukuyama, H. (2006). Direct and fast detection of
neuronal activation in the human brain with diffusion MRI. Proceedings of the National Academy of
Sciences of the United States of America *103*, 8263-8268.

Li, J., Jia, Z., Xu, W., Guo, W., Zhang, M., Bi, J., Cao, Y., Fan, Z., and Li, G. (2019). TGN-020 alleviates
edema and inhibits astrocyte activation and glial scar formation after spinal cord compression injury in
rats. Life sciences *222*, 148-157.

Li, X., Xie, Z., Zhou, Q., Tan, X., Meng, W., Pang, Y., Huang, L., Ding, Z., Hu, Y., Li, R.*, et al.* (2023). TGN-020
Alleviate Inflammation and Apoptosis After Cerebral Ischemia-Reperfusion Injury in Mice Through
Glymphatic and ERK1/2 Signaling Pathway. Mol Neurobiol.

Lucchinetti, C.F., Guo, Y., Popescu, B.F., Fujihara, K., Itoyama, Y., and Misu, T. (2014). The pathology of
an autoimmune astrocytopathy: lessons learned from neuromyelitis optica. Brain Pathol *24*, 83-97.

MacAulay, N. (2021). Molecular mechanisms of brain water transport. Nat Rev Neurosci *22*, 326-344.

MacAulay, N., Hamann, S., and Zeuthen, T. (2004). Water transport in the brain: role of cotransporters.
Neuroscience *129*, 1031-1044.

MacVicar, B.A., and Hochman, D. (1991). Imaging of synaptically evoked intrinsic optical signals in
hippocampal slices. J Neurosci *11*, 1458-1469.

Madisen, L., Mao, T., Koch, H., Zhuo, J.M., Berenyi, A., Fujisawa, S., Hsu, Y.W., Garcia, A.J., 3rd, Gu, X.,
Zanella, S.*, et al.* (2012). A toolbox of Cre-dependent optogenetic transgenic mice for light-induced
activation and silencing. Nat Neurosci *15*, 793-802.

Mestre, H., Hablitz, L.M., Xavier, A.L., Feng, W., Zou, W., Pu, T., Monai, H., Murlidharan, G., Castellanos
Rivera, R.M., Simon, M.J.*, et al.* (2018a). Aquaporin-4-dependent glymphatic solute transport in the
rodent brain. eLife *7*.

Mestre, H., Tithof, J., Du, T., Song, W., Peng, W., Sweeney, A.M., Olveda, G., Thomas, J.H., Nedergaard,
M., and Kelley, D.H. (2018b). Flow of cerebrospinal fluid is driven by arterial pulsations and is reduced in
hypertension. Nature communications *9*, 4878.

Mireles-Ramirez, M.A., Pacheco-Moises, F.P., Gonzalez-Usigli, H.A., Sanchez-Rosales, N.A., Hernandez-
Preciado, M.R., Delgado-Lara, D.L.C., Hernandez-Cruz, J.J., and Ortiz, G.G. (2022). Neuromyelitis optica
spectrum disorder: pathophysiological approach. Int J Neurosci, 1-13.

Mola, M.G., Sparaneo, A., Gargano, C.D., Spray, D.C., Svelto, M., Frigeri, A., Scemes, E., and Nicchia, G.P.
(2016). The speed of swelling kinetics modulates cell volume regulation and calcium signaling in
astrocytes: A different point of view on the role of aquaporins. Glia *64*, 139-154.

Murphy, T.R., Davila, D., Cuvelier, N., Young, L.R., Lauderdale, K., Binder, D.K., and Fiacco, T.A. (2017).
Hippocampal and Cortical Pyramidal Neurons Swell in Parallel with Astrocytes during Acute
Hypoosmolar Stress. Front Cell Neurosci *11*, 275.

Noda, Y., Sohara, E., Ohta, E., and Sasaki, S. (2010). Aquaporins in kidney pathophysiology. Nat Rev
Nephrol *6*, 168-178.

Ochoa-de la Paz, L.D., and Gulias-Canizo, R. (2022). Glia as a key factor in cell volume regulation
processes of the central nervous system. Front Cell Neurosci *16*, 967496.

Oosuka, S., Kida, T., Oku, H., Horie, T., Morishita, S., Fukumoto, M., Sato, T., and Ikeda, T. (2020). Effects
of an Aquaporin 4 Inhibitor, TGN-020, on Murine Diabetic Retina. Int J Mol Sci *21*.

Papadopoulos, M.C., and Verkman, A.S. (2013). Aquaporin water channels in the nervous system. Nat
Rev Neurosci *14*, 265-277.

Pham, C., Moro, D.H., Mouffle, C., Didienne, S., Hepp, R., Pfrieger, F.W., Mangin, J.M., Legendre, P.,
Martin, C., Luquet, S.*, et al.* (2020). Mapping astrocyte activity domains by light sheet imaging and
spatio-temporal correlation screening. Neuroimage *220*, 117069.

Rasmussen, M.K., Mestre, H., and Nedergaard, M. (2022). Fluid transport in the brain. Physiol Rev *102*,
1025-1151.

Risher, W.C., Andrew, R.D., and Kirov, S.A. (2009). Real-time passive volume responses of astrocytes to
acute osmotic and ischemic stress in cortical slices and in vivo revealed by two-photon microscopy. Glia
57, 207-221.

Salman, M.M., Kitchen, P., Yool, A.J., and Bill, R.M. (2022). Recent breakthroughs and future directions in
drugging aquaporins. Trends Pharmacol Sci *43*, 30-42.

Slezak, M., Goritz, C., Niemiec, A., Frisen, J., Chambon, P., Metzger, D., and Pfrieger, F.W. (2007).
Transgenic mice for conditional gene manipulation in astroglial cells. Glia *55*, 1565-1576.

Smith, A.J., and Verkman, A.S. (2019). CrossTalk opposing view: Going against the flow: interstitial solute
transport in brain is diffusive and aquaporin-4 independent. J Physiol *597*, 4421-4424.

Smith, A.J., Yao, X., Dix, J.A., Jin, B.J., and Verkman, A.S. (2017). Test of the 'glymphatic' hypothesis
demonstrates diffusive and aquaporin-4-independent solute transport in rodent brain parenchyma.
eLife *6*.

Solenov, E., Watanabe, H., Manley, G.T., and Verkman, A.S. (2004). Sevenfold-reduced osmotic water
permeability in primary astrocyte cultures from AQP-4-deficient mice, measured by a fluorescence
quenching method. Am J Physiol Cell Physiol *286*, C426-432.

Stokum, J.A., Gerzanich, V., and Simard, J.M. (2016). Molecular pathophysiology of cerebral edema.
Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral
Blood Flow and Metabolism *36*, 513-538.

Sun, C., Lin, L., Yin, L., Hao, X., Tian, J., Zhang, X., Ren, Y., Li, C., and Yang, Y. (2022). Acutely Inhibiting
AQP4 With TGN-020 Improves Functional Outcome by Attenuating Edema and Peri-Infarct Astrogliosis
After Cerebral Ischemia. Front Immunol *13*, 870029.

Tait, M.J., Saadoun, S., Bell, B.A., Verkman, A.S., and Papadopoulos, M.C. (2010). Increased brain edema
in aqp4-null mice in an experimental model of subarachnoid hemorrhage. Neuroscience *167*, 60-67.

Taniguchi, H., He, M., Wu, P., Kim, S., Paik, R., Sugino, K., Kvitsiani, D., Fu, Y., Lu, J., Lin, Y.*, et al.* (2011). A
resource of Cre driver lines for genetic targeting of GABAergic neurons in cerebral cortex. Neuron *71*,
995-1013.

Toft-Bertelsen, T.L., Larsen, B.R., Christensen, S.K., Khandelia, H., Waagepetersen, H.S., and MacAulay,
N. (2021). Clearance of activity-evoked K(+) transients and associated glia cell swelling occur
independently of AQP4: A study with an isoform-selective AQP4 inhibitor. Glia *69*, 28-41.

Tofts, P.S., Lloyd, D., Clark, C.A., Barker, G.J., Parker, G.J., McConville, P., Baldock, C., and Pope, J.M.
(2000). Test liquids for quantitative MRI measurements of self-diffusion coefficient in vivo. Magn Reson
Med *43*, 368-374.

Tsurugizawa, T., Ciobanu, L., and Le Bihan, D. (2013). Water diffusion in brain cortex closely tracks
underlying neuronal activity. Proceedings of the National Academy of Sciences of the United States of
America *110*, 11636-11641.

Verkman, A.S., Ratelade, J., Rossi, A., Zhang, H., and Tradtrantip, L. (2011). Aquaporin-4: orthogonal
array assembly, CNS functions, and role in neuromyelitis optica. Acta Pharmacol Sin *32*, 702-710.

Verkman, A.S., Smith, A.J., Phuan, P.W., Tradtrantip, L., and Anderson, M.O. (2017). The aquaporin-4
water channel as a potential drug target in neurological disorders. Expert opinion on therapeutic targets
21, 1161-1170.

Vindedal, G.F., Thoren, A.E., Jensen, V., Klungland, A., Zhang, Y., Holtzman, M.J., Ottersen, O.P., and
Nagelhus, E.A. (2016). Removal of aquaporin-4 from glial and ependymal membranes causes brain water
accumulation. Mol Cell Neurosci *77*, 47-52.

Woo, J., Kim, J.E., Im, J.J., Lee, J., Jeong, H.S., Park, S., Jung, S.Y., An, H., Yoon, S., Lim, S.M*., et al.* (2018).
Astrocytic water channel aquaporin-4 modulates brain plasticity in both mice and humans: a potential
gliogenetic mechanism underlying language-associated learning. Mol Psychiatry *23*, 1021-1030.

Wu, Y.C., and Alexander, A.L. (2007). A method for calibrating diffusion gradients in diffusion tensor
imaging. J Comput Assist Tomogr *31*, 984-993.

Xiao, M., Hou, J., Xu, M., Li, S., and Yang, B. (2023). Aquaporins in Nervous System. Adv Exp Med Biol
1398, 99-124.

Yang, J., Vitery, M.D.C., Chen, J., Osei-Owusu, J., Chu, J., and Qiu, Z. (2019). Glutamate-Releasing SWELL1
Channel in Astrocytes Modulates Synaptic Transmission and Promotes Brain Damage in Stroke. Neuron.

Yao, X., Smith, A.J., Jin, B.J., Zador, Z., Manley, G.T., and Verkman, A.S. (2015). Aquaporin-4 regulates the
velocity and frequency of cortical spreading depression in mice. Glia *63*, 1860-1869.

Zhao, H.H., Du, H., Cai, Y., Liu, C., Xie, Z., and Chen, K.C. (2019). Time-resolved quantification of the
dynamic extracellular space in the brain: study of cortical spreading depression. Journal of
neurophysiology *121*, 1735-1747.

FIGURE 1

Fig. 1. Astrocyte aquaporin mediates a tonic water efflux. (**A**) *In vivo* chemical labelling of astrocytes. Sulforhodamine B (SRB, 10 mg/ml) was intraperitoneally injected in awake mice (10 µl/g). Monochrome images show representative astrocyte labeling in living acute brain slices from cortex under low (×20; scale bar, 50 µm) and high magnification (×63; scale bar, 20 µm) by epifluorescence. *Below*, SRB labeling was confirmed to be astrocyte-specific in acute brain slices of the astrocyte reporter line GFAP-EGFP, where light sheet imaging was used to gain optical sectioning (Materials and methods; Fig. S1). The light gray circle indicates the astrocyte regions used for fluorescence analysis. Scale bar, 50 µm. (**B**) Optical imaging of astrocyte water transport in acute brain slices. Transmembrane water transport was triggered with hypo- and hypertonic solution, inducing water inflow and outflow that were respectively reflected by SRB fluorescence decrease (*above*) and increase (*below*; expressed as dF/F$_0$). The hypo- or hypertonic solution was applied to slices over different lengths of duration displayed in colors corresponding to the time courses of SRB fluorescence (n = 52 astrocytes, 4 mice). (**C**) Acutely blocking astrocyte AQP4 with TGN-020 caused intracellular water accumulation and swelling. The downward change in the SRB fluorescence was respectively calculated for the phases prior and post to vehicle or TGN application. *Left*, while no effect was seen under CTR condition (vehicle only, n = 23 astrocytes, 3 mice), TGN-020 (20 µM) significantly decreased astrocyte SRB fluorescence (n = 30, 6 mice). Imaging was performed in acute brain slices of layer II/III S1 cortex. *Middle*, the downward slope was compared between the periods before and after the application of TGN-020. *Right*, illustration shows astrocyte aquaporin sustaining a tonic water efflux. Its blockade causes water accumulation and cell swelling. (**D**) *In vivo* validation of the effect of TGN-020 application on astrocyte water homeostasis. *Left*, fiber photometry was used for real-time recording of SRB fluorescence of astrocyte population in S1 cortex in freely moving mice, with saline (CTR) or TGN-020 being intraperitoneally injected when SRB was trapped in astrocytes. Fiber photometry recording shows that *in vivo* SRB injection resulted in rapid entry into mouse cortex and, in about one hour, led to astrocyte labeling (inset scale bar, 50 µm). *Middle*, example response to saline and TGN-020. The change of SRB fluorescence relative to the photobleaching tendency delineated by line fitting (dotted line) was examined. *Right*, relative to CTR, TGN administration led to a decrease in astrocyte SRB fluorescence and its oscillation range (n = 8 recordings per condition, 5 mice).

FIGURE 2

Fig. 2. Acutely blocking astrocyte aquaporin induces swelling-associated Ca^{2+} oscillation. (**A**) *In vivo* expression in astrocytes of the genetically encoded fluorescent Ca^{2+} indicator GCaMP6 for imaging astrocyte Ca^{2+} in acute brain slices. Angular light sheet microscopy (Fig. S1) was used to capture transient Ca^{2+} signals of local regions. As a positive control, astrocyte swelling was induced by hypotonic solution (100 mOsM) that caused Ca^{2+} changes from their homeostatic level. *Left*, representative time courses of swelling-induced Ca^{2+} changes in detected response regions; *middle to left*, the histogram distribution and bar representation showing the signal strengths that were derived from the temporal integral of individual Ca^{2+} time courses normalized per minute, before and after cell swelling (n = 15 ROIs, from three mice). (**B**) Astrocyte Ca^{2+} oscillation caused by acute aquaporin blocking with TGN-020 (20 Imaging perivascular endfeet of GCaMP6-expressing astrocytes.µM, n = 31 ROIs, from five mice), due to the inhibition of the tonic water efflux that led to astrocyte swelling as illustrated. Time scale and Ca^{2+} rise scale (dF/F$_0$) are the same as in *A*. Scale bar, 50 µm. (**C**) Intercellular effect on SST interneurons of blocking astrocyte aquaporin water efflux. TGN-020 (20 µM) or the equal molar vehicle (CTR) was bath applied to acute cortical slices of SST-GCaMP6 mice (n = 7-12 regional measurements from four mice). Scale bar, 50 µm.

FIGURE 3

Fig. 3. Tonic water efflux via aquaporin modulates phasic transmembrane water transport and astrocyte volume response. (**A**) *Left*, in basal condition astrocyte aquaporin mediates a tonic water efflux. *Right*, protocol to induce water outflow from astrocytes, therefore their shrinking, by hypertonic extracellular solution (400 mOsM) in either control condition (CTR) or in the presence of AQP4 inhibitor TGN-020 (20 µM). (**B**) Time courses of astrocyte SRB fluorescence increase upon the phasically induced water outflow, reflecting the occurrence of shrinking. The histograms and bar charts compare the start time, namely the delay between hypertonic solution application and rise in SRB fluorescence, the time to reach the peak of shrinking, and the absolute amplitude of water outflow-induced SRB increase (n = 58 astrocytes for CTR, and 47 astrocytes for TGN-020, four mice). (**C**) *Right*, protocol to trigger water inflow into astrocytes by hypotonic extracellular solution (100 mOsM) in either CTR solution or in the presence of TGN-020 (20 µM). (**D**) Time courses of astrocyte SRB fluorescence decrease caused by water inflow, which also reflects concomitant cell swelling. In contrast to the observation with hypertonicity-induced water outflow and astrocyte shrinking, a reduction was observed for both the start time and the time-to-peak with TGN-020 (n = 12 astrocytes for CTR, and 12 astrocytes for TGN-020, four mice). TGN-020 led to a decrease in the absolute amplitude of astrocyte swelling.

FIGURE 4

Fig. 4. AQP4-mediated tonic water outflow regulates global swelling in cortical parenchyma. CSD-associated general swelling was induced by photostimulating ChR2-expressing pyramidal cells in acute cortical slices, and recorded by imaging intrinsic optical signal (IOS) with infrared illumination. (**A**) Representative recording. The pia surface is on the upper side; green, blue and red squares correspond to regions of interest in Layers I, II-III and IV, respectively. Transmittance signals are represented in pseudocolor images at different time points post photostimulation. Scale bar, 100 µm. (**B-C**) Kymograph and time courses showing the IOS changes following photoactivaiton, derived from the radial line of interest indicated by a black arrow in *A*. The light blue line indicates the 10-s photostimulation that increases the infrared transmittance signal (dT/T_0) across cortical layers, as also illustrated by the time courses; *a*, *b*, *c* and *d* correspond to the time points depicted in *A*. After the onset delay (*a*), the first (*b*) and second (*c*) peak of IOS are characteristic of the CSD and a prolonged general swelling, respectively. (**D-E**) TGN-020 (20 µM) inhibition of AQP4 reduced the initial onset and the maximal amplitude of both the CSD and general swelling (n = 6 - 18 measurements from 13 mice per condition) in layer II/III cortex. Inhibiting spiking activity with TTX (1 µM) only affected the amplitude of the initial CSD response. (**F**) Astrocytes swelling, reflected by SRB fluorescence decrease, monitored in control condition (n = 75 astrocytes) and in the presence of TGN-020 (n = 17 astrocytes) in layer II/III cortex.

FIGURE 5

Fig. 5. Perturbing the tonic water efflux via astrocyte aquaporin alters brain water homeostasis. *In vivo* DW-MRI (7T) was employed to map water diffusion in the entire brain scale following the acute inhibition of astrocytic AQP4 (TGN-020, intraperitoneal injection, 200 mg/kg), with paralleled control experiments performed with saline injection. (**A**) Experimental protocol for DW-MRI. Saline or TGN020 was injected at 10 min after the start of acquisition. DWI was acquired every 5 min. (**B**) Representative image obtained at three different b values to derive the water diffusion rates. (**C**) *Left*, brain water diffusion rate was mapped by the calculation of apparent diffusion coefficient (ADC). *Right*, representative images illustrating the relative changes of ADC at 60 min after injection of saline or TGN-020. (**D**) *Upper panel*, time courses depicting the temporal changes in ADC in the cortex, striatum, and hippocampus, revealing the regional heterogeneity. Arrowhead indicates the injection of saline or TGN-020 (n = 7 mice for saline injection, 7 mice for TGN-020). *p < 0.05 vs baseline at each time-point following the two-way repeated measures ANOVA. *Lower panel*, box-plot of the averaged ADC between 30 and 50 min after the injection. *$p < 0.05$ by paired t-test.