Astrocyte aquaporin mediates a tonic water efflux maintaining brain homeostasis

Cuong Pham[1], Yuji Komaki[2], Anna Deàs-Just[1], Benjamin Le Gac[1], Christine Mouffle[1], Clara Franco[1], Vincent Vialou[1], Tomokazu Tsurugizawa[3,4,#], Bruno Cauli[1,#], Dongdong Li[1,#]

[1]Sorbonne Université, Institute of Biology Paris Seine, Neuroscience Paris Seine, CNRS UMR8246, INSERM U1130, Paris 75005, France

[2]Central Institute for Experimental Animals, Kawasaki, Japan

[3]Human Informatics and Interaction Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Japan

[4]Faculty of Engineering, University of Tsukuba, Tsukuba, Japan

[#]Correspondence: t-tsurugizawa@aist.go.jp, bruno.cauli@upmc.fr, dongdong.li@inserm.fr

Keywords: AQP4, imaging, swelling, diffusion

ABSTRACT

Brain water homeostasis provides not only physical protection, but also determines the diffusion of chemical molecules key for information processing and metabolic stability. As a major type of glia in brain parenchyma, astrocytes are the dominant cell type expressing aquaporin water channel. However, how astrocyte aquaporin contributes to brain water homeostasis in basal physiology remains to be understood. We report that astrocyte aquaporin 4 (AQP4) mediates a tonic water efflux in basal conditions. Acute inhibition of astrocyte AQP4 leads to intracellular water accumulation as optically resolved by fluorescence-translated imaging in acute brain slices, and *in vivo* by fiber photometry in moving mice. We then show that the tonic aquaporin water efflux maintains astrocyte volume equilibrium, astrocyte and neuron Ca^{2+} signaling, and extracellular space remodeling during optogenetically induced cortical spreading depression. Using diffusion-weighted magnetic resonance imaging (DW-MRI), we observed that *in vivo* inhibition of AQP4 water efflux heterogeneously disturbs brain water homeostasis in a region-dependent manner. Our data suggest that astrocyte aquaporin, though bidirectional in nature, mediates a tonic water outflow to sustain cellular and environmental equilibrium in brain parenchyma.

Significance statement - Our brain is immersed, thus protected, in a water environment. It

ensures intra- and extracellular molecular diffusion, which is vital for brain function and health.

Brain water homeostasis is maintained by dynamic water transport between different cell types.

Astrocytes are a main type of glial cell widely distributed in brain parenchyma, expressing the

bidirectional aquaporin water channel. Here we show that in basal conditions, aquaporin channel

mediates a tonic water efflux from astrocytes. This mechanism maintains astrocyte volume

stability, activity-gated brain parenchyma remodeling and brain water homeostasis. Our finding

sheds light on how astrocytes regulate water states in the brain, and will help to understand brain

allostasis in specific life contexts.

1 Introduction

Every aspect of brain function relies on the delicately maintained water environment. It supports

brain structural stability and molecular diffusion, laying the ground for information processing,

metabolite shuttling and adaptation to living environments (Kimelberg, 2004). Water is the

fundamental constituent of the cerebrospinal fluid infiltrating into the central nervous system and

the interstitial fluid distributed in brain parenchyma (Agnati et al., 2017, Brinker et al., 2014).

Brain fluid transport is suggested to support the diffusion of energetic fuels like glucose and

lactate to warrant the metabolic circumstances in the parenchyma, and is also implicated in the

clearance of waste molecules from the brain as described in the glymphatic system (Abbott et al.,

2018, Louveau et al., 2017). In addition, brain water diffusion is the basis for diffusion-weighted

magnetic resonance imaging (DW-MRI) for structural and functional neuroimaging (Le Bihan et

al., 2006), and for clinical diagnostics widely applied for neurological diseases such as ischemia,

brain tumor and edema (Gaddamanugu et al., 2022, Le Bihan & Iima, 2015).

Water equilibrium in the brain is maintained by dynamic transport between different cell

entities. Aquaporin is a family of transmembrane channels facilitating bidirectional water flow,

with aquaporin 4 (AQP4) being the main subtype expressed in the central nervous system. In the

brain, AQP4 is predominantly expressed in astrocytes (Papadopoulos & Verkman, 2013, Xiao et

al., 2023) that are a major type of glial cell in the parenchyma (Barres, 2008). This feature

enables astrocytes to well balance osmotic oscillations imposed by the transmembrane transport

of ions, metabolites, and signaling molecules, all mediated by water. Dysregulation in astrocyte

aquaporin water transport is implicated in pathological scenarios such as cerebral edema and

neuroinflammation (Verkman et al., 2011). Valuable insights have been obtained on the function

of astrocyte aquaporin in specific physiopathological conditions including osmotically evoked

volume responses, ischemia and neurodegenerative diseases (Verkman et al., 2017). Meanwhile, how astrocyte aquaporin contributes to brain water homeostasis in basal physiology has remained elusive. A better understanding will aid to discern astrocytic regulation of cerebral water states, thereby the potential maladaptations in neurological disorders.

Here, combining *in vivo* chemical targeting, optical and MRI imaging of water dynamics, we report that AQP4 sustains a tonic water efflux from brain astrocytes. This mechanism was found to be critical in maintaining astrocyte volume and signaling stability, as well as the extracellular space remodeling in mouse cortex during optogenetically induced cortical spreading depression. Using DW-MRI, we observed that acute inhibition of astrocyte water efflux *in vivo* heterogeneously alters water diffusion across brain regions. Our finding suggests that aquaporin acts as a water export route in astrocytes in physiological conditions, so as to counterbalance the constitutive intracellular water accumulation caused by constant transmitter and ion uptake, as well as the cytoplasmic metabolism processes. This mechanism hence plays a necessary role in maintaining water equilibrium in astrocytes, thereby brain water homeostasis.

Results

Astrocyte aquaporin mediates a tonic water efflux

To optically follow *in situ* astrocyte water transport, we performed fluorescence intensity-translated (FIT) imaging. To this end, mouse brain astrocytes were chemically labeled *in vivo* with a highly water-diffusible and astrocyte-specific red fluorescent dye, sulforhodamine B (SRB) (Fig. 1A) (Appaix et al., 2012, Pham et al., 2020). About one hour after its intraperitoneal injection, astrocytic SRB labeling was widely distributed in the mouse cortex, as seen in living

acute brain slices (Fig. 1A) and validated by colocalization with EGFP-identified astrocytes in slices taken from GFAP-EGFP transgenic mice (Fig. 1A). Net water transport across the astrocyte membrane alters cytoplasmic SRB concentration and cellular volume, which can thus be followed by changes in fluorescence intensity when imaged in a fixed field of view. To validate the FIT imaging of astrocyte water transport, we employed wide-field fluorescence microscopy for single-plane time-lapse imaging in acute brain slices in the primary somatosensory (S1) cortex. This approach allowed us to collect fluorescence from both the focal plane and along the axial extensions, thereby imprinting volumetric fluorescence into the single image plane. Indeed, water influx induced by the applicaiton of hypoosmotic solution over different periods proportionally decreased astrocyte SRB fluorescence, which also reflected cell swelling (Fig. 1B); whereas water export and astrocyte shrinking upon hyperosmotic manipulation increased astrocyte fluorescence (Fig. 1B). Hence, FIT imaging enables real-time recording of astrocyte transmembrane water transport and volume dynamics.

In basal conditions, flat fluorescence time course was recorded in brain slices suggesting equilibrated water transport across the astrocyte membrane (Fig. 1B) and volume homeostasis in the brain parenchyma. To inspect a tonic role of aquaporin in astrocyte water transport, we sought to acutely block astrocytic aquaporin channel, aquaporin 4 (AQP4). We used the synthetic compound 2-(nicotinamido)-1,3,4-thiadiazole (TGN-020) that is derived from the condensation of nicotinamide and thiadiazole derivatives (Burnett et al., 2015, Huber et al., 2009), whose specificity for AQP4 has been validated *in vitro* by ion channel heterologous expression system (Toft-Bertelsen et al., 2021) and *in vivo* using the AQP4 knockout (KO) mice (Harrison et al., 2020, Igarashi et al., 2013). This approach guaranteed the functional pinpointing of astrocyte aquaporin under physiological conditions, while avoiding the chronic compensations

caused by genetic tools and mouse models that have been reported to alter brain water content,

volume and extracellular architecture potentially confounding the functional readouts (Binder et

al., 2004, Gomolka et al., 2023, Haj-Yasein et al., 2011, MacAulay, 2021, Yao et al., 2015).

AQP4 is a bidirectional channel facilitating passive water transport along the osmotic

gradient (Papadopoulos & Verkman, 2013). As an osmotic equilibrium is maintained in the brain

parenchyma in basal state, there might not be net water transport across AQP4. However, we

observed in cortical slices that acute inhibition of astrocyte AQP4 by TGN-020 caused a gradual

decrease in SRB fluorescence intensity, indicating an intracellular water accumulation (Fig. 1C).

This observation suggests that astrocyte aquaporin mediates a tonic water efflux; its blocking

causes intracellular water accumulation and swelling (Fig. 1C, *right*). To corroborate this

observation *in vivo*, we performed fiber photometry recording in moving mice (Fig. 1D). An

optical fiber was implanted into the mouse S1 cortex to follow fluorescence signals of local

astrocyte population post the intraperitoneal injection of SRB (Fig. 1D, *left*). After about one

hour, when astrocytes were well labeled, either saline or TGN-020 was injected intraperitoneally.

While no significant effect was observed with the saline control, TGN-020 induced a decrease in

astrocyte SRB fluorescence and a reduction in its oscillation range (Fig. 1D, *right*), mirroring an

intracellular dilution of fluorescent molecules. This observation confirms that acute inhibition of

astrocyte aquaporin leads to intracellular water accumulation thereby cell swelling.

Aquaporin inhibition perturbs astrocyte and neuron signaling

Astrocyte volume equilibrium not only determines brain architectural stability, but also

associates with dynamic cellular signals. Astrocyte swelling has been shown to alter intracellular

Ca^{2+} signaling (Benfenati et al., 2011, Eilert-Olsen et al., 2019). We performed Ca^{2+} imaging to

confirm that acute aquaporin inhibition induces astrocyte swelling and then Ca^{2+} oscillation. The

genetically encoded Ca^{2+} sensor GCaMP6f was selectively expressed in astrocytes by crossing

the Glast-Cre^{ERT2} and GCaMP6^{floxP} mouse lines (Herrera Moro Chao et al., 2022, Pham et al.,

2020). Because astrocyte Ca^{2+} signals often occur in local domains, we adapted a light sheet

microscope (Pham et al., 2020) for wide-field optical sectioning so as to image them with high

signal-to-noise ratio (Fig. S1). We first confirmed that water accumulation, therefore astrocyte

swelling, induced by hypotonic solution enhances the intracellular Ca^{2+} signal (Fig. 2A). Then

we followed astrocyte Ca^{2+} in isotonic control condition while blocking astrocyte AQP4 with

TGN-020, and observed that Ca^{2+} signaling was similarly augmented following the acute

aquaporin inhibition (Fig. 2B). This result confirms that inhibiting astrocyte tonic water efflux

leads to intracellular water accumulation and swelling, which causes alterations in astrocytic

Ca^{2+} signaling. In addition, astrocyte swelling has been reported to induce the release of

neuroactive molecules such as glutamate thereby influencing nearby neuron activity (Fiacco et

al., 2007, Yang et al., 2019). Disrupting AQP4 tonic water outflow would not only cause

astrocyte swelling but may also influence the activity of neighboring neurons. We then imaged

Ca^{2+} signals as a surrogate for neuronal activity in cortical somatostatin (SST) interneurons

which exhibit high excitability (Karagiannis et al., 2021) and express both ionotropic and

metabotropic glutamate receptors (Cauli et al., 2000), rendering them ideal to sense astrocyte-

released signaling molecules. We expressed GCaMP6f in SST interneurons by crossing

homozygous SST-Cre (Taniguchi et al., 2011) and homozygous GCaMP6^{floxP} mice. As expected,

TGN-020 inhibition of astrocyte AQP4 led to a global Ca^{2+} elevation in SST interneuron

populations (Fig. 2C). These observations support that astrocyte aquaporin mediates a tonic

water efflux, contributing to maintain both the volume and signaling homeostasis.

Aquaporin water efflux regulates astrocyte volume response

We then examined the role of aquaporin water efflux in astrocyte volume response to osmotic

changes. The AQP4 blocker TGN-020 or equimolar vehicle was applied throughout the

recording, namely being continuously present before and after osmotic challenges. We first

followed the evoked water efflux and shrinking induced by hypertonic solution while. As

astrocyte AQP4 supports preferentially water efflux, its inhibition would attenuate hypertonicity-

imposed water extrusion (Fig. 3A). Supporting this, application of TGN-020 slowed down the

hypertonicity-induced water efflux and shrinking, reflected by the longer time to peak in the

SRB fluorescence time course as compared to control (Fig. 3B), though the delay in the initial

onset was not significantly prolonged. The maximal increase in astrocyte SRB fluorescence was

lower in the presence of TGN-020, suggesting that AQP4 blocking reduced the overall amount

of water efflux (Fig. 3B).

We next evoked water influx, thereby astrocyte swelling, by hypotonic solution in the

presence or absence of TGN-020 (Fig. 3C). In contrast to the effects on hypertonicity-evoked

water efflux, AQP4 acute inhibition was observed to accelerate both the initial water

accumulation and the swelling rate in astrocytes, reflected by their earlier onset in SRB

fluorescence decrease and faster reaching to the plateau value relative to the control condition

(Fig. 3D). This observation cross-validates the role of astrocyte aquaporin in supporting water

efflux, whereby its blockade facilitates the initial accumulation of the evoked water influx (Fig.

3C). In contrast, the late-stage maximum decrease in astrocyte SRB fluorescence was observed

to be reduced in the presence of TGN-020 (Fig. 3D), reflecting a reduction in the total amount of

water accumulation and swelling. As TGN-020 was present prior to hypotonic challenge, it

would have slightly swelled astrocytes due to the blockade of tonic water efflux, thereby

constraining the range of the further swelling induced the subsequent hypotonicity. Further,

while astrocyte AQP4 sustains a water outflow in basal physiology, in conditions when

transmembrane water gradient is altered as occurred in hypotonic solutions, the net water flow

through AQP4 should be finally dictated by the osmotic gradient due to its bidirectional nature.

Therefore, along the imposed hypotonicity, AQP4 started to instruct a water influx; its constant

blocking by TGN-020 would have reduced the total amount of water influx and swelling.

Astrocytes are widely distributed throughout the brain parenchyma, orchestrating water

transport and volume homeostasis (Ochoa-de la Paz & Gulias-Canizo, 2022). Water outflow via

astrocyte AQP4 may play a role in the structural remodeling of parenchyma at the global level.

We then induced general cell swelling by triggering cortical spreading depression (CSD)

(Holthoff & Witte, 1996, Zhao et al., 2019). To be orthogonal to the pharmacological control of

astrocyte aquaporin, the CSD depolarization wave was initiated by optogenetically stimulating

ChR2-expressing glutamatergic pyramidal cells in the cortical slices of Emx1-Cre/Ai32ChR2

mice (Gorski et al., 2002, Madisen et al., 2012), termed Opto-CSD (Chung et al., 2019). As cell

swelling increases the transmittance of infrared light, we imaged Opto-CSD by the intrinsic

optical signal (IOS) derived from infrared illumination (Holthoff & Witte, 1996, MacVicar &

Hochman, 1991), which was spectrally distinct from ChR2 photoactivation (detailed in Methods).

The IOS displayed transient increase across cortical layers during the Opto-CSD (Fig. 4A-C),

exhibiting propagating kinetics characteristic of spreading depression (Chung et al., 2019,

Holthoff & Witte, 1996). The biphasic starting of IOS coincided with a sharp response in the

extracellular potential indicating the CSD initiation (Fig. S2A). IOS signal contains a first peak

reflecting the rapid CSD response (Fig. 4C, *b*) followed by a prolonged phase of general cellular

swelling (Fig. 4C, *c*). By combining fluorescence imaging of SRB- labelled astrocytes, whose

spectrum is separated from the IOS infrared signal, we observed that astrocytes swelling (i.e.,

decrease in SRB fluorescence) paralleled CSD swelling (Fig. S3). Consistent with our

observation on astrocyte volume response (Fig. 3D), when pre-incubating slices with TGN-020

to block AQP4 tonic water outflow, the initiation of both the CSD and general swelling was

accelerated while their maximum amplitude reduced (Fig. 4D-E*;* Fig. S2B). Blocking action

potentials with tetrodotoxin (TTX) only reduced the amplitude of the initial CSD response while

the effect on general swelling is insignificant (Fig. 4D-E). We further followed the swelling of

individual SRB-labelled astrocytes during Opto-CSD. In consistence with the result obtained

from hypotonic challenge (Fig. 3D), the presence of TGN-020 reduced the peak amplitude of

astrocyte swelling (i.e., the maximal SRB fluorescence decrease; Fig. 4F). As a post hoc

adaptation to the transiently induced Opto-CSD, astrocyte swelling is followed by a shrinking to

recover cellular volume to the homeostatic level, which was recorded as a rebound in SRB

fluorescence intensity. We noted that TGN-020 presence significantly restrained the recovery of

SRB fluorescence back to the baseline (Fig. 4F), showing that AQP4 inhibition hinders astrocyte

water efflux therefore the shrinking.

Collectively, our data show that astrocyte AQP4 sustains a tonic water outflow regulating

the cellular volume response and the general cell swelling of the parenchyma.

Tonic astrocyte water transport underlies brain homeostasis

Water homeostasis sets the basis for molecular diffusion in the brain, which instructs

neurotransmitter availability, ion recycling and metabolite trafficking. We then examined *in vivo*

the role of astrocyte aquaporin outflow in brain water diffusion using DW-MRI (Le Bihan &

Iima, 2015). It uses the diffusion of water molecules to generate contrast that can be quantified

by the apparent diffusion coefficient (ADC) in the nerve tissue (Beaulieu, 2002, Le Bihan, 2014).

We used a 7T MRI to image global brain water diffusion in mice lightly anesthetized with

medetomidine that normalized the animals under the sedative state. Astrocyte AQP4 outflow was

acutely perturbed by TGN-020 applied via intraperitoneal injection. Brain water diffusion was

mapped every 5 min before and after TGN or saline (as control) administration (Fig. 5A). Each

acquisition sweep was performed with three b values (0, 250, 1800) that accounted for ADC, so

to fit the exponential curve (Fig. 5B). Compared to the control saline, TGN-020 increased basal

water diffusion within multiple regions including the cortex, hippocampus and the striatum in a

heterogeneous manner (Fig. 5C). The temporal features in water diffusion change appeared to be

also different: an early elevation followed by a reversible tendency was observed for cortical

areas, a rapid and long-lasting elevation for the striatum while a delayed and transient increase

for the hippocampus (Fig. 5D). The *in vivo* neuroimaging results confirm that the tonic water

efflux from astrocyte aquaporin contributes to maintain the homeostasis of brain water diffusion,

and also suggest spatiotemporal heterogeneities in brain water handling.

Discussion

20 Water equilibrium underlies brain function, plasticity and dynamics. Astrocytes are the principal

21 brain cell type expressing aquaporin water channels that are highly implicated in regulating local

22 environments in the parenchyma (Xiao et al., 2023). We show that albeit being a bidirectional

channel, astrocyte aquaporin sustains a tonic water outflow in basal states to ensure structural

and functional stability, as well as the water homeostasis at the brain level.

The observation of a basal aquaporin water efflux implies there is a constitutive water accumulation in brain astrocytes. As a ubiquitous vehicle for transporting ions, transmitters and metabolites, water enters astrocytes via a wide range of ion channels, transporters and exchangers. For instance, they express Na^+/K^+ and Na^+/HCO_3^- cotransporters to dynamically regulate intra- and extracellular ion homeostasis (MacAulay, 2021). Standing as a major glial cell type controlling the neuropil environment, astrocytes take up synaptically released K^+ through inwardly rectified K^+ channels and neurotransmitters (e.g., glutamate, GABA) via high-affinity transporters to safeguard synaptic transmission (Dallerac et al., 2018). The transmembrane molecular exchange may lead to continuous water entry into astrocytes. Moreover, astrocytes juxtapose the cerebral vasculature, being a front relay station for brain metabolism. They express glucose transporters and lactate-permeable monocarboxylate transporters facilitating energy substrate uptake and transfer between blood vessels and neuron-glia networks (Cauli et al., 2023). As water is constantly produced during metabolic processes (Bonvento & Bolanos, 2021), astrocyte metabolism would also contribute to the constitutive water accumulation in the cytoplasm. An efficient efflux pathway appears necessary to counterbalance the constitutive water buildup, thereby maintaining astrocyte and brain homeostasis. Our data show that astrocyte AQP4 fulfills such a role by mediating tonic water outflow (Fig. S4), lending functional supports to the previous suggestion (Amiry-Moghaddam et al., 2003). It has been observed that knock out of AQP4 leads to water over accumulation in the brain, likely due to the reduction in water exit from the pial membrane (Haj-Yasein et al., 2011, Vindedal et al., 2016). Our result also echoes that in the principal cells of the kidney collecting

duct, AQP4 is suggested to mainly mediate water exit to balance the AQP2-sustained water entry

(Noda et al., 2010). Moreover, we observed that inhibiting astrocyte basal water efflux alters

Ca^{2+} activities in both astrocytes and neurons, reminiscent of the early finding that inhibition of

AQP4 by TGN-020 enhances cerebral blood flow in mouse cortex (Igarashi et al., 2013).

This study also provides mechanistic hints to understand AQP4-relevant pathologies in

specific contexts. For instance, astrocyte water accumulation and swelling are implicated in the

development of brain edema. Different types of edema have been delineated depending on the

pathological contexts, including such as cytotoxic, vasogenic and hydrocephalic subtypes. The

presence of AQP4 has been found to ameliorate vasogenic (vascular leak) and hydrocephalic

edema (Jeon et al., 2021, Tait et al., 2010), where the excessive water has been suggested to exit

the brain via AQP4-dependent route. This is consistent with our observation that astrocyte AQP4

contributes to basal water extrusion. Hence aquaporin facilitators would help to control

vasogenic and hydrocephalic edema. Differently, inhibiting AQP4 has been observed to

ameliorate cytotoxic edema encountered during cerebral ischemia (Verkman et al., 2011, Tait et

al., 2010). Beneficial effects of TGN-020 on edema have been reported in spinal cord injuries (Li

et al., 2019), in ischemia (Sun et al., 2022) and in retinal edema in diabetic animal models

(Oosuka et al., 2020). These observations suggest that the role of AQP4 in pathological

conditions is context-dependent. Indeed, while our results suggest astrocyte AQP4 mediates a

water outflow in basal states, the direction of water transport in specific contexts will be

determined by the actual transmembrane water gradient. As the origin of edema is variable and

water is among the many other ions involved in edema formation (Stokum et al., 2016), astrocyte

aquaporin may either be the primary cause for cytosolic water accumulation or secondarily

sustain protective water extrusion. In neuromyelitis optica spectrum disorders (NMOSD), an

unusual nervous autoimmune disorder often featured with circulating IgG autoantibody against

AQP4, the impairment in astrocyte water homeostasis has been observed (Hinson et al., 2012,

Lucchinetti et al., 2014). The inhibition of astrocyte AQP4 by autoantibody may initially affect

the basal water efflux. Yet chronic pathological evolution could convert astrocytes into reactive

states, with either or both the aquaporin expression, intra- and extracellular environments being

altered, the role of AQP4 in water transport might be accordingly shifted (Mireles-Ramirez et al.,

2022). Additionally, pro-inflammatory cytokines and chemokines during the immunological

maladaptation could also contribute to shape astrocyte functions including aquaporin-mediated

water transport (Lucchinetti et al., 2014, Mireles-Ramirez et al., 2022). Recent data show that

TGN-020 targeting of AQP4 mitigates the inflammation during ischemia-reperfusion injury, by

enhancing glymphatic amyloid clearance and inhibiting the ERK 1/2 pathway (Li et al., 2023),

suggesting that in pathological conditions, multiple factors could be recruited by astrocyte

aquaporin targeting.

We observed that acutely blocking aquaporin caused basal water accumulation leading to

swelling in astrocytes, and facilitated the initial occurrence of the evoked swelling by

hypoosmoticity and during cortical spreading depression. This recalls early observation by

electron microscopy that knocking out the anchoring protein alpha-syntrophin to disperse AQP4

clustering induces an enlargement of astrocyte endfeet (Amiry-Moghaddam et al., 2003). It

likely reflects astrocyte local swelling caused by the disruption of AQP4-mediated water efflux.

Our observation also parallels the report that AQP4 knock out facilitates astrocytes swelling

induced by hypoosmotic solution (Murphy et al., 2017). Nevertheless, genetic inactivation of

AQP4 also reveals inhibitory effects on astrocyte swelling (Benfenati et al., 2011, Mola et al.,

2016, Woo et al., 2018). Such discrepancy might be due to the variable compensations in brain

water content and structure integrity during chronic genetic manipulations (Binder et al., 2004, Gomolka et al., 2023, Haj-Yasein et al., 2011, MacAulay, 2021, Yao et al., 2015). In addition, though our data suggest that under basal condition, astrocyte AQP4 sustains a tonic water outflow, in conditions when massive water influx persists as exemplified by the application of hypoosmotic solutions, the direction of water transport of AQP4 would follow the imposed water gradient.

Astrocyte aquaporin modulates brain water transport (MacAulay, 2021), though its role in the glymphatic system is under deliberation (Mestre et al., 2018, Rasmussen et al., 2022, Smith & Verkman, 2019, Smith et al., 2017). Our data suggest that astrocyte aquaporin-mediated outflow helps to maintain local water environment in the brain parenchyma. Moreover, besides affecting water transport, targeting AQP4 also impacts astrocyte volume therefore the size of extracellular space. These two factors would need to be considered when evaluating AQP4 involvement in brain fluid transport.

We show by DW-MRI that water transport by astrocyte aquaporin is critical for brain water homeostasis. When blocking AQP4 with TGN-020, therefore its basal water outflow, we observed spatially heterogeneous elevations of diverse temporal kinetics in brain water diffusion rate. Our current data are derived from 5 min apart acquisitions, providing information over the early phase of AQP4 inhibition therefore extending our early report (Debacker et al., 2020). The regional heterogeneity likely reflects the various levels of AQP4 expression; its relative enrichment in the cerebral cortex (Clarke et al., 2018) corresponds to the pronounced effect observed here by DW-MRI. An overall increase in brain water diffusion rate was observed when blocking AQP4 water efflux. This treatment causes water accumulation inside astrocytes and their swelling, which reduces the extracellular space but increases the intracellular space. Water

diffusion would be enhanced inside astrocytes and decreased in extracellular space, respectively.

DW-MRI maps global water diffusion of both intra- and extracellular space. Likely, a net

increase in brain water diffusion may reflect its intracellular increase exceeding the extracellular

decrease. In addition, convective brain fluid flow has been suggested to be present in the

perivascular space (Smith & Verkman, 2019), a scenario extended by the glymphatic system to

the extracellular space of the parenchyma (Mestre et al., 2018). In this sense, transiently

squeezing extracellular space might increase the rate of the water flow. The current MRI data yet

lack sufficient spatial resolution to delineate the spatial compartmentalization of brain fluid flow.

Our study sheds light on the mechanisms by which astrocyte aquaporin contributes to the

water environment in brain parenchyma, and will help to understand the processes underlying

brain homeostasis and adaptation to life conditions.

Materials and methods

Animals and acute brain slice preparation

Experiments were undertaken in accordance with European Community guiding principles on the care and use of animals (86/609/CEE), and the rules of the host institute. For water transport and volume imaging, coronal slices comprising somatosensory cortex were acutely prepared from C57BL/6 mice of both sexes at ages of 4-6 weeks, unless otherwise indicated. Mice were deeply anesthetized by isoflurane (Axcience) evaporation in a closed plexiglass box. The brain was taken out and placed in a modified artificial cerebrospinal fluid (aCSF) for slicing (in mM: 30 NaCl, 4.5 KCl, 1.2 NaH2PO4, 1 MgCl2, 26 NaHCO3, 10 D-Glucose, and 194 sucrose) maintained at 4 °C during sectioning. The brain was cut into 300-µm thick slices with a vibratome (Leica VT1200S). Brain slices were recovered in standard aCSF (mM: 124 NaCl, 4.5 KCl, 1.2 NaH2PO4, 1 MgCl2, 2 CaCl2, 26 NaHCO3, and 10 D-Glucose) at 37 °C for about 1 h, and the same aCSF was used for brain slice imaging at room temperature. For optogenetic cortical spreading depression (opto-CSD) experiments, 18 to 21 day-old mice were deeply anesthetized with 150 µL of isoflurane (ISOVET, isoflurane, Piramal/healthcare). After the decapitation, the brain was quickly removed and placed into ice-colded oxygenated aCSF containing the following (in mM): NaCl 125; NaHCO3 26; Sucrose 15; Glucose 10; KCl 2,5; CaCl2 2; NaH2PO4 1,25; MgCl2 1; kynurenic acid 1 (nonspecific glutamate receptor antagonist, Sigma-Aldrich). Coronal slices from mouse somatosensory cortex (300µm thick) were cut with a vibratome (VT1000, Leica) and allowed to recover at room temperature for at least 45 min in aCSF saturated with O_2/CO_2 (95%/5%).

***In vivo* chemical and genetic targeting of astrocytes**

Mice of similar age were used also for *in vivo* labeling of astrocytes, where the astrocyte-specific

red fluorescent dye sulforhodamine B (10 mg/ml, Sigma) was intraperitoneally injected into

awake mice at a dose of 10 μL/g. The genetically encoded Ca^{2+} indicator GCaMP6 was

expressed in astrocytes *in vivo* by crossing a Cre-dependent GCaMP6f mouse line (Ai95, The

Jackson Laboratory) with an astrocyte-specific Glast-Cre^{ERT2} mouse line (Slezak et al., 2007).

Tamoxifen (T5648, Sigma) was dissolved in ethanol 100 % (at 1 mg/250μL) and vortexed until

tamoxifen is fully dissolved. Corn oil (sigma) were then added to have a final solution at 10%

ethanol and 90% oil. Solution is then heated at 37°c for 15 min and vortexed, and put in

sonicator for 15 min to reach transparency, and then aliquoted. All procedures were made under

light-proof protocols. Aliquots were stored at -20 °C until use. GCaMP6f/Glast-Cre^{ERT2} mice

were injected with tamoxifen at ~3–4 weeks of age, once a day (1 mg) for two consecutive days.

Genotyping of the Cre-dependent GCaMP6f mouse line (Ai95, The Jackson Lab) used the

standard primers and polymerase chain reaction protocols provided by the supplier. Genotyping

of GLAST-Cre^{ERT2} mouse line was performed using the primers for Cre recombinase

(TK139/141) as reported (Slezak et al., 2007). For Ca^{2+} imaging in somatostatin interneurons,

homozygous mice B6J.Cg-Sst$^{tm2.1(cre)Zjh}$/MwarJ (SST-Cre, Jackson Laboratory) (Taniguchi et al.,

2011) have been crossed with homozygous Ai95$^{GCaMP6f/GCaMP6f}$ mice to obtain heterozygous SST-

cre$^{cre/WT}$::Ai95$^{GCaMP6f//WT}$ mice. For opto-CSD, homozygous mice B6;129PEmx1tm1lr/J ((Gorski

et al., 2002), Emx1-Cre$^{cre/cre}$, Jackson Laboratory) have been crossed with homozygous mice

B6;129SGt(ROSA)26Sortm32.1(CAG-COP4*H134R/EYFP)Hze/J ((Madisen et al., 2012)

Ai32$^{ChR2/ChR2}$, Jackson Laboratory) to obtain heterozygous Emx1-cre$^{cre/WT}$::Ai32$^{ChR2/WT}$ mice. All

procedures using animals were carried out in strict accordance with French regulations (Rural

Code R214/87 to R214/130) and conformed to the ethical recommendations of the European

Economic Community (Directive 86/609/EEC) and the National Charter French on ethics in

animal experimentation. All protocols were approved by the Charles Darwin ethics committee

and submitted to the French Ministry of Education and Research (Approval 2015 061011367540

APAFIS#573-2015061011367547 v1).

***Ex vivo* fluorescence imaging**

Light sheet and epifluorescence imaging were both performed using a wide-field upright

microscope (Zeiss Axioskop 50, Germany) equipped with water-immersion objectives.

Epifluorescence illumination was provided by a monochromator light source (Polychrome II,

TILL Photonics, Germany) directly coupled to the imaging objective via an optical fiber. We

used water immersion imaging objectives (ZEISS): ×10 NA0.3 used to have a global view, ×20

NA0.5 used to perform light sheet imaging, and ×63 NA1.0 to image individual astrocytes under

epifluorescence configuration so as to collect their whole-cell SRB fluorescence to follow

astrocyte water transport and volume dynamics. The green fluorescence of GCaMP6 was

separated from the red fluorescence of SRB, by spectrally exclusive double-band filters (Di03-

R488/561-t3 and FF01- 523/610, Semrock). SRB-labeled astrocytes were excited at 550 nm.

Fluorescence signal was collected using a digital electron-multiplying charge-coupled device

(EMCCD Cascade 512B, Photometrics).

A light sheet imaging system was adapted from our previous publication (Pham et al.,

2020), where a chamber with a slightly tilted surface was used so as to spare the agarose

embedding and allow the light sheet to access to any parts of the brain slice (Fig. S1). Briefly,

the in-focus light sheet was introduced laterally from an independent optical module (Alpha3

light sheet add-on, Phaseview, France) equipped with an air objective (Zeiss EC EPIPlan ×10,

0.25NA). This optical module was orthogonal to the upright imaging pathway and coupled via a

wavelength combiner (Thorlabs) to two continuous wave lasers 473-nm and 561-nm (CNI,

China), for the excitation of GCaMP6 and SRB, respectively. The imaging chamber (20 × 30 ×

46 mm width, height and length) was custom-designed using 3D modeling software (Trimble

SketchUP), produced by a Mojo 3D printer (Stratasys) where the horizontal slice-supporting

surface in the chamber was tilted by 8° along the laser entry direction. In this way, the brain slice

could be laid on the slightly tilted surface while being stabilized with a conventional Harp slice

anchor grid (Warner Instruments), whereby the laterally generated laser light sheet could reach

local regions throughout the brain slices. The chamber was mounted on a motorized PI stage

(Physik Instrumente GmbH, Germany) for axial micro-manipulation. Laser excitation and image

acquisition were controlled by MetaMorph (Molecular Devices). Fluorescence dynamics of

individual astrocytes was analyzed with ImageJ software (NIH). Background averaged from non-

fluorescent subregion was subtracted from time-lapse images. Because single astrocytes typically

occupy an area of ~50-100 µm in radius (Appaix et al., 2012), regions of interest of 50-µm

diameter were delineated surrounding identified astrocyte footprint, to comprise the extended

area of individual astrocytes while minimizing interference with neighbors. Time courses

extracted from single regions of interest were normalized to the baseline (F_0) expressed as dF/F_0.

Dynamic imaging of the extracellular space and astrocytes volume changes

Individual slices were transferred to a submerged recording chamber and perfused with

oxygenated aCSF (~2 ml/min) lacking kynurenic acid. Slices were observed using a double port

upright microscope (BX51WI, Olympus, Japan) equipped with Dodt gradient contrast optics

(DGC; Luigs and Neumann), collimated light emitting device (LED; 780 nm, ThorLabs) as

transmitted light sources, ×4 (PlanN, 0.10NA, Olympus), ×20 (UMPlanFL N, 0.50 W, Olympus)

or ×40 (LUMPlanF/IR, 0.80 w, Olympus) objectives, and a digital camera (OrcaFlash 4.0,

Hamamatsu) attached on the front port of the microscope. The observation with ×4 allows to

delimit the cortical layers and to confirm the presence of "barrels" in layer IV. The ×20 objective

was used to observe infrared intrinsic optical signals (IOS) (Holthoff & Witte, 1996, MacVicar

& Hochman, 1991) of the cortical layers I to IV. The ×40 objective was used to image SRB

labelled astrocytes combined with IOS in layers I and II/III.

After one minute of baseline period, a 10-s continuous photostimulation at 470 nm of the

ChR2$_{H134R}$ (30% of the maximal power of the LED) was carried out via the epifluorescence port

of the microscope using a collimated LED system (CoolLED, PreciseExcite) and a set of

multiband filters consisting of an excitation filter (HC 392/474/554/635, Semrock), a dichroic

mirror (BS 409/493/573/652, Semrock) and an emission filter (HC 432/515/595/730, Semrock)

to observe the induced changes of light transmittance. The infrared and epifluorescence images

were acquired each at 1 Hz (exposure 100 ms, binning 2×2) using the Imaging Workbench 6.1

software (Indec Byosystems) for a total duration of 15 min.

***In vivo* fiber photometry**

The optical fiber (200 μm core, NA = 0.37, Neurophotometrics) coupled to a stainless-steel

ferrule (1.25 mm) was stereotaxically implanted in mouse S1 cortex (rostro-caudal 0 mm, lateral

+/- 2.5 mm, vertical -1.2mm), and fixed to the skull with dental cement (SuperBond, Sun

medical). Mice were habituated to the cage and fiber photometry setup for 30 minutes. Then,

fluorescence was recorded in mobile mice. 5-min baseline was recorded before SRB injection,

then SRB was applied to label brain astrocytes via intraperitoneal injection and the fluorescence

time course was continuously recorded. To examine *in vivo* the effect on astrocyte water

transport of acute AQP4 blocking, either TGN-020 (200 mg/kg; Tocris or MedChemExpress) or

saline control was intraperitoneally injected 1 h after SRB injection, a time point when astrocytes

were well labeled as confirmed by brain slice imaging.

Fluorescence signals were recorded using a Neurophotometrics fiber photometry system

(FP3002). A branched fiber optic patch cord (BFP_200/230/900-0.37_FC-MF1.25, Doric Lenses)

connected to the fiber photometry apparatus was attached to the implanted fiber optic cannula

using a cubic zirconia sleeve. To record fluorescence signals from SRB, excitation light from

560-nm LED was band-pass filtered, collimated, reflected by a dichroic mirror and focused by a

×20 objective. The excitation light power at the tip of the patch cord was 50 µW. Emitted

fluorescence was band-pass filtered and focused on digital camera. Signals were collected at a

rate of 1 Hz for SRB and visualized using the open-source software Bonsai 2.4 (http://bonsai-

rx.org) and analyzed in Igor software (Wavemetrics).

***In vivo* diffusion-weighted MRI**

To test the effect of astrocyte AQP4 acute inhibition on brain water homeostasis, we used

diffusion-weighted magnetic resonance imaging (DW-MRI) to calculate the apparent diffusion

coefficient (ADC) of tissue water molecules in the brain. MRI experiment was performed with

medetomidine anesthesia (0.6 mg/kg/h, i.v.). The mice were injected with either the water-

soluble TGN-020 sodium salt (Key Organics, LR-0041) or saline. For the crossover test, the

order of injection of TGN-020 or saline was randomly determined. The respiration was

monitored, and the body temperature was maintained at 36 °C throughout the measurement.

The MRI experiments were conducted on a Bruker 7T scanner with a cryogenic radio

frequency coil. After 10 min from the start of the scanning, either TGN-020 (200 mg/kg body

weight) or Saline was injected. Then, scanning was continued for 60 min. The respiration rate

was monitored and was confirmed within the range of 80-104 /min throughout the experiment.

Mice were injected with 200 mg/kg of TGN-020. We used a standard diffusion-weighted spin

echo EPI sequence, with the following parameters: spatial resolution = 175 × 175 × 500

μm^3/voxel, 3 b-values (b = 0, 250 and 1800 s/mm^2), 6 directions, 1 segment, echo time = 37.1 ms

/ repetition time = 5769 ms, bandwidth = 300 kHz, δ = 3 ms, Δ = 24 ms, scan time = 5 min. The

DWI scan was continued for 70 min (total 14 scans).

The ADC at each time point was calculated from DWI data using DSIstudio (https://dsi-

studio.labsolver.org/). The percentage change in ADC was then calculated according to the

equation: $\mathit{\Delta}ADC_i = (ADC_i / ADC_2 - 1) \times 100$ (%). The ADC_i is the ADC at the i-th timepoint.

We define the ADC at the 2nd time-point as the baseline because saline injection was performed

after the 2nd DWI scan. The averaged ADC changes within the regions of interests (ROIs) were

calculated using a homemade program. The ROIs of the cortex, the hippocampus, and the

striatum were created with reference to Allen mouse brain atlas.

Statistics

Experimental data are expressed as mean ± standard error unless otherwise mentioned. The *t*-test

was performed for two-group comparisons and significant difference was determined by p values

less than 0.05. Mann-Whitney U test was also used to ascertain the significance level, in

considering the deviation of the experimental data from the theoretical symmetrical distribution.

Bonferroni-Holm correction was used for multiple comparisons. Statistical tests were carried out

with Matlab (The MathWorks) and Statistica 6 (Statsoft). The significance levels are shown in

figures by *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Acknowledgements and funding sources

We thank the animal and imaging facilities of the IBPS (Sorbonne Université, Paris, France). We thank Nathalie Rouach for the GFAP-EGFP mice, Hervé Le Corronc and Thomas Panier for the discussion and the 3D printing of the light sheet chamber, respectively. This work was supported by the Agence Nationale de la Recherche (ANR-17-CE37-0010-03; ANR-20-CE14-0025-02), France Alzheimer individual fellowship (2021-2022), Japanese society for promotion of science (JSPS) 2022 summer grant, Grant-in-Aid for Challenging Research (Exploratory) in Japan (grant number 21K19464), the i-Bio initiative grant of Sorbonne Université. BLG was supported by a fellowship from the Fondation pour la Recherche sur Alzheimer.

Supplemental file

There are four supplementary figures (Fig. S1-S4).

Competing interest

The authors declare no competing financial interests.

Data sharing plans

The authors commit to share data, documentation, and code used in analysis.

References:

Abbott NJ, Pizzo ME, Preston JE, Janigro D, Thorne RG (2018) The role of brain barriers in fluid
movement in the CNS: is there a 'glymphatic' system? *Acta Neuropathol* 135: 387-407

Agnati LF, Marcoli M, Leo G, Maura G, Guidolin D (2017) Homeostasis and the concept of 'interstitial
fluids hierarchy': Relevance of cerebrospinal fluid sodium concentrations and brain temperature control
(Review). *Int J Mol Med* 39: 487-497

Amiry-Moghaddam M, Otsuka T, Hurn PD, Traystman RJ, Haug FM, Froehner SC, Adams ME, Neely
JD, Agre P, Ottersen OP, Bhardwaj A (2003) An alpha-syntrophin-dependent pool of AQP4 in astroglial
end-feet confers bidirectional water flow between blood and brain. *Proceedings of the National Academy
of Sciences of the United States of America* 100: 2106-11

Appaix F, Girod S, Boisseau S, Romer J, Vial JC, Albrieux M, Maurin M, Depaulis A, Guillemain I, van
der Sanden B (2012) Specific in vivo staining of astrocytes in the whole brain after intravenous injection
of sulforhodamine dyes. *PLoS One* 7: e35169

Barres BA (2008) The mystery and magic of glia: a perspective on their roles in health and disease.
Neuron 60: 430-40

Beaulieu C (2002) The basis of anisotropic water diffusion in the nervous system - a technical review.
NMR Biomed 15: 435-55

Benfenati V, Caprini M, Dovizio M, Mylonakou MN, Ferroni S, Ottersen OP, Amiry-Moghaddam M
(2011) An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for
cell-volume control in astrocytes. *Proceedings of the National Academy of Sciences of the United States
of America* 108: 2563-8

Binder DK, Papadopoulos MC, Haggie PM, Verkman AS (2004) In vivo measurement of brain
extracellular space diffusion by cortical surface photobleaching. *J Neurosci* 24: 8049-56

Bonvento G, Bolanos JP (2021) Astrocyte-neuron metabolic cooperation shapes brain activity. *Cell
metabolism* 33: 1546-1564

Brinker T, Stopa E, Morrison J, Klinge P (2014) A new look at cerebrospinal fluid circulation. *Fluids
Barriers CNS* 11: 10

Burnett ME, Johnston HM, Green KN (2015) Structural characterization of the aquaporin inhibitor 2-
nicotinamido-1,3,4-thiadiazole. *Acta Crystallogr C Struct Chem* 71: 1074-9

Cauli B, Dusart I, Li D (2023) Lactate as a determinant of neuronal excitability, neuroenergetics and
beyond. *Neurobiol Dis*: 106207

Cauli B, Porter JT, Tsuzuki K, Lambolez B, Rossier J, Quenet B, Audinat E (2000) Classification of
fusiform neocortical interneurons based on unsupervised clustering. *Proceedings of the National Academy
of Sciences of the United States of America* 97: 6144-9

Chung DY, Sadeghian H, Qin T, Lule S, Lee H, Karakaya F, Goins S, Oka F, Yaseen MA, Houben T,
Tolner EA, van den Maagdenberg A, Whalen MJ, Sakadzic S, Ayata C (2019) Determinants of
Optogenetic Cortical Spreading Depolarizations. *Cereb Cortex* 29: 1150-1161

Clarke LE, Liddelow SA, Chakraborty C, Munch AE, Heiman M, Barres BA (2018) Normal aging
induces A1-like astrocyte reactivity. *Proceedings of the National Academy of Sciences of the United
States of America* 115: E1896-E1905

Dallerac G, Zapata J, Rouach N (2018) Versatile control of synaptic circuits by astrocytes: where, when and how? *Nat Rev Neurosci* 19: 729-743

Debacker C, Djemai B, Ciobanu L, Tsurugizawa T, Le Bihan D (2020) Diffusion MRI reveals in vivo and non-invasively changes in astrocyte function induced by an aquaporin-4 inhibitor. *PLoS One* 15: e0229702

Eilert-Olsen M, Hjukse JB, Thoren AE, Tang W, Enger R, Jensen V, Pettersen KH, Nagelhus EA (2019) Astroglial endfeet exhibit distinct Ca(2+) signals during hypoosmotic conditions. *Glia* 67: 2399-2409

Fiacco TA, Agulhon C, Taves SR, Petravicz J, Casper KB, Dong X, Chen J, McCarthy KD (2007) Selective stimulation of astrocyte calcium in situ does not affect neuronal excitatory synaptic activity. *Neuron* 54: 611-26

Gaddamanugu S, Shafaat O, Sotoudeh H, Sarrami AH, Rezaei A, Saadatpour Z, Singhal A (2022) Clinical applications of diffusion-weighted sequence in brain imaging: beyond stroke. *Neuroradiology* 64: 15-30

Gomolka RS, Hablitz LM, Mestre H, Giannetto M, Du T, Hauglund NL, Xie L, Peng W, Martinez PM, Nedergaard M, Mori Y (2023) Loss of aquaporin-4 results in glymphatic system dysfunction via brain-wide interstitial fluid stagnation. *eLife* 12

Gorski JA, Talley T, Qiu M, Puelles L, Rubenstein JL, Jones KR (2002) Cortical excitatory neurons and glia, but not GABAergic neurons, are produced in the Emx1-expressing lineage. *J Neurosci* 22: 6309-14

Haj-Yasein NN, Vindedal GF, Eilert-Olsen M, Gundersen GA, Skare Ã, Laake P, Klungland A, Thorén AE, Burkhardt JM, Ottersen OP, Nagelhus EA (2011) Glial-conditional deletion of aquaporin-4 (Aqp4) reduces blood-brain water uptake and confers barrier function on perivascular astrocyte endfeet. *Proceedings of the National Academy of Sciences of the United States of America* 108: 17815-20

Harrison IF, Ismail O, Machhada A, Colgan N, Ohene Y, Nahavandi P, Ahmed Z, Fisher A, Meftah S, Murray TK, Ottersen OP, Nagelhus EA, O'Neill MJ, Wells JA, Lythgoe MF (2020) Impaired glymphatic function and clearance of tau in an Alzheimer's disease model. *Brain* 143: 2576-2593

Herrera Moro Chao D, Kirchner MK, Pham C, Foppen E, Denis RGP, Castel J, Morel C, Montalban E, Hassouna R, Bui LC, Renault J, Mouffle C, Garcia-Caceres C, Tschop MH, Li D, Martin C, Stern JE, Luquet SH (2022) Hypothalamic astrocytes control systemic glucose metabolism and energy balance. *Cell metabolism* 34: 1532-1547 e6

Hinson SR, Romero MF, Popescu BF, Lucchinetti CF, Fryer JP, Wolburg H, Fallier-Becker P, Noell S, Lennon VA (2012) Molecular outcomes of neuromyelitis optica (NMO)-IgG binding to aquaporin-4 in astrocytes. *Proceedings of the National Academy of Sciences of the United States of America* 109: 1245-50

Holthoff K, Witte OW (1996) Intrinsic optical signals in rat neocortical slices measured with near-infrared dark-field microscopy reveal changes in extracellular space. *J Neurosci* 16: 2740-9

Huber VJ, Tsujita M, Nakada T (2009) Identification of aquaporin 4 inhibitors using in vitro and in silico methods. *Bioorg Med Chem* 17: 411-7

Igarashi H, Tsujita M, Suzuki Y, Kwee IL, Nakada T (2013) Inhibition of aquaporin-4 significantly increases regional cerebral blood flow. *Neuroreport* 24: 324-8

Jeon H, Kim M, Park W, Lim JS, Lee E, Cha H, Ahn JS, Kim JH, Hong SH, Park JE, Lee EJ, Woo CW, Lee S (2021) Upregulation of AQP4 Improves Blood-Brain Barrier Integrity and Perihematomal Edema Following Intracerebral Hemorrhage. *Neurotherapeutics* 18: 2692-2706

Karagiannis A, Gallopin T, Lacroix A, Plaisier F, Piquet J, Geoffroy H, Hepp R, Naude J, Le Gac B, Egger R, Lambolez B, Li D, Rossier J, Staiger JF, Imamura H, Seino S, Roeper J, Cauli B (2021) Lactate is an energy substrate for rodent cortical neurons and enhances their firing activity. *eLife* 10

Kimelberg HK (2004) Water homeostasis in the brain: basic concepts. *Neuroscience* 129: 851-60

Le Bihan D (2014) Diffusion MRI: what water tells us about the brain. *EMBO Mol Med* 6: 569-73

Le Bihan D, Iima M (2015) Diffusion Magnetic Resonance Imaging: What Water Tells Us about Biological Tissues. *PLoS Biol* 13: e1002203

Le Bihan D, Urayama S, Aso T, Hanakawa T, Fukuyama H (2006) Direct and fast detection of neuronal activation in the human brain with diffusion MRI. *Proceedings of the National Academy of Sciences of the United States of America* 103: 8263-8

Li J, Jia Z, Xu W, Guo W, Zhang M, Bi J, Cao Y, Fan Z, Li G (2019) TGN-020 alleviates edema and inhibits astrocyte activation and glial scar formation after spinal cord compression injury in rats. *Life sciences* 222: 148-157

Li X, Xie Z, Zhou Q, Tan X, Meng W, Pang Y, Huang L, Ding Z, Hu Y, Li R, Huang G, Li H (2023) TGN-020 Alleviate Inflammation and Apoptosis After Cerebral Ischemia-Reperfusion Injury in Mice Through Glymphatic and ERK1/2 Signaling Pathway. *Mol Neurobiol*

Louveau A, Plog BA, Antila S, Alitalo K, Nedergaard M, Kipnis J (2017) Understanding the functions and relationships of the glymphatic system and meningeal lymphatics. *J Clin Invest* 127: 3210-3219

Lucchinetti CF, Guo Y, Popescu BF, Fujihara K, Itoyama Y, Misu T (2014) The pathology of an autoimmune astrocytopathy: lessons learned from neuromyelitis optica. *Brain Pathol* 24: 83-97

MacAulay N (2021) Molecular mechanisms of brain water transport. *Nat Rev Neurosci* 22: 326-344

MacVicar BA, Hochman D (1991) Imaging of synaptically evoked intrinsic optical signals in hippocampal slices. *J Neurosci* 11: 1458-69

Madisen L, Mao T, Koch H, Zhuo JM, Berenyi A, Fujisawa S, Hsu YW, Garcia AJ, 3rd, Gu X, Zanella S, Kidney J, Gu H, Mao Y, Hooks BM, Boyden ES, Buzsaki G, Ramirez JM, Jones AR, Svoboda K, Han X et al. (2012) A toolbox of Cre-dependent optogenetic transgenic mice for light-induced activation and silencing. *Nat Neurosci* 15: 793-802

Mestre H, Hablitz LM, Xavier AL, Feng W, Zou W, Pu T, Monai H, Murlidharan G, Castellanos Rivera RM, Simon MJ, Pike MM, Plá V, Du T, Kress BT, Wang X, Plog BA, Thrane AS, Lundgaard I, Abe Y, Yasui M et al. (2018) Aquaporin-4-dependent glymphatic solute transport in the rodent brain. *eLife* 7

Mestre H, Tithof J, Du T, Song W, Peng W, Sweeney AM, Olveda G, Thomas JH, Nedergaard M, Kelley DH (2018) Flow of cerebrospinal fluid is driven by arterial pulsations and is reduced in hypertension. *Nature communications* 9: 4878

Mireles-Ramirez MA, Pacheco-Moises FP, Gonzalez-Usigli HA, Sanchez-Rosales NA, Hernandez-Preciado MR, Delgado-Lara DLC, Hernandez-Cruz JJ, Ortiz GG (2022) Neuromyelitis optica spectrum disorder: pathophysiological approach. *Int J Neurosci*: 1-13

Mola MG, Sparaneo A, Gargano CD, Spray DC, Svelto M, Frigeri A, Scemes E, Nicchia GP (2016) The speed of swelling kinetics modulates cell volume regulation and calcium signaling in astrocytes: A different point of view on the role of aquaporins. *Glia* 64: 139-54

Murphy TR, Davila D, Cuvelier N, Young LR, Lauderdale K, Binder DK, Fiacco TA (2017) Hippocampal and Cortical Pyramidal Neurons Swell in Parallel with Astrocytes during Acute Hypoosmolar Stress. *Front Cell Neurosci* 11: 275

Noda Y, Sohara E, Ohta E, Sasaki S (2010) Aquaporins in kidney pathophysiology. *Nat Rev Nephrol* 6:
168-78

Ochoa-de la Paz LD, Gulias-Canizo R (2022) Glia as a key factor in cell volume regulation processes of
the central nervous system. *Front Cell Neurosci* 16: 967496

Oosuka S, Kida T, Oku H, Horie T, Morishita S, Fukumoto M, Sato T, Ikeda T (2020) Effects of an
Aquaporin 4 Inhibitor, TGN-020, on Murine Diabetic Retina. *Int J Mol Sci* 21

Papadopoulos MC, Verkman AS (2013) Aquaporin water channels in the nervous system. *Nat Rev
Neurosci* 14: 265-77

Pham C, Moro DH, Mouffle C, Didienne S, Hepp R, Pfrieger FW, Mangin JM, Legendre P, Martin C,
Luquet S, Cauli B, Li D (2020) Mapping astrocyte activity domains by light sheet imaging and spatio-
temporal correlation screening. *Neuroimage* 220: 117069

Rasmussen MK, Mestre H, Nedergaard M (2022) Fluid transport in the brain. *Physiol Rev* 102: 1025-
1151

Slezak M, Goritz C, Niemiec A, Frisen J, Chambon P, Metzger D, Pfrieger FW (2007) Transgenic mice
for conditional gene manipulation in astroglial cells. *Glia* 55: 1565-76

Smith AJ, Verkman AS (2019) CrossTalk opposing view: Going against the flow: interstitial solute
transport in brain is diffusive and aquaporin-4 independent. *J Physiol* 597: 4421-4424

Smith AJ, Yao X, Dix JA, Jin BJ, Verkman AS (2017) Test of the 'glymphatic' hypothesis demonstrates
diffusive and aquaporin-4-independent solute transport in rodent brain parenchyma. *eLife* 6

Stokum JA, Gerzanich V, Simard JM (2016) Molecular pathophysiology of cerebral edema. *Journal of
cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow
and Metabolism* 36: 513-38

Sun C, Lin L, Yin L, Hao X, Tian J, Zhang X, Ren Y, Li C, Yang Y (2022) Acutely Inhibiting AQP4
With TGN-020 Improves Functional Outcome by Attenuating Edema and Peri-Infarct Astrogliosis After
Cerebral Ischemia. *Front Immunol* 13: 870029

Tait MJ, Saadoun S, Bell BA, Verkman AS, Papadopoulos MC (2010) Increased brain edema in aqp4-
null mice in an experimental model of subarachnoid hemorrhage. *Neuroscience* 167: 60-7

Taniguchi H, He M, Wu P, Kim S, Paik R, Sugino K, Kvitsiani D, Fu Y, Lu J, Lin Y, Miyoshi G, Shima
Y, Fishell G, Nelson SB, Huang ZJ (2011) A resource of Cre driver lines for genetic targeting of
GABAergic neurons in cerebral cortex. *Neuron* 71: 995-1013

Toft-Bertelsen TL, Larsen BR, Christensen SK, Khandelia H, Waagepetersen HS, MacAulay N (2021)
Clearance of activity-evoked K(+) transients and associated glia cell swelling occur independently of
AQP4: A study with an isoform-selective AQP4 inhibitor. *Glia* 69: 28-41

Verkman AS, Ratelade J, Rossi A, Zhang H, Tradtrantip L (2011) Aquaporin-4: orthogonal array
assembly, CNS functions, and role in neuromyelitis optica. *Acta Pharmacol Sin* 32: 702-10

Verkman AS, Smith AJ, Phuan PW, Tradtrantip L, Anderson MO (2017) The aquaporin-4 water channel
as a potential drug target in neurological disorders. *Expert opinion on therapeutic targets* 21: 1161-1170

Vindedal GF, Thoren AE, Jensen V, Klungland A, Zhang Y, Holtzman MJ, Ottersen OP, Nagelhus EA
(2016) Removal of aquaporin-4 from glial and ependymal membranes causes brain water accumulation.
Mol Cell Neurosci 77: 47-52

Woo J, Kim JE, Im JJ, Lee J, Jeong HS, Park S, Jung SY, An H, Yoon S, Lim SM, Lee S, Ma J, Shin EY,
Han YE, Kim B, Lee EH, Feng L, Chun H, Yoon BE, Kang I et al. (2018) Astrocytic water channel
aquaporin-4 modulates brain plasticity in both mice and humans: a potential gliogenetic mechanism
underlying language-associated learning. *Mol Psychiatry* 23: 1021-1030

Xiao M, Hou J, Xu M, Li S, Yang B (2023) Aquaporins in Nervous System. *Adv Exp Med Biol* 1398: 99-
124

Yang J, Vitery MDC, Chen J, Osei-Owusu J, Chu J, Qiu Z (2019) Glutamate-Releasing SWELL1
Channel in Astrocytes Modulates Synaptic Transmission and Promotes Brain Damage in Stroke. *Neuron*

Yao X, Smith AJ, Jin BJ, Zador Z, Manley GT, Verkman AS (2015) Aquaporin-4 regulates the velocity
and frequency of cortical spreading depression in mice. *Glia* 63: 1860-9

Zhao HH, Du H, Cai Y, Liu C, Xie Z, Chen KC (2019) Time-resolved quantification of the dynamic
extracellular space in the brain: study of cortical spreading depression. *Journal of neurophysiology* 121:
1735-1747

A In vivo chemical labeling for FIT imaging

SRB injection | In vivo/slice imaging

~1hr

SRB (x20) SRB (x63)

SRB GFAP-EGFP Overlay

B

10 s
20 s
30 s
1 min
2 min
3min

H_2O

0.1 | 20 s

2 s
5 s
10 s
20s

H_2O

C

CTR or
TGN-020

**

0.005
20 s

CTR
TGN

0.01

0.00

Basal CTR Basal TGN

Downward (SRB)

AQP4 H_2O

D

In vivo
fiberphotometry

Astrocyte (SRB)

5.0

SRB TGN or CTR

20 min

post-CTR

post-TGN

2 min

1.0

0.0

-1.0

-2.0

**

Intensity
change (dF/F$_0$)

0.0

-0.2

-0.4

*

Oscillation
range (STD)

CTR TGN

FIGURE 1

Fig. 1. Astrocyte aquaporin mediates a tonic water efflux. (**A**) *In vivo* chemical labelling of astrocytes. Sulforhodamine B (SRB, 10 mg/ml) was intraperitoneally injected in awake mice (10 µl/g). Monochrome images show representative astrocyte labeling in living acute brain slices from cortex under low (×20; scale bar, 50 µm) and high magnification (×63; scale bar, 20 µm) by epifluorescence. *Below*, SRB labeling was confirmed to be astrocyte-specific in acute brain slices of the astrocyte reporter line GFAP-EGFP, where light sheet imaging was used to gain optical sectioning (Materials and methods; Fig. S1). Scale bar, 50 µm. (**B**) Optical imaging of astrocyte water transport in acute brain slices. Transmembrane water transport was triggered with hypo- and hypertonic solution, inducing water inflow and outflow that were respectively reflected by SRB fluorescence decrease and increase (expressed as dF/F_0). The hypo- or hypertonic solution was applied to slices over different lengths of duration displayed in colors corresponding to the time courses of SRB fluorescence (n = 52 astrocytes, 4 mice). (**C**) Acutely blocking astrocyte AQP4 with TGN-020 caused intracellular water accumulation and swelling. *Left*, while no effect was seen under CTR condition (vehicle only, n = 23 astrocytes), TGN-020 (20 µM) significantly decreased astrocyte SRB fluorescence (n = 30, 6 mice). Imaging was performed in acute brain slices of layer II/III S1 cortex. *Middle*, the downward slope was compared between the periods before and after the application of TGN-020. *Right*, illustration shows astrocyte aquaporin sustaining a tonic water efflux. Its blockade causes water accumulation and cell swelling. (**D**) *In vivo* validation of the effect of TGN-020 application on astrocyte water homeostasis. *Left*, fiber photometry was used for real-time recording of SRB fluorescence of astrocyte population in S1 cortex in freely moving mice, with saline (CTR) or TGN-020 being intraperitoneally injected when SRB was trapped in astrocytes. Fiber photometry recording shows that *in vivo* SRB injection resulted in rapid entry into mouse cortex and, in about one hour, led to astrocyte labeling (inset scale bar, 50 µm). *Middle*, example response to saline and TGN-020. Relative to CTR, TGN caused a decrease in astrocyte SRB fluorescence and its oscillation range (n = 8 recordings per condition, 5 mice).

FIGURE 2

Fig. 2. Acutely blocking astrocyte aquaporin induces swelling-associated Ca^{2+} oscillation. (**A**) *In vivo* expression in astrocytes of the genetically encoded fluorescent Ca^{2+} indicator GCaMP6 for imaging astrocyte Ca^{2+} in acute brain slices. Light sheet microscopy was used to capture transient Ca^{2+} signals of local regions. As a positive control, astrocyte swelling was induced by hypotonic solution (100 mOsM) that caused Ca^{2+} changes from their homeostatic level. *Left*, representative time courses of swelling-induced Ca^{2+} changes in detected response regions; *middle to left*, the histogram distribution and bar representation showing the signal strengths that were derived from the temporal integral of individual Ca^{2+} time courses normalized per minute, before and after cell swelling (n = 15 astrocytes of three mice). (**B**) Astrocyte Ca^{2+} oscillation caused by acute aquaporin blocking with TGN-020 (n = 31 astrocytes of 5 mice), due to the inhibition of the tonic water efflux that led to astrocyte swelling as illustrated. Time scale and Ca^{2+} rise scale (dF/F$_0$) are the same as in *A*. Scale bar, 50 µm. (**C**) Intercellular effect on SST interneurons of blocking astrocyte aquaporin water efflux. TGN-020 (20 µM) or the equal molar vehicle (CTR) was bath applied to acute cortical slices of SST-GCaMP6 mice (n = 7-12 measurements from 4 mice). Scale bar, 50 µm.

FIGURE 3

Fig. 3. Tonic water efflux via aquaporin modulates phasic transmembrane water transport and astrocyte volume response. (**A**) *Left*, in basal condition astrocyte aquaporin mediates a tonic water efflux. *Right*, protocol to induce water outflow from astrocytes, therefore their shrinking, by hypertonic extracellular solution (400 mOsM) in either control condition (CTR) or in the presence of AQP4 inhibitor TGN-020 (20 μM). (**B**) Time courses of astrocyte SRB fluorescence increase upon the phasically induced water outflow, reflecting the occurrence of shrinking. The histograms and bar charts compare the start time, namely the delay between hypertonic solution application and rise in SRB fluorescence, the time to reach the peak of shrinking, and the absolute amplitude of water outflow-induced SRB increase (n = 58 astrocytes for CTR, and 47 astrocytes for TGN-020, four mice). (**C**) *Right*, protocol to trigger water inflow into astrocytes by hypotonic extracellular solution (100 mOsM) in either CTR solution or in the presence of TGN-020 (20 μM). (**D**) Time courses of astrocyte SRB fluorescence decrease caused by water inflow, which also reflects concomitant cell swelling. In contrast to the observation with hypertonicity-induced water outflow and astrocyte shrinking, a reduction was observed for both the start time and the time-to-peak with TGN-020 (n = 12 astrocytes for CTR, and 12 astrocytes for TGN-020, four mice). TGN-020 led to a decrease in the absolute amplitude of astrocyte swelling.

FIGURE 4

Fig. 4. AQP4-mediated tonic water outflow regulates global swelling in cortical parenchyma. CSD-associated general swelling was induced by photostimulating ChR2-expressing pyramidal cells in acute cortical slices, and recorded by imaging intrinsic optical signal (IOS) with infrared illumination. (**A**) Representative recording. The pia surface is on the upper side; green, blue and red squares correspond to regions of interest in Layers I, II-III and IV, respectively. Transmittance signals are represented in pseudocolor images at different time points post photostimulation. Scale bar, 100 μm. (**B-C**) Kymograph and time courses showing the IOS changes following photoactivaiton, derived from the radial line of interest indicated by a black arrow in *A*. The light blue line indicates the 10-s photostimulation that increases the infrared transmittance signal (dT/T_0) across cortical layers, as also illustrated by the time courses; *a*, *b*, *c* and *d* correspond to the time points depicted in *A*. After the onset delay (*a*), the first (*b*) and second (*c*) peak of IOS are characteristic of the CSD and a prolonged general swelling, respectively. (**D-E**) TGN-020 (20 μM) inhibition of AQP4 reduced the initial onset and the maximal amplitude of both the CSD and general swelling (n = 6 - 18 measurements from 13 mice per condition) in layer II/III cortex. Inhibiting spiking activity with TTX (1 μM) only affected the amplitude of the initial CSD response. (**F**) Astrocytes swelling, reflected by SRB fluorescence decrease, monitored in control condition (n = 75 astrocytes) and in the presence of TGN-020 (n = 17 astrocytes) in layer II/III cortex.

FIGURE 5

Fig. 5. Perturbing the tonic water efflux via astrocyte aquaporin alters brain water homeostasis. *In vivo* DW-MRI (7T) was employed to map water diffusion in the entire brain scale following the acute inhibition of astrocytic AQP4 (TGN-020, intraperitoneal injection, 200 mg/kg), with paralleled control experiments performed with saline injection. (**A**) Experimental protocol for DW-MRI. Saline or TGN020 was injected at 10 min after the start of acquisition. DWI was acquired every 5 min. (**B**) Representative image obtained at three different b values to derive the water diffusion rates. (**C**) *Left*, brain water diffusion rate was mapped by the calculation of apparent diffusion coefficient (ADC). *Right*, representative images illustrating the relative changes of ADC at 60 min after injection of saline or TGN-020. (**D**) Time courses depicting the temporal changes in ADC in the cortex, striatum, and hippocampus, revealing the regional heterogeneity. Arrowhead indicates the injection of saline or TGN-020 (n = 10 mice for saline injection, 9 mice for TGN-020).