Identification of Determinants that Allow Maintenance of High-Level Fluoroquinolone

Resistance in *Acinetobacter baumannii*

Efrat Hamami[1], Wenwen Huo[1], Juan Hernandez-Bird[1], Arnold Castaneda[3], Jinna Bai[2], Sapna

Syal[1], Juan C. Ortiz-Marquez[4,5], Tim van Opijnen[5], Edward Geisinger[1,2] and Ralph R. Isberg[1,*]

[1]Department of Molecular Biology and Microbiology, Tufts University School of Medicine, 150

Harrison Ave., Boston, MA 02111, USA.

[2]Department of Biology, Northeastern University, Boston, MA 02115, USA

[3]Department of Biology, Tufts University, Medford, MA 02155, USA

[4]Department of Biology, Boston College, Chestnut Hill, MA 02135, USA

[5]Innovation Laboratory, Broad Institute, Cambridge, MA 02412, USA

[*]Corresponding Author

Abstract

Acinetobacter baumannii is associated with multidrug resistant (MDR) infections in

healthcare settings, with fluoroquinolones such as ciprofloxacin being currently ineffective.

Clinical isolates largely harbor mutations in the GyrA and TopoIV fluoroquinolone targets, as

well as mutations that increase expression of drug resistance-nodulation-division (RND) efflux

pumps. Factors critical for maintaining fitness levels of pump overproducers are uncharacterized

despite their prevalence in clinical isolates. We here identify proteins that contribute to the

fitness of FQR strains overexpressing three known RND systems using high-density insertion

mutagenesis. Overexpression of the AdeFGH efflux pump caused hypersensitization to defects

in outer membrane homeostatic regulation, including lesions that reduced LOS biosynthesis and

blocked production of the major *A. baumannii* porin. In contrast, AdeAB pump hyperexpression,

in the absence of elevated *adeC* expression (the outer membrane component of the pump), was

relatively tolerant to loss of these functions, consistent with the outer membrane protein being

the primary disruptive component. Surprisingly, overexpression of proton-transporting efflux

pumps had little impact on cytosolic pH, consistent with a compensatory response to pump

activity. The most striking transcriptional changes were associated with AdeFGH pump

overexpression, including the activation of the phenylacetate (PAA) degradation regulon.

Disruption of the PAA pathway resulted in cytosolic acidification and defective expression of

genes involved in protection from oxidative stress. These results indicate that RND efflux pump

overproduction is compensated by maintenance of outer membrane integrity in *A. baumannii* to

facilitate fitness of FQR isolates.

Importance

Acinetobacter baumannii is a pathogen that often causes multidrug resistant (MDR) infections in healthcare settings, presenting a threat to the efficacy of known therapeutic interventions. Fluoroquinolones such as ciprofloxacin are currently ineffective against a majority of clinical A. baumannii isolates, many of which express pumps that remove this antibiotic class from within the bacterium. Three of these pumps can be found in most clinical isolates, with one of the three often hyperproduced at all times. In this study we identify proteins that are necessary for the fitness of pump hyperproducers. The identified proteins are necessary to stabilize the outer membrane and allow the cytoplasm to tolerate the accumulation of ions as a consequence of excess pump activity. These results point to strategies for developing therapies that combine known antibiotics with drugs that target proteins important for survival of strains hyperexpressing efflux pumps.

Introduction

Acinetobacter baumannii is a Gram-negative bacterium primarily associated with diseases in healthcare settings (1-3). The microorganism has the potential to cause a diverse range of diseases including pneumonia, wound, and urinary tract infections (1-3). Of great concern is its role in causing ventilator-associated pneumonia, which may be closely tied to its ability to grow in biofilms on indwelling devices (2-6). Particularly challenging is the high incidence of multidrug resistant (MDR) A. baumannii in hospitalized patients, leaving afflicted individuals with few treatment options (1-3). Although there was a reduction in nosocomial cases

in the US from 2013 to 2019, the SARS-CoV2 pandemic saw a large jump in cases that

illustrates the difficulties in suppressing diseases by this organism (3). Furthermore, the

increasingly common resistance profiles of clinical isolates that are refractory to most

antimicrobial therapy regimens indicate that the pathogen poses a continued threat to successful

treatment (1-3, 7-10). These clinical isolates include a high proportion that are multi-drug

resistant (MDR) or have decreased susceptibility to various single agents (1-3, 7-10). This has

prompted the World Health Organization to report that *A. baumannii* is a high priority candidate

for research and development of new antimicrobials (8). For this reason, there is a high demand

for novel antimicrobials or improved combination therapies with which we can combat this

pathogen.

There are a variety of means by which *A. baumannii* acquires resistance, including the

acquisition of drug resistance elements via horizontal gene transfer (HGT), the selection of

mutations that inactivate drug target sites, and hyper-expression of preexisting drug efflux pumps

(11-15). Pathogens not only harbor a multitude of efflux systems, some of which are acquired by

HGT, but many have broad substrate specificities (11-15), stimulating efforts to discover

effective pump inhibitors (16). Unfortunately, there has been no indication of progress in

bringing these inhibitors to the clinic, although several have proven effective in interfering with

drug action or synergizing with clinically useful antibiotics using laboratory culture conditions or

in animal models (16). Nevertheless, given the relentless patterns of resistance acquisition after

the use (and misuse) of single-target antibiotics, a treatment regimen that uses a combination of

readily available antimicrobials working in synergy could be effective in interfering with

resistance evolution. This study approaches this concept by determining candidate proteins that

could be targeted together with clinically-proven classes of antibiotics.

One drug class that has reduced effectiveness against nosocomial pathogens is the

fluoroquinolones (2, 17). Resistance in clinical isolates primarily selects for mutations that

disrupt drug binding to the targets topoisomerases II and IV, encoded for by *gyrA* and *parC*

genes respectively (17). In *A. baumannii*, the highest levels of fluoroquinolone resistance (FQR)

are found in isolates that have increased expression of drug efflux pumps in addition to

mutations in the drug targets. These drug efflux pumps belong to the resistance-nodulation-

division (RND) family, in which overexpression is commonly driven by mutations in regulatory

elements that result in constitutive expression (18-23). The three major efflux pumps commonly

associated with FQR in *A. baumannii* clinical isolates are encoded for by *adeAB*, *adeFGH*, and

adeIJK, which are controlled by *adeRS* (encoding a two-component system), *adeL* (a LysR type

regulator), and *adeN* (a TetR repressor) regulators respectively (18-23). Among these three, the

AdeN-regulated pump is associated with intrinsic resistance to fluoroquinolones, as this is the

only RND pump in *A. baumannii* that has measurable levels of expression in the absence of

activating mutations (17, 22, 23).

These pump systems traverse both membranes (24-30), and overproduction requires that

the envelope accommodates an elevated population of protein complexes not present in drug-

sensitive isolates, putting potential stress on the inner and outer membrane. Therefore, the

stability and assembly of these proteins, as well as the fitness of *A. baumannii* during exponential

growth may rely upon bacterial functions that work to maintain cell envelope integrity or prevent

intrinsic toxicity of the overexpressed efflux system. Our work focuses on identifying genetic

interactions in these high-level FQR strains that allow them to maximize fitness, with the goal of

identifying proteins necessary to buffer the deleterious effects of pump hyperexpression that

could lead to decreased fitness levels in these strains. Such mutations are predicted to identify

targets for drugs that are specifically toxic for pump overproducers.

In this study, we used dense transposon sequencing and transcriptomics strategies to

identify proteins necessary to maintain fitness of *A. baumannii* efflux pump overexpressing

strains. We demonstrate that strains harboring overexpressing mutations in *adeN* and *adeS* show

surprising robustness and identify targets of vulnerability in overexpressers of the AdeL-

regulated pump. These targets are central players in maintaining cell envelope homeostasis. We

also connect the expression of the phenylacetate catabolic pathway as a response to pump

hyperexpression that leads to maintaining cytosolic pH and maximizing protection from reactive

oxygen species.

Results

Identification of mutations that depress fitness of fluoroquinolone-resistant strains

overexpressing RND efflux pumps. To identify genes that are important for maintaining the

fitness of strains having high-level fluoroquinolone resistance (FQR), we isolated mutations in

the *A. baumannii* ATCC 17978UN strain background that are viable in high concentrations of

ciprofloxacin (CIP; Materials and Methods). These strains were then subjected to a mutagenesis

strategy (Fig. 1A) that identified insertion mutations that had lowered viability in an FQR strain

background compared to the wild type (called strain WT) (Fig. 1B). Three FQR strains were

interrogated, each of which harbored *gyrA*(S81L) and *parC*(S84L) mutations that are responsible

for fluoroquinolone resistance in most Gram-negative organisms and are commonly found in the

clinic (called strain GP). The three strains were distinguished from each other by having distinct

mutations in loci that regulate the expression of clinically relevant *A. baumannii* resistance-

nodulation-division (RND) drug efflux pumps. The mutation in a component of a two-

component regulatory system *adeS*(R152S) upregulates the AdeAB pump, the activating

adeL(ΔI335A336) mutation upregulates the AdeFGH pump, while a knockout in the repressor

adeN::ISAba1 increases expression of the AdeIJK pump (Fig. 1C). It should be noted under

conditions of AdeAB pump overexpression, AdeK is used as the outer membrane component

which is expressed at wild type levels in the presence of the *adeS*(R152S) pump activating

mutation (Fig. 1C) used in this study (31, 32).

Transposon (Tn) mutagenesis was used to determine candidates important for viability or

fitness in either the absence or presence of drug. Between 110,000-150,000 unique transposon

mutants were created in all three FQR and the WT strain backgrounds, reaching pool saturation

levels of about 37-50% based on targets predicted for the *himar-1* transposon (Materials and

Methods). Each bank was sequenced post-harvest and transposon-proximal sequences were

amplified to identify the locations of insertions in each strain background and to determine

overall pool saturation. This sequence data was also utilized to identify insertion site

distributions that were present in each strain, because insertions in genes predicted to be essential

or cause fitness defects in a specific pump overproducer strain are predicted to be

underrepresented relative to the WT strain. Per our first approach (Fig. 1A), insertion abundance

throughout the genome was compared between strain backgrounds using the TRANSIT

resampling method to identify genes that have fewer insertions relative to WT in the absence of

drug (33) (Fig. 2, Data Set S1), as candidates for genes that are putative conditionally essential

targets. To focus on strain-dependent conditional essentiality, 445 genes characterized as

essential by TRANSIT's Gumbel method (34, 35) were excluded from this and downstream

analyses using transposon mutagenesis. Data were displayed by the statistical significance as a

function of the insertion abundance in each of the pump overproducer strains relative to WT

(Fig. 2A-C). Among genes identified in this fashion (passing significance criteria adj. $p<0.05$) in

the absence of fluoroquinolones were *psiE* (*adeS* background), *sodB* (*adeN* background), and

"nuc-diph-s-epim" or "n.d.s.e" (*adeL* background), respectively encoding a phosphate starvation

inducible locus, superoxide dismutase, and a nucleoside diphosphate sugar epimerase more

recently implicated in LOS assembly (36) (Fig. 2D-F).

Prior to the exclusion of essential genes, this analysis revealed that *cyoAB* encoding

components of the essential ubiquinol oxidase complex harbored significantly fewer insertions in

GPL than in WT (Fig. S1A, B). Its function during aerobic respiration as a generator of the

proton motive force may play a role in allowing the bacterium to tolerate proton influx resulting

from the drug-proton antiport activity of the RND efflux pump (37, 38). To provide support that

lowered Cyo activity reduces fitness in the presence of *adeL*ΔI335A336, we used CRISPRi to

knock down expression of the *cyoABCDE* operon in the GPL and WT strains and observed

growth in broth culture. To this end, an anhydrotetracycline (aTc) -inducible dCas9 was inserted

at the *att*Tn7 locus in both strains and transformed with a plasmid harboring a constitutive single

guide RNA (sgRNA) (Materials and Methods). As a control, the non-targeting original vector

(labeled NC-sgRNA; Table S1) was used. Growth was monitored in cultures grown for 5.5 or 6

hours after dilution to low density in LB and aTc. We saw a fitness defect when expression of

the ubiquinol oxidase system was knocked down in both the WT and the GPL strains, with the

defect enhanced in WT treated with 100ng/mL aTc (Fig. S1C). Quantification of *cyoABCDE*

operon transcript levels after aTc addition showed that expression was silenced in both strains at

comparable efficiency compared to controls (Fig. S1D).

Notably, there were very few insertions in both the WT and GPL strains which is an

indication of gene essentiality. As a consequence, although the locus was distinguished among

those showing the most highly significant difference in insertion efficiencies (Fig. S1B), these

differences were measured by degrees of essentiality (Fig. S1E). Furthermore, the average

number of viable insertion mutants per *himar-1* transposon location (TA) among the cumulative

WT pools was approximately 258 and in GPL about 440. While the loci highlighted from these

analyses demonstrate significantly different insertion efficiencies, these differences were likely

artifacts of plating efficiency rather than conditional essentiality. Therefore, a second strategy

was pursued to measure fitness defects during growth in broth culture to attempt to identify more

clearcut differences between the two strain backgrounds.

The FQR mutants have increased vulnerability to disruption of cell envelope

homeostasis. To attack this problem using a second approach, multiple small Tn insertion pools

(~8-16k unique insertions) were constructed in each background, and each was grown for 7-8

generations in the absence of antibiotics to determine the relative fitness levels of individual gene

disruptions (>79,000 unique insertions in total for each background; Materials and Methods).

The difference in fitness levels between an individual pump overproducer and the wild type was

then determined, and statistical significance was plotted as a function of fitness differences in the

two backgrounds (Fig. 3A-C). In the absence of drug, the GPS mutant was particularly tolerant

of insertion mutations relative to the WT control (Fig. 3A; Data Set S2). For the GPN and GPL

strains in the absence of drug, among the proteins identified are those affecting membrane

integrity including those that modulate lipid trafficking and LOS synthesis (Fig. 3B-C; Data Set

S2). Notably, the GPS strain which shows no upregulation of the outer membrane component of

the RND systems was the most tolerant of disruption in these pathways (Fig. 3A).

To determine if antibiotic treatment increased the spectrum of mutations that sensitize

pump overproducers, each of the strains was treated at various CIP concentrations to identify

conditions that resulted in 20-30% growth rate inhibition (Fig. 3D). Using drug concentrations

that reproduced these growth rates, each of the insertion banks was grown 7-8 generations in the

presence of CIP (17), cultures were harvested, gDNA was isolated, insertion sites were

selectively amplified and sequenced, and the abundance of each insertion was determined using

the MAGenTA pipeline (17, 39, 40) (Materials and Methods). From these datasets, the fitness

scores of gene disruptions were normalized to a LOWESS curve to avoid the position bias

exhibited with CIP treatment (Fig. S2). The differences in fitness levels between an individual

pump overexpresser and the wild type were then used to identify mutants that preferentially

hypersensitize the pump overexpressing strain to CIP (Fig. 3E-H; Data Set S2).

In general, treatment with drug increased the number of insertion mutations that resulted

in fitness defects in the pump hyper expressing strains compared to WT (compare Figs. 3A-C

and 3E-H). Primary targets that preferentially sensitize at least two FQR strains during CIP

treatment were concentrated in loci related to envelope homeostasis, including *bfmS* which

affects a collection of cell envelope processes (41) and *pbpG* which is involved in peptidoglycan

degradation (40) (Figs. 3E-H). Even so, the magnitude of defects was consistently higher for

mutations in GPL compared to the GPS background, with mutations in the GPN background

often showing intermediate effects (Fig. 3E). As in the absence of drug, the GPS strain showed

higher tolerance than the GPL and GPN strains for mutations in the same pathways (Fig. 3E-H).

Knockout mutations of candidates were generated as gene deletions in both the FQR and

parent strains via homologous recombination to verify their phenotypes (their genotypes

confirmed by whole genome sequencing, Materials and Methods). The targets for deletion were

chosen based on the following criteria: 1) they resulted in significant fitness defects in at least

two of the FQR strains ($q \leq 0.05$); and 2) they had severe fitness defects in at least one FQR

strain (fitness defect ≥ 0.25). Bacterial growth was then evaluated in the presence of CIP (Figs.

4, 5) at concentrations that yielded comparable growth inhibition across strains (Fig. 3D). Based

on Tn-seq analysis (Figs. 3E-H), insertion mutants in *bfmS* and *ompA$_{Ab}$* showed increased CIP

sensitivity in GPN and GPL compared to the WT strain background (Fig. 4A). Consistent with

these findings, growth defects caused by the deletion mutations were exacerbated in the pump

overexpressing strains after CIP treatment as compared to the WT strain background (Fig. 4B).

Of particular interest was the behavior of each of the mutants in the GPL background. Both the

Δ*bfmS::aacC1* and Δ*ompA* mutants were highly attenuated for growth in the presence of CIP,

with growth plateauing far below that observed for controls (Fig. 4B). For instance, when

Δ*bfmS::aacC1* was crossed into the GPL background, growth plateau was observed at about 10X

lower culture densities than that observed for the GPL parent and 12-20X lower than the same

mutation crossed into WT (Fig. 4B). Similarly, when the Δ*ompA* mutation was crossed into

GPL, plateaus occurred at 5-8X lower culture density than the GPL parent and about 8X lower

than the same mutation crossed into WT (Fig. 4B).

To ensure the growth inhibition was a result of the pump regulator mutation, the

consequences of crossing the mutations into the lower-level resistance GP strain were analyzed

in comparison to GP carrying a pump-activating mutation in each of the pump regulators (Fig.

S3). This allowed a direct comparison between strains differing by a single FQR mutation. The

increase in drug sensitivity was more pronounced in the FQR pump mutants carrying the

$\Delta ompA_{Ab}$ and $\Delta bfmS::aacC1$ mutations relative to GP with the same lesions (Fig. S3A).

The pump-overexpressing mutations alter regulatory elements that could be modulating

the expression of other unidentified genes that cause fitness alterations. To eliminate this

possibility, we determined if the fitness defect of the GPL strain harboring the $\Delta ompA_{Ab}$ and

$\Delta bfmS::aacC1$ mutations was reversed by deleting the *adeFGH* efflux pump genes. Deletion of

adeFGH reversed the fitness defects caused by the $\Delta ompA_{Ab}$ and $\Delta bfmS::aacC1$ mutations,

consistent with pump overproduction causing the observed fitness defects (Figs. S4 F and S5 F

respectively). Furthermore, for each mutation tested, the GP strain background showed

comparable fitness to the GPL$\Delta adeFGH$ strain background (Figs. S4 H & S5 G).

The GPL$\Delta ompA_{Ab}$ strain (Fig. 4B) was found by whole genome sequencing (WGS) to

harbor an additional mutation in *lpxK* (I283V), so we created additional strains harboring

$\Delta ompA_{Ab}$ via homologous recombination in the GPL background (Materials and Methods) and

tested their phenotypes relative to that of GP. The heightened sensitivity of GPL to CIP after

introduction of the $\Delta ompA_{Ab}$ lesion was retained (Fig. S4 B-D), indicating that the fitness defect

was due to the loss of OmpA. Similarly, the GP$\Delta bfmS::aacC1$ strain in Fig. S3A was found to

have an additional mutation upstream of the *bfmR* gene. This mutation dampened the expression

of *bfmR* (Fig. S5H; strain #272 GP$\Delta bfmS::aacC1$ (Fig. S3A) compared to strains #271

GP$\Delta bfmS::aacC1$ (Fig. S5B) and #304 $\Delta bfmS::aacC1$ (Fig. 4B)). Additional clones cured of this

bfmR promotor mutation were created in the GP strain background, and knockout of *bfmS*

reproducibly showed minimal fitness alterations of the GP parent in the presence of CIP, while

still sensitizing the *adeL* variant that overexpresses the efflux pump (Fig. S5B-E).

Given the strong defects observed with these two mutants, other candidate mutations

predicted to interfere with envelope biogenesis were interrogated, including *lpsB* and *pbpG*,

which showed lowered insertion mutation fitness in the pump hyper-expressors when compared

to WT, particularly in the GPL strain background (Fig. 5A). Deletion mutations were constructed

in WT, GP, GPN and GPL backgrounds, and growth of the eight mutant strains was monitored in

the presence of CIP. Loss of PbpG from either the GPN or GPL strain background was

particularly detrimental to growth in the presence of low CIP concentrations (Fig. 5). In both

GPN and GPL backgrounds harboring Δ*pbpG*, strains initiated growth but culture densities

stopped increasing after about 3 hours (Fig. 5B; Fig. S3B). In contrast, the parental GP mutant

harboring Δ*pbpG* did not show this phenotype (Fig. S3B). In the case of the Δ*lpsB* mutant,

growth plateaued at lower levels when the deletion was analyzed in the GPL mutant (8X lower

than parent GPL) compared to GPN (5-7X lower than GPN), consistent with the

adeL(ΔI335A336) mutation being particularly susceptible to envelope stress (Fig. 5B).

To ensure that this phenotype was due to these null mutations, both deletions were

complemented *in trans*. The presence of plasmids harboring these genes restored the fitness of

both pump overexpression strains, with growth in the presence of CIP indistinguishable from the

appropriate parents harboring empty vector (Fig. 5C). WGS of these strains uncovered an *mrcB*

(G539C) mutation in the GPNΔ*lpsB* background. Complementation of the GPNΔ*lpsB* strain,

however, restored growth to levels of the parental strain, indicating that the growth defect was

caused by loss of *lpsB* and not due to the secondary missense mutation (Fig. 5C). Furthermore,

when a deletion of the *adeFGH* efflux pump operon was introduced into the GPL strain

harboring either the Δ*lpsB* or the Δ*pbpG* mutations, both growth defects were alleviated,

indicating that defective growth was linked to the overproduction of the efflux pump (Figs. S6-

7).

The absence of PbpG or LpsB has been associated with alterations in *A. baumannii* LOS

homeostasis (36, 40). Therefore, LOS profiles of strains disrupted in *pbpG* and *lpsB* were

determined by fractionation on SDS-polyacrylamide gels (Fig. S8; Materials and Methods). No

significant differences in LOS profiles were detected between the WT or FQR strain

backgrounds (Fig. S8A). For the various strains harboring the Δ*pbpG* mutant, there were

decreased levels of a species unrelated to LOS, but otherwise the migration and relative

concentration of LOS species appeared unaffected (Fig. S8B). In contrast, as expected, Δ*lpsB*

mutations resulted in truncated LOS, with similar profiles in each of the efflux pump

backgrounds (Fig. S8C). Therefore, LOS is altered by these mutations as previously described,

and these alterations were independent of the strain background tested (40).

As reduced A_{600} yields could be indicative of morphological differences between mutants,

the various strains were imaged. As expected, the *bfmS*::*aacC1* and Δ*pbpG* mutants showed

morphological alterations that have been previously documented, ranging from stubbier to

slightly rounded and enlarged, dependent on the mutation (41, 42). The latter was observed with

Δ*lpsB* as well. These alterations were independent of pump overproduction (Fig. S9). In contrast,

the Δ*ompA* mutation resulted in no morphological changes (Fig. S9). In response to CIP, the

parental strain elongated as anticipated (43, 44), while the mutations that showed morphological

changes continued to appear different from their parent strain, with changes ranging from

reduced length to wider cells (Fig. S10). While structural deformation from loss of PbpG appears

slightly more severe in the pump overexpressing strain background (Fig. S11), the specific

morphological alterations caused by the mutants were observed both in the presence and absence

of pump hyperexpression (Figs. S11-14). Critically, what differentiated the morphotypes of these

lesions between the pump overexpressing strain and parent strain is that cells in the pump

overexpressing strain background consistently displayed larger volumes (Figs. S11-14). We

conclude that although the mutations that resulted in enhanced CIP sensitivity in pump

overproducer strains resulted in altered morphologies, it is unlikely these changes in morphology

explain for the reduced culture densities observed in growth curves of strains that overproduce

efflux pumps (Figs. 4 and 5).

Maintenance of pH homeostasis in the presence of pump overproduction. The RND-

class efflux pumps are known to function as drug-proton antiporters (45). With this in mind, we

investigated whether FQR pump hyper-expression causes *A. baumannii* cytosolic pH to become

more acidic in the presence or absence of CIP substrate. WT, the high-level FQR strains, and the

GP precursor strains were grown to mid-exponential phase and loaded with BCECF-AM dye to

determine internal pH (pHi). The pHi values were then determined from a standard curve of cells

incubated in pH-adjusted buffers in the presence of CCCP to equilibrate external and internal pH

(Materials and Methods). In the absence of antibiotic, the cytosol of GPN, the AdeIJK pump

over-expressor, was significantly more basic than the GP parent strain (Fig. 6A), while the

cytosolic pH values of the other pump overproducers were otherwise comparable to GP or WT

controls (Fig. 6A; one-way ANOVA, multiple comparisons between all strains and GP, * where

$p<0.05$). To determine if the presence of CIP, which should result in proton-CIP antiport, affects

internal pH, the experiment was repeated after exposure to CIP at concentrations yielding

comparable levels of growth inhibition across strains. CIP treatment lowered the pHi compared

to the untreated conditions for all strains. The cytosols of the GPS and GPL strains were

fractionally more acidic than the parental GP strain after CIP addition, but the differences did not

rise to the level of statistical significance (Fig. 6B; same tests as in 6A). Imaging shows these

strains exhibit similar cell morphologies both in the absence (Fig. S15) and presence of CIP (Fig.

S16). These data indicate there may be homeostatic mechanisms to neutralize the effects of

pump overexpression that act at either transcriptional or post-transcriptional levels, so the global

transcriptional effects of pump overexpression were analyzed in the presence and absence of CIP

(Fig. 1A, third approach).

Large transcriptional changes in response to pump overproduction are limited to the

AdeL-regulated pump. To determine if pump overproduction results in transcriptional changes

that support fitness, the FQR, low-level resistance GP, and WT strains were cultured in the

absence or presence of CIP at concentrations causing 20-30% growth inhibition and analyzed by

RNAseq (Materials and Methods). Samples were collected during exponential phase (A_{600} ~0.5)

as well as early post-exponential phase (A_{600} ~1), and differentially regulated genes were

identified by comparing the high-level FQR strains and the GP parent strain (Fig. 6C-F; GP vs

GPL (C, +CIP) and (D, -CIP), GP vs GPN (E), and GP vs GPS (F), Data Sets S3 and S4). In

either the presence or absence of CIP, the GPL strain had the greatest number of differentially

expressed genes relative to the control GP strain when compared to the other pump-

overexpressing strains (Fig. 6C-F). In fact, hyperexpression of either the AdeS- or AdeN-

regulated pumps only altered the gene expression patterns of very few genes in the presence or

absence of CIP (Fig. 6E, F). This result echoes the Tn-seq analysis of the three overproducer

strains, as the *adeL* mutant had the largest set of mutations that lowered fitness relative to WT

(Figs. 2, 3, and Figs. S17-19).

Among the various transcriptional changes resulting from AdeL-regulated pump hyper-

expression, the most notable was upregulation of the *paa* operon, which encodes proteins

involved in degradation of phenylacetate (Fig. 6C, D). In the absence of drug, 14/46 genes that

were differentially regulated in the two backgrounds were from the *paa* regulon (Fig. 6D).

Furthermore, in early post-exponential phase, genes rank-ordered as the 12 most highly

overexpressed in the GPL strain compared to the GP parent were also in the *paa* regulon

(Supplemental Dataset 4), so the functional importance of this upregulation in the absence of

drug was pursued further. As confirmation of the overexpression of *paa*, RNA was isolated from

parallel cultures of GP and GPL grown in the absence of drug and analyzed by q-rtPCR

(Materials and Methods). Consistent with the RNAseq results, transcript levels of the first gene

of the operon, *paaA*, were significantly elevated in the GPL strain compared to the GP parent,

with a 23-fold increase during exponential phase and a 6-fold increase in early post-exponential

phase (Fig. 7A, B). Another transcriptomic change of note in the GPL strain was the

downregulation of the *atpE* gene encoding a component of the F_1F_0 proton translocating

ATPase/synthase in the absence of drug (Fig. 6D). This parallels other work, which shows that

depletion of the ATPase/synthase increases fitness of the GPL strain in the presence of CIP (35).

Depletion of *paaA* alters expression of stress-related genes and lowers intracellular

pH in the GPL pump hyperexpressor. The phenylacetate catabolic pathway breaks down the

substrate phenylacetate to generate end products that feed into the TCA cycle (46). Previous

studies indicate that induction of the *paa* pathway in *A. baumannii* may be a response to stress

observed after antibiotic treatment (47). Therefore, upregulation of this operon may similarly

support overproduction of the AdeL-regulated pump. It should be emphasized that *paa* mutants

showed no significant fitness defect in our TnSeq screen. Consistent with the Tn-seq findings,

targeting *paaA* by CRISPRi caused no growth defect in the GPL strain as compared to a parallel

non-targeting control strain. Depletion was efficient, as there was a 20-fold knockdown in

exponential phase and almost 200-fold in early post-exponential phase compared to the negative

control (Fig. 7C, D). This argues that in the absence of phenylacetate degradation, other

compensatory pathways may allow fitness to be maintained.

The *paa* pathway has been shown to promote resistance to hydrogen peroxide (47, 48), so

enhanced expression of the *paa* pathway in the AdeFGH pump hyperexpresser may facilitate a

response to oxidative stress and/or other stressors. To test this idea, transcription of a set of

oxidative stress genes was probed in the GPL strain by q-rtPCR analysis. Knockdown of *paaA*

expression in early post exponential phase yielded reduced transcript levels of *katG, katE*, and

sodC (encoding two peroxidases and superoxide dismutase, respectively; Fig. 7D). This

diminished expression is consistent with previous observations that *paa* null mutants are more

sensitive to hydrogen peroxide stress through some previously uncharacterized mechanism (47,

48).

Internal pH was determined after *paaA* knockdown in the GPL mutant (Fig. 7E, F).

Knockdown of *paaA,* the first gene of the large operon, resulted in a significant acidification

during exponential growth compared to a non-targeting control (Fig. 7E) (A_{600} ~0.52-0.62). By

early post-exponential phase (A_{600}~1.22-1.39), the effect of the knockdown was no longer

statistically significant compared to the non-targeting control (Fig. 7F). When the *paa*

knockdown cultures were grown to exponential phase and exposed to sub-MIC levels of CIP for

the last hour, the pHi was lower than the non-targeting control (Fig. 7G; significance determined

via unpaired t tests). These data indicate a role for this pathway in maintaining pH homeostasis

during exponential growth of strains overexpressing the AdeL-regulated pump.

Given that the *paa* operon has been documented to be associated with a drug stress

response, we determined if the expression of the *paa* operon is altered in response to CIP in each

of the strains. Based on the RNAseq data in exponential (Fig. 8A) and early post-exponential

(Fig. 8B) phase, *paaA* transcript levels in all the strains were significantly upregulated in

response to CIP exposure as compared to growth in the absence of drug (Fig. 8A, B). These

results are consistent with previous results showing that upregulation of the *A. baumannii paa*

catabolon occurs in response to drug stress (47). Notably, upregulation of *paa* transcript levels in

the absence of drug is a special feature of the GPL strain when compared to the GP parent, and

this is not observed in the GPS or GPN strains (Fig. 6D).

To determine if this upregulation is in response to overexpression of the RND efflux

pump, rather than a consequence of a mutation in the regulatory element, the AdeL single mutant

and AdeL mutant with the AdeFGH pump deleted were grown to mid-exponential phase with

WT, GP, and GPL cultured in parallel. RNA was extracted and used for q-rtPCR to quantify

paaA transcript levels. As anticipated, *paaA* transcript levels were elevated in the AdeL mutant.

These levels dropped, however, in the absence of the AdeFGH RND efflux pump (Fig. 8C). This

result supports the hypothesis that the *paa* operon is involved in a stress response to efflux pump

hyperexpression.

Previous work has shown that pump hyper-expression and the oxidative stress response

are contributors to the development of persistence in the presence of supra-MIC levels of

antibiotics (49-52). In particular, the single AdeL mutant promotes persistence in the presence of

about 20X MIC levels of CIP (52). We, therefore, determined if depleting *paaA* expression

would interfere with persistence. Depleting *paaA* transcript did not alter the ability of a single

adeL mutation in the WT background to persist in high levels of CIP when compared to the non-

targeting control in AdeL (Fig. 8D). These results indicate that during exponential growth,

upregulation of phenylacetate metabolism plays a role in maintaining cellular homeostasis in

response to pump expression, mimicking the response of *A. baumannii* to antibiotic exposure.

Discussion

In this work, we set out to identify proteins that were required to maintain the fitness of

Acinetobacter baumannii drug efflux pump-activated strains. The most surprising result was that

in the AdeN and AdeS mutants, efflux pump overexpression was well-tolerated, with few

mutations causing defects specific to these strain backgrounds (Fig. 3) and little compensatory

transcriptional changes (Fig. 6). In contrast, the high-level fluoroquinolone resistant (FQR) strain

expressing the AdeL-regulated pump showed a range of mutations that lowered fitness relative to

the WT control and a more pronounced shift in its transcriptome relative to a *gyrAparC* FQR

mutant (GP). An obvious difference between these mutants is that the *adeL*-activating mutations

resulted in a higher level of expression of the AdeH outer membrane component than the

corresponding AdeS- and AdeN-regulated outer membrane components (Fig. 6). The AdeS-

regulated pump that was the most tolerant of insertion mutations has an outer membrane

component that is not subject to AdeS-mediated upregulation (31, 32).

We previously observed that AdeL mutations confer persistence during high-level CIP

exposure (52). In the mouse pneumonia model, *adeL*-activating mutants are outcompeted by

strains that have increased fitness in the absence of antibiotic (52), consistent with *adeL*-

activating mutations imparting a fitness cost. We suspect that the lowered growth rate is a

consequence of envelope stress (Fig. 3) and could have clinical corollaries. The inability of *adeL*

mutants to efficiently compete in immunocompetent animals (52) may mirror drug resistance

evolution in hospitals, where immunocompromised patients may act as reservoirs for these

variants that eventually act as evolutionary precursors to more fit and higher resistance variants.

In this fashion, evolutionary precursors that cause cell envelope disruptions may be compensated

by further mutations that allow spread in individuals with intact immune systems.

Fitness costs and resistance drivers are important for the evolution of MDR clinical

isolates and it is common for a pathogen to employ multiple drug efflux systems at a time, which

has the potential for accumulated envelope stress (19). Of note, while it appears there is no

advantage to overexpressing both AdeIJK and AdeFGH efflux systems simultaneously during

CIP challenge (Figs. 3H, S17, GPNL mutant in Fig. S20), this was not investigated further in the

presence of other drugs nor with AdeAB. This work does emphasize, however, that there is an

interplay between fitness costs and fractional differences in drug resistance, and that strains

having multiple antibiotic efflux systems may require compensatory mutations to reverse fitness

costs. The work described here identified points of vulnerability in the high-level FQR strains

that may be exploited to sensitize MDR strains to drug. It is evident that GPL heavily relies on

determinants that stabilize the cell envelope and an effective alarmone response in the presence

of drug. Furthermore, it may explain why *adeS* and *adeN*-regulated pump overproducers appear

to be particularly common in clinical isolates.

BfmS, identified in our Tn-seq screen, is the histidine kinase sensor that negatively

regulates the BfmR response regulator in the BfmRS two-component regulatory system (TCS).

BfmR controls a large transcriptional network including proteins linked to cell wall biogenesis,

capsule production, and outer membrane protein assembly (41). It is intriguing that disruption of

bfmS is detrimental to the fitness of the RND pump hyper-expressors, because disruption in a

drug-sensitive isolate has been shown to increase drug resistance and the expression of genes

such as *pbpG* that support RND pump expression (41). Presumably, proteins known to be

preferentially downregulated or mis-localized when the TCS is activated by loss of BfmS may,

in bulk, contribute to the fitness of the FQR strains. One such downregulated gene is *ompA*,

encoding a major outer membrane protein of *A. baumannii* (53, 54), which apparently functions

as a low-efficiency non-selective ion channel similar to *E.coli* OmpA and *Pseudomonas* OprF

(55, 56).

If there is a single feature linking together the Tn-seq mutant candidates, it is that they hit

proteins associated with maintenance of outer membrane integrity. Various studies have

demonstrated that defects causing instability to the outer membrane can be overcome by

compensatory changes in envelope components, particularly associated with the outer membrane

(57, 58). The functionality of the RND transporters is likely to be similarly supported by these

components. We postulate that PG, LPS, and other OMPs are central players in allowing

overproduction of the outer membrane component of the RND pump while simultaneously

supporting outer membrane barrier integrity. As it provides a selective permeability barrier

against antimicrobials, any destabilization of this barrier may simultaneously depress drug efflux

pump function or lead to a compromised drug barrier, leading to drug sensitivity. Most of the

insertions that caused fitness defects in pump-overexpressing strains were revealed only after the

addition of CIP (Fig. 3). This means that either of these two mechanisms could come into play,

with high pump concentrations combined with a partially dysfunctional outer membrane barrier

collaborating to drive drug sensitivity.

There were gene disruptions that yielded significantly higher fitness in all FQR strain

backgrounds when compared to the drug-susceptible WT strain, particularly after CIP exposure.

The most striking knockouts were in *apaH,* encoding a dinucleoside polyphosphate hydrolase.

ApaH degrades (p)ppGpp alarmones, which accumulate in response to stress as a consequence of

RelA synthetase activity (59-61). Consistent with this result, we have recently shown that

CRISPRi depletion of SpoT, which also degrades alarmones, increases fitness of GPL during

CIP treatment (35). Further supporting the model that (p)ppGpp accumulation increases fitness

during CIP treatment, transposon insertions in RelA, predicted to eliminate alarmone synthesis,

resulted in a fitness reduction in the GPL background (Fig. S17A, C). Therefore, it is likely that

accumulation of signals activating a stress response increase fitness specifically in the RND

hyper-expressing strains, particularly in GPL.

The necessity of alarmone synthesis for optimum fitness was consistent with our

observation that there were spikes in insertion mutation abundance at two prophage locations

post-drug exposure in all strain backgrounds (Figs. S17-19C). These peaks were more prominent

in the FQR strains, especially GPL, than in the WT strain and were responsible for many of the

overrepresented insertions in these backgrounds (Fig. 3). Our previous work indicates that the

abundance of these sequences is due to prophage induction after CIP exposure (17), and is

associated with an amplified SOS stress response. Therefore, RND overproduction, particularly

in the GPL strain, requires an effective stress response in the presence of CIP under conditions in

which all strains are growing at equivalent rates.

Analysis of the transcriptome of the RND overproducers provides further evidence for

the induction of cell stress as a consequence of AdeL-regulated pump overexpression (Fig. 6).

Most notably, transcription of the phenylacetate operon (*paa*) was induced in response to pump

overexpression in the absence of antibiotic treatment. This catabolon has been documented to

protect against antimicrobial treatment (47) and supports virulence in *A. baumannii*,

Burkholderia cenocepacia, and *Pseudomonas aeruginosa* (47, 62-64). Although the

consequences of loss of *paaA* expression appear mild, there may be redundant pathways that

come into play in the presence of drug pressure to support fitness of the AdeL mutant. The

absence of *paaA* led to lowered cytoplasmic pH (Fig. 7E, G), so the production of substrates

from the Paa pathway, which feed into the TCA cycle, could provide relief from proton influx

produced by the overproduced pump. Alternatively, accumulation of phenylacetic acid could

lead to lowered cytosolic pH upon knockdown of the *paa* operon. The importance of maintaining

pH homeostasis in the presence of pump overproduction is further emphasized by the result that

loss of the proton-pumping ubiquinol oxidase complex was predicted to cause lowered fitness of

the GPL strain.

In summary, this work indicates that targets involved in supporting the integrity of the *A.*

baumannii envelope are associated with maintaining fitness of strains that overexpress efflux

pumps. The most highly expressed pump, regulated by AdeL, was associated with the largest

number of genetic requirements for fitness, and was the most disruptive to the transcriptome. The

genetic loci identified can be viewed as drug targets for synergy with fluoroquinolones as well

(Fig. S21), defining pathways that may buffer the stress that accompanies increased pump

activity. Still unclear is how the phenylacetate degradation pathway is induced in response to the

stress identified here, and what physiological role, if any, it plays in supporting the fitness of the

strains analyzed in this study. Future work will attempt to expand these studies into analyzing the

role of essential proteins in supporting pump overproduction in hopes of identifying potential

targets for multidrug therapy targeting drug-resistant mutants.

Materials and Methods

Bacterial strains and growth conditions. Bacterial strains used in this work are listed in

Table S1. All *Acinetobacter baumannii* strains are derived from ATCC 17978UN. Cultures were

grown in lysogeny broth (LB) (10 g/L tryptone, 5 g/L yeast extract, 10 g/L NaCl) or on LB agar

plates (LB supplemented with 15g/L agarose) and incubated at 37°C with aeration, as well as

shaking or on a roller drum if grown in broth. Unless grown the same day, cultures were

typically diluted, incubated to recover after overnight growth, then diluted to A_{600}=0.1-0.001 into

either 5-10mL broth or into 96-well plates (Costar), and growth was monitored by measuring

A_{600} every hour (broth culture) or every 15min (Epoch 2 or SynergyMx, (Biotek) plate readers.)

Growth is then plotted as log-transformed absorbance values, with the mean ± SEM shown.

Superoptimal broth with catabolite repression (SOC) was used for recovery following all

electroporations of *E. coli* and *A. baumannii* during cloning or library preparations.

Electroporations of *A. baumannii* were performed using 0.1cm gap cuvettes (0.2cm for *E. coli*)

and the Gene Pulser (Bio-Rad) at the settings 200 Ω, 25 µF, and 1.8 kV (at 2.2kVd). As needed,

LB was supplemented with 10% Sucrose or antibiotics at the following concentrations:

ampicillin (Amp): 100µg/mL; carbenicillin (Carb): 50-1000µg/mL; gentamicin (Gent):

100µg/mL; chloramphenicol (Cm): 100µg/mL; kanamycin (Kan): 20µg/mL; or CIP: varying

concentrations (Sigma).

Molecular cloning and mutant construction. Primers (IDT) and plasmids described in

this study as well are listed in Table S1. The AdeS (R152S) mutant was isolated by drug

selection of a precursor strain by plating on LB with 10 µg/mL gentamicin. Single colonies were

purified on LB agar plates, and then *adeR* and *adeS* were amplified by colony PCR and

sequenced (Genewiz) to identify mutations. The selected *adeS* mutation (R152S) was introduced

into *A. baumannii gyrA(S81L)parC(S84L)* (also called GP (GVA37)) to create

gyrA(S81L)parC(S84L)adeS(R152S) in a fashion similar to that described previously (also called

GPS (EHA105)) (65). Briefly, the mutant allele was amplified with upstream and downstream

flanking sequences to generate a product of about 1500bp in length and introduced first into

pUC18 by restriction enzyme digestion and T4 DNA ligation. Once propagated in DH5α and

sequence confirmed (Genewiz), the mutant allele was introduced into the pJB4648 plasmid by

restriction enzyme digestion and T4 DNA ligation and propagated in DH5α λpir. The mutant

allele in pJB4648 was then introduced into *A. baumannii* by electroporation and homologous

recombination with two selection steps, the first selecting for transformants on LB+10 μg/mL

gentamicin on bacteriological plates and the second counterselection for Gent^S on LB+10%

Sucrose plates at either room temperature or at 30°C. The internal, in-frame gene deletion

mutants were previously created or constructed during this study using the same methodology

after three-way restriction enzyme digestions and ligations (40, 41, 65). The GPL (EHA23) strain

(*gyrA(S81L)parC(S84L)adeL(ΔI335A336)*) was constructed using the gentamicin

selection/sucrose counterselection. The *adeL(ΔI335A336)* allele was PCR amplified from gDNA

isolated from P9 Cy3.2 (52) with flanking regions, cloned into pUC18 then moved into pJB4648,

propagated in XL1 Blues and DH5α λpir *E. coli*, sequence confirmed, then introduced into the

GP background by electroporation and homologous recombination. GP was created previously

(17) and *gyrA(S81L)parC(S84L)adeN*::ISAba1 (GVA47), also known as GPN, was generated by

plating GP on LB+32μg/mL CIP.

WGS. Strains used in this study were sequence verified by whole genome sequencing

(WGS) as previously described (52). Illumina Nextera reagents were used for library preparation

and 100bp single-end reads were obtained using the Tufts University Core Facility's HiSeq2500.

BRESEQ 0.35.5 was used to align reads to the *Acinetobacter baumannii* 17978UN reference

genome (GenBank Accession: CP079212.1), and mutations, those of which are absent in the

parent strains, were identified. All sequences are deposited in the SRA database under BioProject

number PRJNA1068501.

LOS analysis. LOS profiles were examined using Tricine gel electrophoresis of cell

lysates as previously described (40), using lysates from cells grown to mid-logarithmic phase.

After fractionation on Novex 16% tricine gels (Invitrogen), the gels were stained with Pro-Q

Emerald 300 (Invitrogen) for LOS staining, and Coomassie Brilliant Blue for total protein

staining for sample quantification normalization (40).

pHi assay. Buffers of varying pH were made using 50mL combinations of 0.2M

Na_2HPO_4 and 0.2M NaH_2PO_4 (66). 5mM BCECF/AM (ThermoFisher) and 0.5mM carbonyl

cyanide 3-chlorophenylhydrazone (CCCP; Sigma) were prepared in DMSO and stored at -20^{o}C

until use. Methods were adapted with modifications from previous protocols (67). Overnight

cultures prepared from individual colonies were diluted for recovery to allow growth, then

diluted again to A_{600} ~0.003 in 8mL LB broth (with or without drug as indicated). In the case of

dCas9 strains, single colonies were inoculated into 2mL LB in capped tubes and incubated to late

exponential phase before diluting into LB + 100ng/mL anhydrotetracycline (aTc). The cultures

that were exposed to sub-MIC levels of CIP were treated with 45 μg/mL of CIP an hour prior to

harvest. Cultures were grown to exponential phase (A_{600}=0.4-0.7) or early post exponential phase

(A_{600}=1.2-1.4) after which samples (about 5mL) were pelleted in 15mL conical tubes at 7500g

for 5 min, resuspended in sterile phosphate-buffered saline (PBS, Thermo Scientific), loaded

with 20μM BCECF/AM in PBS followed by incubation at 30^{o}C for 30 min in a 96-well v-

bottom plate protected from light.

Cells in the 96 well plate were diluted 1:4, pelleted at 2000rpm for 5 min, resuspended in

LB broth and incubated at 37°C for 5 min, before pelleting once more and re-suspending in 200-

400µl previously prepared mono- and di-basic sodium phosphate buffer at neutral pH

(experimental samples) or at a range of pH values (*in situ* standard curve samples). Samples were

diluted 1:1 in the neutral or pH adjusted buffers to a final volume of 200µl in a 96-well black

plate with a clear, flat bottom (Costar). About 10µM CCCP stock was added to the *in situ*

standard curve wells. All samples were mixed via pipetting before fluorescence was read at

490ex/535em and 440ex/535em after 5 minutes (Biotek SynergyMx with Gen5 software).

Background fluorescence from buffer alone was subtracted from each sample before calculating

the ratiometric fluorescence of the calibration curve samples and experimental samples. The

experimental sample internal pH was interpolated from the *in-situ* calibration curve using

standard curve fitting.

Preparation of transposon mutant banks. Transposon banks with *Himar-1* transposon

were created as described (17, 40). 50µl electrocompetent *A. baumannii* cells were

electroporated with 300ng pDL1100 (*Himar-1*-Kan^R). Cells were recovered in 1mL SOC broth,

placed in 14mL snap cap tubes (Corning) and aerated on a drum roller for 15min at 37°C. The

cells were pelleted and resuspended in 400µl SOC, then spread onto two 150 mm culture dishes

containing mixed cellulose ester filters (0.45µm pore size, 137mm diameter; Advantec) that had

been placed on top of the SOC agar. Plates were incubated for 1hr at 37°C before filters were

transferred to 150 mm culture dishes containing LB+20µg/mL kanamycin and incubated

overnight (~16hrs) at 37°C.

Mini-*Tn10* banks were also made in the GPN strain background. GPN electrocompetent

cells were transformed with 75ng pDL1073 (mTn*10-neo*), with subsequent steps following the

same procedure as used for *Himar-1*. CFU counts were determined from plate images using an

Image J counting routine. Mutants were pooled from filters by gentle shaking in 50-75mL sterile

PBS. Glycerol was added for a final concentration of around 10% and aliquots stored at -80oC

until use. About 21 independent subpools were created in each strain background (producing

around 100,000-150,000 cumulative unique mutants).

CIP challenge of transposon banks. Independent subpools comprising of approximately

7,500-16,000 unique transposon mutants were challenged with antibiotic as described (17, 40).

Mutant banks were thawed on ice, diluted in 25mL LB broth to an $A_{600}=0.1$, grown to $A_{600}=0.2$

at 37°C, then diluted to $A_{600}=0.003$ in 10mL LB broth with or without CIP of varied

concentrations added in parallel cultures. The remaining culture (T_1) was harvested to determine

what mutants were present prior to outgrowth and drug challenge. Cultures were incubated for

about 8 generations (around 170min if untreated, 225min if CIP treated), then harvested (T_2).

Drug-treated cultures that exhibited 20-30% growth inhibition were harvested. Cultures were

pelleted and stored at -20°C until all samples were collected. The concentration ranges at which

comparable growth inhibition was met for each strain background as well as the average

population expansions between T_1 and T_2 with each condition (according to A_{600}) are noted in

Data Set S2.

Preparation of transposon libraries for sequencing. As described (17, 40), the gDNA

was extracted using the DNeasy Blood and Tissue Kit (Qiagen). Samples were treated with

proteinase K during resuspension and incubated at 56°C with shaking for 1hr before treatment

with RNase cocktail (Fisher Scientific) for 8min at 62°C. The gDNA was quantified with SYBR

green then diluted to 7ng/µl in UltraPure water (Fisher Scientific). About 28ng of gDNA of each

sample was tagmented using TD buffer and TDE1 enzyme (Illumina) and incubated in a

thermocycler at 55°C for 5min prior to melting at 95°C for 0.5min. Transposon insertion sites

were then amplified using Q5 hot start polymerase (NEB) and 0.6µM Nextera 2A-R and olj638

(for *Himar-1* libraries), or olj928 (for Tn*10* libraries). The 50µl reactions were incubated at 98°C

for 1min, cycled 30 times through 98°C for 10sec, 65°C for 20 sec, and 72°C for 1min, then

finally at 72°C for 2min. 0.5µl of the first PCR products were used for a second PCR reaction

used to introduce indexes with leftward *Himar-1-* or Tn10-specific indexed primers and

rightward Nextera indexed primers. The reaction and settings were the same except cycling was

shortened to 12 cycles. Amplified products were quantified via 2% agarose tris-acetate-EDTA

(TAE) gels using SYBR Safe (Invitrogen) for visualization. The gels were loaded with a known

amount of 100bp DNA ladder (NEB) alongside samples of each reaction to quantify their

concentrations based on band intensity around 250-600bp using a gel imaging system. Equal

measures of all samples were thereafter pooled and purified via PCR column clean-up (Qiagen).

About 17.5 pmol of each pooled sample was reconditioned in an additional 50µl reaction with

primers P1 and P2. Thermocycler settings were set with a heated lid and incubations at 95°C for

1min, with slow ramp of 0.1°C/sec to 64°C then held for 20sec, and finally 72°C for 10min. The

products were purified once more with the PCR column clean-up kit (Qiagen). Primer olj115

(Tn*10*) or mar512 (*Himar-1*) was used for sequencing at the Tufts University Core Facility of

Genomics with the Illumina HiSeq 2500 (SE50, post PippinPrep (Sage Science) size selection

for 250-600bp).

Sequence analysis and assessment of mutant fitness. Sequenced reads were processed

and analyzed as done previously (17, 39, 40, 68). After demultiplexing, the Nextera adapter

sequence was clipped from reads using the fastx_clipper, quality filtered, and reads shorter than

20nt were discarded. Reads were mapped to the *A. baumannii* ATCC 17978 reference genome

(https://genomes.atcc.org/genomes/e1d18ea4273549a0) using Bowtie 1.2.2, formatted so only

unique and best-mapped reads were aligned, allowing up to 1 maximum mismatch. Fitness

scores were then calculated for each transposon mutant using MAGenTA (68), with scores

averaged and aggregated at the gene level across replicates. Insertions within 10% of 3' end of

genes were excluded and fitness scores normalized to a set of neutral/pseudo genes. Of note,

fitness scores of GPN insertion mutants were determined using data from both *Himar-1* and

mini-*Tn10* banks. All other results were from *Himar-1* mutagenesis. Fitness scores of gene

disruptions in all strain backgrounds were normalized to a LOWESS curve to avoid the position

bias exhibited post CIP exposure. The difference in fitness scoring for each gene disruption was

assessed between each FQR strain and the WT drug susceptible strain. Significant fitness

differences yielded from the same gene disruption between the different strain backgrounds were

identified via multiple t test analysis using the original FDR method of Benjamini and Hochberg

with significance falling within the FDR of 5% (q<0.05).

RNAseq experiment to uncover transcriptional differences between FQR and

parent strains. Cultures were grown overnight, diluted to an A_{600} of 0.1 in 10mL LB broth,

grown for about 1.5hrs at 37°C on a drum wheel to approximately mid-log growth, then diluted

once more to A_{600} of 0.003 in fresh LB broth with or without CIP. Some of these were grown to

approximately mid-log phase by incubation for 150-165min (untreated) or for 195-225min (CIP

treated) as similarly done before (17) and reaching A_{600}~0.445-0.68 and A_{600}~0.39-0.66

respectively. Others were grown for 195-210min (untreated) reaching A_{600}~0.85-1.195 or for

255-270min (CIP treated) to A_{600}~0.82-1.125. Of the cultures treated with CIP, those that

exhibited 20-30% growth inhibition (compared to the untreated culture grown in parallel) were

harvested (details provided in Data Sets S3 and S4). About 1-1.5 A_{600} units of all cultures were

harvested and frozen at -80°C until all samples were collected.

Preparing RNA for sequencing. RNA was isolated from the samples using the RNeasy

Mini kit (Qiagen) and prepared for sequencing as described (17, 69, 70) following published

protocols (71). Approximately 600ng of RNA was fragmented at 94°C with 10x FastAP buffer

for 2.5min then treated with TURBO DNase, FastAP (Thermo Scientific) (and murine RNase

inhibitor; NEB) for 30min at 37°C. Samples were then purified using Agencourt RNAClean XP

(Beckman Coulter) and eluted with 12.5µl of DEPC UltraPure water (Fisher Scientific). 5µl of

sample was incubated with 1µl of 100µM barcoded RNA adapters at 72°C for 2 min before

mixed with DMSO (American Bioanalytical), 10x T4 RNA ligase buffer, 100mM ATP, RNase

inhibitor murine, T4 RNA ligase, and 50% PEG8000 (NEB) to allow for 3' linker ligation at

room temperature for 1.5 hours. The 60 samples were pooled into 5 groups of 12 samples each

during purification using RNA Clean & Concentrator (Zymo Research) and all eluted with 16µl

DEPC UltraPure water.

Ribosomal rRNA was depleted using RNase-H as described (72). Probes complementary

to *A. baumannii* ATCC 17978UN 23S, 16S, and 5S rRNA sequences (CP079931.1) were created

using 0.design_probe.py and off-target binding predicted using 2.predict_probe_offtarget.py

(https://github.com/hym0405/RNaseH_depletion). The 89 devised oligonucleotides (IDT) were

resuspended into 100µM in UltraPure water, and a master-mix created with 5µl of each probe.

For each depletion, the RNA was mixed with 1µl of the probe mixture, 0.6µl 1M Tris-HCl pH

7.5, 1.5µl 5M NaCl, and DEPC UltraPure water for a final 15µl reaction. They were incubated in

a thermocycler with a heated lid at 95°C for 2min, temperature reduced to 45°C in 0.1°C/sec

increments and incubated a further 5 min at 45°C. Meanwhile, RNase-H master mix was

prepared for each depletion (3µl Hybridase Thermostable RNase H; Epicentre), 0.5µl 1M Tris-

HCl pH=7.5, 0.2µl 5M NaCl, 0.4µl 1M MgCl2, and 0.9µl DEPC UltraPure water) and preheated

to 45^{o}C. Post probe and RNA hybridization, 5µl prewarmed RNase-H mixture was added to each

probe-RNA mixture and incubated at 45^{o}C for 30min. These were then purified twice with 2x

SPRI RNAClean XP (Agencourt) and eluted with 13µl DEPC UltraPure water.

The RNA preparation was then used as a template for cDNA synthesis. 12µl of sample

was incubated with 2µl of 25µM AR2 primer at 72^{o}C for 2min using 10x AffinityScript RT

buffer and enzyme (Agilent Technologies), 0.1M DTT, 100mM dNTPs, and 40U/µl murine

RNase inhibitor. After 55min at 55^{o}C, RNA was then denatured in a 10% reaction volume

(adding 2µl to the 20µl cDNA reaction product) of 1N freshly prepared NaOH at 70^{o}C for 12min

before neutralization with 4µl of freshly prepared 0.5M acetic acid. The products were then

subjected to Agencourt RNAClean XP, eluted with 5µl DEPC UltraPure water, incubated with

2µl of 40µM 3Tr3 adaptor at 75^{o}C for 3min with the beads still present, and mixed with 10x T4

ligase buffer, DMSO, ATP, PEG8000, and T4 RNA ligase for an overnight 20µl ligation

reaction at room temperature. The products were purified with SPRI RNAClean XP (Agencourt)

beads twice and finally eluted with 25µl H$_2$0. DNA was amplified from cDNA using 10µl of 5x

Q5 reaction buffer and 2.5µl Q5 enzyme (NEB), 0.5µl 100mM dNTPs, 10µl cDNA, 2.5µl

primers RTS_Enr_P5 and RTS_Enr_P7_BCXX, and water for 50µl reaction. The reactions were

incubated at 98^{o}C for 3min, cycled at 98^{o}C for 30 sec, 67^{o}C for 25sec, and 72^{o}C for 15sec for 14

cycles, and completed at 72^{o}C for 2min. PCR products were purified once with AxyPrep

Magnetic PCR beads (Axygen; Corning). DNA was then quantified by Qubit with the dsDNA

HS Assay kit (RNA HS Assay kit was used previously for the quantification of the RNA

samples; Thermo Scientific) and about 100nM of each of the 5 subpools was pooled together

into 1 sample for sequencing.

Throughout sample preparation, 1µl SUPERase-IN RNase inhibitor (Thermo Scientific)

was added to samples if they were stored at -80°C in between steps. RNAse Zap was used

continuously on surfaces until cDNA was synthesized. Samples were routinely analyzed by

submitting samples for QC fragment analysis at the TUCF Genomics Core. The pooled samples

were sequenced at this facility, using a NextSeq550 Single-End High output, with 75 cycles

following nanoQC.

RNAseq analysis. Sequences that had been de-multiplexed for the i7 barcodes were then

de-multiplexed for the i5 barcodes using fastx_barcode_splitter.pl

(https://github.com/agordon/fastx_toolkit/). The 5' anchored adaptor sequences were then

trimmed *in silico* using Cutadapt (73). Truseq adaptor sequences were also trimmed *in silico*,

disallowing any error, and dropping resulting sequences shorter than 20nt. Sequence data were

thereafter analyzed following and modifying published scripts

(https://github.com/huoww07/Bioinformatics-for-RNA-Seq). The

Acinetobacter_baumannii_ATCC_17978.fasta and Acinetobacter_baumannii_ATCC_17978.gbk

files were downloaded from the ATCC genome portal

(https://genomes.atcc.org/genomes/e1d18ea4273549a0?tab=overview-tab) to create the STAR

genome index (via genomeGenerate with STAR). Annotation GTF files were constructed using

Geneious, then cufflinks gffread for conversion to a strict gtf file. Alignments were then

performed using STAR. Subread/1.6.3 was used to get a summary of the sequence data features,

then featurecount_stat.R as well as mapping_percentage.R were adapted to analyze and visualize

features including mapping efficiency and overview of mapped vs. unmapped reads. Using the

Tufts Cluster on-demand R studio app, DESeq2, ggplot2, dplyr, tidyverse, ggrepel, DEGreport,

and pheatmap tools were employed to analyze the quality of the dataset as well as visualize and

deconvolve the contrasting transcriptomic data between the high-level FQR pump-overproducing

strains and the GP parental strain.

771   **qPCR analysis for transcript abundance.** Cultures derived from individual colonies

were grown in LB broth to mid- or early post-exponential phase. About 1 A_{600} unit of culture

was harvested and RNA was extracted using RNeasy (Qiagen). cDNA was synthesized after

ezDNase treatment using the SuperScript VILO cDNA kit (Thermo Fisher), then amplified using

PowerUp SYBR Green Master Mix (Applied Biosystems) and StepOnePlus Real-Time PCR

system (Applied Biosystems) per manufacturer's instructions. The efficiency of target

amplification was tested using a standard curve of cDNA at varying dilutions and consequently

verified to be greater than 95% for each primer pair used. Three biological replicates were

evaluated for each strain and three technical replicates examined per biological sample. To

confirm a lack of gDNA contamination, control samples tested in parallel which lacked reverse-

transcriptase during cDNA synthesis were included. Transcript levels for each indicated target

were compared between each mutant strain and the parental strain (via the comparative $2^{-\Delta\Delta CT}$

method) after normalization to that of 16S, the endogenous control. The data are represented in

bar plots showing the mean fold difference in transcript levels ($2^{-\Delta\Delta CT}$) ± SEM.

785   **MIC determination.** MICs were determined by broth microdilutions. Overnight cultures

were diluted to A_{600}=0.1 in 3mL LB broth, incubated to early exponential phase (A_{600}=0.2-0.3),

then diluted to a final A_{600}=0.003 in 200ul LB broth in the presence of different dilutions of CIP

in a 96-well plate (Costar). Growth was monitored by measuring A_{600} every 15min (Epoch 2 or

SynergyMx, (Biotek) plate readers). At least three replicates were tested for all strains. CIP MIC

was determined as the lowest concentration that prevented growth up to 12 hours, where

geometric mean A_{600} units did not exceed 0.035, and are not significantly different from the LB

broth-only negative controls. CIP MICs were 256µg/mL (GPS and GPN) and 384µg/mL (GPL).

Persistence assay. The *A. baumannii* strains EHA417 (*adeL*ΔI335A336 *P_Tet_::dCas9*

(pWH1266-*paaA*-sgRNA)), EHA418 (*adeL*ΔI335A336 *P_Tet_::dCas9* (pWH1266-NC-sgRNA)),

and EHA254 (WT *P_Tet_::dCas9* (pWH1266-NC-sgRNA)) were incubated in 1.5mL LB in

triplicate for 2-3 hours. They were then diluted 1/1000 in 8mL LB + 100ng/mL

anhydrotetracycline (aTc) to induce dCas9 expression and grown to mid-exponential phase. Each

culture was then normalized to A_{600}~0.5 with fresh LB containing inducer aTc, then treated with

10 µg/mL CIP. Samples of culture (500µl) were removed at timepoints=0, 1, 6, and 24 hours,

washed with PBS, then resuspended in PBS and serial dilutions were spotted on LB agar plates

for overnight incubation at 37°C and CFU enumeration the next day.

Microscopy. Culture samples were placed on 1% agarose in PBS pads and imaged using

DIC at 100X magnification on the Zeiss Axiovert Observer.Z1.

Accession numbers

All analyzed sequences are in the SRA database under BioProject numbers PRJNA1067619

(TnSeq), PRJNA1067535 (RNAseq), and PRJNA1068501 (WGS).

Acknowledgements

This work is supported by NIAID awards U01AI124302 and U19AI158076 (RRI, TvO), U19AI142780 and R21AI128328 (RRI), and R01AI162996 (EG). We thank David Lazinski (Tn plasmids), Germán Vargas-Cuebas (for the GPN mutant), Rebecca Batorsky (bioinformatic help with TRANSIT), Yunfei Dai (for communications regarding dCas9 strain generation, sgRNA design, and TRANSIT, and for the pUC-Tn7-GentR-pTet::dCas9 and pWH1266-sgRNA plasmids), Matthew Hall (for help with RNAseq library preparation protocols), Albert Tai and Irina V. Grinvald for communications regarding sequencing and library preparation, as well as the next-generation sequencing support provided by S10OD032203 via Tufts University Core Facility Genomics Core.

Author Contributions

R.R.I and E.G. designed the TnSeq study and provided training and support with experimental methodologies. S.S. and A.C. created mariner banks in the WT and GPN strains. A.C. performed TnSeq experiments with GPN and WT. J.B. performed the LOS experiments and analyses. W.H. provided bioinformatic support and ran the DESeq2 bioinformatic workflow for the comparative RNAseq study. J.H-B. performed the WGS library prep and sequence analysis for all deletion mutants. J.C.O-M. and TvO provided support with RNAseq library preparation methodologies. E.G. created the vectors used to make two deletion mutants and the backbone vector used for gene complementation. E.H. and R.R.I devised the research, interpreted the results, and wrote the manuscript. E.G reviewed and revised the manuscript.

References

1. Peleg AY, Seifert H, Paterson DL. 2008. *Acinetobacter baumannii*: emergence of a

   successful pathogen. Clin Microbiol Rev 21:538-82.

2. CDC. 2019. Antibiotic Resistance Threats in the United States. Atlanta, GA: US

   Department of Health and Human Services, CDC.

3. CDC. 2022. COVID-19: U.S. Impact on Antimicrobial Resistance, Special Report 2022.

   Atlanta, GA: US Department of Health and Human Services, CDC.

4. Tomaras AP, Dorsey CW, Edelmann RE, Actis LA. 2003. Attachment to and biofilm

   formation on abiotic surfaces by *Acinetobacter baumannii*: involvement of a novel

   chaperone-usher pili assembly system. Microbiology (Reading) 149:3473-3484.

5. Badave GK, Kulkarni D. 2015. Biofilm Producing Multidrug Resistant *Acinetobacter*

   baumannii: An Emerging Challenge. J Clin Diagn Res 9:DC08-10.

6. Sanchez CJ, Jr., Mende K, Beckius ML, Akers KS, Romano DR, Wenke JC, Murray CK.

   2013. Biofilm formation by clinical isolates and the implications in chronic infections.

   BMC Infect Dis 13:47.

7. Gonzalez-Villoria AM, Valverde-Garduno V. 2016. Antibiotic-Resistant *Acinetobacter*

   baumannii Increasing Success Remains a Challenge as a Nosocomial Pathogen. J Pathog

   2016:7318075.

8. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, Pulcini C,

   Kahlmeter G, Kluytmans J, Carmeli Y, Ouellette M, Outterson K, Patel J, Cavaleri M,

   Cox EM, Houchens CR, Grayson ML, Hansen P, Singh N, Theuretzbacher U, Magrini N,

   Group WHOPPLW. 2018. Discovery, research, and development of new antibiotics: the

856 WHO priority list of antibiotic-resistant bacteria and tuberculosis. Lancet Infect Dis

857 18:318-327.

858 9. Lob SH, Hoban DJ, Sahm DF, Badal RE. 2016. Regional differences and trends in

859 antimicrobial susceptibility of *Acinetobacter baumannii*. Int J Antimicrob Agents 47:317-

860 23.

861 10. Giammanco A, Cala C, Fasciana T, Dowzicky MJ. 2017. Global Assessment of the

862 Activity of Tigecycline against Multidrug-Resistant Gram-Negative Pathogens between

863 2004 and 2014 as Part of the Tigecycline Evaluation and Surveillance Trial. mSphere 2.

864 11. Poole K. 2004. Efflux-mediated multiresistance in Gram-negative bacteria. Clin

865 Microbiol Infect 10:12-26.

866 12. Sun J, Deng Z, Yan A. 2014. Bacterial multidrug efflux pumps: mechanisms, physiology

867 and pharmacological exploitations. Biochem Biophys Res Commun 453:254-67.

868 13. Coyne S, Courvalin P, Perichon B. 2011. Efflux-mediated antibiotic resistance in

869 *Acinetobacter* spp. Antimicrob Agents Chemother 55:947-53.

870 14. Blair JM, Webber MA, Baylay AJ, Ogbolu DO, Piddock LJ. 2015. Molecular

871 mechanisms of antibiotic resistance. Nat Rev Microbiol 13:42-51.

872 15. Geisinger E, Huo W, Hernandez-Bird J, Isberg RR. 2019. *Acinetobacter baumannii*:

873 Envelope Determinants That Control Drug Resistance, Virulence, and Surface

874 Variability. Annu Rev Microbiol 73:481-506.

875 16. Compagne N, Vieira Da Cruz A, Muller RT, Hartkoorn RC, Flipo M, Pos KM. 2023.

876 Update on the Discovery of Efflux Pump Inhibitors against Critical Priority Gram-

877 Negative Bacteria. Antibiotics (Basel) 12.

878   17.   Geisinger E, Vargas-Cuebas G, Mortman NJ, Syal S, Dai Y, Wainwright EL, Lazinski D,

879   Wood S, Zhu Z, Anthony J, van Opijnen T, Isberg RR. 2019. The Landscape of

880   Phenotypic and Transcriptional Responses to Ciprofloxacin in *Acinetobacter baumannii*:

881   Acquired Resistance Alleles Modulate Drug-Induced SOS Response and Prophage

882   Replication. mBio 10.

883   18.   Naidu V, Bartczak A, Brzoska AJ, Lewis P, Eijkelkamp BA, Paulsen IT, Elbourne LDH,

884   Hassan KA. 2023. Evolution of RND efflux pumps in the development of a successful

885   pathogen. Drug Resist Updat 66:100911.

886   19.   Pagdepanichkit S, Tribuddharat C, Chuanchuen R. 2016. Distribution and expression of

887   the Ade multidrug efflux systems in *Acinetobacter baumannii* clinical isolates. Can J

888   Microbiol 62:794-801.

889   20.   Magnet S, Courvalin P, Lambert T. 2001. Resistance-nodulation-cell division-type efflux

890   pump involved in aminoglycoside resistance in *Acinetobacter baumannii* strain BM4454.

891   Antimicrob Agents Chemother 45:3375-80.

892   21.   Coyne S, Rosenfeld N, Lambert T, Courvalin P, Perichon B. 2010. Overexpression of

893   resistance-nodulation-cell division pump AdeFGH confers multidrug resistance in

894   *Acinetobacter baumannii*. Antimicrob Agents Chemother 54:4389-93.

895   22.   Yoon EJ, Chabane YN, Goussard S, Snesrud E, Courvalin P, De E, Grillot-Courvalin C.

896   2015. Contribution of resistance-nodulation-cell division efflux systems to antibiotic

897   resistance and biofilm formation in *Acinetobacter baumannii*. mBio 6.

898   23.   Rosenfeld N, Bouchier C, Courvalin P, Perichon B. 2012. Expression of the resistance-

899   nodulation-cell division pump AdeIJK in *Acinetobacter baumannii* is regulated by AdeN,

900   a TetR-type regulator. Antimicrob Agents Chemother 56:2504-10.

24. Wang Z, Fan G, Hryc CF, Blaza JN, Serysheva, II, Schmid MF, Chiu W, Luisi BF, Du D.

2017. An allosteric transport mechanism for the AcrAB-TolC multidrug efflux pump.

Elife 6.

25. Murakami S, Nakashima R, Yamashita E, Yamaguchi A. 2002. Crystal structure of

bacterial multidrug efflux transporter AcrB. Nature 419:587-93.

26. Murakami S, Nakashima R, Yamashita E, Matsumoto T, Yamaguchi A. 2006. Crystal

structures of a multidrug transporter reveal a functionally rotating mechanism. Nature

443:173-9.

27. Koronakis V, Sharff A, Koronakis E, Luisi B, Hughes C. 2000. Crystal structure of the

bacterial membrane protein TolC central to multidrug efflux and protein export. Nature

405:914-9.

28. Yu EW, Aires JR, McDermott G, Nikaido H. 2005. A periplasmic drug-binding site of

the AcrB multidrug efflux pump: a crystallographic and site-directed mutagenesis study.

J Bacteriol 187:6804-15.

29. Yu EW, McDermott G, Zgurskaya HI, Nikaido H, Koshland DE, Jr. 2003. Structural

basis of multiple drug-binding capacity of the AcrB multidrug efflux pump. Science

300:976-80.

30. Higgins MK, Bokma E, Koronakis E, Hughes C, Koronakis V. 2004. Structure of the

periplasmic component of a bacterial drug efflux pump. Proc Natl Acad Sci U S A

101:9994-9.

31. Sugawara E, Nikaido H. 2014. Properties of AdeABC and AdeIJK efflux systems of

Acinetobacter baumannii compared with those of the AcrAB-TolC system of *Escherichia*

coli. Antimicrob Agents Chemother 58:7250-7.

32.   Leus IV, Weeks JW, Bonifay V, Smith L, Richardson S, Zgurskaya HI. 2018. Substrate

925   Specificities and Efflux Efficiencies of RND Efflux Pumps of *Acinetobacter baumannii*.

926   J Bacteriol 200.

927   33.   DeJesus MA, Ambadipudi C, Baker R, Sassetti C, Ioerger TR. 2015. TRANSIT--A

928   Software Tool for Himar1 TnSeq Analysis. PLoS Comput Biol 11:e1004401.

929   34.   Bai J, Dai Y, Farinha A, Tang AY, Syal S, Vargas-Cuebas G, van Opijnen T, Isberg RR,

930   Geisinger E. 2021. Essential Gene Analysis in *Acinetobacter baumannii* by High-Density

931   Transposon Mutagenesis and CRISPR Interference. J Bacteriol 203:e0056520.

932   35.   Hamami E, Huo W, Neal K, Neisewander I, Geisinger E, Isberg RR. 2024. Identification

933   of essential genes that support fitness of *Acinetobacter baumannii* efflux pump

934   overproducers in the presence of fluoroquinolone. bioRxiv

935   doi:10.1101/2024.01.04.574119:2024.01.04.574119.

936   36.   VanOtterloo LM, Macias LA, Powers MJ, Brodbelt JS, Trent MS. 2024. Characterization

937   of *Acinetobacter baumannii* core oligosaccharide synthesis reveals novel aspects of

938   lipooligosaccharide assembly. mBio 15:e0301323.

939   37.   Arai H, Kawakami T, Osamura T, Hirai T, Sakai Y, Ishii M. 2014. Enzymatic

940   characterization and in vivo function of five terminal oxidases in *Pseudomonas*

941   *aeruginosa*. J Bacteriol 196:4206-15.

942   38.   Borisov VB, Verkhovsky MI. 2015. Oxygen as Acceptor. EcoSal Plus 6.

943   39.   van Opijnen T, Bodi KL, Camilli A. 2009. Tn-seq: high-throughput parallel sequencing

944   for fitness and genetic interaction studies in microorganisms. Nat Methods 6:767-72.

945   40.   Geisinger E, Mortman NJ, Dai Y, Cokol M, Syal S, Farinha A, Fisher DG, Tang AY,

946   Lazinski DW, Wood S, Anthony J, van Opijnen T, Isberg RR. 2020. Antibiotic

susceptibility signatures identify potential antimicrobial targets in the *Acinetobacter*

baumannii cell envelope. Nat Commun 11:4522.

41.   Geisinger E, Mortman NJ, Vargas-Cuebas G, Tai AK, Isberg RR. 2018. A global

regulatory system links virulence and antibiotic resistance to envelope homeostasis in

Acinetobacter baumannii. PLoS Pathog 14:e1007030.

42.   Russo TA, MacDonald U, Beanan JM, Olson R, MacDonald IJ, Sauberan SL, Luke NR,

Schultz LW, Umland TC. 2009. Penicillin-binding protein 7/8 contributes to the survival

of *Acinetobacter baumannii* in vitro and in vivo. J Infect Dis 199:513-21.

43.   Bulssico J, Papukashvil II, Espinosa L, Gandon S, Ansaldi M. 2023. Phage-antibiotic

synergy: Cell filamentation is a key driver of successful phage predation. PLoS Pathog

19:e1011602.

44.   Bos J, Zhang Q, Vyawahare S, Rogers E, Rosenberg SM, Austin RH. 2015. Emergence

of antibiotic resistance from multinucleated bacterial filaments. Proc Natl Acad Sci U S

A 112:178-83.

45.   Nikaido H, Takatsuka Y. 2009. Mechanisms of RND multidrug efflux pumps. Biochim

Biophys Acta 1794:769-81.

46.   Teufel R, Mascaraque V, Ismail W, Voss M, Perera J, Eisenreich W, Haehnel W, Fuchs

G. 2010. Bacterial phenylalanine and phenylacetate catabolic pathway revealed. Proc

Natl Acad Sci U S A 107:14390-5.

47.   Hooppaw AJ, McGuffey JC, Di Venanzio G, Ortiz-Marquez JC, Weber BS, Lightly TJ,

van Opijnen T, Scott NE, Cardona ST, Feldman MF. 2022. The Phenylacetic Acid

Catabolic Pathway Regulates Antibiotic and Oxidative Stress Responses in

Acinetobacter. mBio 13:e0186321.

48.   Green ER, Juttukonda LJ, Skaar EP. 2020. The Manganese-Responsive Transcriptional

Regulator MumR Protects *Acinetobacter baumannii* from Oxidative Stress. Infect Immun

88.

49.   Vega NM, Allison KR, Khalil AS, Collins JJ. 2012. Signaling-mediated bacterial

persister formation. Nat Chem Biol 8:431-3.

50.   Wu Y, Vulic M, Keren I, Lewis K. 2012. Role of oxidative stress in persister tolerance.

Antimicrob Agents Chemother 56:4922-6.

51.   Lee HH, Molla MN, Cantor CR, Collins JJ. 2010. Bacterial charity work leads to

population-wide resistance. Nature 467:82-5.

52.   Huo W, Busch LM, Hernandez-Bird J, Hamami E, Marshall CW, Geisinger E, Cooper

VS, van Opijnen T, Rosch JW, Isberg RR. 2022. Immunosuppression broadens

evolutionary pathways to drug resistance and treatment failure during *Acinetobacter*

baumannii pneumonia in mice. Nat Microbiol 7:796-809.

53.   Park JS, Lee WC, Yeo KJ, Ryu KS, Kumarasiri M, Hesek D, Lee M, Mobashery S, Song

JH, Kim SI, Lee JC, Cheong C, Jeon YH, Kim HY. 2012. Mechanism of anchoring of

OmpA protein to the cell wall peptidoglycan of the gram-negative bacterial outer

membrane. FASEB J 26:219-28.

54.   Gribun A, Nitzan Y, Pechatnikov I, Hershkovits G, Katcoff DJ. 2003. Molecular and

structural characterization of the HMP-AB gene encoding a pore-forming protein from a

clinical isolate of *Acinetobacter baumannii*. Curr Microbiol 47:434-43.

55.   Sugawara E, Nikaido H. 2012. OmpA is the principal nonspecific slow porin of

Acinetobacter baumannii. J Bacteriol 194:4089-96.

56.   Iyer R, Moussa SH, Durand-Reville TF, Tommasi R, Miller A. 2018. *Acinetobacter*

baumannii OmpA Is a Selective Antibiotic Permeant Porin. ACS Infect Dis 4:373-381.

57.   Powers MJ, Trent MS. 2018. Phospholipid retention in the absence of asymmetry

strengthens the outer membrane permeability barrier to last-resort antibiotics. Proc Natl

Acad Sci U S A 115:E8518-E8527.

58.   Gwin CM, Prakash N, Christian Belisario J, Haider L, Rosen ML, Martinez LR, Rigel

NW. 2018. The apolipoprotein N-acyl transferase Lnt is dispensable for growth in

Acinetobacter species. Microbiology (Reading) 164:1547-1556.

59.   Haas TM, Laventie BJ, Lagies S, Harter C, Prucker I, Ritz D, Saleem-Batcha R, Qiu D,

Huttel W, Andexer J, Kammerer B, Jenal U, Jessen HJ. 2022. Photoaffinity Capture

Compounds to Profile the Magic Spot Nucleotide Interactomes. Angew Chem Int Ed

Engl 61:e202201731.

60.   Diez S, Ryu J, Caban K, Gonzalez RL, Jr., Dworkin J. 2020. The alarmones (p)ppGpp

directly regulate translation initiation during entry into quiescence. Proc Natl Acad Sci U

S A 117:15565-15572.

61.   Levenson-Palmer R, Luciano DJ, Vasilyev N, Nuthanakanti A, Serganov A, Belasco JG.

2022. A distinct RNA recognition mechanism governs Np(4) decapping by RppH. Proc

Natl Acad Sci U S A 119.

62.   Law RJ, Hamlin JN, Sivro A, McCorrister SJ, Cardama GA, Cardona ST. 2008. A

functional phenylacetic acid catabolic pathway is required for full pathogenicity of

Burkholderia cenocepacia in the *Caenorhabditis elegans* host model. J Bacteriol

190:7209-18.

63.   Musthafa KS, Sivamaruthi BS, Pandian SK, Ravi AV. 2012. Quorum sensing inhibition

1015   in *Pseudomonas aeruginosa* PAO1 by antagonistic compound phenylacetic acid. Curr

1016   Microbiol 65:475-80.

1017   64.   Lightly TJ, Frejuk KL, Groleau MC, Chiarelli LR, Ras C, Buroni S, Deziel E, Sorensen

1018   JL, Cardona ST. 2019. Phenylacetyl Coenzyme A, Not Phenylacetic Acid, Attenuates

1019   CepIR-Regulated Virulence in *Burkholderia cenocepacia*. Appl Environ Microbiol 85.

1020   65.   Geisinger E, Isberg RR. 2015. Antibiotic modulation of capsular exopolysaccharide and

1021   virulence in *Acinetobacter baumannii*. PLoS Pathog 11:e1004691.

1022   66.   Gomori G. 1955. Preparation of buffers for use in enzyme studies. Methods in

1023   Enzymology 1:143.

1024   67.   Clementi EA, Marks LR, Roche-Hakansson H, Hakansson AP. 2014. Monitoring

1025   changes in membrane polarity, membrane integrity, and intracellular ion concentrations

1026   in *Streptococcus pneumoniae* using fluorescent dyes. J Vis Exp

1027   doi:10.3791/51008:e51008.

1028   68.   McCoy KM, Antonio ML, van Opijnen T. 2017. MAGenTA: a Galaxy implemented tool

1029   for complete Tn-Seq analysis and data visualization. Bioinformatics 33:2781-2783.

1030   69.   Jensen PA, Zhu Z, van Opijnen T. 2017. Antibiotics Disrupt Coordination between

1031   Transcriptional and Phenotypic Stress Responses in Pathogenic Bacteria. Cell Rep

1032   20:1705-1716.

1033   70.   Zhu Z, Surujon D, Ortiz-Marquez JC, Huo W, Isberg RR, Bento J, van Opijnen T. 2020.

1034   Entropy of a bacterial stress response is a generalizable predictor for fitness and antibiotic

1035   sensitivity. Nat Commun 11:4365.

71. Shishkin AA, Giannoukos G, Kucukural A, Ciulla D, Busby M, Surka C, Chen J,

Bhattacharyya RP, Rudy RF, Patel MM, Novod N, Hung DT, Gnirke A, Garber M,

Guttman M, Livny J. 2015. Simultaneous generation of many RNA-seq libraries in a

single reaction. Nat Methods 12:323-5.

72. Huang Y, Sheth RU, Kaufman A, Wang HH. 2020. Scalable and cost-effective

ribonuclease-based rRNA depletion for transcriptomics. Nucleic Acids Res 48:e20.

73. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing

reads. 2011 17:3.

74. Benjamini Y, & Hochberg, Y. 1995. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. Journal of the Royal Statistical Society Series B

(Methodological):289-300.

75. Benjamini Y, Krieger, A. M., & Yekutieli, D. 2006. Adaptive linear step-up procedures

that control the false discovery rate. Biometrika 93:491-507.

1049

1050

Figure Legends

Figure 1. Uncovering the limitations of high-level FQR strains with enhanced RND pump

expression. (A) Diagram for the three experimental approaches as described (Materials and

Methods; created with BioRender.com). 1) Approach #1: Transposon insertion libraries were

created in FQR strains overexpressing efflux pumps and the drug susceptible WT strain.

Insertion abundance across the genome between the two strain types was compared to identify

genes with fewer insertions in the FQR strain background than in WT (see B for example). 2)

Approach #2: Transposon insertion banks were challenged with sub-MIC levels of CIP, parallel

to those grown without drug. Fitness of all transposon mutants was calculated and compared

between the strain types. 3) Approach #3: Strains were grown in presence or absence of CIP to

exponential or post-exponential phase in presence or absence of drug and transcriptomic data

was obtained to discover differentially regulated pathways. (B) Example of insertion abundance

in a single gene cluster in two strain backgrounds. (C) Activating mutations in efflux pump

regulatory mutants specifically activate their cognate pumps. RNA was isolated from the noted

strains and relative transcription determined by q-rtPCR (Materials and Methods). Displayed is

mean fold change ± SEM of each target as compared to the parental GP strain (Log2 scale). At

least three independent biological replicates were used for each strain. FQR: fluoroquinolone

resistant. Mid: mid exponential phase. EPE: early post-exponential phase. CIP: ciprofloxacin.

Untx: untreated. WT: wildtype. GPL: *gyrAparCadeL*. GPN: *gyrAparCadeN*. GPS:

gyrAparCadeS. GP: *gyrAparC*.

1071

Figure 2: Discovery of genes with lowered tolerance for insertions in the high-level FQR

strains. (A-C) Volcano plots in which statistical significance is plotted as a function of the

differences in insertion abundance between each FQR strain and WT based on TRANSIT

algorithm (33). All adjusted inverse p-values = 0 were assigned the value of 0.005 to allow

graphing. Black & blue: genes with adjusted p-values<0.05. Values for all other genes are shown

in grey. (D-F) Number of reads plotted as a function of insertion site for noted genes as well as

flanking regions. Shown are: FQR: fluoroquinolone resistant. WT: wildtype. GPL:

gyrAparCadeL. GPN: *gyrAparCadeN*. GPS: *gyrAparCadeS*.

1080

Figure 3: Maintenance of cell envelope homeostasis supports the fitness of efflux pump overproducer strains. (A-C, F-H): Statistical significance plotted as a function of relative fitness comparing identical mutations in FQR and WT strains grown in the absence (A-C) or presence (F-H) of sub-MIC levels of CIP. (D) Growth rates at noted CIP concentrations to identify conditions allowing 20-30% growth inhibition in the FQR and WT strains. (A-C) Blue or (F-H) Red: Mutations above the FDR = 5%, q<0.05 as determined by Benjamini and Hochberg (74). Notable findings are labeled. All q-values < 0.000001 assigned the value of 0.000001 for graphing purposes. (E) Heatmap showing fitness of insertions in each gene in noted strain backgrounds. Selected genes with Tn-disruptions yielding lowered fitness in the FQR strains are shown in the presence of CIP. Gene names are followed by accession numbers which are preceded by "NDEONHPJ_" per the ATCC 17978 strain annotations. This data excludes genes characterized as essential and genes pertaining to prophages or plasmids. Shown are: WT: wildtype. GP: *gyrAparC*. GPS: *gyrAparCadeS*. GPN: *gyrAparCadeN*. GPL: *gyrAparCadeL*.

Figure 4: OmpA and BfmS are critical fitness determinants for *adeL*∆I335A336 strains during CIP exposure. (A) The Single Fitness pipeline of MAGenTA was used to evaluate the fitness of Tn mutants at each position across the loci shown, post CIP treatment. (B) Null mutations in the FQR strain backgrounds grown in the presence of CIP compared to the WT background. Same experimental data for the WT strain and single deletion mutants are shown in each graph for comparative purposes. Growth was monitored by taking absorbance measurements every 15 minutes in 96 well plates in presence of CIP (0.1 µg/ml for the WT, 55 µg/ml for the GPN, and 65 µg/ml for the GPL parent strains and deletion mutants in those respective backgrounds). Displayed are means ± SEM (n≥3). Of note, GPL∆*ompA* has an

1104 additional mutation in *lpxK* (I283V). Ciptx: CIP treated. WT: wildtype. GPL: *gyrAparCadeL*.

1105 GPN: *gyrAparCadeN*.

1106

Figure 5: LpsB and PbpG are required for the AdeN (GPN) and AdeL (GPL) mutants for

optimal fitness in presence of CIP. (A) Transposon mutant fitness post CIP treatment at each

insertion position across the selected loci. (B) Growth of the indicated strains in the presence of

CIP was monitored as in Fig. 4, displayed as mean ± SEM (n≥3). Data for the WT strain and

single deletion mutants from identical cultures are shown in each graph for comparative

purposes. (C) Growth of deletion mutants in LB + 65 µg/ml CIP + 1mM IPTG to induce

expression of the indicated genes complemented *in trans* in GPN (column 2) or GPL (column 4)

backgrounds. Data are means ± SEM (n≥6). Ciptx: CIP treated. IPTG: Isopropyl β-d-1-

thiogalactopyranoside. WT: wildtype. GP: *gyrAparC*. GPL: *gyrAparCadeL*. GPN:

gyrAparCadeN. EV: empty vector.

1117

Figure 6: Transcriptional and physiological response to pump overproduction. (AB)

Alterations in intracellular pH (pHi) in response to pump overproduction. Noted strains grown to

mid-exponential phase in the absence (A) or presence (B) of CIP. Internal pH (pHi) was

measured using 20µM BCECF-AM as described (Materials and Methods). Displayed are the

means ± SEM for each strain from n=5-8 biological replicates, with P values derived from one-

way ANOVA making multiple comparisons. * = $p < 0.05$ (0.0259); ns=not significant. (C-F)

Transcriptional alterations in response to pump overproductions are magnified in presence of

*adeL*ΔI335A336 mutation. Transcriptomic data (n=3) was analyzed comparing high-level FQR

pump hyper-expressing strains and the GP parent strain (Materials and Methods). Differences in

transcript levels for each comparison are shown via the heat maps with GP vs GPL (C, +CIP)

and (D, -CIP), GP vs GPN (E) and GP vs GPS in (F). This data excludes genes pertaining to

prophages or plasmids. A gene is listed if a significant change in gene expression was observed

in at least one condition within the panel. WT: wildtype. GPL: *gyrAparCadeL.* GPN:

gyrAparCadeN. GPS: *gyrAparCadeS.* GP: *gyrAparC.* Mid: mid-exponential phase. EPE: early

post-exponential phase.

1133

Figure 7: The presence of *adeL*ΔI335A336 in GPL results in overexpression of the *paa*

operon. RNA was isolated from GP and GPL strains and transcription of *paaA* gene was

quantified by q-rtPCR. Strains grown to exponential (A) or early post-exponential phase (B) in

absence of drug. Data are mean fold change in transcript levels ($2^{-\Delta\Delta CT}$) ± SEM of each target

(n=3) compared to parental GP strain, analyzed by two-tailed t-test. Significance marked by

asterisks; **, p=0.0099; ***, p=0.0004. (C, D) Alterations in transcript levels in CRISPRi-

depleted *paaA* strain, with knockdown verified as described (Materials and Methods). RNA was

prepared from cultures grown to mid-exponential (C) or early post-exponential phase (D). Data

are transcript levels of each target relative to nontargeting negative control ($2^{-\Delta\Delta CT}$) ± SEM

(n=3), analyzed by multiple unpaired t test analyses following the two-stage step-up Benjamini,

Krieger, and Yekutieli, false discovery rate of 5.00% (75). Significance: *, p=0.020-0.034; **,

p=0.002-0.003; ****, p<0.001. (E-G). Acidification of cytosol as a consequence of *paaA*

depletion. Internal pH (pHi) determined during mid-exponential phase in absence (E, F) or

presence (G) of CIP. (F). Intracellular pH in early post-exponential phase. Data are mean ± SEM

from n=5. P values were derived from an unpaired t test comparing the mean pHi values between

the two strains. (***, p=0.0009 in E; *, p=0.0420 in G) & (ns, p=0.8479 in F). Untx: Untreated.

Ciptx: CIP treated at 45µg/mL. Mid: grown to mid-exponential phase. EPE: grown to early post-

exponential phase. GPL: *gyrAparCadeL*. GP: *gyrAparC*.

1152

Figure 8: *paaA* is significantly upregulated during CIP exposure. (A,B) The *paaA* transcript

(shown with L2FCSE) is upregulated after CIP exposure independently of strain background.

Strains were grown to noted growth phases, with data derived from RNAseq experiments (Fig. 6;

Materials and Methods). Significance determined by s values from triplicate cultures for each (s

$\leq$ 1.11E-03). (C) *paaA* upregulation is pump dependent. Strains were grown to mid exponential

phase and their RNA were used to quantify *paaA* transcript levels by q-rtPCR. Bars show mean

transcript levels ($2^{-\Delta\Delta CT}$) $\pm$ SEM (n=3) relative to parent strains (WT or GP), significance

determined by Ordinary one-way Anova (**, p=0.0027). (D) Persistence of *adeL*ΔI335A336

single mutant (in WT strain background) is unaffected by depletion of *paaA*. The indicated

strains were grown to exponential phase with aTc inducer to deplete *paaA* transcript, then

exposed to CIP at ~20X MIC for the noted times prior to plating on LB agar plates in absence of

drug to determine CFU/mL. The mean CFU/mL $\pm$ SEM are shown (n=3). The dotted line

indicates LOD, the limit of detection for the assay (375 CFU/mL). Mid: grown to mid-

exponential phase. EPE: grown to early post-exponential phase. aTc: anhydrotetracycline. WT:

wildtype. GPL: *gyrAparCadeL*. GPN: *gyrAparCadeN*. GPS: *gyrAparCadeS*. GP: *gyrAparC*.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8