1 SC-Track: a robust cell tracking algorithm for generating accurate single-cell lineages from

2 diverse cell segmentations

Chengxin Li[1,2], Shuang Shuang Xie[2], Jiaqi Wang[2], Septavera Sharvia[4], Kuan Yoow Chan[1,2,3]

[1] Department of Cardiovascular Medicine, The Second Affiliated Hospital, Zhejiang University School

of Medicine, Hangzhou, 310058, P. R. China.

[2] Centre for Cellular Biology and Signalling, Zhejiang University-University of Edinburgh Institute,

Zhejiang University School of Medicine, Zhejiang University, Haining, 314400, P. R. China.

[3] College of Medicine and Veterinary Medicine, The University of Edinburgh, Edinburgh, EH4 2XR,

UK.

[4] Department of Computer Science, University of Hull, Hull, HU6 7RX, UK.

Correspondence to:

kychan@intl.zju.edu.cn

16
17

Abstract

Computational analysis of fluorescent timelapse microscopy images at the single-cell level is a powerful

approach to study cellular changes that dictate important cell fate decisions. Core to this approach is the

need to generate reliable cell segmentations and classifications necessary for accurate quantitative

analysis. Deep learning-based convolutional neural networks (CNNs) have emerged as a promising

solution to these challenges. However, current CNNs are prone to produce noisy cell segmentations and

classifications, which is a significant barrier to constructing accurate single-cell lineages. To address

this, we developed a novel algorithm called Single Cell Track (SC-Track), which employs a hierarchical

probabilistic cache cascade model based on biological observations of cell division and movement

dynamics. Our results show that SC-Track performs better than a panel of publicly available cell

trackers on a diverse set of cell segmentation types. This cell-tracking performance was achieved

without any parameter adjustments, making SC-Track an excellent generalised algorithm that can

maintain robust cell-tracking performance in varying cell segmentation qualities, cell morphological

appearances and imaging conditions. Furthermore, SC-Track is equipped with a cell class correction

function to improve the accuracy of cell classifications in multi-class cell segmentation time series.

These features together make SC-Track a robust cell-tracking algorithm that works well with noisy cell

instance segmentation and classification predictions from CNNs to generate accurate single-cell

lineages and classifications.

Keywords: timelapse microscopy imaging, single-cell tracking, cell division, deep learning,

convolutional neural networks, cell cycle.

Introduction

The analysis of time-resolved fluorescent microscopy images to obtain cellular dynamics at the single-cell level has enabled the detailed study of cellular events previously inaccessible to conventional cell biological approaches [1,2]. This approach has led to the delineation of biological processes that induce a variety of cell fate decisions [3–7]. Core to this analysis is the extensive use of fluorescent markers to mark single cells and to classify cellular states. However, generating single-cell tracks from these fluorescent timelapse microscopy images is often challenging, requiring laborious optimisations of fluorescent markers and imaging conditions [2]. These optimisations are essential as good-quality cell segmentations are critical for accurate lineage tracing while avoiding biological artefacts caused by phototoxicity.

Deep learning-based convolutional neural networks (CNNs) are increasingly employed to overcome the inherent limitations of conventional fluorescence-based microscopy approaches [8]. Among the most successful applications is the use of autoencoder CNNs, enabling computationally efficient image restoration of low-light microscopy images for deconvolution, denoising and generating super-resolution image reconstructions [9]. Another area where CNNs have been successfully deployed is in the automated segmentation and classification of microscopy images [10–13]. Recent work has shown that CNNs perform well in automatically detecting, segmenting and classifying heterogeneous cellular features of microscopy images, a task previously requiring time-consuming manual human annotations [13–16].

However, applying deep learning CNNs in the automated segmentation and classification of fluorescent microscopy images presents another challenge for reliable cell tracking. State-of-the-art CNNs are inherently noisy, with instances where objects fail to be detected or are misclassified [12,13,17,18]. These inaccuracies pose a significant challenge for widely used cell tracking solutions as these algorithms are not designed to confront the various object classification errors derived from CNNs. Thus, extensive finetuning of cell tracking parameters and manual corrections of cell tracking outputs are required, posing a significant barrier for many biologists as they often lack the technical know-how and time to undertake such tasks.

To address this challenge, we developed a robust generalised cell tracking algorithm called

Single Cell Track (SC-Track). SC-Track employs a hierarchical probabilistic cache-cascade model

inspired by biological observations of cell division and movement dynamics of mammalian cells. We

show that SC-Track can generate robust single-cell tracks from segmentation outputs from CNNs that

contain missing segmentations to false detections. Furthermore, SC-Track functions well with whole

cell or nuclear segmentations of diverse morphologies and imaging conditions without parameter

optimisations. Finally, SC-Track can take noisy cell instance classifications and provide smoothed

classification tracks to quantify cellular events accurately.

Materials and methods

Tracking algorithm overview

SC-Track employs a tracking-by-detection approach whereby detected cells are linked between frames.

A TrackTree data structure was used to store the tracking relationships between each segmented cell

temporally and spatially (Figure 1A). Each branch of the TrackTree represents a single-cell lineage of

the tracked instance of a segmented cell, where branch divisions indicate cell division events and the

nodes on the branches represent the segmented instances of individual cells in a specific frame. The

extracted features of the segmented cell are contained in each node of the TrackTree branch.

During the tracking process, SC-Track initialises the TrackTree list with all cells from the initial

frame, representing the initial single-cell tracks for the entire time-lapse sequence. To reduce

computational costs, SC-Track will attempt to connect each segmented instance with its corresponding

cell from the previous frame using a hierarchical tracking approach. SC-Track will initially examine the

intersection over union (IoU) of the area between segmented cells between the current frame and

preceding frame (Figure 2A). Segmented cells with only one overlapping segmentation are assumed to

be high-confidence links and are assigned to the corresponding TrackTree. In situations with multiple

segmented cells with overlapping IoUs, SC-Track will determine the correct cells by maximising the

similarity index between candidate cells between frames. When no segmented cell in the current frame

overlaps with a segmented cell from the previous frame, SC-Track will expand the search area to

identify possible candidates.

By recursively searching for candidate segmented cells from the previous frame, virtually all

segmented cells can be accurately assigned to the correct TrackTree. If there are more segmented cells

than the number of cached TrackTrees, three possible scenarios will be considered: (1) The orphan cell

is a false detection; (2) The orphan cell is an actual detection of a cell recently migrated into the field

of view; (3) A cell division event has occurred leading to the generation of daughter cells.

Detecting and assigning cell division events

In scenario three above, SC-Track will determine if a cell division occurred by searching for a potential

mother cell in the mitotic state in the preceding frame (Figure 2B). When the segmented cell contains

cell cycle classifications, SC-Track will allow cell division events at the TrackTree nodes where the

mother cell is classified as in mitosis (M phase). However, a cell cycle-independent approach will be

applied if no cell cycle information is available. In such cases, SC-Track will attempt to determine if a

cell division event has occurred by matching an orphan segmented cell in the current frame to a potential

mother cell in the previous frame using morphological data of the segmented cells.

To enable robust detection of cell division events without cell cycle data, SC-Track applies a

series of rules based on well-established principles observed from mammalian cells undergoing cell

division [19,20]. When assigning a potential mother-daughter association from a possible cell division

event, the following criteria must be met: (1) At least one unlinked segmented cell was found; (2) The

segmented mother cell in the previous frame must be at least $1.3 \times$ the size of the segmented daughter

cell in the following frame. (3) The candidate mother cell has not undergone a cell division event in the

previous 20 frames. (4) A candidate mother cell is identified in the expanded search area of the unlinked

segmented cell. If a suitable candidate mother cell was identified in the previous frame for the orphan

segmented cell, the TrackTree will be branched accordingly. However, if no appropriate candidate

mother cells are found, SC-Track will assume a new detection event may have occurred. The following

section will discuss the algorithm employed to address the possibility of new detection events and the

initialisation of new TrackTrees.

Cache matching frames to address false and missing detection events

The stochastic nature of CNNs may lead to instances where some cells fail to be detected or are false

detections [12,13,17,18], resulting in TrackTrees that contain inaccurate cell segmentations. We

developed a cache-matching algorithm to address the stochastic loss of detected instances in the

segmented cells (Figure 3A). In this process, SC-Track will follow a tiered response based on the

likelihood that the instance segmentations are accurate. Initially, SC-Track will focus on generating

high-confidence TrackTrees with cell segmentations that appear consecutively across multiple frames.

When cell segmentations cannot be linked to an initialised TrackTree from the previous frame, and no

suitable candidate mother cell is identified, SC-Track will search for up to five preceding frames for

potential orphan TrackTrees to link (Figure 3A). If an unlinked TrackTree is found within five frames,

SC-Track will assign the matched segmented cell to the corresponding TrackTree. When no matches

are found, SC-Track will assume a new detection event may have occurred, and a new TrackTree will

be initialised.

After all segmented frames are analysed, SC-Track will review all TrackTrees to filter high-

confidence TrackTree initialisations (Figure 3B). SC-Track assumes that TrackTrees that contain only

one node are false detections and will exclude these TrackTrees in the linking process. For TrackTrees

containing at least two nodes, SC-Track will assume that some of these TrackTrees are incomplete

TrackTree fragments. To identify and repair these incomplete TrackTree fragments, SC-Track will

survey the terminal ends of all TrackTrees to determine instances where a terminal end of one TrackTree

is spatially and temporally proximal to a terminal end of another TrackTree. When such instances are

detected, SC-Track will attempt to pair the first instance of an unlinked TrackTree with the last instance

of another unlinked TrackTree. The linking of TrackTrees will follow two requirements: (1) The last

segmented cell of the preceding unlinked TrackTree is within the expanded search area of the first

segmented cell of the unlinked TrackTree; (2) The last segmented cell of the preceding TrackTree is

within three subsequent frames of the first segmented cell of the unlinked TrackTree. If more than one

candidate TrackTree is available for linkage, the similarity index of the last instances of both competing

candidate TrackTrees will be calculated, and the candidate TrackTree with the highest similarity will be

linked. After the linking of terminal ends of track trees is completed, short TrackTree initialisations that

span less than 10 frames are discarded to remove false detection instances. Finally, the intervening gaps

between the linked TrackTrees will be filled with the preceding cached cell segmentation in the linked

TrackTree.

Instance classification smoothing

Instance classification of multi-class cell segmentations is often noisy [21]. We have implemented a

class smoothing function to smooth out noisy classification of cells that transition from one cellular

state to another. We developed the TrackTree Class Smoothing (TCS) algorithm to automatically correct

the predicted results of cell type classifications (Figure 3C). TCS assumes that a cell classification

change is more likely to occur in a time series when the same cell maintains the same cell classification

over several frames. To evaluate the accuracy of the cell class change, TCS adopts a probabilistic cached

search model. This search process is confined to the individual branch of the TrackTree and does not

extend beyond the cell division branch.

The TCS probabilistic cached search model functions with the following logic: During the

initialisation of the TrackTree, TCS will automatically adopt the initial classification of the detected cell

instance as the default class. When TCS detects an instance where the tracked cell undergoes a cell

classification change to Type A, the algorithm will undertake a forward search on the TrackTree to count

the number of Type A classifications in the subsequent 9 frames. When the number of nodes classified

as Type A exceeds a probability threshold of 60%, TCS will conclude that a change in cell classification

has occurred and will update the default classification as Type A. Otherwise, the node where Type A

was first detected will be corrected to the default cell classification. Exiting the default Type A

classification occurs when a Type B class change is detected. To determine if a class change has

occurred, TCS performs a forward TrackTree search in the subsequent 9 frames for the Type B

classification. When the number of nodes of Type B classifications exceeds the probability threshold of

60%, the new default Type B classification is adopted. This process can be repeated to multiple cell

classifications.

Calculating similarity index when connecting segmented cells between frames

When there is more than one segmented cell overlapping with the previous frame, SC-Track will select

the segmented cell with the highest similarity value with the segmented cell in the previous frame. SC-

Track will employ the following formula to determine the similarity value of the possible candidate

pairs between frames:

$$S_i = \{P_1, P_2, \dots P_n\}, \text{where } P_n = (x_n, y_n)$$

$$sm_{i,j} = IoU(S_i, S_j) + Dis(S_i, S_j) + Sps(S_i, S_j) + Sas(S_i, S_j) + \Delta(S_i, S_j)$$

S_i represents the set of contour points in a 2D space defined by x_n, y_n for points $P_{1 \to n}$ of a cell. $sm_{i,j}$

represents the similarity index between the segmented cell i in the previous frame and the segmented

cell j in the subsequent frame. Dis is the calculated distance between the centroid of the segmented cell

i in the previous frame and the centroid of the segmented cell j in the current frame. IoU represents the

intersection over union of the contours of cells i and j. Sps represents the shape similarity value [22],

and Sas represents the area similarity of the two cells. $\Delta(S_i, S_j)$ represents additional supplementary

features, such as the similarity in the variance or total intensity of fluorescent signals from segmented

cells. To calculate $IoU(S_i, S_j)$, $Dis(S_i, S_j)$, $Sps(S_i, S_j)$, $Sas(S_i, S_j)$, and $\Delta(S_i, S_j)$, the following

formula was employed:

$$IoU(S_i, S_j) = \frac{intersection(S_i, S_j)}{union(S_i, S_j)}$$

$$Dis(S_i, S_j) = \frac{1}{10^{-5} + \sqrt{(s_{ix} - s_{jx})^2 + (s_{iy} - s_{jy})^2}}$$

$$Sps(S_i, S_j) = \sum_{n=1\dots7} \left| \frac{1}{m_n^{S_i}} - \frac{1}{m_n^{S_j}} \right|, \text{ where } m_n^{S_i} \text{ represents the seven Hu Moments.}$$

$$Sas(S_i, S_j) = \frac{min(S_{i_{Area}}, S_{j_{Area}})}{max(S_{i_{Area}}, S_{j_{Area}})}, \text{ where } S_{n\ Area} = \frac{1}{2} \left| \sum_{i=1}^{n} (x_i \cdot y_{i+1} - y_i \cdot x_{i+1}) \right|$$

$$\Delta(S_i, S_j) = \{\frac{mean(s_i)}{mean(s_j)}, \frac{var(s_i)}{var(s_j)}\}, \text{ where } mean(s_i) < mean(s_j) \text{ and } var(s_i) < var(s_j)$$

198   Bounding box search area expansion

199   When no segmented cell in the preceding frame is detected in the segmented area of a cell in the current

200   frame, SC-Track will expand its search area to search for potential candidates. The expansion of the

201   search area utilises the bounding box of the segmented cell, which is expended with the following

202   formula:

$$Bc = Pos(x_1, x_2, y_1, y_2), where\ x_1 < x_2\ and\ y_1 < y_2$$

203    $$Ec = \alpha \cdot Bc$$

$$= Pos(x_1 - \alpha \cdot (x_2 - x_1), x_2 + \alpha \cdot (x_2 - x_1), y_1 - \alpha \cdot (y_2 - y_1), y_2 + \alpha \cdot (y_2 - y_1))$$

Bc represents the bounding box of a cell. Pos represents the position of the bounding box with the

minimum value of the segmented cell in the x-axis and y-axis represented by x_1 and y_1 while the

maximum value as x_2 and y_2 respectively. Ec represents the expanded bounding box where potential

cell candidates located in the current frame can be matched to the previous frame, α represents the

coefficient for expanding the bounding box. By default, α is set to 1.

Evaluation of cell tracking performance using MOTA and IDF1

To evaluate the performance of cell trackers in accurately tracking segmented cells, we used

performance measures established in the Multiple Object Tracking (MOT) framework, which includes

IDF1 [23] and MOTA [24,25]. IDF1 measures how long a tracker accurately assigns segmented cells

to the correct single-cell lineage over time. It represents the ratio of correctly identified detections over

the average ground-truth and computed detections [23]. IDF1 is calculated from the following formula:

$$IDF1 = \frac{2IDTP}{2IDTP + IDFP + IDFN}, where\ IDP = \frac{IDTP}{IDTP + IDFP}\ and\ IDR = \frac{IDTP}{IDTP + IDFN}$$

IDP represents the cell tracking identification precision. It is computed as the average ratio of accurately

identified true positives divided by the sum of accurately identified true positives and inaccurately

classified false positives. IDR represents the identification recall computed as the average ratio of

accurately identified true positives divided by the sum of accurately identified true positives and failed

detections of each single cell track.

The Multiple objects tracking accuracy (MOTA) measures the overall accuracy of the tracker

performance by measuring how often a mismatch occurs between the tracking results and the ground

truth [24,25]. The MOTA score is determined from the total number of errors for false positives (*FP*),

missed targets (*FN*), and identity switches (*IDsw*) normalised over the total number of ground-truth (*GT*)

tracks. This measure is computed using the following formula:

227
$$MOTA = 1 - \frac{\sum_t (FN_t + FP_t + IDsw_t)}{\sum_t GT_t}$$

The cell tracking outputs used to calculate the IDF1 and *MOTA* values are available from Zenodo:

https://zenodo.org/record/8284987. The Python scripts used to analyse the cell tracking results can be

found on GitHub: https://github.com/chan-labsite/SC-Track-evaluation.

Evaluating cell division detection events using the CDF1 score

Generating single-cell lineages over multiple cell division events requires the reliable detection of cell

division events and the accurate assignment of mother-daughter relationships. We established the Cell

Division *F1* score (CDF1) as IDF1 and MOTA do not measure cell division events reliably. This metric

is based on the principles of the widely used *F1* score and is defined by the formula:

237
$$CDF1 = \frac{2CDTP}{2CDTP + CDFP + CDFN}$$

CDTP indicates an actual positive cell division event, where both daughter cells of a cell division event

are accurately identified and assigned to the correct TrackTree. *CDFP* indicates a false positive cell

division event, where daughter cells are incorrectly assigned to a TrackTree and classified as a cell

division event. *CDFN* indicates a false negative cell division event, where a cell division event occurred

but is not detected or the mother-daughter cells were inaccurately assigned to the wrong TrackTree. The

cell   tracking   outputs   used   to   benchmark   the   CDF1   results   are   available   from   Zenodo:

https://zenodo.org/record/8284987. The Python scripts used to analyse the cell tracking results can be

found on GitHub: https://github.com/chan-labsite/SC-Track-evaluation.

Generation of datasets for cell tracking

Two cell lines with distinct morphological appearances were used to generate the imaging data to

develop and test SC-Track. hTERT-RPE1 cells endogenously tagged with fluorescent mScarlet-PCNA

were grown in DMEM/F-12 (Sigma, D6421) supplemented with 10% FBS (ExCell Bio, FSP500), 1×

GlutaMAX (Gibco, 35050-061), 7.5% sodium bicarbonate (Sigma). MCF10A cells endogenously

tagged with fluorescent mScarlet-PCNA were grown in DMEM/F-12 (Sigma, D6421) supplemented

with 5% heat-inactivated horse serum (Biological Industries, 04-124-1A), 1× GlutaMAX (Gibco,

35050-061), 10 µg/ml insulin (Biological Industries, 41-975-100), 10 ng/ml cholera toxin (Sigma-

Aldrich, #C-8052), 20 ng/ml EGF-β (Thermo Fisher, PHG0311), 0.5 mg/ml Hydrocortisone (MCE,

HY-N0583). These cells were seeded in 8-Well chambered glass bottom slides (Cellvis, C8-1.5H-N) for

two days before being imaged under a Nikon Ti2 inverted widefield fluorescence microscope equipped

with a Lumencor Sola SE 365 as a light source. The cells were placed in an Okolab stage incubator

(OKO) at 37°C with 5% CO_2 and 80% humidity. The cells were observed under a 20× plan apo objective

(NA 0.75), and images were captured using a Photometrics Prime BSI camera with a pixel resolution

of 2048×2048. The following filter sets were used (mCherry: 560/40 nm EX, 630/75 nm EM). A single

widefield image was taken in the mCherry channel (1% power, 200ms) at each stage at 5-minute

intervals for up to 48 hours. A DIC image was captured at each time point (5% power, 100ms).

The timelapse microscopy images used to develop SC-Track were generated from cells cultured

under the conditions described above. The images were saved as individual multi-channel TIFF files.

Four time-lapse movies with varying cell densities were generated (Supplementary Table 1). These

datasets were automatically segmented using a custom pre-trained StarDist model [12] and manually

corrected using the VGG Image Annotator (VIA) [26] to remove false and inaccurate classifications.

The annotated files contain the cell contour and "cell cycle phase" class information. The contour

information was converted into a mask with values ranging from 1 to 255. The uncorrected and

corrected mask images, along with the original mCherry channel image, constitute the datasets used to

finetune the tracking parameters of SC-Track.

For the testing datasets, three RPE1 microscopy timelapse image series and two MCF10A

microscopy timelapse image series were automatically segmented using our custom-trained StarDist

model and manually corrected to ensure accuracy of the instance segmentations, cell classifications and

identity of single-cell lineages (Supplementary Table 2). The imaging conditions were described above

with a sampling frequency of 5 minutes. To test the reliability of SC-Track to track segmented cells

with missing or false positive instances accurately, we utilised the uncorrected segmentations of the

testing dataset (Supplementary Table 3). In addition, to assess how SC-Track can cope with varying

levels of missing cell segmentations, we randomly deleted additional segmented cells from each frame

to varying degrees to simulate higher levels of missed segmentations (Supplementary Table 4). The in-

house generated segmentation masks, custom-trained StarDist model, and ground truth tracking results

used in testing SC-Track can be obtained from Zenodo: https://zenodo.org/record/8284987.

Cell Tracking Challenge dataset

We used the silver reference segmentation results from the Cell Tracking Challenge to test the SC-Track

performance with various cell types and segmentation modes [27]. The silver reference datasets are

derived from the uncorrected segmentation results obtained from various custom-trained CNNs applied

on mammalian cell lines with different morphological appearances and imaging conditions

(Supplementary Table 5). The silver reference masks and ground truth tracking results were obtained

from the Cell Tracking Challenge website (http://celltrackingchallenge.net/2d-datasets/).

Generation of single-cell lineages from segmentation masks

The segmentation results from the various evaluation datasets were used to measure the cell-tracking

performance of SC-Track and three other trackers: pcnaDeep [28], Deepcell-tracking [29], and

TrackMate [30,31]. For cell tracking experiments involving in-house generated testing datasets, the

segmentation results in the form of a VGG image annotator (VIA2) compatible JSON file containing

cell cycle class information of each segmented cell was used [26]. The data in the JSON files were read

directly by SC-Track and pcnaDeep to generate the cell lineage tables. The cell segmentation data in

the JSON files were converted into greyscale multi-TIFF image files before being read by TrackMate

and Deepcell-tracking, as both software packages lack the function to read JSON files directly.

Default tracking settings were applied to SC-Track, pcnaDeep and Deepcell-tracking. The Lap

tracker algorithm was used with default tracking settings with TrackMate. We could not perform the

cell tracking experiments with the Cell Tracking Challenge dataset on pcnaDeep as it requires cell cycle

class information to function [28]. The scripts used to generate the cell tracking results are available

from GitHub: https://github.com/chan-labsite/SC-Track-evaluation.

Evaluating automated cell cycle class correction

To evaluate the cell cycle class accuracy from the TCS function of SC-Track, we computed the F1 score

of uncorrected raw cell classifications obtained from our custom pre-trained StarDist model and

compared it with the cell cycle corrected F1 score of SC-Track (Supplementary Table 3). The F1 score

is calculated with the following formula:

$$F1 = 2 \cdot \frac{Precision \cdot Recall}{Precision + Recall} \text{ where } Precision = \frac{TP}{TP+FP} \text{ and } Recall = \frac{TP}{TP+FN}$$

The ground truth cell cycle classification used to compute the F1 score was obtained by manually

correcting the automated cell cycle classifications from our custom StarDist model. The JSON file

containing the raw uncorrected cell segmentations and the cell cycle classification data used to compute

the F1 results is available on Zenodo: https://zenodo.org/record/8284987. The scripts used to compute

the F1 scores of individual cell cycle phases are available on GitHub: https://github.com/chan-

labsite/SC-Track-evaluation.

Runtime and multi-platform compatibility testing

We conducted runtime efficiency tests on Windows, Linux, and macOS platforms. All tests were

performed using the same dataset and repeated three times. The Windows platform was equipped with

an AMD R7 3700X CPU, RTX 2080 GPU, and 16GB of RAM. The Linux platform was equipped with

an Intel i7 11800H CPU, RTX 3050Ti GPU, and 16GB of RAM. The macOS platform is a 2021

MacBook Pro with an M1 processor and 8GB of RAM.

Results

Evaluation of SC-Track cell tracking performance

The overall performance of SC-Track in cell tracking was measured using two metrics: the Multi-Object

Tracking Accuracy (MOTA)[24,25] and the harmonic mean of Identification Precision and Recall

(IDF1)[23]. We employed two cell tracking metrics because IDF1 is more sensitive to the total duration

of incorrect track assignments, while MOTA is more sensitive to the total number of track switches [23].

We also introduced a new metric, the Cell Division F1 score (CDF1), to measure the cell tracker's

ability to reliably detect cell division events and accurately assign mother-daughter cell relationships.

The CDF1 score is essential for measuring single-cell lineage reconstructions over multiple cell division

events, which the IDF1 and MOTA scores cannot reliably capture. For comparison, we benchmarked

SC-Track against three other freely available cell tracking algorithms that provide similar functionalities:

TrackMate [30,31], Deepcell-tracking [29], and pcnaDeep [28]. The initial tests focused on generating

single-cell tracks from nuclear masks obtained under ideal conditions, using manually corrected nuclear

segmentation masks with accompanying cell cycle classifications with a 5-minute temporal resolution

(Table 1). The results show that with ideal segmentation results, SC-Track performed best in all three

measured metrics (Table 1).

To further test SC-Track in generating accurate single-cell lineages, we resampled our original

test dataset to mimic imaging time intervals of 10, 15 and 20 minutes. The increase in time intervals

poses a more challenging cell tracking problem as each cell has more time to migrate and change

morphologically between successive frames. Our results show that SC-Track gives the best overall

IDF1, MOTA and CDF1 scores in the 5-minute interval, but as expected, its performance is reduced at

longer intervals (Figure 4). Despite the reduced performance, SC-Track maintained the best overall

performance in all three metrics across all time intervals tested. These results indicate that SC-Track

works well with different temporal resolutions (Figure 4).

CNN instance segmentations generally display errors ranging from missing to inaccurate

segmentations [12,13,17,18]. To assess the various tracker's ability to generate reliable single-cell

lineages from noisy CNN-based cell segmentations, we repeated the tests with uncorrected image

segmentations from our custom-trained StarDist models (Supplementary Table 3). Our results show

decreased tracking accuracy for all tested trackers, while SC-Track again gave the best overall

performance, maintaining the highest average MOTA, IDF1 and CDF1 scores throughout (Table 2). To

further examine SC-Track's ability to overcome missing instances of cell segmentations, we generated

a synthetic test dataset where cell instances were randomly removed at varying degrees (Supplementary

Table 4). Our results show that SC-Track's cache matching algorithm can compensate for the loss of

instance detections while maintaining a high IDF1 and MOTA score throughout all tested conditions

(Figure 5A and 5B). Furthermore, despite increasing missing instance detections, SC-Track remains

reliable in detecting cell division events (Figure 5C).

To examine the generalisability of SC-Track, we used several publicly available microscopy

datasets containing diverse cell segmentations extracted from different cell types and imaging

conditions (Supplementary Table 5). We decided to use the silver reference segmentation results from

the Cell Tracking Challenge (CTC), as they were derived from the segmentation results of CNN models

[27]. These cell segmentations are based on a collection of timelapse microscopy images taken with

various imaging settings, cell morphologies and a mixture of nuclear to whole-cell segmentations [27].

Furthermore, the silver reference segmentations are accompanied by ground truth tracking results,

making these datasets an impartial real-life test for SC-Track. Our results show that SC-Track

consistently displayed the best cell tracking performance measured by MOTA and IDF1 scores for

nearly all the CTC datasets (Table 3). Using only the silver reference segmentation results, SC-Track

can reliably detect cell division events on most CTC datasets (Table 3). These results suggest that SC-

Track is a general cell-tracking algorithm which performs equally well on various cell segmentation

types and can be used under challenging conditions where the cell segmentation dataset exhibits high

levels of detection loss.

Instance classification smoothing of single cell tracks and runtime evaluations.

It is well established that CNNs occasionally misclassify multi-class instance segmentations. Instance

misclassifications are caused by objects that partially fit into a particular class or suboptimal imaging

conditions. The inherent noise in the cell classifications can pose a problem if accurate classifications

of cellular states are essential, such as in the quantification of cell cycle phases in an image time series

[28]. To overcome this problem in SC-Track, we developed a TrackTree Class Smoothing (TCS)

algorithm that employs a probabilistic cached class smoothing approach to identify cell phase transition

points accurately. To evaluate the utility of TCS, we measured the F1 scores of our custom-trained

StarDist model to classify the cells in our test dataset using the fluorescent PCNA cell cycle reporter

signal (Table 4). The results indicate that TCS can improve the F1 classification scores of all cell cycle

classes.

Finally, we conducted runtime tests for SC-Track to determine how long SC-Track takes to

generate single-cell tracks from cell segmentations. We measured the time taken to analyse cell

segmentations from microscopy timelapse series of varying lengths (50-500 frames) and compared it

with TrackMate, Deepcell-tracking, and pcnaDeep. Our results show that when working with small to

moderate imaging datasets, SC-Track had the best performance (Figure 6). Increasing the number of

analysed frames decreases processing speed (Figure 6), mainly because of the duration required to load

the time-lapse microscopy images before generating the single-cell lineages by SC-Track.

Discussion

We introduced SC-Track, a novel cell-tracking algorithm employing a hierarchical probabilistic cache

cascade model inspired by biologically relevant observations of cell division and movement dynamics.

SC-Track can generate highly accurate single-cell tracks from nuclear and cell segmentations of diverse

morphologies and imaging conditions. To better assess the ability of cell trackers to generate single-cell

lineages over multiple cell division events, we introduced a new metric called the Cell Division F1

(CDF1) score. SC-Track outperformed other tested tools in detecting cell division events under all

conditions tested.

SC-Track's hierarchical probabilistic cache-cascade model tolerates false or missing cell

segmentations caused by the stochastic nature of CNNs, reducing the need for extensive and time-

consuming manual corrections of image segmentations. We further implemented a cache smoothing

algorithm to reduce the stochastic noise in cell classifications from CNNs while increasing the accuracy

of the cell classifications in time series. These functionalities are computationally efficient, allowing

SC-Track to be run locally without requiring access to high-performance computing clusters.

Despite SC-Track advantages in cell tracking, there are some limitations. Firstly, SC-Track

experiences a reduction in computational efficiency as the number of frames increases. This could be

an issue for users who require cell tracking on high-throughput timelapse microscopy experiments.

Secondly, SC-Track currently supports only 2D image analysis. Thirdly, the SC-Track class smoothing

algorithm makes generalised assumptions that the degree of cell classification errors is equal across all

classes and that the frequency for each cell class appearing in a timelapse series is similar. These

assumptions, while reasonable, may not be valid under all biological conditions.

In summary, SC-Track solves a longstanding problem involving the use of CNNs in automated

segmentation and classification of cells from time-lapse microscopy images. By encoding biological

intuition in SC-Track's cell tracking algorithm, SC-Track performs well in diverse segmentation

qualities without finetuning parameters. A Python implementation of SC-Track is available, requiring

only cell segmentation data and microscopy images to function, making it easy to integrate into existing

analytical pipelines. These features together make SC-Track a valuable tool for biologists to help

generate robust single-cell lineages and cell classifications from their timelapse microscopy images.

Key points

- Deep learning-based convolution neural networks produce image segmentation errors that often

confound well-established cell tracking algorithms.

- SC-Track is a hierarchical cache-matching algorithm inspired by biological observations of cell

division and movement dynamics.

- SC-Track generates accurate single-cell lineages without parameter tuning from cell

segmentations of varying qualities, morphological appearances, and imaging conditions.

- SC-Track works with features extracted from cell segmentation masks, making it easy to

integrate into existing image analysis pipelines.

Acknowledgements

The authors thank Mikael Björklund for critically reading our manuscript and for the helpful

suggestions.

Funding

This work was supported by funds from Zhejiang University and the Fundamental Research Funds for

the Central Universities (2019QN30001) to KYC. Zhejiang University-University of Edinburgh (ZJE)

Institute of Zhejiang University funds SSX postdoctoral fellowship.

Author contributions

KYC conceived the study. KYC, CL, SSX and JW conceived the experiments. CL, SSX and JW

conducted the experiments. KYC, CL and SS wrote the manuscript. KYC, CL, SSX, JW and SS

analysed the results. All authors reviewed the manuscript.

Data and code availability

The raw and corrected cell segmentation results, raw cell tracking outputs, and raw analysis results are

available from Zenodo: https://zenodo.org/record/8284987. The silver reference masks and ground truth

tracking   results   are   available   from   the   Cell   Tracking   Challenge   website

(http://celltrackingchallenge.net/2d-datasets/). A Python implementation of SC-Track with its

corresponding usage documentation is available at GitHub: https://github.com/chan-labsite/SC-Track.

The scripts used to analyse the raw data and generate the figures presented in this manuscript are

available on GitHub: https://github.com/chan-labsite/SC-Track-evaluation.

Author Biographies

Chengxin Li is an MSc student at Zhejiang University-University of Edinburgh (ZJE) Institute at

Zhejiang University, China. His research interest is bioinformatics and computer science.

Shuang Shuang Xie is a postdoctoral researcher at ZJE Institute at Zhejiang University, China. Her

research interests include cancer biology, cell biology, molecular biology and biochemistry.

Jiaqi Wang is a PhD student at ZJE Institute at Zhejiang University, China. Her research interests

primarily focus on cell biology, cancer, and molecular biology.

Septavera Sharvia is a lecturer at the Department of Computer Science at the University of Hull, UK.

Her research interests span dependability analysis, explainable deep learning, and computational

modelling.

Kuan Yoow Chan is an assistant professor at ZJE Institute at Zhejiang University, China. His research interests span computational biology, cell biology, cell cycle research and cancer biology.

References

1. Skylaki S, Hilsenbeck O, Schroeder T. Challenges in long-term imaging and quantification of single-cell dynamics. Nat. Biotechnol. 2016; 34:1137–1144

2. Cooper S, Bakal C. Accelerating Live Single-Cell Signalling Studies. Trends Biotechnol. 2017; 35:422–433

3. Min M, Rong Y, Tian C, et al. Temporal integration of mitogen history in mother cells controls proliferation of daughter cells. Science (80-.). 2020; 368:1261–1265

4. Spencer SL, Cappell SD, Tsai F-C, et al. The Proliferation-Quiescence Decision Is Controlled by a Bifurcation in CDK2 Activity at Mitotic Exit. Cell 2013; 155:369–383

5. Cappell SD, Mark KG, Garbett D, et al. EMI1 switches from being a substrate to an inhibitor of APC/CCDH1 to start the cell cycle. Nature 2018; 558:313–317

6. Arora M, Moser J, Hoffman TE, et al. Rapid adaptation to CDK2 inhibition exposes intrinsic cell-cycle plasticity. Cell 2023; 186:2628-2643.e21

7. Barr AR, Cooper S, Heldt FS, et al. DNA damage during S-phase mediates the proliferation-quiescence decision in the subsequent G1 via p21 expression. Nat. Commun. 2017; 8:14728

8. Meijering E. A bird's-eye view of deep learning in bioimage analysis. Comput. Struct. Biotechnol. J. 2020; 18:2312–2325

9. Weigert M, Schmidt U, Boothe T, et al. Content-aware image restoration: pushing the limits of fluorescence microscopy. Nat. Methods 2018; 15:1090–1097

10. Ronneberger O, Fischer P, Brox T. U-Net: Convolutional Networks for Biomedical Image Segmentation. Lect. Notes Comput. Sci. (including Subser. Lect. Notes Artif. Intell. Lect. Notes Bioinformatics) 2015; 9351:234–241

11. Moen E, Bannon D, Kudo T, et al. Deep learning for cellular image analysis. Nat. Methods 2019; 16:1233–1246

12. Schmidt U, Weigert M, Broaddus C, et al. Cell detection with star-convex polygons. Lect.

Notes Comput. Sci. (including Subser. Lect. Notes Artif. Intell. Lect. Notes Bioinformatics)

2018; 11071 LNCS:265–273

13.   Pachitariu M, Stringer C. Cellpose 2.0: how to train your own model. Nat. Methods 2022 1912

2022; 19:1634–1641

14.   Falk T, Mai D, Bensch R, et al. U-Net: deep learning for cell counting, detection, and

morphometry. Nat. Methods 2019; 16:67–70

15.   Bilodeau A, Delmas CVL, Parent M, et al. Microscopy analysis neural network to solve

detection, enumeration and segmentation from image-level annotations. Nat. Mach. Intell.

2022 45 2022; 4:455–466

16.   Meijering E. Cell segmentation: 50 Years down the road [life Sciences]. IEEE Signal Process.

Mag. 2012; 29:140–145

17.   Caicedo JC, Roth J, Goodman A, et al. Evaluation of Deep Learning Strategies for Nucleus

Segmentation in Fluorescence Images. Cytom. Part A 2019; 95:952–965

18.   Stringer C, Wang T, Michaelos M, et al. Cellpose: a generalist algorithm for cellular

segmentation. Nat. Methods 2020 181 2020; 18:100–106

19.   Xie S, Skotheim JM. A G1 Sizer Coordinates Growth and Division in the Mouse Epidermis.

Curr. Biol. 2020; 30:916-924.e2

20.   Fantes PA, Grant WD, Pritchard RH, et al. The regulation of cell size and the control of

mitosis. J. Theor. Biol. 1975; 50:213–244

21.   Guerrero-Pena FA, Marrero Fernandez PD, Ren TI, et al. Multi-class Weighted Loss for

Instance Segmentation of Cluttered Cells. Proc. - Int. Conf. Image Process. ICIP 2018; 2451–

2455

22.   Hu MK. Visual Pattern Recognition by Moment Invariants. IRE Trans. Inf. Theory 1962;

8:179–187

23.   Ristani E, Solera F, Zou R, et al. Performance measures and a data set for multi-target, multi-

camera tracking. Lect. Notes Comput. Sci. (including Subser. Lect. Notes Artif. Intell. Lect.

Notes Bioinformatics) 2016; 9914 LNCS:17–35

24.   Bernardin K, Stiefelhagen R. Evaluating Multiple Object Tracking Performance: The CLEAR

MOT Metrics. EURASIP J. Image Video Process. 2008 20081 2008; 2008:1–10

25.   Leal-Taixé L, Taixé T, Milan A, et al. MOTChallenge 2015: Towards a Benchmark for Multi-

Target Tracking. 2015;

26.   Dutta A, Zisserman A. The VIA Annotation Software for Images, Audio and Video. Proc. 27th

ACM Int. Conf. Multimed. 2019; 2276–2279

27.   Maška M, Ulman V, Delgado-Rodriguez P, et al. The Cell Tracking Challenge: 10 years of

objective benchmarking. Nat. Methods 2023 207 2023; 20:1010–1020

28.   Gui Y, Xie S, Wang Y, et al. pcnaDeep: a fast and robust single-cell tracking method using

deep-learning mediated cell cycle profiling. Bioinformatics 2022; 38:4846–4847

29.   Moen E, Borba E, Miller G, et al. Accurate cell tracking and lineage construction in live-cell

imaging experiments with deep learning. bioRxiv 2019; 803205

30.   Tinevez JY, Perry N, Schindelin J, et al. TrackMate: An open and extensible platform for

single-particle tracking. Methods 2017; 115:80–90

31.   Ershov D, Phan MS, Pylvänäinen JW, et al. TrackMate 7: integrating state-of-the-art

segmentation algorithms into tracking pipelines. Nat. Methods 2022 197 2022; 19:829–832

Fig. 1: Schematic illustration providing an overview of SC-Track, the TrackTree data structure and analysis pipeline.

(A) Summary of the TrackTree data structure. Each linked segmented cell is tracked in a TrackTree. A node in a TrackTree branch represents an instance of the segmented cell in a particular frame with its accompanying cell segmentation information. A branching of a TrackTree represents a cell division event.

(B) Simplified overview of the analysis pipeline of SC-Track. Instance segmentation of cells from each frame is sequentially added to their respective TrackTrees. The hierarchical probabilistic cache-cascade model of SC-Track determines the assignment of each instance segmentation. If cell classification

information is encoded in the TrackTrees, SC-Track will employ the TrackTree Class Smoothing (TCS) algorithm to correct the noisy cell classifications.

A

Evaluated frames

Previous frame t = 1

Next frame t = 2

Step 1: IoU matching

Step 2: Similarity maximisation for segmented cells with multiple overlapping IoU

Similarity calculation

t = 1   t = 2

Step 3: Candidate search enlargement via bounding box expansion of unlinked cell (d)

Unlinked cell

Unlinked cell

B

Evaluated frames

Previous frame t = 1

Next frame t = 2

Resolution of unlinked cells in frame 1 and 2

Candidate mother cell identified

No candidate mother cell identified

Cell division event

New cell instance

Fig. 2: Schematic illustration summarising the hierarchical tracking approach for single-cell tracking and identifying cell division events.

(A) SC-Track employs a hierarchical cell tracking approach to minimise computational costs. The overlap between the segmented cells of the preceding and subsequent frame determines the linking of segmented cells between frames. If only one cell segmentation overlaps, the segmented cell in the subsequent frame is automatically linked to the respective TrackTree in the preceding frame. When there are multiple overlapping cell segmentations, the identification of the linked cells will be determined by the similarity value of the overlapping cell segmentations. If no overlapping candidate segmented cell was identified, the bounding box of the unlinked segmented cell will be expended to identify possible candidates. (B) When a segmented cell instance that cannot be linked to available TrackTrees is identified, SC-Track will attempt to determine if a cell division event has occurred. SC-Track will perform a search in the preceding frame to determine if a compatible candidate mother cell is available. When a compatible mother cell is detected, the segmented cell instance will be branched to the corresponding TrackTree, and a cell division event is recorded. If no compatible mother cell is identified, SC-Track will assume that the segmented cell is due to a recent appearance of a cell in the microscope field of view and a new TrackTree is initialised.

Fig. 3: Schematic illustration describing SC-Track cached matching algorithm and TrackTree Class Smoothing (TCS).

(A) Diagram describing the cache matching algorithm. When a segmented instance of a cell is detected but no compatible TrackTree is available in the previous frame, SC-Track will perform a cached search of up to five preceding frames to identify a compatible unlinked TrackTree. If a compatible TrackTree is found, the segmented instance will be linked to the TrackTree. Missing gaps in the TrackTree will be filled with the last cached cell segmentation in the preceding frame. (B) Diagram describing the TrackTree linking process. When all segmented cells are analysed, SC-Track will attempt to link TrackTrees that contain more than a single node. The linking process is based on the last segmentation of the TrackTree and the first segmentation of a separate TrackTree. When more than one compatible TrackTree is available for linkage, the cell segmentation with the highest similarity will be linked. Missing gaps in the linked TrackTrees are filled with the last cached cell segmentation in the preceding frame. (C) When a multi-class cell segmentation is performed, it is often observed that erroneous cell classifications would occur stochastically. The TCS algorithm employs a probabilistic cached search algorithm to determine if a class switch has occurred for the respective segmented cell in a time series. Note: The vertical dotted line represents a frame of a timelapse image series, and the blue circles represent an instance of a segmented cell. An established track tree is represented by a blue circle with a blue line running through the circles.

Fig. 4: Evaluation metrics of the cell tracking accuracy based on ground truth segmentations.

Box plots of (A) IDF1, (B) MOTA and (C) CDF1 scores for all four cell trackers in varying imaging time intervals. Each point displayed on the boxplots represents the respective scores of the five test datasets. The line in the boxplot represents the median. The results for Deepcell-tracking CDF1 scores were not included in (C) as the tracker failed to detect any cell division instances in all the datasets tested.

Fig. 5: Evaluation metrics of the cell tracking accuracy based on varying cell segmentation loss.

Box plots of (A) IDF1, (B) MOTA and (C) CDF1 scores for all four cell trackers challenged with the ground truth segmentation dataset with randomly deleted segmentations to mimic varying levels of cell segmentation loss. Each point displayed on the boxplots represents the respective scores of the five test datasets. The line in the boxplot represents the median. The results for Deepcell-tracking CDF1 scores were not included in (C) as the tracker failed to detect any cell division instances in all the datasets tested.

Fig. 6: Runtime evaluation comparisons.

The number of frames each tracker can process in one second is displayed on the y-axis, while the x-axis represents the varying number of processed image frames. The solid line represents the average performance, with the shaded area representing the 95% confidence interval for each cell tracker on three different computer systems running Windows, Linux, or macOS, respectively.

Table 1: IDF1, MOTA and CDF1 test results based on ground truth segmentation masks.

Dataset	SC-Track			pcnaDeep			TrackMate			Deepcell-tracking		
	IDF1	MOTA	CDF1	IDF1	MOTA	CDF1	IDF1	MOTA	CDF1	IDF1	MOTA	CDF1
RPE1-01	**0.994**	**1.000**	**0.965**	0.950	0.957	0.638	0.902	0.955	0.308	0.916	0.983	-
RPE1-02	**0.997**	**1.000**	**0.959**	0.957	0.998	0.533	0.927	0.975	0.229	0.935	0.990	-
MCF10A-01	**0.997**	**0.997**	**0.906**	0.973	0.992	0.496	0.943	0.874	0.022	0.962	0.981	-
MCF10A-02	**0.982**	**0.998**	**0.971**	**0.982**	0.978	0.657	0.945	0.885	0.000	0.952	0.967	-
RPE1-03	**0.989**	**0.994**	**0.957**	0.975	0.995	0.308	0.856	0.923	0.024	0.931	0.976	-
Average	**0.992**	**0.998**	**0.952**	0.967	0.984	0.526	0.915	0.922	0.116	0.939	0.979	-

Note: The best scores for each respective dataset and the best average score are highlighted in bold. The results for Deepcell-tracking CDF1 scores were not included as the tracker failed to detect any cell division instances in all the datasets tested.

Table 2: IDF1, MOTA and CDF1 test results based on uncorrected segmentation masks.

Dataset	SC-Track			pcnaDeep			TrackMate			Deepcell-tracking		
	IDF1	MOTA	CDF1	IDF1	MOTA	CDF1	IDF1	MOTA	CDF1	IDF1	MOTA	CDF1
RPE1-01	**0.959**	**0.996**	**0.917**	0.953	0.982	0.391	0.754	0.876	0.193	0.929	0.974	-
RPE1-02	**0.939**	**0.997**	**0.768**	0.914	0.980	0.480	0.746	0.948	0.088	0.919	0.971	-
MCF10A-01	**0.954**	**0.971**	**0.841**	0.942	0.957	0.346	0.755	0.829	0.108	0.942	0.955	-
MCF10A-02	**0.959**	**0.978**	**0.968**	0.944	0.953	0.286	0.787	0.874	0.091	0.948	0.950	-
RPE1-03	0.964	**0.998**	**0.952**	**0.967**	0.995	0.308	0.771	0.963	0.081	0.890	0.958	-
Average	**0.955**	**0.988**	**0.889**	0.944	0.973	0.362	0.763	0.898	0.112	0.926	0.962	-

Note: The best scores for each respective dataset and the best average score are highlighted in bold. The results for Deepcell-tracking CDF1 scores were not included as the tracker failed to detect any cell division instances in all the datasets tested.

Table 3: Cell Tracking Challenge imaging dataset test results for SC-Track, TrackMate and Deepcell-tracking.

Dataset	Temporal res (min)	Image	Mask	Division events	SC-Track			TrackMate			Deepcell-tracking		
					IDF1	MOTA	CDF1	IDF1	MOTA	CDF1	IDF1	MOTA	CDF1
DIC-C2DH-HeLa	10	DIC	Cell	2	**0.992**	**0.992**	-	0.568	0.639	-	0.206	-0.020	-
PhC-C2DH-U373	15	Phase	Cell	0	**1.000**	**1.000**	-	0.719	0.954	-	0.761	0.604	-
Fluo-C2DL-MSC	20	Fluor	Cell	1	**0.979**	**0.993**	-	0.445	0.685	-	0.588	0.599	-
Fluo-N2DH-GOWT1	5	Fluor	Nuclear	1	**0.991**	**0.999**	-	0.971	0.948	-	0.902	0.855	-
Fluo-N2DH-SIM+	29	Fluor	Nuclear	16	**0.963**	**0.991**	**0.896**	0.690	0.634	0.151	0.747	0.925	-
PhC-C2DL-PSC	10	Phase	Cell	44	**0.989**	**1.000**	**0.977**	0.903	0.945	0.301	0.879	0.991	-
Fluo-N2DL-HeLa	30	Fluor	Nuclear	56	**0.992**	**0.998**	**0.991**	0.881	0.855	0.061	0.890	0.968	-
BF-C2DL-MuSC	5	BF	Cell	5	**1.000**	**1.000**	-	0.143	-0.348	-	0.540	0.096	-
				Average	**0.988**	**0.997**	**0.955**	0.665	0.664	0.171	0.689	0.627	-

Note: We were unable to evaluate pcnaDeep's cell tracking performance on the CTC dataset because pcnaDeep requires cell cycle data encoded in the cell segmentations to generate single cell tracks. The results for Deepcell-tracking CDF1 scores were not included as the tracker failed to detect any cell division instances in all the datasets tested. Only timelapse movies containing more than 10 cell division events are included in the analysis of CDF1 scores to ensure a reliable measure of cell division tracking performance.

Table 4: F1 cell cycle classification test results obtained from raw StarDist cell classification predictions compared with TCS corrected cell classifications.

Dataset	Raw F1 values			TCS corrected F1 values		
	G1/G2	S	M	G1/G2	S	M
RPE1-01	0.956	0.775	0.659	**0.995**	**0.981**	**0.867**
RPE1-02	0.949	0.809	0.748	**0.982**	**0.922**	0.691
MCF10A-01	0.931	0.690	0.469	**0.982**	**0.923**	**0.580**
MCF10A-02	0.881	0.660	0.479	**0.963**	**0.848**	**0.768**
RPE1-03	0.965	0.795	0.236	**0.999**	**0.969**	**0.719**
Average	0.936	0.746	0.518	**0.984**	**0.929**	**0.725**

Note: TCS corrected F1 values that show an improvement over the Raw F1 values are highlighted in each category.

Supplementary Table 1: Development dataset used in the development and finetuning of SC-Track tracking parameters.

Dataset	Cell Type	Number of image frames	Total Cells	Average cells per frame	Temporal resolution (minutes)
sub-rpe-1	RPE1	100	5374	53.74	5
sub-rpe-2	RPE1	100	6056	60.56	5
sub-rpe-3	RPE1	100	7296	72.96	5
sub-rpe-4	RPE1	69	5673	82.22	5

Supplementary Table 2: SC-Track evaluation dataset summary.

Dataset	Original filename	Cell Type	Number of image frames	Average cells per frame	Temporal resolution (minutes)	G1/G2	S	M	Cell division events
RPE1-01	copy_of_1_xy01	RPE1	369	63.41	5	20760	2082	552	44
RPE1-02	copy_of_1_xy19	RPE1	369	66.16	5	22624	1202	567	32
MCF10A-01	MCF10A_copy02	MCF10A	101	119.89	5	11147	541	345	29
MCF10A-02	MCF10A_copy11	MCF10A	101	70.75	5	6236	572	211	17
RPE1-03	src06	RPE1	577	34.50	5	18459	1395	114	11

Supplementary Table 3: SC-Track evaluation dataset with uncorrected raw segmentations and classifications.

Dataset	Detected instances			Cell classifications			G1/G2 phase classification			S phase classification			Mitosis phase classification		
	TP	FP	FN	TP	FP	FN	TP	FP	FN	TP	FP	FN	TP	FP	FN
RPE1-01	23121	245	279	21742	1624	1658	19474	493	1321	1935	952	172	333	179	165
RPE1-02	24031	499	382	22088	2442	2325	19145	1495	1190	2342	641	711	601	306	424
MCF10A-01	11418	324	691	11088	654	1021	10008	418	756	562	35	124	518	201	141
MCF10A-02	6812	120	334	5549	1383	1597	4818	414	1199	588	782	125	143	187	273
RPE1-03	19651	158	255	18724	1085	1182	17392	366	916	1282	455	206	50	264	60

Supplementary Table 4: SC-Track synthetic evaluation dataset with randomly deleted segmented images.

Dataset	Detection loss ratio (%)	Average cells per frame	Detected cells per frame	G1/G2 phase classification			S phase classification			M phase classification		
				TP	FP	FN	TP	FP	FN	TP	FP	FN
RPE1-01	5	63.41463	59.6802	19569	0	1226	1978	0	129	475	0	23
	10	63.41463	56.523	18535	0	2260	1873	0	234	449	0	49
	20	63.41463	50.3008	16491	0	4304	1680	0	427	390	0	108
	30	63.41463	43.9756	14424	0	6371	1458	0	649	345	0	153
	50	63.41463	31.4607	10317	0	10478	1043	0	1064	249	0	249
RPE1-02	5	66.15989	62.4526	19167	0	1168	2905	0	148	972	0	53
	10	66.15989	59.1247	18176	0	2159	2734	0	319	906	0	119
	20	66.15989	52.5474	16180	0	4155	2412	0	641	797	0	228
	30	66.15989	45.8753	14096	0	6239	2126	0	927	705	0	320
	50	66.15989	32.8482	10099	0	10236	1557	0	1496	464	0	561
MCF10A-01	5	119.8911	113.4455	10187	0	577	648	0	38	623	0	36
	10	119.8911	107.4356	9644	0	1120	611	0	75	596	0	63
	20	119.8911	95.5743	8574	0	2190	559	0	127	520	0	139
	30	119.8911	83.4554	7487	0	3277	471	0	215	471	0	188
	50	119.8911	59.6535	5355	0	5409	342	0	344	328	0	331
MCF10A-02	5	70.75248	66.6139	5662	0	355	676	0	37	390	0	26
	10	70.75248	63.1881	5370	0	647	635	0	78	377	0	39
	20	70.75248	56.198	4754	0	1263	572	0	141	350	0	66
	30	70.7525	49.0495	4169	0	1848	491	0	222	294	0	122
	50	70.7525	35.1584	2974	0	3043	378	0	335	199	0	217
RPE1-03	5	34.4991	32.4974	17243	0	1065	1406	0	82	102	0	8
	10	34.4991	30.5546	16232	0	2076	1304	0	184	94	0	16
	20	34.4991	27.2704	14503	0	3805	1146	0	342	86	0	24
	30	34.4991	23.6118	12536	0	5772	1017	0	471	71	0	39
	50	34.4991	16.9636	9005	0	9303	731	0	757	52	0	58

Supplementary Table 5: Cell Tracking Challenge Silver Reference datasets.

Dataset	Cell Type	Number of frames	Image Size (pixel, width x height)	Total Cells	Average cells per frame	Cell division events	Temporal resolution (minutes)	Image Type	Mask Type	Microscope	Microscope Objective
DIC-C2DH-HeLa	HeLa	24	512x512	279	11.63	2	10	DIC	Cell Mask	Zeiss LSM 510 Meta	Plan-Apochromat 63x/1.4 (oil)
PhC-C2DH-U373	Glioblastoma-astrocytoma U373	115	696x520	880	7.65	0	15	Phase Contrast	Cell Mask	Nikon	Plan Fluor DLL 20x/0.5
Fluo-C2DL-MSC	Rat mesenchymal stem cell	48	992x832	475	9.90	1	20	Fluorescence	Cell Mask	PerkinElmer UltraVIEW ERS	Plan-Neofluar 10x/0.3 (Plan-Apo 20x/0.75)
Fluo-N2DH-GOWT1	GFP-GOWT1 mouse stem cell	92	1024x1024	2146	23.33	1	5	Fluorescence	Nuclear Mask	Leica TCS SP5	Plan-Apochromat 63x/1.4 (oil)
Fluo-N2DH-SIM+	Simulated nuclei of HL60 cell	100	773x739	1642	16.42	16	29	Fluorescence	Nuclear Mask	Zeiss Axiovert 100S with a Micromax 1300-YHS camera	Plan-Apochromat 40x/1.3 (oil)
PhC-C2DL-PSC	Pancreatic stem cell	100	576x720	9208	92.08	44	10	Phase Contrast	Cell Mask	Olympus ix-81	UPLFLN 4XPH
Fluo-N2DL-HeLa	HeLa	92	700x110	6894	94.50	0	30	Fluorescence	Nuclear Mask	Olympus IX81	Plan 10x/0.4
BF-C2DL-MuSC	Mouse muscle stem cell	1100	1036x1070	2594	2.36	5	5	Bright Field	Cell Mask	Zeiss PALM/ AxioObserver Z1	EC Plan-Neofluar 10x/0.30 Ph1