The archerfish uses motor adaptation in shooting to correct for changing physical conditions

Svetlana Volotsky[1, 2,3], Opher Donchin[1,2], Ronen Segev[1, 2, 3]

[1] Department of Biomedical Engineering, Ben-Gurion University of the Negev

[2] School of Brain Sciences and Cognition, Ben-Gurion University of the Negev

[3] Department of Life Sciences, Ben-Gurion University of the Negev

Abstract

The archerfish is unique in its ability to hunt by shooting a jet of water from its mouth that

hits insects situated above the water's surface. To aim accurately, the fish needs to overcome

physical factors including changes in light refraction at the air-water interface. Nevertheless,

archerfish can still hit the target with a high success rate under changing conditions. One

possible explanation for this extraordinary ability is that it is learned by trial and error

through a motor adaptation process. We tested this possibility by characterizing the ability

of the archerfish to adapt to perturbations in the environment to make appropriate

adjustments to its shots. We introduced a perturbing airflow above the water tank of

archerfish trained to shoot at a target. For each trial shot, we measured the error; i.e., the

distance between the center of the target and the center of the water jet produced by the fish.

Immediately after the airflow perturbation, there was an increase in shot error. Then, over

the course of several trials, the error was reduced and eventually plateaued. After the

removal of the perturbation, there was an aftereffect, where the error was in the opposite

direction but washed out after several trials. These results indicate that archerfish can adapt

to the airflow perturbation. Testing the fish with two opposite airflow directions indicated

that adaptation took place within an egocentric frame of reference. These results thus suggest

that the archerfish is capable of motor adaptation, as indicated by data showing that the fish

produced motor commands that anticipated the perturbation.

Introduction

In a 1764 letter to the Royal Society of London, John Schlosser reported his discovery of a unique species (Schlosser. 1764). In this letter, Schlosser describes an amazing fish from Southeast Asia that can hunt insects above the water level by shooting a squirt of water from its mouth without ever missing. The archerfish, as it is called today, may not be as accurate as claimed (in fact, it misses quite often), but its behavior is nevertheless considered one of the most remarkable hunting strategies in nature. Of the many intriguing questions it raises, one remains conspicuously unresolved: how does the archerfish achieve accurate shooting despite various physical factors that affect the water jet?

This question has been asked with regard to many types of motor behavior: how do animals make accurate saccades (Shadmehr et al. 2010,Wallman and Fuchs. 1998)? how do humans and other primates make accurate reaching movements (Scheidt et al. 2005,Kluzik et al. 2008)? In many of these cases, mechanisms of adaptation that respond to error and success manage aiming and calibration of the motor system (Shadmehr et al. 2010,Donchin et al. 2012). However, the question of maintenance of accurate motor behavior in fish has rarely been asked (Tsvilling et al. 2012,Rossel et al. 2002).

This study tests whether accurate shooting in the archerfish manifests characteristics that suggest underlying adaptation mechanisms similar to those seen in other systems. That is, first, can we show gradual change in the shot direction that reduces error in response to a perturbation? Second, will there be an aftereffect after the perturbation is removed?

During a shot, the fish's mouth is in the air, while its eyes remain well below the water level. Thus, in order to hit the target, the fish needs to compensate for multiple physical factors that affect the

Figure 1 Volotsky et al.

visible target

θ_1

Wind

actual target

Light refraction

$$\frac{\sin \theta_2}{\sin \theta_1} = 1.33$$

θ_2

water jet

θ_1 θ_2

Figure 1| The archerfish needs to correct for physical factors in shooting. The viewing angle of a target above the water level is shifted towards the zenith due to Snell's law. The fish needs to correct for this shift to hit the target in its actual position. The jet itself is affected by gravity and it does not proceed straight from the fish's mouth to the target. In addition, wind can affect the trajectory of the shot.

shot. This includes light refraction at the air-water interface, wind, and angle (Figure 1). Even the

altitude of the target affects the shot due to gravitation (Rossel et al. 2002,Timmermans and

Vossen. 2000a).

The mechanism behind this ability to compensate for the different physical factors is unclear (Dill.

1977,Temple. 2007). Possible hypotheses as to the mechanism the fish uses for this correction is

based on positing that the compensation is pre-programmed and hardwired through innate models

of the physics. While this explanation is conceptually simple, it is hard to believe that it is

sufficient, since during growth of the fish, the physical characteristics of its motor and visual

systems change continuously, which would require a prewired program to take an unlikely number

of highly interacting variables into account.

An opposing hypothesis posits that the archerfish does not have any innate physical knowledge

and instead has a neural mechanism that develops an internal model of the current conditions

through experience (Shadmehr et al. 2010). In this case, accurate shooting would depend on

ongoing adaptation to consistent errors in a manner similar to the adaptation mechanisms that have

been extensively investigated in the study of other vertebrate motor systems (Martin et al.

1996a,Fernandez-Ruiz and Diaz. 1999a,Donchin et al. 2012). In this way, through a series of trials

and errors, the fish would correct for misses while trying to hit the target. One advantage of an

adaptation mechanism is that it can overcome the parameter differences caused by the physical

changes that occur throughout the fish's life cycle.

These two approaches may be complementary. Innate models rely on underlying neural circuits,

and these neural circuits probably have mechanisms of adaptation and calibration. Thus, the

question of how the archerfish corrects for refraction and other physical factors can in fact be

reformulated as 'to what extent is this ability innate, and how is it adapted?'

Indeed, the process of compensation itself can reveal aspects of the fish's innate internal models.

For instance, if compensation generalizes in an egocentric reference frame, this implies that the

fish associates errors with factors connected to the body of the fish such as muscle fatigue or

physical factors that do not change with reference frame such as refraction index. If the

compensation generalizes allocentrically, this implies that the fish associates errors with factors

connected to the environment reference frame such as wind.

One of the key concepts in investigations of the associations between vison and motor output is

visuomotor adaptation (Martin et al. 1996a,Fernandez-Ruiz and Diaz. 1999b), which is a broad

term for any practice-related change or improvement in motor performance in response to a

perturbation in a task based on visual input. Visuomotor adaptation returns behavior to baseline

levels of performance usually within tens or a few hundred trials, depending on the particulars of

the task. A classic example of visuomotor adaptation is prism adaptation (Kitazawa et al.

1995,Martin et al. 1996b): a sensory-motor adaptation that occurs after the visual field has been

artificially shifted laterally or vertically by a prism. During prism adaptation studies, individuals

wear special prismatic goggles made of prism wedges. The individuals then engage in a

perceptual motor task such as pointing to a visual target. After performing the task repeatedly,

the individuals improve over time until their performance is comparable to performance before

the prism goggles were worn. Visuomotor adaptation has a few defining characteristics that differentiate it from other mechanisms of error correction; namely, it is gradual, error-driven, and shows an "aftereffect": a tendency for the corrected behavior to persist for some time after the perturbation has been removed (Fernandez-Ruiz and Diaz. 1999a).

Here, we explored visuomotor adaptation in the archerfish, based on the extensive literature studying visuomotor adaptation in humans (Izawa and Shadmehr. 2011,Shadmehr and Mussa-Ivaldi. 1994a,Krakauer and Mazzoni. 2011,Fernandez-Ruiz and Diaz. 1999a). We examined whether the archerfish can adapt to perturbations in the environment which result in distortion to the shot by generating an airflow above the water level to mimic the perturbation to the properties of air-water refraction. We show that the archerfish is indeed capable of visuomotor adaptation, which is consistent with the hypothesis that this adaptation partially contributes to the ability of the fish to overcome the changing conditions that affects the shot.

Method

Animals. Wild-caught adult fish (6-14 cm in length; 10-18 gr) were purchased from a local supplier. The fish were kept separately in 100-liter aquaria filled with brackish water (salinity 6-8 ppt) at 25-29^{0}C on a 12/12 hour light-dark cycle. Fish care and experimental procedures were approved by the Ben-Gurion University of the Negev Institutional Animal Care and Use Committee and were in accordance with government regulations of the State of Israel.

Experimental scenery. The experiments were conducted in a water tank with a cover above the water that forced the fish to shoot through a slit in the cover such that the shot was aligned with the airflow perturbation when present (Figure 2A,B).

After delivery to the lab, the fish went through a period of acclimatization to the lab environment.

Then, the fish were gradually trained to shoot at a black circle on a metal net placed 35-40 cm

above the water, depending on the water level in the tank. After the fish were able to make 20-25

shots in 10 minutes, a plastic cover with a slot was placed above the water tank. The fish were

trained to shoot at a target above the water tank to ensure uniform direction of the shot in relation

to the direction of the later perturbation. The target was placed at the distance of 10 cm from the

air nozzle. The speed of the airflow above the water was measured using an anemometer (SP-

82AM, Lutron Electronics, Taiwan) just below the target. The air speed was 7.5 m/s ±0.3 m/s.

Training

Overall, 39 candidate fish were tested for the study. We first screened for the ability to persist

shooting at least 20 shots per session through a slot in a cover above the water tank and remain

accurate throughout the session. This was crucial since it enabled the collection of enough data for

statistical analysis. Overall, seven fish passed the screening test and continued to the perturbation

experiments.

Two of the seven fish were able to complete the 15 sessions. One fish completed 8 sessions, two

fish - 7 sessions, one fish - 2 sessions and one fish - 1 session in the first experiment. For the second

experiment, the fish were required to perform at least 30 consecutive shots through a slot with a

change in direction in the middle of the session. Only three fish met this criterion and were used

in the experiment: one fish competed 15 sessions, one fish – 4 sessions and one fish – 8 sessions.

Experiment 1. The experiment consisted of three stages (Figure 2C). First, the fish performed 5

to 10 trials with no air flow. The number of trials under baseline conditions was generated from a

uniform distribution so that the fish would not anticipate the change in conditions after a fixed

number of trials. Then, we introduced a perturbing airflow horizontally to the water's surface that

deflected the fish's shot trajectory and caused the fish to miss the target. The airflow was either in

the direction of the shot or opposite to the direction of the shot. The fish performed 10 to 15 trials.

Then the air flow was removed and the fish shot again under the baseline condition. Thus, the

errors in the first and final stage permitted comparisons of fish behavior in normal and post-

adaptation conditions in the same physical environment.

Experiment 2. We tested whether the fish adapt to the airflow perturbation within an egocentric

or allocentric frame of reference. In the egocentric frame of reference, any perturbation is

perceived as constant regardless of the subject position; for example, refraction index on the water

surface. In the allocentric frame of reference, the perturbation perception changes with the change

in position; an example for this scenario is a wind above the water surface that affects the water

shot differently depending on the fish position relative to the wind direction The second

experiment consisted of four stages: first, the fish performed five to 10 trials without the airflow

to estimate the baseline; then, 10 trials with the airflow in one direction were administered (Figure

2D). Next, the fish were forced to reverse the direction of their shot by rotating the cover with the

slot by 180 degrees with no change in the airflow direction. The airflow was turned off after 15 to

20 trials and the fish performed five additional shots with no perturbation. Throughout the

experiments, the fish were given no clues about the initiation or termination of the airflow, nor

about its direction.

Video recording. The experiments were recorded using 3 high-resolution cameras (ISG

LightWise Allegro, Imaging Solutions, US) at a frame rate of 190 frames per second. One camera

was focused on the target and its immediate surrounding area. Two other cameras recorded the

fish from different angles to capture their behavior. The video clips were analyzed offline.

Figure 2 Volotsky et al.

A

B

Air flow

C

| Baseline no air flow | Perturbation air flow | Washout no air flow |

Trial number

D

| Baseline no air flow | Perturbation air flow direction I | Reverse Perturbation air flow direction II | Washout no air flow |

Trial number

Figure 2| Schematic view of the experimental setup. A. Water tank with a cover and a target above it. Airflow applied horizontally to the water's surface deflects the fish's shot. **B.** An example from a video capturing the experiment. The airflow was oriented from right to left. The water jet is visible just before the impact at the target. **C.** Experimental timeline of the first experiment: 5 to 10 shots before the introduction of the airflow, 10 to 15 shots with the airflow, 5 to 10 shots after the removal of the airflow. **D.** Experimental timeline of the second experiment: 5 to 10 shots before the introduction of the airflow, 8 to 12 shots with the airflow in one direction, 15 to 20 shots with the airflow in the opposite direction, 5 to 10 shots after the removal of the airflow.

Measuring shot error. We characterized the adaptation process in terms of the shooting error

measured on each trial shot. The error was defined as the distance between the center of a target

and the center of the water jet produced by the fish. The data were extracted from the movie frames

of the camera that was focused on the target and its immediate surroundings.

Statistical analysis.

We used a Bayesian approach to analyze the adaptation process. To analyze the results of the first

experiment, we defined 6 epochs in the timeline of the experiment: 1 – baseline trials; 2 – first two

trials after the initiation of the air flow perturbation; 3 – trial from the third up to the two trials

preceding the end of the perturbation; 4 -the last two trials before the termination of the

perturbation; 5 – the first two trials after the termination of the perturbation; 6 – the remainder of

the trials until the end of the session.

To analyze the results of the second experiment we defined 9 epochs in the timeline of the

experiment: 1 – baseline trials; 2 – first two trials after the initiation of the air flow perturbation; 3

– trial from the third up to the two trials before the direction changed; 4 – the last two trials before

the direction changed; 5 – first two trials after the direction changed; 6 - trial from the third up to

the two trials preceding termination of the perturbation; 7 – last two trials before the termination

of the perturbation; 8 – first two trials after the termination of the perturbation; 9 – the remainder

of the trials until the end of the session.

We performed a hierarchical Bayesian analysis to evaluate the behavior of the fish in response to

the perturbation. The statistical analysis was conducted using R 4.0.4 programming language

(https://www.r-project.org/) and the JAGS 4.3.0 statistical package (https://mcmc-

jags.sourceforge.io/) (Plummer. 2003). JAGS was used to produce samples from the posterior

probability distribution based on the data for the parameters of the statistical model described

below (Kruschke. 2014).

The central parameter of interest was the shooting error of each fish in each time epoch. In our

model, this error was drawn from a normal distribution. The mean of this distribution was modeled

hierarchically as a linear combination of parameters that depended on the trial epoch, the subject,

and the interaction between them. That is, the model of the error was written in the following way:

$$error_{trial} \sim Normal(a_{baseline} + a_{epoch} + a_{subject} + a_{epoch\&subject}, \sigma^2) \quad (1)$$

179   where $a_{baseline}$ is the average all baseline errors, a_{epoch} is the average contribution to the error

180   during a specific epoch for all fish, $a_{subject}$ is the fish's unique characteristics and $a_{epoch\&subject}$

181   is contribution to the error due to the interaction between subject and epoch. Finally, σ^2 is the

182   global variance. The model was hierarchical in the sense that the prior distributions of these

183   parameters were themselves modeled using parametric distributions as follows:

$$a_{baseline} \sim N(\mu_{all\ trials}, \sigma^2_{all\ trials})$$

$$a_{epoch} \sim N(\mu_{epoch}, \sigma^2_{epoch})$$

$$a_{subject} \sim N(\mu_{subject}, \sigma^2_{subject}) \tag{2}$$

$$a_{epoch\&subject} \sim N(\mu_{epoch\&subject}, \sigma^2_{epoch\&subject})$$

185 The posterior distributions of the hyper-parameters ($\mu_\bullet, \sigma^2_\bullet$) were determined simultaneously and

186 jointly with the posterior distribution of the parameters in the linear model. The priors of the hyper-

187 means were broad normal distributions and the priors of the hyper-variances were broad gamma

188 distributions. The prior for the global variance parameter in the linear model was a broad uniform

189 distribution.

190 Bayesian data analysis combines the data and the model to generate samples of the posterior

191 distribution of the parameters given the data. JAGS carries out this sampling using a version of the

192 Markov Chain Monte Carlo (MCMC) algorithm for sampling arbitrary target distributions. We

193 generated 3 chains of 10,000 MCMC samples from the joint posterior probability distribution of

194 all the parameters. The standard procedure is to use 3 or 4 chains to show that they converge to a

195 similar result to verify the robustness of the outcome. Convergence of the algorithm and sampling

196 properties were tested using both graphical and quantitative methods (Kruschke. 2014).

197 ***Testing for significance.*** We used the 10,000 MCMC samples to calculate the 95% Highest

198 Density Interval (HDI). HDI is the range of values for which there is a 95% posterior probability

199 of finding the parameter and where all the values within the interval are more probable than any

200 value outside of it. By examining the 95% HDI of each fish in each time epoch, we could describe

201 the different stages of the adaptation process in the archerfish. If the 95% HDI of the difference in

202 the error between two epochs was greater than the region of practical equivalence (ROPE) of 5%

around zero, the error was considered to be significantly different. For every comparison we report

the HDI of the difference between the epochs in mm.

Determination of effect size. Effect size was calculated for the difference between transition stages

of the experiments: for experiment 1 – for the differences in error between epochs 1 and 2, 2 and

4, 4 and 5. For the second experiment, we compared error between epochs 1 and 2, 2 and 4, 4 and

5, 5 and 7, 7 and 8. For every MCMC sample, Cohen's d effect size was calculated as the difference

between the mean values of the two compared instances divided by the pooled standard deviation.

Then, 95% HDI was calculated for the values of effect size. Effect sizes are reported for all

comparisons as the HDI of Cohen's d.

Results

To test the capability of the archerfish for motor adaptation, we examined its ability to adjust to a

consistent perturbation generated by airflow above the water level (Figure 2A). First, we trained

the fish to shoot at a food pellet placed on a metal net. Then, an airflow was directed such that it

provided a backwind to the jet produced by the fish and thus generated an error in the shot (which

we defined as a positive error). The direction of the error in the shot was measured manually from

images taken with a high-resolution video camera (see Methods).

Archerfish exhibit motor adaptation

In the first experiment the fish had to shoot at the target so that we could obtain a baseline for the

accuracy of its shooting. Then, the airflow perturbation was turned on and the fish had to

compensate using motor adaptation to the deflection of the jet (Figure 2A, see Methods). Finally,

there was a washout period where the airflow was turned off. The error for each shot was measured.

Figure 3 Volotsky et al.

A   Shot with the airflow

B   Shot against the airflow

C   One example session of 3 fish

D   One example session of 4 fish

E   All sessions of one example fish

F   All sessions of one example fish

Figure 3| Examples of fish responses to the perturbation. A. Experiment 1 setup: airflow in direction 1 - with the fish shot. **B.** Experiment 1 setup: airflow in direction 2 - against the fish shot. **C.** Example sessions for three fish that had to adapt to the perturbation in direction 1. Error was around zero during the baseline condition, increased with the introduction of the perturbation, and diminished with time. After the removal of the perturbation, the error was in the opposite direction. **D.** Example sessions for the fish that had to adapt to the perturbation in direction 2. **E.** All sessions for one example fish with the perturbation in direction 1. **F.** All sessions of one example fish with the perturbation in direction 2.

Figure 3C, D presents the errors for seven different fish in representative sessions - three with the

direction and four – against the direction of the airflow, and all sessions for two fish in Figure 3E,F

– one in each direction.

For further analysis, we used all sessions from the seven fish. We analyzed individual fish

performance and the average performance of all fish (Figure 4). As expected, when the

perturbation was not present, the errors were distributed around zero. This is indicated in the raw

traces (Figure 4A, B) and in the average responses of the individual fish (Figure 4C, D) as well as

the average response of all fish (Figure 4C, D).

After the airflow was turned on, the fish adapted to this perturbation. Immediately after the

introduction of the airflow perturbation, there was a significant increase in shot errors (Figure 4,

Perturbation, difference between epochs HDI 4.28-5.52 mm, Cohen's d HDI 1.23-1.58). Then,

over the course of several trials, the error became smaller and eventually plateaued. For all fish,

the error at the beginning differed significantly from the error at the end of the perturbation period

(difference between epochs HDI 3.1-4.33 mm, Cohen's d HDI 0.86-1.25).

After the airflow perturbation stopped (Figure 4, washout), the fish exhibited an aftereffect (Figure

4, Washout, difference between epochs HDI 2.65-3.99 mm, Cohen's d HDI 0.76-1.15). On the

first trials after the airflow perturbation was removed, there was an error in the opposite direction

Figure 4| The archerfish can correct for perturbation using motor adaptation. A. Response of three fish to the perturbation. Direction of the airflow is with the fish shots. Black dots are the error values across the sessions, the black line is the mean error value, and the SE is highlighted in green. **B.** Response of three fish to the perturbation. Direction of the airflow is against the fish shots. **C.** For three fish that shoot in direction 1: Mean value of the error and 95% HDI for epoch 1 – the baseline, epoch 2 – first two trials after the introduction of the airflow, epoch 4 - last two trials before the termination of the perturbation, epoch 5 – first two trials after the termination of the perturbation. **D.** For the four fish that shot in direction 2: Mean value of the error and 95% HDI.

that also decayed after several trials. This suggests that the fish generated motor commands that

anticipated the perturbing airflow. The results were consistent for all seven fish in the experiment

(Figure 4).

In our examination of the data, we sought out evidence of session-to-session 'saving' effect.

However, the inherent noise within the collected data prevented us from drawing any definite

conclusions.

Motor adaptation is situated in fish egocentric frame of reference

Next, we assessed whether archerfish motor adaptation is based on an egocentric or allocentric

frame of reference. The experiment was designed to let the fish adapt to airflow in one direction

and then force the fish to switch direction by rotating the cover with the slot above the water tank,

then shoot with the airflow in the opposite direction. There are no cues about the presence or

direction of the airflow perturbation above the water tank; the activation of the air blower is done

outside of the fish vision field. But the fish can see the visual landmarks inside the water tank, such

as filter, plant décor and thermostat. And they demonstrate a behavioral indication of position and

direction within the tank in their shooting behavior (Ben-Tov et al. 2018,Tsvilling et al.

2012,Dewenter et al. 2017); when the cover with the slot above the water tank is rotated, the fish

change their body position to shoot through the slot from the new angle.

This experiment was motivated by the fact that in our experimental setup, we introduced a

perturbation, thus creating an environmental condition that remained constant while the change in

direction relative to the perturbation was done by the fish. We hypothesized two outcome scenarios

in response to the change. In the first, the fish would perceive the shift in allocentric reference

frame: the change occurring in its own position and the airflow perturbation remaining constant.

The fish could use this information to correct for the true direction of the airflow and produce a

smaller error.

In the second scenario, the fish would make a motor adaptation based on an egocentric reference

frame: the environment changes together with the body position. In this case, the correction after

the switch would present a large error since the correction for the previous perturbation would be

added to the effect of the airflow. Motor adaptation in an egocentric reference frame is consistent

with using the process to overcome refraction or other body-linked physical factors since they will

be identical in both directions. Other physical factors, such as wind, would not be compensated

correctly in an egocentric reference frame.

We put this hypothesis to the test and forced the fish to reverse the direction of their first shot while

the direction of an airflow remained constant (Figure 5A). As in the first experiment, initially the

response to the perturbation was a decrease in the error such that the fish hit the target towards the

end of the trials (Figure 5B-E, Perturbation, difference between epochs HDI 2.47-5.05 mm,

Cohen's d HDI 0.45-0.71). Then, immediately after the change in direction, the magnitude of the

error increased considerably (difference between epochs HDI 14.32-16.89 mm, Cohen's d HDI

2.02-2.38), and adaptation took more trials than the initial response (Figure 5B-E, Reverse

Perturbation). Thus, in the absence of clues about the airflow direction, the fish apparently did not

perceive the airflow perturbation as constant. After the switch, the fish continued to correct for the

perturbation in the original direction relative to their body position.

We reanalyzed the results of the experiments by building mixed-effects models, with fixed-effects

factor for the epoch and random-effects grouping factor for the fish. In ANOVA test for the

models, the effect of the epoch factor was found to be significant and large (Experiment 1:

$p<0.001$, $\eta^2 = 0.67$; Experiment 2: $p<0.001$, $\eta^2 = 0.31$). We performed post-hoc analysis of the

models to compare errors in different epochs of the experiments. The significance was evaluated

using t-test with Bonferroni adjustment to correct for multiple comparisons. Cohen's d was used

for effect size evaluation. In both experiments, all transitions between the stages of the experiments

affected the error (Experiment 1: epochs 1-2 – $p = 0.001$, Cohen's d $= 1.46$; epochs 2-4 – $p = 0.003$,

Cohen's d $= 1.04$; epochs 4-5 – $p = 0.002$, Cohen's d $= 1.74$. Experiment 2: epochs 1-2 – $p = 0.005$,

Cohen's d $= 1.55$; epochs 2-4 – $p = 0.036$, Cohen's d $= 0.93$; epochs 4-5 – $p = 0.0004$, Cohen's d

$= 4.12$; epochs 5-7 – $p = 0.003$, Cohen's d $= 2.21$; epochs 7-8 – $p = 0.007$, Cohen's d $= 1.95$).

Thus, we concluded that the fish engaged in motor adaptation of the shot in an egocentric reference

frame. The egocentric adaptation process indicates that the fish did not use information about the

unchanged nature of the perturbation; in this case airflow. Rather, the fish motor adaptation was

consistent with adaptation to the air-water interface, which is isotropic in nature; i.e. it does not

change with the shooting direction of the fish.

A

Fish switch direction

All sessions

B

Fish 3

C

Fish 6

D

Fish 7

Trial

E

Figure 5| Adaptation in two directions sequentially reveals that motor adaptation is performed in the fish's egocentric reference frame. A. Experimental setup: Airflow applied horizontally to the water's surface deflects the fish's shot. The fish first adapted to the perturbation in one direction and then enforced the switch direction of the shoot. **B.** For the fish that changed direction from shooting in the direction with the perturbation to against the perturbation: mean error value and SE for the baseline trials, trials at the beginning and at the end of the adaptation period, trials at the beginning and the end of the adaptation period in the opposite direction and the beginning and the end of the washout period. **C-D.** For the fish that changed direction from shooting in the direction against the perturbation to shooting with the perturbation: mean error value and SE for the baseline trials, trials at the beginning and the end of the adaptation period, trials at the beginning and the end of the adaptation period with reversed direction and at the beginning and the end of the washout period. **E.** Mean value of the error and 95% HDI for epoch 1 – the baseline, epoch 2 – first two trials after the introduction of the airflow, epoch 4 - last two trials before the change in the direction, epoch 5 – first two trials after the direction change, epoch 7 – last two trials before the termination of the perturbation, epoch 8 – first two trials after the termination of the perturbation.

Discussion

We tested the existence of the motor adaptation process in archerfish as a possible explanation for

its ability to overcome changes in the environment during shooting. We introduced a perturbing

airflow above the water tank on a fish trained to shoot at a target. The airflow deflected the

trajectory of the water jet and as a result the fish's water jet missed the target initially. Over the

course of several trials, all the fish managed to modify their shooting direction and successfully

hit the target, demonstrating their ability to adapt to changes in the environment. After the removal

of the perturbation, we observed an aftereffect, where the fish made errors in the opposite direction.

This result suggests the formation of internal model in the fish brain that anticipates the presence

of the perturbation.

We also tested whether the distortion by the airflow perturbation was consistent with egocentric

or allocentric adaptation. For instance, the distortion expected by light refraction at the interface

would require egocentric adaptation where wind would require allocentric adaptation. Thus, in the

second experiment, we kept the airflow constant while the fish was forced to rotate its direction of

shooting. We reasoned that if the fish perceived the perturbation as constant, they would correct

for the opposite direction after the position change, and we would see small errors and quick

improvement. Alternatively, if the fish adapted to egocentric factors, it would not be affected by

the direction of the view. Thus, there would be no need to make additional corrections after the

initial adaptation. Hence, immediately after the directional switch, the fish should not change its

shooting strategy and we should observe large error. As shown in Figure 5B-E, after the change in

direction, there was a pronounced increase in the error, and the decrease was slower than in the

initial adaptation stage. This response to the change in the environment was consistent with the

response to a change in the refraction index or other egocentric factors.

It is important to note that although it has been demonstrated that other fish species possess

orientation and spatial memory (Rodriguez et al. 1994,Cohen et al. 2023,Givon et al. 2022,Lee et

al. 2012,Givon et al. 2023,Vinepinsky et al. 2020), there is a lack of such information regarding

archerfish specifically. The knowledge about other fish provides supportive evidence that

archerfish may be sensitive to their relative positions in the water tank. However, it is essential to

keep in mind that the current results do not completely rule out the possibility that archerfish are

unaware of changes in their body position. Instead, they may continue with previously successful

actions, which could appear as an egocentric generalization.

The two most important studies of archerfish ability to compensate for the distortion to vision at

the air-water interface have focused on the phenomenology of compensation. Specifically, it was

shown both experimentally (Dill. 1977,Timmermans and Vossen. 2000b) and theoretically (Barta

and Horváth. 2003) that the archerfish needs to compensate for the large distortion in the apparent

elevation. Clearly, additional insights into how the archerfish aims at the target can be obtained by

observing fish when they hatch and start shooting. To the best of our knowledge, there have been

no reports of breeding archerfish in the lab or observation of the emergence of shooting behavior

in the wild.

Our findings are consistent with previous work in the archerfish which evaluated the adaptability

of the shoot behavior. These studies showed that the archerfish can predict the future position of

targets (Ben-Simon et al. 2012), the future landing position of prey (Rossel et al. 2002,Tsvilling et

al. 2012), and adjust their shot force during underwater behavior (Dewenter et al. 2017). Our

findings extend these results to changes in the physical environment. Since most studies have used

the archerfish visual system as a model for visual processing in general (Newport et al.

2014,Newport et al. 2015,Newport et al. 2018,Ben-Simon et al. 2012,Mokeichev et al.

2010,Vasserman et al. 2010,Timmermans and Vossen. 2000b,Temple et al. 2013,Volotsky et al.

2019,Volotsky et al. 2022,Segev et al. 2007,Ben-Tov et al. 2018,Ben-Tov et al. 2015,Gabay et al.

2013), the current study contributes to a better understanding of a unique trait of the archerfish:

the shooting mechanism and its control.

The need to correct for physical factors such as refraction at the interface is not unique to

archerfish. For example, dwarf gouramis can shoot at targets above water and need to correct for

refraction in the air-water interface (Jones et al. 2021,Mann and Patterson. 2013,Jearld and Miller.

1983). Other examples in the opposite direction of the water-air interface include works on the

widespread hunting behavior implemented by birds to submerge prey by diving (Capuska et al.

2011,Machovsky-Capuska et al. 2012).

This response profile in the archerfish is also consistent with the results of multiple studies of

adaptation process in mammals (Tseng et al. 2007,Shadmehr and Mussa-Ivaldi. 1994a,Darmohray

et al. 2019,Kitazawa et al. 1995). When the sensory feedback to the executed movement differs

from the expected feedback, animals can learn to adjust the strength and the direction of their

subsequent movements until they are able to perform the task successfully. For instance, in studies

of prism adaptation, subjects are asked to throw a ball at a target while wearing glasses with prisms

that shift their visual field horizontally. For all subjects the initial error was shown to decrease

after several trials; removal of the prisms caused disorientation and error in the opposite direction

that also decreased with time (Fernandez-Ruiz and Diaz. 1999a,Kitazawa et al. 1995). Another

important category of motor adaptation studies has investigated reaching movements when

subjected to either visual distortions or opposing force fields (Shadmehr et al. 2010,Shadmehr and

Mussa-Ivaldi. 1994b,Kluzik et al. 2008). The results with respect to the error magnitude were

similar and exhibited a gradual decrease with time and the existence of an aftereffect. Finally, the

study of visuo-motor adaptation in mammals underscores the importance of the cerebellum as the

neural site for this type of adaptation (Darmohray et al. 2019,Donchin et al. 2012,Popa et al.

2016). While the neuroanatomy of the archerfish has been mapped (Karoubi et al. 2016), it remains

unclear whether the archerfish homologue is similar.

Finally, it is important to note that not all fish subjects of the current study advanced to the final

test, due to the challenging nature of the task and the difficulty in maintaining the fish's motivation

throughout the test, which relies on feeding. Such challenges are common in behavioral studies.

Consequently, this situation raises the question of whether all individuals are capable of solving

the motor adaptation task. Addressing this question is a task for future studies.

Our work also suggests how the archerfish corrects for physical factors such as refraction through

the use of motor adaptation as part of the correction process. In so doing, it sheds light on motor

adaptation in vertebrates in general.

Acknowledgements

We thank Mor Ben-Tov and Ehud Vinepinksy for help in the initial stages of this project. We

gratefully acknowledge financial support from THE ISRAEL SCIENCE FOUNDATION—FIRST

Program (Grant no. 281/15), THE ISRAEL SCIENCE FOUNDATION (Grant

no. 824/21), and The Human Frontiers Science Foundation Grant RGP0016/2019.

Competing interests

The authors declare no competing interests

References

Barta A, Horváth G (2003) Underwater binocular imaging of aerial objects versus the position of
eyes relative to the flat water surface. JOSA A 20:2370-7

Ben-Simon A, Ben-Shahar O, Vasserman G, Ben-Tov M, Segev R (2012) Visual acuity in the
archerfish: behavior, anatomy, and neurophysiology. Journal of Vision 12:18

Ben-Tov M, Ben-Shahar O, Segev R (2018) What a predator can teach us about visual
processing: a lesson from the archerfish. Curr Opin Neurobiol 52:80-7

Ben-Tov M, Donchin O, Ben-Shahar O, Segev R (2015) Pop-out in visual search of moving
targets in the archer fish. Nature Communications 6:1-11

Capuska GEM, Vaughn RL, Würsig B, Katzir G, Raubenheimer D (2011) Dive strategies and
foraging effort in the Australasian gannet Morus serrator revealed by underwater videography.
Mar Ecol Prog Ser 442:255-61

Cohen L, Vinepinsky E, Donchin O, Segev R (2023) Boundary vector cells in the goldfish
central telencephalon encode spatial information. PLoS Biology 21:e3001747

Darmohray DM, Jacobs JR, Marques HG, Carey MR (2019) Spatial and temporal locomotor
learning in mouse cerebellum. Neuron 102:217,231. e4

Dewenter J, Gerullis P, Hecker A, Schuster S (2017) Archerfish use their shooting technique to
produce adaptive underwater jets. J Exp Biol 220:1019-25

Dill L (1977) Refraction and the spitting behavior of the archerfish (Toxotes chatareus). Behav
Ecol Sociobiol 2:169-84

Donchin O, Rabe K, Diedrichsen J, Lally N, Schoch B, Gizewski ER, Timmann D (2012)
Cerebellar regions involved in adaptation to force field and visuomotor perturbation. J
Neurophysiol 107:134-47

Fernandez-Ruiz J, Diaz R (1999a) Prism adaptation and aftereffect: specifying the properties of a
procedural memory system. Learn Mem 6:47-53

Fernandez-Ruiz J, Diaz R (1999b) Prism adaptation and aftereffect: specifying the properties of a
procedural memory system. Learn Mem 6:47-53

Gabay S, Leibovich T, Ben-Simon A, Henik A, Segev R (2013) Inhibition of return in the archer
fish. Nature communications 4:1-5

Givon S, Altsuler-Nagar R, Oring N, Vinepinsky E, Segev R (2023) Lateral and medial
telencephalic pallium lesions impair spatial memory in goldfish. Brain Res Bull 204:110802

Givon S, Pickholtz R, Pickholtz EY, Ben-Shahar O, Kiflawi M, Segev R (2022) Toward
naturalistic neuroscience of navigation: opportunities in coral reef fish. Frontiers in Neural
Circuits 16:895381

Izawa J, Shadmehr R (2011) Learning from sensory and reward prediction errors during motor
adaptation. PLoS Comput Biol 7:e1002012. doi: 10.1371/journal.pcbi.1002012 [doi]

Jearld A, Miller RJ (1983) Behavior and phylogeny of fishes of the genus Colisa and the family
Belontiidae. Behaviour 83:155-85

Jones NA, Klump BC, Abaurrea TM, Harrower S, Marr C, Scott L, Rendell L, Webster MM
(2021) Short-range hunters: exploring the function and constraints of water shooting in dwarf
gouramis. J Exp Biol 224:jeb243477

Karoubi N, Segev R, Wullimann MF (2016) The Brain of the Archerfish Toxotes chatareus: A
Nissl-Based Neuroanatomical Atlas and Catecholaminergic/Cholinergic Systems. Frontiers in
neuroanatomy 10

Kitazawa S, Kohno T, Uka T (1995) Effects of delayed visual information on the rate and
amount of prism adaptation in the human. Journal of Neuroscience 15:7644-52

Kluzik J, Diedrichsen J, Shadmehr R, Bastian AJ (2008) Reach adaptation: what determines
whether we learn an internal model of the tool or adapt the model of our arm? J Neurophysiol
100:1455-64. doi: 10.1152/jn.90334.2008 [doi]

Krakauer JW, Mazzoni P (2011) Human sensorimotor learning: adaptation, skill, and beyond.
Curr Opin Neurobiol 21:636-44. doi: 10.1016/j.conb.2011.06.012 [doi]

Kruschke J (2014) Doing Bayesian data analysis: A tutorial with R, JAGS, and Stan. Academic
Press

Lee SA, Vallortigara G, Ruga V, Sovrano VA (2012) Independent effects of geometry and
landmark in a spontaneous reorientation task: a study of two species of fish. Animal cognition
15:861-70

Machovsky-Capuska GE, Howland HC, Raubenheimer D, Vaughn-Hirshorn R, Würsig B,
Hauber ME, Katzir G (2012) Visual accommodation and active pursuit of prey underwater in a
plunge-diving bird: the Australasian gannet. Proceedings of the Royal Society B: Biological
Sciences 279:4118-25

Mann J, Patterson EM (2013) Tool use by aquatic animals. Philosophical Transactions of the
Royal Society B: Biological Sciences 368:20120424

Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT (1996a) Throwing while looking
through prisms: I. Focal olivocerebellar lesions impair adaptation. Brain 119:1183-98

Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT (1996b) Throwing while looking
through prisms: I. Focal olivocerebellar lesions impair adaptation. Brain 119:1183-98

Mokeichev A, Segev R, Ben-Shahar O (2010) Orientation saliency without visual cortex and
target selection in archer fish. Proc Natl Acad Sci U S A 107:16726-31. doi:
10.1073/pnas.1005446107 [doi]

Newport C, Wallis G, Siebeck UE (2018) Object recognition in fish: accurate discrimination
across novel views of an unfamiliar object category (human faces). Anim Behav 145:39-49

Newport C, Wallis G, Siebeck UE (2015) Same/different abstract concept learning by archerfish
(Toxotes chatareus). PloS one 10:e0143401

Newport C, Wallis G, Siebeck UE (2014) Concept learning and the use of three common
psychophysical paradigms in the archerfish (Toxotes chatareus). Frontiers in neural circuits 8:39

Plummer M (2003) JAGS: A program for analysis of Bayesian graphical models using Gibbs
sampling. 124:1-10

Popa LS, Streng ML, Hewitt AL, Ebner TJ (2016) The errors of our ways: understanding error
representations in cerebellar-dependent motor learning. The Cerebellum 15:93-103

Rodriguez F, Duran E, Vargas JP, Torres B, Salas C (1994) Performance of goldfish trained in
allocentric and egocentric maze procedures suggests the presence of a cognitive mapping system
in fishes. Anim Learn Behav 22:409-20

Rossel S, Corlija J, Schuster S (2002) Predicting three-dimensional target motion: how archer
fish determine where to catch their dislodged prey. J Exp Biol 205:3321-6

Scheidt RA, Conditt MA, Secco EL, Mussa-Ivaldi FA (2005) Interaction of visual and
proprioceptive feedback during adaptation of human reaching movements. J Neurophysiol
93:3200-13

Schlosser JA (1764) An account of a fish from Batavia, called jaculator: In a letter to Mr. Peter
Collinson, FRS from John Albert Schlosser, MDFR S. Philosophical Transactions of the Royal
Society of London :89-91

Segev R, Schneidman E, Goodhouse J, Berry MJ (2007) Role of eye movements in the retinal
code for a size discrimination task. J Neurophysiol 98:1380-91

Shadmehr R, Mussa-Ivaldi FA (1994a) Adaptive representation of dynamics during learning of a
motor task. J Neurosci 14:3208-24

Shadmehr R, Mussa-Ivaldi FA (1994b) Adaptive representation of dynamics during learning of a
motor task. J Neurosci 14:3208-24

Shadmehr R, Smith MA, Krakauer JW (2010) Error correction, sensory prediction, and
adaptation in motor control. Annu Rev Neurosci 33:89-108. doi: 10.1146/annurev-neuro-
060909-153135 [doi]

Temple S (2007) Effect of salinity on the refractive index of water: considerations for archer fish
aerial vision. J Fish Biol 70:1626-9

Temple S, Manietta D, Collin SP (2013) A comparison of behavioural (Landolt C) and
anatomical estimates of visual acuity in archerfish (Toxotes chatareus). Vision Res 83:1-8

Timmermans P, Vossen J (2000a) Prey catching in the archer fish: does the fish use a learned
correction for refraction? Behav Processes 52:21-34

Timmermans P, Vossen J (2000b) Prey catching in the archer fish: does the fish use a learned
correction for refraction? Behav Processes 52:21-34

Tseng Y, Diedrichsen J, Krakauer JW, Shadmehr R, Bastian AJ (2007) Sensory prediction errors
drive cerebellum-dependent adaptation of reaching. J Neurophysiol 98:54-62

Tsvilling V, Donchin O, Shamir M, Segev R (2012) Archer fish fast hunting maneuver may be
guided by directionally selective retinal ganglion cells. Eur J Neurosci 35:436-44

Vasserman G, Shamir M, Simon AB, Segev R (2010) Coding "what" and "when" in the archer
fish retina. PLoS computational biology 6:e1000977

Vinepinsky E, Cohen L, Perchik S, Ben-Shahar O, Donchin O, Segev R (2020) Representation
of edges, head direction, and swimming kinematics in the brain of freely-navigating fish.
Scientific reports 10:14762

Volotsky S, Ben-Shahar O, Donchin O, Segev R (2022) Recognition of natural objects in the
archerfish. J Exp Biol 225:jeb243237

Volotsky S, Vinepinsky E, Donchin O, Segev R (2019) Long-range neural inhibition and
stimulus competition in the archerfish optic tectum. Journal of Comparative Physiology A
205:537-52

Wallman J, Fuchs AF (1998) Saccadic gain modification: visual error drives motor adaptation. J
Neurophysiol 80:2405-16