**Title: Inhibition of miR-199b-5p reduces pathological alterations in Osteoarthritis by tar-**

**geting *Fzd6* and *Gcnt2*.**

**Authors:** Tong Feng[1]†, Qi Zhang[1,2]†, Si-Hui Li[1], Shuan-hu Zhou[3,4], Xin Wang[5], Jun-Meng

Wang[1], Fan-Rong Liang[1], Shu-Guang Yu[1,6]*, Qiao-Feng Wu[1, 6]*

**Affiliations:**

1 Acupuncture and Tuina College, Chengdu University of Traditional Chinese Medicine,

Chengdu, Sichuan, 610075, China

2 Chongqing Hospital of Traditional Chinese Medicine, Chongqing, 400011, China

3 Department of Orthopedic Surgery, Brigham and Women's Hospital, Harvard Medical

School, Boston, Massachusetts, 02115, USA

4 Harvard Stem Cell Institute, Harvard University, Cambridge, Massachusetts, 02138, USA

5 Departments of Neurosurgery, Brigham and Women's Hospital, Harvard Medical School,

Boston, Massachusetts, 02115, USA

6 Acupuncture & Chronobiology Key Laboratory of Sichuan Province, Chengdu, Sichuan,

610075, China

*Corresponding authors.

Qiao-feng Wu: wuqiaofeng@cdutcm.edu.cn

ORCID: 0000-0002-8308-2843

Shu-guang Yu: ysg@cdutcm.edu.cn

ORCID：0000-0002-1413-8989

Address: No.37, Road Shi-Er-Qiao, Jinniu District, Chengdu, Sichuan, 610075, China

† These authors contributed equally to this manuscript.

**Abstract**

Osteoarthritis (OA) is a degenerative disease with a high prevalence in the elderly population,

but our understanding of its mechanisms remains incomplete. Analysis of serum exosomal small

RNA sequencing data from clinical patients and gene expression data from OA patient serum

and cartilage obtained from the GEO database revealed a common dysregulated miRNA, miR-

NA-199b-5p. In vitro cell experiments demonstrated that miRNA-199b-5p inhibits chondrocyte

vitality and promotes extracellular matrix degradation. Conversely, inhibition of miRNA-199b-

5p under inflammatory conditions exhibited protective effects against damage. Local viral injec-

tion of miRNA-199b-5p into mice induced a decrease in pain threshold and OA-like changes. In

an OA model, inhibition of miRNA-199b-5p alleviated the pathological progression of OA. Fur-

thermore, bioinformatics analysis and experimental validation identified *Gcnt2* and *Fzd6* as tar-

get genes of miRNA-199b-5p. Thus, these results indicated that miRNA-199b-5p/*Gcnt2* and

*Fzd6* axis might be a novel therapeutic target for the treatment of OA.

**Keywords: Osteoarthritis; Extracellular matrix; miRNA-199b-5p; Animal**

**Introduction**

Osteoarthritis (OA)  is a degenerative disease characterized by the deterioration of articular carti-

lage and affecting all components of the joint, including the synovium, subchondral bone, and

meniscus[1]. Knee OA (KOA) is the leading cause of disability and increased living costs among

elderly patients[2]. Currently, the treatment options for osteoarthritis are limited to symptomatic

relief and joint replacement. However, the insufficient relief of symptoms, potential medication

side effects, and the economic burden and complications associated with joint replacement sur-

geries have prompted the need to identify new, effective, and safe treatment methods and explore

unknown targets for individuals with KOA[3].

Exosomes are extracellular vesicles that are distributed in body fluids such as serum and play a

crucial biological role by delivering molecules such as MicroRNAs (miRNAs)[4]. They serve as

vital carriers for intercellular communication and transfer of genetic information. MiRNAs are a

class of non-coding RNAs ranging from 18 to 24 nucleotides in length, which exert inhibitory

effects on the expression of target genes through interactions with mRNA. Studies have demon-

strated that miRNAs play a crucial role as a pathogenic factor in osteoarthritis (OA).[5]. For in-

stance,  miR-199b-5p was found to contribute to the osteogenic differentiation of bone marrow

stromal cells [6]; another miR-140 showed cartilage-specific expression, and its expression was

significantly reduced in OA cartilage [7;8].recently, intra-articular injection of antisense oligonu-

cleotides of miR-181a-5p produced chondroprotective effects in OA mice [9].

In this study, we initially screened miR-199b-5p as a potential key miRNA based on clinical data

by detecting differentially expressed exosomal miRNAs. To elucidate the role and function of

miR-199b-5p, we investigated its impact on chondrocytes, which are crucial pathogenic cells in

KOA, and identified its molecular targets through in vitro experiments. Furthermore, we ex-

plored its role in vivo. Hence, this article not only identifies miR-199b-5p as a potential micro-

target for KOA but also provides a potential strategy for future identification of new molecular

drugs.

**Results**

**Identification and enrichment analysis of differentially expressed miRNAs in serum exosomes**

To investigate the dysregulated miRNAs in serum exosomes of KOA patients, we extracted exosomal miRNAs from serum and performed sequencing. Serum samples from 15 patients with KOA and 10 healthy subjects were collected (Supplementary Table 2). After extraction, the serum exosomes were observed under transmission electron microscopy and nanoparticle tracking analysis, revealing a diameter ranging from approximately 70 to 150 nm (Fig. S1A, B). Furthermore, the surface marker proteins CD9, CD63, and CD81 were found to be expressed on the exosomes (Fig. S1C). Next, we performed sequencing of miRNAs in serum exosomes.

The results showed that 88 miRNAs were up-regulated and 89 miRNAs were downregulated in KOA patients compared with the control group based on fold change > 1.5 and $p < 0.05$ (Fig. 1A, B). Afterwards, we performed bioinformatics Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses on these differentially expressed genes. Among the top 5 enriched results of up-regulated differentially expressed miRNAs, the CC (Cell, intracellular, and cytoplasm) GO terms were notable. The MF (Translation activator activity, binding to ubiquitin-conjugating enzymes, and possessing transferase activity) GO terms and BP (Cell differentiation and the stabilization of mRNA through 3'-UTR mediation) GO terms were also significant (Fig. 1C). The top 10 enriched KEGG pathways, PI3K-AKT signaling, Ras signaling, apoptosis, and other types of O-glycan biosynthesis are considered to be of significant importance (Fig. 1D).The analysis of the top 5 down-regulated differential miRNAs reveals interesting findings. the CC (Cell, intracellular) GO terms were particularly noteworthy. The MF (RNA polymerase II transcription factor activity and sequence-specific DNA binding) GO terms, as well as the BP (collagen-activated signaling pathway) GO terms, were also deemed significant (Fig. 1E). More-

over, the analysis of the top 10 enriched KEGG pathways revealed a strong association with the

Hedgehog signaling pathway, pluripotency of stem cells, and glycosaminoglycan biosynthesis-

keratan sulfate (Fig. 1F).

It is noteworthy that in the aforementioned results, both the up-regulated and down-regulated dif-

ferential miRNA enrichment analyses are involved in processes related to cartilage synthesis,

including O-glycan biosynthesis and glycosaminoglycan biosynthesis-keratan sulfate[10;11].

**96 Dysregulated serum exosomal miRNAs were found to be expressed in both serum and car-**

**97 tilage**

Subsequently, we aimed to investigate whether these dysregulated miRNAs in serum exosomes

show differential expression in other affected sites or tissues of KOA. Leveraging the KOA pa-

tient cartilage and serum sequencing data available in the GEO database, we compared our

dysregulated miRNA list with the datasets. Remarkably, we identified 169 miRNAs that exhibit-

ed differential expression in the serum of KOA patients (Fig. 2A), suggesting their involvement

in the disease. Moreover, these miRNAs were also found to be expressed in KOA patient carti-

lage (Fig. 2B). This robust validation reinforces the reliability of our data. Consequently, we pro-

ceeded with further screening of differentially expressed genes.

**107 miRNA-199b-5p has been identified as a potential target molecule in KOA.**

Based on the P-value and exosomal expression, five miRNAs（hsa-mir-3168, hsa-mir-1296-5p,

hsa-mir-15b-3p, hsa-mir-338-3p and hsa-mir-199b-5p) were selected to further research and val-

idated in independent human samples by RT-qPCR (hsa-mir-199b-5p, $p < 0.01$; hsa-mir-3168,

$p$=0.33; hsa-mir-15b-3p, $p < 0.01$; hsa-mir-338-3p, $p$=0.20 and hsa-mir-1296-5p, $p < 0.01$) (Fig.

2C). To further explore the functional roles of these miRNAs, we established a mouse model of

KOA to evaluate their expression in the joints. Due to species limitations, we examined the ex-

pression of miR-199b-5p ($p$ <0.01), miR-15b-3p ($p$ <0.05) and miR-338-3p ($p$ <0.05) in joint

tissue samples (control $vs$ M-0.5) (Fig. 2D). The results showed that only the expression trend of

miR-199b-5p was consistent between the clinical samples and the mouse arthritis model. There-

fore, we selected miR-199b-5p as the target for our subsequent research.

**miR-199b-5p mimic inhibits the cell viability and extracellular matrix (ECM) of chondro-**

**cytes and inhibitor restores LPS-induced chondrocytes damage.**

We first extracted mouse primary chondrocytes (Fig. 3A-C). In vitro experiments showed that

overexpression miR-199b-5p inhibited the viability of chondrocytes($p<0.01$) (Fig. 3D, Fig. S2).

we also find miR-199b-5p mimic increased the mRNA expressions of *MMP3* ($p$=0.09) and

*ADAMTS5* ($p<0.05$) and decreased the mRNA expression of *COL2A1*($p$=0.05), *AGGRECAN*

($p$=0.20) and *SOX9* ($p$=0.22), which are often used as the biomarkers of chondrocytes ECM met-

abolic balance. In contrast, miR-199b-5p inhibitor decreased the mRNA expression of *MMP3*

($p<0.01$) and *ADAMTS5* ($p$=0.11) and increased the mRNA expression of *COL2A1* ($p$=0.07),

*AGGRECAN* ($p<0.01$) and *SOX9* ($p<0.01$) (Fig. 3E–I). While some gene expression changes

may not be significant in statistic, but the modulation of miR-199b-5p expression has been ob-

served to exert an influence on the metabolic alterations of chondrocytes.

To explore the effect of miR-199b-5p under pathological conditions, an LPS-stimulated inflam-

mation chondrocyte cell model was established. We examined the effect of LPS at 5, 10, and 15

μg/ml on cell viability and found that cell viability was significantly decreased at 15

μg/ml($p<0.001$) (Fig. 2J). Next, we chose 15 μg/ml of LPS to establish a chondrocyte injury

model. The CCK-8 assay showed that miR-199b-5p inhibitor reversed the decrease of cell viabil-

ity caused by LPS ($p<0.01$) (Fig. 3K). Also, we revealed that LPS elevated *MMP3* ($p<0.01$) and

*ADAMTS5* ($p<0.01$) and decreased *COL2A1*($p=0.06$), *AGGRECAN* ($p<0.01$), and *SOX9* ($p=0.13$)

expression. In the presence of miR-199b-5p inhibitor, the changes in mRNA levels were re-

versed (Fig. 3L-P). These results suggest that miR-199b-5p overexpression reduces cell viability

and miR-199b-5p inhibition partly restores LPS-induced cell damage and ECM degradation.

**miR-199b-5p mimic induces inflammation in normal mice, and miR-199b-5p inhibitor alle-**

**viates symptoms in KOA mice.**

Now，we want to know the vivo role of miRNA-199b-5p. Firstly, we screened Adneovirus(AD)

and utilized High-AD as a vector to either overexpress or inhibit the expression of miR-199b-

5p(Fig. S3). Then, miR-199b-5p mimic was injected to the joint of mice (Fig. 4A) and a decrease

in pain threshold was found ($p<0.01$) (Fig. 4B). Four weeks later, serum *IFN-γ* ($p<0.01$) and

*TNF-α* ($p<0.01$) were also significantly increased (Fig. 4C, D). The cartilage of mice injected

with the miR-199b-5p mimic was slightly degraded ($p=0.02$) (Fig. 4E, F). Additionally, the ar-

ticular surface showed insect erosion (Fig. 4G). interestingly, the level of serum inflammation in

the miR-199b-5p inhibitor injection group was significantly decreased compared with the mimic

group (*IFN-γ*, $p<0.01$; *TNF-α*, $p<0.01$). These results indicated that intra-articular injection of

miR-199b-5p mimic induced inflammation response in mice.

Followingly, we established an MIA-induced KOA model to further investigate the role of miR-

199b-5p (Fig. S4). We observed a decrease in pain threshold in the model group, and recovery

was observed on the 10th day in the inhibitor group ($p<0.01$) (Fig. 5A, B). Joint tissues were tak-

en at the fourth week, and revealed that the expression of *IFN-γ* ($p<0.01$) and *TNF-α* ($p<0.01$)

decreased after inhibiting miRNA-199b-5p (Fig. 5C, D). Safranin-fast green staining of joints

showed recovery of articular cartilage degradation ($p<0.05$) (Fig. 5E, F), and the moth erosion of

the cartilage μCT was partly improved (Fig. 5G). These results proved that intra-articular injec-

tion of the miR-199b-5p inhibitor partly recovered pain, inflammation, and cartilage degenera-

tion in KOA mice.

**_Gcnt2_ and _Fzd6_ are two target genes of miR-199b-5p**

In order to investigate the underlying mechanism of miRNA-199b-5p, we utilized five widely

used miRNA target gene prediction tools, namely miRWalk, miRDB, TarBase, starbase, and

TargetScan, to identify potential target genes. Consequently, we identified six putative target

genes of miRNA-199b-5p (Fig. 6A). Bioinformatics analysis of the six possible target genes

showed that BP is in posttranscriptional regulation of gene expression and angiogenesis; CC is in

cytoplasmic vesicle and cell leading edge; and MF is in ubiquitin protein ligase binding and pro-

tein domain specific binding (Fig. 6B).

After Mimic infected cells, we found decreases expression in *Fzd6* (P<0.01), *Gcnt2* (P<0.01),

and *Caprin1* (P=0.024) in the chondrocytes (Fig. 6C). Conversely, after inhibitor infection, *Hif-*

*1α* (P=0.048), *Fzd6* (P<0.01) and *Gcnt2* (P<0.01) were increased and *Atg14* (P=0.042) increased

(Fig. 6D). Notably, we observed corresponding changes in the expression of *Fzd6* and *Gcnt2* up-

on miR-199b-5p overexpression and under-expression. Moreover, we predicted potential binding

sites for miRNA-199b-5p within the 3'-untranslated region (UTR) of these two target genes (Fig.

6E, F) and luciferase reporter assays confirmed that miR-199b-5p can bind to and suppress the

expression of both *Gcnt2* (P < 0.01) and *Fzd6* (P < 0.01) *via* their complementary sequences

(Fig. 6G, H). Furthermore, we found differential expression of *Fzd6* in both synovial tissue data

(GDS5401) and chondrocyte data (GDS3758) from KOA patients in the GEO profile (Fig. 6I, J).

Similarly, differential expression tendency of *Gcnt2* was observed in both the synovial tissue da-

ta (GDS5403) and GDS3758 from the same KOA patients (Fig. 6K, L). These findings provide

further validation to our results. Therefore, *Fzd6* and *Gcnt2* was confirmed as an important

downstream target for further research.

**Discussion**

In this study, we initially performed sequencing of serum exosomal miRNAs from clinical pa-

tients and identified 177 dysregulated miRNAs. Subsequently, through comparison with GEO

data, we found that 169 miRNAs were expressed in both KOA serum and cartilage. Following a

screening process, miR-199b-5p was selected for further experiments. In cell-based assays, we

discovered that miR-199b-5p can influence the viability of chondrocytes and cytokine-mediated

extracellular matrix metabolism. Moreover, in vitro experiments demonstrated that it can induce

inflammation and abnormal pain threshold in normal mice. Importantly, inhibition of miR-199b-

5p alleviated the pathological symptoms of KOA. Finally, these effects were achieved by target-

ing *Gcnt2* and *Fzd6* (Fig. 7). Thus, our findings demonstrated that miR-199b-5p might be a novel

potential therapeutic target for OA prevention and treatment.

miRNAs function as regulators of gene expression in biological processes by regulate mRNA

translation by specifically binding to the 3′UTRs of target mRNAs [12]. Exosomes are rich in

miRNAs, and various cells can secrete exosomes to target cells under physiological and patho-

logical conditions, which function as delivery vehicles for miRNAs [13;14]. There is evidence sug-

gesting that miRNAs can participate in various cellular processes (inflammation, cell viability,

ECM dysregulation) and signaling pathways (Hedgehog signaling, PI3K-AKT signaling) rele-

vant to OA[15;16]. Our differential miRNA enrichment analysis also strongly supports the correla-

tion with these findings.

In cancer research, overexpression of miR-199b-5p can inhibit the proliferation, migration and

invasion of prostate cancer cells in vitro and tumor growth and metastasis in vivo by targeting

DDR1[17]. In addition, exogenous miR-199b-5p inhibited the growth of bone marrow mesenchy-

mal stem cells (MSCs) and promoted the differentiation of bone MSCs into chondrocytes by tar-

geting the *JAG1* pathway [18;19]. Our functional experiments showed that overexpression of miR-

199b-5p reduced chondrocyte viability and the expression of anabolic factors such as COL2A1,

AGGREGN, and SOX9. It also increased inflammation levels and decreased pain thresholds in

control mice. Although no pathological changes were observed in the articular cartilage of con-

trol mice, a similar study demonstrated that pathological cartilage changes only occurred after six

months in mice with miR-211 and miR-204 knockout[20]. Therefore, we speculate that the time of

overexpression of miR-199b-5p in our experiment was too short, so it only caused inflammation

and pain threshold response. To best of our knowledge, this is the initial investigation document-

ing the involvement of miR-199b-5p in KOA.

*Fzd6* is known to be up-regulated during the osteogenic differentiation of MSCs and can be regu-

lated by miR-194-5p to activate the WNT signaling pathway, thereby promoting osteogenic dif-

ferentiation of MSCs[21]. *Gcnt2* has been found to induce epithelial-mesenchymal transition and

enhance migration and invasion of esophageal squamous cell carcinoma cells[22]. Our findings in-

dicate that miR-199b-5p plays a crucial role in KOA by targeting *Fzd6* and *Gcnt2*. Future re-

search should further explore the roles and mechanisms of *Fzd6* and *Gcnt2* in the context of

KOA.

This study also has some limitations. We initially detected serum exosomes miRNA and later

examined the miRNA through in vitro experiment and animal study. However, whether the

miRNAs targeting the joints have the same role as the serum exosome miRNAs or blood miRNA

has not been clear. We also performed direct intra-articular injection of adenoviral vectors. The

intra-articular injection can reduce the exposure of extra-articular tissue, thereby minimizing the

associated side effects and improving targeting. Besides, more in-vivo study such as gene

knockout mice study can be used in the future study.

In conclusion, we found that miR-199b-5p is elevated in osteoarthritis and may affect cell viabil-

ity and related cytokines by targeting *Gcnt2* and *Fzd6*. Overexpression of miR-199b-5p induced

OA-like pathological changes in normal mice and inhibiting miR-199b-5p alleviated symptoms

in KOA mice, suggesting it may be a target for treatment of OA. Through human clinical trials,

cell experiments, and animal models, we not only identified a new OA-related miR-199b-5p but

also examined the biological function of miR-199b-5p in OA.

**MATERIALS AND METHODS**

**Human samples**

Study participants were recruited from the Hospital of Chengdu University of Traditional Chi-

nese Medicine and the surrounding communities. The study was approved by the Ethics Review

Committee of the Hospital at Chengdu University of Traditional Chinese Medicine (2016KL-

017) and conformed to the ethical guidelines of the 1975 Declaration of Helsinki. Serum was

collected from all participants and serum samples were stored at -80°C.

**Human subjects**

Patients were enrolled if they fulfilled the following criteria: (1) diagnosed with KOA according

to the ACR; (2) age between 40 and 65 years; (3) agreed to cooperate with researchers in all re-

search procedures after enrollment; (4) provided with written informed consent. Patients with

any of the following conditions were excluded: (1) accompanied with other diseases such as

rheumatoid arthritis, bone tumors, and bone tuberculosis; (2) treated with intra-articular gluco-

corticoid or viscoelastic supplementation in the last six months; (3) knee replacement history; (4)

had complicated cardiovascular disease, diabetes, skin disease, and liver or kidney impairment;

(5) pregnant or breastfeeding women; (6) accompanied with mental and intellectual disabilities;

or (7) undergoing other clinical trials.

**Extraction and sequencing of serum exosome miRNAs**

Serum samples were filtered using 0.22 μM filters. Exosomes were isolated from the serum

sample using ExoQuick™ Exosome Precipitation Solution (System Biosciences) following the

manufacturer's instructions. Briefly, serum was thawed on ice and centrifuged at 4000 rpm for

15 min to remove any cells or cellular debris. Next, 50 μL ExoQuick Solution was added into the

200 μL serum sample and mixed thoroughly. The exosomes were suspended in PBS.

Exosomes were characterized by electron microscopy (Tecnai G2 Spirit 120KV, FEI), nanopar-

ticle tracking analysis (NTA) (NTA 3.2 Dev Build 3.2.16), and western blot analysis (for CD9,

CD63 and CD81). Sequencing was performed by single-end sequencing (1×150 bp) on Illumina

NextSeq 500. The libraries were sequenced on an Agilent 2100 Bioanalyzer platform. The

mirdeep2 software (https://www.mdc-berlin.de/content/mirdeep2-documentation) was used to

analyze the miRNA sequences and quantification. The heatmap was plotted based on the log2

(fold change), using Heatmap Illustrator software (Heml 1.0).

**Total RNA was isolated from human serum, cell and mice**

Chondrocytes using a kit (Yeasen,Shanghai, China) according to the manufacturer's instructions.

Reverse transcription was performed using 1000 ng total RNA and a Prime Script RT Reagent

Kit(Yeasen, Shanghai, China) or Prime Script RT Master Mix (Yeasen, Shanghai, China), which

were used for miRNA and mRNA, respectively. For miRNA, the reactions were incubated at

42°C for 15 min followed by inactivation at 85°C for 5 s. qRT-PCR amplification was assessed

in a CFX Connection Real-Time System (Bio-Rad) using the SYBR Premix Ex Taq II kit

(Yeasen, Shanghai,China). The following cycling conditions were used: 95°C for 30 s, followed

by 40 cycles of 95°C for 5 s and 60°C for 30 s. All reactions were performed in duplicate and

normalized to the internal reference U6 for miRNA and GAPDH mRNA for mRNAs. The $2^{-\Delta\Delta CT}$

CT method was used to evaluate the relative mRNA/miRNA expression levels.

**RNA primer**

Primers are listed in Supplementary Table 1.

**Bioinformatics analysis**

GO analysis was performed, involving the biological process (BP), cellular component (CC), and

molecular function (MF), using DAVID[23] (https://david.ncifcrf.gov/home.jsp).We used the

miRNA target gene prediction websites miRDB[24](http://mirdb.org/), miR-

Walk[25](http://mirwalk.umm.uni-heidelberg.de/), Starbase[26] (https://starbase. sysu.edu.cn/), DI-

ANA-TarBase[27] (http://diana.imis.athena-innovation.gr/DianaTools/ index.php?r=tarbase/index),

Targetscan[28] (https://www.targetscan.org/vert_72/). The target genes of mmu-miR-199b-5p were

predicted, and the intersection of the target gene prediction list results of the five sites was taken.

The dysregulated miRNAs were compared to relevant published miRNA data from human KOA

patients. The dataset GSE105027 was obtained from serum samples of KOA patients, while

GSE175961 comprised sequencing data from KOA patient cartilage. Finally, we validated the

expression of the target genes Gcnt2 and Fzd6 in KOA patients using synovial data (GDS5401,

GDS5403) and cartilage data (GDS3758).

**Primary mouse chondrocyte culture**

296  Primary mouse cartilage was extracted following the protocol of Gosset et al.[29]. The cells were

297  cultured in F12 medium containing 10% fetal bovine serum and 1% penicillin-streptomycin at

298  37°C and a 5% $CO_2$. The medium was changed every three days, and cells were cultured to the

299  second passage for experiments. Cells were stimulated with LPS for 6 h before subsequent inves-

300  tigation.

**Animal model of KOA**

Animal experiments were performed using total of 108 8-week-old male C57BL/6 mice. The

protocol was approved by the Committee on the Ethics of Animal Experiments of Chengdu Uni-

versity of Traditional Chinese medicine (2019-04). After randomization, the KOA mouse model

was established by orthotopic injection of MIA (sigma) into the knee joint of mice. After the an-

imals were anesthetized with isoflurane in a small animal anesthesia machine, the right knee

joint of the mouse was shaved, the knee joint was flexed 90°, and 10 μL of MIA solution was

injected into the knee joint cavity using a 28G microsyringe. The syringe needle was slowly

withdrawn and the knee joint was gently moved. Before model establishment and at 3, 7, 10, 14,

21, and 28 days after model establishment. After the 3rd day and 14th day after the start of the

experiment, the experimental mice were injected with adenovirus (total volume 10μL) in the

knee joint, and the control mice were injected with empty adenovirus as a control (Hanbio tech,

Shanghai, China)[30].

**Thermal pain threshold detection**

A pain threshold detection instrument was used to measure the latency of the right plantar leg

raising reflex of mice under heat radiation. Three days before the experiment, mice were cut off

from water for half a day and adapted to the environment for 1 h. A certain intensity of pyrogen

was used to irradiate the plantar position of the mouse's right limb, and a machine was used to

automatically record the time when the mouse moved the limb from thermal pain. The resting

intensity of the thermal pain stimulator was set to 10%, and the maximum duration was set to 20

s to avoid prolonged heating and burning of skin. After the light source is aimed at the sole of the

mouse, we observed the mouse in real time. If paw raising, paw licking, or paw retraction was

observed, the irradiation was stopped and the irradiation time of the instrument was recorded.

The right limb of each mouse was tested five times, with an interval of 10 min each time; outliers

were eliminated, and the average value was calculated and included in the statistics[31].

**ELISA and μCT scanning**

Mice were sacrificed after four weeks, and serum was collected for ELISA assay (Jiangsu Jing-

mei Biotechnology Co., Ltd., Yancheng, China). The knee joint specimens of the right hindlimb

of mice were scanned using a high-resolution micro-CT skysan1267 instrument (Bruker, Germa-

ny). The sample was removed from fixative and dried. The scanning parameters were as follows:

voltage 55 KV, current 200 μA, and filter 0.25AL. After scanning, three-dimensional reconstruc-

tion was performed using NRecon1.7.4.2 software.

**Safranin Fast green staining**

334 Mouse knee joint staining was performed using the Safranin Fast Green Staining Kit (Servicebio

335 Biological Technology, Wuhan, China). The sample was examined under a microscope, and the

336 cartilage integrity was scored according to the OARSI grading system, in which the score ranged

337 from 0 to 6 points[32].

**Luciferase assay**

The wild-type (WT) and mutant-type (MUT) sequences (according to the predicted binding site)

were inserted into the pmiRGLO plasmid. HEK-293T cells were seeded in 6-well plates at 24 h

before transfection. GCNT2 and FZD6 3′UTR-wt and gcnt2 and fzd6 3′UTR-mut plasmids (500

ng) and 20 nmol miR-199b-3p and NC were co-transfected with Lipofectamine 3000 (Hanbio

Biotechnology, China) following the manufacturer's instructions. After 48h, firefly and Renilla

luciferase activities were calculated using the Promega Dual-Luciferase system following the

manufacturer's instructions (Hanbio tech, Shanghai, China). Firefly/Renilla luciferase was meas-

ured to evaluate relative luciferase activity.

**Statistical analysis**

Data are reported as the means ± SD. Normal distribution and homogeneity of variance of data

were first tested. Shapiro-Wilk test was used to verify data normality, while Levene's test or

Browne-Forsythe test was adopted for assessment of variance equality. Statistical analysis was

performed by unpaired two $t$-test and Tukey-corrected one-way analysis of variance (ANOVA)

for comparisons between groups. Tukey-corrected two-way ANOVA for the comparison of mice

thermal pain threshold data. $P < 0.05$ was considered statistically significant for all statistical

calculations. PRISM 8.0 (GraphPad Software, San Diego, CA, USA) was used for data analysis.

**Contributions:**

**Contributions:**

F.T., Z.Q., and Q.-F.W. designed the study.

F.T. performed most in vitro and in vivo experiments.

F.T. and J.-M.W. analyzed data.

Z.Q., Y.J. and S.-H.L. performed the miRNA sequencing and interpreted the data.

Z.Q., and Y.J collected and inspected human patient samples.

F.T. and Q.-F.W. wrote the manuscript.

All authors read and edited the manuscript.

F.-R.L., S.-G.Y., and Q.-F.W. supervised the study.

**Competing interests**

The authors declare that they have no known competing financial interests or personal relation-

ships that could have appeared to influence the work reported in this paper

**Acknowledgements:**

This work was supported by the National Key R&D Program of China (No.2019YFC1709001);

Innovation Team and Talents Cultivation Program of National Administration of Traditional

Chinese Medicine (No. ZYYCXTD-D-202003); Fund of Science and Technology Department of

Sichuan Province, China (No. 2021ZYD0081); and National Natural Science Foundation of Chi-

na (No. 82174512, 81873383).

**Data Sharing Statement:**

All data are available in the main text or the supplementary materials.

**List of Supplementary Materials:**

Present a list of the Supplementary Materials in the following format.

Materials and Methods

Fig S1. Identification of exosome. (A) Transmission electron microscope scanning of isolated

exosomes from the serum of participants. (B) The Nano sight particle analysis of isolated exo-

somes from the serum of participants. (C) The Western blot analysis of symbolic surface markers

of isolated exosomes from the serum of participants.

Fig S2. Fluorescence of Adenovirus-Infected Chondrocytes.

Fig S3. Expression of adenovirus in mouse knee joint.

Fig S4. Establishment of KOA model in mice by injection of MIA. (A) Behavioral detection of

animal thermal pain threshold. (B) Microscopic observation of the surface of the mouse knee

joint. (C) Safranin-fast green staining and semiquantitative scoring of articular cartilage. (D) 3D

reconstruction images of joints from μCT scans. *$P<0.05$, **$P<0.01$, *n*=6.

Table S1  Primers list

Table S2 Basic information of participants

**References**

1.      Chen D, Shen J, Zhao W, et al. Osteoarthritis: toward a comprehensive understanding of
pathological mechanism. *Bone research.* 2017;5:16044.
2.      Hunter D, Bierma-Zeinstra S. Osteoarthritis. *Lancet (London, England).*
2019;393(10182):1745-1759.
3.      Ito Y, Matsuzaki T, Ayabe F, et al. Both microRNA-455-5p and -3p repress hypoxia-
inducible factor-2α expression and coordinately regulate cartilage homeostasis. *Nature*
*communications.* 2021;12(1):4148.
4.      Xu W, Yang Z, Lu N. From pathogenesis to clinical application: insights into exosomes
as transfer vectors in cancer. *Journal of Experimental & Clinical Cancer Research : CR.*
2016;35(1):156.
5.      Ali S, Peffers M, Ormseth M, Jurisica I, Kapoor M. The non-coding RNA interactome in
joint health and disease. *Nature reviews Rheumatology.* 2021;17(11):692-705.
6.      Zhao R, Li Y, Lin Z, et al. miR-199b-5p modulates BMSC osteogenesis via suppressing
GSK-3β/β-catenin signaling pathway. *Biochemical and biophysical research*
*communications.* 2016;477(4):749-754.

7. Yamashita S, Miyaki S, Kato Y, et al. L-Sox5 and Sox6 proteins enhance chondrogenic miR-140 microRNA expression by strengthening dimeric Sox9 activity. *The Journal of biological chemistry.* 2012;287(26):22206-22215.

8. Miyaki S, Nakasa T, Otsuki S, et al. MicroRNA-140 is expressed in differentiated human articular chondrocytes and modulates interleukin-1 responses. *Arthritis and rheumatism.* 2009;60(9):2723-2730.

9. Nakamura A, Rampersaud Y, Nakamura S, et al. microRNA-181a-5p antisense oligonucleotides attenuate osteoarthritis in facet and knee joints. *Annals of the rheumatic diseases.* 2019;78(1):111-121.

10. Wu J, Yang F, Wu X, Liu X, Zheng D. Comparison of genome-wide DNA methylation patterns between antler precartilage and cartilage. *Molecular Genetics and Genomics : MGG.* 2023;298(2):343-352.

11. Yang X, Yip J, Harrison M, Brockhausen I. Primary human osteoblasts and bone cancer cells as models to study glycodynamics in bone. *The International Journal of Biochemistry & Cell Biology.* 2008;40(3):471-483.

12. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell.* 2004;116(2):281-297.

13. Kalluri R, LeBleu V. The biology function and biomedical applications of exosomes. *Science (New York, NY).* 2020;367(6478).

14. Sun H, Hu S, Zhang Z, Lun J, Liao W, Zhang Z. Expression of exosomal microRNAs during chondrogenic differentiation of human bone mesenchymal stem cells. *Journal of Cellular Biochemistry.* 2019;120(1):171-181.

15. Ali SA, Peffers MJ, Ormseth MJ, Jurisica I, Kapoor M. The non-coding RNA interactome in joint health and disease. *Nature Reviews Rheumatology.* 2021;17(11):692-705.

16. Jiang Y, Zhang L, Tian H. MicroRNA-149 improves osteoarthritis via repression of VCAM-1 and inactivation of PI3K/AKT pathway. *Experimental Gerontology.* 2023;174:112103.

17. Zhao Z, Zhao S, Luo L, et al. miR-199b-5p-DDR1-ERK signalling axis suppresses prostate cancer metastasis via inhibiting epithelial-mesenchymal transition. *British Journal of Cancer.* 2021;124(5):982-994.

18. Zhang M, Yuan S, Sun H, Sun L, Zhou D, Yan J. miR-199b-5p promoted chondrogenic differentiation of C3H10T1/2 cells by regulating JAG1. *Journal of tissue engineering and regenerative medicine.* 2020;14(11):1618-1629.

19. Qu X, Chen Z, Fan D, et al. MiR-199b-5p inhibits osteogenic differentiation in ligamentum flavum cells by targeting JAG1 and modulating the Notch signalling pathway. *Journal of cellular and molecular medicine.* 2017;21(6):1159-1170.

20. Huang J, Zhao L, Fan Y, et al. The microRNAs miR-204 and miR-211 maintain joint homeostasis and protect against osteoarthritis progression. *Nature communications.* 2019;10(1):2876.

21. Lu Y, Liu Y, Wan F, Shi S, Tao R. CircSmg5 stimulates the osteogenic differentiation of bone marrow mesenchymal stem cells by targeting the miR-194-5p/Fzd6 axis and beta-catenin signaling. *Environmental toxicology.* 2022;37(3):593-602.

22. Peng F, He Q, Cheng C, Pan J. GCNT2 induces epithelial-mesenchymal transition and promotes migration and invasion in esophageal squamous cell carcinoma cells. *Cell biochemistry and function.* 2019;37(1):42-51.

23.   Sherman BT, Hao M, Qiu J, et al. DAVID: a web server for functional enrichment
analysis and functional annotation of gene lists (2021 update). *Nucleic Acids Research.*
2022.
24.   Chen Y, Wang X. miRDB: an online database for prediction of functional microRNA
targets. *Nucleic acids research.* 2020;48:D127-D131.
25.   Dweep H, Sticht C, Pandey P, Gretz N. miRWalk--database: prediction of possible
miRNA binding sites by "walking" the genes of three genomes. *Journal of biomedical*
*informatics.* 2011;44(5):839-847.
26.   Li J, Liu S, Zhou H, Qu L, Yang J. starBase v2.0: decoding miRNA-ceRNA, miRNA-
ncRNA and protein-RNA interaction networks from large-scale CLIP-Seq data. *Nucleic*
*acids research.* 2014;42:D92-97.
27.   Karagkouni D, Paraskevopoulou M, Chatzopoulos S, et al. DIANA-TarBase v8: a
decade-long collection of experimentally supported miRNA-gene interactions. *Nucleic*
*acids research.* 2018;46:D239-D245.
28.   Agarwal V, Bell G, Nam J, Bartel D. Predicting effective microRNA target sites in
mammalian mRNAs. *eLife.* 2015;4.
29.   Pitcher T, Sousa-Valente J, Malcangio M. The Monoiodoacetate Model of Osteoarthritis
Pain in the Mouse. *Journal of visualized experiments : JoVE.* 2016(111).
30.   Choi W-S, Lee G, Song W-H, et al. The CH25H-CYP7B1-RORα axis of cholesterol
metabolism regulates osteoarthritis. *Nature.* 2019;566(7743):254-258.
31.   Cheah M, Fawcett J, Andrews M. Assessment of Thermal Pain Sensation in Rats and
Mice Using the Hargreaves Test. *Bio-protocol.* 2017;7(16).
32.   Glasson S, Chambers M, Van Den Berg W, Little C. The OARSI histopathology
initiative - recommendations for histological assessments of osteoarthritis in the mouse.
*Osteoarthritis and cartilage.* 2010:S17-23.

**Figure Legends**

**Figure 1** MiRNA expression in KOA patients. **(A, B)** Volcano plot and heatmap showing dif-

ferential miRNAs between KOA and healthy groups (healthy, $n = 10$; KOA, $n = 15$, $|$ log2 Fold

Change$| \geq 0.585$, $P < 0.05$). **(C)** GO and KEGG enrichment analyses of upregulated genes. **(D)**

GO and KEGG enrichment analyses of downregulated genes.

**Figure 2** Verification with GEO data and miRNA screening. **(A, B)** Venn plot showing GEO

dataset and our result. **(C)** RT-qPCR in clinical samples to verify the expression (HC, $n = 5$,

KOA, n = 5). **(D)** RT-qPCR results of mouse joint samples ($n = 6$). Data are shown as means $\pm$

SD. *$P<0.05$, **$P<0.01$.

**Figure 3** Chondrocyte proliferation and marker expression changes after miR-199b-5p mimic or inhibitor treatment. **(A)** Second generation primary mouse chondrocytes. **(B)** Toluene blue staining. **(C)** Type II collagen immunoassay. **(D)** CCK-8 assay for cell viability ($n$=6). **(E, F)** RT-qPCR detection of *MMP-3* and *ADAMTS5* mRNA expression ($n$=3). **(G-I)** RT-qPCR detection of *COL-2A1*, *AGGRECAN*, and *SOX9* mRNA expression ($n$=3). **(J)** CCK-8 cell viability assay after different doses of LPS induction ($n$=5). **(K)** CCK-8 cell viability assay after virus infection ($n$=6). **(L, M)** RT-qPCR detection of *MMP-3* and *ADAMTS5* mRNA expression ($n$=3). **(N-P)** RT-qPCR detection of *COL2A1*, *AGGRECAN*, and *SOX9* mRNA expression ($n$=3). Data are shown as means ± SD. *$P$<0.05, **$P$<0.01.

**Figure 4** Injection of adenovirus expressing miR-199b-5p mimic results in inflammation and pain threshold sensitivity in mice. **(A)** Animal experiment schematic. **(B)** Behavioral detection of animal thermal pain threshold. **(C, D)** Detection of serum levels of *IFN-γ* and *TNF-α* in controls by ELISA. **(E, F)** Safranin-fast green staining and semiquantitative scoring of articular cartilage. **(G)** 3D reconstruction and 2D images of joints from μCT scans. Data are shown as mean ± SD. *$P$<0.05, **$P$<0.01, $n$=6.

**Figure 5** Injection of adenovirus expressing miR-199b-5p inhibitor partly recovered pathological changes in KOA mice. **(A)** Animal experiment schematic. **(B)** Behavioral detection of animal thermal pain threshold. **(C, D)** Detection of serum levels of *IFN-γ* and *TNF-α* in controls by ELISA. **(E, F)** Safranin-fast green staining and semiquantitative scoring of articular cartilage. **(G)** 3D reconstruction and 2D images of joints from μCT scans. Data are shown as mean ± SD. *$P$<0.05, **$P$<0.01, $n$=6.

**Figure 6** Validation of the miR-199b-5p target gene. **(A)** Prediction of target genes of miR-199b-5p using search sites. **(B)** Target gene GO analysis. **(C, D)** Detection of the expression of

target genes under different conditions (*n*=3). **(E, F)** Predicting the binding site of miR-199b-5p

and target genes. **(G, H)** Validation by luciferase reporter gene assay (n=3). Data are show as

mean ± SD.\**P*<0.05, \*\**P*<0.01, *n*=6. **(I-L)** The expression of FZD6 and GCNT2 in the synovial

membrane and chondrocytes of GEO Profiles KOA.

**Figure 7** miR-199b-5p exerts its effects on in vitro cells and in vivo mice by targeting *Fzd6* and

*Gcnt2*.

**Supplement table 1: Primer List**

| gene | sequence |
|---|---|
| U6 | GSP:5'GCTTCGGCAGCACATATACTAAAAT3' |
| | R:5'CGCTTCACGAATTTGCGTGTCAT3' |
| hsa-miR-199b-5p | GSP:5'GGGACCCCAGTGTTTAGACTAT3' |
| | R: 5'GTGCGTGTCGTGGAGTCG3' |
| hsa-miR-338-3p | GSP:5'GGGGGTCCAGCATCAGTGA3' |
| | R:5'GTGCGTGTCGTGGAGTCG3' |
| hsa-miR-15b-3p | GSP:5'GGGGCGAATCATTATTTGCT3' |
| | R:5' GTGCGTGTCGTGGAGTCG3' |
| hsa-miR-1296-5p | GSP:5'GGTATTAGGGCCCTGGCTC3' |
| | R:5'GTGCGTGTCGTGGAGTCG3' |
| hsa-miR-3168 | GSP:5'TTGGGGGGGAGTTCTACAG3' |
| | R:5'GTGCGTGTCGTGGAGTCG3' |
| mmu-miR-199b-5p | GSP:5'GGGACCCCAGTGTTTAGACTAT3' |
| | R: 5'GTGCGTGTCGTGGAGTCG3' |
| mmu-miR-15b-3p | GSP:5'GGGGCGAATCATTATTTGCT3' |
| | R:5' GTGCGTGTCGTGGAGTCG3 |
| mmu-miR-338-3p | GSP:5'GGGGGTCCAGCATCAGTGA3' |
| | R:5'GTGCGTGTCGTGGAGTCG3' |
| Gapdh | F: TGTTTCCTCGTCCCGTAGA |
| | R: ATCTCCACTTTGCCACTGC |
| Hif-1α | F: GGGGAGGACGATGAACATCAA |
| | R: GGGTGGTTTCTTGTACCCACA |
| Fzd6 | F: AGAGGTGAAAGCGGACGGA |
| | R: AGAGAGTCTGGAGATGGATGCT |
| Gcnt2 | F: TCCTGGACGGGTAACCTCAG |
| | R: CTGCAAGTCTCCGTTTCCATAG |
| Myh9 | F: AGAAGTTGGTATGGGTGCCTT |
| | R: CCCTGAGTAGTATCGCTCCTTG |
| Caprin1 | F: GAAGCAGATTCTCGGCGTAAT |
| | R: TCCCCTTTATTCATTCGTTCCTG |
| Atg14 | F: GAGGGCCTTTACGTGGCTG |
| | R: AATAGACGAAATCACCGCTCTG |
| Mmp3 | F: ACTGTGTCCCAAGGAGAGGAG |
| | R: AAACCATCTACACAGTTCAGACAC |
| Adamts5 | F: ATGCAGCCATCCTGTTCACC |
| | R: AAGGCCAAGTAGATGCCCAATTT |
| Aggrecan | F: CACTGTCAAAGCACCATGCC |
| | R: TAGGCTGGCTCCCATTCAGT |

|  |  |
|---|---|
| Sox9 | F: TAATTCCCCAGGCTCTTGGAT |
|  | R: GCAGCCGGGATTTAAGGCTC |
| Col2a1 | F: CACGCATGAGCCGAAGCTA |
|  | R: GGGTTTCCACGTCTCACCA |

3 **Supplement table 2: Basic information for recruiting patients**

| **Baseline** | **KOA（N=15）** | **HC（N=10）** | *P* **value** |
|---|---|---|---|
| Age | 52.67 ± 7.35 | 51.50 ± 7.71 | 0.706 |
| Height | 1.63±0.07 | 1.60±0.05 | 0.201 |
| Weight | 65.03±9.53 | 61.30±8.47 | 0.327 |
| BMI | 24.28 ± 2.39 | 23.90± 2.65 | 0.715 |

**Fig 1**

**Fig 2**

Fig 3

**Fig 4**

**Fig 5**

**Fig 6**