1 **Palatal segment contributions to midfacial growth in three inbred mouse strains**

2 Short running page heading: **Palatal segment contributions to midfacial growth**

3 Ian C. Welsh @X – 1) Program in Craniofacial Biology, 2) Department of Orofacial
4   Sciences and 3) Department of Anatomy, University of California at San Francisco, San
5   Francisco, California 94143, USA
6
7 Maria E. Feiler – Interdepartmental Doctoral Program in Anthropological Sciences, Stony Brook
8   University, Stony Brook, NY 11790
9
10 Danika Lipman @Y – Department of Cell Biology and Anatomy, University of Calgary
11
12 Isabel Mormile – Interdepartmental Doctoral Program in Anthropological Sciences, Stony Brook
13   University, Stony Brook, NY 11790
14
15 Karissa Hansen – 1) Program in Craniofacial Biology, 2) Department of Orofacial Sciences and
16   3) Department of Anatomy, University of California San Francisco, San Francisco, CA
17   94143
18
19 Christopher J. Percival* – Department of Anthropology, Stony Brook University, Stony Brook
20   NY 11794
21
22 * = Corresponding Author: Department of Anthropology, Circle Road, SBS Building S-501,
23   Stony Brook, NY 11994, christopher.percival@stonybrook.edu
24 @X Present Address: The Jackson Laboratory, 600 Main St. Bar Harbor, ME 04609
25 @Y Present Address: Department of Mathematics and Statistics, 612 Campus Place NW,
26   University of Calgary, Calgary, AB
27

Abstract

Following facial prominence fusion, anterior-posterior (A-P) elongation of the palate is a

critical aspect of palatogenesis and integrated midfacial elongation. Reciprocal epithelial-

mesenchymal interactions drive secondary palate elongation and periodic signaling center

formation within the rugae growth zone (RGZ). However, the relationship between RGZ

dynamics and the morphogenetic behavior of underlying palatal bone mesenchymal precursors

has remained enigmatic. Our results indicate that cellular activity at the RGZ simultaneously

drives rugae formation and elongation of the maxillary bone primordium within the anterior

secondary palate, which more than doubles in length prior to palatal shelf fusion. The first

formed palatal ruga, found just posterior to the RGZ, represents a consistent morphological

boundary between anterior and posterior secondary palate bone precursors, being found at the

future maxillary-palatine suture. These results suggest a model where changes in RGZ-driven A-

P growth of the anterior secondary palate may produce interspecies and intraspecies differences

in facial prognathism and differences in the proportional contribution of palatal segment-

associated bones to total palate length. An ontogenetic comparison of three inbred mouse strains

indicated that while RGZ-driven growth of the anterior secondary palate is critical for early

midfacial outgrowth, subtle strain-specific bony contributions to adult palate length are not

present until after this initial palatal growth period. This multifaceted illustration of normal

midfacial growth dynamics confirms a one-to-one relationship between palatal segments and

upper jaw bones during the earliest stages of palatal growth, which may serve as the basis for

evolutionary change in upper jaw morphology. Additionally, identified mouse strain-specific

differences in palatal segment elongation provide a useful foundation for understanding the

impact of background genetic effects on facial morphogenesis.

Keywords: palatogenesis, secondary palate, rugae, inbred mouse strains, face elongation,

RRID:IMSR_JAX:003715, RRID:IMSR_JAX:001976, RRID:IMSR_JAX:000664

Introduction

Morphological variation of the midfacial complex, which consists of the nose, upper jaw, cheek, and palate, is a defining aspect of both intra- and inter-specific differences in facial shape. Evolutionary differences in midfacial prognathism can be quantified as variation in upper jaw bone elongation and in the proportional contribution of specific bones to total upper jaw length. Species-specific differences in the size, shape, and growth trajectories of cranial neural crest (CNC)-derived facial prominences, the frontonasal process (FNP) and maxillary processes (MxP), are identifiable during the earliest phases of facial morphogenesis, prior to skeletal differentiation, often presaging interspecies differences in adult facial shape (Selleri and Rijli, 2023). Notably, after initial facial prominence fusion, early facial growth often includes substantial anterior-posterior (A-P) midfacial elongation, leading to variable degrees of midfacial prognathism (Young et al., 2014); possibly based on the balance of CNC progenitor self-renewal versus osteoblast differentiation (Hall et al., 2014; Morris and Abzhanov, 2021; Schneider, 2015). Although characteristic interspecies differences in craniofacial morphology are often present at birth (Ponce de León and Zollikofer, 2001; Zumpano and Richtsmeier, 2003; Cobb and O'Higgins, 2004; Singleton, 2012; Smith et al., 2015; Woronowicz and Schneider, 2019; Kundrát et al., 2020), postnatal bone deposition and remodeling are also required to produce the highly differentiated adult morphology, which often includes more extreme midfacial prognathism (Enlow and Bang, 1965; Sarnat, 1997; Cobb and O'Higgins, 2004; Leigh, 2006;

Martinez-Maza et al., 2013; Maga, 2016). Thus, both embryonic and postnatal developmental

processes play a role in determining adult upper jaw size and shape.

Basic facial shape is the result of facial prominence outgrowth and a series of fusion

events, which occur between embryonic day (E) 10 and E15 in mice (Fig. 1). Initial fusion

between the medial nasal processes (mnp) of the FNP and the anterior MxP gives rise to the

primary palate and lip while also producing a unified upper jaw from previously separated

tissues. Disproportional growth of FNP or MxP prior to fusion can result in cleft lip and palate,

leading to reduced evolutionary fitness. Conversely, differential growth of FNP and MxP derived

tissues after facial prominence fusion is a major basis for evolutionary differences in midfacial

elongation; where facial elongation is driven primarily by MxP growth in mammals and by FNP

growth in birds (Young et al., 2014).

Figure 1 – Tissue origins of the midfacial complex and rugae position during secondary palate morphogenesis – The

upper and lower jaws are formed from the frontonasal process (FNP - tan), and branchial arch 1 derived maxillary

and mandibular processes (MxP - pale blue and Mnd – magenta, respectively). From E11.5 to E15.5, outgrowth and

fusion of the medial nasal process (mnp) with the superficial portion of the MxP frame out the lip and primary (1°)

palate, while the A-P elongation and medially directed growth of the palatal shelves from the internal portion of the

MxP gives rise to the secondary palate (2° pal – light purple). *Shh* expression (dark blue) highlights the dynamics of

rugae formation and illustrates regional expansion of the anterior secondary palate. At E11.5, ruga 1 (red

arrowheads) forms at the anterior extent of the nascent palatal shelf and subsequently defines the caudal end of the

rugae growth zone (asterisk) where new rugae form prior to being displaced anteriorly. Additional abbreviation:

lateral nasal process (lnp).

Soon after primary palate fusion at E11.5 in mice, secondary palate morphogenesis

begins through outgrowth of the nascent palatal shelf along the medial aspect of the MxP.

Significant palatal growth along the A-P axis is accompanied by vertical growth of the palatal

shelves prior to their elevation and medial fusion dorsal to the tongue, after which the palatal

shelf separates the oral and nasal cavities (Bush and Jiang, 2012; Hammond and Dixon, 2022). A

lack of coordination between the multiple growth axes of MxP-derived tissues can prevent

secondary palate closure even when the palatal shelves maintain competency to fuse.

The mammalian palate is composed of three major longitudinal segments called the

primary palate, anterior secondary palate, and posterior secondary palate. The boundary between

primary palate and anterior secondary palate is the posterior edge of the fused mnps. The A-P

organization of the secondary palate can be defined by various criteria including the presence of

boney versus muscular tissue (i.e. the hard versus soft palate), modes of palatal shelf elevation

and closure, and molecular differences in tissue patterning and cell signaling competence

(Hilliard et al., 2005; Yu and Ornitz, 2011; Hammond and Dixon, 2022). We use mutually

exclusive patterns of gene expression and the position of epithelial appendages called rugae to

define the boundary between anterior secondary and posterior secondary palate, as described

below.

From a morphogenetic regulatory perspective, the spatial expression of cell type specific transcription factors (TFs) and intercellular signaling molecules is particularly informative to understanding A-P organization of palate growth. For example, the CNC-derived mesenchymal expression of *Msx1* and *Shox2* is exclusive to the anterior secondary palate during palate morphogenesis, while *Barx1* and *Tbx22* expression is exclusive to the posterior secondary palate (Hilliard et al., 2005; Okano et al., 2006; Welsh et al., 2018). Mutation of *Shox2* or *Tbx22* results in midline clefts of the anterior or posterior palate, respectively (Yu et al., 2005; Pauws et al., 2009), although normal expression of these factors is maintained in genetically engineered mice with severe segmental growth defects (Welsh et al., 2018). The A-P expression boundary of these and other factors is found at the first formed palatal ruga (Hilliard et al., 2005; Li and Ding, 2007; Welsh et al., 2007; Welsh and O'Brien, 2009; Welsh et al., 2018; Hammond and Dixon, 2022), one of multiple parallel epithelial thickenings on the anterior secondary palate.

The first ruga (ruga 1 numbered according to (Welsh et al., 2007; Welsh and O'Brien, 2009, although see Pantalacci et al., 2008) forms at E11.5 on the anterior extent of the nascent secondary palate (Fig. 1). Remaining rugae form sequentially within the rugae growth zone (RGZ) located just anterior to ruga 1 (Welsh and O'Brien, 2009), via a Turing type mechanism (Economou et al., 2012; Kawasaki et al., 2018). Rugae are centers of Sonic Hedgehog signaling (SHH) that is critical for anterior secondary palatal patterning and A-P growth. As the secondary palate completes midline fusion at E15.5, the last formed ruga appears adjacent to ruga 1 in the middle palate while each precedingly formed ruga is found at an increasingly anterior position along the anterior secondary palate.

Based upon the dynamics of rugae formation, palate growth, and patterning gene expression, Pantalacci (et al., 2008) hypothesized that ruga 1 represents a distinct morphological

boundary related to either the future hard versus soft palate boundary or the maxillary-palatine

suture. However, the dynamics of rugae formation and A-P patterns of molecular and

morphological variation in the secondary palate have not been explicitly defined in relation to the

earliest stages of facial bone development. One goal of our research was to further clarify the

degree to which early ontogenetic processes in the palate are coordinated with skeletal

differentiation. To address this knowledge gap, we defined the anatomical and molecular

relationship of the anterior and posterior secondary palate during anterior secondary palate

outgrowth with reference to boundary ruga 1, the rugae forming RGZ, and the broader context of

midfacial anatomical structures. We systematically mapped the initial specification and early

differentiation of osteoblasts relative to this anatomically defined segmental organization across

a developmental time course spanning mouse secondary palate morphogenesis and upper jaw

outgrowth (E11.5-E15.5). Visualization of molecular markers during the earliest phases of

palatal morphogenesis reveals the segmental origins of upper jaw bone formation and a

previously unappreciated coupling of maxillary formation to rugae morphogenesis during

expansion of the anterior secondary palate.

Given that 1) there is strong early one-to-one association between upper jaw bones and

the three major palatal segments (primary, anterior secondary, posterior secondary), 2)

morphogenic and differentiative dynamics are distinct along the A-P axis of the elongating

midface, and 3) complex embryonic processes at cellular and tissue levels are often critical for

the production of typical facial shape (Hilliard et al., 2005; Kaucka et al., 2018; Yuan et al.,

2020), early differences in the longitudinal growth of the three palatal segments might underlie

inter- and intraspecies differences in upper jaw morphology. Specifically, we hypothesized that

genetically based differences in palate segment-specific A-P outgrowth would explain previously

identified variation in the contribution of the premaxilla, maxilla, and palatine bones to adult

upper jaw morphology of inbred mouse strains (Percival et al., 2016). For example, a minor

segment-specific reduction in the growth rate of the anterior secondary palate between E11.5 and

E15.5 could result in a relatively short anterior secondary palate and relatively short adult

maxillary bone, as identified within the C57BL/6J (C57; RRID:IMSR_JAX:000664) strain.

To test our hypothesis and shed light on the overall developmental basis of adult palatal

morphology, we measured the facial morphology of an ontogenetic series of common inbred

strains of mice between E11 and E15, one day after birth, and in adults (8-12 weeks old). The

C57 and NOD/ShiLtJ (NOD; RRID:IMSR_JAX:001976) inbred strains were chosen because

C57 adults have a relatively long premaxilla and NOD adults have a relatively long maxilla

(Percival et al., 2016). The PWK/PhJ (PWK; RRID:IMSR_JAX:003715) wild-derived inbred

strain was chosen as a mouse with a smaller adult body size, a relatively long maxilla compared

to the premaxilla, and a relatively long upper jaw for its body size when compared to the other

strains (Percival et al., 2016). If variation in early A-P elongation of longitudinal palatal shelf

segments is the causal basis for strain specific differences in upper jaw morphology, we

predicted that C57 mice would have a relatively long primary palate and a relatively short

anterior secondary palate by E15 and that the other two strains would display relatively long

anterior secondary palates at the same developmental stage.

Understanding how genetically based differences in early craniofacial development

produce morphological variation within model organisms is an important step in determining

how these developmental processes can be modified to produce evolutionary differences in upper

jaw morphology (Boughner et al., 2008; Hallgrímsson et al., 2009; Fish, 2019; Morris and

Abzhanov, 2021), including facial length and the relative contribution of specific bones to the

upper jaw. The subtle strain-associated differences in palatal morphology we report are

reminiscent of brachycephalic versus dolichocephalic dog breeds and species-level differences

between primates, including the reduced facial length of humans compared to other apes and the

increased facial length of baboons compared to other cercopithecines. Overall, our multifaceted

analysis of ontogenetic trends in palatal and facial elongation provides a novel contextual

framework and developmental perspective within which to evaluate the impact of both non-

pathogenic as well as pathogenic genetic differences on midfacial growth and differentiation.

Methods

Specimen and Image Acquisition

Animal breeding, specimen collection, and tissue fixation were performed in accordance

with the protocols of the University of California, San Francisco Institutional Animal Care and

Use Committee under protocol approval number AN192776-01F. Mice were socially housed

under a twelve-hour light-dark cycle with food and water *ad libitum*. Additional enrichment was

provided when single housing was required for breeding purposes. Mice were euthanized by CO_2

inhalation followed by cervical dislocation or decapitation. Embryos of C57BL/6J

(RRID:IMSR_JAX:000664; C57), NOD/ShiLtJ (RRID:IMSR_JAX:001976; NOD), and

PWK/PhJ (RRID:IMSR_JAX:003715; PWK) strains (Jackson Labs, Bar Harbor, ME) were

collected between gestational days E11.5 and E15.5, as determined from copulatory plug

occurrence. Postnatal day one (P1) C57 (n=10) and NOD (n=7) specimens were collected one

day after birth. Embryo and P1 specimens were fixed in 4% PFA and stored in 1x PBS for

micro-computed tomography (μCT) imaging or dehydrated through a graded PBST-MeOH

series and stored at -20°C until use for in situ hybridization.

Specimens for μCT scanning were received, stored, and imaged at the University of

Calgary in accordance with the protocols of the University of Calgary Institutional Care and Use

Committee under approval number AC13-0268. After approximately an hour of soaking in

Cysto-Conray II (Liebel-Flarsheim Canada), μCT images of embryo heads were acquired with a

Scanco μ35 with 45kV/177μA for images of 0.012 mm^3 voxel size. μCT images of P1 heads

were acquired similarly, but with 0.021 mm^3 voxel size. Photographs of embryo hindlimb buds

were collected using a dissecting microscope for developmental age estimation. Adult specimens

were previously collected and μCT imaged, as described by Percival et al., 2016.

In Situ Hybridization

In situ hybridization was performed as described in Welsh et al., 2018. cDNA probes for

Phospho1 (nucleotides 274–1739 of RefSeq NM_153104), *Runx2* (nucleotides 1275–2704 of

RefSeq NM_001271627), *Shh* (full length cDNA clone from A. McMahon, University of

Southern California, Los Angeles CA, USA), *Shox2* (nucleotides 870–1603 of RefSeq

NM_130458), *Sp7* (nucleotides 431–1715 of RefSeq NM_130458), *Tbx22* (nucleotides 307–

1547 of RefSeq NM_013665) were linearized and in vitro transcribed to label with either

digoxygenin (DIG) or dintrophenol (DNP). Colorimetric detection of probes used BM purple

(dark blue), BCIP (teal), or MagentaPhos (magenta).

DMDD HREM Data

High resolution episcopic microscopy (HREM) data (Geyer et al., 2009) was generated as

part of the deciphering mechanisms of developmental disorders (DMDD) program (Wilson et al.,

2016) and is available from the BioImage Archive at:

https://www.ebi.ac.uk/biostudies/bioimages/studies/S-BIAD490?query=DMDD

Embryonic Palatal Segment Growth Trajectories

<u>Developmental Age Estimation</u>

Given that common mouse strains vary in gestation length and there is variation in

developmental timing within litters, it was necessary to standardize our cross-strain

morphometric analysis by embryonic developmental age rather than gestational age.

Developmental age was estimated for each μCT scanned embryonic specimen using eMOSS, an

application that predicts developmental age from hindlimb bud outlines, based on a previous

analysis of C57 mice (Musy et al., 2018). The resulting limb-based estimates of developmental

age were reported as days since conception, up to two decimal places. We combined similar

developmental age estimates into developmental age categories so that specimens were assigned

to a whole- or half-day developmental age within 0.25 days of their initial eMOSS estimate

(Table 1). For example, an eMOSS estimated E12.20 specimen was assigned to the E12.0 limb-

based developmental age category.

Table 1 –Sample size of μCT imaged and morphometrically analyzed embryo specimens by strain and

forelimb based developmental age estimation, as estimated using eMOSS (Musy et al., 2018).

Limb-Based Developmental Age	C57	NOD	PWK
E11	8	12	4
E11.5	2	0	0
E12	7	3	6
E12.5	4	0	2
E13	9	3	6
E13.5	0	0	2
E14	26	6	6
E14.5	1	0	0
E15	9	4	0

Additional error may be expected when estimating the developmental age of NOD and

PWK embryos using a predictive model built on C57 strain limb bud ontogeny, given the

possibility of differences in the correlation of limb and head development between strains.

Therefore, we also utilized the first principal component of embryonic head shape variation

(PC1), based on the landmark analysis described below, as a proxy for developmental age in

some analyses. PC1 score consistently tracked ontogenetic change in facial and palatal shape

across strains (see Results).

<u>Anatomical Landmark Collection</u>

All embryo facial and palate landmarks were collected within Meshlab (Cignoni et al.,

2008) on minimum threshold-based superficial tissue surfaces (downsampled x2) produced from

the μCT images. Landmarks that could be identified consistently in anatomically homologous

positions across the embryonic period of facial growth were chosen around the nose and whisker

region, the eyes, and along the palate (Fig. 2, Supporting Table 1). Care was taken to define

epithelial tissue palatal landmarks in embryos that could be associated with the bony morphology

of the P1 and adult mouse upper jaw.

Figure 2 – Anatomically homologous epithelial landmarks collected for the quantification of midfacial and palatal

shape across limb based developmental stages, from (a-c) palatal and (d-f) oblique facial views. See also Supporting

Table 1.

<u>Prenatal Geometric Morphometric Analysis</u>

A Procrustes superimposition-based geometric morphometric analysis was used to

quantify the ontogenetic shape change of the palate and face for the embryonic C57 sample and

deviations of the NOD and PWK strains from this C57 baseline. Embryonic specimens with

limb-based developmental age estimates between E11 and E15 were included in the embryonic

shape analysis (Table 1). We performed a geometric morphometric analysis of facial landmarks

using *geomorph* (Adams et al., 2020) and RRPP (Collyer and Adams, 2018) libraries in R

Statistical Software (R Core Team, 2021). We first performed a generalized Procrustes analysis

(GPA) to align specimen landmark sets by translating, scaling, and rotating their landmark

coordinates (reviewed by Zelditch et al., 2012). Embryonic specimen shape analyses were

completed using the symmetric component of Procrustes-aligned specimen landmark coordinate

variation, as we assumed that most bilateral shape differences between the left and right sides of

a specimen's face are due to random effects associated with developmental noise and tissue

fixation (Palmer and Strobeck, 1986). Thus, symmetrized landmark coordinate data were

interpreted to represent the facial shape defined by a given inbred strain genotype.

The mean facial shape of specimens within each developmental age category were

estimated for each strain. Differences between the mean shapes of a subset of developmental age

categories within the C57 sample were plotted to illustrate typical facial/palatal shape growth

trajectories during this important period of secondary palate elongation and midline fusion.

A principal components analysis (PCA) was completed to identify the major axes of

shape covariation across the embryonic sample. As with most PCA of ontogenetic series, the first

principal component (PC1) was strongly associated with overall specimen size and specimen

developmental age. Therefore, we treated it as a proxy for ontogenetic growth and

developmental age. C57-specific segmented linear regressions represented a baseline of overall

facial/palatal shape change to which the NOD and PWK specimens were compared. Due to clear

inflection points in the slopes of embryonic specimen PC scores when plotted against PC1,

segmented linear regressions were completed separately for PC2, PC3, and PC4 versus PC1

values for the C57 sample.

After removing three NOD specimens with PC1 scores that were substantially lower than

the minimum C57 PC1 score that the C57 regression was based on, we predicted PC2, PC3, and

PC4 scores from each C57, NOD, and PWK specimen's PC1 score based on the C57 baseline

segmented regressions. Then we calculated the PC score residuals of each specimen as the

difference between predicted and measured PC scores. We completed Wilcoxon rank sum tests

to identify significant genotypic differences for PC2, PC3, and PC4 residual values. A significant

difference in mean PC score residual values between strains was interpreted as a significant

difference in the ontogeny of facial shape for that pair of strains.

Prenatal Palate Segment Length

The anterior-posterior length of the three major palatal segments (i.e., primary palate,

anterior secondary palate, and posterior secondary palate) were estimated from landmark

coordinates without any superimposition or scaling. To estimate the length of these segments

along the anterior-posterior axis of the palate, the mean position of bilateral pairs of palatal

landmarks were calculated (i.e., the midline projection of the bilateral landmarks). Primary

palate, anterior secondary palate, and posterior secondary palate lengths were calculated in

millimeters from these landmark positions. Proportional palatal segment lengths were calculated

as the length of a single segment divided by the sum of all three segment lengths for a given

specimen. LM15 represented the midline anterior extent of the primary palate, the midpoint

between LM18 and LM22 represented the border between primary palate and the anterior

secondary palate, the midpoint between LM20 and LM24 represented the border between the

anterior and posterior secondary palate, the midpoint between LM21 and LM25 represented the

posterior boundary of the posterior secondary palate.

Small strain-specific sample sizes at multiple limb-based developmental ages meant it

was not possible to statistically test whether the genetic strains had different proportional palatal

segment contributions at single developmental ages. Instead, separate second-degree polynomial

regressions of proportional palate segment lengths versus specimen facial shape PC1 score were

calculated for the C57 embryos. Because PC1 score was considered a proxy for developmental

age, these regressions define ontogenetic trajectories of palatal segment contributions to overall

palatal length, which we used as a baseline for comparison with NOD and PWK proportional

segment lengths. After removing three NOD specimens with PC1 scores that were substantially

lower than the minimum C57 PC1 score, we predicted the proportional length of each palatal

segment for all specimens based on the C57 regressions. The residual differences in proportional

palatal segment lengths between predicted and measured values were compared across the three

strains using Wilcoxon rank sum tests. A significant difference in mean residual values was

interpreted to indicate a generally higher or lower proportional contribution of a given palate

segment to overall palatal length across the embryonic period being measured.

Postnatal Palate Segment Length Comparisons

To determine if embryonic differences in proportional palate segment length remained

consistent postnatally and to investigate the degree of correspondence between surface epithelial

and skeletal measures of palatal variation, postnatal specimen landmarks were collected on

minimum threshold-based surfaces produced from μCT scans. Superficial epithelial surfaces of

P1 specimens were produced using Amira (downsampled x2), and epithelial landmarks

(Supporting Table 1) were collected within Meshlab (Cignoni et al., 2008).

Minimum threshold-based skeletal surfaces of P1 and adult specimens were produced

using 3D Slicer (Fedorov et al., 2012) after Gaussian blur image filtering (sigma set to 0.01 for

P1; sigma set to 0.02 for adult). We identified skeletal anatomical landmarks that most closely

matched the palate segment landmarks defined on surface epithelium (Supporting Table 2).

Landmarks were collected for C57 (n=10) and NOD (n=8) P1 specimens and for C57 (n=20),

NOD (n=18), and PWK (n=18) adult specimens. Midline palate segment lengths and associated

upper jaw bone lengths were measured using landmark coordinates without scaling or

superimposition. Midline points were calculated from pairs of bilateral landmarks defining each

palate segment boundary, midline projected palate segment lengths were estimated for each

specimen, and total palate length was calculated as the sum of the three midline palate segment

lengths. Proportional palate segment lengths were calculated as individual segment lengths

divided by total palate length. Wilcoxon rank sum tests were used to identify significant

differences in proportional palate segment lengths between strains within single age and tissue

type categories.

<u>Results</u>

Defining an Anatomical and Molecular Reference Framework of Secondary Palate A-P

Growth

363   We defined the anatomical relationship between the boundary ruga 1, the surrounding

midfacial anatomy, and the A-P expression of genes critical for palatal shelf patterning as a

foundation for assessing rugae morphogenesis and segmental growth dynamics of the secondary

palate. A-P secondary palate elongation was defined with reference to the first ruga's position at

the boundary between the anterior and posterior secondary palate and the A-P expression of

genes critical for palatal shelf patterning, including *Shox2* and *Tbx22*. Ruga 1 formed within the

MxP at the anterior extreme of the nascent secondary palate (Fig. 3, red arrowheads). Expression

of *Shox2* overlapped the medial portion of the *Shh* expression domain in forming ruga 1, while

Tbx22 palatal shelf expression was found just posterior to ruga 1. Sequential rugae formation

occurred anterior to ruga 1 within the RGZ and accompanied expansion of the anterior secondary

palate. *Shox2* expression remained restricted to the rugae-bearing anterior secondary palate,

while *Tbx22* expression was restricted posterior to ruga 1 throughout this critical period of

secondary palate elongation (Fig. 3a,c,e).

Figure 3 – A-P molecular heterogeneity and anatomical relationships during secondary palate morphogenesis. (a, c,

e) Whole mount in situ hybridization for *Shh*, *Shox2*, and *Tbx22* expression, and (b, d, f) sagittal sections of wildtype

embryos imaged with high resolution episcopic microscopy to provide histological resolution of the primary (1°

pal), anterior secondary (ant 2°), and posterior secondary palate (post 2°) relative to surrounding facial structure

during midfacial outgrowth at (a, b) E11.5 (c, d) E13.5, and (e, f) E14.5. Throughout midfacial outgrowth, ruga 1

(red arrowheads) marks the shared A-P expression boundary of *Shox2* and *Tbx22* in the anterior and posterior

secondary palate, respectively. Double label WISH for *Shox2* (magenta) and *Tbx22* (blue) at E14.5 highlights

mutually exclusive anterior and posterior expression domains organized relative to ruga 1 (oral view left, sagittal

view right). White arrowheads mark the choanae, black asterisk marks the RGZ. Red dashed lines mark the primary-

secondary palate junction and posterior wall of the nasal capsule to highlight coordinated elongation of the anterior

secondary palate and overlying sinus cavity (sc). Regions in dashed boxes of HREM images shown enlarged to the

right. Additional abbreviations: mandible (mnd), molar tooth bud (mtb), maxillary process (MxP), medial nasal

process (mnp), tongue (t), greater palatine nerve (gpn), palatine (p).

Close relationships between different anatomical domains underlying midfacial

outgrowth are found centered at ruga 1, which is at the boundary between the anterior and

posterior secondary palate. Elongation of the anterior secondary palate is coordinated with the

formation of the overlying sinus cavity, and disrupted morphogenesis of either structure can

impact the other (Pauws et al., 2009; Kaucka et al., 2018; Welsh et al., 2018). We found that

ruga 1 formed adjacent to the choanae, the posterior openings of the forming sinus cavity (Fig.

3a, white arrowheads). Expansion of the secondary palate anterior to ruga 1 between E11.5 and

E15.5 anteriorly displaced the primary palate and external nares. Ruga 1 and the posterior wall of

the nasal capsule remained approximately in the same coronal plane during elongation of the

anterior secondary palate and nasal capsule (Fig.3b,d,f), suggesting that they are positioned at a

proximal source of directional growth. Ruga 1 was also coincident with a gross morphological

inflection point of the palatal tissue where the anterior secondary palate and overlying sinus

cavity exhibited a more inferiorly accentuated angle relative to the cranial base than the posterior

secondary palate did (Fig. 3d-e, Supporting Fig. 1). Lastly, between E12.0 and E12.5, the greater

palatine nerve and artery were found entering the palatal shelves from a location immediately

dorsal to ruga 1. This spatial relationship was maintained until the greater palatine foramen

forms prior to E15.5 by the articulation of maxilla and palatine bones (Fig. 3f, Supporting Fig.

1). These anatomical relationships collectively demonstrated that ruga 1 represents an important

stable anatomical boundary and point of reference within the context of early midfacial

outgrowth and segmental organization of the palate.

We conducted a complementary 3D morphometric analysis of epithelial landmarks across

the midface, including several defined by the palatal anatomy described above, to quantify the

integrated morphogenetic processes occurring across the facial region during this critical early

developmental period. The first major principal axis of facial shape variation (PC1) for all limb

bud staged E11-E15 specimens represented the majority of shape variance present in the sample

(70%) and was associated with general ontogenetic growth during this time window, as

illustrated for the C57 sample (Fig. 4) and for all three inbred strains (Supporting Fig. 2).

Additional principal axes of variation suggested shifts in the direction of ontogenetic shape

changes over developmental time. For example, the PC2 scores of C57 embryos increased from

E11 to E12.5, but then decreased again from E12.5 to E15 (Fig. 4a). The direction of shape

change shifted twice along PC3 (Fig. 4b) and three times along PC4 (Fig. 4c).

Figure 4 – Major axes of embryonic facial shape variation - C57 embryo specimens plotted along the (a) first and

second principal component axes of shape (i.e., PC1 and PC2), (b) along PC1 and PC3, and (c) along PC1 and PC4.

Circle color indicates the limb based developmental age categories of each specimen, with the yellow bordered

circles indicating the mean PC scores for the C57 mice of that age category. The proportions of facial shape variance

associated with each principal component are provided. The principal component axes were estimated from the full

sample of E11-E15 developmental age embryos, including C57, NOD, and PWK samples, although only C57

samples are plotted in this figure. The same plot, but incorporating all three inbred strains is available as Supporting

Figure 2.

Average developmental age category shape differences were illustrated as landmark

specific ontogenetic shifts in palate shape (Fig. 5) and overall midfacial shape (Supporting Fig.

3). Mouse embryos grew substantially between limb-based developmental ages E11 and E15,

including increased facial size and oral cavity length. Within C57 mice at E11, the mnp and the

MxP fused to create the primary palate, while the cell populations that will give rise to the

posterior secondary palate were already visible. Compared to E11, E12.5 specimens had a

relatively longer primary palate and a relatively short distance between the left and right

secondary palatal shelves. After E12.5, secondary palate length (relative to overall facial size)

increased noticeably with each developmental stage. As the specimens approached E15, the most

substantial changes in palatal shape were related to 1) the previously described A-P elongation of

the secondary palate and 2) the medial growth and ultimate fusion of the palatal shelves.

Figure 5 – C57 palatal landmark growth trajectories – The positions of palatal landmarks are identified on

representative C57 specimens at limb-derived developmental ages (a) E11, (b) E12.5, (c) E14, and (d) E15. (e) The

average palatal landmark positions are plotted for each developmental age category that had more than one C57

specimen. These landmark positions represent palatal shape after the removal of overall facial scale. The arrows

indicate the trajectory of shape change for each landmark between ages. Black=E11; Light Blue = E12; Orange =

E12.5; Green = E13; Dark Blue = E14; Red = E15.

Between developmental stages E11 and E15, each of the three major palatal segments

grew in length, but the anterior secondary palate grew proportionally more than the primary

palate and posterior secondary palate (Figs. 3 & 5). Midline epithelial measurements of C57

embryos soon after the start of A-P palatal elongation indicated that the primary palate

represented ~20% of the total palate length, the anterior secondary palate ~20%, and the

posterior secondary palate ~60%. In contrast, by the time of midline palatal fusion at E15.5, the

primary palate represented ~25% of palate length, the anterior secondary palate represented

>40%, while the posterior secondary palate represented <35%. Therefore, between E11 and E15,

the proportional contribution of the primary palate increased by a quarter, while posterior

secondary palate contribution decreased by nearly a third. Strikingly, the proportional

contribution of the anterior secondary palate doubled during this period.

Segmental Differences in Skeletal Specification, Differentiation, and Growth During

Secondary Palate Morphogenesis

Cellular changes in gene expression represent one of the most proximal steps in the

morphogenetic cascade of osteogenic differentiation and bone formation. We therefore sought to

evaluate the earliest stages of midfacial skeleton specification and differentiation in relation to

the pattern of segmental growth described above. We analyzed the expression of *Runx2*, *Sp7*, and

Phospho1 using whole mount in situ hybridization (WISH) on mouse embryos from E11.5 to

E15.5, a developmental time course spanning the emergence of the secondary palate until

midline palatal shelf fusion. These molecular markers were chosen because *Runx2* expression

overlaps the initial specification of osteoprogenitors during intramembranous bone formation;

subsequent expression of *Sp7* is required for osteoblast precursor differentiation (Galea et al.,

2021; Hojo and Ohba, 2022); and the phosphatase *Phospho1* is essential for inorganic phosphate

incorporation and mineral deposition in the forming osteoid of differentiating osteoblasts (Dillon

et al., 2019).

Comparing the spatial expression dynamics for these marker genes showed that while

Runx2 signal does highlight some osteogenic domains, particularly in the premaxilla within the

primary palate, it tended to be more broadly expressed within the early facial prominences.

Within the developing secondary palate, expression of *Runx2* remained medial throughout this

period, being restricted to domains adjacent to the molar tooth bud and medially along the

adjoining palatal shelves (Fig. 6a).

Figure 6 – Segmental organization of skeletal specification and growth during A-P morphogenesis of the midface –

WISH time course for (a) *Runx2*, (b) *Sp7*, and (c) *Phospho1* between E11.5 and E15.5. Expression of *Runx2* in

osteoprogenitors highlights facial domains with osteogenic potential during midfacial morphogenesis. Expression of

Sp7 and *Phospho1* in committed and differentiating osteoblasts delineates the growth dynamics of individual

skeletal anlagen during midfacial morphogenesis. Abbreviations: maxilla (mx), palatine (p), premaxilla (pmx),

pterygoid (ptg), palatal process of the maxilla (ppmx), palatal process of the palatine (ppp)

The osteoblast-specific expression of both *Sp7* and *Phospho1* more precisely delineated

the morphology of the emerging midfacial skeleton during this early period of midfacial

outgrowth. At E11.5, *Sp7* was strongly expressed in bilateral domains within the mnp of the

forming primary palate. Between E12.5 and E15.5, presaging premaxilla morphology, *Sp7*

expression formed a cup-like region surrounding the developing incisors (Fig. 6b). At E11.5, *Sp7*

expression in the MxP was observed in two domains. The anterio-lateral domain, interpreted as

the presumptive maxilla, abutted the point of fusion with the mnp and extended laterally away

from the oral cavity and choanae. The posterior domain spanned the length of the forming

secondary palate, from the anterior limit abutting the choanae to the posterior-most palatal edge,

and likely comprised the presumptive palatine and pterygoid bones. The antero-lateral domain of

Sp7 maintained proximity to the primary palate throughout the growth period, formed a new

medial expansion (E12.5-E13.5) that extended posteriorly (E13.5-E15.5) within the growing

anterior secondary palate (Fig. 6b). Between E12.5 and E15.5, anterior secondary palate

elongation displaced the original posterior *Sp7* domain away from the primary palate. This

posterior domain separated into two subdomains; the anterior subdomain remained associated

with the posterior wall of the sinus cavity and gave rise to the palatine bone, while the posterior

subdomain formed the pterygoid. Likely reflecting a temporal lag between initial osteoblast

commitment and later differentiation, *Phospho1* was not detected in the midface at E11.5 and

was observed only in the forming premaxilla at E12.5, followed by expression in progressively

more posterior bone anlagen between E13.5 and E15.5 (Fig 6c).

These osteoblast-specific expression patterns indicated that the premaxilla, palatine, and

pterygoid anlagen grew to acquire their characteristic adult morphology within the same domains

where they were initially specified. However, the maxillary anlage was initially specified

external to the oral cavity at the site of the future zygomatic plate. Only after anterior secondary

palate elongation was initiated (E11.5-12.5) does a new portion of the maxillary anlage grow

medially into the anterior secondary palate (E12.5-13.5) and extend posteriorly towards the

palatine bone anlage (E13.5-15.5). The fact that *Phospho1* expression closely followed *Sp7*

spatial dynamics within the maxillary anlage further supported the idea that *Sp7* expression

patterns represent bone anlagen formation and expansion. Lastly, it is notable that the palatal

processes of the maxilla and palatine bones formed within the medial region of the elevated

palatal shelves, a domain that maintained high levels of *Runx2* expression during secondary

palate growth (Fig. 6a), suggesting maintenance of a less differentiated osteoprogenitor

population until palatal fusion was complete.

Rugae morphogenesis provided a set of temporally ordered anatomical features that can

be used to visualize A-P growth dynamics of the secondary palate. WISH double labeling of *Shh*

and *Sp7* between E12.5 and E15.5 indicated the position of specific rugae relative to developing

bone anlagen (Fig. 7a). These data further demonstrated that the initial site of maxillary

specification (i.e. the future zygomatic plate) maintained a position external to the oral cavity and

adjacent to the anterior most rugae during secondary palate elongation (Fig. 7a). Notably, medial

maxillary anlage expansion into the secondary palate occurred around E13.5, after initial A-P

growth of the anterior secondary palate and formation of 3-4 rugae anterior to ruga 1, and thus

uniquely coupled to RGZ growth dynamics. The maxillary anlage grew posteriorly towards ruga

1 and the palatine anlage at the same time as subsequent rugae formation (E13.5-E14.5; Fig. 7a).

Figure 7.

A

E12.5 **E13.5** **E14.0** **E14.5** **E15.5**

Shh/Sp7

2->3 rugae   4->5 rugae   5->6 rugae   6->7 rugae   8 rugae

B — summary of A-P palate morphogenesis and growth of skeletal anlagen

E11.5 **E12.5** **E13.5** **E14.5** **E15.5**

C — surface landmarks and expansion of A-P domains

skeletal anlage/palate segment

- premaxilla/ 1⁰ palate
- maxilla/ anterior 2⁰ palate
- palatine, pterygoid/ posterior 2⁰ palate

Figure 7 – Double label RNA WISH time course for *Shh* (cyan) and *Sp7* (dark purple) between E12.5 and E15.5. (a)

Expression of *Sp7* in committed and differentiating osteoblasts delineates the growth dynamics of individual skeletal

anlagen and *Shh* expression in rugae provides a temporally ordered set of A-P landmarks (regions in dashed boxes

enlarged below). (b) Summary model of skeletal growth dynamics during midfacial outgrowth. Growth of the

premaxilla (yellow) and palatine (pale blue) anlagen towards their characteristic shape occurs largely at the site of

initial specification. Following initial specification external to the oral cavity, the maxilla (purple) grows into the

anterior of the anterior secondary palate (black dashed arrows) towards the position of ruga 1 (red arrowhead) and

palatine as expansion of the anterior secondary palate (double headed red arrow) separates the primary and posterior

secondary palate. (c) Summary of the position of epithelial landmarks (see also Fig. 2) selected to capture segmental

growth dynamics of the primary palate (yellow) and anterior secondary palate (purple) and posterior secondary

palate (pale blue) during midfacial outgrowth. Abbreviations: maxilla (mx), palatine (p), premaxilla (pmx),

pterygoid (ptg), palatal process of the maxilla (ppmx), palatal process of the palatine (ppp)

In contrast, ruga 1 and the palatine anlage initially formed at the E11.5 anterior extent of

the secondary palate and maintained their proximity to each other throughout palatal

development even as they were displaced posteriorly between E11.5 and E15.5 (Fig. 7a). The

palatine and pterygoid bones later formed within the portion of the secondary palate that was

already present at the onset of secondary palate morphogenesis. These results indicated that the

posterior portion of the secondary palate and its associated anlage were present at the E11.5

onset of secondary palate morphogenesis, while the anterior secondary palate and associated

anlage did not contribute substantially to the secondary palate until later (Fig. 7b). The

combination of these morphogenetic results indicated that the meeting of the maxilla and

palatine anlagen was achieved predominantly by posterior growth of the maxilla towards the

palatine, which was itself being displaced posteriorly by anterior secondary palate elongation. By

E15.5, the maxilla and palatine anlagen met to form the maxillary-palatine suture subjacent to

ruga 1 (see also Supporting Fig. 1).

Our results collectively revealed that the segmental relationship between three palatal

regions and associated bone anlagen was already present at the onset of palatal morphogenesis.

Additionally, these results highlighted the previously unappreciated correlation of maxillary

growth to rugae formation as well as the greater proportional expansion of the anterior secondary

palate between E11.5 and E15.5 (summarized in Fig 7C). Based upon the spatial growth and

differential A-P dynamics revealed by our marker analysis, we defined a set of surface

landmarks intended to quantify the baseline characteristics and segmental differences in A-P

tissue domains underlying midfacial growth, within the context of overall midfacial elongation.

Interstrain Comparisons of Facial and Palatal Growth

Based on our findings that each upper jaw bone developed within one of the three palatal

segments and these segments contributed somewhat independently to midfacial A-P elongation,

we hypothesized that early differences in the proportional growth of the three palatal segments

produces strain specific differences in the contributions of specific bones to upper jaw length. If

true, previously identified differences in adult upper jaw morphology should be present by the

end of the critical embryonic A-P elongation period (i.e., by E15.5). We completed a

morphometric comparison of C57, NOD, and PWK strain embryonic midfacial shapes and a

comparison of proportional palatal segment lengths across embryonic and postnatal stages (Fig.

8) based on the direct relationship between segmental borders identified in the epithelial surfaces

and the specific bones that develop within the palatal segments.

Figure 8 – Palate segment length measurements – Identification of landmarks used to estimate midline lengths of the

primary palate (yellow), anterior secondary palate (blue), and posterior secondary palate (green) on (a) E11.0, (b)

E14.0, (c) E15.0, (d) P1 epithelial palatal surfaces, (e) P1 skeletal surfaces, and (f) adult skeletal surfaces. See also

Supporting Tables 1 and 2.

Since there is known variation in the speed of development between mouse strains and

within litters, we standardized our cross-strain ontogenetic analysis by embryonic developmental

age rather than gestational age. Limb bud outline-based developmental ages correlated strongly

with the number of days after plug identification for each measured strain. Though the

correlation coefficients for C57 mice (0.98), NOD (0.93) and PWK (0.90) were high, there was a

larger difference between limb age estimates and copulatory plug-based estimates of embryonic

age for our PWK samples. The C57 specimen limb-based developmental ages were an average

of 0.26 days younger than our copulatory plug estimates, while NOD and PWK limb ages were

an average of 0.30 days and 1.09 days younger, respectively. The major divergence for PWK

was likely driven by a slower speed of embryonic development in this wild-derived strain

compared to the two common lab strains (Murray et al., 2010). Even so, standardization by

developmental age should allow for improved direct comparisons between all strains.

We compared the midfacial shape of NOD and PWK specimens with C57 specimens

along the principal axes of facial shape variation. This comparison indicated that mice with

similar limb-based developmental age estimates tended to fall near each other along the first and

second principal components, regardless of strain. As these axes represented 82% of facial shape

variance across our sample, all mouse strains displayed major similarities in ontogenetic shape

change across this period (Supporting Fig. 2A). However, across ontogenetic time (as

represented by PC1 score) NOD and PWK specimens had significantly lower PC3 values

compared to the C57 baseline (Fig. 9b). PWK specimens also had significantly lower PC4 values

(Fig. 9c).

Figure 9 – Specimen PC score residuals – The principal component (PC) score residuals for each specimen, relative

to the scores predicted from C57 strain specific segmented linear regressions of (a) PC2, (b) PC3, and (c) PC4 on the

specimen's PC1 score. In this way, the C57 facial shape growth trajectory serves as a baseline with which to

compare the shape of NOD and PWK specimens, based on the fact that the PC1 axis is strongly associated with

facial shape ontogeny from developmental age E11 to E15.

We visualized how these identified significant differences in PC3 or PC4 scores were

reflected within facial and palatal morphology by comparing shapes represented at the minimum

and maximum end of these major shape axes (Fig. 10). A minimum PC3 score was associated

with the two nostril landmarks (LM7 & 8) being relatively distant in the superior/inferior and

lateral directions. Lower PC3 score was also associated with relative proximity of the posterior

whisker margin (LM6) to the anterior canthus (LM4) (Fig. 10h). Within the palate, lower PC3

scores were associated with a relatively medial anterior secondary palatal shelf, suggesting more

medial palatal outgrowth (proportional to overall palatal length) at a given developmental stage

(Fig. 10b). The superior-inferior position of the palatal landmarks indicated that the secondary

palate was more highly angled (less flat) for specimens with low PC3 scores (Fig. 10e). Because

of their generally lower PC3 scores, we anticipated more vertical nostrils in NOD and PWK mice

with more highly angled palatal shelves in greater medial proximity to each other, when

compared to C57 mice within a given developmental age.

Figure 10 – Facial shape variation associated with PC3 and PC4 – Representation of the pattern of shape variation

associated with the third major axis and fourth major axis of embryonic facial/palatal shape variation, from (a-c)

palatal, (d-f) parasagittal, and (g-i) facial views. Superficial tissue landmarks are identified on an example E14.5

specimen to assist with interpretation of the landmark vectors that represent differences in minimum (circle) and

maximum (end of line) PC scores. See Supporting Fig. 2 for a plot of embryonic specimens along these major axes

of embryonic shape variation.

A minimum PC4 score was associated with the whole nasal region (LM1, LM8) being

positioned more inferiorly and the posterior edge of the whisker region (LM3, LM6) more

superiorly when compared to the other facial and palatal features (Fig. 10i, Supporting Fig. 4).

Within the palate, a low PC4 score was associated with a relatively anterior location of ruga 1

between anterior and posterior secondary palate (LM20 & 24) and a relatively posterior position

of the posterior edge of the posterior secondary palate (LM21 & 25 within Fig. 10c). Low PC4

scores were also associated with more medially positioned anterior secondary palate landmarks

that are closer to the boundary with the primary palate. Because of their generally lower PC4

scores, this suggested that PWK embryos display a relatively low superior nasal region, a

medially expanded anterior secondary palatal shelf with a relatively short anterior secondary

palate and relatively long posterior secondary palate, when comparisons are made within a given

developmental age.

A comparison of embryonic palatal segment lengths (Fig. 8a-c) across C57, NOD and

PWK samples indicated that all three mouse strains display parallel ontogenetic changes 1) in

overall palatal length and 2) in the proportional contributions of palatal regions to overall palatal

length during the period of early palatal morphogenesis (Fig. 11). However, comparisons of

regression residuals of proportional segment length versus facial shape PC1 score (as a proxy for

developmental age) indicated small but significant differences in palatal segment proportions

between mouse strains. The NOD and PWK primary palate proportional length values were

significantly lower than for C57 (p-values: 0.012 (NOD vs C57), <0.001 (PWK vs C57)) (Fig.

12a), while they also had significantly higher posterior secondary palate values than C57 (p-

values <0.001) (Fig. 12c). This indicated that the primary palate contributed relatively less and

the posterior secondary palate contributed proportionally more to total embryonic palate length

in these two mouse strains, although visual assessment suggested these differences do not occur

at the earliest measured developmental ages (Fig. 11b). Significantly lower anterior secondary

palate values for NOD and PWK (p-values: 0.029 (NOD vs C57), 0.023 (PWK vs C57) indicated

that the anterior secondary palate represented a lower proportion of the total palate length within

these strains (Fig. 12b), although these differences seemed less pronounced than those noted for

the other two segments (Fig. 11b). Taken together, these general trends suggested that C57 mice

had proportionally longer primary palates and proportionally shorter posterior secondary palates

than NOD and PWK mice, with a trend towards longer anterior secondary palates by the time of

medial palatal shelf fusion (Fig. 12).

Figure 11 – Embryonic palate segment length across ontogeny – The midline projected strain-specific mean lengths

of the primary palate, anterior secondary palate, and posterior secondary palate, presented for each limb-based

developmental stage category (E11-E15). Raw length values (in millimeters) are presented on the left, while palatal

segment length relative to overall midline palate length are presented on the right.

Figure 12 - Specimen palate segment length residuals – The proportional contributions of each palate length

segment (shapes), as compared to the proportional contributions predicted from C57 strain specific linear

regressions (curved lines) of proportional (a) primary palate, (b) anterior secondary palate, and (c) posterior

secondary palate midline lengths versus the specimen's PC1 score. Vertical lines represent the residual of each

specimen from C57 regression. In this way, the C57 palatal segment growth trajectory serves as a baseline that NOD

and PWK specimens were compared to, based on the fact that the PC1 axis is strongly associated with facial shape

ontogeny from E11 to E15.

Interstrain Comparisons of Postnatal Palatal Segments

We compared C57 and NOD proportional palatal segment lengths at P1 based on

epithelial surface landmarks and skeletal landmarks of associated bony elements. Based on the

P1 epithelial surface landmarks (Fig. 8d), C57 had a proportionally shorter anterior secondary

palate and a proportionally longer posterior secondary palate (p-values <0.01) than NOD (Fig.

13; Table 2). This differed from the embryonic pattern where C57 specimens generally displayed

proportionally shorter posterior secondary palates than NOD specimens. The significant

differences in epithelially-measured proportional segment contributions between NOD and C57

mice at P1 appeared largely based on differences in the measured length (in mm) of the posterior

secondary palate (Fig. 13).

Table 2 – Average proportional palatal segment lengths for postnatal specimens, presented as

percentages of total palatal length.

	C57			NOD			PWK
	P1 epithelial	P1 bone	Adult bone	P1 epithelial	P1 bone	Adult bone	Adult bone
Primary Palate / Premaxilla	18.4%	23.3%	32.4%	19.7%	24.4%	31.7%	30.7%
Ant. Sec. Palate / Maxilla	45.7%	47.5%	39.9%	49.5%	47.4%	41.8%	42.8%
Post. Sec. Palate / Palatine & Pterygoid	35.9%	29.2%	27.7%	30.8%	28.3%	26.5%	26.5%

Figure 13 – Interstrain comparisons of palate segment lengths in older specimen – The midline projected strain-

specific mean lengths of the primary palate, anterior secondary palate, and posterior secondary palate, based on

epithelial landmark measures for limb based developmental stages E14 and E15, as well as for P1 specimens.

Analogous length measures of primary palate derived premaxilla, anterior secondary palate derived maxilla, and

posterior secondary palate derived palatine/pterygoid bones are presented for P1 and adult specimens. Raw length

values (in millimeters) are presented on the left, while palatal segment length relative to overall midline palate

length are presented on the right.

709   A comparison of proportional palatal bone lengths in P1 samples (Fig. 8e) indicated that

710 C57 has a proportionally shorter premaxilla (associated with primary palate) and proportionally

711 longer palatine/pterygoid bones (associated with posterior secondary palate) (p-values <0.01)

712 than NOD (Fig. 13; Table 2). A similar comparison of proportional palatal bone lengths (Fig. 8f)

713 in 8-12 week old adult specimens of all three strains revealed that the premaxillary (primary

714 palate) contribution to total palate length is significantly lower in PWK (p-value: < 0.001) and

715 NOD (p-value: 0.044) compared to C57, with the opposite true for the maxillary (anterior

716 secondary palate) portion of the palate (p-values: < 0.001) (Fig. 13). The palatine/pterygoid

717 (posterior secondary palate) portion was proportionally shorter in PWK and NOD when

718 compared to C57 (p-values: < 0.001) (Fig. 13). The PWK average proportional measures of

719 premaxilla and maxilla showed a more extreme divergence from C57 than in the NOD mouse

720 comparison.

721   Overall, there were some ontogenetic changes related to strain differences for

722 proportional palatal segment lengths from the embryonic period to P1 and then again between P1

723 and adulthood. NOD and C57 showed parallel postnatal changes in palatal bone contributions

724 across the postnatal period, where the posterior secondary palate contribution remained fairly

725 stable (~27-29%), but the primary palate contribution increased from ~24% to ~32% and the

726 anterior secondary palate contribution decreased from ~47% to ~40-42% (Fig. 13; Table 2).

727 During the postnatal period, the entire mouse and its palate grew substantially, but premaxilla

728 growth contributed more than the maxilla growth to overall postnatal palatal elongation.

729 Although the A-P growth of the anterior secondary palate was a critical driver for midfacial

730 growth between E11.5 and E15.5, outgrowth of the premaxilla may have played a larger role in

731 postnatal midfacial outgrowth.

<u>Discussion</u>

We provided here a multifaceted illustration of normal murine midfacial growth

dynamics within the secondary palate and upper jaw bones during the earliest stages of palatal

growth, followed by subsequent changes in upper jaw bone proportions in newborn and adult

mice. By tracking the position of the RGZ, each palatal ruga, and palatal bone precursor

populations across the earliest phases of palatal A-P elongation, our results confirmed that the

first-formed palatal ruga, which sits at an important border of regulatory gene expression, also

represents a consistent morphological boundary between the presumptive maxilla of the anterior

secondary palate and the presumptive palatine and pterygoid bones of the posterior secondary

palate. The process of rugae morphogenesis is coupled to maxillary osteogenesis during anterior

secondary palate expansion. Also, from the time of its formation, ruga 1 represents the future

position of the maxillary-palatine suture.

We tested the hypothesis that proportional differences in the earliest embryonic phases of

palatal segment A-P elongation underlie strain-specific differences in the adult contributions of

the premaxilla, maxilla, and palatine/pterygoid bones to upper jaw length. While these early

morphogenetic processes are critical for normal midfacial outgrowth, our ontogenetic

comparison of epithelial and skeletal palatal anatomy across three mouse strains indicated that

adult-like strain-specific differences in upper jaw morphology are not present before or at birth.

However, the significant strain-specific differences that were identified during embryonic palatal

morphogenesis are a useful foundation for understanding the impact of background genetic

effects on facial shape and elongation.

Normal Upper Jaw and Palatal Elongation

Between mouse E11.5 and E15.5, as overall head size increases, the fused facial prominences grow dramatically outward to form an elongated snout. It is well known that A-P elongation of the palate is a necessary component of the integrated processes of midfacial outgrowth. However, we previously lacked a detailed picture of the earliest spatiotemporal dynamics of intramembranous skeletal specification and differentiation in the mammalian midface. Our combined morphometric and whole mount gene expression analysis spanned the period of secondary palate morphogenesis, providing a detailed description of early intramembranous skeletal specification and differentiation within the context of the surrounding palatal segments and rugae formation. This work indicated that the relationship between three longitudinal palate segments and associated upper jaw bones was already established during the earliest phases of palatal morphogenesis. These results also illustrated how differential growth of the palatal segments contribute to overall palate length in mice and likely in other mammals. While all three palatal segments elongated between E11.5 and E15.5, the proportional elongation of the anterior secondary palate was notably more pronounced than that of the primary or posterior secondary palate and was accompanied by rugae formation at the RGZ and elongation of the maxillary bone primordium.

Our results verified that ruga 1 forms at an important gene expression boundary between the anterior and posterior secondary palate. They also clarified previous hypotheses regarding the anatomical relationship between the position of ruga 1, the hard verus soft palate, and the maxillary-palatine suture (Pantalacci et al., 2008). Our results established that throughout palate morphogenesis, ruga 1 represents a stable morphological boundary positioned at the future maxillary-palatine suture and not the more posterior soft palate. Posteriorly, ruga 1 maintained a

position at the anterior edge of the palatine anlage as both are posteriorly displaced between

E11.5 and E15.5. This implied that at E11.5 the secondary palate was initially comprised of only

the posterior secondary palate domains that will give rise to the palatine and pterygoid bones.

Anterior to ruga 1, growth of the maxillary anlage into the secondary palate only occurred as

anterior secondary palate expansion displaced the original site of maxillary specification

anteriorly away from ruga 1. This suggested that cellular cues in the proximity of ruga 1 may

initially inhibit maxillary bone formation. Expansion of the palatal portion of the maxillary

anlage appeared uniquely coupled to dynamics within the RGZ of anterior secondary palate

elongation, meaning this critical period of RGZ associated morphogenesis likely plays a major

role in determining the proportional contribution of the maxilla to the upper jaw within a broad

evolutionary context.

Differences in the formation and anatomy of two major upper jaw sutures may also be

related to segmental differences in early bone growth. The premaxilla and maxilla were specified

in adjacent tissue domains and maintained close apposition throughout palate morphogenesis,

suggesting equal contributions to the formation of the premaxillary-maxillary suture. Conversely,

the palatal portion of the maxilla grew posteriorly towards the palatine anlage at rugae 1,

suggesting that maxillary growth (and likely RGZ dynamics) played a larger role than palatine

growth in determining maxillary-palatine suture position. Different growth dynamics at these

sutures may lead to a more vertical premaxillary-maxillary suture versus a more oblique

maxillary-palatine suture with substantial A-P overlap of the maxilla and palatine bones. Suture

formation also provides a critical niche for facial skeletal progenitors (Zhao et al., 2015). Thus, it

is possible that early differences in segmental contributions to palate and upper jaw sutures may

also contribute to differences in postnatal dynamics of midfacial growth and remodeling (Enlow

and Bang, 1965; Kurihara et al., 1980; Sarnat, 1997; Martinez-Maza et al., 2013; Vora et al.,

2015; Maga, 2016). Further support for this hypothesis comes from recent work that showed the

widespread presence of Gli1+ mesenchymal stem cells and nascent bone formation (via calcein

labeling) within the center of the adult maxillary-palatine suture and also in the maxillary bone,

but a lack of stem cells and only sparse bone formation in the palatine bone (Luo et al., 2019).

Given the established role of calvarial sutures in directing bone growth of the skull, further

investigation of osteoprogenitor dynamics during premaxillary-maxillary and maxillary-palatine

suture formation is warranted.

Interstrain Palatal Growth Comparison

Our data illustrated 1) direct early associations between palatal segments and upper jaw

bones morphogenesis and 2) that anterior secondary palate growth makes a central contribution

towards total palatal elongation. This suggested a model where changes in embryonic A-P

growth of the anterior secondary palate at the RGZ may produce the range of prognathism

observed amongst mammals (Young et al., 2014). Given that A-P growth of the anterior

secondary palate was associated with rugae formation, support for this mechanism is further

strengthened by the fact that flat faced humans typically have 3-6 palatal rugae (Hauser et al.,

1989; Jayasankar et al., 2016), mice have 8-9 rugae (Peterkova et al., 1987), and prognathic pigs

have 20-25 rugae (Tonge and McCance, 1965). Based on this potential evo-devo relationship, we

hypothesized that intra-species differences in the contribution of individual bones to upper jaw

length would be produced by strain-specific differences in early embryonic A-P palatal segment

elongation. Based on previously reported adult phenotypes (Percival et al., 2016), we

hypothesized that C57 strain mice would have a relatively long primary palate and relatively

short anterior secondary palate by E15, while NOD and PWK mice would display the opposite

phenotype.

Although patterns of embryonic and postnatal palatal growth were broadly similar across

our three inbred mouse strains (Figs. 11 & 13), we identified significant strain-associated

differences in facial shape and palatal segment length. Embryonic facial shape differences (Figs.

4, 9, & 10; Table 2), as identified along major axes of variation, may represent more exaggerated

versions of normal ontogenetic shifts in the relationship of various facial structures or shifts in

the relative timing of normal developmental events occurring within different parts of the face.

Significant strain-specific differences in palatal segments proportions were identified across the

embryonic samples, with NOD and PWK mice having proportionally longer posterior secondary

palates and shorter primary palates than C57 mice. This result broadly matched our hypothesis

that embryonic C57 would have a relatively long primary palate, although without support for

initially predicted differences in anterior secondary palate proportions. Therefore, these results

demonstrated that there were significant embryonic differences in palatal proportions between

the strains, but they do not coincide with the differences observed in adults.

The results also indicated that differences in proportional segment lengths between NOD

and C57 did not remain consistent between the embryonic, neonatal (P1), and adult specimens.

At P1, NOD had a relatively short posterior secondary palate (i.e. palatine and pterygoid)

compared to C57, based on epithelial and skeletal measures. NOD also displayed a relatively

long anterior secondary palate based on epithelial measures and relatively long premaxilla (i.e.

primary palate) based on skeletal measures. This indicated a swap in proportional contributions

for primary palate and posterior secondary palate between E15 and P1. Strain-specific

premaxillary proportions flipped again between P1 and adult measurements.

Although changes within the RGZ growth dynamics of anterior secondary palate growth

may be critical for major species-specific differences in prognathism, our results did not support

the idea that this mechanism is responsible for the adult intraspecies differences in upper jaw

morphology between our mouse strains. This may indicate that later prenatal (after E15) and

postnatal growth processes play a major role in determining adult upper jaw bone proportions.

Our results suggested that variation in postnatal growth of the primary palate derived premaxilla

contributed more to postnatal elongation of the upper jaw than growth of the other palatal

segments did. A larger sample of later embryonic ages (i.e., between E15.5 and P0) would allow

us to determine when the initial NOD pattern of short primary and long posterior secondary

palate changes to a pattern of long primary and short posterior secondary palate by P1.

Concluding Statement

Our multifaceted illustration of normal midfacial growth dynamics confirmed a one-to-

one relationship between palatal segments and upper jaw bones during the earliest stages of

palatal growth, suggesting that the first formed ruga represents a consistent morphological

boundary between anterior and posterior secondary palate bone precursors, being found at the

future maxillary-palatine suture. Our results also indicated that interactions at the RGZ

simultaneously drives rugae formation and elongation of the maxillary bone primordium within

the anterior secondary palate, which more than doubles in length prior to palatal shelf fusion.

Although it is likely that RGZ-driven A-P growth of the anterior secondary palate contributes to

evolutionary changes in facial upper jaw morphology, this process was not solely responsible for

the subtle strain-specific upper jaw bone proportions measured across three inbred mouse strains.

However, significant strain-specific differences in early palatal segment elongation provide a

useful foundation for understanding the impact of background genetic effects on facial

morphogenesis. Our multifaceted analysis of ontogenetic trends in palatal and facial elongation

provides a novel contextual framework and developmental perspective within which to evaluate

the impact of both non-pathogenic and pathogenic genetic differences on midfacial growth and

differentiation.

Acknowledgements

We are indebted to Benedikt Hallgrímsson for the use of his laboratory for uCT and microscopy

image collection. This work was supported by start-up funds to Christopher J. Percival from

Stony Brook University; NIH/NIDCR R01 grant DE028753 and UCSF Chancellor Funds from

Recruitment Package to Licia Selleri who supported the personnel and all the research completed

in her lab for this study; and NIH/NIDCR F32 Fellowship DE026950 to Ian Welsh. The authors

of this manuscript have no conflicts of interest to declare.

Data Availability Statement

μCT images, limb bud microcopy photographs, landmark coordinate data, and associated

morphometric analysis R scripts will be made available publicly as a dataset within Dryad

(https://doi.org/10.5061/dryad.ghx3ffbvb; Welsh et al., 2023) after publication of this manuscript

within a peer-reviewed journal.

Supporting Information

Supporting Tables and Figures are found as a Supporting .pdf

Author Contributions

ICW contributed to study design, acquisition and interpretation of histological images, and

drafting of the manuscript. MEF contributed to acquisition, analysis, and interpretation of

morphometric data, and critical revision of the manuscript. DL contributed to the design of

morphometric methods, acquisition of morphometric data, measurement and analysis of limb

bud images, and critical revision of the manuscript. IM contributed to acquisition of μCT images,

acquisition, measurement and analysis of limb bud images, and critical revision of the

manuscript. KH contributed to mouse breeding, specimen collection, histological data collection,

and critical revision of the manuscript. CJP contributed to study design, acquisition of μCT

images, acquisition, analysis, and interpretation of morphometric data, and drafting of the

manuscript.

References

Adams, D.C., Collyer, M.L., Kaliontzopoulou, A., 2020. geomorph: Software for geometric
morphometric analyses. R package version 3.3.1.https://cran.r-
project.org/package=geomorph
Boughner, J.C., Wat, S., Diewert, V.M., Young, N.M., Browder, L.W., Hallgrímsson, B., 2008.
Short-faced mice and developmental interactions between the brain and the face. Journal
of Anatomy 213, 646–662.
Bush, J.O., Jiang, R., 2012. Palatogenesis: morphogenetic and molecular mechanisms of
secondary palate development. Development 139, 231–243.
Cignoni, P., Callieri, M., Corsini, M., Dellepiane, M., Ganovelli, F., Ranzuglia, G., 2008.
Meshlab: an open-source mesh processing tool., in: Eurographics Italian Chapter
Conference. pp. 129–136.
https://doi.org/10.2312/LocalChapterEvents/ItalChap/ItalianChapConf2008/129-136
Cobb, S.N., O'Higgins, P., 2004. Hominins do not share a common postnatal facial ontogenetic
shape trajectory. Journal of Experimental Zoology Part B: Molecular and Developmental
Evolution 302, 302–321.
Collyer, M.L., Adams, D.C., 2018. RRPP: An R package for fitting linear models to high-
dimensional data using residual randomization. Methods in Ecology and Evolution.
Dillon, S., Staines, K.A., Millán, J.L., Farquharson, C., 2019. How To Build a Bone:
PHOSPHO1, Biomineralization, and Beyond. JBMR Plus 3, e10202.
https://doi.org/10.1002/jbm4.10202

Economou, A.D., Ohazama, A., Porntaveetus, T., Sharpe, P.T., Kondo, S., Basson, M.A., et al.,
2012. Periodic stripe formation by a Turing mechanism operating at growth zones in the
mammalian palate. Nat Genet 44, 348–351. https://doi.org/10.1038/ng.1090
Enlow, D.H., Bang, S., 1965. Growth and remodeling of the human maxilla. Am J Orthod 51,
446–464. https://doi.org/10.1016/0002-9416(65)90242-3
Fedorov, A., Beichel, R., Kalpathy-Cramer, J., Finet, J., Fillion-Robin, J.-C., Pujol, S., et. al.,
2012. 3D Slicer as an image computing platform for the Quantitative Imaging Network.
Magnetic Resonance Imaging 30, 1323–1341. https://doi.org/10.1016/j.mri.2012.05.001
Fish, J.L., 2019. Evolvability of the vertebrate craniofacial skeleton. Semin Cell Dev Biol 91,
13–22. https://doi.org/10.1016/j.semcdb.2017.12.004
Galea, G.L., Zein, M.R., Allen, S., Francis-West, P., 2021. Making and shaping endochondral
and intramembranous bones. Dev Dyn 250, 414–449. https://doi.org/10.1002/dvdy.278
Geyer, S.H., Mohun, T.J., Weninger, W.J., 2009. Visualizing vertebrate embryos with episcopic
3D imaging techniques. ScientificWorldJournal 9, 1423–1437.
https://doi.org/10.1100/tsw.2009.154
Hall, J., Jheon, A.H., Ealba, E.L., Eames, B.F., Butcher, K.D., Mak, S.-S., et. al., 2014.
Evolution of a developmental mechanism: Species-specific regulation of the cell cycle
and the timing of events during craniofacial osteogenesis. Dev Biol 385, 380–395.
https://doi.org/10.1016/j.ydbio.2013.11.011
Hallgrímsson, B., Jamniczky, H., Young, N.M., Rolian, C., Parsons, T.E., Boughner, J.C., et. al.,
2009. Deciphering the Palimpsest: Studying the Relationship between Morphological
Integration and Phenotypic Covariation. Evolutionary Biology 36, 355–376.
Hammond, N.L., Dixon, M.J., 2022. Revisiting the embryogenesis of lip and palate
development. Oral Dis 28, 1306–1326. https://doi.org/10.1111/odi.14174
Hauser, G., Daponte, A., Roberts, M., 1989. Palatal rugae. Journal of Anatomy 165, 237–249.
Hilliard, S.A., Yu, L., Gu, S., Zhang, Z., Chen, Y.P., 2005. Regional regulation of palatal growth
and patterning along the anterior–posterior axis in mice. Journal of Anatomy 207, 655–
667. https://doi.org/10.1111/j.1469-7580.2005.00474.x
Hojo, H., Ohba, S., 2022. Sp7 Action in the Skeleton: Its Mode of Action, Functions, and
Relevance to Skeletal Diseases. Int J Mol Sci 23, 5647.
https://doi.org/10.3390/ijms23105647
Jayasankar, P., Banker, A., Bhattacharya, A., Gandhi, R., Patel, N., Parikh, S., 2016. Quantitative
and qualitative analysis of palatal rugae patterns in Gujarati population: A retrospective,
cross-sectional study. J Forensic Dent Sci 8, 126. https://doi.org/10.4103/0975-
1475.195110
Kaucka, M., Petersen, J., Tesarova, M., Szarowska, B., Kastriti, M.E., Xie, M., et. al., 2018.
Signals from the brain and olfactory epithelium control shaping of the mammalian nasal
capsule cartilage. Elife 7, e34465. https://doi.org/10.7554/eLife.34465
Kawasaki, M., Kawasaki, K., Meguro, F., Yamada, A., Ishikawa, R., Porntaveetus, T., et. al.,
2018. Lrp4/Wise regulates palatal rugae development through Turing-type reaction-
diffusion mechanisms. PLoS One 13, e0204126.
https://doi.org/10.1371/journal.pone.0204126
Kundrát, M., Coria, R.A., Manning, T.W., Snitting, D., Chiappe, L.M., Nudds, J., et. al., 2020.
Specialized Craniofacial Anatomy of a Titanosaurian Embryo from Argentina. Current
Biology 30, 4263-4269.e2. https://doi.org/10.1016/j.cub.2020.07.091

Kurihara, S., Enlow, D.H., Rangel, R.D., 1980. Remodeling reversals in anterior parts of the
human mandible and maxilla. Angle Orthod 50, 98–106. https://doi.org/10.1043/0003-
3219(1980)050<0098:RRIAPO>2.0.CO;2
Lee, S.-H., Bédard, O., Buchtová, M., Fu, K., Richman, J.M., 2004. A new origin for the
maxillary jaw. Developmental biology 276, 207–224.
Leigh, S.R., 2006. Cranial ontogeny of Papio baboons (Papio hamadryas). American Journal of
Physical Anthropology 130, 71–84. https://doi.org/10.1002/ajpa.20319
Li, Q., Ding, J., 2007. Gene expression analysis reveals that formation of the mouse anterior
secondary palate involves recruitment of cells from the posterior side. Int. J. Dev. Biol.
51, 167–172.
Luo, W., Yi, Y., Jing, D., Zhang, S., Men, Y., Ge, W.-P., et. al., 2019. Investigation of Postnatal
Craniofacial Bone Development with Tissue Clearing-Based Three-Dimensional
Imaging. Stem Cells and Development 28, 1310–1321.
https://doi.org/10.1089/scd.2019.0104
Maga, A.M., 2016. Postnatal Development of the Craniofacial Skeleton in Male C57BL/6J Mice.
J Am Assoc Lab Anim Sci 55, 131–136.
Martinez-Maza, C., Rosas, A., Nieto-Díaz, M., 2013. Postnatal changes in the growth dynamics
of the human face revealed from bone modelling patterns. J Anat 223, 228–241.
https://doi.org/10.1111/joa.12075
Morris, Z.S., Abzhanov, A., 2021. Heading for higher ground: Developmental origins and
evolutionary diversification of the amniote face. Curr Top Dev Biol 141, 241–277.
https://doi.org/10.1016/bs.ctdb.2020.12.003
Murray, S.A., Morgan, J.L., Kane, C., Sharma, Y., Heffner, C.S., Lake, J., et. al., 2010. Mouse
Gestation Length Is Genetically Determined. PLoS ONE 5, e12418.
https://doi.org/10.1371/journal.pone.0012418
Musy, M., Flaherty, K., Raspopovic, J., Robert-Moreno, A., Richtsmeier, J.T., Sharpe, J., 2018.
A quantitative method for staging mouse embryos based on limb morphometry.
Development 145, 1–7.
Okano, J., Suzuki, S., Shiota, K., 2006. Regional heterogeneity in the developing palate:
morphological and molecular evidence for normal and abnormal palatogenesis.
Congenital Anomalies 46, 49–54. https://doi.org/10.1111/j.1741-4520.2006.00103.x
Palmer, A.R., Strobeck, C., 1986. Fluctuating Asymmetry: Measurement, Analysis, Patterns.
Annu. Rev. Ecol. Syst. 17, 391–421.
https://doi.org/10.1146/annurev.es.17.110186.002135
Pantalacci, S., Prochazka, J., Martin, A., Rothova, M., Lambert, A., Bernard, L., et. al., 2008.
Patterning of palatal rugae through sequential addition reveals an anterior/posterior
boundary in palatal development. BMC Developmental Biology 8, 116.
https://doi.org/10.1186/1471-213X-8-116
Pauws, E., Hoshino, A., Bentley, L., Prajapati, S., Keller, C., Hammond, P., et. al., 2009.
Tbx22null mice have a submucous cleft palate due to reduced palatal bone formation and
also display ankyloglossia and choanal atresia phenotypes. Hum Mol Genet 18, 4171–
4179. https://doi.org/10.1093/hmg/ddp368
Percival, C.J., Liberton, D.K., Pardo-Manuel de Villena, F., Spritz, R., Marcucio, R.,
Hallgrímsson, B., 2016. Genetics of murine craniofacial morphology: diallel analysis of
the eight founders of the Collaborative Cross. Journal of Anatomy 228, 96–112.
https://doi.org/10.1111/joa.12382

Peterkova, R., Klepacek, I., Peterka, M., 1987. Prenatal development of rugae palatinae in mice:
scanning electron microscopic and histologic studies. Journal of craniofacial genetics and
developmental biology 7, 169–89.
Ponce de León, M.S.P., Zollikofer, C.P.E., 2001. Neanderthal cranial ontogeny and its
implications for late hominid diversity. Nature 412, 534–538.
R Core Team, 2021. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria [WWW Document]. URL https://www.R-
project.org/
Sarnat, B.G., 1997. Some methods of assessing postnatal craniofaciodental growth: a
retrospective of personal research. Cleft Palate Craniofac J 34, 159–172.
https://doi.org/10.1597/1545-1569_1997_034_0159_smoapc_2.3.co_2
Schneider, R.A., 2015. Regulation of Jaw Length During Development, Disease, and Evolution.
Curr Top Dev Biol 115, 271–298. https://doi.org/10.1016/bs.ctdb.2015.08.002
Selleri, L., Rijli, F.M., 2023. Shaping faces: genetic and epigenetic control of craniofacial
morphogenesis. Nature Reviews Genetics 24, 610–626. https://doi.org/10.1038/s41576-
023-00594-w
Singleton, M., 2012. Postnatal Cranial Development in Papionin Primates: An Alternative Model
for Hominin Evolutionary Development. Evolutionary Biology 39, 499–520.
https://doi.org/10.1007/s11692-011-9153-4
Smith, F.J., Percival, C.J., Young, N.M., Hu, D., Schneider, R.A., Marcucio, R.S., et. al., 2015.
Divergence of craniofacial developmental trajectories among avian embryos.
Developmental Dynamics. https://doi.org/10.1002/dvdy.24262
Tonge, C.H., McCance, R.A., 1965. Severe undernutrition in growing and adult animals: 15. The
mouth, jaws and teeth of pigs. British Journal of Nutrition 19, 361–372.
https://doi.org/10.1079/BJN19650034
Vora, S.R., Camci, E.D., Cox, T.C., 2015. Postnatal Ontogeny of the Cranial Base and
Craniofacial Skeleton in Male C57BL/6J Mice: A Reference Standard for Quantitative
Analysis. Front Physiol 6, 417. https://doi.org/10.3389/fphys.2015.00417
Welsh, I.C., Feiler, M.E., Lipman, D., Mormile, I., Hansen, K., Percival, C.J., 2023. Palatal
segment contributions to midfacial growth in three inbred mouse strains.
https://doi.org/10.5061/dryad.ghx3ffbvb
Welsh, I.C., Hagge-Greenberg, A., O'Brien, T.P., 2007. A dosage-dependent role for Spry2 in
growth and patterning during palate development. Mechanisms of Development 124,
746–761. https://doi.org/10.1016/j.mod.2007.06.007
Welsh, I.C., Hart, J., Brown, J.M., Hansen, K., Rocha Marques, M., Aho, R.J., et. al., 2018. Pbx
loss in cranial neural crest, unlike in epithelium, results in cleft palate only and a broader
midface. Journal of Anatomy 233, 222–242. https://doi.org/10.1111/joa.12821
Welsh, I.C., O'Brien, T.P., 2009. Signaling integration in the rugae growth zone directs
sequential SHH signaling center formation during the rostral outgrowth of the palate.
Developmental Biology 336, 53–67. https://doi.org/10.1016/j.ydbio.2009.09.028
Wilson, R., McGuire, C., Mohun, T., DMDD Project, 2016. Deciphering the mechanisms of
developmental disorders: phenotype analysis of embryos from mutant mouse lines.
Nucleic Acids Res 44, D855-861. https://doi.org/10.1093/nar/gkv1138
Woronowicz, K.C., Schneider, R.A., 2019. Molecular and cellular mechanisms underlying the
evolution of form and function in the amniote jaw. EvoDevo 10, 17.
https://doi.org/10.1186/s13227-019-0131-8

Young, N.M., Hu, D., Lainoff, A.J., Smith, F.J., Diaz, R., Tucker, A.S., et. al., 2014. Embryonic
bauplans and the developmental origins of facial diversity and constraint. Development
141, 1059–1063. https://doi.org/10.1242/dev.099994
Yu, K., Ornitz, D.M., 2011. Histomorphological study of palatal shelf elevation during murine
secondary palate formation. Developmental Dynamics 240, 1737–1744.
https://doi.org/10.1002/dvdy.22670
Yu, L., Gu, S., Alappat, S., Song, Y., Yan, M., Zhang, X., et. al., 2005. Shox2-deficient mice
exhibit a rare type of incomplete clefting of the secondary palate. Development 132,
4397–4406. https://doi.org/10.1242/dev.02013
Yuan, Y., Loh, Y.-H.E., Han, X., Feng, J., Ho, T.-V., He, J., et. al., 2020. Spatiotemporal cellular
movement and fate decisions during first pharyngeal arch morphogenesis. Sci Adv 6,
eabb0119. https://doi.org/10.1126/sciadv.abb0119
Zelditch, M., Swiderski, D., Sheets, D.H., 2012. Geometric morphometrics for biologists: a
primer, 2nd ed. Elsevier Academic Press, San Diego.
Zhao, H., Feng, J., Ho, T.-V., Grimes, W., Urata, M., Chai, Y., 2015. The suture provides a niche
for mesenchymal stem cells of craniofacial bones. Nat Cell Biol 17, 386–396.
https://doi.org/10.1038/ncb3139
Zumpano, M.P., Richtsmeier, J.T., 2003. Growth-related shape changes in the fetal craniofacial
complex of humans (Homo sapiens) and pigtailed macaques (Macaca nemestrina): a 3D-
CT comparative analysis. American Journal of Physical Anthropology 120, 339–351.