1   **Palatal segment contributions to midfacial anterior-posterior growth**

2   Short running page heading: **Palatal segment contributions to midfacial growth**

3   Ian C. Welsh @X – 1) Program in Craniofacial Biology, 2) Department of Orofacial
4   Sciences and 3) Department of Anatomy, University of California at San Francisco, San
5   Francisco, California 94143, USA
6
7   Maria E. Feiler – Interdepartmental Doctoral Program in Anthropological Sciences, Stony Brook
8   University, Stony Brook, NY 11790
9
10   Danika Lipman @Y – Department of Cell Biology and Anatomy, University of Calgary
11
12   Isabel Mormile – Interdepartmental Doctoral Program in Anthropological Sciences, Stony Brook
13   University, Stony Brook, NY 11790
14
15   Karissa Hansen – 1) Program in Craniofacial Biology, 2) Department of Orofacial Sciences and
16   3) Department of Anatomy, University of California San Francisco, San Francisco, CA
17   94143
18
19   Christopher J. Percival* – Department of Anthropology, Stony Brook University, Stony Brook
20   NY 11794
21
22   * = Corresponding Author: Department of Anthropology, Circle Road, SBS Building S-501,
23   Stony Brook, NY 11994, christopher.percival@stonybrook.edu
24   @X Present Address: The Jackson Laboratory, 600 Main St. Bar Harbor, ME 04609
25   @Y Present Address: Department of Mathematics and Statistics, 612 Campus Place NW,
26   University of Calgary, Calgary, AB
27

Abstract

Anterior-posterior (A-P) elongation of the palate is a critical aspect of integrated midfacial morphogenesis. Reciprocal epithelial-mesenchymal interactions drive secondary palate elongation that is coupled to the periodic formation of signaling centers within the rugae growth zone (RGZ). However, the relationship between RGZ-driven morphogenetic processes, the differentiative dynamics of underlying palatal bone mesenchymal precursors, and the segmental organization of the upper jaw has remained enigmatic. A detailed ontogenetic study of these relationships is important because palatal segment growth is a critical aspect of normal midfacial growth, can produce dysmorphology when altered, and is a likely basis for evolutionary differences in upper jaw morphology. We completed a combined whole mount gene expression and morphometric analysis of normal murine palatal segment growth dynamics and resulting upper jaw morphology. Our results demonstrated that the first formed palatal ruga (ruga 1), found just posterior to the RGZ, maintained an association with important nasal, neurovascular and palatal structures throughout early midfacial development. This suggested that these features are positioned at a proximal source of embryonic midfacial directional growth. Our detailed characterization of midfacial morphogenesis revealed a one-to-one relationship between palatal segments and upper jaw bones during the earliest stages of palatal elongation. Growth of the maxillary anlage within the anterior secondary palate is uniquely coupled to RGZ-driven morphogenesis. This may help drive the unequaled proportional elongation of the anterior secondary palate segment prior to palatal shelf fusion. Our results also demonstrated that the future maxillary-palatine suture, approximated by the position of ruga 1 and consistently associated with the palatine anlage, formed predominantly via the posterior differentiation of the maxilla within the expanding anterior secondary palate. Our ontogenetic analysis provides a

novel and detailed picture of the earliest spatiotemporal dynamics of intramembranous midfacial

skeletal specification and differentiation within the context of the surrounding palatal segment A-

P elongation and associated rugae formation.

Keywords: craniofacial, palatogenesis, secondary palate, rugae, face elongation, *Mus musculus,*

gene expression, RRID:IMSR_JAX:000664

Abbreviations: anterior-posterior (A-P), cranial neural crest (CNC), frontonasal process (FNP),

High resolution episcopic microscopy (HREM), maxillary processes (MxP), medial nasal

processes (mnp), micro-computed tomography (μCT), rugae growth zone (RGZ), transcription

factors (TFs), Whole mount in situ hybridization (WISH)

Introduction

Morphological variation of the midfacial complex, which consists of the nose, upper jaw,

cheek, and palate, is a defining aspect of both intra- and inter-specific differences in facial shape.

Facial shape morphogenesis begins with the specification of cranial neural crest (CNC) cells

during neurulation, which then delaminate around mouse embryonic day (E) 8.5 and migrate to

drive facial prominence outgrowth between E9 and E10 (Minoux and Rijli, 2010; Dash and

Trainor, 2020). By mouse E11.5, fusion between the medial nasal processes (mnp) of the

frontonasal process (FNP) and the anterior maxillary processes (MxP) gives rise to the primary

palate and lip, producing a unified upper jaw from previously separated tissues (Fig. 1).

Secondary palate morphogenesis then begins through outgrowth of the nascent palatal shelf

along the medial aspect of the MxP. Significant palatal growth along the anterior-posterior (A-P)

axis is accompanied by vertical growth of the palatal shelves prior to their elevation and medial

fusion dorsal to the tongue, after which the palate separates the oral and nasal cavities (Bush and

Jiang, 2012; Hammond and Dixon, 2022). Disproportional growth of the FNP or MxP prior to

fusion can result in cleft lip and palate (Young et al., 2014), while poor coordination between the

multiple growth axes of MxP-derived tissues can prevent secondary palate closure even when the

palatal shelves maintain competency to fuse (Kouskoura, et al., 2013).

The three longitudinal segments of the palate are called the primary palate, anterior

secondary palate, and posterior secondary palate (Fig. 1). The boundary between primary palate

and anterior secondary palate is the posterior edge of the fused mnp. Although A-P organization

of the secondary palate can be defined by the presence of boney versus muscular tissue (i.e., the

hard versus soft palate) or modes of palatal shelf elevation and closure (reviewed by Yu and

Ornitz, 2011), we define secondary palate segments by molecular differences in tissue patterning

and cell signaling competence (Hilliard et al., 2005; Hammond and Dixon, 2022).

The A-P expression domains of certain transcription factors (TFs) and intercellular

signaling molecules within CNC-derived mesenchyme are important for A-P organization and

growth during palatal morphogenesis. For example, within this mesenchyme, *Msx1* and *Shox2*

are exclusively expressed in the anterior secondary palate while *Barx1* and *Tbx22* are exclusive

to the posterior secondary palate (Hilliard et al., 2005; Okano et al., 2006; Welsh et al., 2018).

Mutations of *Shox2* or *Tbx22* have resulted in midline clefts of the anterior or posterior palate,

respectively (Yu et al., 2005; Pauws et al., 2009), although regional expression is maintained in

genetically engineered mice with severe segmental growth defects (Welsh et al., 2018). The A-P

expression boundary of these and other factors is found at the first formed palatal ruga (Hilliard

et al., 2005; Li and Ding, 2007; Welsh et al., 2007; Welsh and O'Brien, 2009; Welsh et al., 2018;

Hammond and Dixon, 2022), one of multiple rugae, which are parallel epithelial thickenings on

the anterior secondary palate.

This first ruga (numbered according to Welsh et al., 2007, Welsh and O'Brien, 2009;

although Pantalacci et al., 2008 use a different numbering system) forms at E11.5 on the anterior

extent of the nascent secondary palate (red arrow in Fig. 1). Remaining rugae form sequentially

within the rugae growth zone (RGZ) located just anterior to ruga 1 (Welsh and O'Brien, 2009),

via a Turing type mechanism (Economou et al., 2012; Kawasaki et al., 2018). Rugae are centers

of Sonic Hedgehog (SHH) signaling, which is critical for anterior secondary palate patterning

and A-P growth. As the secondary palate completes midline fusion at E15.5, the last formed ruga

appears adjacent to ruga 1 in the middle of the secondary palate while each precedingly formed

ruga is found at an increasingly anterior position along the anterior secondary palate (Welsh and

O'Brien, 2009).

Based upon the dynamics of rugae formation, palate growth, and patterning gene

expression, Pantalacci (et al., 2008) hypothesized that ruga 1 represents a distinct morphological

boundary related to either the future hard versus soft palate boundary or the maxillary-palatine

suture. We formally test these alternate hypotheses within our analysis of tissue relationships and

cell identity markers across a developmental time course spanning mouse secondary palate

morphogenesis and upper jaw outgrowth (E11.5-E15.5). More broadly, our analysis explores the

degree to which rugae formation and regional A-P patterns of secondary palate gene expression

are coordinated with palate skeletal differentiation and elongation across this critical

developmental period.

Our results establish one-to-one palatal segment origins of upper jaw bones, highlighting

a previously unappreciated coupling of maxillary bone formation to rugae morphogenesis during

initial elongation of the anterior secondary palate (E11-E15). Strikingly, the proportional

contribution of anterior secondary palate to overall palate length doubles during this ontogenetic

period. Further, the one-to-one relationship between palatal segments and bones also confirms

that ruga 1 is coincident with the future position of the maxillary-palatine suture, rather than the

future boundary of the hard and soft palate. Overall, our multifaceted analysis of ontogenetic

trends in palatal and midfacial elongation provides a novel contextual framework and

developmental perspective within which to evaluate the impact of both non-pathogenic and

pathogenic genetic differences on palatal segment contributions to midfacial growth and

differentiation.

Methods

Specimen and Image Acquisition

Animal breeding, specimen collection, and tissue fixation were performed in accordance

with the protocols of the University of California, San Francisco Institutional Animal Care and

Use Committee under protocol approval number AN192776-01F. Mice were socially housed

under a twelve-hour light-dark cycle with food and water *ad libitum*. Additional enrichment was

provided when single housing was required for breeding purposes. Mice were euthanized by CO_2

inhalation followed by cervical dislocation or decapitation. The C57BL/6J strain

(RRID:IMSR_JAX:000664; Jackson Labs, Bar Harbor, ME) was chosen for analysis because it

is the most widely utilized inbred strain for craniofacial genetic studies and because it was the

strain originally used to train the eMOSS limb bud staging algorithm (Musy et al., 2018) that we

employed as a basis for standardized developmental staging.

143   C57BL/6J embryos were collected between gestational days E11.5 and E15.5, as

144 determined from copulatory plug occurrence. Ten postnatal day one (P1) specimens were

145 collected. Embryo and P1 specimens were fixed in 4% PFA and stored in 1x PBS for micro-

146 computed tomography (μCT) imaging or dehydrated through a graded PBST-MeOH series and

147 stored at -20°C until use for in situ hybridization. A flowchart provides an overview of our study

148 design (Fig. 2).

149   Specimens for μCT scanning were received, stored, and imaged at the University of

150 Calgary in accordance with the protocols of the University of Calgary Institutional Care and Use

151 Committee under approval number AC13-0268. After approximately an hour of soaking in

152 Cysto-Conray II (Liebel-Flarsheim Canada), embryo heads were placed upside down on cheese

153 wax and immediately μCT scanned in air, a method that leads to minimal dehydration and good

154 tissue surface quality of surface anatomical structures (Schmidt, et al. 2010). These μCT images

155 were acquired with a Scanco μ35 with 45kV/177μA for images of 0.012 mm^3 voxel size. μCT

156 images of P1 heads were acquired similarly, but with 0.021 mm^3 voxel size. Photographs of

157 embryo hindlimb buds were collected using a dissecting microscope for developmental age

158 estimation. Adult specimens were previously collected and μCT imaged as described by Percival

159 et al., 2016.

161 **In Situ Hybridization**

162   In situ hybridization was performed as described in Welsh et al., 2018. cDNA probes for

163 *Phospho1* (nucleotides 274–1739 of RefSeq NM_153104), *Runx2* (nucleotides 1275–2704 of

164 RefSeq NM_001271627), *Shh* (full length cDNA clone from A. McMahon, University of

165 Southern California, Los Angeles CA, USA), *Shox2* (nucleotides 870–1603 of RefSeq

NM_130458), *Sp7* (nucleotides 431–1715 of RefSeq NM_130458), and *Tbx22* (nucleotides 307–

1547 of RefSeq NM_013665) were linearized and in vitro transcribed to label with either

digoxigenin (DIG) or dinitrophenol (DNP). Colorimetric detection of probes used BM purple

(dark blue), BCIP (teal), or MagentaPhos (magenta). Minimally, 3-6 embryos per time point

were processed for each probe analyzed.

DMDD HREM Data

High resolution episcopic microscopy (HREM) data (Geyer et al., 2009) was generated as

part of the deciphering mechanisms of developmental disorders (DMDD) program (Wilson et al.,

2016) and is available from the BioImage Archive at:

https://www.ebi.ac.uk/biostudies/bioimages/studies/S-BIAD490?query=DMDD

Embryonic Palatal Segment Growth

<u>Developmental Age Estimation</u>

Given that specimen developmental timing varies within litters and between litters of the

same estimated embryonic day, we categorized embryonic specimens by developmental age

rather than gestational age prior to morphometric analysis. Developmental age was estimated for

each µCT scanned embryonic specimen using eMOSS, an application that predicts

developmental age from hindlimb bud outlines, based on a previous analysis of C57BL/6J mice

(Musy et al., 2018). The resulting limb-based estimates of developmental age were reported as

days since conception, up to two decimal places. eMOSS developmental age estimates correlated

strongly with the number of days after plug identification (correlation coefficient: 0.98). Limb-

based developmental ages were an average of 0.26 days younger than our copulatory plug

estimates. We combined similar developmental age estimates within whole- or half-day

developmental age categories, which include specimens within 0.25 days of their initial eMOSS

estimate (Table 1).

<u>Anatomical Landmark Collection</u>

All embryo midfacial and palate landmarks were collected within Meshlab (Cignoni et

al., 2008) on minimum threshold-based epithelial tissue surfaces (downsampled x2) produced

from the µCT images. Landmarks that could be identified consistently in anatomically

homologous positions across the embryonic period of facial growth were chosen around the nose

and whisker region, the eyes, and along the palate (Fig. 3, Supporting Table 1). The midfacial

landmarks we collected were previously defined to track developmentally homologous structures

across multiple stages of facial development (Percival, et al., 2014) and our new set of epithelial

surface palatal landmarks were defined similarly.

To determine if differences in proportional palate segment length contributions were

consistent across embryonic and postnatal ages, a subset of epithelial palatal landmarks

(Supporting Table 2) were collected on epithelial surfaces produced from ten P1 specimen □CT

scans. Minimum threshold-based skeletal surfaces of the same ten P1 and twenty adult (70-73

days old; 9 male and 11 female) specimens were produced using 3D Slicer (Fedorov et al., 2012)

after Gaussian blur image filtering (sigma set to 0.01 for P1; sigma set to 0.02 for adult). Great

care was taken to identify skeletal anatomical landmarks that closely and homologously matched

the palate segment landmarks defined on surface epithelium (Supporting Table 2).

All epithelial landmarks were collected by a single observer to remove the possibility of

interobserver error. All skeletal landmarks were collected by a single observer for the same

reason. A reliability study was completed to quantify intraobserver error in landmark placement.

Two trials were completed by the epithelial landmark observer on three E11.5, three E12.5, and

three E14.5 specimens. The second trials were completed more than one year after the first trials

were completed, providing a realistic estimate of error across the entire period of landmark data

collection. There is no significant effect of trial or trial by age interaction on midfacial and

palatal shape, indicating there is no systematic difference in how landmarks were placed between

trial 1 and trial 2. The median millimeter difference in specific landmark location between trials

varies across landmarks (Supporting Fig. 1), but the overall mean landmark placement difference

is 0.034mm with a standard deviation of 0.022mm. This minor intraobserver error has a

negligible impact on our results, given that average total palate length is 1.23mm at E11 and

2.79mm at E14.

<u>Prenatal Geometric Morphometric Analysis</u>

A Procrustes superimposition-based geometric morphometric analysis was used to

quantify the ontogenetic shape change of the palate and face for embryonic specimens with limb-

based developmental age estimates between E11 and E15 (Table 1). This analysis of midfacial

and palatal landmark ontogeny was performed using *geomorph* (Adams et al., 2020) and RRPP

(Collyer and Adams, 2018) libraries in R Statistical Software (R Core Team, 2021). Generalized

Procrustes analysis (GPA) aligned specimen landmark sets by translating, scaling, and rotating

their landmark coordinates (reviewed by Zelditch et al., 2012). Subsequent shape analyses were

completed using the symmetric component of Procrustes-aligned specimen landmark coordinate

variation, assuming that most bilateral shape differences between the left and right sides of a

specimen's face are due to random effects associated with developmental noise and tissue

fixation (Palmer and Strobeck, 1986). Thus, symmetrized landmark coordinate data were

interpreted to represent the midfacial and palatal shape of the C57BL/6J inbred strain genotype.

The mean midfacial shape of specimens within each developmental age category were

estimated. Differences between age-specific mean shapes were plotted to illustrate typical

midfacial/palatal shape ontogenetic changes during this important period of secondary palate

elongation and midline fusion. A principal components analysis (PCA) was completed to

identify the major axes of shape covariation across the embryonic sample.

Palate Segment Length Comparisons

The AP lengths of the three major palatal segments (i.e., primary palate, anterior

secondary palate, and posterior secondary palate) were estimated in millimeters from landmark

coordinates without superimposition or scaling. To estimate the length of these segments along

the anterior-posterior axis of the palate, the mean midline position between bilateral landmark

pairs were calculated (Fig. 4). Proportional palatal segment lengths were calculated as the length

of a single segment divided by the sum of all three segment lengths for a given specimen.

Wilcoxon rank sum tests were used to identify significant differences in proportional palate

segment lengths between age categories and between epithelial and skeletal P1 landmarks.

Results

Secondary Palate A-P Growth and the Position of Ruga 1 at a Facial Growth Center

Close relationships between different anatomical domains underlying midfacial

outgrowth are found centered near ruga 1 (Fig. 5 & Supporting Fig. 2, red arrowheads). Our

mouse embryonic series confirmed that ruga 1 formed within the MxP from a domain of *Shh*

expression at the anterior extreme of the nascent secondary palate. As previously shown, other

rugae then formed anterior to ruga 1 at the RGZ (Fig. 5c, asterisk) as the anterior secondary

palate elongated, anteriorly displacing the primary palate and external nares. A separate domain

of *Shh* expression posterior to ruga 1 gave rise to the geschmacksstreifen (gs) and sensory papilla

arrayed across the most posterior palate. Our embryonic series also confirmed that ruga 1 is

consistently found at the *Shox2/Tbx22* gene expression boundary between the anterior and

posterior secondary palates during this critical embryonic period of palatal development (Fig. 5

& Supporting Fig. 2).

We identified a previously unappreciated association between ruga 1 and the posterior

wall of the nasal capsule. At E11.5, ruga 1 formed adjacent to the primary choanae (Fig. 5a &

Supporting Fig. 2a, white arrowheads), the posterior openings of the nasal passages, which

initially occur at the boundary between the primary and secondary palatal segments (Tamarin,

1982). Elongation of the anterior secondary palate then coincided with the formation and

expansion of the overlying nasal capsule. As the primary choanae and nasal capsule elongated

between E11.5 and E15.5, ruga 1 and the posterior wall of the nasal capsule remained in

approximately the same coronal plane (represented by position of red dashed lines in Fig. 5b, 5d,

5f and Supporting Fig. 2). Ruga 1 was also found coincident with a gross morphological

inflection point of the palatal tissue where the anterior secondary palate and overlying nasal

capsule exhibited an inferiorly accentuated angle relative to the cranial base, while the posterior

secondary palate did not (Fig. 5d-f, Supporting Fig. 2).

The greater palatine neurovascular bundle was also consistently associated with ruga 1

across this embryonic period. The greater palatine nerve and artery were found entering the

palatal shelves from a location immediately dorsal to ruga 1 at E12.5, and this spatial

relationship was maintained until the greater palatine foramen formed by the articulation of

maxilla and palatine bones around the palatine neurovascular bundle at the maxillary-palatine

suture (Fig. 5f, Supporting Fig. 2). In combination, the consistent associations between ruga 1,

the anterior-posterior secondary palate boundary, the posterior margin of the nasal capsule, and

greater palatine neurovascular bundle suggested that these features are positioned at a proximal

source of directional growth that the elongating anterior secondary palate and nasal capsule

extend away from between E11.5 and E15.5.

A complementary 3D morphometric analysis of midfacial and palatal epithelial

landmarks was used to quantify the integrated morphogenetic processes occurring during this

critical developmental period. The first major principal axis of midfacial and palatal shape

variation (PC1) for all limb bud staged E11-E15 specimens represented 73% of shape variance

and was associated with general ontogenetic growth during this time window (Fig. 6). Mice with

similar limb-based developmental age estimates fell near each other along the first principal

component of shape variation. Additional principal axes of embryonic midfacial shape variation

suggested shifts in the direction of ontogenetic changes over developmental time, with one

inflection noted along PC2 (Fig. 6a), two along PC3 (Fig. 6b), and three along PC4 (Fig. 6c).

Average developmental age category shape differences were illustrated as landmark specific

ontogenetic shifts in palate shape (Fig. 7) and overall midfacial shape (Supporting Fig. 3).

Between developmental stages E11 and E15, each of the three major palatal segments

grew in length, with significant differences identified for almost all segment-specific consecutive

age comparisons (Figs. 8a, 9a). However, the anterior secondary palate grew proportionally more

than the primary palate and posterior secondary palate (Table 2, Figs. 8c, 9c). Soon after the start

of embryonic A-P palate elongation, the primary palate represented ~20% of the total palate

length, the anterior secondary palate ~20%, and the posterior secondary palate ~60%. After

E12.5, anterior secondary palate length increased proportionally faster than total palate length

between each measured developmental day. By the time of midline palatal fusion, the primary

palate represented ~25% of palate length, the anterior secondary palate represented >40%, and

the posterior secondary palate represented <35%. Therefore, between E11 and E15, the

proportional contribution of the primary palate increased by a quarter while posterior secondary

palate contribution decreased by nearly a third (Table 2, Fig. 9c). Strikingly, the proportional

contribution of the anterior secondary palate to overall palate length doubled during this period.

Segmental Differences in Skeletal Specification, Differentiation, and Growth During

Secondary Palate Morphogenesis

We performed a WISH analysis of *Runx2*, *Sp7*, and *Phospho1* expression (Fig. 10)

alongside double labeled WISH of *Shh* and *Sp7* between E12.5 and E15.5 (Fig. 11a) to

characterize bone specification and differentiation in relation to A-P growth dynamics of

secondary palate segments. The position of rugae, which provided an ordered set of temporally

specific anatomical features, allowed for the identification of previously unappreciated

associations between the palatal segments and developing palatal bones.

Broad *Runx2* expression in early facial prominences included osteogenic domains (e.g.,

premaxilla within the primary palate) and later became more restricted to regions adjacent to the

molar tooth bud and medially along the adjoining palatal shelves (Fig. 10a). Osteoblast

expression of both *Sp7* and *Phospho1* more precisely delineated the morphology of the emerging

midfacial skeleton (Fig. 10b-c). Palatal processes of the maxilla and palatine bones formed

within the medial region of the elevated palatal shelves, a domain that maintained high levels of

Runx2 expression during secondary palate growth (Fig. 10a), suggesting maintenance of a less

differentiated osteoprogenitor population until palatal elevation and fusion was complete.

At E11.5, *Sp7* was strongly expressed in bilateral domains within the mnp of the forming

primary palate. Between E12.5 and E15.5, presaging premaxilla morphology, *Sp7* expression

formed a cup-like region surrounding the developing incisors (Fig. 10b). *Phospho1* was first

observed in the forming premaxilla at E12.5 (Fig. 10c), reflecting a temporal lag between initial

osteoblast commitment and later differentiation. The premaxilla acquired its characteristic adult

morphology within the primary palate segment in which it was initially specified.

Within the secondary palate, this simple developmental sequence was repeated for the

palatine and pterygoid bones within the posterior secondary palate, but development of the

maxilla within the anterior secondary palate was unique. At E11.5, *Sp7* expression in the MxP

was observed in two domains. The antero-lateral domain, which abutted the point of fusion with

the mnp (i.e., the primary palate) and extended laterally away from the oral cavity and choanae,

corresponded to the maxillary anlage. The posterior domain that spanned the length of the

forming secondary palate, from the anterior limit abutting the choanae to the posterior-most

palatal edge, gave rise to the palatine and pterygoid anlagen.

The antero-lateral domain of *Sp7* expression adjacent to the primary palate was initially

external to the secondary palate at the site of the future zygomatic plate. However, this

expression domain later expanded medially (E12.5-E13.5) and then posteriorly (E13.5-E15.5)

within the growing anterior secondary palate (white curved dashed arrow in Fig. 10b). This

expansion occurred after initial A-P growth of the anterior secondary palate and formation of 3-4

rugae anterior to ruga 1, suggesting that it is coupled with RGZ growth dynamics (Fig. 11a). The

fact that *Phospho1* expression closely followed *Sp7* spatial dynamics further supported the idea

that these *Sp7* expression patterns represent maxillary bone anlagen formation and expansion

(white curved dashed arrow in Fig. 10c). The maxillary anlage continued growing posteriorly

towards ruga 1 concomitant with rugae formation (white and black curved dashed arrows in Fig.

11a-b), within the elongating secondary palate.

The posterior domain of *Sp7* expression that initially spanned the length of the secondary

palate at E11.5 was displaced posteriorly by expansion of the anterior secondary palate between

E12.5 and E15.5. During this period, it separated into two subdomains. The anterior subdomain

remained associated with the posterior wall of the nasal capsule and gave rise to the palatine

bone, while the posterior subdomain formed both the medial and lateral pterygoid processes (Fig.

10b-c). Ruga 1 and the palatine anlage initially formed at the anterior extent of the secondary

palate at E11.5 and maintained proximity to each other throughout palatal development even as

both structures were displaced posteriorly (Fig. 11a).

The palatine and pterygoid anlagen grew to acquire their characteristic adult morphology

within the posterior secondary palate segment where their precursor domains were initially

specified, mirroring premaxillary anlagen formation within the primary palate. This differs from

growth of the maxillary anlage that was initially specified external to the secondary palate at the

site of the future zygomatic plate and only later extended into the anterior secondary palate,

following a period of initial anterior secondary palate elongation, to give rise to the majority of

the maxillary bone.

These results indicated that the posterior portion of the secondary palate and its

associated anlage were present within the secondary palate at the onset of its morphogenesis at

E11.5, while the anlage associated with the anterior secondary palate (i.e., the maxilla) did not

contribute substantially to the secondary palate until later (Fig. 11b). They also indicated that the

eventual meeting of the maxilla and palatine anlagen was an asymmetric process achieved

predominantly by posterior growth of the maxilla towards the palatine (dashed black curved

arrows in Fig. 11b). By E15.5, the maxilla and palatine anlagen met to form the maxillary-

palatine suture subjacent to ruga 1 (see also Supporting Fig. 2).

Our results collectively revealed that the segmental relationship between three palatal

regions and associated bone anlagen remained consistent throughout the period of palatal

elongation, although the maxillary anlage was not present within the anterior secondary palate

until partway through this period. Importantly, these results also highlighted the previously

unappreciated correlation of maxillary growth to rugae formation as well as to the greater

proportional expansion of the anterior secondary palate between E11.5 and E15.5 (summarized

in Fig. 11c).

Ontogeny of Postnatal Palatal Segment Length

We compared palatal segment lengths 1) between E15.5 and P1 based on epithelial

surface landmarks and 2) between P1 and adult specimens based on skeletal landmarks of bony

elements in 1:1 association with each palatal segment. Unsurprisingly, the length of each palate

segment increased significantly between E15 and P1 based on epithelial measures, and between

P1 and adult specimens based on skeletal measures (Figs. 8b, 9b). Proportional palate measures

of all palate segments also changed significantly for the same age group comparisons (Table 2,

Figs. 8c, 9c). Continuing the trend previously identified between E11.5 and E15.5, the anterior

secondary palate grew proportionally more than the other segments between E15 and P1.

However, based on palatal bone measurements, the primary palate derived premaxilla elongated

proportionally more than the other segments between P1 and adult specimens, (Table 2, Fig. 9c).

The posterior secondary palate contribution remained stable (~27-29%) across the postnatal

period, but the primary palate contribution increased from ~24% to ~32% and the anterior

secondary palate contribution decreased from ~47% to ~40% (Table 2). Although the A-P

growth of the anterior secondary palate was the largest proportional contributor to midfacial

growth between E11.5 and P1, outgrowth of the premaxilla plays a larger role during postnatal

midfacial outgrowth.

Discussion

Our results provided a multifaceted characterization of 1) normal murine midfacial

growth dynamics within the secondary palate and upper jaw bones during the earliest stages of

palatal growth and 2) subsequent changes in upper jaw bone proportions in newborn and adult

mice. By tracking the position of the RGZ, sequential palatal rugae formation, and palatal bone

precursor populations across the earliest phases of palatal A-P elongation, our results illustrated

that the first-formed palatal ruga (i.e., ruga 1), which sits at an important border of regulatory

gene expression, is coincident with important nasal, neurovascular and palatal structures

throughout early midfacial development. For example, ruga 1 represents a consistent

morphological boundary between the presumptive maxilla of the anterior secondary palate and

the presumptive palatine and pterygoid bones of the posterior secondary palate. This association

suggested that the process of rugae morphogenesis is coupled to maxillary osteogenesis during

anterior secondary palate elongation, which contributes disproportionately to embryonic palate

growth. It also verifies that ruga 1 approximates the future position of the maxillary-palatine

suture from the time of its formation, rather than being associated with the boundary between the

hard and soft palates.

Normal Upper Jaw and Palatal Elongation

421   It is well known that A-P elongation of the palate is a necessary component of the

integrated processes of midfacial outgrowth. However, within mammals, we previously lacked a

detailed picture of the earliest spatiotemporal dynamics of intramembranous midfacial skeletal

specification and differentiation within the context of the surrounding palatal segments and rugae

formation. Our analysis indicated that the relationship between three longitudinal palate

segments and associated upper jaw bones was already established during the earliest phases of

palatal morphogenesis. These results illustrated the substantial contribution of embryonic

anterior secondary palate elongation to overall palate length in mice and likely in other

mammals. Anterior secondary palate elongation was also directly associated with rugae

formation at the RGZ and elongation of the maxillary bone primordium.

        Beyond verifying that ruga 1 forms at an important gene expression boundary between

the anterior and posterior secondary palate, our results clarified that ruga 1 represented a stable

morphological boundary positioned near the future maxillary-palatine suture. Ruga 1 formed at

the earliest stages of anterior secondary palate elongation, then maintained a position at the

anterior edge of the palatine bone anlage as it was posteriorly displaced by expansion of the

anterior secondary palate. Anterior to ruga 1, the maxillary bone osteogenic population expanded

from the site of initial specification (i.e., the zygomatic plate) into the secondary palate after ruga

1 was posteriorly displaced during anterior secondary palate elongation. Given that the

secondary palate initially only contained osteogenic domains of the posterior secondary palate

(palatine and pterygoid precursors), these results suggested that cellular cues in the proximity of

ruga 1 may initially inhibit maxillary bone formation within the secondary palate. Additionally,

expansion of the palatal portion of the maxillary anlage was uniquely associated with RGZ

dynamics and anterior secondary palate elongation, suggesting that RGZ morphogenesis plays a

major role in determining the proportional contribution of the maxilla to the upper jaw.

Differences in upper jaw suture anatomy may also be related to segmental differences in

early bone growth. The premaxilla and maxilla were specified in adjacent tissue domains that

maintained close apposition throughout palate morphogenesis, suggesting equal contributions to

the formation of the premaxillary-maxillary suture. Conversely, the palatal portion of the maxilla

grew posteriorly towards the palatine anlage at ruga 1, suggesting that maxillary growth (and

likely RGZ dynamics) played a larger role than palatine growth in determining maxillary-

palatine suture position. Different growth dynamics at these sutures may lead to a vertical

premaxillary-maxillary suture versus an oblique maxillary-palatine suture with substantial A-P

overlap of the maxilla and palatine bones.

Because suture formation provides a critical niche for skeletal progenitors (Zhao et al.,

2015), the way palatal segments contribute to palatal sutures may contribute to variation in

postnatal dynamics of midfacial growth and remodeling (Enlow and Bang, 1965; Kurihara et al.,

1980; Sarnat, 1997; Martinez-Maza et al., 2013; Vora et al., 2015; Maga, 2016). The widespread

presence of Gli1+ mesenchymal stem cells and nascent bone within the adult maxillary-palatine

suture and the maxillary bone, but not in the palatine bone (Luo et al., 2019) provides further

support for this hypothesis. Given the established role of calvarial sutures in directing postnatal

craniofacial bone growth, further comparisons of osteoprogenitor dynamics during premaxillary-

maxillary and maxillary-palatine suture formation are warranted.

Palatal Growth as a Basis for Morphological Variation

Given its central contribution to total palatal elongation, variation in RGZ-regulated A-P

growth of the anterior secondary palate may play a role in determining the proportional

contribution of the maxilla to the upper jaw and contribute substantially to the range of

prognathism observed amongst mammals (Young et al., 2014). Support for this idea comes from

the fact that more prognathic species tend to have more rugae: flat-faced humans typically have

3-6 palatal rugae (Hauser et al., 1989; Jayasankar et al., 2016), mice have 8-9 rugae (Peterkova et

al., 1987), and prognathic pigs have 20-25 rugae (Tonge and McCance, 1965). Variation in

interspecies rugae number and palatal segment proportions may stem from ontogenetic shifts in

the relationship of various midfacial structures or shifts in the relative timing of normal

developmental events occurring within different parts of the face.

This hypothesis could be tested within an evo-devo context by characterizing variation in

palatal *Shh* expression between mammals and avians that have drastic differences in maxillary

morphology. These differences include MxP-driven palatal growth in mammals and FNP-driven

palatal growth in birds (Young et al., 2014) as well as an obligate avian cleft palate.

Alternatively, the ontogenetic basis for variation in palatal segment contributions could be

dissected genetically via analysis of *Shh* expression and rugae morphogenesis in mice with

altered A-P skeletal patterning, such as the *Pbx* CNCC mouse mutants (Welsh et al., 2018).

Although changes within the RGZ growth dynamics of anterior secondary palate growth may be

important for major species-specific differences in prognathism, postnatal growth processes also

likely play an important role in determining adult upper jaw bone proportions.

Concluding Statement

Our multifaceted illustration of normal midfacial growth dynamics confirmed a one-to-

one relationship between palatal segments and upper jaw bones during the earliest stages of

palatal growth, suggesting that the first formed ruga represents a consistent morphological

boundary between anterior and posterior secondary palate bone precursors, thus approximating

the position of the future maxillary-palatine suture. In addition to driving rugae formation, our

results suggested that interactions at the RGZ coordinate elongation of the maxillary bone

primordium within the anterior secondary palate, which more than doubles in length prior to

palatal shelf fusion. Overall, our integrated analysis of spatial gene expression dynamics and

palatal ontogenesis during midfacial elongation provides a critical new perspective within which

to consider palatal segment contributions to both normal developmental processes and the

etiological basis of genetic perturbations.

Acknowledgements

We are indebted to Benedikt Hallgrímsson for the use of his laboratory for uCT and microscopy

image collection. This work was supported by start-up funds to Christopher J. Percival from

Stony Brook University; NIH/NIDCR R01 grant DE028753 and UCSF Chancellor Funds from

Recruitment Package to Licia Selleri who supported the personnel and all the research completed

in her lab for this study; and NIH/NIDCR F32 Fellowship DE026950 to Ian Welsh. The authors

of this manuscript have no conflicts of interest to declare.

Data Availability Statement

μCT images, limb bud microscopy photographs, landmark coordinate data, and associated

morphometric analysis R scripts will be available publicly as a dataset within Dryad

(https://doi.org/10.5061/dryad.ghx3ffbvb; Welsh et al., 2024) when this manuscript is accepted

for publication.

Author Contributions

ICW contributed to study design, acquisition and interpretation of histological images, and

drafting of the manuscript. MEF contributed to acquisition, analysis, and interpretation of

morphometric data, and critical revision of the manuscript. DL contributed to the design of

morphometric methods, acquisition of morphometric data, measurement and analysis of limb

bud images, and critical revision of the manuscript. IM contributed to acquisition of μCT images,

acquisition, measurement and analysis of limb bud images, and critical revision of the

manuscript. KH contributed to mouse breeding, specimen collection, histological data collection,

and critical revision of the manuscript. CJP contributed to study design, acquisition of μCT

images, acquisition, analysis, and interpretation of morphometric data, and drafting of the

manuscript.

References

Adams, D.C., Collyer, M.L., Kaliontzopoulou, A., 2020. geomorph: Software for geometric

morphometric analyses. R package version 3.3.1.https://cran.r-

project.org/package=geomorph

Bush, J.O., Jiang, R., 2012. Palatogenesis: morphogenetic and molecular mechanisms of

secondary palate development. Development 139, 231–243.

Cignoni, P., Callieri, M., Corsini, M., Dellepiane, M., Ganovelli, F., Ranzuglia, G., 2008.

Meshlab: an open-source mesh processing tool., in: Eurographics Italian Chapter

Conference. pp. 129–136.

https://doi.org/10.2312/LocalChapterEvents/ItalChap/ItalianChapConf2008/129-136

Collyer, M.L., Adams, D.C., 2018. RRPP: An R package for fitting linear models to high-

dimensional data using residual randomization. Methods in Ecology and Evolution.

Dash, S., Trainor, P.A., 2020. The Development, Patterning, and Evolution of Neural Crest Cell

Differentiation into Cartilage and Bone. Bone 137, 115409.

https://doi.org/10.1016/j.bone.2020.115409

Economou, A.D., Ohazama, A., Porntaveetus, T., Sharpe, P.T., Kondo, S., Basson, M.A., et al.,

2012. Periodic stripe formation by a Turing mechanism operating at growth zones in the

mammalian palate. Nat Genet 44, 348–351. https://doi.org/10.1038/ng.1090

Enlow, D.H., Bang, S., 1965. Growth and remodeling of the human maxilla. Am J Orthod 51,

446–464. https://doi.org/10.1016/0002-9416(65)90242-3

Fedorov, A., Beichel, R., Kalpathy-Cramer, J., Finet, J., Fillion-Robin, J.-C., Pujol, S., et. al.,

2012. 3D Slicer as an image computing platform for the Quantitative Imaging Network.

Magnetic Resonance Imaging 30, 1323–1341. https://doi.org/10.1016/j.mri.2012.05.001

Geyer, S.H., Mohun, T.J., Weninger, W.J., 2009. Visualizing vertebrate embryos with episcopic

3D imaging techniques. ScientificWorldJournal 9, 1423–1437.

https://doi.org/10.1100/tsw.2009.154

Hammond, N.L., Dixon, M.J., 2022. Revisiting the embryogenesis of lip and palate

development. Oral Dis 28, 1306–1326. https://doi.org/10.1111/odi.14174

Hauser, G., Daponte, A., Roberts, M., 1989. Palatal rugae. Journal of Anatomy 165, 237–249.

Hilliard, S.A., Yu, L., Gu, S., Zhang, Z., Chen, Y.P., 2005. Regional regulation of palatal growth

556   and patterning along the anterior–posterior axis in mice. Journal of Anatomy 207, 655–

557   667. https://doi.org/10.1111/j.1469-7580.2005.00474.x

558   Jayasankar, P., Banker, A., Bhattacharya, A., Gandhi, R., Patel, N., Parikh, S., 2016. Quantitative

559   and qualitative analysis of palatal rugae patterns in Gujarati population: A retrospective,

560   cross-sectional study. J Forensic Dent Sci 8, 126. https://doi.org/10.4103/0975-

561   1475.195110

562   Kawasaki, M., Kawasaki, K., Meguro, F., Yamada, A., Ishikawa, R., Porntaveetus, T., et. al.,

563   2018. Lrp4/Wise regulates palatal rugae development through Turing-type reaction-

564   diffusion mechanisms. PLoS One 13, e0204126.

565   https://doi.org/10.1371/journal.pone.0204126

566   Kouskoura, T., Kozlova, A., Alexiou, M., Blumer, S., Zouvelou, V., Katsaros, C., Chiquet, M.,

567   Mitsiadis, T.A., Graf, D., 2013. The Etiology of Cleft Palate Formation in BMP7-

568   Deficient Mice. PLos One 8, e59463. https://doi.org/10.1371/journal.pone.0059463

569   Kurihara, S., Enlow, D.H., Rangel, R.D., 1980. Remodeling reversals in anterior parts of the

570   human mandible and maxilla. Angle Orthod 50, 98–106. https://doi.org/10.1043/0003-

571   3219(1980)050<0098:RRIAPO>2.0.CO;2

572   Li, Q., Ding, J., 2007. Gene expression analysis reveals that formation of the mouse anterior

573   secondary palate involves recruitment of cells from the posterior side. Int. J. Dev. Biol.

574   51, 167–172.

575   Luo, W., Yi, Y., Jing, D., Zhang, S., Men, Y., Ge, W.-P., et. al., 2019. Investigation of Postnatal

576   Craniofacial Bone Development with Tissue Clearing-Based Three-Dimensional

Imaging. Stem Cells and Development 28, 1310–1321.

https://doi.org/10.1089/scd.2019.0104

Maga, A.M., 2016. Postnatal Development of the Craniofacial Skeleton in Male C57BL/6J Mice.

J Am Assoc Lab Anim Sci 55, 131–136.

Martinez-Maza, C., Rosas, A., Nieto-Díaz, M., 2013. Postnatal changes in the growth dynamics

of the human face revealed from bone modelling patterns. J Anat 223, 228–241.

https://doi.org/10.1111/joa.12075

Minoux, M., Rijli, FM., 2010. Molecular Mechanisms of Cranial Neural Crest Cell Migration

and Patterning in Craniofacial Development. Development 137, 2605–2621.

https://doi.org/10.1242/dev.040048

Musy, M., Flaherty, K., Raspopovic, J., Robert-Moreno, A., Richtsmeier, J.T., Sharpe, J., 2018.

A quantitative method for staging mouse embryos based on limb morphometry.

Development 145, 1–7.

Okano, J., Suzuki, S., Shiota, K., 2006. Regional heterogeneity in the developing palate:

morphological and molecular evidence for normal and abnormal palatogenesis.

Congenital Anomalies 46, 49–54. https://doi.org/10.1111/j.1741-4520.2006.00103.x

Palmer, A.R., Strobeck, C., 1986. Fluctuating Asymmetry: Measurement, Analysis, Patterns.

Annu. Rev. Ecol. Syst. 17, 391–421.

https://doi.org/10.1146/annurev.es.17.110186.002135

Pantalacci, S., Prochazka, J., Martin, A., Rothova, M., Lambert, A., Bernard, L., et. al., 2008.

Patterning of palatal rugae through sequential addition reveals an anterior/posterior

boundary in palatal development. BMC Developmental Biology 8, 116.

https://doi.org/10.1186/1471-213X-8-116

Pauws, E., Hoshino, A., Bentley, L., Prajapati, S., Keller, C., Hammond, P., et. al., 2009.

Tbx22null mice have a submucous cleft palate due to reduced palatal bone formation and

also display ankyloglossia and choanal atresia phenotypes. Hum Mol Genet 18, 4171–

4179. https://doi.org/10.1093/hmg/ddp368

Percival, C.J., Green, R., Marcucio, R., Hallgrímsson, B., 2014. Surface landmark quantification

of embryonic mouse craniofacial morphogenesis. BMC Developmental Biology 14, 1-12.

https://doi.org/10.1186/1471-213X-14-31

Percival, C.J., Liberton, D.K., Pardo-Manuel de Villena, F., Spritz, R., Marcucio, R.,

Hallgrímsson, B., 2016. Genetics of murine craniofacial morphology: diallel analysis of

the eight founders of the Collaborative Cross. Journal of Anatomy 228, 96–112.

https://doi.org/10.1111/joa.12382

Peterkova, R., Klepacek, I., Peterka, M., 1987. Prenatal development of rugae palatinae in mice:

scanning electron microscopic and histologic studies. Journal of craniofacial genetics and

developmental biology 7, 169–89.

R Core Team, 2021. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria [WWW Document]. URL https://www.R-

project.org/

Sarnat, B.G., 1997. Some methods of assessing postnatal craniofaciodental growth: a

retrospective of personal research. Cleft Palate Craniofac J 34, 159–172.

https://doi.org/10.1597/1545-1569_1997_034_0159_smoapc_2.3.co_2

Schmidt, E.J., Parsons, T.E., Jamniczky, H.A., Gitelman, J., Trpkov, C., Boughner, J.C., et al.,

2010. Micro-computed tomography-based phenotypic approaches in embryology:

procedural artifacts on assessments of embryonic craniofacial growth and development.

BMC Developmental Biology 10, 1-14. https://doi.org/10.1186/1471-213X-10-18

Tamarin, A., 1982. The formation of the primitive choanae and the junction of the primary and

secondary palates in the mouse. American Journal of Anatomy 165, 319–337.

https://doi.org/10.1002/aja.1001650308

Tonge, C.H., McCance, R.A., 1965. Severe undernutrition in growing and adult animals: 15. The

mouth, jaws and teeth of pigs. British Journal of Nutrition 19, 361–372.

https://doi.org/10.1079/BJN19650034

Vora, S.R., Camci, E.D., Cox, T.C., 2015. Postnatal Ontogeny of the Cranial Base and

Craniofacial Skeleton in Male C57BL/6J Mice: A Reference Standard for Quantitative

Analysis. Front Physiol 6, 417. https://doi.org/10.3389/fphys.2015.00417

Welsh, I.C., Feiler, M.E., Lipman, D., Mormile, I., Hansen, K., Percival, C.J., 2024. Dryad

dataset: Palatal segment contributions to midfacial anterior-posterior growth.

https://doi.org/10.5061/dryad.ghx3ffbvb

Welsh, I.C., Hagge-Greenberg, A., O'Brien, T.P., 2007. A dosage-dependent role for Spry2 in

growth and patterning during palate development. Mechanisms of Development 124,

746–761. https://doi.org/10.1016/j.mod.2007.06.007

Welsh, I.C., Hart, J., Brown, J.M., Hansen, K., Rocha Marques, M., Aho, R.J., et. al., 2018. Pbx

loss in cranial neural crest, unlike in epithelium, results in cleft palate only and a broader

midface. Journal of Anatomy 233, 222–242. https://doi.org/10.1111/joa.12821

Welsh, I.C., O'Brien, T.P., 2009. Signaling integration in the rugae growth zone directs

sequential SHH signaling center formation during the rostral outgrowth of the palate.

Developmental Biology 336, 53–67. https://doi.org/10.1016/j.ydbio.2009.09.028

Wilson, R., McGuire, C., Mohun, T., DMDD Project, 2016. Deciphering the mechanisms of

developmental disorders: phenotype analysis of embryos from mutant mouse lines.

Nucleic Acids Res 44, D855-861. https://doi.org/10.1093/nar/gkv1138

Young, N.M., Hu, D., Lainoff, A.J., Smith, F.J., Diaz, R., Tucker, A.S., et. al., 2014. Embryonic

bauplans and the developmental origins of facial diversity and constraint. Development

141, 1059–1063. https://doi.org/10.1242/dev.099994

Yu, K., Ornitz, D.M., 2011. Histomorphological study of palatal shelf elevation during murine

secondary palate formation. Developmental Dynamics 240, 1737–1744.

https://doi.org/10.1002/dvdy.22670

Yu, L., Gu, S., Alappat, S., Song, Y., Yan, M., Zhang, X., et. al., 2005. Shox2-deficient mice

exhibit a rare type of incomplete clefting of the secondary palate. Development 132,

4397–4406. https://doi.org/10.1242/dev.02013

Zelditch, M., Swiderski, D., Sheets, D.H., 2012. Geometric morphometrics for biologists: a

primer, 2nd ed. Elsevier Academic Press, San Diego.

Zhao, H., Feng, J., Ho, T.-V., Grimes, W., Urata, M., Chai, Y., 2015. The suture provides a niche

for mesenchymal stem cells of craniofacial bones. Nat Cell Biol 17, 386–396.

https://doi.org/10.1038/ncb3139

<u>Supporting Information</u>

Supporting Tables and Figures are found as a Supporting .pdf file in association with this

manuscript on the publisher's website.

Tables

Table 1 –Sample size of µCT imaged and morphometrically analyzed embryo specimens by forelimb

based developmental age estimation, as estimated using eMOSS (Musy et al., 2018).

Limb-Based Developmental Age	C57BL/6J
E11	8
E11.5	2
E12	7
E12.5	4
E13	9
E13.5	0
E14	26
E14.5	1
E15	9

676   Table 2 – Median proportional palatal segment lengths for limb-based embryonic ages and

677   postnatal age categories, presented as percentages of total palatal length with Q1 and Q3 in

678   parentheses. Categories with enough samples for pairwise statistical comparisons are included.

679   See Fig. 9 for associated boxplots.

Age Category and Tissue Measured	Sample Size	Primary Palate /Premaxilla	Ant. Sec. Palate / Maxilla	Post. Sec. Palate / Palatine & Pterygoid
E11 epithelial	8	21.8% (19.2, 23.7)	16.9% (15.8, 18.2)	60.9% (60.1, 63.6)
E12 epithelial	7	26.3% (25.6, 27.9)	28.9% (28.1, 29.4)	44.3% (43.3, 45.5)
E13 epithelial	9	28.8% (26.8, 29.6)	31.8% (31.1, 34.3)	39.3% (39.0, 41.5)
E14 epithelial	26	25.0% (24.2, 25.9)	35.3% (34.1, 36.3)	40.1% (39.0, 40.7)
E15 epithelial	9	24.1% (23.5, 24.7)	42.7% (41.7, 43.8)	33.2% (32.5, 34.0)
P1 epithelial	10	18.5% (17.9, 18.9)	45.9% (45.0, 46.9)	35.3% (33.9, 38.2)
P1 skeletal	10	23.2% (23.1, 23.8)	47.4% (46.8, 48.1)	29.4% (28.9, 29.6)
Adult skeletal	20	32.4% (31.7, 33.3)	39.8% (38.5, 40.8)	28.0% (27.2, 28.4)

Figure Legends

Figure 1 – Tissue origins of the midfacial complex and rugae position during secondary palate

morphogenesis – The upper and lower jaws are formed from the frontonasal process (FNP - tan),

and branchial arch 1 derived maxillary and mandibular processes (MxP - pale blue and Mnd –

magenta, respectively). From E11.5 to E15.5, outgrowth and fusion of the medial nasal process

(mnp) with the superficial portion of the MxP frame out the lip and primary (1$^{\text{o}}$) palate, while the

A-P elongation and medially directed growth of the palatal shelves from the internal portion of

the MxP gives rise to the secondary palate (2^o pal – light purple). *Shh* expression (dark blue)

highlights the dynamics of rugae formation and illustrates regional expansion of the anterior

secondary palate. At E11.5, ruga 1 (red arrowheads) forms at the anterior extent of the nascent

palatal shelf and subsequently defines the caudal end of the rugae growth zone (asterisk) where

new rugae form prior to being displaced anteriorly. Additional abbreviation: lateral nasal process

(lnp).

Figure 2 - Flowchart illustrating this study's research design

Figure 3 – Anatomically homologous epithelial landmarks - Landmarks collected for the

quantification of midfacial and palatal shape across limb based embryonic (E) developmental

stages, from (a-c) palatal and (d-f) oblique facial views. Scale bars: 0.5mm. See also Supporting

Table 1.

Figure 4 – Palate segment length measurements – Identification of landmarks used to estimate

midline lengths of the primary palate (yellow), anterior secondary palate (dark purple), and

posterior secondary palate (light purple) on (a) E11.0, (b) E14.0, (c) E15.0 epithelial, (d) P1

epithelial, (e) P1 skeletal (after removing some neurocranial and facial bones to help with

visualization), and (f) adult skeletal surfaces. Scale bars: 1mm. See also Supporting Tables 1 and

2.

Figure 5 – A-P molecular heterogeneity and anatomical relationships during secondary palate

morphogenesis - (a, c, e) RNA WISH for *Shh*, *Shox2*, and *Tbx22* expression, and (b, d, f) sagittal

sections of wildtype embryos imaged with high resolution episcopic microscopy (HREM) to

provide histological resolution of the primary (1^o pal), anterior secondary (ant 2^o), and posterior

secondary palate (post 2^o) relative to surrounding facial structure during midfacial outgrowth at

(a, b) E11.5 (c, d) E13.5, and (e, f) E14.5. Throughout midfacial outgrowth, ruga 1 (red

arrowheads) marks the shared A-P expression boundary of *Shox2* and *Tbx22* in the anterior and

posterior secondary palate, respectively. Double label WISH for *Shox2* (magenta) and *Tbx22*

(blue) at E14.5 highlights mutually exclusive anterior and posterior expression domains

organized relative to ruga 1 (oral view left, sagittal view right). White arrowheads mark the

choanae, black asterisk marks the RGZ. Red dashed lines mark position of coronal planes

passing through the primary-secondary palate junction and posterior wall of the nasal capsule to

highlight coordinated elongation of the anterior secondary palate and overlying sinus cavity (sc).

Regions in dashed boxes of (a) and in HREM images shown enlarged below or to the right,

respectively. Additional abbreviations: mandible (mnd), molar tooth bud (mtb), maxillary

process (MxP), medial nasal process (mnp), tongue (t), greater palatine nerve (gpn),

geschmacksstreifen (gs), palatine (p), posterior domain of *Shh* expression (pd). Scale bars:

250um

Figure 6 – Major axes of embryonic midfacial and palatal shape variation - C57BL/6J embryo

specimens plotted along the (a) first and second principal component axes of shape (i.e., PC1 and

PC2), (b) along PC1 and PC3, and (c) along PC1 and PC4. Circle color indicates the limb based

developmental age categories of each specimen, with the yellow bordered circles indicating the

mean PC scores for each age category with more than one specimen. The proportions of

midfacial and palatal shape variance associated with each principal component are provided.

Figure 7 – Palatal landmark growth trajectories – The positions of palatal landmarks are

identified on representative C57BL/6J specimens at limb-derived developmental ages (a) E11,

(b) E12.5, (c) E14, and (d) E15. (e) The average palatal landmark positions are plotted for each

developmental age category that had more than one specimen. These landmark positions

represent palatal shape after the removal of overall midfacial and palatal scale (as measured by

centroid size) during Procrustes superimposition. The arrows indicate the trajectory of shape

change for each landmark between ages. Black=E11; Light Blue = E12; Orange = E12.5; Green

= E13; Dark Blue = E14; Red = E15. Scale bars: 0.5mm

Figure 8 – Palate segment length across ontogeny – The midline projected mean lengths (in mm)

of the primary palate, anterior secondary palate, and posterior secondary palate for all (a) E11-

E15 specimens and (b) postnatal specimens. (c) Palate segment length relative to overall midline

palate length for the all specimens. Whole-day limb bud derived age categories and postnatal age

categories used for pairwise statistical comparisons are indicated by brackets above the bars.

Comparable segment lengths were collected from epithelial and skeletal surfaces for a single set

of P1 specimens.

Figure 9 - Palate segment length comparisons - Boxplots of palate segment specific lengths (in

mm) for (a) whole-day limb dub derived embryonic age categories and (b) postnatal age

categories. (c) Palate segment length relative to overall midline palate length for all specimens.

Outliers are specimen values that are 1.5 times the interquartile range outside of the 25th or 75th

quartile of that segment's measures. Significant differences are indicated for each age that a

given segment's median value is different than the next younger age category (* when p-value is

< 0.05; ** when p-value < 0.001). Significant differences for a given segment's P1 skeletal

measurement median when compared to the same segment's P1 epithelial measurement (^ when

p-value < 0.05; ^^ when p-value < 0.001). Dashed lines illustrate ontogenetic trends of palatal

segment proportional contributions to overall palatal length.

Figure 10 – Segmental organization of skeletal specification and growth during A-P

morphogenesis of the midface – WISH time course for (a) *Runx2*, (b) *Sp7*, and (c) *Phospho1*

between E11.5 and E15.5. Expression of *Runx2* in osteoprogenitors highlights facial domains

with osteogenic potential during midfacial morphogenesis. Expression of *Sp7* and *Phospho1* in

committed and differentiating osteoblasts delineates the growth dynamics of individual skeletal

anlagen during midfacial morphogenesis. Curved white dashed arrows in (b) and (c) indicate

medial expansion of maxillary anlage. Abbreviations: maxilla (mx), palatine (p), premaxilla

(pmx), pterygoid (ptg), palatal process of the maxilla (ppmx), palatal process of the palatine

(ppp). Scale bars: 250um

Figure 11 – Double label RNA WISH time course for *Shh* (cyan) and *Sp7* (dark purple) between

E12.5 and E15.5 - (a) Expression of *Sp7* in committed and differentiating osteoblasts delineates

the growth dynamics of individual skeletal anlagen and *Shh* expression in rugae provides a

temporally ordered set of A-P landmarks (regions in dashed boxes enlarged below). (b)

Summary model of skeletal growth dynamics during midfacial outgrowth. Growth of the

premaxilla (yellow) and palatine (pale blue) anlagen towards their characteristic shape occurs

largely at the site of initial specification. Following initial specification external to the oral

cavity, the maxilla (purple) grows into the anterior of the anterior secondary palate (curved white

and black dashed arrows in a and b, respectively) towards the position of ruga 1 (red arrowhead)

and palatine as expansion of the anterior secondary palate (double headed red arrow) separates

the primary and posterior secondary palate. (c) Summary of the position of epithelial landmarks

(see also Fig. 3) selected to capture segmental growth dynamics of the primary palate (yellow)

and anterior secondary palate (purple) and posterior secondary palate (pale blue) during

midfacial outgrowth. Abbreviations: geschmacksstreifen (gs), maxilla (mx), palatine (p),

premaxilla (pmx), pterygoid (ptg), palatal process of the maxilla (ppmx), palatal process of the

palatine (ppp). Scale bars: 250um

| E11.0 | E14.0 | E15.0 |

(a) E11.0

(b) E14.0

(c) E15.0

(d) P1 Epithelial

(e) P1 Skeletal

(f) P1 Adult

(b) E12.5
0.5 mm

(c) E14.0
0.5 mm

(d) E15.0
0.5 mm

(a) E11.0
0.5 mm

(e) Shape Changes

(a)

Shh/*Sp7*

E12.5	E13.5	E14.0	E14.5	E15.5
2->3 rugae	4->5 rugae	5->6 rugae	6->7 rugae	8 rugae

Labels: pmx, mx, p, ptg, gs

(b) summary of A-P palate morphogenesis and growth of skeletal anlagen

E11.5   E12.5   E13.5   E14.5   E15.5

Labels: pm, max, ppmx, ppp, pal, ptg

(c) surface landmarks and expansion of A-P domains

skeletal anlage/palate segment

premaxilla/ 1⁰ palate

maxilla/ anterior 2⁰ palate

palatine,pterygoid/ posterior 2⁰ palate

Shh expression

ruga 1

Molecular and histological characerization of palate segmental morphogenesis

E11.5   E13.5   E15.5

pm
max
ppmx
ppp
pal
ptg

skeletal anlage/palate segment

premaxilla/1° palate

maxilla/anterior 2° palate

palatine, pterygoid/posterior 2° palate

Shh expression

ruga 1

Morphometric comparison of palatal segment contribution to jaw length from microCT

E11.0   E14.0   E15.0

Significant differences in palate segment growth dynamics

Proportional Segment Length

E11   E12   E13   E14   E15

Limb Bud Age Estimate (Embryonic Day)