1 **Palatal segment contributions to midfacial anterior-posterior growth**

2 Short running page heading: **Palatal segment contributions to midfacial growth**

3 Ian C. Welsh @X – 1) Program in Craniofacial Biology, 2) Department of Orofacial
4   Sciences and 3) Department of Anatomy, University of California at San Francisco, San
5   Francisco, California 94143, USA
6
7 Maria E. Feiler – Interdepartmental Doctoral Program in Anthropological Sciences, Stony Brook
8   University, Stony Brook, NY 11790
9
10 Danika Lipman @Y – Department of Cell Biology and Anatomy, University of Calgary
11
12 Isabel Mormile – Interdepartmental Doctoral Program in Anthropological Sciences, Stony Brook
13   University, Stony Brook, NY 11790
14
15 Karissa Hansen – 1) Program in Craniofacial Biology, 2) Department of Orofacial Sciences and
16   3) Department of Anatomy, University of California San Francisco, San Francisco, CA
17   94143
18
19 Christopher J. Percival* – Department of Anthropology, Stony Brook University, Stony Brook
20   NY 11794
21
22 * = Corresponding Author: Department of Anthropology, Circle Road, SBS Building S-501,
23   Stony Brook, NY 11994, christopher.percival@stonybrook.edu
24 @X Present Address: The Jackson Laboratory, 600 Main St. Bar Harbor, ME 04609
25 @Y Present Address: Department of Mathematics and Statistics, 612 Campus Place NW,
26   University of Calgary, Calgary, AB
27

Abstract

Anterior-posterior (A-P) elongation of the palate is a critical aspect of integrated

midfacial morphogenesis. Reciprocal epithelial-mesenchymal interactions drive secondary palate

elongation that is coupled to the periodic formation of signaling centers within the rugae growth

zone (RGZ). However, the relationship between RGZ driven morphogenetic processes, the

differentiative dynamics of underlying palatal bone mesenchymal precursors, and the segmental

organization of the upper jaw has remained enigmatic. A detailed ontogenetic study of these

relationships is important, because palatal segment growth is a critical aspect of normal midfacial

growth, can be modified to produce dysmorphology, and is a likely basis for evolutionary

differences in upper jaw morphology. Variation in palatal-segment specific growth may also

underlie known differences in palatal segment proportions between inbred mouse strains. We

completed a combined whole mount gene expression and morphometric analysis of normal

murine palatal growth dynamics and their association with palatal segment elongation and

resulting upper jaw morphology. Our results demonstrated that the first formed palatal ruga (ruga

1), found just posterior to the RGZ, maintained an association with important nasal,

neurovascular and palatal structures throughout early midfacial development; suggesting that

these features are positioned at a proximal source of embryonic midfacial directional growth.

Our detailed characterization of midfacial morphogenesis revealed a one-to-one relationship

between palatal segments and upper jaw bones during the earliest stages of palatal elongation.

Growth of the maxillary anlage within the anterior secondary palate is uniquely coupled to RGZ-

driven morphogenesis that more than doubles the length of this palatal segment prior to palatal

shelf fusion. Our results also demonstrate that the future maxillary-palatine suture, approximated

by the position ruga 1 and consistently associated with the palatine anlage, forms predominantly

via the posterior differentiation of the maxilla within the expanding anterior secondary palate.

Our complementary ontogenetic comparison of three inbred mouse strains identified small but

significant strain-specific differences in early embryonic palatal segment contributions to the

upper jaw. Although early palatal segment specific growth is not primarily responsible for adult

differences in upper jaw morphology between these strains, our ontogenetic series of

measurements provide a useful foundation for understanding the impact of background genetic

effects on facial shape and elongation. In combination, our results provide a novel and

particularly detailed picture of the earliest spatiotemporal dynamics of intramembranous

midfacial skeletal specification and differentiation within the context of the surrounding palatal

segment A-P elongation and associated rugae formation.

Keywords: palatogenesis, secondary palate, rugae, face elongation, *Mus musculus,*

RRID:IMSR_JAX:003715, RRID:IMSR_JAX:001976, RRID:IMSR_JAX:000664

Abbreviations: anterior-posterior (A-P), cranial neural crest (CNC), frontonasal process (FNP),

High resolution episcopic microscopy (HREM), maxillary processes (MxP), medial nasal

processes (mnp), micro-computed tomography (µCT), rugae growth zone (RGZ), transcription

factors (TFs), Whole mount in situ hybridization (WISH)

<u>Introduction</u>

Morphological variation of the midfacial complex, which consists of the nose, upper jaw,

cheek, and palate, is a defining aspect of both intra- and inter-specific differences in facial shape.

Basic facial shape is the result of cranial neural crest (CNC)-derived facial prominence

outgrowth and fusion, which occur between embryonic day (E) 10 and E15 in mice (Fig. 1).

Initial fusion between the medial nasal processes (mnp) of the frontonasal process (FNP) and the

anterior maxillary processes (MxP) gives rise to the primary palate and lip, producing a unified

upper jaw from previously separated tissues. Secondary palate morphogenesis then begins

through outgrowth of the nascent palatal shelf along the medial aspect of the MxP. Significant

palatal growth along the anterior-posterior (A-P) axis is accompanied by vertical growth of the

palatal shelves prior to their elevation and medial fusion dorsal to the tongue, after which the

palatal shelf separates the oral and nasal cavities (Bush and Jiang, 2012; Hammond and Dixon,

2022). Disproportional growth of FNP or MxP prior to fusion can result in cleft lip and palate

(Young et al., 2014), while poor coordination between the multiple growth axes of MxP-derived

tissues can prevent secondary palate closure even when the palatal shelves maintain competency

to fuse (Kouskoura, et al., 2013).

Figure 1 – Tissue origins of the midfacial complex and rugae position during secondary palate morphogenesis – The upper and lower jaws are formed from the frontonasal process (FNP - tan), and branchial arch 1 derived maxillary and mandibular processes (MxP - pale blue and Mnd – magenta, respectively). From E11.5 to E15.5, outgrowth and fusion of the medial nasal process (mnp) with the superficial portion of the MxP frame out the lip and primary (1º) palate, while the A-P elongation and medially directed growth of the palatal shelves from the internal portion of the MxP gives rise to the secondary palate (2º pal – light purple). *Shh* expression (dark blue) highlights the dynamics of rugae formation and illustrates regional expansion of the anterior secondary palate. At E11.5, ruga 1 (red arrowheads) forms at the anterior extent of the nascent palatal shelf and subsequently defines the caudal end of the rugae growth zone (asterisk) where new rugae form prior to being displaced anteriorly. Additional abbreviation: lateral nasal process (lnp).

Species-specific differences in the size, shape, and growth trajectories of facial

prominences are identifiable during the earliest phases of facial morphogenesis, prior to skeletal

differentiation, often presaging interspecies differences in adult facial shape (Selleri and Rijli,

2023). After initial facial prominence fusion, differential A-P growth of FNP and MxP derived

tissues is a major basis for evolutionary differences in midfacial prognathism (Young et al.,

2014); possibly based on the balance of CNC progenitor self-renewal versus osteoblast

differentiation (Hall et al., 2014; Morris and Abzhanov, 2021; Schneider, 2015). The regional

basis for facial elongation varies across taxa; primarily driven by MxP growth in mammals and

by FNP growth in birds (Young et al., 2014), indicating different palatal segment contributions.

The three longitudinal segments of the palate are called the primary palate, anterior

secondary palate, and posterior secondary palate (Fig. 1). The boundary between primary palate

and anterior secondary palate is the posterior edge of the fused mnp. Although A-P organization

of the secondary palate can be defined by the presence of boney versus muscular tissue (i.e. the

hard versus soft palate) or modes of palatal shelf elevation and closure (reviewed by Yu and

Ornitz, 2011), we define secondary palate segments by molecular differences in tissue patterning

and cell signaling competence (Hilliard et al., 2005; Hammond and Dixon, 2022).

The A-P expression domains of certain transcription factors (TFs) and intercellular

signaling molecules within CNC-derived mesenchyme are important for A-P organization and

growth during palatal morphogenesis. For example, *Msx1* and *Shox2* are exclusively expressed in

the anterior secondary palate while *Barx1* and *Tbx22* are exclusive to the posterior secondary

palate within this mesenchyme (Hilliard et al., 2005; Okano et al., 2006; Welsh et al., 2018).

Mutations of *Shox2* or *Tbx22* have resulted in midline clefts of the anterior or posterior palate,

respectively (Yu et al., 2005; Pauws et al., 2009), although regional expression is maintained in

genetically engineered mice with severe segmental growth defects (Welsh et al., 2018). The A-P

expression boundary of these and other factors is found at the first formed palatal ruga (Hilliard

et al., 2005; Li and Ding, 2007; Welsh et al., 2007; Welsh and O'Brien, 2009; Welsh et al., 2018;

Hammond and Dixon, 2022), one of multiple parallel epithelial thickenings on the anterior

secondary palate.

This first ruga (ruga 1 numbered according to Welsh et al., 2007, Welsh and O'Brien,

2009; although Pantalacci et al., 2008 use a different numbering system) forms at E11.5 on the

anterior extent of the nascent secondary palate (red arrow in Fig. 1). Remaining rugae form

sequentially within the rugae growth zone (RGZ) located just anterior to ruga 1 (Welsh and

O'Brien, 2009), via a Turing type mechanism (Economou et al., 2012; Kawasaki et al., 2018).

Rugae are centers of Sonic Hedgehog signaling (SHH), which is critical for anterior secondary

palatal patterning and A-P growth. As the secondary palate completes midline fusion at E15.5,

the last formed ruga appears adjacent to ruga 1 in the middle of the secondary palate while each

precedingly formed ruga is found at an increasingly anterior position along the anterior

secondary palate (Welsh and O'Brien, 2009).

Based upon the dynamics of rugae formation, palate growth, and patterning gene

expression, Pantalacci (et al., 2008) hypothesized that ruga 1 represents a distinct morphological

boundary related to either the future hard versus soft palate boundary or the maxillary-palatine

suture. However, the degree to which rugae formation and regional A-P patterns of secondary

palate gene expression are coordinated with palate skeletal differentiation remain unclear. To

address this knowledge gap, we mapped markers associated with the specification and

differentiation of osteoblasts within the context of secondary palate segmental organization,

across a developmental time course spanning mouse secondary palate morphogenesis and upper

jaw outgrowth (E11.5-E15.5). Our results indicate that ruga 1 is coincident with the future

position of the maxillary-palatine suture, illustrate the one-to-one palatal segment origins of

upper jaw bones, and reveal a previously unappreciated coupling of maxillary formation to rugae

morphogenesis during elongation of the anterior secondary palate.

Given 1) the strong early one-to-one association between forming upper jaw bones and

palatal segment specific growth factor expression and 2) that embryonic morphogenesis is

critical for the production of typical facial shape (e.g., Hilliard et al., 2005; Kaucka et al., 2018;

Yuan et al., 2020), it's possible that early differences in the longitudinal growth of the three

palatal segments underlie inter- and intraspecies differences in upper jaw morphology.

Specifically, we hypothesized that genetically based differences in palate segment-specific A-P

outgrowth would explain previously identified variation in the contribution of the premaxilla,

maxilla, and palatine bones to adult upper jaw morphology of inbred mouse strains.

C57BL/6J (RRID:IMSR_JAX:000664) inbred strain adults have a relatively long

premaxilla, while NOD/ShiLtJ (RRID:IMSR_JAX:001976) have a relatively long maxilla,

making them a straightforward comparison. The PWK/PhJ (RRID:IMSR_JAX:003715) wild-

derived inbred strain also has a relatively long maxilla, but a proportionally long upper jaw for

its small body size (Percival et al., 2016), allowing a test of whether these NOD/ShiLtj-like bone

length proportions are the result of similar palatal segment growth trajectories. One possible

mechanism for the C57BL/6J strain's relatively short adult maxillary bone might be a slower rate

of anterior secondary palate elongation between E11.5 and E15.5. If this type of variation in

early A-P elongation of longitudinal palatal shelf segments is the causal basis for strain specific

differences in upper jaw morphology, we predicted that C57BL/6J mice would have a relatively

long primary palate and a relatively short anterior secondary palate by E15 and that the other two

strains would both display relatively long anterior secondary palates at the same developmental

stage.

Understanding how genetically based differences in early craniofacial development

produce morphological variation within model organisms is an important step in determining

how these developmental processes can be modified to produce evolutionary differences in upper

jaw morphology (Boughner et al., 2008; Hallgrímsson et al., 2009; Fish, 2019; Morris and

Abzhanov, 2021), including facial length and the relative contribution of specific bones to the

upper jaw. Overall, our multifaceted analysis of ontogenetic trends in palatal and facial

elongation provides a novel contextual framework and developmental perspective within which

to evaluate the impact of both non-pathogenic and pathogenic genetic differences on midfacial

growth and differentiation.

Methods

Specimen and Image Acquisition

Animal breeding, specimen collection, and tissue fixation were performed in accordance

with the protocols of the University of California, San Francisco Institutional Animal Care and

Use Committee under protocol approval number AN192776-01F. Mice were socially housed

under a twelve-hour light-dark cycle with food and water *ad libitum*. Additional enrichment was

provided when single housing was required for breeding purposes. Mice were euthanized by CO_2

inhalation followed by cervical dislocation or decapitation. Embryos of C57BL/6J

(RRID:IMSR_JAX:000664; hereafter referred to as C57), NOD/ShiLtJ

(RRID:IMSR_JAX:001976; hereafter referred to as NOD), and PWK/PhJ

(RRID:IMSR_JAX:003715; hereafter referred to as PWK) strains (Jackson Labs, Bar Harbor,

ME) were collected between gestational days E11.5 and E15.5, as determined from copulatory

plug occurrence. Postnatal day one (P1) C57 (n=10) and NOD (n=7) specimens were collected

one day after birth. Embryo and P1 specimens were fixed in 4% PFA and stored in 1x PBS for

micro-computed tomography (μCT) imaging or dehydrated through a graded PBST-MeOH

series and stored at -20°C until use for in situ hybridization. A flowchart provides an overview of

our study design (Fig. 2).

Figure 2 - Flowchart illustrating this study's research design

In Situ Hybridization

In situ hybridization was performed as described in Welsh et al., 2018. cDNA probes for

Phospho1 (nucleotides 274–1739 of RefSeq NM_153104), *Runx2* (nucleotides 1275–2704 of

RefSeq NM_001271627), *Shh* (full length cDNA clone from A. McMahon, University of

Southern California, Los Angeles CA, USA), *Shox2* (nucleotides 870–1603 of RefSeq

NM_130458), *Sp7* (nucleotides 431–1715 of RefSeq NM_130458), *Tbx22* (nucleotides 307–

1547 of RefSeq NM_013665) were linearized and in vitro transcribed to label with either

digoxygenin (DIG) or dintrophenol (DNP). Colorimetric detection of probes used BM purple

(dark blue), BCIP (teal), or MagentaPhos (magenta). Minimally, 3-6 embryos per time point

were processed for each probe analyzed.

DMDD HREM Data

High resolution episcopic microscopy (HREM) data (Geyer et al., 2009) was generated as

part of the deciphering mechanisms of developmental disorders (DMDD) program (Wilson et al.,

2016) and is available from the BioImage Archive at:

https://www.ebi.ac.uk/biostudies/bioimages/studies/S-BIAD490?query=DMDD

Embryonic Palatal Segment Growth Trajectories

Developmental Age Estimation

Given that common mouse strains vary in gestation length and there is variation in

developmental timing within litters, it was necessary to standardize our cross-strain

morphometric analysis by embryonic developmental age rather than gestational age.

Developmental age was estimated for each μCT scanned embryonic specimen using eMOSS, an

application that predicts developmental age from hindlimb bud outlines, based on a previous

analysis of C57 mice (Musy et al., 2018). The resulting limb-based estimates of developmental

age were reported as days since conception, up to two decimal places. We combined similar

developmental age estimates whole- or half-day developmental age categories that include

specimens within 0.25 days of their initial eMOSS estimate (Table 1).

Additional error may be expected when estimating the developmental age of NOD and

PWK embryos using a predictive model built on C57 strain limb bud ontogeny, given the

possibility of differences in the correlation of limb and head development between strains.

Therefore, we also utilized the first principal component of embryonic head shape variation

(PC1), based on the landmark analysis described below, as a proxy for developmental age in

some analyses. PC1 score consistently tracked ontogenetic change in facial and palatal shape

across strains (see Results).

<u>Anatomical Landmark Collection</u>

All embryo facial and palate landmarks were collected within Meshlab (Cignoni et al.,

2008) on minimum threshold-based superficial tissue surfaces (downsampled x2) produced from

the μCT images. Landmarks that could be identified consistently in anatomically homologous

positions across the embryonic period of facial growth were chosen around the nose and whisker

region, the eyes, and along the palate (Fig. 3, Supporting Table 1). The facial landmarks we

collected were previously defined to track developmentally homologous structures across

multiple stages of facial development (Percival, et al., 2014) and our new set of palatal

landmarks were defined similarly. Similar care was taken to define embryonic epithelial tissue

palatal landmarks that could be associated with the bony morphology of the P1 and adult mouse

upper jaw.

<u>Prenatal Geometric Morphometric Analysis</u>

A Procrustes superimposition-based geometric morphometric analysis was used to

quantify the ontogenetic shape change of the palate and face for the embryonic C57 sample and

Figure 3 – Anatomically homologous epithelial landmarks - Landmarks collected for the quantification of midfacial and palatal shape across limb based embryonic (E) developmental stages, from (a-c) palatal and (d-f) oblique facial views. Scale bars: 0.5mm. See also Supporting Table 1.

deviations of the NOD and PWK strains from this C57 baseline. The C57 strain was chosen as a

baseline because it is the most widely utilized inbred strain for genetic analysis and because it

was the strain originally used to train the eMOSS limb bud staging algorithm (Musy et al., 2018).

Embryonic specimens with limb-based developmental age estimates between E11 and E15 were

included in the embryonic shape analysis (Table 1). We performed a geometric morphometric

analysis of facial landmarks using *geomorph* (Adams et al., 2020) and RRPP (Collyer and

Adams, 2018) libraries in R Statistical Software (R Core Team, 2021). Generalized Procrustes

analysis (GPA) aligned specimen landmark sets by translating, scaling, and rotating their

landmark coordinates (reviewed by Zelditch et al., 2012). Embryonic specimen shape analyses

were completed using the symmetric component of Procrustes-aligned specimen landmark

coordinate variation, assuming that most bilateral shape differences between the left and right

sides of a specimen's face are due to random effects associated with developmental noise and

tissue fixation (Palmer and Strobeck, 1986). Thus, symmetrized landmark coordinate data were

interpreted to represent the facial shape defined by a given inbred strain genotype.

The mean facial shape of specimens within each developmental age category were

estimated for each strain. Differences between C57 strain age-specific mean shapes were plotted

to illustrate typical facial/palatal shape growth trajectories during this important period of

secondary palate elongation and midline fusion. A principal components analysis (PCA) was

completed to identify the major axes of shape covariation across the embryonic sample. As with

most PCAs of ontogenetic series, the first principal component (PC1) was strongly associated

with overall specimen size and specimen developmental age. Therefore, we treated it as a proxy

for ontogenetic growth and developmental age. C57-specific regressions represented a baseline

of overall facial/palatal shape change to which the NOD and PWK specimens were compared.

Due to clear inflection points in the slopes of embryonic specimen PC scores when plotted

against PC1, segmented linear regressions were completed separately for PC2, PC3, and PC4

versus PC1 values for the C57 sample.

After removing three NOD specimens with PC1 scores that were substantially lower than

the minimum C57 PC1 score that the C57 regression was based on, we predicted PC2, PC3, and

PC4 scores from each C57, NOD, and PWK specimen's PC1 score based on the C57 baseline

regressions. Then we calculated the PC score residuals of each specimen as the difference

between predicted and measured PC scores. We completed Wilcoxon rank sum tests to identify

significant genotypic differences for PC2, PC3, and PC4 residual values representing the whole

embryonic period at once (i.e. statistical tests were not performed at specific embryonic ages

within the sampled age range). A significant difference in mean PC score residual values

between strains was interpreted as a significant difference in the ontogeny of facial shape for that

pair of strains.

<u>Prenatal Palate Segment Length</u>

The anterior-posterior length of the three major palatal segments (i.e., primary palate,

anterior secondary palate, and posterior secondary palate) were estimated in millimeters from

landmark coordinates without superimposition or scaling. To estimate the length of these

segments along the anterior-posterior axis of the palate, the mean midline position between

bilateral landmark pairs were calculated. Proportional palatal segment lengths were calculated as

the length of a single segment divided by the sum of all three segment lengths for a given

specimen (Fig. 4).

Figure 4 – Palate segment length measurements – Identification of landmarks used to estimate midline lengths of the primary palate (yellow), anterior secondary palate (blue), and posterior secondary palate (green) on (a) E11.0, (b) E14.0, (c) E15.0 epithelial, (d) P1 epithelial, (e) P1 skeletal (after removing some neurocranial and facial bones to help with visualization), and (f) adult skeletal surfaces. Scale bars: 1mm. See also Supporting Tables 1 and 2.

Small strain-specific sample sizes at multiple limb-based developmental ages meant it

was not statistically appropriate to compare proportional palate segment contributions between

inbred strains at single developmental ages. Instead, second-degree polynomial regressions of

proportional palate segment lengths versus specimen facial shape PC1 score were calculated for

the entire C57 embryo sample. Because PC1 score was considered a proxy for developmental

age, these regressions define ontogenetic trajectories of palatal segment contributions to overall

palatal length, which were used as a baseline for comparison with NOD and PWK proportional

segment lengths. After removing three NOD specimens with PC1 scores that were substantially

lower than the minimum C57 PC1 score that the C57 regressions were based on, we predicted

the proportional length of each palatal segment for all specimens based on the C57 regressions.

The residual differences in proportional palatal segment lengths between predicted and measured

values representing the whole embryonic period at once were compared across the three strains

using Wilcoxon rank sum tests. Statistical tests were not performed at specific embryonic ages

within the sampled age range. A significant difference in mean residual values was interpreted to

indicate a higher or lower proportional contribution of a given palate segment to overall palatal

length, either across the entire embryonic period or during some major portion of this embryonic

period.

Postnatal Palate Segment Length Comparisons

To determine if embryonic differences in proportional palate segment length remained

consistent postnatally and to investigate the degree of correspondence between surface epithelial

and skeletal measures of palatal variation, postnatal specimen landmarks were collected on

minimum threshold-based surfaces produced from μCT scans. Superficial epithelial surfaces of

P1 specimens were produced using Amira (downsampled x2), and epithelial landmarks

(Supporting Table 1) were collected within Meshlab (Cignoni et al., 2008).

Minimum threshold-based skeletal surfaces of P1 and adult specimens were produced

using 3D Slicer (Fedorov et al., 2012) after Gaussian blur image filtering (sigma set to 0.01 for

P1; sigma set to 0.02 for adult). We identified skeletal anatomical landmarks that most closely

matched the palate segment landmarks defined on surface epithelium (Supporting Table 2).

Landmarks were collected for C57 (n=10) and NOD (n=8) P1 specimens and for C57 (n=20),

NOD (n=18), and PWK (n=18) adult specimens. Midline palate segment lengths and associated

upper jaw bone lengths were measured from mean midline positions of bilateral landmark pairs

without scaling or superimposition. Proportional palate segment lengths were calculated as

individual segment lengths divided by sum of the three midline palatal segment lengths (Fig. 4).

Wilcoxon rank sum tests were used to identify significant differences in proportional palate

segment lengths between strains within single age and tissue type categories.

<u>Results</u>

Secondary Palate A-P Growth and Ruga 1's Position at a Facial Growth Center

Close relationships between different anatomical domains underlying midfacial

outgrowth are found centered near ruga 1 (& Supporting Fig. 1, red arrowheads). Our embryonic

series confirmed that ruga 1 formed within the MxP from a domain of *Shh* expression at the

anterior extreme of the nascent secondary palate. As previously shown, other rugae then formed

anterior to ruga 1 at the RGZ (Fig. 5c, asterisk) as the anterior secondary palate elongated,

anteriorly displacing the primary palate and external nares. A separate domain of *Shh* expression

posterior to ruga 1 gave rise to the geschmacksstreifen (gs) and sensory papilla arrayed across

the most posterior palate. Our embryonic series also confirmed that ruga 1 is consistently found

at the *Shox2/Tbx22* gene expression boundary between the anterior and posterior secondary

palates during this critical embryonic period of palatal development (Fig. 5 & Supporting Fig. 1).

We identified a previously underappreciated association between ruga 1 and the posterior

wall of the nasal capsule. At E11.5, ruga 1 formed adjacent to the primary choanae (Fig. 5a &

Supporting Fig. 1a, white arrowheads); the posterior openings of the nasal passages, which

Figure 5 – A-P molecular heterogeneity and anatomical relationships during secondary palate morphogenesis - (a, c, e) RNA WISH for *Shh*, *Shox2*, and *Tbx22* expression, and (b, d, f) sagittal sections of wildtype embryos imaged with high resolution episcopic microscopy (HREM) to provide histological resolution of the primary (1° pal), anterior secondary (ant 2°), and posterior secondary palate (post 2°) relative to surrounding facial structure during midfacial outgrowth at (a, b) E11.5 (c, d) E13.5, and (e, f) E14.5. Throughout midfacial outgrowth, ruga 1 (red arrowheads) marks the shared A-P expression boundary of *Shox2* and *Tbx22* in the anterior and posterior secondary palate, respectively. Double label WISH for *Shox2* (magenta) and *Tbx22* (blue) at E14.5 highlights mutually exclusive anterior and posterior expression domains organized relative to ruga 1 (oral view left, sagittal view right). White arrowheads mark the choanae, black asterisk marks the RGZ. Red dashed lines mark position of coronal planes passing through the primary-secondary palate junction and posterior wall of the nasal capsule to highlight coordinated elongation of the anterior secondary palate and overlying sinus cavity (sc). Regions in dashed boxes of (a) and in HREM images shown enlarged below or to the right, respectively. Additional abbreviations: mandible (mnd), molar tooth bud (mtb), maxillary process (MxP), medial nasal process (mnp), tongue (t), greater palatine nerve (gpn), geschmacksstreifen (gs), palatine (p), posterior domain of *Shh* expression (pd). Scale bars: 250um

initially occur at the boundary of primary and secondary palatal segments (Tamarin, 1982).

Elongation of the anterior secondary palate then coincided with the formation and expansion of

the overlying nasal capsule. As the primary choanae and nasal capsule elongated between E11.5

and E15.5, ruga 1 and the posterior wall of the nasal capsule remained in approximately the same

coronal plane (represented by position of red dashed lines in Fig. 5b, 5d, 5f and Supporting Fig.

1). Ruga 1 was also found coincident with a gross morphological inflection point of the palatal

tissue where the anterior secondary palate and overlying nasal capsule exhibited an inferiorly

accentuated angle relative to the cranial base, while the posterior secondary palate did not (Fig.

5d-f, Supporting Fig. 1).

The greater palatine neuro-vascular bundle was also consistently associated with ruga 1

across this embryonic period. The greater palatine nerve and artery were found entering the

palatal shelves from a location immediately dorsal to ruga 1 at E12.5 and this spatial relationship

was maintained until the greater palatine foramen formed by the articulation of maxilla and

palatine bones around the palatine neuro-vascular bundle at the maxillary-palatine suture (Fig.

5f, Supporting Fig. 1). In combination, the consistent associations between ruga 1, the anterior-

posterior secondary palate boundary, the posterior margin of the nasal capsule, and greater

palatine neurovascular bundle suggested that these features are positioned at a proximal source

of directional growth that the elongating anterior secondary palate and nasal capsule extend away

from between E11.5 and E15.5.

A complementary 3D morphometric analysis of midfacial and palatal epithelial

landmarks was used to quantify the integrated morphogenetic processes occurring during this

critical developmental period. The first major principal axis of facial shape variation (PC1) for

all limb bud staged E11-E15 specimens represented 70% of shape variance and was associated

with general ontogenetic growth during this time window, as illustrated for the C57 sample (Fig.

6) and for all three inbred strains (Supporting Fig. 2). Additional principal axes of variation for

C57 embryos suggested shifts in the direction of ontogenetic shape changes over developmental

time, with one inflection noted along PC2 (Fig. 6a), two along PC3 (Fig. 6b), and three along

PC4 (Fig. 6c). Average developmental age category shape differences were illustrated as

landmark specific ontogenetic shifts in palate shape (Fig. 7) and overall midfacial shape

(Supporting Fig. 3).

Between developmental stages E11 and E15, each of the three major palatal segments

Figure 6 – Major axes of embryonic facial shape variation - C57 embryo specimens plotted along the (a) first and second principal component axes of shape (i.e., PC1 and PC2), (b) along PC1 and PC3, and (c) along PC1 and PC4. Circle color indicates the limb based developmental age categories of each specimen, with the yellow bordered circles indicating the mean PC scores for the C57 mice of that age category. The proportions of facial shape variance associated with each principal component are provided. The principal component axes were estimated from the full sample of E11-E15 developmental age embryos, including C57, NOD, and PWK samples, although only C57 samples are plotted in this figure. The same plot, but incorporating all three inbred strains is available as Supporting Figure 2.

grew in length, but the anterior secondary palate grew proportionally more than the primary

palate and posterior secondary palate (Figs. 5 & 7). The primary palate represented ~20% of the

total palate length, the anterior secondary palate ~20%, and the posterior secondary palate ~60%

for C57 embryos soon after the start of A-P palate elongation. After E12.5, secondary palate

length (relative to overall facial size) increased noticeably with each developmental stage. By the

time of midline palatal fusion, the primary palate represented ~25% of palate length, the anterior

Figure 7 – C57 palatal landmark growth trajectories – The positions of palatal landmarks are identified on representative C57 specimens at limb-derived developmental ages (a) E11, (b) E12.5, (c) E14, and (d) E15. (e) The average palatal landmark positions are plotted for each developmental age category that had more than one C57 specimen. These landmark positions represent palatal shape after the removal of overall facial scale during Procrustes superimposition. The arrows indicate the trajectory of shape change for each landmark between ages. Black=E11; Light Blue = E12; Orange = E12.5; Green = E13; Dark Blue = E14; Red = E15. Scale bars: 0.5mm

secondary palate represented >40%, while the posterior secondary palate represented <35%.

Therefore, between E11 and E15, the proportional contribution of the primary palate increased

by a quarter while posterior secondary palate contribution decreased by nearly a third. Strikingly,

the proportional contribution of the anterior secondary palate to overall palate length doubled

during this period.

Segmental Differences in Skeletal Specification, Differentiation, and Growth During

Secondary Palate Morphogenesis

389   We performed a WISH analysis of *Runx2*, *Sp7*, and *Phospho1* expression () alongside

390   double labeled WISH of *Shh* and *Sp7* between E12.5 and E15.5 () to characterize bone

391   specification and differentiation in relation to A-P growth dynamics of secondary palate

392   segments. The position of rugae, which provided an ordered set of temporally specific

Figure 8 – Segmental organization of skeletal specification and growth during A-P morphogenesis of the midface – WISH time course for (a) *Runx2*, (b) *Sp7*, and (c) *Phospho1* between E11.5 and E15.5. Expression of *Runx2* in osteoprogenitors highlights facial domains with osteogenic potential during midfacial morphogenesis. Expression of *Sp7* and *Phospho1* in committed and differentiating osteoblasts delineates the growth dynamics of individual skeletal anlagen during midfacial morphogenesis. Curved white dashed arrows in (b) and (c) indicate medial expansion of maxillary anlage. Abbreviations: maxilla (mx), palatine (p), premaxilla (pmx), pterygoid (ptg), palatal process of the maxilla (ppmx), palatal process of the palatine (ppp). Scale bars: 250um

anatomical features, allowed for the identification of previously unappreciated associations

between the palatal segments and developing palatal bones.

Broad *Runx2* expression in early facial prominences included osteogenic domains (e.g.,

premaxilla within the primary palate) and later became more restricted to regions adjacent to the

molar tooth bud and medially along the adjoining palatal shelves (Fig. 8a). Osteoblast expression

of both *Sp7* and *Phospho1* more precisely delineated the morphology of the emerging midfacial

skeleton (Fig. 8b-c). Palatal processes of the maxilla and palatine bones formed within the

medial region of the elevated palatal shelves, a domain that maintained high levels of *Runx2*

expression during secondary palate growth (Fig. 8a), suggesting maintenance of a less

differentiated osteoprogenitor population until palatal elevation and fusion was complete.

At E11.5, *Sp7* was strongly expressed in bilateral domains within the mnp of the forming

primary palate. Between E12.5 and E15.5, presaging premaxilla morphology, *Sp7* expression

formed a cup-like region surrounding the developing incisors (Fig. 8b). *Phospho1* was first

observed in the forming premaxilla at E12.5 (Fig 8c), reflecting a temporal lag between initial

osteoblast commitment and later differentiation. The premaxilla acquired its characteristic adult

morphology within the primary palate segment in which it was initially specified.

Within the secondary palate, this simple developmental sequence was repeated for the

palatine and pterygoid bones within the posterior secondary palate, but development of the

maxilla within the anterior secondary palate was unique. At E11.5, *Sp7* expression in the MxP

was observed in two domains. The anterio-lateral domain, which abutted the point of fusion with

the mnp (i.e., the primary palate) and extended laterally away from the oral cavity and choanae,

corresponded to the maxillary anlage. The posterior domain that spanned the length of the

Figure 9 – Double label RNA WISH time course for *Shh* (cyan) and *Sp7* (dark purple) between E12.5 and E15.5 - (a) Expression of *Sp7* in committed and differentiating osteoblasts delineates the growth dynamics of individual skeletal anlagen and *Shh* expression in rugae provides a temporally ordered set of A-P landmarks (regions in dashed boxes enlarged below). (b) Summary model of skeletal growth dynamics during midfacial outgrowth. Growth of the premaxilla (yellow) and palatine (pale blue) anlagen towards their characteristic shape occurs largely at the site of initial specification. Following initial specification external to the oral cavity, the maxilla (purple) grows into the anterior of the anterior secondary palate (curved white and black dashed arrows in a and b, respectively) towards the position of ruga 1 (red arrowhead) and palatine as expansion of the anterior secondary palate (double headed red arrow) separates the primary and posterior secondary palate. (c) Summary of the position of epithelial landmarks (see also Fig. 3) selected to capture segmental growth dynamics of the primary palate (yellow) and anterior secondary palate (purple) and posterior secondary palate (pale blue) during midfacial outgrowth. Abbreviations: geschmacksstreifen (gs), maxilla (mx), palatine (p), premaxilla (pmx), pterygoid (ptg), palatal process of the maxilla (ppmx), palatal process of the palatine (ppp). Scale bars: 250um

forming secondary palate, from the anterior limit abutting the choanae to the posterior-most

palatal edge, gave rise to the palatine and pterygoid anlagen.

The antero-lateral domain of *Sp7* expression adjacent to the primary palate was initially

external to the secondary palate at the site of the future zygomatic plate. However, this

expression domain later expanded medially (E12.5-E13.5) and then posteriorly (E13.5-E15.5)

within the growing anterior secondary palate (white curved dashed arrows in Fig. 8b). This

expansion occurred after initial A-P growth of the anterior secondary palate and formation of 3-4

rugae anterior to ruga 1, suggesting that it is coupled with RGZ growth dynamics (Fig. 9a). The

fact that *Phospho1* expression closely followed *Sp7* spatial dynamics further supported the idea

that these *Sp7* expression patterns represent maxillary bone anlagen formation and expansion

(white curved dashed arrow in Fig. 8c). The maxillary anlage continued growing posteriorly

towards ruga 1 concomitant with rugae formation (white and black curved dashed arrows in Fig.

9a-b), within the elongating secondary palate.

The posterior domain of *Sp7* expression that initially spanned the length of the secondary

palate at E11.5 was displaced posteriorly by expansion of the anterior secondary palate between

E12.5 and E15.5. During this period, it separated into two subdomains. The anterior subdomain

remained associated with the posterior wall of the nasal capsule and gave rise to the palatine

bone, while the posterior subdomain formed both the medial and lateral pterygoid processes (Fig.

8b-c). Ruga 1 and the palatine anlage initially formed at the anterior extent of the secondary

palate at E11.5 and maintained proximity to each other throughout palatal development even as

both structures were displaced posteriorly (Fig. 9a).

The palatine and pterygoid anlagen grew to acquire their characteristic adult morphology

within the posterior secondary palate segment where their precursor domains were initially

specified, mirroring premaxillary anlagen formation within the primary palate. This differs from

growth of the maxillary anlage that was initially specified external to the secondary palate at the

site of the future zygomatic plate and only later extended into the anterior secondary palate,

following a period of initial elongation, to give rise to the majority of the maxillary bone.

These results indicated that the posterior portion of the secondary palate and its

associated anlage were present within the secondary palate at the onset of its morphogenesis at

E11.5, while the anlage associated with the anterior secondary palate (i.e. the maxilla) did not

contribute substantially to the secondary palate until later (Fig. 9b). They also indicated that the

eventual meeting of the maxilla and palatine anlagen was an asymmetric process achieved

predominantly by posterior growth of the maxilla towards the palatine (white and black curved

dashed arrows in Fig. 9-b). By E15.5, the maxilla and palatine anlagen met to form the

maxillary-palatine suture subjacent to ruga 1 (see also Supporting Fig. 1).

Our results collectively revealed that the segmental relationship between three palatal

regions and associated bone anlagen remained consistent throughout the period of palatal

elongation, although the maxillary anlage was not present within the anterior secondary palate

until part way through this period. Importantly, these results also highlighted the previously

unappreciated correlation of maxillary growth to rugae formation as well as to the greater

proportional expansion of the anterior secondary palate between E11.5 and E15.5 (summarized

in Fig 9C).

Interstrain Comparisons of Facial and Palatal Growth

We completed a morphometric comparison of C57, NOD, and PWK strain embryonic

midfacial shapes and a comparison of proportional palatal segment lengths across embryonic and

postnatal stages to test the hypothesis that early differences in palatal segment growth contribute

to variation in adult midfacial shape, including strain specific differences in proportional bone

contributions to upper jaw length.

Because developmental speed varies between mouse strains and within litters, we

standardized our cross-strain ontogenetic analysis using limb bud outline-based embryonic

developmental age rather than gestational age. Limb bud developmental ages correlated strongly

with the number of days after plug identification for each strain. Though the correlation

coefficients for C57 mice (0.98), NOD (0.93) and PWK (0.90) were high, there was a larger

difference between these two embryonic age estimates for our PWK samples. The C57 specimen

limb-based developmental ages were an average of 0.26 days younger than our copulatory plug

estimates, while NOD and PWK limb ages were an average of 0.30 days and 1.09 days younger,

respectively. The major divergence for PWK was likely driven by a slower speed of embryonic

development in this wild-derived strain compared to the two common lab strains (Murray et al.,

2010). Even so, standardization by developmental age should allow for improved direct

comparisons between embryos of all strains.

We compared the midfacial shape of NOD and PWK specimens with C57 specimens

along the principal axes of facial shape variation. This comparison indicated that mice with

similar limb-based developmental age estimates tended to fall near each other along the first and

second principal components, regardless of strain. As these axes represented 82% of facial shape

variance across our sample, all mouse strains displayed major similarities in ontogenetic shape

change across this period (Supporting Fig. 2a). However, across ontogenetic time (as represented

by PC1 score) NOD and PWK specimens had significantly lower PC3 values compared to the

C57 baseline, based on a comparison of regression residual values representing the entire

sampled embryonic period (Fig. 10b). PWK specimens also had significantly lower PC4 values

(Fig. 10c).

We visualized how these identified significant differences in PC3 or PC4 scores were

reflected within facial and palatal morphology by comparing shapes represented at the minimum

and maximum end of these major shape axes (Fig. 11), with strain representative whole mount

(Supporting Fig. 4) and μCT surface (Supporting Fig. 5) images provided for reference. A

minimum PC3 score was associated with the two nostril landmarks (LM7 & 8) being relatively

Figure 10 – Specimen PC score residuals – The principal component (PC) score residuals for each specimen, relative to the scores predicted from C57 strain specific segmented linear regressions of (a) PC2, (b) PC3, and (c) PC4 on the specimen's PC1 score. In this way, the C57 facial shape growth trajectory serves as a baseline with which to compare the shape of NOD and PWK specimens, based on the fact that the PC1 axis is strongly associated with facial shape ontogeny from developmental age E11 to E15.

distant in the superior/inferior and lateral directions. Lower PC3 score was also associated with

relative proximity of the posterior whisker margin (LM6) to the anterior canthus (LM4) (h).

Within the palate, lower PC3 scores were associated with a relatively medial anterior secondary

palatal shelf, suggesting more medial palatal outgrowth (proportional to overall palatal length) at

a given developmental stage (Fig. 11b). The superior-inferior position of the palatal landmarks

indicated that the secondary palate was more highly angled (less flat) for specimens with low

PC3 scores (Fig. 11e). Because of their generally lower PC3 scores, we anticipated more vertical

nostrils in NOD and PWK mice with more highly angled palatal shelves in greater medial

proximity to each other, when compared to C57 mice within a given developmental age.

A minimum PC4 score was associated with the whole nasal region (LM1, LM8) being

positioned more inferiorly and the posterior edge of the whisker region (LM3, LM6) more

superiorly when compared to the other facial and palatal features (Fig. 11i, Supporting Figs. 4 &

5). Within the palate, a low PC4 score was associated with a relatively anterior location of ruga 1

Figure 11 – Facial shape variation associated with PC3 and PC4 – Representation of the pattern of shape variation associated with the third major axis and fourth major axis of embryonic facial/palatal shape variation, from (a-c) palatal, (d-f) parasagittal, and (g-i) facial views. Superficial tissue landmarks are identified on an example E14.5 specimen to assist with interpretation of the landmark vectors that represent differences in minimum (circle) and maximum (end of line) PC scores. See Supporting Fig. 2 for a plot of embryonic specimens along these major axes of embryonic shape variation. Scale bars: 0.5mm

between anterior and posterior secondary palate (LM20 & 24) and a relatively posterior position

of the posterior edge of the posterior secondary palate (LM21 & 25 within Fig. 11c). Low PC4

scores were also associated with more medially positioned anterior secondary palate landmarks

that are closer to the boundary with the primary palate. Because of their generally lower PC4

scores, this suggested that PWK embryos display a relatively low superior nasal region, a

medially expanded anterior secondary palatal shelf with a relatively short anterior secondary

palate and relatively long posterior secondary palate, when comparisons are made within a given

developmental age.

A comparison of embryonic palatal segment lengths (Fig. 4a-c) across C57, NOD and

PWK samples indicated that all three mouse strains display parallel ontogenetic changes 1) in

overall palatal length and 2) in the proportional contributions of palatal regions to overall palatal

length during the period of early palatal morphogenesis (, Supporting Fig. 6). This includes

generally increased proportional contributions of the anterior secondary palate and decreased

proportional contributions of the posterior secondary palate between E11 to E15, as described

above for C57 mice. However, small but significant differences in palatal segment proportions

between mouse strains were identified within a comparison of proportional palatal segment

length versus facial shape PC1 score (a proxy for developmental age). These comparisons were

completed based on regression residuals representing the entire sampled embryonic age range

and were not completed at specific embryonic ages. The NOD and PWK primary palate

proportional length residuals were significantly lower than for C57 (p-values: 0.012 (NOD vs

C57), <0.001 (PWK vs C57)) (Fig. 13a), while they also had significantly higher posterior

secondary palate residuals than C57 (p-values <0.001) (Fig. 13c). This indicated that the primary

palate contributed relatively less and the posterior secondary palate contributed proportionally

more to total embryonic palate length in these two mouse strains, although visual assessment

suggested these differences may not be apparent at the earliest measured developmental ages

(Fig. 12) or for specimens with the lowest PC1 scores (Supporting Fig. 6). Significantly lower

anterior secondary palate residuals for NOD and PWK (p-values: 0.029 (NOD vs C57), 0.023

(PWK vs C57)) indicated that the anterior secondary palate represented a lower proportion of the

total palate length within these strains (Fig. 13b), although these differences seemed less

Figure 12 – Embryonic palate segment length across ontogeny – The midline projected strain-specific mean lengths of the primary palate, anterior secondary palate, and posterior secondary palate, for all E11-E15 specimens binned into 0.1 day limb bud derived developmental age categories and sorted by genotype. Raw length values (in millimeters) are presented above, while palatal segment length relative to overall midline palate length are presented below.

pronounced than those noted for the other two segments (Fig. 12, Supporting Fig. 6). Taken

together, these trends suggested that C57 mice typically had proportionally longer primary

palates and proportionally shorter posterior secondary palates than NOD and PWK mice during

this embryonic period, with a trend towards longer anterior secondary palates by the time of

medial palatal shelf fusion (Fig. 13).

Interstrain Comparisons of Postnatal Palatal Segments

540   We compared C57 and NOD proportional palatal segment lengths at P1 based on

epithelial surface landmarks and skeletal landmarks of associated bony elements. Based on the

epithelial surface landmarks (Fig. 4d), C57 had a proportionally shorter anterior secondary palate

and a proportionally longer posterior secondary palate (p-values <0.01) than NOD at P1 (Fig. 14;

Table 2). This differed from the embryonic pattern where C57 specimens generally displayed

proportionally shorter posterior secondary palates than NOD specimens. The significant

differences in epithelially-measured proportional segment contributions between NOD and C57

mice at P1 appeared largely based on differences in the measured length (in mm) of the posterior

secondary palate (Fig. 14).

A comparison of proportional palatal bone lengths in P1 samples (Fig. 4e) indicated that

C57 has a proportionally shorter premaxilla (associated with primary palate) and proportionally

longer palatine/pterygoid bones (associated with posterior secondary palate) (p-values <0.01)

than NOD (Fig. 14; Table 2). A similar comparison of proportional palatal bone lengths (Fig. 4f)

in 8-12 week old adult specimens of all three strains revealed that the premaxillary (primary

palate) contribution to total palate length is significantly lower in PWK (p-value: < 0.001) and

Figure 13 - Specimen palate segment length residuals – The proportional contributions of each palate length
segment (shapes), as compared to the proportional contributions predicted from C57 strain specific linear
regressions (curved lines) of proportional (a) primary palate, (b) anterior secondary palate, and (c) posterior
secondary palate midline lengths versus the specimen's PC1 score. Vertical lines represent the residual of each
specimen from C57 regression. In this way, the C57 palatal segment growth trajectory serves as a baseline that
NOD and PWK specimens were compared to, based on the fact that the PC1 axis is strongly associated with
facial shape ontogeny from E11 to E15.

NOD (p-value: 0.044) compared to C57, with the opposite true for the maxillary (anterior

secondary palate) portion of the palate (p-values: < 0.001) (Fig. 14). The palatine/pterygoid

(posterior secondary palate) portion was proportionally shorter in PWK and NOD when

compared to C57 (p-values: < 0.001) (Fig. 14). The PWK average proportional measures of

premaxilla and maxilla showed a more extreme divergence from C57 than in the NOD mouse

comparison.

561     Overall, there were some ontogenetic changes in how strains differed in proportional

palate lengths, either between the embryonic period and P1 or between P1 and adulthood. NOD

and C57 showed parallel changes in palatal bone contributions across the postnatal period, where

the posterior secondary palate contribution remained fairly stable (~27-29%), but the primary

palate contribution increased from ~24% to ~32% and the anterior secondary palate contribution

decreased from ~47% to ~40-42% (Fig. 14; Table 2). During the postnatal period, the entire

mouse and its palate grew substantially, but premaxilla growth contributed more than the maxilla

growth to overall postnatal palatal elongation. Although the A-P growth of the anterior

secondary palate was a critical driver for midfacial growth between E11.5 and E15.5, outgrowth

of the premaxilla plays a larger role in postnatal midfacial outgrowth.

Discussion

573     Our results provided a multifaceted characterization of 1) normal murine midfacial

574   growth dynamics within the secondary palate and upper jaw bones during the earliest stages of

575   palatal growth and 2) subsequent changes in upper jaw bone proportions in newborn and adult

576   mice. By tracking the position of the RGZ, sequential palatal rugae formation, and palatal bone

577   precursor populations across the earliest phases of palatal A-P elongation, our results illustrated

Figure 14 – Interstrain comparisons of palate segment lengths in older specimens – The midline projected lengths of the primary palate, anterior secondary palate, and posterior secondary palate, based on epithelial landmark measures for all specimens of limb based developmental stages E14 and E15, as well as for P1 specimens. Analogous length measures of primary palate derived premaxilla, anterior secondary palate derived maxilla, and posterior secondary palate derived palatine/pterygoid bones are presented for all P1 and adult specimens. Raw length values (in millimeters) are presented above, while palatal segment length relative to overall midline palate length are presented below.

that the first-formed palatal ruga (i.e., ruga 1), which sits at an important border of regulatory

gene expression is coincident with important nasal, neurovascular and palatal structures

throughout early midfacial development. For example, ruga 1 represents a consistent

morphological boundary between the presumptive maxilla of the anterior secondary palate and

the presumptive palatine and pterygoid bones of the posterior secondary palate. This association

suggested that the process of rugae morphogenesis is coupled to maxillary osteogenesis during

anterior secondary palate expansion and that ruga 1 approximates the future position of the

maxillary-palatine suture from the time of its formation.

Our ontogenetic comparison of epithelial and skeletal palatal anatomy across three mouse

strains identified small, but significant strain-specific differences in early palatal morphogenesis

and normal midfacial outgrowth. These differences provide a useful foundation for

understanding the impact of background genetic effects on facial shape and elongation.

However, embryonic differences in palatal segment proportions did not match adult strain-

specific differences in the contributions of the premaxilla, maxilla, and palatine/pterygoid bones

to upper jaw length.

Normal Upper Jaw and Palatal Elongation

It is well known that A-P elongation of the palate is a necessary component of the

integrated processes of midfacial outgrowth. However, within mammals, we previously lacked a

detailed picture of the earliest spatiotemporal dynamics of intramembranous midfacial skeletal

specification and differentiation within the context of the surrounding palatal segments and rugae

formation. Our combined whole mount gene expression and morphometric analysis indicated

that the relationship between three longitudinal palate segments and associated upper jaw bones

was already established during the earliest phases of palatal morphogenesis. These results

illustrated the substantial contribution of embryonic anterior secondary palate elongation to

overall palate length in mice and likely in other mammals. Anterior secondary palate elongation

was also directly associated with rugae formation at the RGZ and elongation of the maxillary

bone primordium.

Beyond verifying that ruga 1 forms at an important gene expression boundary between

the anterior and posterior secondary palate, our results clarified that ruga 1 represented a stable

morphological boundary positioned near the future maxillary-palatine suture, as opposed to the

alternate hypothesis that it is positioned between hard and soft palate (Pantalacci et al., 2008).

Ruga 1 formed at the earliest stages of anterior secondary palate elongation, then maintained a

position at the anterior edge of the palatine bone anlage as it was posteriorly displaced by

expansion of the anterior secondary palate. Anterior to ruga 1, the maxillary bone osteogenic

population expanded from the site of initial specification (i.e. the zygomatic plate) into the

secondary palate after ruga 1 was posteriorly displaced during anterior secondary palate

elongation. Given that the secondary palate initially only contained osteogenic domains of the

posterior secondary palate (palatine and pterygoid precursors), these results suggested that

cellular cues in the proximity of ruga 1 may initially inhibit maxillary bone formation within the

secondary palate. Additionally, expansion of the palatal portion of the maxillary anlage was

uniquely associated with RGZ dynamics and anterior secondary palate elongation, suggesting

that RGZ morphogenesis plays a major role in determining the proportional contribution of the

maxilla to the upper jaw within a broad evolutionary context. This hypothesis could be tested

within an evo-devo context by characterizing variation in palatal *Shh* expression between

mammals and avians that have drastic differences in maxillary morphology, including an

obligate avian cleft palate. Alternatively, it could be dissected genetically via analysis of *Shh*

expression and rugae morphogenesis in mice with altered A-P skeletal patterning, such as the

Pbx CNCC mouse mutants (Welsh et al., 2018).

Differences in upper jaw suture anatomy may also be related to segmental differences in

early bone growth. The premaxilla and maxilla were specified in adjacent tissue domains that

maintained close apposition throughout palate morphogenesis, suggesting equal contributions to

the formation of the premaxillary-maxillary suture. Conversely, the palatal portion of the maxilla

grew posteriorly towards the palatine anlage at ruga 1, suggesting that maxillary growth (and

likely RGZ dynamics) played a larger role than palatine growth in determining maxillary-

palatine suture position. Different growth dynamics at these sutures may lead to a more vertical

premaxillary-maxillary suture versus a more oblique maxillary-palatine suture with substantial

A-P overlap of the maxilla and palatine bones.

Because suture formation provides a critical niche for facial skeletal progenitors (Zhao et

al., 2015), the way palatal segments contribute to palatal sutures may contribute to variation in

postnatal dynamics of midfacial growth and remodeling (Enlow and Bang, 1965; Kurihara et al.,

1980; Sarnat, 1997; Martinez-Maza et al., 2013; Vora et al., 2015; Maga, 2016). The widespread

presence of Gli1+ mesenchymal stem cells and nascent bone within the adult maxillary-palatine

suture and the maxillary bone, but not in the palatine bone (Luo et al., 2019) provides further

support for this hypothesis. Given the established role of calvarial sutures in directing postnatal

craniofacial bone growth, further comparisons of osteoprogenitor dynamics during premaxillary-

maxillary and maxillary-palatine suture formation are warranted.

Palatal Growth as a Basis for Morphological Variation

Given its central contribution to total palatal elongation, variation in RGZ regulated A-P

growth of the anterior secondary palate may contribute substantially to the range of prognathism

observed amongst mammals (Young et al., 2014). Support for this idea comes from the fact that

more prognathic species tend to have more rugae: flat faced humans typically have 3-6 palatal

rugae (Hauser et al., 1989; Jayasankar et al., 2016), mice have 8-9 rugae (Peterkova et al., 1987),

and prognathic pigs have 20-25 rugae (Tonge and McCance, 1965). Based on 1) the expectation

that interspecies differences are extensions of milder intra-species differences and 2) the direct

early associations between palatal segments and upper jaw bone morphogenesis, we

hypothesized that strain-specific contributions of individual bones to upper jaw length (Percival

et al., 2016) would be produced by strain-specific differences in early embryonic A-P palatal

segment elongation.

We identified small, but significant strain-associated differences in facial shape and

palatal segment length. Embryonic facial shape differences between strains (Figs. 12 & 13; Table

2; Supporting Fig. 6) may represent ontogenetic shifts in the relationship of various facial

structures or shifts in the relative timing of normal developmental events occurring within

different parts of the face. Subtle differences in embryonic palatal segment proportions were

identified between the strains when considering the entire embryonic series at once, but these

differences did not match the inter-strain differences observed in adult specimens. Additionally,

some proportional palatal contribution differences for NOD and C57 changed between E15 and

P1 as well as between P1 and adult samples. A larger sample of later embryonic ages (i.e.,

between E15.5 and P0) would allow us to determine when the initial NOD pattern of short

primary and long posterior secondary palate changes to the P1 pattern of long primary and short

posterior secondary palate.

Although changes within the RGZ growth dynamics of anterior secondary palate growth

may be critical for major species-specific differences in prognathism, our results did not support

the idea that this mechanism is primarily responsible for the adult intraspecies differences in

upper jaw morphology between our mouse strains. This suggests that later prenatal (after E15)

and postnatal growth processes play a major role in determining adult upper jaw bone

proportions between inbred mouse strains. Broad embryonic growth patterns shared across our

strains indicated that elongation of the anterior secondary palate contributes most to palatal A-P

growth between E11 and E15, but that growth of the primary palate derived premaxilla

contributed most to postnatal elongation of the upper jaw.

Concluding Statement

Our multifaceted illustration of normal midfacial growth dynamics confirmed a one-to-

one relationship between palatal segments and upper jaw bones during the earliest stages of

palatal growth, suggesting that the first formed ruga represents a consistent morphological

boundary between anterior and posterior secondary palate bone precursors, thus approximating

the position of the future maxillary-palatine suture. In addition to driving rugae formation, our

results suggested that interactions at the RGZ simultaneously coordinate elongation of the

maxillary bone primordium within the anterior secondary palate, which more than doubles in

length prior to palatal shelf fusion. Although RGZ-driven A-P growth of the anterior secondary

palate likely contributes to evolutionary changes in facial upper jaw morphology, this process

was not responsible for the small but significant strain-specific differences in adult upper jaw

bone proportions measured across three inbred mouse strains. However, measured differences in

early palatal segment elongation provide a useful foundation for understanding the impact of

background genetic effects on facial morphogenesis. Our multifaceted analysis of ontogenetic

trends in palatal and facial elongation provides a novel contextual framework and developmental

perspective within which to evaluate the impact of both non-pathogenic and pathogenic genetic

differences on midfacial growth and differentiation.

Acknowledgements

We are indebted to Benedikt Hallgrímsson for the use of his laboratory for uCT and microscopy

image collection. This work was supported by start-up funds to Christopher J. Percival from

Stony Brook University; NIH/NIDCR R01 grant DE028753 and UCSF Chancellor Funds from

Recruitment Package to Licia Selleri who supported the personnel and all the research completed

in her lab for this study; and NIH/NIDCR F32 Fellowship DE026950 to Ian Welsh. The authors

of this manuscript have no conflicts of interest to declare.

Data Availability Statement

µCT images, limb bud microscopy photographs, landmark coordinate data, and associated

morphometric analysis R scripts are available publicly as a dataset within Dryad

(https://doi.org/10.5061/dryad.ghx3ffbvb; Welsh et al., 2024).

Author Contributions

ICW contributed to study design, acquisition and interpretation of histological images, and

drafting of the manuscript. MEF contributed to acquisition, analysis, and interpretation of

morphometric data, and critical revision of the manuscript. DL contributed to the design of

morphometric methods, acquisition of morphometric data, measurement and analysis of limb

bud images, and critical revision of the manuscript. IM contributed to acquisition of µCT images,

acquisition, measurement and analysis of limb bud images, and critical revision of the

manuscript. KH contributed to mouse breeding, specimen collection, histological data collection,

and critical revision of the manuscript. CJP contributed to study design, acquisition of µCT

images, acquisition, analysis, and interpretation of morphometric data, and drafting of the

manuscript.

<u>References</u>

724   Adams, D.C., Collyer, M.L., Kaliontzopoulou, A., 2020. geomorph: Software for geometric

725   morphometric analyses. R package version 3.3.1.https://cran.r-

726   project.org/package=geomorph

727   Boughner, J.C., Wat, S., Diewert, V.M., Young, N.M., Browder, L.W., Hallgrímsson, B., 2008.

728   Short-faced mice and developmental interactions between the brain and the face. Journal

729   of Anatomy 213, 646–662.

730   Bush, J.O., Jiang, R., 2012. Palatogenesis: morphogenetic and molecular mechanisms of

731   secondary palate development. Development 139, 231–243.

732   Cignoni, P., Callieri, M., Corsini, M., Dellepiane, M., Ganovelli, F., Ranzuglia, G., 2008.

733   Meshlab: an open-source mesh processing tool., in: Eurographics Italian Chapter

734   Conference. pp. 129–136.

735   https://doi.org/10.2312/LocalChapterEvents/ItalChap/ItalianChapConf2008/129-136

736   Collyer, M.L., Adams, D.C., 2018. RRPP: An R package for fitting linear models to high-

737   dimensional data using residual randomization. Methods in Ecology and Evolution.

738   Economou, A.D., Ohazama, A., Porntaveetus, T., Sharpe, P.T., Kondo, S., Basson, M.A., et al.,

739   2012. Periodic stripe formation by a Turing mechanism operating at growth zones in the

740   mammalian palate. Nat Genet 44, 348–351. https://doi.org/10.1038/ng.1090

741   Enlow, D.H., Bang, S., 1965. Growth and remodeling of the human maxilla. Am J Orthod 51,

742   446–464. https://doi.org/10.1016/0002-9416(65)90242-3

Fedorov, A., Beichel, R., Kalpathy-Cramer, J., Finet, J., Fillion-Robin, J.-C., Pujol, S., et. al.,

2012. 3D Slicer as an image computing platform for the Quantitative Imaging Network.

Magnetic Resonance Imaging 30, 1323–1341. https://doi.org/10.1016/j.mri.2012.05.001

Fish, J.L., 2019. Evolvability of the vertebrate craniofacial skeleton. Semin Cell Dev Biol 91,

13–22. https://doi.org/10.1016/j.semcdb.2017.12.004

Geyer, S.H., Mohun, T.J., Weninger, W.J., 2009. Visualizing vertebrate embryos with episcopic

3D imaging techniques. ScientificWorldJournal 9, 1423–1437.

https://doi.org/10.1100/tsw.2009.154

Hall, J., Jheon, A.H., Ealba, E.L., Eames, B.F., Butcher, K.D., Mak, S.-S., et. al., 2014.

Evolution of a developmental mechanism: Species-specific regulation of the cell cycle

and the timing of events during craniofacial osteogenesis. Dev Biol 385, 380–395.

https://doi.org/10.1016/j.ydbio.2013.11.011

Hallgrímsson, B., Jamniczky, H., Young, N.M., Rolian, C., Parsons, T.E., Boughner, J.C., et. al.,

2009. Deciphering the Palimpsest: Studying the Relationship between Morphological

Integration and Phenotypic Covariation. Evolutionary Biology 36, 355–376.

Hammond, N.L., Dixon, M.J., 2022. Revisiting the embryogenesis of lip and palate

development. Oral Dis 28, 1306–1326. https://doi.org/10.1111/odi.14174

Hauser, G., Daponte, A., Roberts, M., 1989. Palatal rugae. Journal of Anatomy 165, 237–249.

Hilliard, S.A., Yu, L., Gu, S., Zhang, Z., Chen, Y.P., 2005. Regional regulation of palatal growth

and patterning along the anterior–posterior axis in mice. Journal of Anatomy 207, 655–

667. https://doi.org/10.1111/j.1469-7580.2005.00474.x

Jayasankar, P., Banker, A., Bhattacharya, A., Gandhi, R., Patel, N., Parikh, S., 2016. Quantitative

and qualitative analysis of palatal rugae patterns in Gujarati population: A retrospective,

cross-sectional study. J Forensic Dent Sci 8, 126. https://doi.org/10.4103/0975-

1475.195110

Kaucka, M., Petersen, J., Tesarova, M., Szarowska, B., Kastriti, M.E., Xie, M., et. al., 2018.

Signals from the brain and olfactory epithelium control shaping of the mammalian nasal

capsule cartilage. Elife 7, e34465. https://doi.org/10.7554/eLife.34465

Kawasaki, M., Kawasaki, K., Meguro, F., Yamada, A., Ishikawa, R., Porntaveetus, T., et. al.,

2018. Lrp4/Wise regulates palatal rugae development through Turing-type reaction-

diffusion mechanisms. PLoS One 13, e0204126.

https://doi.org/10.1371/journal.pone.0204126

Kouskoura, T., Kozlova, A., Alexiou, M., Blumer, S., Zouvelou, V., Katsaros, C., Chiquet, M.,

Mitsiadis, T.A., Graf, D., 2013. The Etiology of Cleft Palate Formation in BMP7-

Deficient Mice. PLos One 8, e59463. https://doi.org/10.1371/journal.pone.0059463

Kurihara, S., Enlow, D.H., Rangel, R.D., 1980. Remodeling reversals in anterior parts of the

human mandible and maxilla. Angle Orthod 50, 98–106. https://doi.org/10.1043/0003-

3219(1980)050<0098:RRIAPO>2.0.CO;2

Li, Q., Ding, J., 2007. Gene expression analysis reveals that formation of the mouse anterior

secondary palate involves recruitment of cells from the posterior side. Int. J. Dev. Biol.

51, 167–172.

Luo, W., Yi, Y., Jing, D., Zhang, S., Men, Y., Ge, W.-P., et. al., 2019. Investigation of Postnatal

Craniofacial Bone Development with Tissue Clearing-Based Three-Dimensional

Imaging. Stem Cells and Development 28, 1310–1321.

https://doi.org/10.1089/scd.2019.0104

Maga, A.M., 2016. Postnatal Development of the Craniofacial Skeleton in Male C57BL/6J Mice.

J Am Assoc Lab Anim Sci 55, 131–136.

Martinez-Maza, C., Rosas, A., Nieto-Díaz, M., 2013. Postnatal changes in the growth dynamics

of the human face revealed from bone modelling patterns. J Anat 223, 228–241.

https://doi.org/10.1111/joa.12075

Morris, Z.S., Abzhanov, A., 2021. Heading for higher ground: Developmental origins and

evolutionary diversification of the amniote face. Curr Top Dev Biol 141, 241–277.

https://doi.org/10.1016/bs.ctdb.2020.12.003

Murray, S.A., Morgan, J.L., Kane, C., Sharma, Y., Heffner, C.S., Lake, J., et. al., 2010. Mouse

Gestation Length Is Genetically Determined. PLoS ONE 5, e12418.

https://doi.org/10.1371/journal.pone.0012418

Musy, M., Flaherty, K., Raspopovic, J., Robert-Moreno, A., Richtsmeier, J.T., Sharpe, J., 2018.

A quantitative method for staging mouse embryos based on limb morphometry.

Development 145, 1–7.

Okano, J., Suzuki, S., Shiota, K., 2006. Regional heterogeneity in the developing palate:

morphological and molecular evidence for normal and abnormal palatogenesis.

Congenital Anomalies 46, 49–54. https://doi.org/10.1111/j.1741-4520.2006.00103.x

Palmer, A.R., Strobeck, C., 1986. Fluctuating Asymmetry: Measurement, Analysis, Patterns.

Annu. Rev. Ecol. Syst. 17, 391–421.

https://doi.org/10.1146/annurev.es.17.110186.002135

Pantalacci, S., Prochazka, J., Martin, A., Rothova, M., Lambert, A., Bernard, L., et. al., 2008.

Patterning of palatal rugae through sequential addition reveals an anterior/posterior

810   boundary in palatal development. BMC Developmental Biology 8, 116.

811   https://doi.org/10.1186/1471-213X-8-116

812 Pauws, E., Hoshino, A., Bentley, L., Prajapati, S., Keller, C., Hammond, P., et. al., 2009.

813   Tbx22null mice have a submucous cleft palate due to reduced palatal bone formation and

814   also display ankyloglossia and choanal atresia phenotypes. Hum Mol Genet 18, 4171–

815   4179. https://doi.org/10.1093/hmg/ddp368

816 Percival, C.J., Green, R., Marcucio, R., Hallgrímsson, B., 2014. Surface landmark quantification

817   of embryonic mouse craniofacial morphogenesis. BMC Developmental Biology 14, 1-12.

818   https://doi.org/10.1186/1471-213X-14-31

819 Percival, C.J., Liberton, D.K., Pardo-Manuel de Villena, F., Spritz, R., Marcucio, R.,

820   Hallgrímsson, B., 2016. Genetics of murine craniofacial morphology: diallel analysis of

821   the eight founders of the Collaborative Cross. Journal of Anatomy 228, 96–112.

822   https://doi.org/10.1111/joa.12382

823 Peterkova, R., Klepacek, I., Peterka, M., 1987. Prenatal development of rugae palatinae in mice:

824   scanning electron microscopic and histologic studies. Journal of craniofacial genetics and

825   developmental biology 7, 169–89.

826 R Core Team, 2021. R: A language and environment for statistical computing. R Foundation for

827   Statistical Computing, Vienna, Austria [WWW Document]. URL https://www.R-

828   project.org/

829 Sarnat, B.G., 1997. Some methods of assessing postnatal craniofaciodental growth: a

830   retrospective of personal research. Cleft Palate Craniofac J 34, 159–172.

831   https://doi.org/10.1597/1545-1569_1997_034_0159_smoapc_2.3.co_2

Schneider, R.A., 2015. Regulation of Jaw Length During Development, Disease, and Evolution.

Curr Top Dev Biol 115, 271–298. https://doi.org/10.1016/bs.ctdb.2015.08.002

Selleri, L., Rijli, F.M., 2023. Shaping faces: genetic and epigenetic control of craniofacial

morphogenesis. Nature Reviews Genetics 24, 610–626. https://doi.org/10.1038/s41576-

023-00594-w

Tamarin, A., 1982. The formation of the primitive choanae and the junction of the primary and

secondary palates in the mouse. American Journal of Anatomy 165, 319–337.

https://doi.org/10.1002/aja.1001650308

Tonge, C.H., McCance, R.A., 1965. Severe undernutrition in growing and adult animals: 15. The

mouth, jaws and teeth of pigs. British Journal of Nutrition 19, 361–372.

https://doi.org/10.1079/BJN19650034

Vora, S.R., Camci, E.D., Cox, T.C., 2015. Postnatal Ontogeny of the Cranial Base and

Craniofacial Skeleton in Male C57BL/6J Mice: A Reference Standard for Quantitative

Analysis. Front Physiol 6, 417. https://doi.org/10.3389/fphys.2015.00417

Welsh, I.C., Feiler, M.E., Lipman, D., Mormile, I., Hansen, K., Percival, C.J., 2024. Dryad

dataset: Palatal segment contributions to midfacial anterior-posterior growth.

https://doi.org/10.5061/dryad.ghx3ffbvb

Welsh, I.C., Hagge-Greenberg, A., O'Brien, T.P., 2007. A dosage-dependent role for Spry2 in

growth and patterning during palate development. Mechanisms of Development 124,

746–761. https://doi.org/10.1016/j.mod.2007.06.007

Welsh, I.C., Hart, J., Brown, J.M., Hansen, K., Rocha Marques, M., Aho, R.J., et. al., 2018. Pbx

loss in cranial neural crest, unlike in epithelium, results in cleft palate only and a broader

midface. Journal of Anatomy 233, 222–242. https://doi.org/10.1111/joa.12821

Welsh, I.C., O'Brien, T.P., 2009. Signaling integration in the rugae growth zone directs

sequential SHH signaling center formation during the rostral outgrowth of the palate.

Developmental Biology 336, 53–67. https://doi.org/10.1016/j.ydbio.2009.09.028

Wilson, R., McGuire, C., Mohun, T., DMDD Project, 2016. Deciphering the mechanisms of

developmental disorders: phenotype analysis of embryos from mutant mouse lines.

Nucleic Acids Res 44, D855-861. https://doi.org/10.1093/nar/gkv1138

Young, N.M., Hu, D., Lainoff, A.J., Smith, F.J., Diaz, R., Tucker, A.S., et. al., 2014. Embryonic

bauplans and the developmental origins of facial diversity and constraint. Development

141, 1059–1063. https://doi.org/10.1242/dev.099994

Yu, K., Ornitz, D.M., 2011. Histomorphological study of palatal shelf elevation during murine

secondary palate formation. Developmental Dynamics 240, 1737–1744.

https://doi.org/10.1002/dvdy.22670

Yu, L., Gu, S., Alappat, S., Song, Y., Yan, M., Zhang, X., et. al., 2005. Shox2-deficient mice

exhibit a rare type of incomplete clefting of the secondary palate. Development 132,

4397–4406. https://doi.org/10.1242/dev.02013

Yuan, Y., Loh, Y.-H.E., Han, X., Feng, J., Ho, T.-V., He, J., et. al., 2020. Spatiotemporal cellular

movement and fate decisions during first pharyngeal arch morphogenesis. Sci Adv 6,

eabb0119. https://doi.org/10.1126/sciadv.abb0119

Zelditch, M., Swiderski, D., Sheets, D.H., 2012. Geometric morphometrics for biologists: a

primer, 2nd ed. Elsevier Academic Press, San Diego.

Zhao, H., Feng, J., Ho, T.-V., Grimes, W., Urata, M., Chai, Y., 2015. The suture provides a niche

for mesenchymal stem cells of craniofacial bones. Nat Cell Biol 17, 386–396.

https://doi.org/10.1038/ncb3139

<u>Supporting Information</u>

Supporting Tables and Figures are found as a Supporting .pdf file in association with this

manuscript on the publisher's website.

Tables

Table 1 –Sample size of μCT imaged and morphometrically analyzed embryo specimens by strain and

forelimb based developmental age estimation, as estimated using eMOSS (Musy et al., 2018).

Limb-Based Developmental Age	C57	NOD	PWK
E11	8	12	4
E11.5	2	0	0
E12	7	3	6
E12.5	4	0	2
E13	9	3	6
E13.5	0	0	2
E14	26	6	6
E14.5	1	0	0
E15	9	4	0

889   Table 2 – Average proportional palatal segment lengths for postnatal specimens, presented as

890   percentages of total palatal length.

	C57			NOD			PWK
	P1 epithelial	P1 bone	Adult bone	P1 epithelial	P1 bone	Adult bone	Adult bone
Primary Palate / Premaxilla	18.4%	23.3%	32.4%	19.7%	24.4%	31.7%	30.7%
Ant. Sec. Palate / Maxilla	45.7%	47.5%	39.9%	49.5%	47.4%	41.8%	42.8%
Post. Sec. Palate / Palatine & Pterygoid	35.9%	29.2%	27.7%	30.8%	28.3%	26.5%	26.5%