Structure-guided mutagenesis of OSCAs reveals differential activation to mechanical stimuli

Sebastian Jojoa-Cruz[1], Adrienne E. Dubin[2], Wen-Hsin Lee[1], Andrew Ward[1]†

Affiliations
[1] Department of Integrative Structural and Computational Biology, Scripps Research, La Jolla,

California 92037, USA.

[2] Department of Neuroscience, Scripps Research, La Jolla, California 92037, USA.

†Corresponding author
Correspondence to: A.B.W.: andrew@scripps.edu

Abstract (150-200 words)

The dimeric two-pore OSCA/TMEM63 family has recently been identified as mechanically

activated ion channels. Previously, based on the unique features of the structure of OSCA1.2, we

postulated the potential involvement of several structural elements in sensing membrane

tension[1]. Interestingly, while OSCA1, 2, and 3 clades are activated by membrane stretch in cell-

attached patches (i.e., they are stretch-activated channels), they differ in their ability to

transduce membrane deformation induced by a blunt probe (poking). In an effort to understand

the domains contributing to mechanical signal transduction, we used cryo-electron microscopy

to solve the structure of *Arabidopsis thaliana* (At) OSCA3.1, which, unlike AtOSCA1.2, only

produced stretch- but not poke-activated currents in our initial characterization[2]. Mutagenesis

and electrophysiological assessment of conserved and divergent putative mechanosensitive

features of OSCA1.2 reveal a selective disruption of the macroscopic currents elicited by poking

without considerable effects on stretch-activated currents (SAC). Our results support the

involvement of the amphipathic helix and lipid-interacting residues in the membrane

fenestration in the response to poking. Our findings position these two structural elements as

potential sources of functional diversity within the family.

Introduction

Mechanotransduction allows organisms to gather information about their external surroundings as well as internal processes. Mechanically-activated (MA) ion channels are capable of sensing mechanical stimuli and transducing this information to the cell as electrochemical signals[3]. MA ion channels are diverse and capable of responding to different stimuli such as osmotic stress, membrane deformation, touch, and hearing[3,4].

In our previous work, we identified OSCA/TMEM63 as a conserved family of MA ion channels spanning several eukaryotic clades and having considerable diversity in the plant kingdom, where species commonly have more than one paralog. For example, *Arabidopsis thaliana* has 15 members[2] and *Oryza sativa* L. *japonica* (rice), 11[5]. We demonstrated these channels can respond to membrane stretch even more robustly than hypertonicity, originally thought to be the relevant stimulus, with some members (OSCA1.1 and OSCA1.2) also able to respond to poke[2]. We used cryo-electron microscopy (cryo-EM) to solve the structure of OSCA1.2 in a closed state in both detergent (LMNG) and nanodiscs at high-resolution, identifying a new architecture for MA ion channels that is structurally homologous to TMEM16 proteins[1]. Results from other groups are consistent with ours[6-8]. Similar to the TMEM16 family, OSCAs are dimeric and each subunit has a pore lined by transmembrane helices (TMs) 3 to 7, instead of a single pore along the symmetry axes[1,6-8]. A hydrophobic gate on the extracellular half of the pore blocks ion flow. Additionally, molecular dynamics (MD) simulations support our placement of the pore and position lipids along an opening of the pore pathway in the transmembrane domain (TMD)[1].

Based on the unique structural characteristics of OSCA1.2, we proposed that two features might be involved in sensing membrane deformation: the amphipathic helix (AH) located in the first intracellular loop, and the Beam-Like Domain (BLD), composed of the two cytoplasmic membrane-parallel helices and a hydrophobic hook that connects them and re-inserts itself into the membrane[1]. Due to the close association of these two elements to the membrane, we reasoned that they could serve as sensors of mechanical stimuli and transmit the signal to the pore through interactions with the TMD. Additionally, a fenestration allows the membrane

access to the pore, and MD simulations placed lipids in close association with four positively

charged residues in this area suggesting a potential role in gating[1]. We sought to determine

whether these regions contributed to mechanotransduction.

Here, we used cryo-EM to determine a high-resolution structure of AtOSCA3.1, the first reported

OSCA gene[9], in nanodiscs. In our initial characterization, OSCA3.1 responded to stretch but not

poke[2]. Comparison of the OSCA3.1 structure reported here to previous structures of OSCA1.1[6]

and OSCA1.2[1] revealed subtle differences. Structure-function analyses revealed the involvement

of the AH and the lipid-interacting residues in the pore pathway of OSCA1.2 in the ability of the

channel to respond to membrane deformation by poking. Our results suggest that these features

are not only important for mechanotransduction or activation of OSCA1.2 by poking in OSCAs but

may serve to tune the response of OSCA/TMEM63 members to diverse stimuli.

Results

Overall architecture of OSCA3.1 is similar to OSCA1.1 and OSCA1.2

We recombinantly tagged OSCA3.1 with EGFP at its C-terminus, expressed the construct in

HEK293F cells, and purified it using Lauryl Maltose Neopentyl Glycol (LMNG) supplemented with

cholesteryl hemisuccinate (CHS). The EGFP was removed by PreScission Protease cleavage,

leaving a stretch of ten residues on the C-terminus. Purified protein was then reconstituted into

nanodiscs and subjected to cryo-EM analysis, resulting in a 2.6 Å resolution symmetric

reconstruction (Fig. 1a, Supplementary Fig. 1) sufficient to build a model of the majority of the

protein (537 out of 724 residues) (Fig. 1b-c, Supplementary Fig. 2, Supplementary Table 1). Due

to increased flexibility of the BLD relative to previous OSCA structures, we were unable to assign

residue identity, and instead modeled it as two poly-Ala helices.

Overall, there is good agreement between OSCA3.1, OSCA1.1 and OSCA1.2 protomers[1,6] ($C\alpha$

RMSD OSCA3.1-OSCA1.1: 1.243Å, OSCA3.1-OSCA1.2: 1.152Å, Fig. 1c). The pore profile of these

OSCAs is maintained and the small radii towards the extracellular side suggests these channels

are in a closed/non-conductive state, as suggested by previous MD simulations[1,6] (Fig. 1d).

Likewise, the two π-helical turns in TM5 and TM6a seen in previous structures near the neck of

the pore are present at similar positions in OSCA3.1[1,8]. We identified multiple lipid-like densities

in one of our postprocessed maps (Fig. 1e, Supplementary Fig. 3). Notably, one of these densities

is located at the pore fenestration, at a position similar to that observed in previous MD

simulations of OSCA1.2[1], supporting the proposed occupation of the pore pathway by lipids.

Recent cryo-EM structures of AtOSCA1.2[10], and AtOSCA1.1 and human TMEM63A[11] also place a

lipid at a similar position.

Previously, the structure of OSCA3.1 was solved in a digitonin micelle[6], and more recently, in

LMNG[11]. The similarity of our nanodisc sample to these detergent structures suggests that the

use of detergent did not affect the conformation or oligomerization of OSCA3.1, similar to

previous results with OSCA1.2[1,7] (Supplementary Fig. 4a-b). However, the inter-subunit cleft in

the nanodisc structure is wider due to outward movements of innermost TMs ranging from ~4-

8Å (Supplementary Fig. 4c) relative to the digitonin sample, which was recently denoted as an

"extended" state[11]. Thus, our nanodisc structure represents a further "extended" state than the

one found both in digitonin and LMNG micelles[11] (Supplementary Fig. 4d). Whether these

differences represent biologically meaningful states, an effect of the lipid environment

(detergent vs lipids), or inherent flexibility of the channel remains to be determined. Additionally,

our structure enabled a higher degree of certainty in the residue assignment relative to the initial

digitonin structure, particularly for TM0 where sidechains were not previously modelled[6].

Consequently, we have shifted the registry of TM0 by two residues relative to the detergent

structure, in agreement with the LMNG structures[11].

Mutation of key residues in the amphipathic helix abrogates poke but not stretch responses in

OSCA1.2

Structures of AtOSCA1.1, rice and At OSCA1.2, and OSCA3.1 (Fig. 2a, Supplementary Fig. 5), as

well as sequence alignment, suggest that the AH between TM0 and TM1 of OSCAs is a conserved

feature of these channels. AHs can serve as both membrane anchors and sensors of membrane

deformation[12], and they are present in many MA ion channels[1,13-15]. There is evidence for their

involvement in mechanical force transduction; for example, in the bacterial mechanosensitive

channel of large conductance MscL, an AH in the cytoplasmic side of the bilayer links membrane

tension to protein conformation (Supplementary Fig. 6a,b)[14]. Interestingly, structures of

members of the TMEM16 family, structural homologs of OSCAs, have a similarly placed helix in

this region; however, it is located in the cytosol and does not present amphipathic properties

(Supplementary Fig. 6c-e)[16-18]. Involvement of TMEM16A in the detection of mechanical forces

in myocytes and bile ducts was dependent on calcium influx (potentially caused by an upstream

MA cation channel) and not a direct response to the mechanical stimulus[19,20]. To our knowledge,

characterization of TMEM16 proteins as MA ion channels has not been reported. Based on these

observations, we mutated the membrane-facing residues at each end of the AH of OSCA1.2 in an

effort to decrease its propensity to interact with the membrane. Single (W75K or L80E) and

double (W75K/L80E) mutants in OSCA1.2 strongly abrogated the response to indentation of the

membrane with a blunt glass probe (poke) (WT: 47 of 51 cells produced MA currents; W75K: 3 of

24; L80E: 2 of 16; W75K/L80E: 2 of 24; Fig. 2b, top panel). Moreover, in OSCA1.2$_{W75K/L80E}$, a larger

displacement of the membrane by the probe relative to the WT appeared to be needed to

observe currents ("apparent threshold"; WT: 5.3 ± 0.5 μm (N = 46; mean ± S.E.M.), W75K/L80E:

15.3 ± 2.3 μm (N = 2), with the none-responsive W75K/L80E cells rupturing at a 13.0 ± 0.5 μm (N

= 22); Fig. 2b, bottom panel). The insensitivity to the poking stimulus was not due to poor

membrane trafficking of these constructs (see below). The loss of sensitivity of cells expressing

these mutant constructs to poking suggests that the AH of OSCA1.2 plays an important role in

this mechanically-activated response.

We then tested whether these mutations in the AH affected the response of OSCA1.2 to stretch.

Contrary to what was observed during poking, all cell-attached patches revealed stretch-

activated currents (WT: 25 of 26 cells; W75K: 3 of 3; L80E: 5 of 5; W75K/L80E: 11 of 11; Fig. 2c,

top panel; electrode resistances were similar in all cases (see Figure legend)). Despite differences

in the amplitude of macroscopic currents, which could be explained by variation in expression

levels, all mutant channels were activated at similar thresholds (WT: -48 ± 5 mmHg (N = 25);

W75K: -56 ± 11 mmHg (N = 3), L80E: -41 ± 6 mmHg (N = 5), W75K/L80E: -53 ± 8 mmHg (N = 11))

(Fig. 2c, bottom panel). In all cases, there was variability in the rate of inactivation from

moderately fast to extremely slow (data not shown). Overall, the observed stretch response of

OSCA1.2 was apparently not affected by the mutations we introduced.

Subsequently, we mutated the proline in the OSCA1.2 AH to the corresponding arginine in

OSCA3.1 to determine whether this difference impacts the ability of OSCA3.1 to robustly

transduce the poke stimulus; this change (OSCA1.2$_{P77R}$) had no effect on the poke response

(Supplementary Fig. 7).

It is worth noting that in our initial characterization, OSCA3.1-expressing cells produced SAC while

poke did not induce observable currents[2]. However, in the present experiments we were able to

record poke-activated currents mediated by OSCA3.1 on a few occasions (OSCA1.2$_{WT}$: 6 of 6 cells;

OSCA3.1$_{WT}$: 3 of 10) (Supplementary Fig. 8a). The apparent threshold for OSCA3.1 activation by

poke-induced membrane displacement tended to be higher than for OSCA1.2 but the difference

was not statistically significant (OSCA1.2$_{WT}$: 10.4 ± 1.2 µm (N = 6; mean ± S.E.M.); OSCA3.1$_{WT}$:

13.8 ± 0.9 µm (N = 3)) (Supplementary Fig. 8b), perhaps due to the low numbers of responsive

cells analyzed. These data suggest that the poking stimulus is capable of activating OSCA3.1 but

the stimulus intensity required is close to the rupture point of the HEK293T-Piezo1-knockout

(HEK-P1KO) cells used in our assay.

Replacing the OSCA3.1 Beam-Like Domain (BLD) in OSCA1.2 had no effect on stretch-induced

currents but decreased the apparent sensitivity of the cells to poke

One of the unique features of OSCAs is the presence of the BLD, a hydrophobic hook that, by

inserting itself into the membrane, may serve as an anchor that could potentially be displaced

under tension[1,7,8]. Hydrogen/deuterium exchange mass spectrometry (HDXMS) of rice OSCA1.2

indicated that the helix of the BLD closest to the membrane undergoes less deuterium exchange

and potentially remains associated to a neighboring surface[8], hinting at an important interaction

with TM6b. In OSCA3.1, the BLD is poorly resolved, likely due to flexibility (Fig. 1c). To test

whether the increased flexibility in this region affects MA responses, we replaced the BLD of

OSCA1.2 with the BLD of OSCA3.1 (Fig. 3a). HEK-P1KO cells expressing the chimera OSCA1.2$_{OSCA3.1-}$

$_{BLD}$ produced similar MA currents in both poke and stretch assays (Fig. 3b,c) with the exception

that a slightly stronger displacement stimulus was required to activate the mutant channels (Fig.

3b, bottom panel). These data suggest that the BLD is not responsible on its own for the ability

of the channels to be activated by poking, however, it appears to play a role. As expected for

similar SAC from cells expressing OSCA1.2 and OSCA3.1, there was no obvious change in SAC

when the OSCA1.2 BLD was replaced with the OSCA3.1 BLD. It should be noted that the BLD

tolerates a certain degree of mutation as sequence alignment in this region shows higher

variability than other secondary structures of OSCAs (Supplementary Fig. 5). Thus, we cannot rule

out that in the OSCA1.2 background, the BLD of either channel is capable of performing a similar

role.

Substitution of potential lipid-interacting lysine residues for isoleucine abrogates the poke

response in OSCA1.2

Lipids play an important role in the stability and function of ion channels[21]. In TRAAK, a eukaryotic

mechanosensitive channel, occlusion of the pore by a lipid acyl chain has been proposed as a

gating mechanism[13,22]. In the volume-regulated channel SWELL1 lipids block the pore in the

closed state[23]. In the bacterial mechanosensitive channel MscS, the occupation of lipid pockets

may determine channel conformation[24,25], and in FLYC1, a Venus flytrap homolog of MscS,

conformational changes may allow lipids to access the pore and occlude ion conduction[26].

Moreover, exposure of inside-out patches from OSCA1.1-expressing HEK cells to lyso-

phosphatyidylcholine (LPC) enhanced channel response to stretch[6].

Previously, we identified four lysine residues in OSCA1.2 located in TM4 and TM6b at the pore

fenestration. These residues interacted with lipid phosphate head groups throughout the

duration of our MD simulations[1]. The OSCA3.1 structure revealed that two of these lysine

residues (K435 and K536 of OSCA1.2) have been replaced by isoleucine (I423 and I525,

respectively), which are expected to reduce the interaction with lipid phosphate head groups

(Fig. 4a). To test for the contribution of these lipid-interacting residues to channel

mechanosensitivity, we measured poke- and stretch-induced currents in the double mutant

OSCA1.2$_{K435I/K536I}$ (Fig. 4b,c). The double mutant responded poorly to poking, with only 1 out of

20 cells producing currents upon mechanical stimulation (K435I/K536I: -365 pA (N = 1)) (Fig. 4b,

comparison made to WT data acquired on the same days). This finding was not due to poor

expression of the double mutant in HEK-P1KO cells; the SAC I$_{max}$ observed in cell-attached patches

was similar to WT (Fig. 4c, top panel). Furthermore, the mutant could be activated by stretch at

similar thresholds to WT (WT: -43 ± 10 mmHg (N = 6); K435I/K536I: -60 ± 10 mmHg (N = 7)) (Fig.

4c, bottom panel), and similar stretch-response dependence (i.e., similar pressures required to

elicit half-maximal currents (P$_{50}$) and slope) (Fig. 4d). Although macrocurrent amplitudes in SAC

recordings tended to be lower than WT tested on the same day, the ~2-fold difference unlikely

accounts for the ~20-fold fewer responsive cells (Fig. 4b,c). Taken together, these results suggest

that loss of positively charged sidechains in the fenestration selectively impairs poke-induced

responses of OSCA1.2 under physiological conditions, but not its responsiveness to the stretch

stimulus.

Discussion

Here we performed structure-function studies to determine the molecular underpinnings of

mechanical activation of OSCA ion channels. We solved the structure of OSCA3.1 at a resolution

of 2.6 Å in a non-conductive state and revealed a marked structural conservation to previous

OSCA orthologs. The final map presented several lipid-like densities surrounding the channel,

including one density close to the pore and several densities populating the inter-subunit cleft.

This ion channel was reported to be activated by stretch in cell-attached patches but insensitive

to a poke stimulus using a blunt glass probe[2]. Comparisons between the structures of OSCA1.2,

which is robustly activated by both poke and stretch, to OSCA3.1, which is only robustly

activated by stretch, suggested 3 domains that might contribute to gating during a poke

stimulus. First, a conserved short amphipathic helix (AH) which sits at the membrane interface

reminiscent of a similar feature in other MA ion channels like MscL[14], Piezo1[15], MscS[27], and

TRAAK[28]. Second, side chains in TM4 and TM6b in the vicinity of the membrane fenestration

may have differential interactions with lipids as indicated by MD simulations of OSCA1.2. Third,

the beam-like domain (BLD) that contains 2 helices and linker that hooks around the

cytoplasmic surface into the membrane in OSCA1.2 but is less well-resolved in the OSCA3.1

map presumably because it is not intimately associated with the channel/membrane.

Mutagenesis of the AH and residues at the fenestration in OSCA1.2 largely abrogated its

response to poking without affecting its activation by stretch; replacing the BLD with that from

OSCA3.1 only slightly impaired the response to poking, having no effect on stretch-induced

activation.

Residues in the AH were mutated to either confer more of an OSCA3.1 character or tend to

make the domain less likely to interact with the membrane, similar to the TMEM16 family that

are not thought to be directly activated by mechanical stimuli. Swapping the divergent residue

in the AH of OSCA1.2 with its equivalent in OSCA3.1 (OSCA1.2$_{P77R}$), had no negative impact on

the sensitivity to poking. These data suggest that the difference in poke sensitivity between

OSCA1.2 and OSCA3.1 is not due to the AH itself. However, when mutations were incorporated

to presumably destabilize the interaction of the AH domain with the membrane

(OSCA1.2$_{W75K/L80E}$), fewer cells could be activated by poking to similar stimulus strengths while

stretch-activation of channels was not impaired. Although a WT OSCA1.2 AH is critical for

normal poke-induced channel opening, these mutations do not hinder gating by membrane

stretch.

Substituting the BLD of OSCA1.2 for that of OSCA3.1 had little effect on poke- or stretch-

activated responses. Although these results suggest that the BLD may not be involved in

modulating the MA response of OSCA1.2, a recent report suggested evolutionary coupling of 3

sets of residues in the BLD and TM6b that may form electrostatic interactions between these

domains[10]. The evolutionarily favored residue identities at these sites constitute three potential

salt bridges: **R/K**257-**E**557, **E**260-**R**553, and **R/K**261-**E/D**554 (based on OSCA1.2 numbering).

These sites in OSCA1.2$_{OSCA3.1-BLD}$ chimera correspond to N257**K**-**E**557, D260**A**-**K**553, and Y261**R**-

D554 (mutation-interacting residue), perhaps leaving intact 2 out of 3 potential salt bridges and

maintaining its influence (Supplementary Fig. 5). It is worth noting that a similar chimera of

AtOSCA1.2 with the IL2 domain of the human ortholog TMEM63A (OSCA1.2$_{63A\ IL2}$), containing

most of the intracellular domain which includes the BLD as well as the dimerization domain,

presented little inactivation of SAC akin to monomeric TMEM63A$_{WT}$[29]. This behavior was not

due to an inability of OSCA1.2$_{63A\ IL2}$ to dimerize, as the monomeric OSCA1.2$_{5Mu}$, obtained by

disrupting the dimer interface with 5 mutations while maintaining OSCA1.2$_{WT}$ BLD, exhibited

pressure-induced SAC gating similar to OSCA1.2$_{WT}$[29]. This new evidence, taken together with

the decrease in the apparent poking sensitivity of the OSCA1.2$_{OSCA3.1-BLD}$ chimera in this study,

suggests that the BLD plays a role in the force-induced gating of the OSCA family.

The structure of OSCA3.1 exhibits an open fenestration similar to OSCA1.2 and lipids are expected

to populate this area[1], similar to lipid occlusion of the pore in other MA ion channels such as

TRAAK[13] and MscS[24,25]. Our results suggest that TM4 and TM6b interactions with lipids may

contribute to activation during poking. Modification of OSCA1.2 residues to match those of

OSCA3.1 (OSCA1.2$_{K435I/K536I}$) produced an OSCA3.1-like phenotype (MA currents are observed in

response to stretch but rarely to poke). These mutant channels are not completely insensitive to

poking but appear to have a very high threshold to activate, one that is often beyond the point

of cell rupture by the stimulus (similar to that seen with OSCA1.2$_{W75K/L80E}$). Notably, TM4 in

TMEM16 lipid scramblases and TM6 in both TMEM16 scramblases and channels undergo

conformational changes during activation[16,17,30]. The loss of positive charges in OSCA3.1 may

weaken the interactions between residues and lipids, similar to MscS where charge

neutralization of the lipid facing residue R59 by the introduction of a leucine strongly altered

gating, potentially by disturbing a salt bridge with a lipid phosphate[31,32]. A weaker interaction

could decrease the tension generated by the lipids upon mechanical stimulus, hence increasing

the force needed to open the channel, as seen in our poke experiments. A similar argument could

be made for the AH of OSCAs: by disturbing its coupling with the lipid bilayer, a greater stimulus

might be needed to activate the channel. It is also important to note that the membrane of a

plant cell contains a different lipid composition than that of HEK293 cells used in our assays, and

thus these lipids, or association with the plant cell wall, may alter how these channels respond
to physiological stimuli in vivo.

The phenotypes of OSCA1.2$_{W75K/L80E}$ and OSCA1.2$_{K435I/K536I}$ suggest a mechanism recruited by
poking is no longer able to adequately activate OSCA1.2$_{W75K/L80E}$ and OSCA1.2$_{K435I/K536I}$ channels.
Since the factors contributing to channel activation in the poking assay are not understood, we
can only speculate as to mechanisms underlying the loss of sensitivity to poking in these mutants.
Our results suggest that these stimuli are sensed by different features of OSCA channels, akin to
the well-studied ThermoTRP channels[33-35]. Polymodal activation mechanisms have been recently
reported for the MA channel PIEZO2 where a mutagenesis approach revealed differential effects
on the ability to respond to poking and stretch[36]. Nonetheless, the discrepancy could be due to
inherent methodological differences between these two assays, as whole-cell recordings during
poking involve channels in inaccessible membranes (at the cell-substrate interface) and channel
interactions with extracellular and intracellular components[37], while the stretch assay is limited
to recording channels inside the patch.

Overall, the analysis of these mutants demonstrates the involvement of the AH and residues at
the membrane fenestration in the MA response of OSCAs, as we initially proposed[1]. The
participation of residues from TM4 and TM6 hint at further similarities with TMEM16 family, but
to what extent is unknown. Perhaps the most surprising result is the selective disruption of poke-
activated currents of OSCA1.2 without affecting its response to membrane stretch, which
suggests that changes in these features could serve to fine-tune the response to certain
mechanical stimuli and may be a source of functional diversity within this large family. The
structures and characterizations presented here and in our previous work[1,2] will prove
fundamental in assessing the physiological consequences of these differences and further
understanding the role of the OSCA/TMEM63 family.

Materials and methods:

Expression constructs

For structural studies, the OSCA3.1 (UniProt ID: Q9C8G5) coding sequence was synthesized using

optimized codons for expression in human cells and subsequently cloned into the pcDNA3.1

vector used in our previous report[1]. In this plasmid, the C terminus of the protein sequence is

followed by a PreScission Protease cleavage site, EGFP and a FLAG tag.

Protein expression and purification

Samples of OSCA3.1 in nanodisc were obtained following the same method reported previously

for OSCA1.2 in nanodiscs[1], with the exception of the detergent used. Instead of beta-D-

maltopyranoside (DDM), we used Lauryl Maltose Neopentyl Glycol (LMNG) due to increased

yields. Thus, the detergent composition of the solubilization and wash buffers was 1%/0.1% and

0.01%/0.001% LMNG/cholesteryl hemisuccinate (CHS), respectively. Fractions from Size

Exclusion Chromatography (SEC) corresponding to the OSCA3.1 in nanodiscs peak were

concentrated to 3.2mg/mL (corrected using the extinction coefficient of OSCA3.1).

Cryo-EM sample preparation and data collection

3.5 μL of OSCA3.1 at 3.2mg/mL were frozen using the same conditions as OSCA1.2 samples[1].

Grids were imaged on a Titan Krios (ThermoFisher) operating at 300 kV with a K2 Summit direct

electron detector (Gatan), with pixel size of 1.03Å (nominal magnification of 29,000x). A total of

38 frames were collected per movie, adding to a total dose of ~50 electrons per $Å^2$. Automated

image collection was done through Leginon[38] using a defocus range of -0.4 to -1.5 μm. A total of

8,375 movies were collected.

Cryo-EM image processing

During collection, movies were aligned and dose-weighted using MotionCor2[39]. The resulting

micrographs were imported into cryoSPARCv2.5[40] and CTF values were estimated through Gctf[41].

~300 particles were manually picked from ~10 micrographs and subjected to 2D classification to

obtain templates. Template picking was tested on 20 micrographs and particles used to generate

better 2D templates. These templates were later used to pick particles on ~1600 micrographs.

The resulting ~900k particles were subjected to, in order, 2D classification, *ab initio*

reconstruction, 2 rounds of heterogeneous refinement, homogeneous refinement and non-

uniform refinement to obtain an initial 3D reference, which reached high-resolution. Unless

otherwise specified, C2 symmetry was applied to all homogeneous, non-uniform and RELION 3D

refinements.

For the full dataset, 8,375 movies were imported into cryoSPARCv2 and subjected to full-frame

motion correction followed by Gctf estimation. Given the large size of the dataset, and the fact

that initial processing had returned a high-resolution structure, we decided to apply a 2.6 Å CTF

cutoff to reduce the number of micrographs to 3,068 and speed up processing. Templates were

generated based on the initial 3D reference. Template picking, followed by local motion

correction and 2D classification resulted in 1,913,316 selected particles for our initial stack. *Ab*

initio reconstruction produced 3 classes, and heterogeneous refinement was performed on the

two worst classes. The resulting best class was combined with the best class from *ab initio* and

used for two rounds of heterogeneous and homogeneous refinements. The heterogeneous

refinement of the first round was done without symmetry, from that point onwards, C2

symmetry was imposed unless otherwise specified. Only particles belonging to the best class

were selected for further processing (318,249). Non-uniform refinement, followed by 2 rounds

of local refinement (without imposing symmetry), were performed on these particles before

exporting them into RELION-3.1[42,43]. The particle stack underwent the following process in

RELION. 3D refinement, CTF refinement, 3D refinement with global angular searches, 3D

refinement using SIDESPLITTER (SS)[44], and two rounds of CTF and 3D refinements using SS. Unless

specified, all 3D refinements were limited to local angular searches. 3D classification without

alignment (no imposed symmetry) was performed on the particle stack and the best class,

comprising 197,944 particles, was subjected to a final 3D refinement with SS. The map was

sharpened separately with LocalDeblur[45] through Scipion[46-48] and DeepEMhancer using the

'highRes' model[49]. The FSC and local resolution of the map was calculated through RELION.

Model building and refinement

A homology model for OSCA3.1 was obtained through SWISS-MODEL[50], using as template the

structure of OSCA1.2 in nanodiscs (PDB: 6MGV)[1]. Iterative rounds of building in Coot[51,52] and real

space refinement in Phenix[53,54] and Rosetta[55] were used to generate the final model. SMILES

codes for the ligands were used to obtain the appropriate restrains using eLBOW[56]. Model

validation was carried out with MolProbity[57] and EMringer[58] to the LocalDeblur map. Phenix

mtriage[59] was used to calculate the map to model FSC. The final model comprises residues 2-102,

154-231, 307-388, 410-476, and 489-697, for a total of 537 out of 724 residues in OSCA3.1

sequence. A segment of the BLD was modelled as poly-A helices and registry was tentatively

assigned based on sequence and structure alignment to OSCA1.2[1]: residues 234-246 and 280-

303 assigned as IL2H2 and IL2H3, respectively. Pore profiles were predicted by CHAP[60,61].

Structure figures were made with PyMOL[62], UCSF Chimera[63] or UCSF ChimeraX[64]. Amino acid

sequence alignment was obtained from Clustal omega[65] and represented with ESPript3[66].

Sequences used for the alignment were the same as previous publication[1], but the display was

limited to the *Arabidopsis thaliana* OSCAs with solved structure.

Generation of mutants, cell culture and transfections

A vector containing OSCA1.2 pIRES2-mCherry was used for electrophysiology experiments[2]. The

W75K substitution was introduced using Q5 site-directed mutagenesis kit (New England Biolabs,

NEB). Substitutions L80E, K435I, and K536I were introduced using QuikChange multi site-directed

mutagenesis kit (Agilent). For the BLD chimera, OSCA1.2 pIRES2-mCherry vector and the BLD of

OSCA3.1 were separately amplified by PCR using Q5 High-Fidelity 2X Master Mix (NEB), followed

by fragment assembly using Gibson Assembly Master Mix (NEB). Kits were used according to

manufacturer's instructions.

Cell culture and transfection of HEK-P1KO cells for electrophysiology experiments were

conducted as previously reported[1].

Electrophysiology

Patch-clamp experiments in transiently transfected HEK-P1KO cells were performed in standard

whole-cell and cell-attached mode using a Multi-clamp700A amplifier (Axon Instruments) and

followed the procedures described in our previous report[2].

Data availability

Cryo-EM maps of OSCA3.1 in nanodiscs has been deposited to the Electron Microscopy Data Bank

(EMDB) under accession number EMD-41911 with the LocalDeblur map as the primary map. The

corresponding atomic coordinates have been deposited to the Protein Data Bank (PDB) under ID

8U53.

Author contributions:

S.J.C. and W.H.L. cloned expression constructs and mutants. S.J.C. prepared protein samples and

acquired cryo-EM data. S.J.C. processed data and built and refined the atomic structures. A.E.D.

carried out electrophysiology experiments and analyzed data. A.B.W. supervised the project.

S.J.C. and A.E.D. drafted the manuscript, which was edited and finalized with contributions from

all authors.

Acknowledgements:

We thank W. Anderson for management the electron microscopy facility at Scripps Research, J.

Torres for help with data collection, and C. Bowman, L. Dong and J.C. Ducom for assistance with

computation. We acknowledge members of the Ward laboratory. This work was supported by

NIH grant R01 HL143297 and a Ray Thomas Edwards Foundation grant to A.B.W. A.E.D was

supported by grant R35 NS105067.

Molecular graphics and analyses performed with UCSF Chimera and UCSF ChimeraX, developed

by the Resource for Biocomputing, Visualization, and Informatics at the University of California,

San Francisco, with support from National Institutes of Health R01-GM129325 and P41-

GM103311, and the Office of Cyber Infrastructure and Computational Biology, National Institute

of Allergy and Infectious Diseases.

References

1   Jojoa-Cruz, S. *et al.* Cryo-EM structure of the mechanically activated ion channel OSCA1.2.
Elife **7** (2018). https://doi.org:10.7554/eLife.41845
2   Murthy, S. E. *et al.* OSCA/TMEM63 are an Evolutionarily Conserved Family of Mechanically
Activated Ion Channels. *Elife* **7** (2018). https://doi.org:10.7554/eLife.41844
3   Ranade, S. S., Syeda, R. & Patapoutian, A. Mechanically Activated Ion Channels. *Neuron*
87, 1162-1179 (2015). https://doi.org:10.1016/j.neuron.2015.08.032
4   Haswell, E. S., Phillips, R. & Rees, D. C. Mechanosensitive Channels: What Can They Do
and   How   Do   They   Do   It?   *Structure*   **19**,   1356-1369   (2011).
https://doi.org:10.1016/j.str.2011.09.005
5   Li, Y. *et al.* Genome-wide survey and expression analysis of the OSCA gene family in rice.
BMC Plant Biol **15**, 261 (2015). https://doi.org:10.1186/s12870-015-0653-8
6   Zhang, M. *et al.* Structure of the mechanosensitive OSCA channels. *Nat Struct Mol Biol* **25**,
850-858 (2018). https://doi.org:10.1038/s41594-018-0117-6
7   Liu, X., Wang, J. & Sun, L. Structure of the hyperosmolality-gated calcium-permeable
channel OSCA1.2. *Nat Commun* **9**, 5060 (2018). https://doi.org:10.1038/s41467-018-
07564-5
8   Maity, K. *et al.* Cryo-EM structure of OSCA1.2 from Oryza sativa elucidates the mechanical
basis of potential membrane hyperosmolality gating. *Proc Natl Acad Sci U S A* **116**, 14309-
14318 (2019). https://doi.org:10.1073/pnas.1900774116
9   Kiyosue, T., Yamaguchi-Shinozaki, K. & Shinozaki, K. Cloning of cDNAs for genes that are
early-responsive to dehydration stress (ERDs) in Arabidopsis thaliana L.: identification of
three   ERDs   as   HSP   cognate   genes.   *Plant Mol Biol* **25**,   791-798   (1994).
https://doi.org:10.1007/BF00028874
10   Jojoa-Cruz, S., Burendei, B., Lee, W. H. & Ward, A. B. Structure of mechanically activated
ion channel OSCA2.3 reveals mobile elements in the transmembrane domain. *Structure*
(2023). https://doi.org:10.1016/j.str.2023.11.009
11   Zhang, M., Shan, Y., Cox, C. D. & Pei, D. A mechanical-coupling mechanism in
OSCA/TMEM63   channel   mechanosensitivity.   *Nat Commun* **14**,   3943   (2023).
https://doi.org:10.1038/s41467-023-39688-8
12   Drin, G. & Antonny, B. Amphipathic helices and membrane curvature. *FEBS Lett* **584**,
1840-1847 (2010). https://doi.org:10.1016/j.febslet.2009.10.022
13   Brohawn, S. G., Campbell, E. B. & MacKinnon, R. Physical mechanism for gating and
mechanosensitivity of the human TRAAK K+ channel. *Nature* **516**, 126-130 (2014).
https://doi.org:10.1038/nature14013
14   Bavi, N. *et al.* The role of MscL amphipathic N terminus indicates a blueprint for bilayer-
mediated gating of mechanosensitive channels. *Nat Commun* **7**, 11984 (2016).
https://doi.org:10.1038/ncomms11984
15   Saotome, K. *et al.* Structure of the mechanically activated ion channel Piezo1. *Nature* **554**,
481-486 (2018). https://doi.org:10.1038/nature25453
16   Paulino, C., Kalienkova, V., Lam, A. K. M., Neldner, Y. & Dutzler, R. Activation mechanism
of the calcium-activated chloride channel TMEM16A revealed by cryo-EM. *Nature* **552**,
421-425 (2017). https://doi.org:10.1038/nature24652

17 Alvadia, C. *et al.* Cryo-EM structures and functional characterization of the murine lipid
scramblase TMEM16F. *Elife* **8** (2019). https://doi.org:10.7554/eLife.44365
18 Bushell, S. R. *et al.* The structural basis of lipid scrambling and inactivation in the
endoplasmic reticulum scramblase TMEM16K. *Nat Commun* **10**, 3956 (2019).
https://doi.org:10.1038/s41467-019-11753-1
19 Bulley, S. *et al.* TMEM16A/ANO1 channels contribute to the myogenic response in
cerebral   arteries.   *Circ   Res*   **111**,   1027-1036   (2012).
https://doi.org:10.1161/CIRCRESAHA.112.277145
20 Dutta, A. K., Woo, K., Khimji, A. K., Kresge, C. & Feranchak, A. P. Mechanosensitive Cl-
secretion in biliary epithelium mediated through TMEM16A. *Am J Physiol Gastrointest
Liver Physiol* **304**, G87-98 (2013). https://doi.org:10.1152/ajpgi.00154.2012
21 Duncan, A. L., Song, W. & Sansom, M. S. P. Lipid-Dependent Regulation of Ion Channels
and G Protein-Coupled Receptors: Insights from Structures and Simulations. *Annu Rev
Pharmacol Toxicol* **60**, 31-50 (2020). https://doi.org:10.1146/annurev-pharmtox-010919-
023411
22 Brohawn, S. G., Su, Z. & MacKinnon, R. Mechanosensitivity is mediated directly by the
lipid membrane in TRAAK and TREK1 K+ channels. *Proc Natl Acad Sci U S A* **111**, 3614-3619
(2014). https://doi.org:10.1073/pnas.1320768111
23 Kern, D. M. *et al.* Structural basis for assembly and lipid-mediated gating of LRRC8A:C
volume-regulated   anion   channels.   *Nat   Struct   Mol   Biol* **30**, 841-852 (2023).
https://doi.org:10.1038/s41594-023-00944-6
24 Pliotas, C. *et al.* The role of lipids in mechanosensation. *Nat Struct Mol Biol* **22**, 991-998
(2015). https://doi.org:10.1038/nsmb.3120
25 Zhang, Y. *et al.* Visualization of the mechanosensitive ion channel MscS under membrane
tension. *Nature* **590**, 509-514 (2021). https://doi.org:10.1038/s41586-021-03196-w
26 Jojoa-Cruz, S. *et al.* Structural insights into the Venus flytrap mechanosensitive ion
channel Flycatcher1. *Nat Commun* **13**, 850 (2022). https://doi.org:10.1038/s41467-022-
28511-5
27 Reddy, B., Bavi, N., Lu, A., Park, Y. & Perozo, E. Molecular basis of force-from-lipids gating
in   the   mechanosensitive   channel   MscS.   *Elife*   **8**   (2019).
https://doi.org:10.7554/eLife.50486
28 Brohawn, S. G., del Marmol, J. & MacKinnon, R. Crystal structure of the human K2P TRAAK,
a lipid- and mechano-sensitive K+ ion channel. *Science* **335**, 436-441 (2012).
https://doi.org:10.1126/science.1213808
29 Zheng, W. *et al.* TMEM63 proteins function as monomeric high-threshold
mechanosensitive   ion   channels.   *Neuron*   (2023).
https://doi.org:10.1016/j.neuron.2023.07.006
30 Kalienkova, V. *et al.* Stepwise activation mechanism of the scramblase nhTMEM16
revealed by cryo-EM. *Elife* **8** (2019). https://doi.org:10.7554/eLife.44364
31 Flegler, V. J. *et al.* Mechanosensitive channel gating by delipidation. *Proc Natl Acad Sci U
S A* **118** (2021). https://doi.org:10.1073/pnas.2107095118
32 Rasmussen, T. *et al.* Interaction of the Mechanosensitive Channel, MscS, with the
Membrane Bilayer through Lipid Intercalation into Grooves and Pockets. *J Mol Biol* **431**,
3339-3352 (2019). https://doi.org:10.1016/j.jmb.2019.05.043

33 Fernandez-Ballester, G., Fernandez-Carvajal, A. & Ferrer-Montiel, A. Progress in the
Structural Basis of thermoTRP Channel Polymodal Gating. *Int J Mol Sci* **24** (2023).
https://doi.org:10.3390/ijms24010743

34 Jordt, S. E. & Julius, D. Molecular basis for species-specific sensitivity to "hot" chili
peppers. *Cell* **108**, 421-430 (2002). https://doi.org:10.1016/s0092-8674(02)00637-2

35 Bandell, M. *et al.* High-throughput random mutagenesis screen reveals TRPM8 residues
specifically required for activation by menthol. *Nat Neurosci* **9**, 493-500 (2006).
https://doi.org:10.1038/nn1665

36 Verkest, C. *et al.* Intrinsically disordered intracellular domains control key features of the
mechanically-gated ion channel PIEZO2. *Nat Commun* **13**, 1365 (2022).
https://doi.org:10.1038/s41467-022-28974-6

37 Richardson, J., Kotevski, A. & Poole, K. From stretch to deflection: the importance of
context in the activation of mammalian, mechanically activated ion channels. *FEBS J*
(2021). https://doi.org:10.1111/febs.16041

38 Suloway, C. *et al.* Automated molecular microscopy: the new Leginon system. *J Struct Biol*
151, 41-60 (2005). https://doi.org:10.1016/j.jsb.2005.03.010

39 Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).
https://doi.org:10.1038/nmeth.4193

40 Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).
https://doi.org:10.1038/nmeth.4169

41 Zhang, K. Gctf: Real-time CTF determination and correction. *J Struct Biol* **193**, 1-12 (2016).
https://doi.org:10.1016/j.jsb.2015.11.003

42 Scheres, S. H. RELION: implementation of a Bayesian approach to cryo-EM structure
determination. *J Struct Biol* **180**, 519-530 (2012).
https://doi.org:10.1016/j.jsb.2012.09.006

43 Zivanov, J., Nakane, T. & Scheres, S. H. W. Estimation of high-order aberrations and
anisotropic magnification from cryo-EM data sets in RELION-3.1. *IUCrJ* **7**, 253-267 (2020).
https://doi.org:10.1107/S2052252520000081

44 Ramlaul, K., Palmer, C. M., Nakane, T. & Aylett, C. H. S. Mitigating local over-fitting during
single particle reconstruction with SIDESPLITTER. *J Struct Biol* **211**, 107545 (2020).
https://doi.org:10.1016/j.jsb.2020.107545

45 Ramirez-Aportela, E. *et al.* Automatic local resolution-based sharpening of cryo-EM maps.
Bioinformatics **36**, 765-772 (2020). https://doi.org:10.1093/bioinformatics/btz671

46 Vilas, J. L. *et al.* MonoRes: Automatic and Accurate Estimation of Local Resolution for
Electron Microscopy Maps. *Structure* **26**, 337-344 e334 (2018).
https://doi.org:10.1016/j.str.2017.12.018

47 de la Rosa-Trevin, J. M. *et al.* Xmipp 3.0: an improved software suite for image processing
in electron microscopy. *J Struct Biol* **184**, 321-328 (2013).
https://doi.org:10.1016/j.jsb.2013.09.015

48 de la Rosa-Trevin, J. M. *et al.* Scipion: A software framework toward integration,
reproducibility and validation in 3D electron microscopy. *J Struct Biol* **195**, 93-99 (2016).
https://doi.org:10.1016/j.jsb.2016.04.010

49   Sanchez-Garcia, R. *et al.* DeepEMhancer: a deep learning solution for cryo-EM volume
post-processing. *Commun Biol* **4**, 874 (2021). https://doi.org:10.1038/s42003-021-02399-
1
50   Waterhouse, A. *et al.* SWISS-MODEL: homology modelling of protein structures and
complexes.   *Nucleic   Acids   Res   **46**,   W296-W303   (2018).
https://doi.org:10.1093/nar/gky427
51   Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta
Crystallogr D* **60**, 2126-2132 (2004). https://doi.org:10.1107/S0907444904019158
52   Casanal, A., Lohkamp, B. & Emsley, P. Current developments in Coot for macromolecular
model building of Electron Cryo-microscopy and Crystallographic Data. *Protein Sci* **29**,
1069-1078 (2020). https://doi.org:10.1002/pro.3791
53   Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and
electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861-877
(2019). https://doi.org:10.1107/S2059798319011471
54   Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and crystallography.
*Acta   Crystallogr   D   Struct   Biol   **74**,   531-544   (2018).
https://doi.org:10.1107/S2059798318006551
55   Wang, R. Y. R. *et al.* Automated structure refinement of macromolecular assemblies from
cryo-EM maps using Rosetta. *Elife* **5** (2016). https://doi.org:ARTN e17219
10.7554/eLife.17219
56   Moriarty, N. W., Grosse-Kunstleve, R. W. & Adams, P. D. electronic Ligand Builder and
Optimization Workbench (eLBOW): a tool for ligand coordinate and restraint generation.
*Acta   Crystallogr   D   Biol   Crystallogr   **65**,   1074-1080   (2009).
https://doi.org:10.1107/S0907444909029436
57   Williams, C. J. *et al.* MolProbity: More and better reference data for improved all-atom
structure validation. *Protein Sci* **27**, 293-315 (2018). https://doi.org:10.1002/pro.3330
58   Barad, B. A. *et al.* EMRinger: side chain-directed model and map validation for 3D cryo-
electron   microscopy.   *Nat   Methods   **12**,   943-946   (2015).
https://doi.org:10.1038/nmeth.3541
59   Afonine, P. V. *et al.* New tools for the analysis and validation of cryo-EM maps and atomic
models.   *Acta   Crystallogr   D   Struct   Biol   **74**,   814-840   (2018).
https://doi.org:10.1107/S2059798318009324
60   Rao, S., Klesse, G., Stansfeld, P. J., Tucker, S. J. & Sansom, M. S. P. A heuristic derived from
analysis of the ion channel structural proteome permits the rapid identification of
hydrophobic gates. *Proc Natl Acad Sci U S A* **116**, 13989-13995 (2019).
https://doi.org:10.1073/pnas.1902702116
61   Klesse, G., Rao, S., Sansom, M. S. P. & Tucker, S. J. CHAP: A Versatile Tool for the Structural
and Functional Annotation of Ion Channel Pores. *J Mol Biol* **431**, 3353-3365 (2019).
https://doi.org:10.1016/j.jmb.2019.06.003
62   Schrödinger, L. The PyMOL Molecular Graphics System, Version 2.1 Schrödinger, LLC.
63   Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612 (2004). https://doi.org:10.1002/jcc.20084
64   Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. *Protein Sci* **27**, 14-25 (2018). https://doi.org:10.1002/pro.3235

65   Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence
alignments using Clustal Omega. *Mol Syst Biol* **7**, 539 (2011).
https://doi.org:10.1038/msb.2011.75
66   Robert, X. & Gouet, P. Deciphering key features in protein structures with the new
ENDscript server. *Nucleic Acids Res* **42**, W320-324 (2014).
https://doi.org:10.1093/nar/gku316
626

Figures

Figure 1: Cryo-EM structure and pore of OSCA3.1.
a, Cryo-EM map of OSCA3.1 dimer colored by subunit. Nanodisc density in grey corresponds to the unsharpened map (gaussian-filtered to 1.5 σ). **b,** Front (left) and top (right) view of atomic model. Ex.: Extracellular, In.: Intracellular. **c,** Superposition of OSCA1.1, OSCA1.2 (in nanodiscs) and OSCA3.1 protomers. **d,** Pore profile of OSCAs in **c**. **e,** View of the pore pathway (blue) of OSCA3.1. Pore facing residues colored in yellow, with selected residues labeled. π-helical turns in pink. Putative pore lipid in green. Backbone of TM3 and TM4 helices hidden for clarity.

Figure 2: Amphipathic helix mutants of OSCA1.2.
a, Superposition of OSCA1.2 (grey) and OSCA3.1 (red). Insets: close-up view of amphipathic helix. Residues substituted in OSCA1.2 for electrophysiology experiments are underlined. **b,** Few cells expressing mutations in the amphipathic helix respond to poking compared to WT controls. Top panel: Maximum poke-induced currents observed in whole-cell mode for cells exposed to displacements up to 8.9 ± 0.6 μm (N=51; mean ± S.E.M.), 12.8 ± 0.7 μm (N=24), 12.4 ± 0.7 μm (N=16), and 12.8 ± 0.5 μm (N=24) above touching for WT, OSCA1.2$_{W75K}$, OSCA1.2$_{L80E}$, and OSCA1.2$_{W75K/L80E}$, respectively. Bottom panel: the apparent threshold in μm above touching the cell for this cohort. **c,** SAC maximal current (I_{max}) (top) and mmHg threshold (bottom) from WT and OSCA1.2$_{W75K/L80E}$-expressing cells reveal no significant differences in the ability of negative pressure to activate channels in cell-attached patches (Student's *t*-test). Also shown are data from single mutants OSCA1.2$_{W75K}$ and OSCA1.2$_{L80E}$. Too few patches were obtained for OSCA1.2$_{W75K}$ to compare I_{max}. Electrode resistances for **c** were similar in all cases (WT: 2.3 ± 0.1 MΩ (N=26); W75K: 2.1 ± 0.4 MΩ (N=3), L80E: 2.2 ± 0.1 MΩ (N=5), W75K/L80E: 1.8 ± 0.1 MΩ (N=11)). For panels **b-c**: individual cells are represented as scatter points; mean and S.E.M. are displayed.

Figure 3: OSCA1.2$_{OSCA3.1\text{-}BLD}$ chimera.
a, Amino acid sequence alignment of the BLD region of OSCA1.2 and OSCA3.1. Full alignment in **Supplementary Fig. 3.** Blue line at bottom of sequences denotes the sequence swapped in OSCA1.2$_{OSCA3.1\text{-}BLD}$ chimera. **b,** Poke-induced responses are observed in HEK-P1KO cells expressing OSCA1.2$_{3.1\text{-}BLD}$. Top panel: Maximum poke-induced currents observed in whole-cell mode for cells exposed to displacements up to 9.8 ± 0.8 µm (N=11; mean ± S.E.M.) and 11.8 ± 0.6 µm (N=19) above touching for WT and OSCA1.2$_{3.1\text{-}BLD}$, respectively. The percentage of cells with responses are shown above. Bottom panel: the apparent threshold in µm above touching the cell for this cohort. **c,** SAC I$_{max}$ (top) and mmHg threshold (bottom) from WT and OSCA1.2$_{3.1BLD}$-expressing cells reveal similar activity induced by negative pressure in cell-attached patches. Data shown were obtained from the same experiments. Electrode resistances for **c** were similar in all cases (WT: 2.4 ± 0.1 MΩ (N=12); OSCA1.2$_{OSCA3.1\text{-}BLD}$: 2.5 ± 0.1 MΩ (N=13)).

Figure 4: Probing the functional role of the potential lipid-interacting residues.
a, Superposition of OSCA1.2 and OSCA3.1 around the pore fenestration. Residues of OSCA1.2 predicted to interact with lipids and corresponding residues in OSCA3.1 are shown. **b,** Few OSCA1.2$_{K435I/K536I}$ -expressing HEK-P1KO cells respond to the poke stimulus while nearly all cells expressing OSCA1.2 WT channels respond (percentages shown above under the N of cells tested). Top panel: Maximum poke-induced currents observed in whole-cell mode for cells exposed to displacements up to 9.7 ± 1 μm (N=11; mean ± S.E.M.) and 12.1 ± 0.6 μm (N=20) above touching for WT and OSCA1.2$_{K435I/K536I}$, respectively. Bottom panel: the apparent threshold in μm above touching the cell for this cohort. **c,** SAC Imax (top) and mmHg threshold (bottom) from WT and OSCA1.2$_{K435I/K536I}$-expressing cells reveal no significant differences in the ability of negative pressure to activate channels in cell-attached patches (Student's *t*-test). Electrode resistances were similar in all cases (WT: 2.1 ± 0.2 MΩ (N=10); K435I/K546I: 1.7 ± 0.3 MΩ (N=7)). Data were collected within 15 min of exposure of cells to high K+ used in this assay. **d,** Analysis of stimulus-response relationships reveal no significant differences (Student's *t*-test) in pressure to half-maximal activation (P$_{50}$; top) or the slope of the curve (bottom). Whole cell and SAC data were obtained from cells transfected at the same time.