Manuscript title: Genome-wide associations of leaf spectral variation in MAGIC lines of
Nicotiana attenuata

Authors: Cheng Li[1*], Ewa A. Czyż[1†], Bernhard Schmid[1], Rishav Ray[2‡], Rayko Halitschke[2], Ian
T. Baldwin[2], Michael E. Schaepman[1], Meredith C. Schuman[1,3]

[1] Department of Geography, University of Zürich, Winterthurerstrasse 190, 8057 Zürich,
Switzerland
[2] Department of Molecular Ecology, Max Planck Institute for Chemical Ecology, Hans-Knoell-
Strasse 8, 07745 Jena, Germany
[3] Department of Chemistry, University of Zürich, Winterthurerstrasse 190, 8057 Zürich,
Switzerland
*Corresponding author: cheng.li@geo.uzh.ch

Keywords: Field spectroscopy | Genome-wide association studies (GWAS) | Hierarchical

Spectral Clustering with Parallel Analysis | Leaf reflectance | Multiparent Advanced Generation

Inter-Cross (MAGIC) | *Nicotiana attenuata*

[†] Present address: Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, 91109 Pasadena, California, USA
[‡] Present address: Department of Plant Biology, University of California, Davis, 605 Hutchison Drive, Green Hall 1002, 95616 Davis, California, USA

ABSTRACT

The application of in-field and aerial spectroscopy to assess functional and phylogenetic

variation in plants has led to novel ecological insights and supports global assessments of plant

biodiversity. Understanding how plant genetic variation influences reflectance spectra will help

harness this potential for biodiversity monitoring and improve understanding of why plants differ

in functional responses to environmental change. Here, we use a well-resolved genetic mapping

population derived from Multi-parent Advanced Generation Inter-cross (MAGIC) lines of

Nicotiana attenuata to associate genetic differences with differences in leaf spectra between

plants in a field experiment in their natural environment. We analyzed the leaf reflectance

spectra using a hand-held spectroradiometer (350–2500 nm) on 616 fully genotyped plants of *N.*

attenuata grown in a randomized block design. We tested three approaches to conducting

Genome-Wide Association Studies on spectral variants. We introduce a new Hierarchical

Spectral Clustering with Parallel Analysis (HSC-PA) method. This method efficiently captured

the variation in our high-dimensional dataset and allowed us to discover a novel association,

between a locus on chromosome 1 and the 734-1143 nm spectral range, spanning the red-edge

and near-infrared regions that are sensitive to leaf structure and photosynthetic activity. This

locus contains a candidate gene annotated as carbonic anhydrase, an enzyme involved in CO_2

hydration and regulation of photosynthetic efficiency, suggesting a physiological link between

variation in leaf optical properties and carbon assimilation. In contrast, an approach treating

single wavelengths as phenotypes identified the same associations as HSC-PA, but without the

statistical power to pinpoint significant associations. An index-based approach, which reduces

complex spectra to a few dimensionless variables, detected two significant associations for

$ARDSI\_C_w$ (a water-content-related index) with loci on chromosome 1 near genes annotated as a

Zeta toxin domain-containing protein, and an Exocyst subunit Exo70 family protein. While these

findings are biologically plausible, they represent a very narrow subset of the spectral variation

captured by HSC-PA. The HSC-PA approach supports a comprehensive understanding of the

genetic determinants of leaf spectral variation which is data-driven but human-interpretable, and

lays a robust foundation for future research in linking plant genetics with biodiversity

monitoring, large-scale ecological assessment and remote-sensing applications.

1   INTRODUCTION

Genetic diversity is essential for the survival and adaptation of species to changing

environmental conditions. Radiation reflected, absorbed, and transmitted by plants constitutes

the basis for remote sensing of vegetation, commonly in the range from visible to infrared

wavelengths corresponding to solar radiation (Knipling 1970; Thenkabail, et al. 2018). Remote

sensing may revolutionize the way we study genetic diversity by offering non-invasive methods

with the potential of repeated measurement on large spatial scales (Madritch et al. 2014). Remote

sensing technologies have not only facilitated the study of Earth's surface features, including

vegetation, water bodies, and land-use patterns, but are also gaining applications in biodiversity

research. Notably, high-throughput hyperspectral phenotyping has long been used in crop

quantitative genetics to link genetic variation with spectral traits, demonstrating the heritability

of multiple biochemical and physiological traits estimated from spectral signatures—thus

providing valuable methodological precedents for biodiversity and ecological studies

(Grzybowski et al., 2021). Wang and Gamon recently discussed concrete ways in which remote

sensing makes unique contributions to monitoring plant biodiversity (Wang and Gamon 2019).

For example, Féret and Asner showcased the capabilities of high-fidelity imaging spectroscopy

for mapping the diversity of tropical forest canopies (Asner et al., 2014). Although they do not

capture the structural variation of whole canopies, reflectance spectra from single leaves, which

can be captured with controlled lighting and background from plants in field conditions, also

offer insights into plant physiology, stress responses, and genetic variation, with recent studies

establishing a correlation between genetic diversity and leaf reflectance spectra (Serbin et al.,

2014; Cavender-Bares et al., 2016; Wang et al., 2018; Asner et al., 2015; Stasinski et al., 2021;

Czyż et al., 2023). We recently showed that genetically variable plant populations are also

spectrally more variable in comparison to replicates of an inbred genotype grown under either

glasshouse or field conditions, while isogenic plants differing primarily in their gene expression

are spectrally similar to replicates of the inbred genotype from which they were derived (Li et al.,

2023).

A Genome-Wide Association Study (GWAS) is a powerful tool used to identify genetic variants

associated with specific traits or diseases in populations. By examining the entire genome,

researchers can pinpoint specific genetic markers that correlate with phenotypic variation. A

primary advantage of GWAS is that it requires no prior knowledge of potential candidate genes

and can be used for the discovery of novel genetic associations (Visscher et al., 2017; Santure et

al., 2018). The corresponding disadvantage of GWAS is that it does not provide causal links and

thus the mechanisms underlying statistical associations must be dissected and tested. A primary

concern of GWAS is the influence of population structure, which, if not addressed, may produce

misleading associations (Price et al., 2006). Population structure refers to the presence of

subgroups within a population that differ in allele frequencies due to shared ancestry. In GWAS,

population structure can be confounding, because genetic variants associated with the subgroup,

rather than the trait of interest, may result in false-positive associations. To mitigate this, various

methods, such as principal component analysis (PCA) and mixed linear models (Zhang et al.,

2010), are employed to correct for population structure in GWAS analyses. Phenotyping remains

a substantial bottleneck in genetic studies due to its labor-intensive and time-consuming nature.

High-throughput phenotyping, especially hyperspectral imaging, offers a powerful solution—

enabling rapid, scalable, and non-invasive measurements of plant traits that can accelerate

genotype–phenotype association studies (Furbank & Tester, 2011; Zhang & Zhang, 2018).

Nicotiana attenuata Torr. ex S. Watson is a model wild plant for molecular ecology native to the

Great Basin Desert of the southwestern USA. It is primarily found in large ephemeral

populations following fires in sagebrush and pinyon-juniper ecosystems. Its unique germination

behavior is stimulated by cues found in wood smoke and the removal of inhibitors from

unburned litter, allowing it to thrive in post-fire environments (Bahulikar et al., 2004).

Ecologically, *N. attenuata* is a compelling subject for its intricate defense mechanisms against

herbivores, its tissue-specific diurnal metabolic rhythms, and its ability to acclimate to varying

environmental conditions, including high UVB radiation (Glawe et al., 2003; Kim et al., 2011; Li

et al., 2016; INH, Galis, and Baldwin 2013). Further, it offers unique opportunities for genetic

research due to a well-defined MAGIC (Multiparent Advanced Generation Inter-Cross)

population (Ray et al., 2019, 2023). This design comprises 26 phenotypically and genetically

differentiated parental lines (PLs) intercrossed to produce two replicate sets each of 325

recombinant inbred lines (RILs; 616 genotyped plants of the total of 650 used in the present

study), allowing for the study of phenotype-genotype associations without the confounding

influence of population structure. More details on plant material, field site, and generation of

MAGIC RIL Population can be found in Appendix S1: Supporting Text.

In this study, we explore the genetic diversity of the MAGIC population to identify associations

between genetic variants and leaf spectral traits, advancing our understanding of genetic

influences on complex spectral characteristics. Our methodology involves three approaches:

analyzing spectral indices, single wavelengths (SW), and implementing Hierarchical Spectral

Clustering with Parallel Analysis (HSC-PA). Through GWAS using 168,903 SNPs that showed

variation within the population, we investigate genetic links with genome annotation. Our

methodological approach and the tools utilized are summarized in Figure 1.

2   MATERIALS AND METHODS

This study's methodology extends from our previous work (Li et al., 2023), with detailed

description available in Appendix S1: Supporting Text. Briefly, our study utilizes two replicate

sets each of 325 *N. attenuata* MAGIC RILs, derived from 26 parental lines and six generations

of inbreeding, yielding a total of 650 genetically different plants of which 616 were planted in

the current experiment. The RILs sequencing was performed at Novogene HK (Ray et al., 2023),

and leaf optical properties (350-2500 nm) were measured using a FieldSpec 4 spectroradiometer

(Li et al., 2023).

Data analysis was performed using R (R Core Team 2023), employing the spectrolab package

(version 0.0.10, Meireles, et al., 2017) and a three-step filtering approach for outlier removal (Li

et al., 2023). We employed a linear model to remove known environmental effects from the raw

spectral data. Specifically, for each wavelength, we fitted the model:

$$Y \sim LN + REP + BA + LN:(REP + BA) + DT:CT + MO$$

where LN represents the influence of the measurement technique (the number of measured

leaves), REP accounts for potential differences between the two replicate sets of RILs (that

occupied different sectors of the field plot and were measured in series), BA represents spatial

patch effects characterized from *N. attenuata* reference measurements (in each square of 2×2

planting positions one was occupied by a "phytometer" plant of a single tester genotype of the

species), $LN: (REP + BA)$ models the interaction between measurement technique and spatial

effects, $DT: CT$ captures temporal effects (DT for day and CT for time of day), and MO denotes

the maternal cytoplasmic effects. The residuals from each fitted model were taken as the adjusted

spectral values, corrected for (i.e., removing) the modeled environmental influences.

We tested three ways of treating the spectral phenotypic data. First, we used six selected spectral

indices, which are single values derived from ratios of reflectance at specific wavelengths

representing spectral features (details see Table 1 and Appendix S1: Supporting Text). These

indices span pigment-related (e.g., chlorophyll a/b, carotenoids), water-related, and structural

spectral features across the VIS, red-edge, NIR, and SWIR ranges, thus capturing major axes of

leaf functional variation. The indices were first calculated from the raw spectral data and

subsequently adjusted using the same linear model as described above. Second, we used single

wavelengths (SW), i.e., treating each wavelength in the adjusted spectrum as a phenotype. Third,

we used Hierarchical Spectral Clustering with Parallel Analysis (HSC-PA), a new data-driven

dimension reduction approach, developed from a method for the analysis of genetic associations

with human facial features (Claes et al., 2018; Sero et al., 2019), which accounts for correlations

within spectral data while retaining interpretable features. We applied HSC-PA to the adjusted

spectral data for downstream analyses. We associated the resulting phenotypic data with 168,903

SNPs using multiple GWAS models based on the Genome Association and Prediction Integrated

Tool (GAPIT) version 3 (Wang and Zhang 2021). The GWAS models include: General Linear

Model (GLM, Price et al., 2006), Mixed Linear Model (MLM, Yu et al., 2006), Fixed and

random model Circulating Probability Unification (FarmCPU, Liu et al., 2016), and Bayesian

information and Linkage-disequilibrium Iteratively Nested Keyway (BLINK, Huang et al.,

2019). Kinship matrices were automatically calculated by GAPIT using the Zhang method

(Zhang et al., 2010) from the genotype data; these matrices were applied only in models that

incorporate kinship (i.e., MLM). See more details of the four models in Appendix S1 Supporting

Text.

Multiple testing adjustments were conducted on 168,903 SNPs. Initially, an exploratory

threshold of $p < 1 \times 10^{-5}$ was used for preliminary SNP selection. This was followed by a false

discovery rate (FDR) adjustment (Benjamini and Hochberg 1995) and the application of the

PhenoSpD method (Nyholt 2004; Zheng et al., 2018) to determine the significance threshold

$(0.05/M_{eff})$, taking into account the effective number of independent trait calculations. The

PhenoSpD method first derives the pairwise phenotypic correlation matrix from GWAS

summary statistics using cross-trait LD Score regression. Specifically, for two traits t_1 and t_2, the

regression

$$Z_{1j} Z_{2j} = a + b \cdot LDscore_j + \epsilon_j$$

165   is calculated across SNPs j, where Z_{1j} and Z_{2j} are the z-scores from GWAS summary statistics

166   for each trait, and the intercept a provides an estimate of the phenotypic correlation between

167   traits. This phenotypic correlation matrix is then subjected to spectral decomposition (SpD), in

168   which the eigenvalues (λ_i) are used to calculate the effective number of independent tests:

$$M_{eff} = \frac{\left(\sum_{i=1}^{M} \lambda_i\right)^2}{\sum_{i=1}^{M} \lambda_i^2}$$

where M is the total number of phenotypes. We used $0.05/M_{eff}$ as the significant threshold.

SNPs with FDR-adjusted p-values below the significance threshold were considered significantly

associated. For candidate gene selection, we extracted genes within ±100 kb of significant SNPs

from the *N. attenuata* genome, following the approach used in closely related *Nicotiana tabacum*

(Lai et al., 2021). We then refined the candidate gene list using the *N. attenuata* Data Hub

(http://nadh.ice.mpg.de/NaDH/) to filter out genes lacking transcript accumulation evidence in

leaves under various conditions. Genes with relevant functions were then identified based on

annotation information.

We also conducted GWAS using indices, single wavelengths, and HSC-PA segments with raw

spectral data as a general comparison to assess the effect of removing known environmental

effects (see Appendix S1: Supporting Text, Results from raw spectral data).

3   RESULTS

3.1   Phenotypic analysis

3.1.1   Variance partitioning and adjustment of raw spectra

To advance our understanding of the phenotypic variation in leaf spectral traits and to remove

the known environmental effects, we employed linear models. These models partitioned variance

across the spectral range of 350–2500 nm, assessing contributions to cumulative multiple R^2

values (i.e. percent sum of squares) from measurement technique (LN), replicate and spatial

effects and their interactions with measurement technique ($REP + BA + LN:(REP + BA)$),

measurement time ($DT:CT$), and maternal cytoplasm (MO) within the 616 plants. Figure 2

illustrates these components for each wavelength as stacked bars, which—given the large

number of wavelengths—merge into a continuous stacked profile that collectively accounts for

the total variance. Analysis of variance results indicated that residual phenotypic variance

between plants, representing genetic and residual environmental contributions accounted from

71.7% to 89.5% of the total phenotypic variance.

3.1.2   Spectral characteristics of leaves

Our analyses focus on calculated leaf reflectance spectra spanning from 400 to 2500 nm, as the

initial region (350–399 nm) has a poor signal-to-noise ratio and is commonly excluded from

downstream analyses (Cavender-Bares et al., 2016). This range includes the visible (VIS), near

infrared (NIR), and shortwave infrared (SWIR) regions. This broad spectral range captured

variation among the 616 plants, shown in Figure 3a as an orange shaded region indicating the

range of adjusted reflectance spectra for all samples. The blue shaded region indicates the range

of raw reflectance spectra before adjustment with linear models. The NIR region showed more

consistency in reflectance values among the plants, contrasting with greater variability in the VIS

and SWIR regions.

The distribution of six selected spectral indices (Figure 3b), encapsulating attributes like water

content and chlorophyll concentration, provided more specific information about leaf

physiology. Note that the y-axis represents the index values, which are specific to each index

calculation and therefore cannot be directly compared across indices.

Our subsequent application of the HSC-PA segmented the adjusted spectra into 38 distinct

patterns. The blank (NA) areas are regions where further clustering was halted due to the

retention of a singular Principal Component (PC) from a previous level. This approach allowed

for a finer granularity in the spectral data representation while accounting for correlations among

wavelengths. The hierarchical clustering process concluded at different levels, ranging from

level 3 to level 8, indicating the varying depths of spectral similarities among the segments.

Table 2 shows the detailed spectral region and potential associated features of the segments used

in HSC-PA-GWAS. Appendix S1: Table S1 shows the spectral regions of all 38 segments.

By visualizing the 20 segments that each retained only a single PC, we were able to span the

entire spectrum from 400 to 2500 nm. These segments, parsed into several interpretable regions,

were then used for HSC-PA-GWAS approach, offering phenotypic data for uncovering

associations with underlying genetic variations.

The distributions of the six indices derived from raw spectra, along with the HSC-PA results

based on raw spectral data, are shown in Appendix S1: Figure S1.

3.1.3   Effective number of independent traits (M_{eff})

To account for multiple testing due to the analysis of multiple phenotypes in our GWAS

approaches, we determined the effective number of independent traits, denoted as M_{eff}, using

the PhenoSpD method (Zheng et al., 2018). For the Index-GWAS approach, which employed six

distinct spectral indices, the effective number of independent tests, $M_{eff_{ind}}$, was determined to

be 5.0. In the Single Wavelength GWAS (SW-GWAS) approach, the effective number, $M_{eff_{sw}}$,

was notably higher at 999.9. Lastly, for the Hierarchical Spectral Clustering with Parallel

Analysis GWAS (HSC-PA-GWAS) approach, which clusters the spectra based on similarity, the

effective number, $M_{eff_{hsc-pa}}$, was found to be 10.5, reflecting the reduced dimensionality and

the focus on major patterns of variation. These calculated M_{eff} values are used to set a threshold

($0.05/M_{eff}$) for significant associations.

3.2   SNP marker analysis

The kinship relationships among the 616 plants are visualized in Appendix S1: Figure S2.

Through a heatmap, the genetic relatedness between individuals is presented. Most values in the

histogram gravitate towards zero, indicating a consistent and minimal genetic difference between

plants. This uniform distribution of kinship values affirms the thorough genetic mixing of the

parental lines in the MAGIC population and is consistent with the finding that this population

does not display obvious structure (Ray et al., 2023).

Appendix S1: Figure S3 presents the frequency of heterozygosity of individual and markers,

while Appendix S1: Figure S4 illustrates the linkage disequilibrium (LD) decay over distance.

3.3   Results of genome-side association studies (GWAS)

3.3.1   Index-GWAS results

The Manhattan plot presented in Figure 4a, shows the associations between the SNPs and the six

indices ($NDWI$, CCI, CI_{re}, $ARDSI\_C_{ab}$, $ARDSI\_C_w$, and $ARDSI\_C_m$) using four models (GLM,

MLM, FarmCPU, and BLINK). Several genomic regions displayed distinct associations

according to the $-\log_{10}$(p-values). An exploratory threshold, defined at 1×10^{-5} (denoted by

the grey horizontal line), was adopted to shortlist SNPs for a detailed examination.

In Figure 4b, each of the 6 indices is plotted against the SNPs that exceeded the exploratory

threshold. The gradient color scale indicates the association's intensity, with deeper hues

indicating smaller FDR-adjusted p-values. This representation provides a concise overview of

the genomic regions that are most strongly correlated with each spectral index. In total, 12 SNPs

exhibited p-values less than 1×10^{-5}: five with $ARDSI\_C_w$, three with CCI, two each with

$NDWI$ and CI_{re} and one each with $ARDSI\_C_m$ and $ARDSI\_C_{ab}$. The index $ARDSI\_C_w$ displayed

two associations lower than the significance threshold: $\frac{0.05}{5.0} = 0.01$, with two SNPs in

chromosome 1. The first SNP was located at chr1_193505583, near Niat3g_08138, annotated as

a Zeta toxin domain-containing protein. The second was at chr1_438155938, close to

Niat3g_17277 (TPR_REGION domain-containing protein) and Niat3g_17282 (Exocyst subunit

Exo70 family protein). Furthermore, $ARDSI\_C_w$ was chosen as a representative example for the

QQ plots in Figure 4c and 4d. This plot serves as a diagnostic tool for the GWAS results. Each

model is represented by a unique color: red for GLM, yellow for MLM, green for FarmCPU, and

blue for BLINK. The alignment of the observed points with the diagonal line indicates that the

GWAS results are mostly consistent with the expectations under the null hypothesis. However,

deviations can pinpoint regions with stronger genetic signals. Figure 4e and 4f show the

phenotypic distributions for two significant markers, where 0 represents a homozygous reference

allele, 1 a heterozygous allele, and 2 a homozygous alternative allele.

3.3.2   SW-GWAS and HSC-PA-GWAS results

In the SW-GWAS analysis, the BLINK model was selected to investigate the genetic

correlations with 2101 spectral wavelengths spanning from 400 to 2500 nm. Figure 5a highlights

the genetic associations for the spectral wavelength at 952 nm, which displayed the strongest

association among all phenotypes, serving as an example. The diagnostic QQ plot for this

wavelength is also presented in Figure 5c. Figure 5b offers an in-depth depiction of associations

that exceeded the exploratory threshold. The X-axis represents the spectral wavelengths, the Y-

axis the associated SNPs, providing a holistic view of the spectral regions correlated with each

SNP. A total of 27 SNPs are displayed, with the majority linked to one or multiple continuous

spectral regions. This overlap indicates the high correlation inherent in the leaf reflectance

spectra data, underscoring the need for methods that can account for these correlations.

Furthermore, correlations with the same SNP spanning more distant spectral regions may also

reflect underlying biological processes and warrant further investigation. The associations span

nearly the entire spectral range and are distributed across most chromosomes. The significance

level, adjusted by M_{eff}, stands at 5.0×10^{-5}. Notably, no SNP met this threshold.

In the HSC-PA-GWAS, Figure 6a shows the associations that surpassed our exploratory

threshold of 1×10^{-5}. A total of 20 SNP associations distributed over 20 segments emerged

from the analysis, all of which were also found in the SW-GWAS. Focusing on segment

associations obscures spectral information, and so Figure 6b maps these associations back to

their corresponding spectral wavelengths for better interpretation. This visualization facilitates

the identification of specific spectral regions linked to each SNP. It is noteworthy that these

associations span a range of chromosomal regions, indicating a complex genetic basis of the

spectral traits. The significance level, after adjusting for M_{eff}, is set at 4.7×10^{-3}. Two

associations, the SNP chr1_131409683 with Segments 37 and 38 (734-1143 nm), surpassed this

threshold. The QQ plots for Segments 37 and 38, as depicted in Figure 6c and d, serve a dual

purpose of validation and diagnostic assessment. Figure 6e and f show the phenotypic

distributions of the significant marker (chr1_131409683) affiliated with Segments 37 and 38. In

this distribution, the numbers 0, 1, and 2 represent homozygous alleles as the reference,

heterozygous alleles, and homozygous alternative alleles, respectively. The majority of the RILs

are assigned reference or heterozygous alleles at this site, as evidenced by the greater number of

points in classes 0 and 1. A clear difference in the mean values of these three classes is evident in

the boxplot.

To further elucidate the genetic association indicated by HSC-PA-GWAS, we searched for

potential candidate genes within a $\pm$ 100 kb window surrounding the SNP chr1_131409683. This

led to the identification of one gene of interest: Niat3g_06527 (more annotation information can

be found in Appendix S1: Table S2). This gene is annotated as Carbonic anhydrase (CA).

We broadened our analysis to include the top 10 SNPs from HSC-PA GWAS with the smallest

FDR-adjusted P-values (see Appendix S1: Table S2). The top six associations are linked to

Segment 37 or Segment 38 in the red-edge and near-infrared regions. In addition to the

significantly associated locus (chr1_131409683), these associations involve two SNPs:

chr12_4508375 and chr10_23318853. Furthermore, the significantly associated locus

(chr1_131409683) is also associated with Segment 34 (1144–1310 nm), which lies immediately

after Segments 37 and 38 and remains within the NIR region.

4   DISCUSSION

4.1   HSC-PA-GWAS for discovering genetic associations with spectral variation

Understanding the genetic foundation of leaf spectral diversity will facilitate advances in

biodiversity monitoring and phenotyping, but determining appropriate approaches to extract

genetic information from complex, context-dependent spectral phenotypes is a critical step (Li et

al., 2023). SW-GWAS, while straightforward, is limited by the high dimensionality of spectral

data and the spectral resolution of instruments, typically ranging from 3-10 nm. Subsampling can

simplify data and enhance signal-to-noise ratios, yet risks omitting key data points, and the

choice of bands for subsampling and the resulting number of traits require consideration and

testing. It is common to subsample the continuous spectral data to 1 nm intervals, as done here

for the SW-GWAS and as a basis for other analyses, because 1 nm is the subsampling interval

provided in the calibrated output data from the ASD FieldSpec and other field spectrometers.

Cavender-Bares and colleagues (Cavender-Bares et al., 2016) subsampled leaf spectra at a larger

interval of 5 nm prior to PLS-DA and evolutionary analysis and stated that the decision

regarding the interval width (1, 2, or 10 nm) did not significantly impact the results. Czyż and

colleagues used a coarser subsampling interval of 10 nm to estimate the predictive power of

spectral bands for genetic structure (Czyż et al., 2020). Dimensionality reduction methods like

PCA yield complex components challenging to biological interpretation (Li et al., 2023).

Similarly, spectral indices and methods like varimax rotation have limitations due to continuous

spectra and multiple influencing factors. Partial least squares regression (PLSR) is another

alternative, often combined with discriminant analysis or beta regression models, but it requires

orthogonal trait measurements for accurate associations and carries the risk of overfitting

(Cavender-Bares et al., 2016; Wang et al., 2020; Stasinski et al., 2021).

Here, we developed HSC-PA-GWAS as a new approach that retains the information in spectra in

a biologically interpretable form without losing meaningful information, and without requiring

an *a priori* knowledge to target specific traits, while maintaining sufficient statistical power to

identify potentially meaningful associations. This method stands out for its combination of

sensitivity and statistical power when handling spectral phenotypes. Associations identified

using HSC-PA-GWAS were consistent with the SW-GWAS method, as every SNP identified by

HSC-PA-GWAS was also detected by SW-GWAS. However, the HSC-PA method surpasses

SW-GWAS in several key aspects. One of its strengths is its ability to reduce data dimensionality

while accounting for phenotypic correlations within spectra, by transforming multiple correlated

wavelengths into a single value, typically represented by the first principal component. This

mitigates the challenges of multiple testing and correlated phenotypic data and thus increases

statistical power. By aggregating information across multiple wavelengths, the HSC-PA method

offers a more comprehensive and potentially more accurate representation of the underlying

traits with its ability to discover the correlation structure among wavelengths from data. The

resulting genetic associations encompass a broader portion of the spectrum than those derived

from indices, which focus on specific wavelength combinations, or from single wavelengths,

which do not behave independently. While indices remain valuable for highlighting proxies of

relevant biological processes or capturing relative changes between spectral regions, the HSC-

PA approach retains a wider spectral context for interpretation. Moreover, compared to spectral

subsampling (e.g., sampling or averaging every 10 nm), which typically does not allow for

(varying) covariation patterns of spectral bands and focuses only on neighboring bands, the

HSC-PA method provides a more data-driven approach that accounts for empirical internal

correlations.

Comparing the HSC-PA method with established methods, such as an approach combining

partial least squares regression (PLSR) and linear regression to interpret spectral features in

terms of other traits (Verrelst et al., 2019), indicates that HSC-PA has an advantage in at least

two ways: supporting human interpretation of data-derived spectral features and thus genetic

associations, and allowing for experimental designs aimed at discovering associations with novel

trait variation or where there are limitations on producing appropriate validation datasets

required for regression. Verrelst and colleagues (Verrelst et al., 2019) also emphasized the

importance of nonparametric regression and machine learning methods, like the Kernel Ridge

Regression (KRR) and Gaussian Processes Regression (GPR), for their ability to capture

nonlinear relationships in spectral data. These methods, while powerful, often require careful

tuning, can be computationally intensive, and carry the danger of overfitting. The HSC-PA

method offers a data-driven yet human-interpretable approach with the caveat that its criterion

for simplification is linear (dimensionality in a PCA). As discussed in studies on human facial

shapes, from which the HSC-PA method was adapted (Claes et al., 2018), this methodological

shift allows for a transition from a global to a local understanding of spectral variation, and a

holistic view of its genetic basis.

4.2   Spectra adjustment and genetic associations

In this study, we used replicated plants of a single tester genotypes of *N. attenuate* as

"phytometers" to account for small-scale (2x2 planting positions) spatial variation across an

experimental field. In addition, we accounted for differences in other environmental variables

such as measurement time as well as for maternal effects. By applying a linear model across the

full spectral range and using the residuals as adjusted spectra for downstream analyses, we were

able to reduce the influence of confounding environmental variables. This correction, which

could not have been incorporated directly into a mixed-model GWAS analysis, substantially

altered many results, including derived phenotypes and GWAS associations, leading to different

segment definitions and candidate genes in comparison to directly conducting GWAS on spectral

data without prior correction for non-genetic effects (Appendix S1: Supporting Text, Results

from raw spectral data). Such shifts underscore the importance of accounting for non-genetic

effects in spectral-genomic studies, as failing to do so can lead to markedly different—and

potentially misleading—biological conclusions. Here, we were able to do so both by using a

phytometer and by accounting for environmental factors dictated by the experimental and

measurement design; while a genetically invariable phytometer is rarely an option in ecological

studies, it should usually be feasible to account for quantitative environmental and experimental

effects on phenotype values.

Advanced genetic association models considering epistasis, akin to HSC-PA's handling of

spectral correlations, could enhance GWAS power, sensitivity, and interpretability, with

advances in machine learning opening new possibilities (Libbrecht and Noble 2015; Wu et al.,

2021). Random forest models have already been used for analyzing complex genetic effects on

multivariate phenotypes, though their significance interpretation differs (Brieuc et al., 2018;

Wang et al., 2013). The "Epi-MEIF" approach by Saha et al., (2022), using conditional inference

forests for epistasis in complex phenotypes, offers a method for significant association

identification, accommodating multiple loci testing as an add-on to conventional single-locus

models like those in GAPIT.

4.3   Ecological significance of findings in *N. attenuata*

We discovered a significant association between the SNP chr1_131409683 and Segments 37 and

38 (734-1143 nm) in the HSC-PA-GWAS. One interesting candidate gene was identified within

100 kb of this SNP: Niat3g_06527, is annotated as a carbonic anhydrase (CA). Carbonic

anhydrases catalyze the reversible hydration of CO_2 to bicarbonate and protons ($CO_2 + H_2O \rightleftharpoons$

$HCO_3^- + H^+$) (Supuran, 2008). In plants, multiple CA isoforms are found in the chloroplast

stroma, thylakoid membranes, cytosol, and other compartments (Fabre et al., 2007; Rudenko et

al., 2015). Chloroplastic CAs are thought to facilitate the rapid supply of CO_2 to Rubisco by

accelerating the interconversion between dissolved inorganic carbon species, thereby supporting

the Calvin–Benson cycle (Badger & Price, 1994). In C_4 and CAM plants, CA activity in

mesophyll cells is also critical for providing bicarbonate to phosphoenolpyruvate carboxylase

during the initial fixation step (Hatch & Burnell, 1990). In C_3 species such as *N. attenuata*,

stromal and thylakoid-associated CAs have additionally been implicated in maintaining

photosynthetic efficiency under fluctuating light and CO_2 conditions, and in the regulation of pH

and ion balance in the thylakoid lumen (Tiwari et al., 2016).

The associated spectral range, 734-1143 nm, spans the red-edge and near-infrared regions, which

are sensitive to changes in leaf internal structure and chlorophyll content. Genetic variation in a

chloroplast-localized CA could influence CO_2 assimilation rates and downstream effects on

chlorophyll concentration, mesophyll cell structure, or water status, any of which can alter

reflectance in this part of the spectrum. The link between Niat3g_06527 and these spectral

regions therefore suggests that natural allelic differences in CA function may contribute to

variation in photosynthetic performance and leaf optical properties in *N. attenuata*.

In the context of the ecology of *N. attenuata*, understanding the genetic basis of spectral

variation can provide insights into the adaptive mechanisms of this wild plant. The significant

association of a SNP close to a gene annotated as carbonic anhydrase, with a spectral range

spanning the red-edge and near-infrared regions—wavelengths closely linked to chlorophyll

content, canopy structure, and photosynthetic capacity—may reflect evolutionary pressures that

have shaped the species' genetic architecture to optimize carbon fixation and water-use

efficiency, enhancing its survival in the variable and often harsh environments of its native

habitat. Our extended analysis of the top 10 SNPs has unveiled additional genetic associations

that may play important significant roles in the ecological adaptation and survival of *N.*

attenuata.

Two other statistically significant associations were discovered with the $ARDSI\_C_w$ index, an

index of water content that uses the wavelengths 1080, 1240, 1360, and 1560 nm, which are

mostly (except for 1080 nm) in the range of wavelengths longer, *i.e.*, further into the infrared

region, than those comprising segments 37 and 38. The two associated SNPs, both also on

chromosome 1, were located in the vicinity of the gene Niat3g_08138 (chr1_193505583),

annotated as a Zeta toxin domain-containing protein, and the genes Niat3g_17277 annotated as

TPR_REGION domain-containing protein and Niat3g_17282 annotated as Exocyst subunit

Exo70 family protein (chr1_438155938). Proteins with these domains are documented to serve a

wide variety of functions in defense, growth and development, and cellular function.

It is interesting that the strongest associations were all identified between loci on chromosome 1

and wavelengths longer than those of visible light, in the red edge to infrared portions of the leaf

spectrum, that are influenced by micromorphology, structure, abundant non-pigment organic

constituents, and water content. The strongest associations were within regions of high

signal:noise in spectral reflectance, *i.e.*, where the detection of phenotypic differences should be

most robust.

4.4 Limitations and outlook

One limitation of our study lies in the phenotypic interpretation for biological meanings specific

to the samples we used. The six spectral indices we selected are derived from existing literature

and have not been validated in *N. attenuata* generally, or for this dataset. Without chemical or

other analyses to specifically quantify the constituents represented in the indices, such as

chlorophyll, our interpretations remain speculative. Establishing causal links between candidate

genes and specific biological processes is inherently difficult, particularly in ecological systems

where multiple interacting factors may influence spectral traits (Hoban et al. 2016). Moreover,

while the HSC-PA-GWAS approach has identified potential candidate genes, any influence of

these genes on the traits of interest remains to be verified, e.g. through the generation and testing

of knock-out, knock-down, and overexpression lines.

While GWAS with a MAGIC population does not require replication of genotypes, the absence

of such replicates prevented us from testing variation between genotypes directly using the

analysis of variance approach. Nevertheless, by removing small-scale environmental variation

using replicated phytometers of a single genotype as well as other environmental variation and

maternal effects, the residual phenotypic differences between plants was likely due to their

genetic differences rather than further environmental causes.

Our study noted unexpectedly high heterozygosity levels (mean around 0.5, see Appendix S1:

Figure S3) in the dataset, potentially due to the low-pass sequencing (mean 0.5x coverage) of the

RILs (Ray et al., 2023). The significant data gaps (about 95% missing data) necessitated

imputation from parental lines, which weren't highly inbred, likely inflating heterozygosity

estimates in the RILs. While the imputed genotype matrix was accurate for known QTLs (Ray et

al., 2023), the sequencing and imputation approach introduces uncertainty in true heterozygosity

levels. Factors like residual heterozygosity and heterozygote hotspots, seen in species like maize

(Liu et al., 2018), may also influence these high heterozygosity readings in the *N. attenuata*

MAGIC RIL population. Future studies with higher sequencing depth could more accurately

determine homozygosity and heterozygosity in this population, aiding in dissecting genetic

architecture and validating GWAS findings.

Looking forward, there is great potential for deepening our understanding of leaf spectral data

using the *N. attenuata* system. Our analysis indicates relatively large variance in blue light

reflectance and perhaps related photosynthetic traits among the wild genotypes contributing to

the *N. attenuata* MAGIC population, as discussed previously (Li et al., 2023). Variation in

conserved and essential traits, which can be mediated by variants of regulatory genes embedded

in coexpression networks (Ray et al., 2023), may lend different genotypes a competitive

advantage under changing environmental conditions. The significant associations identified here

thus indicate genotype and phenotype targets for novel and interesting in-depth functional studies

that could help us to better understand plant survival strategies. A combination of natural

variants, genetic modification, and laboratory assays, such as gene expression analyses (Schena

et al., 1995), protein-protein interactions (Fields and Song 1989), and chromatin

immunoprecipitation assays (Solomon et al., 1988), comprise the gold standard for functional

testing and validation of discovered genetic associations. At this stage, the associations reported

here represent hypotheses that should be subjected to these tests.

Unraveling the genetic basis of leaf spectral properties has far-reaching implications and could

for example revolutionize plant biodiversity monitoring, ecophysiology, and adaptation research,

and help to identify genotypes resilient to stresses associated with global change. Our study sets

the stage for future research aimed at deciphering these links by presenting a widely applicable

new method for data-driven discovery of associations which can be integrated with existing and

novel tools to discover genetic associations and support moving from association to prediction or

causation.

5   CONCLUSIONS

In our study, we explored the genetic basis of leaf spectral properties in *N. attenuata* by

associating spectral phenotypes with SNP genotypes in GWAS during a field experiment. We

analyzed leaf reflectance spectra from 400 to 2500 nm, identifying regions correlating with leaf

composition and function. We developed the Hierarchical Spectral Clustering with Parallel

Analysis (HSC-PA) method to manage the data's high dimensionality and correlation, proving

more effective than other methods like spectral indices or individual wavelengths. This approach

offered greater statistical power and maintained interpretable features, despite reduced

dimensionality.

Our findings included a significant SNP association in the red-edge to near-infrared spectral

range (734–1143 nm), a region sensitive to chlorophyll content, canopy structure, and

photosynthetic efficiency. The most significant SNP lies near the Niat3g_06527 gene, annotated

as carbonic anhydrase, an enzyme central to photosynthetic carbon fixation and stomatal

regulation. These results require further validation, but they raise the intriguing possibility that

variation in red-edge/NIR reflectance could be linked to differences in carbon assimilation and

water-use strategies in *N. attenuata*, potentially reflecting adaptations to its native, resource-

variable environments.

This study introduces a method and framework that can be broadly applied to attain deeper

understanding of the genetic underpinnings of leaf spectral properties. The associations identified

here using an advanced genetic toolset built for a native plant, and studied in a field experiment

within the plant's natural range, have the potential to advance our understanding of the genetic

mechanisms underlying photosynthetic efficacy and related growth and defense strategies in

plants, and could support new insights in the application of remote sensing to biodiversity

assessment.

520 6   ACKNOWLEDGMENTS

The authors acknowledge funding from the NOMIS Foundation and the University of Zurich,

including the University Research Priority Program on Global Change and Biodiversity, and the

Max Planck Society. We thank our division in the Department of Geography at the University of

Zurich, the Remote Sensing Laboratories, for support and for the shared use of equipment. We

thank the local managers at WCCER, N. Carlson and R. Carlson, and the 2019 field crews for

plant cultivation and plot management. We thank the WCCER for supporting these experiments.

527 7   AUTHOR'S CONTRIBUTIONS

CRediT taxonomy roles are listed with authors in alphabetical order. Conceptualization: I.T.B.,

M.C.S., M.E.S. Data curation: B.S., C.L., R.R. Formal analysis: B.S., C.L. Funding acquisition:

I.T.B., M.E.S. Investigation: E.A.C., M.C.S. Methodology: B.S., C.L., E.A.C., M.C.S. Project

administration: M.C.S. Resources: I.T.B., M.E.S., R.H. Supervision: M.C.S. Visualization: C.L.

Writing—original draft: C.L. Writing—review and editing: B.S., C.L., E.A.C., I.T.B., M.C.S.,

M.E.S., R.H., R.R.

8   CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

9   REFERENCES

Adam, Elhadi, Onisimo Mutanga, and Denis Rugege. 2010. "Multispectral and Hyperspectral

Remote Sensing for Identification and Mapping of Wetland Vegetation: A Review." *Wetlands*

Ecology and Management 18: 281–96.

Asner, Gregory P, Roberta E Martin, Christopher B Anderson, and David E Knapp. 2015.

"Quantifying Forest Canopy Traits: Imaging Spectroscopy Versus Field Survey." *Remote*

Sensing of Environment 158: 15–27.

Asner, Gregory P, Roberta E Martin, Loreli Carranza-Jiménez, Felipe Sinca, Raul Tupayachi,

Christopher B Anderson, and Paola Martinez. 2014. "Functional and Biological Diversity of

Foliar Spectra in Tree Canopies Throughout the Andes to Amazon Region." *New Phytologist*

204 (1): 127–39.

Badger, M. R., & Price, G. D. (1994). The role of carbonic anhydrase in photosynthesis. *Annual*

review of plant physiology and plant molecular biology, *45*(1), 369-392.

Bahulikar, Rahul A, Dominic Stanculescu, Catherine A Preston, and Ian T Baldwin. 2004. "ISSR

and AFLP Analysis of the Temporal and Spatial Population Structure of the Post-Fire Annual,

Nicotiana Attenuata, in SW Utah." *BMC Ecology* 4 (1): 1–13.

Bandillo, Nonoy, Chitra Raghavan, Pauline Andrea Muyco, Ma Anna Lynn Sevilla, Irish T

Lobina, Christine Jade Dilla-Ermita, Chih-Wei Tung, et al. 2013. "Multi-Parent Advanced

Generation Inter-Cross (MAGIC) Populations in Rice: Progress and Potential for Genetics

Research and Breeding." *Rice* 6 (1): 1–15.

Benjamini, Yoav, and Yosef Hochberg. 1995. "Controlling the False Discovery Rate: A Practical

and Powerful Approach to Multiple Testing." *Journal of the Royal Statistical Society: Series B*

(Methodological) 57 (1): 289–300.

Brieuc, Marine S. O., Charles D. Waters, Daniel P. Drinan, and Kerry A. Naish. 2018. "A

Practical Introduction to Random Forest for Genetic Association Studies in Ecology and

Evolution." *Molecular Ecology Resources* 18 (4): 755–66. Cavender-Bares, Jeannine, Jose

Eduardo Meireles, John J Couture, Matthew A Kaproth, Clayton C Kingdon, Aditya Singh,

Shawn P Serbin, et al. 2016. "Associations of Leaf Spectra with Genetic and Phylogenetic

Variation in Oaks: Prospects for Remote Detection of Biodiversity." *Remote Sensing* 8 (3): 221.

Claes, Peter, Jasmien Roosenboom, Julie D White, Tomek Swigut, Dzemila Sero, Jiarui Li,

Myoung Keun Lee, et al. 2018. "Genome-Wide Mapping of Global-to-Local Genetic Effects on

Human Facial Shape." *Nature Genetics* 50 (3): 414–23.

Czyż, Ewa A, Carla Guillén Escribà, Hendrik Wulf, Andrew Tedder, Meredith C Schuman,

Fabian D Schneider, and Michael E Schaepman. 2020. "Intraspecific Genetic Variation of a

Fagus Sylvatica Population in a Temperate Forest Derived from Airborne Imaging Spectroscopy

Time Series." *Ecology and Evolution* 10 (14): 7419–30.

Czyż, Ewa A, Bernhard Schmid, Andreas Hueni, Maarten B Eppinga, Meredith C Schuman,

Fabian D Schneider, Carla Guillén-Escribà, and Michael E Schaepman. 2023. "Genetic

Constraints on Temporal Variation of Airborne Reflectance Spectra and Their Uncertainties over

a Temperate Forest." *Remote Sensing of Environment* 284: 113338.

Fabre, N., Reiter, I. M., BECUWE-LINKA, N. O. E. L. L. E., Genty, B., & Rumeau, D. (2007).

Characterization and expression analysis of genes encoding α and β carbonic anhydrases in

Arabidopsis. *Plant, cell & environment*, *30*(5), 617-629.

Dechant, Benjamin, Youngryel Ryu, Grayson Badgley, Yelu Zeng, Joseph A Berry, Yongguang

Zhang, Yves Goulas, et al. 2020. "Canopy Structure Explains the Relationship Between

Photosynthesis and Sun-Induced Chlorophyll Fluorescence in Crops." *Remote Sensing of*

Environment 241: 111733.

DeLancey, Evan R, John F Simms, Masoud Mahdianpari, Brian Brisco, Craig Mahoney, and

Jahan Kariyeva. 2019. "Comparing Deep Learning and Shallow Learning for Large-Scale

Wetland Classification in Alberta, Canada." *Remote Sensing* 12 (1): 2.

Fields, Stanley, and Ok-kyu Song. 1989. "A Novel Genetic System to Detect Protein–Protein

Interactions." *Nature* 340 (6230): 245–46.

Furbank, R. T., & Tester, M. (2011). Phenomics–technologies to relieve the phenotyping

bottleneck. *Trends in plant science*, *16*(12), 635-644.

Gamon, John A, K Fred Huemmrich, Christopher YS Wong, Ingo Ensminger, Steven Garrity,

David Y Hollinger, Asko Noormets, and Josep Peñuelas. 2016. "A Remotely Sensed Pigment

Index Reveals Photosynthetic Phenology in Evergreen Conifers." *Proceedings of the National*

Academy of Sciences 113 (46): 13087–92.

Gao, Bo-Cai. 1996. "NDWI—a Normalized Difference Water Index for Remote Sensing of

Vegetation Liquid Water from Space." *Remote Sensing of Environment* 58 (3): 257–66.

Gitelson, Anatoly A, Yuri Gritz, and Mark N Merzlyak. 2003. "Relationships Between Leaf

Chlorophyll Content and Spectral Reflectance and Algorithms for Non-Destructive Chlorophyll

Assessment in Higher Plant Leaves." *Journal of Plant Physiology* 160 (3): 271–82.

Gitelson, Anatoly A, Mark N Merzlyak, and Olga B Chivkunova. 2001. "Optical Properties and

Nondestructive Estimation of Anthocyanin Content in Plant Leaves¶." *Photochemistry and*

Photobiology 74 (1): 38–45.

Gitelson, Anatoly A, Andrés Viña, Verónica Ciganda, Donald C Rundquist, and Timothy J

Arkebauer. 2005. "Remote Estimation of Canopy Chlorophyll Content in Crops." *Geophysical*

Research Letters 32 (8).

Glawe, Grit A, Jorge A Zavala, André Kessler, Nicole M Van Dam, and Ian T Baldwin. 2003.

"Ecological Costs and Benefits Correlated with Trypsin Protease Inhibitor Production in

Nicotiana Attenuata." *Ecology* 84 (1): 79–90.

Grabska, Ewa, Patrick Hostert, Dirk Pflugmacher, and Katarzyna Ostapowicz. 2019. "Forest

Stand Species Mapping Using the Sentinel-2 Time Series." *Remote Sensing* 11 (10): 1197.

Grzybowski, M., Wijewardane, N. K., Atefi, A., Ge, Y., & Schnable, J. C. (2021). Hyperspectral

reflectance-based phenotyping for quantitative genetics in crops: Progress and challenges. *Plant*

Communications, *2*(4).

Gu, Yingxin, Jesslyn F Brown, James P Verdin, and Brian Wardlow. 2007. "A Five-Year

Analysis of MODIS NDVI and NDWI for Grassland Drought Assessment over the Central Great

Plains of the United States." *Geophysical Research Letters* 34 (6).

Hatch, M. D., & Burnell, J. N. (1990). Carbonic anhydrase activity in leaves and its role in the

first step of C4 photosynthesis. *Plant Physiology*, *93*(2), 825-828.

Helfenstein, Isabelle S., Fabian D. Schneider, Michael E. Schaepman, and Felix Morsdorf. 2022.

"Assessing Biodiversity from Space: Impact of Spatial and Spectral Resolution on Trait-Based

Functional Diversity." Remote Sensing of Environment 275: 113024. Elsevier.

Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., ... &

Whitlock, M. C. (2016). Finding the genomic basis of local adaptation: pitfalls, practical

solutions, and future directions. *The American Naturalist*, *188*(4), 379-397.

Huang, Meng, Xiaolei Liu, Yao Zhou, Ryan M Summers, and Zhiwu Zhang. 2019. "BLINK: A

Package for the Next Level of Genome-Wide Association Studies with Both Individuals and

Markers in the Millions." *Gigascience* 8 (2): giy154.

Huete, Alfredo R. 2012. "Vegetation Indices, Remote Sensing and Forest Monitoring."

Geography Compass 6 (9): 513–32.

Huynh, Bao-Lam, Jeffrey D Ehlers, Bevan Emma Huang, María Muñoz-Amatriaín, Stefano

Lonardi, Jansen RP Santos, Arsenio Ndeve, et al. 2018. "A Multi-Parent Advanced Generation

Inter-Cross (MAGIC) Population for Genetic Analysis and Improvement of Cowpea (Vigna

Unguiculata l. Walp.)." *The Plant Journal* 93 (6): 1129–42.

INH, SƠN TRƯỜNG, Ivan Galis, and Ian T Baldwin. 2013. "UVB Radiation and 17-

Hydroxygeranyllinalool Diterpene Glycosides Provide Durable Resistance Against Mirid

(Tupiocoris Notatus) Attack in Field-Grown Nicotiana Attenuata Plants." *Plant, Cell &*

Environment 36 (3): 590–606.

Jacquemoud, Stéphane, and Susan Ustin. 2019. *Leaf Optical Properties*. Cambridge University

Press.

Karthikeyan, L, Ila Chawla, and Ashok K Mishra. 2020. "A Review of Remote Sensing

Applications in Agriculture for Food Security: Crop Growth and Yield, Irrigation, and Crop

Losses." *Journal of Hydrology* 586: 124905.

Kim, Sang-Gyu, Felipe Yon, Emmanuel Gaquerel, Jyotasana Gulati, and Ian T Baldwin. 2011.

"Tissue Specific Diurnal Rhythms of Metabolites and Their Regulation During Herbivore Attack

in a Native Tobacco, Nicotiana Attenuata." *PloS One* 6 (10): e26214.

Knipling, Edward B. 1970. "Physical and Physiological Basis for the Reflectance of Visible and

Near-Infrared Radiation from Vegetation." *Remote Sensing of Environment* 1 (3): 155–59.

Li, Cheng, Ewa A Czyz, Rayko Halitschke, Ian T Baldwin, Michael E Schaepman, and Meredith

C Schuman. 2023. "Evaluating Potential of Leaf Reflectance Spectra to Monitor Plant Genetic

Variation."

Li, Cheng, Ewa A Czyż, Rayko Halitschke, Ian T Baldwin, Michael E Schaepman, and Meredith

C Schuman. 2023. "UZH SG Nicotiana Attenuata ASD Data." SPECCHIO.

http://sc22.geo.uzh.ch:8080/SPECCHIO_Web_Interface/search.

Li, Dapeng, Sven Heiling, Ian T Baldwin, and Emmanuel Gaquerel. 2016. "Illuminating a

Plant's Tissue-Specific Metabolic Diversity Using Computational Metabolomics and

Information Theory." *Proceedings of the National Academy of Sciences* 113 (47): E7610–18.

Li, Xu, Qin Wang, Xuhong Yu, Hongtao Liu, Huan Yang, Chenxi Zhao, Xuanming Liu, et al.

2011. "Arabidopsis Cryptochrome 2 (CRY2) Functions by the Photoactivation Mechanism

Distinct from the Tryptophan (Trp) Triad-Dependent Photoreduction." *Proceedings of the*

National Academy of Sciences 108 (51): 20844–49.

Libbrecht, Maxwell W, and William Stafford Noble. 2015. "Machine Learning Applications in

Genetics and Genomics." *Nature Reviews Genetics* 16 (6): 321–32.

Liu, Nannan, Jianxiao Liu, Wenqiang Li, Qingchun Pan, Jie Liu, Xiaohong Yang, Jianbing Yan,

and Yingjie Xiao. 2018. "Intraspecific Variation of Residual Heterozygosity and Its Utility for

Quantitative Genetic Studies in Maize." *BMC Plant Biology* 18: 1–15.

Liu, Xiaolei, Meng Huang, Bin Fan, Edward S Buckler, and Zhiwu Zhang. 2016. "Iterative

Usage of Fixed and Random Effect Models for Powerful and Efficient Genome-Wide

Association Studies." *PLoS Genetics* 12 (2): e1005767.

Madritch, Michael D, Clayton C Kingdon, Aditya Singh, Karen E Mock, Richard L Lindroth,

and Philip A Townsend. 2014. "Imaging Spectroscopy Links Aspen Genotype with Below-

Ground Processes at Landscape Scales." *Philosophical Transactions of the Royal Society B:*

Biological Sciences 369 (1643): 20130194.

Meireles, Jose Eduardo, Jeannine Cavender-Bares, Philip A. Townsend, Susan Ustin, John A.

Gamon, Anna K. Schweiger, Michael E. Schaepman et al. "Leaf reflectance spectra capture the

evolutionary history of seed plants." New Phytologist 228, no. 2 (2020): 485-493.

Meireles, Jose Eduardo, Anna K Schweiger, and Jeannine M Cavender-Bares. 2017. "Spectrolab:

Class and Methods for Hyperspectral Data. R Package Version 0.0. 2."

Miller, JR, MD Steven, and TH Demetriades-Shah. 1992. "Reflection of Layered Bean Leaves

over Different Soil Backgrounds: Measured and Simulated Spectra." *International Journal of*

Remote Sensing 13 (17): 3273–86.

Nyholt, Dale R. 2004. "A Simple Correction for Multiple Testing for Single-Nucleotide

Polymorphisms in Linkage Disequilibrium with Each Other." *The American Journal of Human*

Genetics 74 (4): 765–69.

Price, Alkes L, Nick J Patterson, Robert M Plenge, Michael E Weinblatt, Nancy A Shadick, and

David Reich. 2006. "Principal Components Analysis Corrects for Stratification in Genome-Wide

Association Studies." *Nature Genetics* 38 (8): 904–9.

Qian, X, L Liu, H Croft, and J Chen. 2021. "Relationship Between Leaf Maximum

Carboxylation Rate and Chlorophyll Content Preserved Across 13 Species." *Journal of*

Geophysical Research: Biogeosciences 126 (2): e2020JG006076.

R Core Team. 2023. *R: A Language and Environment for Statistical Computing*. Vienna,

Austria: R Foundation for Statistical Computing. *https://www.R-project.org/*.

Ray, R. et al. 2022. "Whole Genome Sequence of MAGIC Parents and RIL Population." NCBI

Sequence Read Archive. *https://www.ncbi.nlm.nih.gov/bioproject/PRJNA907539*.

Ray, Rishav, Rayko Halitschke, Klaus Gase, Sabrina M Leddy, Meredith C Schuman, Nathalie

Rodde, and Ian T Baldwin. 2023. "A Persistent Major Mutation in Canonical Jasmonate

Signaling Is Embedded in an Herbivory-Elicited Gene Network." *Proceedings of the National*

Academy of Sciences 120 (35): e2308500120.

Ray, Rishav, Dapeng Li, Rayko Halitschke, and Ian T Baldwin. 2019. "Using Natural Variation

to Achieve a Whole-Plant Functional Understanding of the Responses Mediated by Jasmonate

Signaling." *The Plant Journal* 99 (3): 414–25.

Rudenko, N. N., Ignatova, L. K., Fedorchuk, T. P., & Ivanov, B. N. (2015). Carbonic anhydrases

in photosynthetic cells of higher plants. *Biochemistry (Moscow)*, *80*(6), 674-687.

Saha, Saswati, Laurent Perrin, Laurence Röder, Christine Brun, and Lionel Spinelli. 2022. "Epi-

MEIF: Detecting Higher Order Epistatic Interactions for Complex Traits Using Mixed Effect

Conditional Inference Forests." *Nucleic Acids Research* 50 (19): e114.

Sannemann, Wiebke, Bevan Emma Huang, Boby Mathew, and Jens Léon. 2015. "Multi-Parent

Advanced Generation Inter-Cross in Barley: High-Resolution Quantitative Trait Locus Mapping

for Flowering Time as a Proof of Concept." *Molecular Breeding* 35: 1–16.

Santure, A. W., & Garant, D. (2018). Wild GWAS—association mapping in natural populations.

Molecular ecology resources, *18*(4), 729-738.

Schena, Mark, Dari Shalon, Ronald W Davis, and Patrick O Brown. 1995. "Quantitative

Monitoring of Gene Expression Patterns with a Complementary DNA Microarray." *Science* 270

(5235): 467–70.

Serbin, Shawn P, Aditya Singh, Brenden E McNeil, Clayton C Kingdon, and Philip A

Townsend. 2014. "Spectroscopic Determination of Leaf Morphological and Biochemical Traits

for Northern Temperate and Boreal Tree Species." *Ecological Applications* 24 (7): 1651–69.

Sero, Dzemila, Arslan Zaidi, Jiarui Li, Julie D White, Tomás B González Zarzar, Mary L

Marazita, Seth M Weinberg, et al. 2019. "Facial Recognition from DNA Using Face-to-DNA

Classifiers." *Nature Communications* 10 (1): 2557.

Solomon, Mark J, Pamela L Larsen, and Alexander Varshavsky. 1988. "Mapping proteinDNA

Interactions in Vivo with Formaldehyde: Evidence That Histone H4 Is Retained on a Highly

Transcribed Gene." *Cell* 53 (6): 937–47.

Stasinski, Lance, Dawson M White, Peter R Nelson, Richard H Ree, and José Eduardo Meireles.

2021. "Reading Light: Leaf Spectra Capture Fine-Scale Diversity of Closely Related,

Hybridizing Arctic Shrubs." *New Phytologist* 232 (6): 2283–94.

Supuran, C. T. (2004). Carbonic anhydrases: catalytic and inhibition mechanisms, distribution

and physiological roles. In *Carbonic anhydrase* (pp. 13-36). CRC press.

Taiz, Lincoln, Eduardo Zeiger, Ian Max Møller, Angus Murphy, et al. 2015. *Plant Physiology*

and Development. Ed. 6. Sinauer Associates Incorporated.

Thenkabail, Prasad S, John G Lyon, and Alfredo Huete. 2018. "Advances in Hyperspectral

Remote Sensing of Vegetation and Agricultural Crops." In *Fundamentals, Sensor Systems,*

Spectral Libraries, and Data Mining for Vegetation, 3–37. CRC press.

Tiwari, A., Mamedov, F., Grieco, M., Suorsa, M., Jajoo, A., Styring, S., ... & Aro, E. M. (2016).

Photodamage of iron–sulphur clusters in photosystem I induces non-photochemical energy

dissipation. *Nature Plants*, *2*(4), 1-9.

Verrelst, Jochem, Zbyněk Malenovskỳ, Christiaan Van der Tol, Gustau Camps-Valls, Jean-

Philippe Gastellu-Etchegorry, Philip Lewis, Peter North, and Jose Moreno. 2019. "Quantifying

Vegetation Biophysical Variables from Imaging Spectroscopy Data: A Review on Retrieval

Methods." *Surveys in Geophysics* 40: 589–629.

Visscher, Peter M, Naomi R Wray, Qian Zhang, Pamela Sklar, Mark I McCarthy, Matthew A

Brown, and Jian Yang. 2017. "10 Years of GWAS Discovery: Biology, Function, and

Translation." *The American Journal of Human Genetics* 101 (1): 5–22.

Wan, Liang, Zheng Tang, Jiafei Zhang, Shuobo Chen, Weijun Zhou, and Haiyan Cen. 2021.

"Upscaling from Leaf to Canopy: Improved Spectral Indices for Leaf Biochemical Traits

Estimation by Minimizing the Difference Between Leaf Adaxial and Abaxial Surfaces." *Field*

Crops Research 274: 108330.

Wang, Jiabo, and Zhiwu Zhang. 2021. "GAPIT Version 3: Boosting Power and Accuracy for

Genomic Association and Prediction." *Genomics, Proteomics & Bioinformatics* 19 (4): 629–40.

Wang, Ran, and John A Gamon. 2019. "Remote Sensing of Terrestrial Plant Biodiversity."

Remote Sensing of Environment 231: 111218.

Wang, Ran, John A Gamon, Anna K Schweiger, Jeannine Cavender-Bares, Philip A Townsend,

Arthur I Zygielbaum, and Shan Kothari. 2018. "Influence of Species Richness, Evenness, and

Composition on Optical Diversity: A Simulation Study." *Remote Sensing of Environment* 211:

218–28.

Wang, Yue, Wilson Goh, Limsoon Wong, Giovanni Montana, and the Alzheimer's Disease

Neuroimaging Initiative. 2013. "Random Forests on Hadoop for Genome-Wide Association

Studies of Multivariate Neuroimaging Phenotypes." *BMC Bioinformatics* 14 (16): S6.

Wang, Zhihui, Adam Chlus, Ryan Geygan, Zhiwei Ye, Ting Zheng, Aditya Singh, John J

Couture, Jeannine Cavender-Bares, Eric L Kruger, and Philip A Townsend. 2020. "Foliar

Functional Traits from Imaging Spectroscopy Across Biomes in Eastern North America." *New*

Phytologist 228 (2): 494–511.

Weigelt, Alexandra, Liesje Mommer, Karl Andraczek, Colleen M. Iversen, Joana Bergmann,

Helge Bruelheide, Grégoire T. Freschet, et al. 2023. "The Importance of Trait Selection in

Ecology." *Nature* 618 (7967): E29–30.

Wu, Di, Deepti S Karhade, Malvika Pillai, Min-Zhi Jiang, Le Huang, Gang Li, Hunyong Cho,

Jeff Roach, Yun Li, and Kimon Divaris. 2021. "Machine Learning and Deep Learning in

Genetics and Genomics." *Machine Learning in Dentistry*, 163–81.

Yu, Jianming, Gael Pressoir, William H Briggs, Irie Vroh Bi, Masanori Yamasaki, John F

Doebley, Michael D McMullen, et al. 2006. "A Unified Mixed-Model Method for Association

Mapping That Accounts for Multiple Levels of Relatedness." *Nature Genetics* 38 (2): 203–8.

Zhang, F, and G Zhou. 2019. "Estimation of Vegetation Water Content Using Hyperspectral

Vegetation Indices: A Comparison of Crop Water Indicators in Response to Water Stress

Treatments for Summer Maize." *BMC Ecology* 19: 1–12.

Zhang, Zhiwu, Elhan Ersoz, Chao-Qiang Lai, Rory J Todhunter, Hemant K Tiwari, Michael A

Gore, Peter J Bradbury, et al. 2010. "Mixed Linear Model Approach Adapted for Genome-Wide

Association Studies." *Nature Genetics* 42 (4): 355–60.

Zheng, J., Richardson, T. G., Millard, L. A., Hemani, G., Elsworth, B. L., Raistrick, C. A., ... &

Gaunt, T. R. (2018). PhenoSpD: an integrated toolkit for phenotypic correlation estimation and

multiple testing correction using GWAS summary statistics. *Gigascience*, *7*(8), giy090.

10 TABLES

Table 1. Six Spectral Indices Used in Indices-GWAS

Index	Formula	Example of Validated Systems	Citations
$NDWI$	$\dfrac{R_{865} - R_{1614}}{R_{865} + R_{1614}}$	Crops, forests, wetlands, grasslands	(Gao 1996; F. Zhang and Zhou 2019; Karthikeyan, Chawla, and Mishra 2020; Huete 2012; Adam, Mutanga, and Rugege 2010; Gu et al. 2007; Helfenstein et al., 2022)
CI_{re}	$\dfrac{R_{783}}{R_{704}} - 1$	Crops, trees and vines, wetlands	(Gitelson, Merzlyak, and Chivkunova 2001; Qian et al. 2021; Gitelson et al. 2005; Gitelson, Gritz, and Merzlyak 2003; DeLancey et al. 2019; Helfenstein et al., 2022)
CCI	$\dfrac{R_{560} - R_{664}}{R_{560} + R_{664}}$	Crops, trees, forests	(Gamon et al. 2016; Dechant et al. 2020; Grabska et al. 2019; Helfenstein et al., 2022)
$ARDSI\_C_{ab}$	$\dfrac{R_{750} - R_{730}}{R_{770} + R_{720}}$	Crops	(Wan et al. 2021)
$ARDSI\_C_w$	$\dfrac{R_{1360} - R_{1080}}{R_{1560} + R_{1240}}$	Crops	(Wan et al. 2021)
$ARDSI\_C_m$	$\dfrac{R_{2200} - R_{1640}}{R_{2240} + R_{1720}}$	Crops	(Wan et al. 2021)

782 Table 2. Spectral Ranges of Segments Used in GWAS with Potential Associated Features

Segment	Spectral range (nm)	Potentially associated physiological features*
5	518–614, 695–709	VIS–NIR: Chlorophyll a, chlorophyll b, carotenoids, anthocyanins; leaf structure
10	400–432	VIS: Chlorophyll a, chlorophyll b, carotenoids, anthocyanins
12	433–517	VIS: Chlorophyll a, chlorophyll b, carotenoids, anthocyanins
13	648–690	VIS: Chlorophyll a, chlorophyll b, carotenoids, anthocyanins
14	615–647, 691–694	VIS: Chlorophyll a, chlorophyll b, carotenoids, anthocyanins
19	1894–2000	SWIR: Water content
20	1880–1893, 2001–2096	SWIR: Non-pigment organic composition (e.g., lignin, cellulose, starch, protein, sugar, oil)
22	1400–1401, 1864–1879, 2097–2311	NIR–SWIR: Water content; leaf structure; non-pigment organic composition
23	2397–2500	SWIR: Water content
24	2312–2396	SWIR: Non-pigment organic composition
25	1412–1484	SWIR: Water content
26	1402–1411, 1485–1522	SWIR: Non-pigment organic composition
28	1359–1380	SWIR: Water content
29	1381–1385, 1590–1754	SWIR: Non-pigment organic composition
30	1386–1399, 1523–1589, 1755–1863	SWIR: Non-pigment organic composition
33	1311–1358	NIR: Water content
34	1144–1310	NIR: Leaf structure
36	710–733	NIR: Leaf structure
37	734–744, 946–1014, 1129–1143	NIR: Leaf structure
38	745–945, 1015–1128	NIR: Leaf structure

783 Note: The associated features are based on common findings (Jacquemoud and Ustin 2019) and

784 are intended to indicate physiological properties of leaves known to affect the given spectral

785 ranges. These interpretations are subject to various factors and should not be considered

786 definitive.

 11   **FIGURE CAPTIONS**

Figure 1. Approaches and toolbox of this study. Photo credit: Meredith C. Schuman.

Figure 2. Variance partitioning across the spectral range of 350 to 2500 nm. Each column

represents the cumulative percentage of variance for a given wavelength.

Figure 3. Spectra as phenotypes. (a) Leaf reflectance spectra for all 616 plants examined in this

study. The orange shaded region illustrates the range of adjusted spectra, and the blue shaded

region the range of raw spectra. The orange dotted line represents the coefficient of variation for

the entire sample set. (b) Distribution of the six spectral indices used in the Index-GWAS

approach. (c) Results from the Hierarchical Spectral Clustering with Parallel Analysis (HSC-

PA). A total of 38 segments, differentiated by color, emerged from the clustering. Regions left

blank (NA) indicate segments that retained one Principal Component (PC) in the prior level,

halting further clustering. (d) Presentation of the 20 segments, shaped as example spectra, each

maintaining one retained PC. Spanning the entire spectrum from 400 to 2500 nm, these segments

served as phenotypes in the HSC-PA-GWAS approach to pinpoint associations with genetic

variances.

Figure 4. Index-GWAS results. (a) Manhattan plot showing the $-log_{10}(p\_value)$ of SNP

associations for the six indices across various models. The grey horizontal line indicates a p-

value of 1×10^{-5}, which is used to select associations to generate figure (b). (b) Visualization of

associations that have p-values less than 1×10^{-5}. The x-axis represents the six indices, the y-

axis the corresponding associated SNPs. The gradient red color scale represents the

FDR_adjusted p-values. (c) QQ plot contrasting the observed versus the expected

$-log_{10}(p\_value)$ for the GWAS analysis on the $ARDSI\_C_w$ index. The diagonal line represents

the expected distribution under the null hypothesis. The different color schemes depict the

models used: red for GLM, yellow for MLM, green for FarmCPU, and blue for BLINK. (d)-(e)

Phenotypic distribution for two significant markers, chr1_438155938(d) or chr1_193505583(e),

in association with $ARDSI\_C_w$. 0 represents a homozygous allele from the reference, 1 a

heterozygous allele, and 2 a homozygous alternative allele.

Figure 5. SW-GWAS results. (a) Manhattan plot showing the $-log_{10}(p\_value)$ of SNP

associations with BLINK for wavelength 952 with the strongest association. (b) Visualization of

associations that have p-values less than 1×10^{-5}. The x-axis represents the spectral

wavelengths from 400 to 2500 nm, while the y-axis displays the corresponding associated SNPs.

The gradient red color scale represents the FDR_adjusted p-values. (c) QQ plot of the BLINK

model contrasting the observed versus the expected $-log_{10}(p\_value)$ for the GWAS analysis on

the wavelength 952 nm. The diagonal line represents the expected distribution under the null

hypothesis.

Figure 6. HSC-PA-GWAS results. (a) Detailed visualization of associations that surpass the

exploratory threshold. The X-axis enumerates the 20 segments, while the Y-axis displays the

corresponding associated SNPs. (b) An alternative representation of the associations from (a)

with the X-axis showcasing the spectral wavelengths, providing insights into the specific spectral

regions of the associations. (c)-(d) QQ plots highlighting the observed versus expected

$-log_{10}(p\_value)$ for GWAS analysis on Segment 37(c) or Segment 38 (d). (e)-(f) Phenotypic

distribution for the significantly associated marker (chr1_131409683) in association with

Segment 37(e) or Segment 38(f). 0 represents a homozygous allele from the reference, 1 a

heterozygous allele, and 2 a homozygous alternative allele.

  12   FIGURES

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.