1

Key determinants of the dual clamp/activator function of Complexin

Mazen Makke, Alejandro Pastor Ruiz, Antonio Yarzagaray, Surya Gaya, Michelle

Zimmer, Walentina Frisch, Dieter Bruns

Center for Integrative Physiology and Molecular Medicine, School of Medicine,
University of Saarland, 66424 Homburg/Saar, Germany

Running title: Complexin in exocytosis

Keywords: neurotransmitter release, complexin, synaptotagmin, membrane fusion,
exocytosis

Corresponding author:
Dieter Bruns, Ph.D.
Center for Integrative Physiology and Molecular Medicine
School of Medicine
Univ. of Saarland
Dept. of Physiology
Kirrbergerstreet 8
D-66424 Homburg/Saar
Tel: +49 6841 162 6495
Fax: +49 6841 162 6492
E-mail: dieter.bruns@uks.eu

Abstract

Complexin determines magnitude and kinetics of synchronized secretion, but the underlying molecular mechanisms remained unclear. Here, we show that the hydrophobic face of the amphipathic helix at the C-terminus of Complexin II (CpxII, amino acids 115-134) binds to fusion-promoting SNARE proteins, prevents premature secretion and allows vesicles to accumulate in a release-ready state. Specifically, we demonstrate that an unrelated amphipathic helix functionally substitutes for the CTD of CpxII and that amino acid substitutions on the hydrophobic side compromise the arrest of the prefusion intermediate. To facilitate synchronous vesicle fusion, the N-terminal domain (NTD) of CpxII (amino acids 1-27) specifically cooperates with synaptotagmin I, but not with synaptotagmin VII. Expression of CpxII rescues the slow release kinetics of the Ca^{2+}-binding mutant SytI R233Q, whereas the N-terminally truncated variant of CpxII further delays it. These results indicate that the CpxII NTD regulates mechanisms which are governed by the forward rate of Ca^{2+} binding to SytI. Overall, our results shed new light on key molecular properties of CpxII that hinder premature exocytosis and accelerate synchronous exocytosis.

Introduction

The accumulation of vesicles in a release ready state is an essential property to meet the speed requirements of synchronized SNARE mediated exocytosis (Wickner and Schekman, 2008; Sudhof and Rothman, 2009; Jahn and Fasshauer, 2012). One of the key questions is how the assembly of SNARE complexes is coordinately paused to allow rapid synchronous neurotransmitter release upon intracellular Ca^{2+} increase. Mechanistically, the required metastable pre-fusion intermediate could be maintained by the action of a SNARE-interacting protein acting as a transient fusion "clamp". However, the existence and identity of the putative "fusion clamp" factor has long been debated (for review see (Mohrmann et al., 2015; Trimbuch and Rosenmund, 2016; Brunger et al., 2019). Complexin (Cpx) is a small cytosolic SNARE regulatory protein that prevents premature and asynchronous vesicle fusion, and accelerates Ca^{2+}-triggered synchronized exocytosis (Xue et al., 2007; Dhara et al., 2014). Several studies have suggested an inhibitory effect of the accessory alpha-helix of CpxII, although various mechanisms have been proposed, including direct binding to SNAREs or other proteins (Giraudo et al., 2009; Lu et al., 2010; Yang et al., 2010; Krishnakumar et al., 2011; Kummel et al., 2011; Bykhovskaia et al., 2013; Cho et al., 2014; Malsam et al., 2020), electrostatic membrane interactions (Trimbuch et al., 2014), or stabilization of the secondary structure of the central helix (Radoff et al., 2014). The C-terminal region of Cpx also prevents spontaneous fusion in neurons (Cho et al., 2010; Martin et al., 2011; Kaeser-Woo et al., 2012) and premature secretion in neuroendocrine cells (Dhara et al., 2014). In a previous study, we presented evidence that the C-terminus of CpxII may compete with the SNAP-25 SN1 region for binding to SNARE partners, thus disrupting

the progressive SNARE complex formation prior to the actual Ca^{2+} stimulus (Makke et al.,

2018). Furthermore, unlike the accessory α-helix (Radoff et al., 2014), the CTD of CpxII

cannot be functionally replaced by an unrelated helical sequence (Makke et al., 2018),

suggesting that specific structural features or key residues within the CTD are required

for the clamping function of Cpx.

Independent of the clamping function of Cpx, knock-out and knock-down studies of Cpx

have shown, even in the absence of premature spontaneous fusion, a significant

reduction in evoked release, indicating a supporting role of Cpx in synchronous

neurotransmitter release (Reim et al., 2001; Xue et al., 2007; Cai et al., 2008; Maximov

et al., 2009; Strenzke et al., 2009; Xue et al., 2009; Cho et al., 2010; Hobson et al., 2011;

Martin et al., 2011; Jorquera et al., 2012; Lin et al., 2013; Yang et al., 2013; Dhara et al.,

2014; Kurokawa et al., 2015; Lopez-Murcia et al., 2019). Furthermore, there is increasing

evidence that the major fusion-promoting function of complexin in vertebrates is mediated

by its very N-terminus, an action that is mechanistically independent and even separable

from the clamping function of complexin (Dhara et al., 2014). Yet, no consensus has been

reached on the mechanism by which the N-terminus facilitates secretion, including

changes in the Ca^{2+}-affinity of the release machinery (Dhara et al., 2014), a stabilizing

role for SNARE proteins (Xue et al., 2010) and/or direct membrane binding (Lai et al.,

2016).

Using viral expression of CpxII or its mutants in chromaffin cells, we identify the

amphipathic character of the helical domain at the end of CpxII´s CTD as the critical

molecular determinant for the protein´s inhibitory function. Furthermore, our experiments

indicate that the cluster of glutamate residues upstream of the amphipathic α helix may

assist synaptotagmin I (SytI) in releasing Cpx´s clamp to trigger fast exocytosis.

Moreover, our results show that the NTD of CpxII specifically modulates SytI- but not

synaptotagmin VII (SytVII)-mediated exocytosis. In this way, the N-terminus of complexin

speeds up exocytosis triggering that is controlled by the forward rate of Ca^{2+} binding to

the calcium sensor SytI (Voets et al., 2001; Sorensen et al., 2003). Overall, our results

provide new insights into the molecular determinants required for the inhibitory function

of CpxII and elucidate the mechanisms facilitating fusion through the CpxII NTD.

Results

CpxII inhibits premature exocytosis via the amphipathic α helix of the CTD

CpxII is the only Cpx isoform expressed in chromaffin cells (Cai et al., 2008). To elucidate

the mode of action of CpxII in controlling Ca-triggered exocytosis, synchronous secretion

of isolated mouse chromaffin cells was recorded by membrane capacitance

measurements (CM) in response to photolytic uncaging of intracellular Ca^{2+} [Ca]i using

the caged Ca^{2+}-compound Nitrophyenly-EGTA. Changes in [Ca]i were monitored using a

combination of Ca^{2+}-indicators (fura-2 and furaptra). The results show that CpxII knock

out strongly reduced Ca^{2+} triggered synchronized vesicle fusion **(Figure 1B)**, which was

accompanied by a profound increase in asynchronous exocytosis during the loading

phase with NP-EGTA (free [Ca]i ~ 700 nM, **Figure 1E, F**) to allow for Ca^{2+}-dependent

priming of chromaffin granules. Expression of the wt protein (CpxII ko + CpxII) fully

restored the typical flash-evoked response with a prominent exocytotic burst (EB, **Figure**

1B) and prevented any granule loss by premature exocytosis (**Figure 1E, F**). Thus, CpxII

hinders premature vesicle exocytosis, that would otherwise outpace phasic secretion,

agreeing with our previous observations (Dhara et al., 2014; Makke et al., 2018). Both components of the EB, the readily releasable pool (RRP) and the slowly releasable pool (SRP) were similarly affected, without altering the sustained rate (SR) of secretion (**Figure 1C**). Furthermore, loss of CpxII conferred a slower time constant of the RRP response and a longer secretory delay (**Figure 1D**). Overall, CpxII not only hinders premature secretion but also accelerates exocytosis timing. The inhibitory function of CpxII resides within the last 34 amino acids of its CTD (Makke et al., 2018), which is characterized by an amphipathic helix and an upstream located cluster of glutamate residues (**Figure 1A**). Yet, the precise molecular mechanism of its clamp action remained unclear. Truncation of the amphipathic helix (CpxII 1-115) strongly reduced the RRP component of the exocytotic burst (**Figure 1B**) and elevated premature secretion (**Figure 1E, F**) when compared with the wt protein. Surprisingly, the CpxII 1-115 amp helix mutant, in which the last 19 amino acids were replaced with an unrelated amphipathic helix (**Figure 1A**), fully restored the flash-evoked response and the ability of CpxII to prevent premature secretion (**Figure 1B-F**). As the CpxII NTD determines the stimulus-secretion coupling (Dhara et al., 2014; Makke et al., 2018), both the RRP kinetics as well as the secretory delay was fully rescued by the mutant variants **(Figure 1D).** Therefore, the amphipathic helix of CpxII's CTD is a key determinant for the protein to prevent premature secretion.

The hydrophobic face of C-terminal amphipathic helix of CpxII is crucial for its inhibitory function

To investigate whether specific amino acids within the CpxII CTD amphipathic α helix are

essential for its inhibitory function, we introduced point mutations on either the

hydrophobic or hydrophilic face of the helix and tested their impact on regulated

exocytosis. Perturbation of the hydrophobic face of the CpxII CTD amphipathic helix by

replacing the two hydrophobic leucine amino acids L124 and L128 with charged

glutamate residues (CpxII L124E-L128E) mimicked the phenotype of the truncated CpxII

1-115 variant (**Figure 2A-C** compare with **Figure 1**). The CpxII L124E-L128E mutant only

partially restored the EB **(Figure 2B)** and largely failed to clamp premature vesicle

exocytosis **(Figure 2C).** In contrast, replacing the same leucine residues with hydrophobic

tryptophan amino acids (CpxII L124W-L128W) fully rescued the exocytotic burst size

(Figure 2E), and prevented premature exocytosis like wt CpxII **(Figure 2F)**. No changes

in the kinetics of synchronized exocytosis were detected for either mutant variant **(Figure**

2 figure supplement 1). In the same line, mutating L117 and L121 to either glutamate

(CpxII L117E-L121E) or tryptophan (CpxII L117W-L121W) confirmed the result that

preservation of the hydrophobic face of the amphipathic helix is a prerequisite for

arresting premature fusion **(Figure 2 figure supplement 2)**. We have previously pointed

out that the CpxII CTD shares a high degree of structural similarity with the C-terminal

half of the SNAP25-SN1 domain and showed that corresponding CpxII-SNAP25-SN1

chimeras (residues 44 to 77) fully restore the function in CpxII deficient cell (Makke et al.,

2018). In addition to the identical pattern of hydrophobic amino acids (heptad repeat), the

sequence comparison between CpxII CTD and SNAP-25-SN1 reveals similar or even

identical amino acids on the polar side of the amphipathic helix **(Figure 2 figure**

supplement 3A). As a first approach, we replaced these amino acids with alanine

residues (D118A, Q129A, D130A, K133A) to test their functional significance. However,

none of these point mutants showed an obvious phenotype, neither in the synchronous

nor in the asynchronous secretion response (**Figure 2 figure supplement 3**). Similarly,

the double point mutant D118K-D130K, with an intended charge reversal, had no serious

functional consequences compared to the response with the wild-type protein (**Figure 2**

figure supplement 4). These results agree with our observation that even an unrelated

amphipathic helix can functionally replace the CTD of CpxII. Collectively, they indicate

that preserving the hydrophobic face of the amphipathic helix at the end of the CTD of

CpxII is necessary to prevent premature exocytosis.

The ability of the CpxII CTD to rescue secretion parallels its interaction with

SNAREs and SytI

Since the CpxII CTD transiently obstructs the assembly of SDS-resistant SNARE

complexes (Makke et al., 2018), it is possible that the CpxII C-terminus with its SN1

mimetic region competes with the SN1 motif (membrane-proximal layers) of SNAP25 for

binding to its cognate SNARE partners, thereby arresting exocytosis. To probe whether

CpxII CTD and its mutant variants display altered SNARE interactions, immobilized GST-

CpxII CTD or its mutants were incubated with detergent extract from wt mouse brain.

Interaction partners were eluted by thrombin cleavage (STH-fraction). While no binding

to GST alone was observed, the GST-CpxII CTD pulled down Synaptotagmin I (SytI),

Syntaxin 1A (Syx1) and traces of SynaptobrevinII (SybII) **(Figure 3A)**. CpxII CTD-SNARE

interactions and binding to SytI were abolished for the "loss of clamp" mutant CpxII CTD

L124E-L128E (**Figure 3A**). In contrast, they were preserved or even enhanced for the

CpxII CTD L124W-L128W mutant protein with undiminished 'clamp' function.

Furthermore, the CpxII 1-115 amphipathic helix mutant efficiently co-precipitated SytI,

Syx1 and SybII **(Figure 3B)**. Accompanying immunofluorescence analyses reveal similar

expression levels for CpxII and its mutant variants **(Figure 3 figure supplement 1)** and

show that the latter colocalize with SybII like the wt protein, indicating unperturbed sorting

to chromaffin granules and lipid binding **(Figure 3 figure supplement 2)**. Taken together,

the inhibition of premature exocytosis by CpxII CTD and its mutants correlates with their

ability to interact with SytI and SNARE proteins, suggesting that these interactions are a

prerequisite for CpxII CTD to prevent premature exocytosis.

The cluster of glutamates in the CpxII CTD speeds up exocytosis timing

Further truncation of the C-terminus between amino acid position 100 to 115 exacerbates

the 'unclamping' phenotype (Makke et al., 2018), indicating that this region provides

structural elements that contribute to or stabilize the fusion-arresting function of the

amphipathic helix. The region upstream to the amphipathic helix is characterized by a

cluster of glutamate residues (aa 108-114) which is conserved across the animal kingdom

and, according to biochemical experiments, may serve as a putative SytI interaction site

(Tokumaru et al., 2008). However, the functional role of the glutamate cluster in regulated

exocytosis remains to be elucidated. To follow up on this, we replaced the glutamate

cluster with corresponding stretch of alanine residues (CpxII E-A). When expressed in

CpxII ko chromaffin cells, the mutant CpxII E-A **(Figure 4A)** only supported a reduced

synchronized exocytotic burst **(Figure 4B)** with a specifically reduced RRP **(Figure 4C)**.

In addition, the remaining EB exhibited slower fusion kinetics for both phases of the

synchronous exocytosis, the RRP and the SRP, and an increased stimulus-secretion

delay when compared to the secretion response of wt CpxII **(Figure 4D)**. Nevertheless,

premature exocytosis was effectively suppressed by the CpxII E-A mutant, as was

observed for the wt protein **(Figure 4E, F)**. As the CpxII E-A mutation has effects on the

timing of rapid exocytosis, it may interfere with the efficacy of SytI in triggering vesicle

fusion. Notably, no comparable slowing of exocytosis was observed when both protein

domains, the glutamate cluster and the amphipathic helix, were truncated, such as with

the CpxII 1-100 mutant (Makke et al., 2018). Thus, with an intact CTD of CpxII, the

glutamate cluster may be required as a contact site for SytI to release the molecular clamp

of the downstream amphipathic helix and thus to trigger the rapid secretion response.

The functional interplay of SytI and CpxII in Ca^{2+}-triggered exocytosis

To study the potential interplay of SytI and CpxII in triggering secretion, we comparatively

analyzed the phenotypic consequences of single and compound deficiencies of these

proteins. Consistent with previous work (Voets et al., 2001; Nagy et al., 2006; Dhara et

al., 2014), loss of SytI reduced the EB size, specifically by decreasing the RRP

component (**Figure 5A-C**), and prolonged the kinetics of SRP secretion as well as the

secretory delay when compared to the wt response (**Figure 5D**). In stark contrast,

additional loss of SytI in the absence of CpxII did not further aggravate the phenotype,

suggesting that all SytI functions are critically dependent on the presence of CpxII.

Furthermore, SytI deficiency did not alter premature secretion when compared to the

response of wt cells or with the elevated secretion in the absence of CpxII (**Figure 5E, F**).

Collectively, these results demonstrate a clear dependence of SytI functions on CpxII for

the synchronous secretion response. At submicromolar [Ca]i, changes in the CpxII

expression systematically altered the amount of premature secretion, which inversely

correlated with the EB size (**Figure 5 figure supplement 1**). Thus, CpxII inhibits

premature secretion that would otherwise outpace synchronous release. A similar

relationship between tonic secretion and exocytotic burst size was found in the absence

of SytI, indicating that CpxII hinders premature secretion in a SytI-independent manner

(**Figure 5 figure supplement 1**). Genetic loss of SytVII also reduced the exocytotic burst

size (**Figure 5G, H**), agreeing with previous work (Schonn et al., 2008). Both components

of the EB, the RRP and the SRP, were similarly affected (**Figure 5H**). The RRP kinetics

were found to be slightly faster when compared with controls (**Figure 5I (inset), J**).

Strikingly, deletion of SytVII in the CpxII KO background almost completely eliminated the

EB, leaving only a sustained phase of secretion behind (**Figure 5G-I**). The phenotype of

the CpxII-SytVII dko is remarkably similar to the compound deficiency of SytI and SytVII

(Schonn et al., 2008), corroborating the view that CpxII serves as a gatekeeper of SytI

function in the Ca^{2+}-triggered exocytosis of chromaffin granules. In the same line, CpxII

expression in SytVII ko cells enhances exocytosis like in wt cells (**Figure 5 figure**

supplement 2). In contrast to the loss of SytI, SytVII deficiency strongly reduced

premature secretion when compared with wt cells or on the background of CpxII

deficiency (**Figure 6K, L**). Thus, deletion of Syt VII profoundly alters the exocytotic rates

at submicromolar $[Ca^{2+}]$i, a notion that agrees with its proposed function as high-affinity

Ca^{2+}-sensor (MacDougall et al., 2018). Overall the combined set of data reveals a

differential interplay between the two Syt isoforms and CpxII. While function of SytI is

critically dependent on CpxII, SytVII appears to act independently, most likely in the Ca^{2+}-

dependent priming reaction (Tawfik et al., 2021).

Acceleration of synchronized exocytosis by the CpxII N-terminus relies on SytI

Phenotypic cues from many model systems as well as *in vitro* analyses point to a fusion-

promoting effect of complexin due to increased Ca^{2+} sensitivity, suggesting the possibility

of mechanistic crosstalk between complexin and the Ca^{2+} sensor synaptotagmin (Reim

et al., 2001; Tadokoro et al., 2005; Huntwork and Littleton, 2007; Xue et al., 2007; Diao

et al., 2012; Malsam et al., 2012; Dhara et al., 2014). Indeed, expression of an N-

terminally truncated complexin variant (e.g. residues 28–134) in cplxII-deficient

chromaffin cells failed to restore normal release rates (Dhara et al., 2014) and also

reduced the EPSC amplitude of hippocampal neurons in response to the short-lasting

action potential evoked Ca^{2+}-influx (Xue et al., 2009; Xue et al., 2010). Given the different

apparent calcium binding rates of SytI and SytVII (Sugita et al., 2002; Pinheiro et al.,

2016), one could speculate that the NTD of Cpx accelerates the kinetics of vesicle fusion

by preferentially recruiting SytI as a calcium sensor for rapid exocytosis instead of SytVII.

To test this hypothesis, we overexpressed the CpxII ∆N mutant in chromaffin cells lacking

either the calcium sensor SytI or SytVII. Overexpression of either wt CpxII or the mutant

CpxII ∆N had no effect on the kinetics of exocytosis in cells lacking SytI **(Figure 6A-C)**,

which demonstrates that CpxII requires SytI to accelerate exocytosis. In contrast, in SytVII

ko cells, CpxII ∆N slowed down synchronous secretion and increased the secretory delay

(Figure 6D-F) as previously reported for overexpression in wt cells (Dhara et al., 2014).

Furthermore, both, CpxII and CpxII ∆N, hindered premature exocytosis independent of

the present Syt isoform, corroborating the view that CpxII hinders SNARE action rather

than blocking one of the main Ca^{2+}-sensors in chromaffin cells **(Figure 6E, F and K, L)**.

Together, these findings support a scenario in which CpxII NTD and SytI are functionally

interdependent in the activation of fast synchronous exocytosis. Because the CpxII ΔN

mutant slows down secretion in SytVII ko-cells as in wt cells (Dhara et al., 2014), the CpxII

NTD directly regulates SytI rather than acting as a molecular switch between these Syt

isoforms.

CpxII NTD accelerates exocytosis timing by enhancing the apparent Ca^{2+} affinity of

the release machinery

Since the promotive role of the NTD of CpxII can only be tested in the presence of SytI,

we took advantage of the knock in SytI R233Q mutant, which carries the substitution

mutation in the calcium binding pocket of the C2A domain and lowers the affinity for Ca^{2+}-

binding (Fernandez-Chacon et al., 2001). Intriguingly, the phenotype of the R233Q ki

mutant in chromaffin cells is kinetically similar to that of the CpxII ΔN mutant, as it also

slows down timing of exocytosis compared to wt cells (Sorensen et al., 2003; Dhara et

al., 2014). Consistent with previous work by Soerensen and colleagues (Sorensen et al.,

2003), we found that the SytI R233Q mutation slowed down the rate of exocytosis,

prolonged the secretory delay and enhanced the EB size when compared to wt cells

(Figure 7A-D). Overexpression of the CpxII ΔN mutant in SytI R233Q ki cells, which is

expected to outcompete the function of endogenous CpxII (Dhara et al., 2014), further

slowed down the rate of synchronized release and restored the EB size to the wt level

(Figure 7C, D). Remarkably, overexpression of the wt CpxII significantly accelerated the

kinetics of synchronous exocytosis in SytI R233Q ki cells, restoring them to the level of

wt cells. Furthermore, CpxII wt and its mutant showed an undiminished suppression of

premature exocytosis in the SytI R233Q ki cells (**Figure 7E, F**). Thus, CpxII with its NTD

is able to compensate for the functional deficiencies of the SytI R233Q mutation. Given

that exocytosis timing in chromaffin cells is largely determined by the kinetics of Ca^{2+}-

binding to SytI (Voets et al., 2001; Sorensen et al., 2003), the opposite effects of the wt

protein and its ∆N mutant on the kinetics of secretion indicate that the N-terminal domain

of CpxII accelerates mechanisms regulated by the Ca^{2+} binding kinetics of SytI.

CpxII NTD supports the forward rate of calcium binding to SytI in accelerating

exocytosis

To obtain release kinetics over a wider range of [Ca]i, we made use of a calcium-ramp

protocol, in which calcium was progressively released from the calcium cage by

alternating 340 and 380 nm illumination allowing us to combine slow calcium uncaging

with ratiometric measurements of [Ca]i (Sorensen et al., 2003). Simultaneous CM

measurements revealed a sigmoid-shaped capacitance increase during the calcium ramp

protocol indicating progressive depletion of the primed vesicle pool and thus providing an

estimate of the exocytotic burst size (**Figure 8A**). To measure fusion kinetics, the slope

of the capacitance increase, the remaining pool, and the corresponding exocytosis rates

were determined during the Ca^{2+}-ramp **(Figure 8B)**. Overexpression of CpxII wt protein

in SytI R233Q ki cells clearly shifted the half-maximal concentration for pool depletion

(P50) to lower [Ca]i, whereas overexpression of the CpxII ∆N mutant increased it **(Figure**

8C). Evidently, wt CpxII rescued the secretion deficit of the Syt R233Q mutant, as

observed in the flash experiment **(Figure 7)**. For the SytI R233Q ki cells, fusion rates

increase as a function of [Ca]i and are shifted with similar slope to either lower [Ca^{2+}]i

(SytI R233Q + CpxII) or higher [Ca]i (SytI R233Q + CpxII ΔN), indicating unchanged

cooperativity of Ca^{2+} binding **(Figure 8D)**. The fusion rate-[Ca]i relationships are

reasonably well approximated by Hill equations **(Figure 8D, dashed lines)** with similar

coefficients (n), but different KD-values (SytI R233Q + CpxII, n=2.3, KD=17.4 µM; SytI

R233Q, n=2.03, KD=38.2 µM; SytI R233Q + CpxII ΔN, n=2.07, KD=58.13 µM). It should

be noted that the rate of exocytosis in the ramp experiment could only be reliably

determined at [Ca]i levels below 3, 4, and 6 µM for SytI R233Q + CpxII, SytI R233Q, and

SytI R233Q + CpxII ΔN cells, respectively, so that the accuracy of the measurement is

not affected by excessive depletion of the pool of primed vesicles. As the Ca^{2+}-sensor is

expected to be in a "quasi steady-state" during the slow Ca^{2+}-ramp, it is not possible to

disentangle whether the observed changes in Ca^{2+}-affinity are due to increased forward

or decreased backward rates in Ca^{2+}-binding to SytI. For the flash-evoked response,

instead, the fusion rate is dominated by the forward rate in Ca^{2+}-binding. As similar

quantitative differences in exocytosis timing were observed under these conditions, the

results indicate that the NTD of CpxII and the forward rate of Ca^{2+}-binding to SytI act

synergistically to accelerate exocytosis timing.

Discussion

In many secretory systems, Complexin plays a dual role in the regulation of SNARE-

mediated vesicle fusion. On the one hand, Cpx inhibits asynchronous exocytosis, thereby

enhancing Ca^{2+}-triggered synchronized secretion, and on the other hand, it accelerates

fusion of primed vesicles upon elevation of intracellular Ca^{2+}. Here, we have investigated

the molecular determinants required for the CTD of CpxII to clamp premature vesicle

exocytosis and delineated molecular mechanisms by which CpxII's N-terminal domain

accelerates exocytosis timing. Our findings show that the hydrophobic character of the

amphipathic helix at the very C terminus of CpxII is crucial for inhibiting premature vesicle

fusion. We also provide evidence that the group of glutamate residues within the CTD of

CpxII accelerates exocytosis by lifting the molecular clamp of the downstream

amphipathic α helix, most likely through interactions with SytI. Furthermore, we show that

CpxII cooperates exclusively with SytI and modulates the apparent Ca^{2+} affinity of

secretion by regulating processes controlled by the forward rate of Ca^{2+} binding to the

Ca^{2+} sensor.

364   The clamping action of CpxII in Ca^{2+}-triggered secretion

The ability to build up a pool of primed vesicles is a central property of many secretory

cells. To ensure adequate stimulus-secretion coupling and to prevent premature

exocytosis of these vesicles, a molecular mechanism must exist that arrests these

vesicles in their docked state. Recently, we provided the first evidence that the CTD of

CpxII (amino acids 101-134) plays a central role in the molecular clamp of the secretion

machinery of chromaffin cells (Makke et al., 2018). Thus, it became clear that the C-

terminal domain (CTD) of CpxII is essential and rate-limiting to prevent premature fusion

and to build up a pool of primed vesicles. Structural similarities between the CTD of CpxII

and the C-terminal half of the SNAP25-SN1 domain, as well as the observation that

CpxII:SNAP25-SN1 chimeras (C-terminal half) fully rescued function in CpxII knock-out

cells, suggested that the C-terminus of CpxII may compete with SNAP25-SN1 for binding

to the SNARE complex, thereby halting the progressive assembly of the SNARE complex

prior to the triggering Ca^{2+} stimulus. However, it remained unclear which structural

determinants of the CTD of CpxII mediate its actual inhibitory potential. Our new

experiments show that substitution of the CTD with an unrelated amphipathic helix fully

restores the function of the CpxII protein (**Figure 1**), suggesting that amphipathicity and

most likely preservation of the hydrophobic face is crucial for the inhibitory effect of the

CTD. In good agreement, substitution of single hydrophobic leucine residues with

charged glutamate residues almost completely abolished the inhibitory phenotype of CTD

(Figure 2), whereas replacement with equally hydrophobic tryptophan residues was

functionally tolerated. Furthermore, replacement of structurally similar amino acids

between the CTD of CpxII and the SNAP25-SN1 domain on the polar side of the

amphipathic helix had no functional consequences **(Figure 2 figure supplement 2)**.

Thus, an unperturbed hydrophobic face of the amphipathic helix is essential for the fusion-

inhibiting effect of the CpxII CTD. In particular, experiments on the *C. elegans* NMJ

suggested that membrane binding of the CpxII CTD via its amphipathic helix sequesters

the protein to synaptic vesicles, thereby, concentrating other SNARE-binding regions of

the protein (e.g. the accessory helix at the site of exocytosis for efficient molecular

clamping (Wragg et al., 2013; Snead et al., 2014; Gong et al., 2016). However, other

structure-function analyses with Cpx from C. elegans have shown that membrane binding

is important but does not suffice for the inhibitory effect of Cpx (Snead et al., 2017; Wragg

et al., 2017). Our biochemical pull-down experiments firstly show, that the isolated CTD

of CpxII binds to SNARE proteins (i.e. syntaxin, SybII) and SytI **(Figure 3)**. Second, they

demonstrate that the differential restoration of exocytosis by CpxII and its mutant variants

mirrors its altered binding to SNAREs and SytI. Furthermore, compared to the wild-type

protein, none of these CTD mutants exhibited defects in binding to vesicular membranes

(Figure 3 figure supplement 2). Thus, mutants with unaltered membrane binding

showed a profound deficit in the prevention of premature exocytosis and the formation of

a primed vesicle pool rendering the presumed necessary concentration of CpxII on

secretory organelles as a genuine mechanism for the inhibitory action of the CTD unlikely.

Evidently, protein-protein rather than protein-membrane interactions of the CpxII CTD

determine the mechanism of molecular clamping in chromaffin cells. Nevertheless, our

findings confirm membrane binding by the CpxII CTD and do not exclude additional

membrane sculpting activity of Cpx, as observed by nanodisc-black lipid membrane

electrophysiology (Courtney et al., 2022). As Cpx binds with much higher affinity to binary

acceptor complexes than to membranes (Zdanowicz et al., 2017), the Cpx CTD may

switch from a membrane-bound to a protein-bound state in the course of vesicle docking

to the plasma membrane. Furthermore, an unaltered fusion-inhibitory phenotype of the

CpxII-CTD can be observed in the absence of SytI **(Figure 5 figure supplement 1)** or

SytVII **(Figure 6)** and in the joint absence of both Ca^{2+} sensors in chromaffin cells (Dhara

et al., 2014) which counters alternative mechanisms such as a functional antagonism of

the Ca^{2+} sensor SytI or of secondary Ca^{2+} sensors like SytVII in chromaffin cells (cf. Yang

et al., 2010). Indeed, recent structural analyses of the formation of individual SNARE

complexes using optical tweezers confirm that the CTD of CpxII plays an active role in

the arrest of assembling SNARE complexes (Hao et al., 2023). We have previously

shown, that acute infusion of the isolated C-terminal peptide of CpxII into wt cells

significantly diminishes premature vesicle fusion, but fails to clamp tonic secretion in CpxII

ko cells, indicating that other domains of the Cpx protein are required to cooperate with

the CTD (Makke et al., 2018). As the accessory helix of Cpx has been found to bind to

membrane proximal cytoplasmic regions of SNAP-25 and SybII (Malsam et al., 2012;

Bykhovskaia et al., 2013; Vasin et al., 2016; Bera et al., 2022), an attractive scenario

could be that both domains of CpxII, the CTD and the accessory helix, synergistically

cooperate to stall final SNARE assembly. In summary, our studies provide new functional

insights into fundamental mechanisms of secretion control and support a model wherein

the CpxII CTD, with the hydrophobic face of its amphipathic helix, prevents untimely

fusion by arresting the progressive assembly of membrane-bridging SNARE complexes

prior to the actual Ca^{2+} stimulus.

Lifting the clamp

Locking of fusion-promoting SNARE complexes requires a fast-acting mechanism to

release the clamp in order to meet the speed requirements of rapid Ca^{2+}-driven

exocytosis. While recent models propose a mechanism by which concurrent binding of

SytI and Cpx to partially-assembled SNARE complexes defines the primed vesicle state,

no consensus has been reached, despite tremendous efforts, on the precise mechanism

by which Ca^{2+}-binding to the C2-domains of SytI triggers stimulus-secretion coupling (for

review see Brunger et al., 2018, Rizo, 2022). Previous biochemical experiments have

shown that SytI interacts with Cpx by binding to a polyacidic glutamate cluster

immediately upstream of the amphipathic helix (Tokumaru et al., 2008). Even though the

long acidic stretch is conserved across the animal kingdom (Lottermoser and Dittman,

2022), no functional significance has been ascribed to it. Our observations show that

substitution of the glutamate residues with a corresponding stretch of alanine residues

reduced the EB size and most remarkably slowed down exocytosis timing **(Figure 4)**. The

latter result indicates that Cpx's CTD cooperates with SytI in stimulus-secretion coupling

and suggests that SytI interaction with the polyacidic glutamate cluster is instrumental in

lifting the clamp imposed by Cpx's amphipathic helix. Notably, a comparable slowing in

exocytosis timing was not observed in the absence of the entire CTD of CpxII (i. e. by

deleting the amino acid positions 101-134, Makke et al., 2018), suggesting that this SytI-

CpxII interaction site is specifically required for lifting the clamp signal imposed by Cpx's

downstream amphipathic helix. While CpxII aligns with its central helix in an antiparallel

orientation on the SNARE complex, downstream protein regions of CpxII may provide

sufficient structural flexibility for the far C-terminus to fold back in a parallel orientation on

membrane-proximal layers of the partially assembled SNARE complex (Bowen et al.,

2005; Makke et al., 2018, Hao et al., 2023). Under these conditions, the glutamate cluster

of Cpx's CTD may come into close proximity to the position of crucial amino acids of

SNAP-25 (i.e. D51, E52, E55), which together with R398, R399 and neighboring basic

residues (K288, R281) in the C2B of SytI form the SytI-SNARE primary interface (Zhou

et al., 2015, Wang et al., 2016). The primary interface is critical for SytI´s function to

efficiently trigger release in cultured neurons and chromaffin cells (Gaffaney et al., 2008;

Mohrmann et al., 2013; Rickman et al., 2006; Schupp et al., 2016; Xue et al., 2008; Zhou

et al., 2015). Indeed, mutations of amino acids of SNAP25 SN2 (such as E55, D51, E52)

that line the SytI-SNARE primary interface resulted in a phenotype similar to that of the

CpxII glutamate cluster mutation (Mohrmann et al., 2013). Furthermore, mutations within

SNAP25 (e.g. D166A, E170E, Mohrmann et al., 2013), which contribute to the primary

SytI-SNARE interface and are located on the opposite side of the Cpx-bound SNARE

complex (Chen et al., 2002), impair Cpx-mediated clamping of liposome fusion *in vitro*

(Schupp et al., 2016), an observation consistent with the hypothesis of an independent

binding site of the CpxII CTD on the SNARE complex. Collectively, these results point to

a structurally plausible colocalisation of 'triggering' and 'unclamping' mechanisms on the

surface of the SNARE complex.

SytI – CpxII interplay

By analyzing the impact of single and combined deficiencies for CpxII and SytI (CpxII-/-;

SytI-/-; CpxII-/-:SytI-/- dko), we found that CpxII functions interdependently with SytI

(Figure 5). Additional loss of SytI in the absence of CpxII had no impact on either the

magnitude or the kinetics of exocytosis. This suggests that all functions of SytI depend

on CpxII. In sharp contrast, additional loss of SytVII in the absence of CpxII further

aggravated the secretion deficit (CpxII-/-:SytVII-/- dko, **Figure 5**). Moreover, CpxII

expression experiments show that CpxII fails to boost exocytosis in the absence of SytI,

but is able to do so in the absence of SytVII **(Figure 5 figure supplement 2)**. Remarkably,

the profound loss of secretion in the CpxII:SytVII double deficiency closely resembles the

phenotype of the SytI:SytVII deficiency (Schonn et al., 2008; Dhara et al., 2014),

indicating the CpxII serves as a gatekeeper for all SytI functions. In murine hippocampal

neurons, loss of CpxI and Syt1 has additive effects on fast synchronous release,

suggesting independent mechanisms (Xue et al., 2010). On the other hand, the same

study also showed that Syt1 heterozygosity fails to reduce release probability in wild-type

neurons, but does so in the absence of Cpx, again suggesting that Cpx and Syt1 may

functionally interact in Ca^{2+}-triggered release. Recent structural evidence provides

potential cues explaining the differential impact of SytI or SytVII deficiencies in the

absence of CpxII (Zhou et al., 2015; Zhou et al., 2017). For instance, core residues in SytI

that are essential for SNARE binding at the 'primary' interface are not conserved in SytVII,

which has been implicated as Ca^{2+} sensor for asynchronous synaptic vesicle exocytosis

(Wen et al., 2010; Bacaj et al., 2013; Liu et al., 2014; Jackman et al., 2016). Thus, it

seems unlikely that SytVII engages the SNARE complex in the same manner as SytI. SytI

and Cpx, instead, form a separated but continuous α-helix at the 'tripartite' SytI-Cpx-

SNARE interface, which may serve as pre-fusion intermediate for the primed vesicle state

(Zhou et al., 2017). Furthermore, sequence alignment reveals that amino acids involved

in specific side-chain interactions of the tripartite interface are only highly conserved in

SytI, SytII and SytIX and thus in the isoforms involved in rapid evoked release. Taken

together, our results show that SytI and CpxII acts as interdependent allies in rapid

secretion, whereas SytVII acts autonomously and thereby additively as Ca^{2+}-sensor that

accelerates vesicle recruitment during train stimulation, resulting in sustained release (Liu

et al., 2014; Tawfik et al., 2021).

Despite extensive research, there is still no consensus on the mechanisms underlying the

fusion-promoting function of Cpx. Using hippocampal microisland cultures, it was found

that the N-terminal domain of CpxI (amino acid positions 1-26) is required to restore basal

evoked transmitter release (Xue et al., 2007) and that the NTD, by binding to the C-

terminal end of the SNARE complex, provides conformational support to the SNARE

machinery (Xue et al., 2010). In the same line, individual liposome-liposome content

mixing experiments with the minimal fusion machinery showed that the Cpx-N terminus

decisively improves the fusion fidelity (Lai et al., 2014). On the other hand, Cpx deficiency

was found to be associated with a reduction in the Ca^{2+} sensitivity of evoked release in

various preparations, including neurons and endocrine cells (Reim et al., 2001; Tadokoro

et al., 2005; Huntwork and Littleton, 2007; Xue et al., 2007; Jorquera et al., 2012; Cho et

al., 2014; Dhara et al., 2014). In the present work, we show that the CpxII-NTD

accelerates synchronous exocytosis by enhancing its apparent Ca^{2+} affinity in a SytI

dependent manner **(Figures 7 & 8)**. While an excess of CpxII rescues the slow release

kinetics of the SytI R233Q mutant (Sorensen et al., 2013), this is further slowed by

expression of the CpxII ΔN mutant **(Figure 7)**. In contrast, in the absence of SytI, no

corresponding changes were observed with expression of the CpxII ΔN mutant **(Figure**

6). The latter result confirms the strongly interdependent action of CpxII and SytI and is

difficult to reconcile with a potential direct action of the CpxII NTD on SNARE assembly

(Xue et al., 2010). Moreover, in response to slow ramp-like Ca^{2+}-stimuli, CpxII and its ΔN-

mutant led to opposite shifts in the Ca^{2+} dependence of secretion **(Figure 8C)** and to

corresponding changes in the fusion rate as the function of [Ca]i **(Figure 8D)**. The results

favor a model wherein the CpxII NTD either directly regulates the biophysical properties

of the Ca^{2+}-sensor by increasing the forward rate of Ca^{2+}-binding or indirectly affects SytI-

SNARE or SytI-membrane interactions, thereby, lowering the energy barrier of Ca^{2+}-

triggered fusion.

Taken together, Cpx impedes SNARE assembly via the hydrophobic face of the

amphipathic helix at its C-terminus and simultaneously provides structures with the

upstream glutamate cluster that adjust SytI-dependent triggering of the fusion

mechanism. Complexin thus maintains a delicate balance between preventing premature

release and triggering rapid exocytosis by manipulating the assembly of SNARE motifs

and assisting SytI in triggering efficient rapid exocytosis.

Materials and methods

Mutagenesis and viral constructs

Briefly, mutations in CpxII were generated by overlap extension polymerase chain

reaction (PCR). For expression in chromaffin cells, cDNAs encoding for CpxII or its

mutant variants were sub-cloned into the first open reading frame (ORF) of a bicistronic

Semliki Forest vector (pSFV1, Invitrogen, San Diego, CA). Enhanced-GFP expression

from the second ORF allowed for the identification of infected cells.

Culture of chromaffin cells and electrophysiological recordings

All experiments were performed on mouse chromaffin cells prepared at postnatal day 0–

1 from knockout /knockin pups and their littermate control (+/+ or +/-) which were identified

by genotyping. Preparation of adrenal chromaffin cells was done as described previously

(Borisovska et al., 2005). Electrophysiological recordings were carried out at DIV2, 5.5h

after addition of viral particles to the cultured cells.

Electrophysiological recordings were done at room temperature. Cells were recorderded

in an extracellular Ringer's solution containing (in mM): 130 NaCl, 4 KCl, 2 $CaCl_2$, 1

$MgCl_2$, 30 glucose, 10 HEPES-NaOH, pH 7.3, 310 mOsm. Recordings of membrane

capacitance (reflecting vesicle fusion) and ratiometric $[Ca^{2+}]_i$ changes (using Fura-2 and

Furaptra) were performed as described previously (Borisovska et al., 2005). The

intracellular solution for Ca^{2+}-uncaging experiments contained (in mM): 110 Cs-

glutamate, 8 NaCl, 3.5 $CaCl_2$, 5 NP-EGTA, 0.2 fura-2, 0.3 furaptra, 2 MgATP, 0.3

Na_2GTP, 40 HEPES-CsOH, pH 7.3, 300 mOsm. Membrane capacitance was recorded

with the Pulse software (HEKA, Lambrecht, Germany) and capacitance measurements

were performed according to the Lindau–Neher technique (sine wave stimulus: 1000 Hz,

35 mV peak-to-peak amplitude, DC-holding potential -70 mV). Synchronized vesicle

exocytosis was stimulated by a brief UV-flash that led to Ca^{2+}-uncaging upon photolysis

of NP-EGTA. Priming of secretory vesicles was promoted by infusing cells for two minutes

with the intracellular solution containing 500 nM free $[Ca^{2+}]_i$. During this phase premature

vesicle secretion was assessed. The flash-evoked capacitance increase was

approximated with the function: $f(x)=A0+A1(1-exp[-t/τ1])+A2(1-exp[-t/τ2])+kt$ (Rettig

and Neher, 2002). For calcium ramp experiments, after vesicle priming, intracellular Ca^{2+}

was gradually uncaged by the continuous UV illumination from the monochromator. The

resulting membrane capacitance increase was approximated with a polynomial function.

The size of the synchronized exocytotic burst (EB), the remaining pool and the slope of

the capacitance trace were determined at fixed time intervals of 40 ms ($\square$Cm/0.04s) for

each cell to match the time interval of the Ca^{2+}-measurement. The secretion rate for each

interval was determined by rate [1/sec] = slope [fF/sec] / remaining pool [fF]. For

averaging data from different cells, the corresponding rates were plotted against the Ca^{2+}

rise interpolated at 50 nM Ca^{2+} intervals. All signals were analyzed with customized

IgorPro routines (Wavemetrics, Lake Oswego, OR).

Immunocytochemistry

Chromaffin cells were processed either 5.5 hr (**Figure 3 Suppl. 1**) or for 2.5 hr (**Figure 3**

Suppl. 2) after virus infection as described previously (Makke et al., 2018). A homemade,

affinity purified rabbit polyclonal antibody against CpxII (epitope: amino acids 1–100 of

CpxII) was used for all immunofluorescence experiments described in the manuscript. In

colocalization experiments (**Figure 3 Suppl. 2**), chromaffin cells were co-stained with

rabbit polyclonal CpxII and mouse monoclonal SybII antibodies (clone 69.1, antigen

epitope amino acid position 1–14, kindly provided by R. Jahn, MPI for Biophysical

Chemistry, Göttingen, Germany). Immunopositive signals were determined after

threshold adjustment (4x background signal) and cytofluorgram as well as Pearson's co-

localization coefficient were analyzed with ImageJ (JACoP plugin).

Biochemistry

Recombinant N-terminal tagged GST fusion proteins (pGEX-KG-vector with an internal

thrombin cleavage site) were expressed in the *E. coli* strain BL21DE3 and purified using

glutathione-agarose according to the manufacturer's instructions. 250µg recombinant

GST-CpxII-CTD or its mutant variants were incubated with 100µL GST bead slurry for 1

hour at room temperature while shaking at 1000 rpm (Vibrax, IKA, Staufen, Germany).

Beads were then centrifuged (200 rpm, 3 mins) and washed 3 times with assay buffer

containing: 130mM NaCl, 50 mM HEPES, 1 mM EDTA, 1mM DTT, 1% Triton X100, and

complete protease inhibitor. Integrity and purity of bound GST protein was verified by

Coomassie staining. 170µl Triton X-100 extract of mouse brain homogenate (0.7 mg/ml,

containing 130 mM NaCl, 50 mM HEPES-NaOH, 1 mM EDTA, 2% Triton X-100, 1 mM

PMSF, pH 7.3) was incubated with 50 µl beads (2 h, room temperature, final assay

volume 250 µl). Immobilized proteins were eluted by thrombin cleavage (thrombin 0.02

UN/µl, Sigma, Germany). The unbound fraction (*Sbh*), the thrombin eluted fraction (*STh*)

and the pellet fraction after thrombin cleavage (*P*) were analyzed by 12% SDS-PAGE and

Western blotting (**Figure 3**). For the detection of Syntaxin 1a, SNAP25, SybII, SytI and

Synapsin1 the following mouse antibodies from Synaptic systems (Göttingen, Germany)

were used: Anti-Syntaxin1 (CL 78.2) No: 110 001; SNAP25 (CL 71.1), No:111 011;

SynaptobrevinII (CL 69.1), No: 104 211; Synaptotagmin 1 (CL 41.1), No:105 011; Anti-

Synapsin 1 (CL 46.1), No:106 001. Primary mouse antibodies were used at a dilution of

1:1000. Immunoreactive bands were visualized with secondary goat-anti mouse or goat

anti-rabbit antibodies conjugated with horseradish peroxidase and with an enhanced

chemiluminescence system (Thermo Fisher Scientific, Schwerte, Germany).

Statistics

Data in bar graphs and X-Y plots present means ± SEM. Statistical analyses were

performed using Prism 7 (GraphPad). All statistical data is summarized in the

corresponding 'Source Data' tables and were tested for normality with Kolmogorov-

Smirnov test. Data from groups with nonparametric distribution was subjected to Kruskal–

Wallis followed by Dunn's post hoc test when at least one group showed a nonparametric

distribution. Data from groups showing parametric distributions was subjected to ordinary

one-way ANOVA followed by Tukey-Kramer post hoc test. Significance levels: '*' $p < 0.05$,

'**' $p < 0.01$, and '***' $p < 0.001$.

Conflict of interest

The authors declare no competing financial interest.

Acknowledgement

We thank Marina Wirth. This work was supported by the Collaborative Research Center 1027 to DB.

Figures and Legends

Figure 1. An unrelated amphipathic helix functionally substitutes for the last 19 amino acids of CpxII CTD. (A) Primary sequence of CpxII´s CTD (left) and the mutant CpxII 1-115 amphipathic α helix (right). **(B)** Mean [Ca]i levels (upper panel) and the corresponding membrane capacitance response (lower panel) from CpxII ko cells (red n=15), and those expressing either wt CpxII (blue n=15), the CpxII 1-115 mutant (purple n=14), or the CpxII 1-115 amphipathic helix mutant (orange n=16). **(C)** Amplitudes of the

readily releasable pool (RRP), the slow releasable pool (SRP), and the rate of sustained
release (SR). CpxII 1-115 amphipathic helix mutant fully supports exocytosis like the wt
protein. **(D)** The time constants for the EB components (τRRP and τSRP), and the
exocytotic delay are rescued by the CpxII mutants like the wt protein. **(E)** Mean premature
secretion of the tested groups at similar submicromolar [Ca]i before triggering the flash
response shown in (B). **(F)** The rate of tonic exocytosis (determined at similar $[Ca^{2+}]i$: in
nM, CpxII ko: 789 ± 120; CpxII ko + CpxII: 777 ± 57; CpxII ko+ CpxII 1-115: 739 ± 84;
CpxII ko+ CpxII1-115 amph helix: 796 ± 64) is significantly reduced with CpxII and CpxII
1-115 amph helix mutant but not with the truncated CpxII 1-115 mutant. ANOVA or
Kruskal-Wallis followed by corresponding post-hoc test. **p<0.01; ***p<0.001. Error bars
indicate mean ± SEM.

Figure 2. Disrupting the hydrophobic face of the amphipathic helix impairs the ability of the CpxII CTD to prevent premature release. (A, D) Primary sequences depicting the point mutations L124E-L128E (A) and L124W-L128W (D) within the amphipathic region of CpxII CTD. **(B, E)** Mean [Ca]i levels (upper panel) and the corresponding ∆CM (lower panel) from CpxII ko cells (red n=25), and those expressing CpxII wt (blue n=25), or the mutants CpxII L124E-L128E (yellow n=26) in (B), and CpxII ko cells (red n=18) or those expressing the CpxII wt protein (blue n=15) or CpxII LL-WW (dark green n=12) in (D). Mean amplitudes of the RRP, SRP and SR show a significant reduction in the size of the EB by the CpxII L124E-L128E mutant, whereas CpxII L124W-L128W rescued the synchronized exocytosis like CpxII wt. **(C, F)** Average tonic exocytosis at similar submicromolar [Ca]i levels. Note that the mutant CpxII L124W-L128W fully clamps premature secretion like the CpxII wt protein, but the CpxII L124E-L128E mutant fails to do so. ANOVA or Kruskal-Wallis followed by corresponding post-hoc test. *p<0.05; **p<0.01; ***p<0.001. Error bars indicate mean ± SEM.

Figure 2 Suppl. 1. The CpxII CTD mutants CpxII L124E-L128E and CpxII L124W-L128W rescue the kinetics of synchronized exocytosis. (A, B) The kinetics of synchronized exocytosis as well as the delay of the stimulus-secretion coupling of CpxII ko cells (red) are fully restored by the CpxII L124E-L128E (yellow) as well as the CpxII L124W-L128W mutant (green) when compared with expression of the CpxII wt protein (blue), consistent with the view that exocytosis timing is determined by the functionally intact NTD. ANOVA or Kruskal-Wallis followed by corresponding post-hoc test. *p<0.05; ***p<0.001. Error bars indicate mean ± SEM.

Figure 2 Suppl. 2. The hydrophobic face of CpxII's amphipathic helix is essential to suppress premature release. (A) Primary sequences showing point mutations L117W/–L121W and L117E/–L121E within the amphipathic α helix of CpxII CTD. **(B)** Mean [Ca]i levels (upper panel) and the corresponding average increase in the membrane capacitance from CpxII ko cells (red, n=26), and those expressing CpxII wt (blue, n=25), or the mutants CpxII L117E-L121E (yellow, n=25), or CpxII L117W-L121W (dark green, n=14). **(C)** Mean amplitudes of the RRP, SRP and SR show a significant reduction for the CpxII L117E-L121E mutant while CpxII L117W-L121W rescues the synchronized exocytosis like the expressed CpxII wt protein. **(D)** All mutant variants restore exocytosis timing of CpII ko cells like the CpxII wt protein. **(E, F)** Average tonic exocytosis at similar submicromolar [Ca]i levels. The CpxII L117E-L121E mutant failed to rescue premature secretion phenotype of CpxII ko cells, but CpxII L117W-L121W mutant hindered it like the CpxII wt protein. ANOVA or Kruskal-Wallis followed by corresponding post-hoc test. ***p<0.001. Error bars indicate mean ± SEM.

Figure 2 Suppl. 3: Alanine substitutions on the polar face of CpxII CTD have no impact on its clamping ability. (A) Primary sequences of SNAP-25 SN1 and CpxII CTD showing the positions of the alanine substitutions (subsequent panels refer to the same color coding). **(B)** Mean [Ca]i levels (upper panel) and the corresponding membrane capacitance increase from CpxII ko cells (red n=42), and those expressing either CpxII wt (blue n=58), or the mutants CpxII D118A (yellow, n=17) CpxII Q129A (pink, n=19), CpxII D130A (violet, n=15), CpxII K133A (green, n=17). **(C)** When compared with expression of the CpxII wt protein, all CpxII mutants fully support an undiminished exocytotic burst, without changing either the RRP, SRP or the SR. **(D)** Kinetic parameters of the synchronized exocytosis are rescued by CpxII and its mutant variants. **(E,F)** CpxII mutants suppress premature vesicle exocytosis at submicromolar calcium concentrations with similar efficacy as the CpxII wt protein. ANOVA or Kruskal-Wallis followed by corresponding post-hoc test.. **p<0.01; ***p<0.001. Error bars indicate mean ± SEM.

Figure 2 Suppl. 4: The charge reversal mutation D118K-D130K does not affect the inhibitory function of the CpxII CTD. (A) Primary sequence of CpxII CTD showing the double point mutation D118K-D130K. **(B)** Mean [Ca]i levels (upper panel) and the corresponding average increase in CM from CpxII ko cells (red n=27), and those expressing either CpxII wt (blue n=29) or the mutant CpxII DD-KK (green n=14). **(C)** The D118K-D130K mutant fully rescued all component of the exocytotic burst and the SR, like the expression of the CpxII wt protein. **(D-F)** Kinetic parameters of the synchronized exocytosis (D) and the suppression of premature vesicle exocytosis (E, F) were unaffected by the CpxII mutant. ANOVA or Kruskal-Wallis followed by corresponding post-hoc test. *p<0.05; ***p<0.001. Error bars indicate mean ± SEM.

Figure 3. CpxII-CTD and its mutant variants differentially interact with SNAREs and Sytl. (A, C) The GST-CpxII CTD fusion protein co-precipitates Sytl and Syx1a from detergent extract of mouse brain. Substitution of leucine residues with glutamate residues (CpxII-CTD L124E-L128E, LE) abolishes any binding to CpxII CTD (A). Substitution with tryptophan residues (CpxII-CTD L124W-L128W, LW), instead, or replacement of the entire CpxII CTD with an unrelated amphipathic helix (CpxII-CTD-ampH, C) is tolerated. Samples were eluted from the column by thrombin cleavage. Equal volumes of supernatant (S_{bh}), the thrombin-eluted (S_{Th}) and the non-eluted fraction (P) were analyzed by SDS-PAGE (12% gel) and Western blotting with antibodies against the indicated antigens. No binding to GST alone could be detected. **(B, D)** Corresponding Coomassie gels documenting the integrity of the GST fusion proteins, remaining in the non-eluted fraction (P) after thrombin cleavage.

Figure 3 Suppl. 1. Expression analysis of CpxII and its mutants reveals similar levels of protein expression. (A) Exemplary images of CpxII ko, wt,or CpxII ko cells expressing CpxII wt or its mutant variants. Exposure times were adjusted to avoid the saturation of the sensor (CpxII ko 2s, wt 300ms CpxII ko over expressing CpxII or the CpxII mutants 30 ms). **(B)** Mean total fluorescence intensity (corrected for the exposure time) shows that CpxII wt and its mutants are expressed to similar levels in CpxII ko cells. Number of cells for each group is indicated in the corresponding bar. Kruskal-Wallis followed by Dunn's post-hoc test. ***p<0.001. Error bars indicate mean ± SEM.

(A)

wt
CpxII 488   5µm   SybII 555   5µm   overlap   5µm

ko+CpxII wt

ko+Cpx 1-115 amph. α helix

ko+CpxII L124E-L128E

ko+CpxII L124W-L128W

ko+CpxII E-A

ko+CpxII ΔN

ko+CpxII DDKK

(B)

wild type (boxed area)
Pearson´s coeff. 0.85

X-axis: 555 SybII flurescence int. x 10^3 a.u.
Y-axis: 488 CpxII flurescence int. x 10^3 a.u.

(C)

*

Y-axis: Pearson´s coefficient

Bars (with n values):
- wt: 32
- CpxII wt: 29
- CpxII 1-115 amph. α helix: 15
- CpxII ΔN: 21
- CpxII E-A: 21
- CpxII L124W-L128W: 21
- CpxII L124E-L128E: 25
- CpxII D118A: 31
- CpxII D130A: 32
- CpxII D118K-D130K: 34
- CpxII K133A: 23
- CpxII Q129A: 23

CpxII ko +

Figure 4. The cluster of glutamate residues in CpxII CTD facilitates Ca²⁺ triggered exocytosis. (A) Schematic representation of CpxII highlighting the position of the glutamate cluster within the CTD. Primary sequence of CpxII CTD showing the exchange of glutamate into alanine residues (CpxII E-A mutant). **(B)** Mean [Ca]i levels (upper panel) and CM response from wt cells (black n=18), CpxII ko cells (red n=19), and CpxII ko cells expressing either CpxII wt (blue n=17), or the mutant CpxII E-A (green n=~~25~~24). **(C)** The mutant CpxII E-A only partially restores the fast component (RRP) of synchronized

exocytosis. **(D)** The CpxII E-A mutation significantly slows down the speed of the RRP and SRP and delays the stimulus secretion coupling. **(E)** Average tonic exocytosis at similar submicromolar [Ca]i levels with the quantification in **(F)** showing that the mutant CpxII E-A effectively prevents premature vesicle loss at submicromolar [Ca]i levels like CpxII wt. ANOVA or Kruskal-Wallis followed by corresponding post-hoc test. *p<0.05; **p<0.01; ***p<0.001. Error bars indicate mean ± SEM.

Figure 5. CpxII and SytI act interdependently in triggering synchronized exocytosis.
(A) Mean [Ca]i levels (upper panel) and CM responses from wt cells (black n=50), SytI ko cells (green n=46), CpxII ko (red n=35), CpxII/SytI dko (dark blue n=26). **(B)** Quantification of the RRP and SRP size shows that SytI deletion in the absence of CpxII does not further aggravate the secretory deficit. **(C)** Normalized CM (as shown in A) scaled to the wt response 1s after the flash. Inset, extended scaling of normalized CM during the first 20ms after flash (arrow) depicting the delayed onset of secretion. **(D)** Kinetics of RRP and SRP exocytosis and the secretory delay. **(E)** Loss of SytI does not alter premature vesicle fusion. Additional loss of CpxII increases premature exocytosis to the level of CpxII ko **(F)**. **(G)** Mean [Ca]i levels (upper panel) and CM responses from wt cells (black n=17), SytVII ko cells (orange n=20), CpxII ko (red n=20), CpxII/SytVII dko (blue n=24). **(H)** The RRP and SRP sizes are reduced by either CpxII or SytVII single ko, while the combined loss of CpxII and SytVII abolishes the EB. **(I)** Normalized CM (as shown in G) scaled to the wt response 1s after the flash. Inset, extended scaling of normalized CM during the first 20ms after flash (arrow). **(J)** Loss of CpxII but not of SytVII slows the time constants of RRP and SRP exocytosis (τRRP, τSRP) and prolongs the secretory delay. **(K, L)** SytVII deficiency diminishes the elevated premature exocytosis of CpxII ko cells (SytVII CpxII dko) and reduces it compared to wt cells. ANOVA or Kruskal-Wallis followed by corresponding post-hoc test. *p<0.05; **p<0.01; ***p<0.001. Error bars indicate mean ± SEM.

Figure 5 Suppl. 1. CpxII clamps premature exocytosis independently of SytI. (A)
Mean time course of premature exocytosis from CpxII ko cells (red n=35), CpxII/SytI dko
(violet n=26), SytI ko (light green n=46), wt (black n=49), SytIko+CpxII (dark green n=22),
CpxII ko+CpxII (blue n=15). **(B, C)** Quantification of the premature exocytosis rate per 10
seconds at similar calcium levels shown in (C) for indicated groups. **(D, E)** The EB size
(RRP+SRP component, E) correlates inversely with the extent of the preceding
premature exocytosis upon changes of CpxII expression in the presence and in the
absence of SytI (solid lines linear regressions, R^2=0.99). ANOVA or Kruskal-Wallis
followed by corresponding post-hoc test. **p<0.01; ***p<0.001. Error bars indicate mean
± SEM.

Figure 5 Suppl. 2. CpxII boosts exocytosis in the absence of SytVII. (A) Mean [Ca]i levels (upper panel) and CM responses from wt cells (black n=17), SytVII ko cells (orange n=18), and SytVIIko cells expressing CpxII (green n=18). **(B)** SytVII deletion reduces the RRP and SRP size without affecting the sustained rate of secretion. **(C)** Kinetics of RRP and SRP exocytosis and the secretory delay. **(D)** Time course of premature exocytosis at submicromolar [Ca]i. **(E, F)** Loss of SytVII reduces premature vesicle fusion (E), which is further diminished upon CpxII expression at nearly comparable [Ca]i (F). ANOVA or Kruskal-Wallis followed by corresponding post-hoc test. *p<0.05; **p<0.01; ***p<0.001. Error bars indicate mean ± SEM.

Figure 6. CpxII NTD exclusively modulates the kinetics of SytI mediated exocytosis.
(A) Mean [Ca]i levels (upper panel) and CM response from wt cells (black n=13), SytI ko
cells (red n=19), and those overexpressing either CpxII wt (dark blue n=16), or the mutant
CpxII ∆N (green n=20). In the absence of SytI, Cpx ∆N fails to slow down exocytosis
timing as observed in SytVII cells (I). **(B-D)** Neither CpxII nor CpxII ∆N expression in SytI
ko cells alters the size of the pools, the SR rate (B) or the kinetics of synchronous
exocytosis (C, normalized CM; D, time constants of RRP and SRP exocytosis and
secretory delay). **(E, F)** Both, CpxII and CpxII ∆N expression, hinders asynchronous
release in the absence of SytI **(F)**. **(G)** Mean [Ca]i levels (upper panel) and CM response
of wt cells (black n=13), SytVII ko cells (orange n=17), and those expressing CpxII ∆N
(Light blue n=29). **(H)** Both the RRP and SRP sizes are reduced in the absence of SytVII.
(I) Normalized CM responses (of data shown in G) scaled to the wt response 1s after the
flash. Note that CpxII ∆N mutant slows kinetics of release in SytVII ko cells. **(J)** Loss of
SytVII speeds up exocytosis timing, which in turn is slowed down by additional expression
of CpxII ∆N. **(K, L)** Analysis of premature exocytosis showing reduced premature
exocytosis in SytVII ko cells compared to wt. CpxII ∆N suppresses asynchronous release
also in the absence of SytVII. ANOVA or Kruskal-Wallis followed by corresponding post-
hoc test. $*p<0.05$; $**p<0.01$; $***p<0.001$. Error bars indicate mean ± SEM.

Figure 7. CpxII overexpression rescues the slow secretory rates of SytI R233Q ki cells. (A) Mean [Ca]i levels (upper panel) and CM responses from wt (black n=35), SytI R233Q ki (violet n=39), and SytI R233Q ki cells expressing either CpxII wt (blue n=18), or the mutant CpxII ΔN (olive green n=11). **(B)** Compared to wt cells, SytI R233Q ki cells show an increased the RRP and SRP size, which are restored by CpxII or CpxII ΔN expression. **(C)** While CpxII speeds up exocytosis timing of SytI R233Q ki cells to the level of wt cells, CpxII ΔN slows it further down. Normalized CM responses (of data shown in A) scaled to the wt response 1s after the flash. **(D)** CpxII and CpxII ΔN expression oppositely regulate the kinetics of exocytosis (τRRP, τSRP) and the secretory delay. **(E, F)** Mean premature exocytosis, determined at similar submicromolar [Ca]i (in nM, SytI R233Q ki:623±31; SytI R233Q ki + CpxII:569±28; SytI R233Q ki + CpxII ΔN:600±26; wt: 582±22), showing that CpxII wt or CpxII ΔN expression effectively hinders premature vesicle secretion. ANOVA or Kruskal-Wallis followed by corresponding post-hoc test *p<0.05; **p<0.01; ***p<0.001. Error bars indicate mean ± SEM.

Figure 8. CpxII NTD increases the Ca²⁺ affinity of secretion. (A) Exemplary capacitance measurement of a chromaffin cell (lower panel) in response to a ramp like increase in intracellular Ca²⁺ (top panel). **(B)** The slope of the ΔCM (top panel), remaining pool (middle panel) and rate of fusion (lower panel) were determined from the data shown under (A) with a temporal interval of 40 ms to match the timing of the ratiometric Ca²⁺-measurement. The rate was determined by dividing the slope [fF/s] by the remaining pool size [fF]. **(C)** Mean profile of pool depletion as the function of [Ca]i (left panel) for SytI R233Q ki cells (violet n=15) and those overexpressing CpxII wt (blue n=14) or CpxII ΔN (olive green n=11). Note that expression of CpxII wt lowers [Ca]i for half-maximal pool depletion (P50), whereas CpxII ΔN increases it (right panel). **(D)** Double-logarithmic plot of the mean exocytotic rate as the function of [Ca]i for the indicated groups. CpxII overexpression increases the rate of secretion from SytI R233Q ki cells, whereas loss of CpxII NTD further slows it down. Hill equations with similar coefficient, but different KD value approximate the data (dashed lines). Extrapolation of the rates at low [Ca]i

coincides with the RRP kinetics (Figure 7D) determined in the flash experiment (hollow circle). ANOVA or Kruskal-Wallis followed by corresponding post-hoc test. *p<0.05; ***p<0.001. Error bars indicate mean ± SEM.

References

Bacaj, T., Wu, D., Yang, X., Morishita, W., Zhou, P., Xu, W., Malenka, R.C., and Sudhof, T.C. (2013). Synaptotagmin-1 and synaptotagmin-7 trigger synchronous and asynchronous phases of neurotransmitter release. Neuron *80*, 947-959.

Bera, M., Ramakrishnan, S., Coleman, J., Krishnakumar, S.S., and Rothman, J.E. (2022). Molecular determinants of complexin clamping and activation function. eLife *11*.

Borisovska, M., Zhao, Y., Tsytsyura, Y., Glyvuk, N., Takamori, S., Matti, U., Rettig, J., Sudhof, T., and Bruns, D. (2005). v-SNAREs control exocytosis of vesicles from priming to fusion. The EMBO journal *24*, 2114-2126.

Brunger, A.T., Choi, U.B., Lai, Y., Leitz, J., White, K.I., and Zhou, Q. (2019). The pre-synaptic fusion machinery. Curr Opin Struct Biol *54*, 179-188.

Bykhovskaia, M., Jagota, A., Gonzalez, A., Vasin, A., and Littleton, J.T. (2013). Interaction of the complexin accessory helix with the C-terminus of the SNARE complex: molecular-dynamics model of the fusion clamp. Biophysical journal *105*, 679-690.

Cai, H., Reim, K., Varoqueaux, F., Tapechum, S., Hill, K., Sorensen, J.B., Brose, N., and Chow, R.H. (2008). Complexin II plays a positive role in Ca2+-triggered exocytosis by facilitating vesicle priming. Proc Natl Acad Sci U S A *105*, 19538-19543.

Chen, X., Tomchick, D.R., Kovrigin, E., Arac, D., Machius, M., Sudhof, T.C., and Rizo, J. (2002). Three-dimensional structure of the complexin/SNARE complex. Neuron *33*, 397-409.

Cho, R.W., Kummel, D., Li, F., Baguley, S.W., Coleman, J., Rothman, J.E., and Littleton, J.T. (2014). Genetic analysis of the Complexin trans-clamping model for cross-linking SNARE complexes in vivo. Proc Natl Acad Sci U S A *111*, 10317-10322.

Cho, R.W., Song, Y., and Littleton, J.T. (2010). Comparative analysis of Drosophila and mammalian complexins as fusion clamps and facilitators of neurotransmitter release. Mol Cell Neurosci *45*, 389-397.

Courtney, K.C., Wu, L., Mandal, T., Swift, M., Zhang, Z., Alaghemandi, M., Wu, Z., Bradberry, M.M., Deo, C., Lavis, L.D.*, et al.* (2022). The complexin C-terminal amphipathic helix stabilizes the fusion pore open state by sculpting membranes. Nat Struct Mol Biol *29*, 97-107.

Dhara, M., Yarzagaray, A., Schwarz, Y., Dutta, S., Grabner, C., Moghadam, P.K., Bost, A., Schirra, C., Rettig, J., Reim, K.*, et al.* (2014). Complexin synchronizes primed vesicle exocytosis and regulates fusion pore dynamics. J Cell Biol *204*, 1123-1140.

Diao, J., Grob, P., Cipriano, D.J., Kyoung, M., Zhang, Y., Shah, S., Nguyen, A., Padolina, M., Srivastava, A., Vrljic, M.*, et al.* (2012). Synaptic proteins promote calcium-triggered fast transition from point contact to full fusion. eLife *1*, e00109.

Fernandez-Chacon, R., Konigstorfer, A., Gerber, S.H., Garcia, J., Matos, M.F., Stevens, C.F., Brose, N., Rizo, J., Rosenmund, C., and Sudhof, T.C. (2001). Synaptotagmin I functions as a calcium regulator of release probability. Nature *410*, 41-49.

Giraudo, C.G., Garcia-Diaz, A., Eng, W.S., Chen, Y., Hendrickson, W.A., Melia, T.J., and Rothman, J.E. (2009). Alternative zippering as an on-off switch for SNARE-mediated fusion. Science *323*, 512-516.

Gong, J., Lai, Y., Li, X., Wang, M., Leitz, J., Hu, Y., Zhang, Y., Choi, U.B., Cipriano, D., Pfuetzner, R.A.*, et al.*
(2016). C-terminal domain of mammalian complexin-1 localizes to highly curved membranes. Proc Natl
Acad Sci U S A *113*, E7590-E7599.
Hao, T., Feng, N., Gong, F., Yu, Y., Liu, J., and Ren, Y.X. (2023). Complexin-1 regulated assembly of single
neuronal SNARE complex revealed by single-molecule optical tweezers. Commun Biol *6*, 155.
Hobson, R.J., Liu, Q., Watanabe, S., and Jorgensen, E.M. (2011). Complexin maintains vesicles in the
primed state in C. elegans. Curr Biol *21*, 106-113.
Huntwork, S., and Littleton, J.T. (2007). A complexin fusion clamp regulates spontaneous neurotransmitter
release and synaptic growth. Nat Neurosci *10*, 1235-1237.
Jackman, S.L., Turecek, J., Belinsky, J.E., and Regehr, W.G. (2016). The calcium sensor synaptotagmin 7 is
required for synaptic facilitation. Nature *529*, 88-91.
Jahn, R., and Fasshauer, D. (2012). Molecular machines governing exocytosis of synaptic vesicles. Nature
490, 201-207.
Jorquera, R.A., Huntwork-Rodriguez, S., Akbergenova, Y., Cho, R.W., and Littleton, J.T. (2012). Complexin
controls spontaneous and evoked neurotransmitter release by regulating the timing and properties of
synaptotagmin activity. J Neurosci *32*, 18234-18245.
Kaeser-Woo, Y.J., Yang, X., and Sudhof, T.C. (2012). C-terminal complexin sequence is selectively required
for clamping and priming but not for Ca2+ triggering of synaptic exocytosis. J Neurosci *32*, 2877-2885.
Krishnakumar, S.S., Radoff, D.T., Kummel, D., Giraudo, C.G., Li, F., Khandan, L., Baguley, S.W., Coleman, J.,
Reinisch, K.M., Pincet, F.*, et al.* (2011). A conformational switch in complexin is required for
synaptotagmin to trigger synaptic fusion. Nat Struct Mol Biol *18*, 934-940.
Kummel, D., Krishnakumar, S.S., Radoff, D.T., Li, F., Giraudo, C.G., Pincet, F., Rothman, J.E., and Reinisch,
K.M. (2011). Complexin cross-links prefusion SNAREs into a zigzag array. Nat Struct Mol Biol *18*, 927-
933.
Kurokawa, A., Narukawa, M., Ohmoto, M., Yoshimoto, J., Abe, K., and Misaka, T. (2015). Expression of the
synaptic exocytosis-regulating molecule complexin 2 in taste buds and its participation in peripheral
taste transduction. J Neurochem *133*, 806-814.
Lai, Y., Choi, U.B., Zhang, Y., Zhao, M., Pfuetzner, R.A., Wang, A.L., Diao, J., and Brunger, A.T. (2016). N-
terminal domain of complexin independently activates calcium-triggered fusion. Proc Natl Acad Sci U
S A *113*, E4698-4707.
Lai, Y., Diao, J., Cipriano, D.J., Zhang, Y., Pfuetzner, R.A., Padolina, M.S., and Brunger, A.T. (2014).
Complexin inhibits spontaneous release and synchronizes Ca2+-triggered synaptic vesicle fusion by
distinct mechanisms. eLife *3*, e03756.
Lin, M.Y., Rohan, J.G., Cai, H., Reim, K., Ko, C.P., and Chow, R.H. (2013). Complexin facilitates exocytosis
and synchronizes vesicle release in two secretory model systems. J Physiol *591*, 2463-2473.
Liu, H., Bai, H., Hui, E., Yang, L., Evans, C.S., Wang, Z., Kwon, S.E., and Chapman, E.R. (2014). Synaptotagmin
7 functions as a Ca2+-sensor for synaptic vesicle replenishment. eLife *3*, e01524.
Lopez-Murcia, F.J., Reim, K., Jahn, O., Taschenberger, H., and Brose, N. (2019). Acute Complexin Knockout
Abates Spontaneous and Evoked Transmitter Release. Cell reports *26*, 2521-2530 e2525.
Lu, B., Song, S., and Shin, Y.K. (2010). Accessory alpha-helix of complexin I can displace VAMP2 locally in
the complexin-SNARE quaternary complex. J Mol Biol *396*, 602-609.
MacDougall, D.D., Lin, Z., Chon, N.L., Jackman, S.L., Lin, H., Knight, J.D., and Anantharam, A. (2018). The
high-affinity calcium sensor synaptotagmin-7 serves multiple roles in regulated exocytosis. J Gen
Physiol *150*, 783-807.
Makke, M., Mantero Martinez, M., Gaya, S., Schwarz, Y., Frisch, W., Silva-Bermudez, L., Jung, M.,
Mohrmann, R., Dhara, M., and Bruns, D. (2018). A mechanism for exocytotic arrest by the Complexin
C-terminus. eLife *7*.

Malsam, J., Barfuss, S., Trimbuch, T., Zarebidaki, F., Sonnen, A.F., Wild, K., Scheutzow, A., Rohland, L.,
Mayer, M.P., Sinning, I.*, et al.* (2020). Complexin Suppresses Spontaneous Exocytosis by Capturing the
Membrane-Proximal Regions of VAMP2 and SNAP25. Cell reports *32*, 107926.
Malsam, J., Parisotto, D., Bharat, T.A., Scheutzow, A., Krause, J.M., Briggs, J.A., and Sollner, T.H. (2012).
Complexin arrests a pool of docked vesicles for fast Ca2+-dependent release. The EMBO journal *31*,
3270-3281.
Martin, J.A., Hu, Z., Fenz, K.M., Fernandez, J., and Dittman, J.S. (2011). Complexin has opposite effects on
two modes of synaptic vesicle fusion. Curr Biol *21*, 97-105.
Maximov, A., Tang, J., Yang, X., Pang, Z.P., and Sudhof, T.C. (2009). Complexin controls the force transfer
from SNARE complexes to membranes in fusion. Science *323*, 516-521.
Mohrmann, R., Dhara, M., and Bruns, D. (2015). Complexins: small but capable. Cell Mol Life Sci *72*, 4221-
4235.
Nagy, G., Kim, J.H., Pang, Z.P., Matti, U., Rettig, J., Sudhof, T.C., and Sorensen, J.B. (2006). Different effects
on fast exocytosis induced by synaptotagmin 1 and 2 isoforms and abundance but not by
phosphorylation. J Neurosci *26*, 632-643.
Pinheiro, P.S., Houy, S., and Sorensen, J.B. (2016). C2-domain containing calcium sensors in
neuroendocrine secretion. J Neurochem *139*, 943-958.
Radoff, D.T., Dong, Y., Snead, D., Bai, J., Eliezer, D., and Dittman, J.S. (2014). The accessory helix of
complexin functions by stabilizing central helix secondary structure. eLife *3*.
Reim, K., Mansour, M., Varoqueaux, F., McMahon, H.T., Sudhof, T.C., Brose, N., and Rosenmund, C. (2001).
Complexins regulate a late step in Ca2+-dependent neurotransmitter release. Cell *104*, 71-81.
Rettig, J., and Neher, E. (2002). Emerging roles of presynaptic proteins in Ca++-triggered exocytosis.
Science *298*, 781-785.
Schonn, J.S., Maximov, A., Lao, Y., Sudhof, T.C., and Sorensen, J.B. (2008). Synaptotagmin-1 and -7 are
functionally overlapping Ca2+ sensors for exocytosis in adrenal chromaffin cells. Proc Natl Acad Sci U
S A *105*, 3998-4003.
Schupp, M., Malsam, J., Ruiter, M., Scheutzow, A., Wierda, K.D., Sollner, T.H., and Sorensen, J.B. (2016).
Interactions Between SNAP-25 and Synaptotagmin-1 Are Involved in Vesicle Priming, Clamping
Spontaneous and Stimulating Evoked Neurotransmission. J Neurosci *36*, 11865-11880.
Snead, D., Lai, A.L., Wragg, R.T., Parisotto, D.A., Ramlall, T.F., Dittman, J.S., Freed, J.H., and Eliezer, D.
(2017). Unique Structural Features of Membrane-Bound C-Terminal Domain Motifs Modulate
Complexin Inhibitory Function. Front Mol Neurosci *10*, 154.
Snead, D., Wragg, R.T., Dittman, J.S., and Eliezer, D. (2014). Membrane curvature sensing by the C-terminal
domain of complexin. Nat Commun *5*, 4955.
Sorensen, J.B., Fernandez-Chacon, R., Sudhof, T.C., and Neher, E. (2003). Examining synaptotagmin 1
function in dense core vesicle exocytosis under direct control of Ca2+. J Gen Physiol *122*, 265-276.
Strenzke, N., Chanda, S., Kopp-Scheinpflug, C., Khimich, D., Reim, K., Bulankina, A.V., Neef, A., Wolf, F.,
Brose, N., Xu-Friedman, M.A.*, et al.* (2009). Complexin-I is required for high-fidelity transmission at the
endbulb of Held auditory synapse. J Neurosci *29*, 7991-8004.
Sudhof, T.C., and Rothman, J.E. (2009). Membrane fusion: grappling with SNARE and SM proteins. Science
323, 474-477.
Sugita, S., Shin, O.H., Han, W., Lao, Y., and Sudhof, T.C. (2002). Synaptotagmins form a hierarchy of
exocytotic Ca(2+) sensors with distinct Ca(2+) affinities. The EMBO journal *21*, 270-280.
Tadokoro, S., Nakanishi, M., and Hirashima, N. (2005). Complexin II facilitates exocytotic release in mast
cells by enhancing Ca2+ sensitivity of the fusion process. J Cell Sci *118*, 2239-2246.
Tawfik, B., Martins, J.S., Houy, S., Imig, C., Pinheiro, P.S., Wojcik, S.M., Brose, N., Cooper, B.H., and
Sorensen, J.B. (2021). Synaptotagmin-7 places dense-core vesicles at the cell membrane to promote
Munc13-2- and Ca(2+)-dependent priming. eLife *10*.

Tokumaru, H., Shimizu-Okabe, C., and Abe, T. (2008). Direct interaction of SNARE complex binding protein
synaphin/complexin with calcium sensor synaptotagmin 1. Brain Cell Biol *36*, 173-189.
Trimbuch, T., and Rosenmund, C. (2016). Should I stop or should I go? The role of complexin in
neurotransmitter release. Nat Rev Neurosci *17*, 118-125.
Trimbuch, T., Xu, J., Flaherty, D., Tomchick, D.R., Rizo, J., and Rosenmund, C. (2014). Re-examining how
complexin inhibits neurotransmitter release. eLife *3*, e02391.
Vasin, A., Volfson, D., Littleton, J.T., and Bykhovskaia, M. (2016). Interaction of the Complexin Accessory
Helix with Synaptobrevin Regulates Spontaneous Fusion. Biophysical journal *111*, 1954-1964.
Voets, T., Moser, T., Lund, P.E., Chow, R.H., Geppert, M., Sudhof, T.C., and Neher, E. (2001). Intracellular
calcium dependence of large dense-core vesicle exocytosis in the absence of synaptotagmin I. Proc
Natl Acad Sci U S A *98*, 11680-11685.
Wen, H., Linhoff, M.W., McGinley, M.J., Li, G.L., Corson, G.M., Mandel, G., and Brehm, P. (2010). Distinct
roles for two synaptotagmin isoforms in synchronous and asynchronous transmitter release at
zebrafish neuromuscular junction. Proc Natl Acad Sci U S A *107*, 13906-13911.
Wickner, W., and Schekman, R. (2008). Membrane fusion. Nat Struct Mol Biol *15*, 658-664.
Wragg, R.T., Parisotto, D.A., Li, Z., Terakawa, M.S., Snead, D., Basu, I., Weinstein, H., Eliezer, D., and
Dittman, J.S. (2017). Evolutionary Divergence of the C-terminal Domain of Complexin Accounts for
Functional Disparities between Vertebrate and Invertebrate Complexins. Front Mol Neurosci *10*, 146.
Wragg, R.T., Snead, D., Dong, Y., Ramlall, T.F., Menon, I., Bai, J., Eliezer, D., and Dittman, J.S. (2013).
Synaptic vesicles position complexin to block spontaneous fusion. Neuron *77*, 323-334.
Xue, M., Craig, T.K., Xu, J., Chao, H.T., Rizo, J., and Rosenmund, C. (2010). Binding of the complexin N
terminus to the SNARE complex potentiates synaptic-vesicle fusogenicity. Nat Struct Mol Biol *17*, 568-
575.
Xue, M., Lin, Y.Q., Pan, H., Reim, K., Deng, H., Bellen, H.J., and Rosenmund, C. (2009). Tilting the balance
between facilitatory and inhibitory functions of mammalian and Drosophila Complexins orchestrates
synaptic vesicle exocytosis. Neuron *64*, 367-380.
Xue, M., Reim, K., Chen, X., Chao, H.T., Deng, H., Rizo, J., Brose, N., and Rosenmund, C. (2007). Distinct
domains of complexin I differentially regulate neurotransmitter release. Nat Struct Mol Biol *14*, 949-
958.
Yang, X., Cao, P., and Sudhof, T.C. (2013). Deconstructing complexin function in activating and clamping
Ca2+-triggered exocytosis by comparing knockout and knockdown phenotypes. Proc Natl Acad Sci U S
A *110*, 20777-20782.
Yang, X., Kaeser-Woo, Y.J., Pang, Z.P., Xu, W., and Sudhof, T.C. (2010). Complexin clamps asynchronous
release by blocking a secondary Ca(2+) sensor via its accessory alpha helix. Neuron *68*, 907-920.
Zdanowicz, R., Kreutzberger, A., Liang, B., Kiessling, V., Tamm, L.K., and Cafiso, D.S. (2017). Complexin
Binding to Membranes and Acceptor t-SNAREs Explains Its Clamping Effect on Fusion. Biophysical
journal *113*, 1235-1250.
Zhou, Q., Lai, Y., Bacaj, T., Zhao, M., Lyubimov, A.Y., Uervirojnangkoorn, M., Zeldin, O.B., Brewster, A.S.,
Sauter, N.K., Cohen, A.E.*, et al.* (2015). Architecture of the synaptotagmin-SNARE machinery for
neuronal exocytosis. Nature *525*, 62-67.
Zhou, Q., Zhou, P., Wang, A.L., Wu, D., Zhao, M., Sudhof, T.C., and Brunger, A.T. (2017). The primed SNARE-
complexin-synaptotagmin complex for neuronal exocytosis. Nature *548*, 420-425.

1062

1063