1
7 # Reading reshapes stimulus selectivity in the visual word form area
8

9 Vassiki S. Chauhan[1*], Krystal C. McCook[1], Alex L. White[1]

17 [1]Department of Neuroscience & Behavior
18 Barnard College, Columbia University
19 76 Claremont Ave
20 New York, NY 10027
21 USA

22
23 *Correspondence: vchauhan@barnard.edu

24
Abbreviated Title: Reading reshapes stimulus selectivity in the VWFA
Number of Figures: 7
Number of Tables: 3
Number of Extended Data Figures: 3
Number of Extended Data Tables: 3
Abstract Word Count: 250
Introduction Word Count: 635
Discussion Word Count: 1462
33

34 Acknowledgments: We are grateful to Nicole Oppenheimer and Kimya Firoozan for help with
35 data collection, Oscar Woolnough for advice on the insula, and to the National Eye Institute for
36 funding (grant R00 EY-029366).

Abstract

Reading depends on a brain region known as the "visual word form area" (VWFA) in left ventral occipito-temporal cortex. This region's function is debated because its stimulus selectivity is not absolute, it is modulated by a variety of task demands, and it is inconsistently localized. We used fMRI to characterize the combination of sensory and cognitive factors that activate word-responsive regions that we precisely localized in 16 adult humans (4 male). We then presented three types of character strings: English words, pseudowords, and unfamiliar characters with matched visual features. Participants performed three different tasks while viewing those stimuli: detecting real words, detecting color in the characters, and detecting color in the fixation mark. There were three primary findings about the VWFA's response: (1) It preferred letter strings over unfamiliar characters even when the stimuli were ignored during the fixation task; (2) Compared to those baseline responses, engaging in the word reading task *enhanced* the response to words but *suppressed* the response to unfamiliar characters. (3) Attending to the stimuli to judge their font color had little effect on the response magnitudes. Thus, the VWFA is uniquely modulated by a cognitive signal that is specific to voluntary linguistic processing and is not additive. Functional connectivity analyses revealed that communication between the VWFA and a left frontal language area increased when the participant engaged in the linguistic task. We conclude that the VWFA is inherently selective for familiar orthography, but it falls under control of the language network when the task demands it.

Significance Statement

The function of the "visual word form area" (VWFA) is controversial. Some researchers emphasize its bottom-up visual selectivity for words, hence the region's common name. Others argue that its activity is explained by feedback from regions that control attention or language. To seek clarity, we investigated what drives the VWFA: seeing words, attending visually to words, or trying to read words. None of those factors was sufficient on its own. Our results support a hybrid model: the VWFA has inherent selectivity for words, but its function is reshaped by voluntary language processing. Thus, with an integrated analysis of sensory inputs, task demands, and network connectivity, we provide some resolution to debates about this important region.

Introduction

Over the course of history, developments in human culture have changed our environment and required our brains to adapt. A unique example is a region in the left ventral occipito-temporal cortex that, in literate individuals, is specialized for processing written words (Nobre et al., 1994; McCandliss et al., 2003; Dehaene and Cohen, 2011). This region is often termed the visual word form area (VWFA). While the VWFA's plasticity during literacy acquisition is well established (Dehaene et al., 2015; Kubota et al., 2019), less is known about how its stimulus selectivity changes in the short term to meet task demands.

The nature of the VWFA's functional specialization is debated. Some researchers characterize it primarily as a visual region tuned for a particular category of stimulus (McCandliss et al., 2003; Dehaene et al., 2005). It may serve as the "orthographic lexicon" that identifies familiar letter strings (Glezer et al., 2009; Dehaene and Cohen, 2011; Yeatman and White, 2021). Other researchers emphasize its integration of linguistic information across sensory modalities (Price and Devlin, 2011; Qin et al., 2021; Dębska et al., 2023). We contribute to this lively discussion by examining how task demands shape the VWFA's stimulus selectivity as well as its communication with the canonical language network.

Like other areas around it, the VWFA is sensitive to visual stimulus properties. While it responds above baseline to many types of images, it prefers text (Ben-Shachar et al., 2007; Muayqil et al., 2015; White et al., 2023). Its response is affected by visual properties of words, such as their length (Woolnough et al., 2021), position (Rauschecker et al., 2012), and contrast (Kay and Yeatman, 2017). Beyond purely visual properties, the VWFA is also sensitive to

statistics of letter combinations (Vinckier et al., 2007; Woolnough et al., 2021; Gagl et al., 2022),

and word frequency (Kronbichler et al., 2004).

Moreover, the VWFA's activity is influenced by attentional allocation and task demands

(Mano et al., 2013). Words evoke larger responses when they are attended than ignored (Kay

and Yeatman, 2017; White et al., 2019, 2023). There is also evidence for top-down language

influences without visual input, such as during spoken language comprehension (Planton et al.,

2019) or reading Braille (Reich et al., 2011; Striem-Amit et al., 2012). Indeed, the VWFA's

function is supported by its connections with language and attention networks (Yeatman et al.,

2012; Stevens et al., 2017; Lerma-Usabiaga et al., 2018; Chen et al., 2019; Li et al., 2020).

To better understand the VWFA's functional specialization, we used functional magnetic

resonance imaging (fMRI) to examine the *interaction* of stimulus features and task demands. On

each trial, we briefly flashed a character string that was either a real English word, a

pronounceable pseudoword, or a string of unfamiliar "false font" characters (see **Figure 1C**). The

false fonts were matched to the Latin fonts in visual features (size, number of strokes,

perimetric complexity, etc.) (Vidal et al., 2017; Vildavski et al., 2022).

The stimulus types were randomly intermingled during three different tasks. The *fixation*

color task was to report whether the fixation dot turned red. This task evaluated the VWFA's

inherent preference for words in the absence of voluntary attention or linguistic processing. The

dot's color change was simultaneous with the character string's brief presentation and set to

individual detection thresholds. The *stimulus color task* was to report whether the character

string was red or gray. This task evaluated the effect of attending to the stimuli, which may

result in automatic linguistic processing. The *lexical decision task* was to report whether the

stimulus was a real word or not. This task required explicit linguistic processing.

By comparing stimulus responses and functional connectivity across the tasks, we ask:

what maximally drives activity in the VWFA? Simply viewing words, attending to the visual

properties of words, or explicitly engaging in a linguistic task?

Materials and Methods

Participants

We recruited 17 participants (Age range 19 to 38, 21.12 ± 4.44, 4 self-identified as male,

1 left-handed) from the Barnard College and Columbia University student body. The study was

approved by the Internal Review Board at Barnard College, Columbia University. All participants

provided written informed consent, acquired digitally, and were monetarily compensated for

their participation. All participants had learned English before the age of five.

We chose the sample size on the basis of a power analysis on a task effect observed in a

recent study with a similar experimental paradigm (White et al., 2023). Specifically, we focused

on the reduced response to non-letter shapes when they were attended than when ignored

(the smallest effect in that study). A simulation determined that a sample size of 17 would be

sufficient to detect, with 80% power, such an effect with a predicted mean (0.037) and standard

deviation (0.052, adjusted for the number of trials per condition).

To ensure high data quality, we used the following criteria for excluding functional runs

and participants. If the participant moved by a distance greater than 2 voxels (4 mm) within a

single run, that run was excluded from analysis. Additionally, if the participant responded in less

than 50% of the trials in the main experiment, that run was removed. Finally, if half or more of

the runs met any of these criteria for a single participant, that participant was dropped from

the dataset. Using these constraints, the analysis reported here is based on data from 16

participants. They ranged in age from 19 to 38 years (mean = 21.12 ± 4.58,). 4 participants self-

identified as male, and 1 was left-handed. A total of 6 runs were removed from three of the

remaining participants due to excessive head motion.

Equipment

We collected MRI data at the Zuckerman Institute, Columbia University, a 3T Siemens

Prisma scanner and a 64-channel head coil. In each MR session, we acquired a T1 weighted

structural scan, with voxels measuring 1 mm isometrically. We acquired functional data with a

T2* echo planar imaging sequences with multiband echo sequencing (SMS3) for whole brain

coverage. The TR was 1.5s, TE was 30 ms and the flip angle was 62°. The voxel size was 2 mm

isotropic.

Stimuli were presented on an LCD screen that the participants viewed through a mirror

with a viewing distance of 142 cm. The display had a resolution of 1920 by 1080 pixels, and a

refresh rate of 60 Hz. We presented the stimuli using custom code written in MATLAB and the

Psychophysics Toolbox (Brainard, 1997; Pelli, 1997). Throughout the scan, we recorded

monocular gaze position using an SR Research Eyelink 1000 tracker. Participants responded

with their right hand via three buttons on an MR-safe response pad.

Figure 1: Experimental Paradigm. (A) Example stimuli in the visual localizer scan for category-selective regions in the ventral temporal cortex. See Extended Data Figure 1-1 for list of all stimuli used in the experiment. **(B)** Regions of interest (ROIs) in one example participant. Regions in color were defined using the localizer in panel A. Regions in black were defined from the language localizer scan, and regions in white were obtained from a previous study (White et al., 2023). See Extended Data Figure 1-2 for ROIs all individual participants. **(C)** Example stimuli in the main experiment. **(D)** Mean accuracy in the three behavioral tasks plotted as a function of the character string's position in degrees of visual angle. Error bars are ±1 SEM. Stars and horizontal lines depict statistically significant task effects (* = p < 0.05, ** = p < 0.01, *** = p < 0.0001)

Main Experiment

Stimuli

Figure 1A shows example stimuli. A small dark fixation dot at screen center was present

throughout all trials against a white background (95% of screen maximum). The dot's diameter

was 0.11 degrees of visual angle, or dva. The stimuli were character strings from three different

categories: real English words (different parts of speech), pseudowords (meaningless but

pronounceable strings) and visually matched false fonts. The pseudowords were generated

using MCWord to match the bigram and trigram statistics of the real English words (Medler and

Binder, 2005). All strings were composed of four characters each. For the full list of stimuli, see

Extended Data Figure 1-1.

Two different fonts were used for the real words and pseudowords: Courier New and

Sloan. The false fonts, BACS2 (Vidal et al., 2017) and PseudoSloan (Vildavski et al., 2022), were

chosen to match the visual properties of the real fonts. The false Courier characters were

matched in size, symmetry, and the number of strokes, junctions and terminations. The false

Sloan characters were matched in ink area and perimetric complexity. All four fonts were scaled

such that the height of the letter 'x' (or its false font sibling) was 0.41 dva. The distance between

centers of neighboring letters was roughly 0.53 dva (ranging 0.52–0.55 across fonts). Each word

was 2 dva wide, and between 0.41 and 0.78 dva tall.

On each trial, a single character string flashed for 150 ms at one of three locations:

centered at fixation, 3 dva left, or 3 dva right). The stimulus was followed by a blank with only

the fixation mark present for 3850 ms, during which the participant had the opportunity to

respond with a button press. After every five trials, there was a rest period (no task except to

fixation on the dot). The duration of the rest period was either 4, 6 or 8 s in duration (randomly

and uniformly selected).

The color of the character string was dark gray on 67% of trials (70% contrast against the

background) or dark red on 33% of trials. The dark red was roughly the same luminance as the

dark gray. Similarly, on an independently selected 33% of trials, the fixation dot turned from

dark gray to dark red during the 150 ms that the character string appeared. On the remaining

trials the fixation dot remained dark gray. See the description of tasks below for how the colors

were controlled.

Each character string was presented exactly once to each participant during the

experiment. The sets of words used for each participant were also balanced to equalize the

orthographic neighborhood size, log lexical frequency, mean reaction time and accuracy

estimated from the English Lexicon Project (Balota et al., 2007) across all conditions in the

experiment. We generated 6 sets of word-to-location assignments, and randomly used one for

each participant.

Task

Participants performed three different tasks during different runs, two of which required

attending to the character strings, and one that encouraged participants to ignore them. In the

lexical decision task, participants reported whether the character string on each trial was a real

word or not. In the ***stimulus color task***, participants reported whether the color of the character

string was red or gray. In the ***fixation color task***, participants reported whether or not the

fixation dot turned red.

The fixation color task served as the baseline for neural responses: the same stimuli

were presented as in the other two tasks, but the participants were required to ignore them as

they tried to detect a subtle, threshold-level color change that occurred at the exact same time

the character string appeared. The 150 ms duration did not allow for attention to switch from

the fixation dot to the character string. For the stimulus color task and the fixation color task,

the saturation of the red color was updated using an adaptive staircase that converged at the

80% correct detection threshold. During runs when the character (or fixation dot) color was not

task-relevant, the saturation levels across trials were replayed from the staircase in another

color task run.

Across all three tasks, the probability of a real word appearing was 0.33. The probability

that the character string was red was 0.33. Similarly, the probability that the fixation dot turned

red was 0.33. All three of those events were independent of each other. Thus, for all three

tasks, the probability of a "target" (a real word, a red character string, or a red fixation dot) was

0.33. All that changed across runs was which stimulus attributes were task-relevant.

Participants responded via a button press with their right index finger for "yes"

responses (word or not, red or not), and right middle finger for "no" responses. Participants

were instructed to respond within 3s after each stimulus and to prioritize accuracy over speed.

<u>Procedure</u>

Before the first scanning session, participants were trained on all tasks in a testing room

outside the scanner. They performed a minimum of 24 trials of each task, and were also trained

to calibrate the eye-tracker and to maintain fixation. Instructions reminded them to respond as

accurately as possible.

The main experiment consisted of fifteen 5-minute runs, with 5 runs for each of the

three tasks. The order in which the tasks were presented was:

LD SC FC SC FC LD FC LD SC LD FC SC SC LD FC

where LD is lexical decision, SC is stimulus color, and FC is fixation color task. This order was

chosen such that each task was equally likely to follow both of the others.

Each run began with written instructions on the screen, informing the participant which

task to perform. They pushed a button to proceed, and then were quizzed on which task they

were to perform in that run. The run would not start until the participant answered correctly, to

ensure that they performed the correct task.

There was no feedback about response accuracy after each trial, but at the end of each

run, text appeared on the screen to report their percent correct, along with the percentage of

trials in which they failed to make a response.

The sequence of trials within each run was counterbalanced and uniquely generated for

each participant. There were 45 trials in each run, and the category of the character string

stimulus was randomized. In summary, there were 27 unique conditions, defined by stimulus

type, stimulus location, and task. We collected 25 trials per condition.

Predictions

If the VWFA is sensitive only to stimulus features, it should respond more strongly to

words than false fonts by the same degree across tasks. If it is sensitive only to linguistic task

demands, it should respond most strongly during the lexical task regardless of stimulus type.

Other hypotheses involve interactions; for instance, the VWFA's preference for words may be

due to top-down feedback and arise only when the stimuli are attended. It is difficult to predict

whether such modulations would occur only during an explicitly linguistic task, or whether they

would occur *automatically* whenever the stimuli are attended (as in the stimulus color task).

Moreover, the response to false fonts may even be suppressed when attended, similar to what

was found during a shape judgement task (White et al., 2023).

Localizer for visual category-selective ventral temporal regions

We examined activity in regions of interest (ROIs) that we defined precisely in each

individual's brain using independent localizer scans (**Figure 1A**). This is necessary because the

VWFA is composed of several small patches that differ across individuals (Centanni et al., 2019;

Caffarra et al., 2021). Averaging in a template brain obscures word-selective processing (Glezer

and Riesenhuber, 2013; Mahowald and Fedorenko, 2016; Stevens et al., 2017). An example

participant's ROIs are shown in **Figure 1B.**

Stimuli

Participants viewed sequences of images, each of which contained 3 items of one

category: words, pseudowords, false fonts, faces, and limbs (examples in **Figure 1A**). This

localizer was also used in White et al., 2023, which describes the stimuli in detail. The major

change here is the addition of the "limbs" category instead of objects. The grayscale limb

images included hands, arms, feet and legs (Stigliani et al., 2015).

Task and Procedure

Participants performed a one-back repetition detection task. On 33% of the trials, the

exact same images flashed twice in a row. The participant's task was to push a button with their

right index finger whenever they detected such a repetition. Each participant performed 4 runs

of the localizer task. Each run consisted of 77 four-second trials, lasting for approximately 6

minutes. Each category was presented 56 times through the course of the experiment.

Language Network Localizer

Stimuli

In order to map the canonical language network, we used a language localizer (Mahowald and Fedorenko, 2016). Details about the localizer are provided by the authors who developed the resource. Briefly, the stimuli on each trial were a sequence of 12 written words or pronounceable pseudowords, presented one at a time. The words were presented as meaningful sentences, while pseudowords formed "Jabberwocky" phrases that served as a control condition.

Task and Procedure

Participants were instructed to read the stimuli silently to themselves, and also to push a button upon seeing the icon of a hand that appeared between trials. Participants performed three runs of the language localizer. Each run included 16 trials and lasted for 6 minutes. Each trial lasted for 6s, beginning with a blank screen for 100ms, followed by the presentation of 12 words or pseudowords for 450ms each (5400s total), followed by a response prompt for 400ms and a final blank screen for 100ms. Each run also included 5 blank trials (6 seconds each).

Study Procedure

The entire study required four sessions per participant. The first session was dedicated to obtaining informed consent, MR safety screening and training the participant on the tasks and eye-tracking. The first two MRI sessions were identical: T1 structural scan, 6 runs of the main experiment, 2 runs of the ventral visual localizer, and finally fieldmap acquisition. One run of the localizer was interspersed after every 2 runs of the main experiment. In the third session,

295   participants performed the final 3 runs of the main experiment, and 3 runs of the language

296   localizer.

Analysis of behavioral performance

299   We calculate accuracy as the proportion of trials with correct responses made within

300   3.85 seconds after stimulus onset. On an average of 8% of trials (SEM=2%), the participant did

301   not make a response within that window. Those trials were excluded from analyses of accuracy

302   and response time.

Preprocessing

We pre-processed our MRI data with *fMRIPrep* 21.0.1 (Esteban et al., 2019), which is

based on *Nipype* 1.6.1 (Gorgolewski et al., 2011).

<u>Structural data preprocessing</u>

For each subject, for each session, we obtained a T1-weighted structural scan. These

images were skull-stripped, corrected for intensity non-uniformity and averaged across sessions.

Cortical surfaces were constructed from the boundaries between gray and white matter using

Freesurfer's implementation (Dale et al., 1999; Reuter et al., 2010). We obtained surfaces in

subjects' anatomical space as well as the template *fsaverage* space.

<u>Functional data preprocessing</u>

Head-motion parameters with respect to the BOLD reference (transformation matrices,

and six corresponding rotation and translation parameters) were estimated before any

spatiotemporal filtering using mcflirt (FSL 6.0.5.1:57b01774, Jenkinson et al. 2002). Each

functional EPI sequence acquired during a single session was corrected for magnetic field

inhomogeneities using a B_0 nonuniformity map collected during the same session. All runs were

aligned to native anatomical (*fsnative*) and template MNI and *fsaverage* spaces using Freesurfer.

The runs were slice-time corrected (using 3dTShift from AFNI) and then co-registered to the T1-

weighted images (using bbregister from Freesurfer) (Greve and Fischl, 2009). The BOLD time-

series were also resampled onto each subject's native space cortical surface (Lanczos, 1964).

BOLD Response Estimation

To estimate the BOLD responses to each trial, we used GLMSingle in Python (Prince et

al., 2022) which optimizes the hemodynamic response function for every voxel or node in the

brain. For the main experiment, the design matrix consisted of a column for each combination

of task, stimulus and presentation location. For the functional localizer, each unique category of

stimulus received its own column. For the language localizer, the design matrix included two

columns corresponding to the sentence and jabberwocky conditions. Before estimating the

BOLD responses, the timeseries for each run were upsampled from 1.5s to 1s using an

interpolation function from pyslicetime (Kay et al., 2020). We used GLMSingle to estimate the

response to each individual trial of each condition for each surface node in each participant's

fsnative space. The algorithm found the optimal hemodynamic responses function for each

surface node, estimated cross-validated nuisance regressors computed from repetitions of the

same experimental condition, and regularized responses for each node to remove correlated

noise across single trials.

Region of Interest Definition

All regions of interest (ROIs) analyzed in this study were obtained using one of three

procedures: (1) The ventral visual localizer scan, used to define the regions depicted in color in

Figure 1B; (2) The language localizer scan, to define regions outlined in black; or (3) from an

independent study using a separate pool of subjects (White et al., 2023), for the regions

outlined in white.

We defined visual category-selective ROIs in the ventral temporal cortex by contrasting

the responses to different image categories. For each category (words, faces, and limbs), we

computed t-statistics for the contrast of that category vs. all others using a paired-sample t-test.

The t-values were visualized on each participant's inflated cortical surfaces, with the threshold

at $t \geq 3$.

We identified up to three text-selection areas (the VWFAs) in each hemisphere. VWFA-1

and VWFA-2 were in the occipitotemporal sulcus, VWFA-1 being anterior to hV4 and VWFA-2 as

a separate region anterior to VWFA-1. The face-selective areas, FFA-1 and FFA-2, were medial to

the VWFAs, within the fusiform gyrus. FFA-2 was anterior to FFA-1 but sometimes contiguous

with it. The limb-selective area was lateral to the VWFAs, defined in the occipito-temporal

sulcus (Grill-Spector and Weiner, 2014) but in some subjects contiguous with a patch on the

lateral side of the brain (encompassing what is sometimes called the extrastriate body area

(Downing et al., 2001)). The ROI for area hV4 was taken from a previous study that conducted

retinotopic mapping (White et al., 2023), defined in fsaverage space and remapped into each

individual's native surface.

We also defined two language-responsive regions in the left frontal lobe ("Broca's area").

First, using our data from the language localizer, we used a similar procedure as the one

described above, with a contrast between sentence and jabberwocky conditions. For each

individual, we defined a region with a peak in the inferior frontal sulcus, extending into the

inferior frontal gyrus as the "Language Broca's Area" (Fedorenko et al., 2010). This is the black

frontal region in **Figure 1B**. The second left frontal area we took from a previous study that

contrasted responses to written words vs. strings of shapes (White et al., 2023). This area was

defined in fsaverage space and remapped into each individual's native surface. We call "Word

Form Broca's, " and it is the white frontal region in **Figure 1B**. It was in the left precentral sulcus,

slightly posterior to "Language Broca's. The names of these two frontal language-related

regions are intended to reflected how these ROIs were defined, and not to make any strong

claims about their function. Given the variety of neuroimaging tasks the frontal operculum is

active in, it is unlikely that these regions serve any singular processing function. Finally, we also

leveraged the language localizer to define an ROI in the superior temporal sulcus (STS).

ROIs on an example subject's cortical surfaces are shown in **Figure 1B,** and all

participants are shown in **Extended Data Figure 1-2**. Details about the numbers of participants

in which we were able to identify each ROI are reported in **Table 1**.

Region of Interest	Hemisphere	Number of Subjects
VWFA1	Left	15
VWFA1	Right	9
VWFA2	Left	16
VWFA2	Right	4
Language Broca's	Left	16
Language Broca's	Right	16
FFA1	Left	16
FFA1	Right	16
FFA2	Left	14
FFA2	Right	16
OTS-Limbs	Left	16
OTS-Limbs	Right	16
STS	Left	16
STS	Right	14
MFUS-Text	Left	8
MFUS-Text	Right	2

378 **Table 1: Number of participants with each ROIs:** Number of participants (N) in whom each ROI could be defined,
379 reported separately for each hemisphere. Two other ROIs (V4 and Word Form Broca's) were defined using data
380 from White et al. 2023 and applied to all participants.
381

To create the group-level ROIs in **Figures 5-7**, we used the following approach. For the

VWFAs, we projected individually-defined contrasts between responses to text vs. all other

categories in the visual localizer to the template fsaverage space. We then concatenated these

contrasts across all subjects, and performed a one sample t-test across subjects to localize the

nodes in which responses between text and other categories were significantly different from

zero. We performed an FDR correction for multiple comparisons, and saved the nodes with

significant t-values. We used this map to define labels for VWFA-1 and -2 in the fsaverage

space. For the Language Broca's area, we used a similar approach but with the contrast

between meaningful sentences and jabberwocky in the language localizer.

Statistical Analysis

We used linear mixed effect models to determine which of our experimental manipulations significantly affected BOLD responses. The dependent variable was the mean beta response for each subject, and the fixed effects included the stimulus type, location and task. The random effects included slopes and intercepts for these variables across participants. The F-statistics and p-values reported were obtained with a type-three sum of squares table upon performing repeated-measures ANOVA using orthogonalized contrasts on the fixed effects. The t-tests for post-hoc tests are reported with FDR corrected p-values. All t-tests conducted were two-tailed. These analyses were implemented using the python package pymer4 (Jolly, 2018). We also report Bayes Factors (BFs) for each pairwise test, to quantify strength of evidence. The BF is the ratio of the probability of the data under the alternate hypothesis (that two conditions differ), relative to the probability of the data under the null hypothesis (Rouder et al., 2009). For example, a BF of 10 indicates that the data are ten times more likely under the alternate hypothesis than the null hypothesis. We computed BFs using the R package "BayesFactor" (Morey et al., 2015).

Whole-cortex Analyses

We performed several analyses of cortex-wide activity to complement our approach of individually-mapped ROIs. The goals were to create maps of the sensitivity to stimulus types or task demands. For each analysis, we first obtained the test statistic of interest for each participant in their native anatomical space (e.g., the mean difference in BOLD response magnitude for text vs. false fonts in a particular task). Next, we transformed this map of

statistics to the *fsaverage* template. Finally, for each cortical surface node in *fsaverage*, we

conducted a one sample t-test over subjects, to find which nodes had a value different from

zero. We then performed FDR correction on these p-values, and thresholded the resulting map

at p<0.05. This was all done with custom Python code, using scipy (Virtanen et al., 2020).

Functional Connectivity Analysis

The goal of this analysis was to quantify the correlations between brain regions in terms

of the *fluctuations* of BOLD response across trials, separately for each experimental condition.

Similar to previous studies (Kay and Yeatman, 2017; White et al., 2023), we evaluated across-

region correlations in the single-trial beta responses, from which low frequency drifts and other

nuisance regressors had already been removed, rather than the raw time series. To do so, for

each participant, for each surface node or ROI, we collated vectors of the responses to trials of

each condition. Conditions are defined by the combination of task, stimulus type, and stimulus

location. (Trials with real words and pseudowords were treated separately). For each condition,

we subtracted out the mean response magnitude from the vector of responses, yielding de-

meaned "residuals".

We then analyzed across-region correlations in these residual responses in two ways.

First, we conducted whole-brain analyses of functional connectivity to a "seed region," in each

individual's native cortical surface. We first averaged responses over the nodes within the seed

region, and then we calculated the Pearson correlation coefficient between the seed's residuals

and the residuals from every other surface node. We then projected the resulting correlation

coefficients from each subject's native space to the *fsaverage* space. With all 16 participants'

correlation maps in the same space, we averaged the correlation coefficients and at each node

computed a one-sample t-test to assess whether the correlation was significantly different from

0. Using the FDR-corrected p-values for each vertex as a threshold, we obtained a map of nodes

that were significantly correlated with the seed region during each condition. We performed

these analyses with the scipy and statsmodels libraries (Virtanen et al., 2020).

Second, we assessed how functional connectivity between pairs of ROIs changed across

experimental conditions. In Figure 6B, we focused on the correlated activity between the

VWFAs and the "Language Broca's area", given the result of the whole-brain analysis and prior

evidence (White et al., 2023). The correlations between residuals were calculated on data first

averaged across nodes within the two ROIs in each individual's native brain, separately for each

condition. We then averaged the correlation coefficients across subjects. For these measures

we conducted similar statistical tests as for the BOLD response magnitudes (described above).

Results

Behavior

The participant's task on each trial was to push one of two buttons to report whether or

not they saw a 'target': a real word in the lexical decision task; red letters in the stimulus color

task; or a brief change in the fixation dot's color from gray to red. In both color tasks, an

adaptive staircase continuously adjusted the saturation of red color to each participant's

detection threshold. The character string's position varied randomly across trials, either

centered on the fixation mark, 3º left, or 3º right. The stimulus types were randomly

intermingled, and the task conditions were blocked.

Figure 1D plots mean task accuracy in each condition. Using linear mixed models with

subjects as random effects, we analyzed accuracy as a function of the task, stimulus type, and

stimulus location. We found a main effect of all three experimental manipulations (Task:

$F_{(2,33)}=37.77$, $p<0.001$, Stimulus Type: $F_{(2,23)}=109.84$, $p<0.001$, Stimulus Location:

$F_{(2,46)}=6.12$, $p=0.004$). There were also and two-way interactions between task and stimulus

type ($F_{(4,375)}=117.09$, $p<0.001$), and between task and location ($F_{(4,375)}=9.8$, $p<0.001$).

Accuracy was highest in the stimulus color task, and lowest in the fixation color task (all pairwise

comparisons $p<0.03$). In the lexical decision and stimulus color tasks, accuracy was higher when

the stimuli were presented at fixation (0º) than ±3º in the parafovea, but the opposite was true

in the fixation task. In other words, fixation task accuracy relatively impaired by characters

appearing at fixation (**Figure 1D**). Post-hoc tests (corrected for false discovery rate) revealed

better lexical decision performance when real words were presented in the right compared to

the left hemifield ($t=2.31$, $p=0.03$), consistent with prior results. Reaction times are plotted in

Figure 7.

The VWFA prefers text over unfamiliar characters even when attention is withdrawn

The mean BOLD responses in the left VWFA are shown in **Figure 2.** In all our analyses, we

found that the two main subregions of the left VWFA (VWFA-1 and VWFA-2) had very similar

response patterns. We therefore average across them here. Results for the separate regions are

reported in **Extended Data Table 2-1.** A third ventral text-selective region, MFUS-text, is

summarized in **Figure 3F.**

The first result concerns the bottom-up selectivity for text in the VWFA. This region was

defined as a patch of cortex that, in an independent localizer scan, responded more strongly to

attended words than to false fonts, faces, and limbs (during a one-back task). We therefore

expected that the VWFA would also prefer words and pseudowords over false font strings in

the main experiment. But other researchers have argued that this region appears to have

selectivity for text due to top-down influence from cerebral networks involved in language and

attention (e.g., Dębska et al 2023; Price & Devlin, 2011). Such an account predicts that during a

non-linguistic task in which attention is totally withdrawn from the stimuli, the VWFA would

respond equivalently to words as to meaningless shapes that have similar visual features.

To test that prediction, we compared the response to text (words and pseudowords

together) vs. false font strings during the fixation color task. In that task, the stimuli were task-

irrelevant and the participant was focused on detecting a threshold-level color change in the

fixation dot. They did not have time to switch attention from the dot to the character strings

that flashed briefly at the same time. Nonetheless, the left VWFAs responded more strongly to

text than false fonts during this fixation task (mean betas = 0.23 vs 0.14; t(15)=4.82, p = 0.0002,

BF=139.87). These data are the two open bars in **Figure 2A.** This text selectivity was maintained

even when the stimuli were presented 3º from the focus of attention at the fixation dot, as

shown in the bottom right panel of **Figure 2D**

We infer that the VWFA has inherent selectivity for a certain category of shapes: strings

of familiar letters. That selectivity is minimally dependent on attentional engagement with the

stimuli or on voluntary linguistic processing. Importantly, however, the selectivity for text was

absent in the FFAs, limb-selective areas, and V4 (see **Figure 3**).

Figure 2: The VWFA's activity reflects an interaction of stimulus features and task demands. (A) Mean percent signal change (p.s.c.) as a function of stimulus type and task condition in left hemisphere VWFAs. "Text" includes both real words and pseudowords (which are separately analyzed in panel **C**). Error bars represent ±1 SEM. Horizontal bars and stars indicate statistical significance for the main effect of stimuli and pairwise t-tests between tasks. **(B)** Task effects computed as the mean difference in percent signal change from the fixation task as baseline. Error bars indicate *95% confidence intervals.* Horizontal bars and stars indicate a significant difference in the effect of engaging in the lexical task (compared to fixation task) on responses to text vs. false fonts. The effect of the stimulus color task did not differ across stimulus types. For task differences in individual participants, see Extended Data Figure 2-1. **(C)** Responses to words vs. pseudowords in the three tasks. The lexicality effect is only observable during the lexical decision task. Error bars are ±1 SEM. **(D)** Responses in the VWFAs as a function of stimulus position, separately for each task and stimulus type (green = text; purple = false fonts). Error bars are ±1 SEM.

The lexical decision task enhances the VWFA's response to text and suppresses its response to false fonts

The next result concerns how the VWFA's activity changed when the participant engaged in the lexical decision task, which required them to attend to each stimulus and judge whether it was a known English word. To start, we found a significant interaction between stimulus type and task on mean BOLD response magnitudes ($F(2, 108)=34.34$, $p=3\times10^{-12}$). As shown in **Figure 2A-B**, the effects of the task differ across stimulus types. To quantify that interaction, for each stimulus type we calculated comparisons of the responses between pairs of tasks. The statistics are reported in **Table 2**. The mean differences from the fixation task baseline are shown in **Figure 2B,** with error bars that represent 95% confidence intervals.

First, compared to when the stimuli were ignored during the fixation color task, the VWFA's response to text (real words and pseudowords) was enhanced during the lexical decision task, but not during the stimulus color task. The dark green bar in Figure 2B shows the enhancement in the lexical task. The mean response during the lexical decision task was 1.6 times the mean response during the fixation color task. All individual participants showed this effect.

Second, for false fonts, we observed the opposite task effect: responses during the

lexical decision task were *lower* compared to the other two tasks. See the dark purple bar in

Figure 2B. This result suggests that VWFA's response to non-preferred stimuli is suppressed

below the sensory baseline level while the brain is engaged in linguistic processing. 12 out of 16

participants showed lower responses for false fonts during the lexical decision task, compared

to the fixation color task baseline. The mean response to false fonts in the lexical task was 54%

of the mean response to false fonts in the fixation task. To characterize the consistency of these

effects, we refer the reader to **Extended Data Figure 2-1**, which compares the VWFA responses

for each task and stimulus type within individual participants.

ROI	Stimulus	Task	Mean ΔPSC	SEM	95% CI	t-value	BF	p-value
VWFA	Text	LD - FC	0.13	0.02	0.08, 0.18	6.85	242.22	<0.01
		LD - SC	0.1	0.02	0.06, 0.14	6.41	391.07	<0.01
		SC - FC	0.03	0.02	0.02, 0.07	1.55	1.05	0.13
	False Fonts	LD - FC	-0.06	0.02	-0.12, -0.003	2.58	8.44	0.02
		LD - SC	-0.09	0.02	-0.14, -0.04	4	26.95	<0.001
		SC - FC	0.03	0.02	-0.03, 0.08	1.15	1.79	0.25
Word Form Broca's	Text	LD - FC	0.11	0.02	0.07, 0.15	6.66	884.14	<0.001
		LD - SC	0.09	0.02	0.06, 0.13	6.05	334.98	<0.001
		SC - FC	0.01	0.02	-0.02, 0.05	0.91	0.88	0.37
	False Fonts	LD - FC	-0.1	0.02	-0.15, 0.04	4.27	56.96	<0.001
		LD - SC	-0.11	0.02	-0.17, -0.06	5.27	156.28	<0.001
		SC - FC	0.02	0.02	-0.03, 0.07	0.9	0.51	0.37
Language Broca's	Text	LD - FC	0.07	0.01	0.04, 0.10	5.73	83.58	<0.001
		LD - SC	0.07	0.01	0.04, 0.10	6.02	544.78	<0.001
		SC - FC	0	0.01	-0.03, 0.03	0.03	0.26	0.98
	False Fonts	LD - FC	-0.05	0.02	-0.09, 0.004	2.66	4.44	0.01
		LD - SC	-0.06	0.02	-0.10, -0.02	3.47	13.49	0.002
		SC - FC	0.01	0.02	-0.03, 0.05	0.75	0.37	0.46

540   **Table 2: Mean task effects on percent signal change:** Across pairs of tasks, separately for text and false font stimuli,
541   in the left VWFA and the two Broca's areas. See Extended Data Table 2-1 for statistics on VWFA-1 and VWFA-2

542 separately. For each ROI, the six p-values are corrected for false discovery rate. Abbreviations: PSC = Percent Signal
543 Change, SEM = Standard Error of Mean, CI = Confidence Interval, LD = lexical decision task; FC = fixation color task;
544 SC = stimulus color task; BF = Bayes Factor.
545

Moreover, **Figure 2C** shows that the VWFA responds differently to real words and

pseudowords, but only during the lexical decision task (word type by task interaction:

$F(2,45)=3.5$, $p=0.038$). Consistent with prior studies, pseudowords evoked larger responses than

real words during the lexical decision task ($t(15)=2.8$, $p = 0.01$, BF=4.58). That may be because

processing pseudowords takes more time than processing familiar words (Vinckier et al., 2007;

Cohen et al., 2008). However, that *lexicality effect* is absent when the stimuli are ignored during

the fixation task ($t(15)=0.6$, $p = 0.56$, BF=0.3). This result suggests a key role for top-down input

in creating the VWFA's sensitivity to higher-level lexical features of the stimuli (in contrast to the

selectivity for letter strings, which was strong in all tasks).

Figure 2D shows that VWFAs are also sensitive to the visual field position of the words

(Rauschecker et al., 2012; Le et al., 2017; White et al., 2023). In general, the VWFA prefers

stimuli at the fovea (0º) over stimuli presented 3º to the left or right (main effect of location:

$F(2,24)=29.7$, $p < 0.001$). However, that spatial sensitivity is greatly reduced for text presented

during the lexical decision task. In **Figure 2D**, compare the dark green line on the left to the

lighter green lines on the right. During the lexical task, there were no significant differences in

pairwise comparisons of stimulus positions (all $t<1$, $p>0.7$), because the enhancement of

attended words was strongest at ±3º. Thus, voluntary effort to read enhances the response

especially when words are in the parafovea where legibility is reduced (Vinckier et al., 2007; Kay

and Yeatman, 2017). Differences between the lexical decision and fixation color tasks were also

present when the stimuli were presented foveally, for text (t(15)=3.75, p=0.0007, BF=43.76) as

well as for false fonts (t(15)=2.69, p=0.02, BF=3.59).

Top-down modulations do not occur automatically when words are visually attended

Word recognition is sometimes described as an "automatic" process (Stroop, 1935; Price

et al., 1996; Brown et al., 2002; Ruthruff et al., 2008; Pattamadilok et al., 2017; Joo et al., 2021).

The Stroop effect is the most famous evidence: participants cannot stop themselves from

processing the meaning of a word when instructed to say aloud the color of its ink. An

automaticity-based account would predict that attending to words to judge the font color would

engage the VWFA just as strongly as when voluntarily reading the words. However, we found

that the VWFA responded quite similarly during the stimulus color task as when the words were

ignored during the fixation task. The light bars in **Figure 2B** represent the mean differences

across those two tasks for text (green bar) and false fonts (purple bar). While engaging in the

lexical decision task greatly increased the response to text, engaging in the stimulus color task

had a negligible effect (not significant after correcting for multiple comparisons; BF=1.05). The

response to false fonts was not suppressed in the stimulus color task, but rather slightly and not

significantly increased (BF=1.79; see **Table 2**).

One potential criticism of the comparison between the lexical and stimulus color tasks is

that the latter could be accomplished by attending to only one letter. However, the character

string was presented for only 150 ms at an unpredictable location, and with the saturation of

the red color set to detection threshold. The optimal strategy therefore was to integrate the

color signal across all the letters.

Thus, the elevation of the VWFA's response to words does not occur automatically when

the words are attended, but requires engagement in an explicitly lexical task. The same applies

to the relative suppression of the response false fonts.

The task and stimulus interactions are absent from other occipital and temporal

regions

We analyzed responses in six other regions of interest in the left occipital and temporal

lobes. See **Figure 3.** Within the ventral occipito-temporal cortex, we analyzed V4, face-selective

regions in the posterior and middle fusiform gyrus (FFA-1 and FFA-2, respectively), a limb-

selective region (OTS-Limbs), and another text-selective region within the mid-fusiform gyrus

(MFUS-Text, which was only detectable in half of our participants). Additionally, we analyzed

responses in a large swath of the superior temporal sulcus (STS) that was activated in the

language localizer.

We did not find a significant effect of task, stimulus type, or an interaction between task

and stimuli within any of these ROIs. Thus, the pattern of activity we found in the VWFA is

unique among the visual regions we analyzed and also absent in the STS. Statistics for all ROIs

are reported in the **Extended Data Figure 3-1**. FFA-2 had a pattern that was qualitatively similar

to the VWFA, but the task effects were less consistent across subjects and not statistically

significant (all ps>0.5).

The only marginally significant effect was observed in the left MFUS-Text (**Figure 3F**),

where responses to text were somewhat higher in the lexical task than the other two tasks

(p=0.07). Left MFUS-text also responded more strongly to text than false fonts overall, but not

significantly after correcting p-values across the ROIs, perhaps because we were able to localize

this region in only 8/16 participants. Right hemisphere regions, including VWFA, are shown in

Extended Data Figure 3-2. There were no significant effects of stimulus or task in any of these

regions.

Figure 3: Task and stimulus effects are absent in other regions of left occipital and temporal lobes. Each panel
plots mean BOLD responses in one ROI, formatted as in Figure 2A. See Extended Data Figure 3-1 for statistics on
these control regions and Extended Data Figure 3-2 for right hemisphere ROIs.
617

Broca's area shows strong task-dependent enhancement and suppression

We also localized two language processing regions in the left frontal cortex. The first we refer to as "*Language Broca's*," because it was defined in each individual by an independent language localizer (see Methods) (Fedorenko et al., 2010; Mahowald and Fedorenko, 2016). It was located in the inferior frontal sulcus, sometimes extending into the inferior frontal gyrus. The second region we refer to as "*Word Form Broca's*" because it was defined (in an average brain) from a previous study contrasting the response to written words and non-letter shape strings (White et al., 2023). It was slightly posterior to Language Broca's, in the precentral sulcus. A similar contrast on our main experiment, contrasting text stimuli with false fonts, produced an overlapping region. We used the Word Form Broca's from the previous study to avoid using the same data for defining an ROI and analyzing mean responses. The locations of the two Broca's regions relative to each other are depicted in **Figure 1B** in an example subject, and on the average brain in **Figure 5B**.

Figure 4 shows that in both these Broca's area sub-regions, we found interactions between stimulus type and task that went in the same direction as in the VWFAs. However, response magnitudes in the more anterior "Language Broca's area" were different: it did not respond positively to stimuli overall (**Figure 4A**). Its response differed from baseline only during the lexical decision task, but in opposite directions for text vs. false fonts: a strong increase for text, and a decrease below baseline for false fonts (**Figure 4C**). These differences between the lexical decision task and the fixation color task were observed for text in 15 out of 16 participants, and for false fonts in 13 out of 16 participants. It also showed sensitivity to lexicality (pseudowords > real words) only during the lexical task (**Figure 4E**).

640   The response magnitudes in Word Form Broca's were more similar to the VWFA, with

641 generally positive trial-evoked responses above baseline (**Figure 4B**). It showed the same task

642 effects: a selective enhancement for text and suppression for false fonts during the lexical

643 decision task (**Figure 4D**). In this region, we observed the task effects for text in all participants,

644 and for false fonts in 14 out of 16 participants. It also responded more strongly to pseudowords,

645 except during the fixation task (**Figure 4F**). Statistics for the task effects in these two regions are

646 reported in **Table 2**.

647   One interpretation of these data is that the "Language Broca's area" functions as a

648 *control region* that applies positive or negative modulations to other parts of the reading

649 network. This region is engaged only during the lexical task. If letter strings are presented, its

650 activity rises and it in turn excites other regions involved recognizing words: the VWFAs as well

651 as the more posterior Word Form Broca's. If it detects false fonts during the lexical task, its

652 activity decreases below baseline and the other regions are suppressed as well.

Figure 4: Task and stimulus effects in two left frontal regions. (A, B) Mean percent signal change as a function of stimulus type and task in two frontal regions in the left hemisphere. Format as in Figure 2A. (C, D) Difference in mean percent signal change from fixation task baseline. Format as in Figure 2C; error bars indicate *95% confidence intervals.* (E, F) Responses to real words vs. pseudowords in each task (format as in Fig 2E).

A whole-cortex analysis of the task-by-stimulus interaction highlights Broca's area and the left VWFAs

The preceding analyses were based on small regions of interest localized precisely in each individual's brain from independent data. This was necessary given the documented shortcomings of first morphing to an average brain. However, we also investigated whether our

primary findings are widespread across the cortex or in fact unique to the primary regions of

interest.

A key result in the VWFA and Broca's ROIs was an interaction between stimulus type

and task: the response to text was stronger in the lexical task than in the fixation color task, but

the response to false fonts was stronger in the fixation task. **Figure 5** shows a whole-cortex map

of this interaction between stimulus type (text vs false fonts) and task (lexical decision vs

fixation task).

Four regions demonstrated a significant interaction: (1) the left occipitotemporal sulcus,

where the VWFA is located; (2) portions of the precental sulcus and inferior frontal sulcus

(bilaterally), where, in the left hemisphere, our "Broca's" ROIs are located; (3) the anterior

insula / medial frontal operculum (bilaterally), and (4) a small portion of the left intraparietal

sulcus (IPS). The first two of those regions overlap closely with our pre-defied ROIs (see Figure 2

and Figure 4), although the right hemisphere homologue of 'Broca's area' did not.

The latter two regions with a significant interaction were not pre-defined ROIs, but they

have previously been implicated in reading. The anterior insula is often highlighted in fMRI

studies of language function (Oh et al., 2014) although in some cases activity localized there

may originate in the overlying frontal operculum (Fedorenko et al., 2015; Woolnough et al.,

2019, 2023). The IPS region is also involved in cognitive processes relevant to reading (Rapp et

al., 2016; Forseth et al., 2018; Woolnough et al., 2022) and is functionally correlated with the

VWFA and Broca's area (Vogel et al., 2012; White et al., 2023), as shown below.

Figure 5: Whole-cortex analysis of the interaction between task and stimulus type. (**A**) Surface map (in the fsaverage template) of the interaction between stimulus type (text - false fonts) and task (lexical decision - fixation color). For each stimulus type, we computed differences in responses during the lexical decision task and the fixation color task. We then computed a difference of these difference maps for the two stimulus types. The resulting contrast map shows all nodes that had t-values > 3. Black outlines represent the patches with FDR-corrected p<0.05. (**B**) Lateral view of the left hemisphere, with the same interaction contrast as the colored blobs (corrected p<0.05). The outlines represent the ROIs analyzed in Figure 4: the two regions outlined in black together form "Language Broca's area" and the more posterior area in whit is "Word Form Broca's".

Modulations of VWFA activity are accompanied by task-dependent functional

connectivity with the language network

To investigate how the areas that are modulated by reading task demands may be

communicating with each other, we performed a functional connectivity analysis. Specifically,

we correlated the *fluctuations* in trial-evoked responses, after subtracting out the mean

response in each condition (White et al., 2023). For each surface vertex, we averaged the

responses for each unique combination of task, stimulus type (real words, pseudowords and

false fonts) and stimulus location. We then removed this mean response from the responses to

all trials of the same unique condition, and performed the subsequent analysis with these

demeaned *residual* responses. Specifically, we computed correlations coefficients in these

residual responses across regions, separately for each condition.

Figure 6A shows a whole-brain map of correlations between each surface vertex and the

left Language Broca's area as the seed region. This represents functional connectivity during the

lexical task (all stimulus types). Autocorrelation within the seed region during the lexical

decision task was r=0.50. We observed positive correlations between the Language Broca's area

and much of the left superior temporal lobe, left intraparietal sulcus (IPS), and bilateral occipito-

temporal sulcus, where the VWFAs are. There was also a hot spot of connectivity with a

homologous 'Broca's area' in the right frontal cortex. This map is consistent with prior results

(White et al., 2023). **Figure 6C** shows analogous analyses with Word Form Broca's area, VWFA-

1, VWFA-2 and as the seed regions. The whole-brain maps of connectivity with the left VWFAs

looked quite similar to the map in **Figure 6A** (with the seed in Broca's area), but with somewhat

less correlated activity in the superior temporal lobe. We also analyzed activity in anatomically-

defined Brodmann's areas 44 and 45, which are typically associated with "Broca's area." Activity

there, in terms of BOLD responses and function connectivity patterns, resembled what we

observed in "Word Form Broca's."

In addition to whole brain functional connectivity, we also computed mean correlations

between the left VWFA and left Language Broca's area, separately for each task and stimulus

condition. For this analysis, we first averaged the single-trial beta responses across all nodes

within each ROI, which reduces noise in the subsequent correlations. The results are in **Figure**

6B. Of primary interest here is how the functional connectivity between the VWFA and Broca's

area significantly *changed* depending on the task demands and stimulus content. We found a

main effect of task ($F_{(2,15)}=21.63$, $p < 0.001$) and stimulus type ($F_{(2,15)}=4.87$, $p = 0.03$) on the

correlation coefficients, and an interaction between them ($F_{(2,396)}=3.09$, $p = 0.04$). We

observed highest correlations between the two regions during the lexical decision task, for both

types of stimuli, and lowest correlations during the fixation task. The correlation between these

two regions in their responses to text were roughly twice as strong during the lexical task as

during the fixation task. Statistics on these comparisons are reported in **Table 3**.

Stimulus	Task	Mean Δ r	SEM	95% CI	t-value	BF	p-value
Text	LD - FC	0.18	0.03	0.11, 0.25	6.33	3.21×10^6	< 0.001
	LD - SC	0.17	0.04	0.08, 0.26	4.91	4.64×10^5	< 0.001
	SC - FC	0.01	0.03	-0.07, 0.09	0.3	0.12	0.75
False Fonts	LD - FC	0.15	0.04	0.05, 0.24	3.68	49.97	0.001
	LD - SC	0.05	0.03	-0.05, 0.16	1.24	0.40	0.22
	SC - FC	0.09	0.04	0.01, 0.2	2.09	1.29	0.06

Table 3: Functional Connectivity between Broca's region and VWFA: Statistics on how functional connectivity
between the language Broca's Area and the left VWFA (collapsing over VWFA-1 and VWFA-2) differed across pairs
of tasks. This is calculated separately for trials with text stimuli (both words and pseudowords) and false fonts.

"Mean Δ r" is the mean difference in the correlation coefficient r across the two tasks listed in each row. P-values
are FDR-corrected for multiple comparisons. Abbreviations: LD = lexical decision task; FC = fixation color task; SC =
stimulus color task.
738

It is noteworthy that functional connectivity between the VWFA and Broca's *increased*

during the lexical task compared to the fixation task, even when false fonts were presented.

This goes in the opposite direction as the mean overall response magnitudes, which were

decreased for false fonts during the lexical task (**Figure 2A-B)**. This suggests that the

suppression of response magnitudes to false fonts in the VWFAs could be caused by

communication with Language Broca's area.

745
Figure 6: Functional connectivity shows task-dependent communication between Broca's area and the VWFA. (A)
Whole-brain connectivity with the seed region in left language Broca's area (black outline) during the lexical
decision task for all stimulus types. The color map represents Pearson's correlation coefficients (minimum r>0.11,
p<0.05, FDR corrected). The two black outlines on the left ventral surface represent the peaks of the contrast
between words and all other categories in the category localizer experiment, in the average brain. Thus, they
roughly correspond to the most likely locations of VWFA-1 and VWFA-2. (**B**) Average correlation coefficients
between left language Broca's area and the left VWFA (collapsing VWFA-1 and VWFA-2). We first averaged
responses for each condition within each of the two ROIs, de-meaned them and then computed pairwise
correlation coefficients. The correlations were highest during the lexical decision task for both types of stimuli.
Error bars are +/- 1 SEM. Horizontal lines and stars indicate statistical significance. (*=p<0.05, **=p<0.01,
***=p<0.001). (**C**) Correlation maps like in panel A, but for three different seed regions: Word Form Broca's Area,
left VWFA-1, and left VWFA-2.
758

Task-driven correlations in BOLD responses and reaction times

Response times (RTs) are known to correlate with BOLD responses in several cerebral

networks (Yarkoni et al., 2009; Domagalik et al., 2014; Mumford et al., 2024). The simplest

explanation is that RTs indicate "time on task," or the duration of time that the brain spends

processing a stimulus. If the neural activity lasts for more time, the BOLD signal (which

sluggishly integrates neural activity over time), yields a response of greater magnitude.

We conducted several analyses to determine whether a general effect of task difficulty

on BOLD responses can explain the primary results reported above. First, we conducted a

whole-cortex analysis of the correlation between BOLD response and RTs on single trials. The

result is shown in the top row of **Figure 7A:** the mean correlation coefficients between

normalized RTs and single-trial beta responses, for all stimulus types during the lexical decision

task. These two measures were both de-meaned within each experimental condition (stimulus

type and location) before entering into the correlation. The regions with significant correlation

overlap partially with the regions highlighted as having functional connectivity with Broca's area

and the VWFA (**Figure 6A and 6C**), including left inferior frontal gyrus, insula/frontal operculum,

pre-central sulcus, intraparietal sulcus, and occipito-temporal sulcus. In the right hemisphere

(not shown) during the lexical decision task we observed hotspots of correlations with RT in the

insula/frontal operculum and anterior cingulate cortex.

The bottom row of **Figure 7A** is the same analysis but for trials during the fixation task,

when the same stimuli were presented and participants also had to make a response on each

trial. Compared to what was observed during the lexical task (top row), the correlations

between BOLD and RT were markedly weaker in the vicinities of Broca's area and the VWFA (left

inferior frontal sulcus and occipito-temporal sulcus). Thus, to the extent that there is a relation

between the difficulty of each trial and the activity in these regions, it is not domain-general but

rather specific to the task that requires reading the words.

 Figure 7: Relationship between reaction times and BOLD responses (A) Whole-cortex correlations between
response times and magnitude of BOLD response. The color map represents Pearson's correlation coefficients
(minimum r>0.03, q<0.05). The data in all panels corresponds to trials with correct behavioral responses only. The
top panel shows correlations for the lexical decision task, and the lower panel for the fixation color task. **(B)** Mean
left VWFA response magnitudes as a function of mean response times in each stimulus and task condition (correct
trials only). Error bars represent ±1 SEM. The top panel averages over all stimulus locations, and the bottom is only
for trials with stimuli at the central (foveal) location.

The second analysis concerns the relation between RTs and activity in the left VWFA

specifically, across all conditions. **Figure 7B** plots the mean BOLD response in the left VWFA as a

function of the mean RTs in each condition, on correct trials only. The top graph is for all

stimulus locations, and the bottom graph is for trials when the stimuli were presented at the

fovea (0º). The three stimulus types are represented by different shapes, and the three tasks are

represented by different shadings. In the top graph, there is clearly a positive correlation

between RT and BOLD. This may provide an explanation for the VWFA's responses during the

lexical decision task: false fonts have the smallest RTs and smallest BOLD responses, whereas

pseudowords have the largest RTs and largest BOLD responses. This could be because when a

false font string is presented, the participant, and their VWFA, quickly recognizes that it is not a

word and stops processing it. In contrast, pseudowords look plausibly like real words, so the

brain must effortfully search through the mental lexicon before deciding to reject the stimulus.

Thus, correct RTs are longer and the BOLD response is higher for pseudowords, compared even

to real words, which can be identified relatively quickly (Woolnough et al., 2021; Gagl et al.,

2022). However, Taylor et al. (2013) demonstrated that even when regressing out the effect of

RT, pseudowords still evoked a larger response in the VVFA than real words.

We also used a model comparison approach to demonstrate that RTs, by themselves, do

not explain all the variance in the BOLD responses. We first constructed a model that included

only response times as a fixed effect predictor, with random effects of response time, stimulus

type and task across subjects. We then constructed a second model with the same random

effects, but added in stimulus type and task and their interactions as additional fixed effects. We

found that the second model fit the VWFA better than the first (χ^2(10) =112.64, p<0.001), even

when penalizing the second model for its greater number of parameters (BIC 9268.3 for the

second model compared to 9289.2 for the first; lower BIC values indicate better model fit). The

weaker fit of the RT-only model is made clear by how the VWFA responds during the fixation

color task (open symbols in Figure 7B). During this task, there is no clear relationship between

RT and BOLD magnitude across stimulus types. Moreover, RTs were generally longer in the

fixation task than the stimulus color task, but BOLD response magnitudes were similar between

them.

Also, if we consider only trials when the stimulus was presented at fixation (excluding

stimuli 3º to the side), the influence of RT is reduced but the task and stimulus effects are still

strong. That is shown in the lower plot in **Figure 7B**. Again, the model that included task and

stimulus type as predictors in addition to reaction time fit the data better than one that only

included reaction times (χ^2(10) =40.82, p<0.001; BICs 3117.2 vs. 3138.0, respectively). Thus, we

conclude that much of the shared variance between RTs and BOLD that produced the pattern

the upper plot was explained by the increased difficulty when the stimuli are in the parafovea.

Without that difficulty, BOLD is less related to RT but still strongly influenced by the interaction

of task and stimulus.

Discussion

Discussion

We investigated the combination of stimulus features and task demands that maximally

engage the VWFA. Consistent with the hypothesis that this region is a module tuned for a

special class of stimulus, we found that it responds more strongly to words than strings of

unfamiliar characters even during a demanding visual task that withdraws attention from the

stimuli. Nonetheless, the results show that the VWFA's activity can neither be solely accounted

for by stimulus selectivity, nor by top-down modulation. The strongest activation requires

specific stimuli (strings of familiar letters) presented during specific tasks (explicitly trying to

read words). Engaging in the lexical task increased the VWFA's BOLD response to words and

pronounceable pseudowords (with an even greater increase for pseudowords), but *suppressed*

the response to visually-matched unfamiliar characters (false fonts). Thus, top-down language

processing does not simply *add* to the VWFA's BOLD activity; rather, the modulation differs

greatly across types of character strings. Moreover, compared to when the stimuli were

ignored, attending to either type of character string to judge its color had little impact on the

VWFA's response.

This interaction between bottom-up and top-down factors was unique to the VWFA

among visually-driven areas. Given the patterns of functional connectivity, which also differed

greatly across tasks, we suggest that the modulations targeted to the VWFA are due to

interactive feedback from regions involved in processing language.

Possible explanations for what drives activity within the VWFA

By fully crossing task demands and stimulus types, our results rule out four hypotheses

about what determines the magnitude of VWFA responses:

(1) Word-likeness of visual stimuli: One hypothesis is that the VWFA's activity is explained

by bottom-up tuning for visual stimuli that are the most like known words. That predicts, in our

study, that responses should be weakest for false fonts, intermediate for pseudowords, and

highest for real words, in all three tasks. Our data only partly match that prediction. The VWFA

did prefer real letter strings over false fonts across all tasks. But that selectivity was greatly

magnified during the lexical task (compared to both other tasks). More importantly, the

stronger response to pseudowords than real words occurred only occurred during the lexical

decision task (**Figure 2C**). These results suggest that the VWFA's sensitivity to *lexical* attributes of

letter strings is contingent on top-down cognitive influences during linguistic tasks.

(2) Top-down boost for attended stimuli: Throughout visual cortex, task-relevant stimuli

evoke larger responses than task-irrelevant or ignored stimuli. These attentional effects could be

described as additive or multiplicative modulations of the stimulus-driven response, with a

magnitude that depends on the degree of similarity between the tuning properties of neurons

and what is attended (Reynolds and Heeger, 2009). A related hypothesis for the VWFA is that

attentional mechanisms boost responses to task-relevant stimuli, but only when the stimuli are

text. That hypothesis would predict a higher response to words when they are task-relevant

than when they are ignored, and no modulation of the response to false fonts.

Our data violate this prediction in two ways. First, the VWFA responded similarly to

attended words during the stimulus color task and to ignored words during the fixation color

task (light green bar in **Figure 2B**). Second, the response to false fonts was *lower* when they

were task-relevant during the lexical decision task than when they were ignored (dark purple

bar in **Figure 2B**). Together, these results show that the top-down modulations in the VWFA

cannot easily be explained by visual attention, consistent with White et al. (2023).

(3) *Automaticity of word recognition:* This hypothesis implies that as long as a word is

legible and some attentional resources are available, its meaning is fully processed (Carr, 1992;

Brown et al., 2002). In its simplest form, such a theory would predict equivalent responses to

written words in the lexical decision and stimulus color tasks. In contrast, we found that

engaging in the stimulus color task had a negligible effect compared to the fixation task.

Therefore, strong VWFA activation requires voluntary effort to read the words.

(4) Task difficulty: This hypothesis is that the VWFA, along with other regions, responds

more strongly on trials when the participant makes a more difficult judgment that requires

more processing time. Several studies of word reading have found correlations between the

behavioral RTs and BOLD magnitudes in the VWFA as well as a wider network of regions (Binder

et al., 2005; Graves et al., 2010; Taylor et al., 2013).

In our data, two patterns suggest that this hypothesis cannot fully explain the

modulations of VWFA activity. First, in the VWFA and Broca's area, the correlation between RT

and BOLD is not a *general* effect of the effort required on each trial. Rather, it is much stronger

during the lexical decision task than the fixation task (see the maps in **Figure 7A**). Second, VWFA

responses are best predicted by a model that includes task and stimulus conditions in addition

to mean RTs (see **Figure 7B**). Nonetheless, during the lexical decision task there is a relation

between BOLD and RT, in terms of the differences between pseudowords, real words, and false

895 fonts. We believe that this is due to a top-down signal that prolongs the VWFA's processing of

896 pseudowords and shuts down processing of false fonts.

898 Our proposal for the role of the VWFA and its functional connectivity

899   Our results support the theory that the VWFA is primarily a visual region devoted to

900 identifying familiar letter strings (Vinckier et al., 2007; Dehaene et al., 2015). Its bottom-up

901 responses are selective for familiar letter strings, even when attention is withdrawn. However,

902 the VWFA is not automatically engaged in word recognition when familiar letter strings are

903 presented and attended. Its selectivity for text is greatly enhanced when the task requires

904 reading, in part due to active suppression of non-letter stimuli. Also, its sensitivity to lexical

905 features of stimuli depends critically on the participant engaging in an explicitly linguistic task.

906   These aspects of its function are dependent on communication with other parts of the

907 language network, as our functional connectivity analyses revealed (**Figure 6;** see also (Vogel et

908 al., 2012; Stevens et al., 2017; Chen et al., 2019; López-Barroso et al., 2020). We also discovered

909 higher correlations between the VWFA and Broca's area during the lexical decision task than the

910 other two tasks. This task-dependent modulation of connectivity was unique to the VWFA

911 within ventral temporal cortex (White et al., 2023). Other researchers have also reported that

912 the connections between VWFA and frontal regions change as a function of familiarity with a

913 language (Boukrina et al., 2014).

914   Importantly, correlated activity with Broca's area was strong both when the VWFA's

915 response magnitudes were enhanced (for text) and when they were suppressed (for false fonts

916 during the lexical decision task). The mean BOLD responses in the seed region used for this

analysis, which we called Language Broca's area, offer clues to the nature of its role. Its BOLD

signal deviated from baseline only during the lexical decision task, with a positive response to

letter strings and a negative response to false fonts (**Figure 4A**). This suggests that the

suppression of the VWFA response to false fonts during the lexical decision task is due to top-

down feedback from Broca's area.

Limitations and future directions

Methods that offer greater temporal resolution, such as magnetoencephalography or

intracranial recordings, will be necessary to reveal *when* the task effects arise during the

processing of a word. Past research gives some clues: intracranial recordings from ventral

temporal cortex showed that responses to attended words differed from ignored words

relatively late, ~400-800 ms after word onset (Nobre et al., 1998). However, the time course of

attention effects may also depend on the task, as engaging in an orthographic task has also

been shown to modulate an earlier evoked potential (the N200; Ruz and Nobre, 2008).

By using fMRI, we were able to demonstrate the spatial specificity of cognitive

modulations in the VWFA, which did not occur in nearby category-selective regions.

Nonetheless, there may be analogous cognitive effects in those other regions during tasks

optimized for them, such as face recognition in the FFA.

Lastly, there is more to learn about differences between word-selective sub-regions. We

found similar activity patterns in VWFA-1 and VWFA-2. While some previous studies have

reported the same (White et al., 2023), others have emphasized the importance of functional

subdivisions of the VWFA (Lerma-Usabiaga et al., 2018). Similar questions remain about sub-

regions in the vicinity of "Broca's area." Other researchers have speculated about functional

segregation within this general region along the dorsal-ventral axis, as well as the anterior-

posterior axis (Hagoort, 2014).

Conclusions

We conclude that activity in the VWFA is determined by an interaction between inherent

stimulus selectivity and top-down input from frontal language regions. The VWFA is maximally

activated when letter strings are presented *and* the observer is voluntarily looking for known

words. It is not sufficient for words to be presented or even attended visually. Moreover,

engagement in a lexical task does not always enhance the VWFA's response; in fact, it

suppresses the response to unfamiliar characters. These findings open the door to future work

on more naturalistic reading tasks, as well as comparisons to other specialized parts of the

visual system.

References

Balota DA, Yap MJ, Cortese MJ, Hutchison KA, Kessler B, Loftis B, Neely JH, Nelson DL, Simpson GB, Treiman R (2007) The English Lexicon Project. Behav Res Methods 39:445–459.

Ben-Shachar M, Dougherty RF, Deutsch GK, Wandell BA (2007) Differential sensitivity to words and shapes in ventral occipito-temporal cortex. Cereb Cortex 17:1604–1611.

Binder JR, Medler DA, Desai R, Conant LL, Liebenthal E (2005) Some neurophysiological constraints on models of word naming. Neuroimage 27:677–693.

Boukrina O, Hanson SJ, Hanson C (2014) Modeling activation and effective connectivity of VWFA in same script bilinguals. Hum Brain Mapp 35:2543–2560.

Brainard DH (1997) The Psychophysics Toolbox. Spat Vis 10:433–436.

Brown TL, Gore CL, Carr TH (2002) Visual attention and word recognition in stroop color naming: is word recognition "automatic"? J Exp Psychol Gen 131:220–240.

Caffarra S, Karipidis II, Yablonski M, Yeatman JD (2021) Anatomy and physiology of word-selective visual cortex: from visual features to lexical processing. Brain Struct Funct 226:3051–3065.

Carr TH (1992) Automaticity and cognitive anatomy: is word recognition "automatic"? Am J Psychol 105:201–237.

Centanni TM, Norton ES, Ozernov-Palchik O, Park A, Beach SD, Halverson K, Gaab N, Gabrieli JDE (2019) Disrupted left fusiform response to print in beginning kindergartners is associated with subsequent reading. Neuroimage Clin 22:101715.

Chen L, Wassermann D, Abrams DA, Kochalka J, Gallardo-Diez G, Menon V (2019) The visual word form area (VWFA) is part of both language and attention circuitry. Nat Commun 10:5601.

Cohen L, Dehaene S, Vinckier F, Jobert A, Montavont A (2008) Reading normal and degraded words: contribution of the dorsal and ventral visual pathways. Neuroimage 40:353–366.

Dale AM, Fischl B, Sereno MI (1999) Cortical surface-based analysis. I. Segmentation and surface reconstruction. Neuroimage 9:179–194.

Dębska A, Wójcik M, Chyl K, Dzięgiel-Fivet G, Jednoróg K (2023) Beyond the Visual Word Form Area – a cognitive characterization of the left ventral occipitotemporal cortex. Front Hum Neurosci 17 Available at: https://www.frontiersin.org/articles/10.3389/fnhum.2023.1199366.

Dehaene S, Cohen L (2011) The unique role of the visual word form area in reading. Trends Cogn
Sci 15:254–262.

Dehaene S, Cohen L, Morais J, Kolinsky R (2015) Illiterate to literate: behavioural and cerebral
changes induced by reading acquisition. Nat Rev Neurosci 16:234–244.

Dehaene S, Cohen L, Sigman M, Vinckier F (2005) The neural code for written words: a proposal.
Trends Cogn Sci 9:335–341.

Domagalik A, Beldzik E, Oginska H, Marek T, Fafrowicz M (2014) Inconvenient correlation - RT-
BOLD relationship for homogeneous and fast reactions. Neuroscience 278:211–221.

Downing PE, Jiang Y, Shuman M, Kanwisher N (2001) A cortical area selective for visual
processing of the human body. Science 293:2470–2473.

Esteban O, Markiewicz CJ, Blair RW, Moodie CA, Isik AI, Erramuzpe A, Kent JD, Goncalves M,
DuPre E, Snyder M, Oya H, Ghosh SS, Wright J, Durnez J, Poldrack RA, Gorgolewski KJ
(2019) fMRIPrep: a robust preprocessing pipeline for functional MRI. Nat Methods
16:111–116.

Fedorenko E, Fillmore P, Smith K, Bonilha L, Fridriksson J (2015) The superior precentral gyrus of
the insula does not appear to be functionally specialized for articulation. J Neurophysiol
113:2376–2382.

Fedorenko E, Hsieh P-J, Nieto-Castañón A, Whitfield-Gabrieli S, Kanwisher N (2010) New method
for fMRI investigations of language: defining ROIs functionally in individual subjects. J
Neurophysiol 104:1177–1194.

Forseth KJ, Kadipasaoglu CM, Conner CR, Hickok G, Knight RT, Tandon N (2018) A lexical
semantic hub for heteromodal naming in middle fusiform gyrus. Brain 141:2112–2126.

Gagl B, Richlan F, Ludersdorfer P, Sassenhagen J, Eisenhauer S, Gregorova K, Fiebach CJ (2022)
The lexical categorization model: A computational model of left ventral occipito-
temporal cortex activation in visual word recognition. PLoS Comput Biol 18:e1009995.

Glezer LS, Jiang X, Riesenhuber M (2009) Evidence for highly selective neuronal tuning to whole
words in the "visual word form area." Neuron 62:199–204.

Glezer LS, Riesenhuber M (2013) Individual variability in location impacts orthographic
selectivity in the "visual word form area." J Neurosci 33:11221–11226.

Gorgolewski K, Burns CD, Madison C, Clark D, Halchenko YO, Waskom ML, Ghosh SS (2011)
Nipype: a flexible, lightweight and extensible neuroimaging data processing framework
in python. Front Neuroinform 5:13.

Graves WW, Desai R, Humphries C, Seidenberg MS, Binder JR (2010) Neural systems for reading
aloud: a multiparametric approach. Cereb Cortex 20:1799–1815.

Greve DN, Fischl B (2009) Accurate and robust brain image alignment using boundary-based
registration. Neuroimage 48:63–72.

Grill-Spector K, Weiner KS (2014) The functional architecture of the ventral temporal cortex and
its role in categorization. Nat Rev Neurosci 15:536–548.

Hagoort P (2014) Nodes and networks in the neural architecture for language: Broca's region
and beyond. Curr Opin Neurobiol 28:136–141.

Jolly E (2018) Pymer4: Connecting R and python for linear mixed modeling. J Open Source Softw
3:862.

Joo SJ, Tavabi K, Caffarra S, Yeatman JD (2021) Automaticity in the reading circuitry. Brain Lang
214:104906.

Kay K, Jamison KW, Zhang R-Y, Uğurbil K (2020) A temporal decomposition method for
identifying venous effects in task-based fMRI. Nat Methods 17:1033–1039.

Kay KN, Yeatman JD (2017) Bottom-up and top-down computations in word- and face-selective
cortex. Elife 6 Available at: http://dx.doi.org/10.7554/eLife.22341.

Kronbichler M, Hutzler F, Wimmer H, Mair A, Staffen W, Ladurner G (2004) The visual word form
area and the frequency with which words are encountered: evidence from a parametric
fMRI study. Neuroimage 21:946–953.

Kubota EC, Joo SJ, Huber E, Yeatman JD (2019) Word selectivity in high-level visual cortex and
reading skill. Dev Cogn Neurosci 36:100593.

Lanczos CAD (1964) Evaluation of Noisy Data. 1:10.

Le R, Witthoft N, Ben-Shachar M, Wandell B (2017) The field of view available to the ventral
occipito-temporal reading circuitry. J Vis 17:6.

Lerma-Usabiaga G, Carreiras M, Paz-Alonso PM (2018) Converging evidence for functional and
structural segregation within the left ventral occipitotemporal cortex in reading. Proc
Natl Acad Sci U S A 115:E9981–E9990.

Li J, Osher DE, Hansen HA, Saygin ZM (2020) Innate connectivity patterns drive the development
of the visual word form area. Sci Rep 10:18039.

López-Barroso D, Thiebaut de Schotten M, Morais J, Kolinsky R, Braga LW, Guerreiro-Tauil A,
Dehaene S, Cohen L (2020) Impact of literacy on the functional connectivity of vision and
language related networks. Neuroimage 213:116722.

Mahowald K, Fedorenko E (2016) Reliable individual-level neural markers of high-level language
processing: A necessary precursor for relating neural variability to behavioral and genetic
variability. Neuroimage 139:74–93.

Mano QR, Humphries C, Desai RH, Seidenberg MS, Osmon DC, Stengel BC, Binder JR (2013) The
role of left occipitotemporal cortex in reading: reconciling stimulus, task, and lexicality
effects. Cereb Cortex 23:988–1001.

McCandliss BD, Cohen L, Dehaene S (2003) The visual word form area: expertise for reading in
the fusiform gyrus. Trends Cogn Sci 7:293–299.

Muayqil T, Davies-Thompson J, Barton JJS (2015) Representation of visual symbols in the visual
word processing network. Neuropsychologia 69:232–241.

Mumford JA, Bissett PG, Jones HM, Shim S, Rios JAH, Poldrack RA (2024) The response time
paradox in functional magnetic resonance imaging analyses. Nat Hum Behav 8:349–360.

Nobre AC, Allison T, McCarthy G (1994) Word recognition in the human inferior temporal lobe.
Nature 372:260–263.

Nobre AC, Allison T, McCarthy G (1998) Modulation of human extrastriate visual processing by
selective attention to colours and words. Brain 121 (Pt 7):1357–1368.

Oh A, Duerden EG, Pang EW (2014) The role of the insula in speech and language processing.
Brain Lang 135:96–103.

Pattamadilok C, Chanoine V, Pallier C, Anton J-L, Nazarian B, Belin P, Ziegler JC (2017)
Automaticity of phonological and semantic processing during visual word recognition.
Neuroimage 149:244–255.

Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming numbers into
movies. Spat Vis 10:437–442.

Planton S, Chanoine V, Sein J, Anton J-L, Nazarian B, Pallier C, Pattamadilok C (2019) Top-down
activation of the visuo-orthographic system during spoken sentence processing.
Neuroimage 202:116135.

Price CJ, Devlin JT (2011) The interactive account of ventral occipitotemporal contributions to
reading. Trends Cogn Sci 15:246–253.

Price CJ, Wise RJ, Frackowiak RS (1996) Demonstrating the implicit processing of visually
presented words and pseudowords. Cereb Cortex 6:62–70.

Prince JS, Charest I, Kurzawski JW, Pyles JA, Tarr MJ, Kay KN (2022) Improving the accuracy of
single-trial fMRI response estimates using GLMsingle. Elife 11 Available at:
http://dx.doi.org/10.7554/eLife.77599.

Qin L, Lyu B, Shu S, Yin Y, Wang X, Ge J, Siok W-T, Gao J-H (2021) A Heteromodal Word-Meaning
Binding Site in the Visual Word Form Area under Top-Down Frontoparietal Control. J
Neurosci 41:3854–3869.

Rapp B, Purcell J, Hillis AE, Capasso R, Miceli G (2016) Neural bases of orthographic long-term
memory and working memory in dysgraphia. Brain 139:588–604.

Rauschecker AM, Bowen RF, Parvizi J, Wandell BA (2012) Position sensitivity in the visual word
form area. Proc Natl Acad Sci U S A 109:E1568-77.

Reich L, Szwed M, Cohen L, Amedi A (2011) A ventral visual stream reading center independent
of visual experience. Curr Biol 21:363–368.

Reuter M, Rosas HD, Fischl B (2010) Highly accurate inverse consistent registration: a robust
approach. Neuroimage 53:1181–1196.

Reynolds JH, Heeger DJ (2009) The normalization model of attention. Neuron 61:168–185.

Rouder JN, Speckman PL, Sun D, Morey RD, Iverson G (2009) Bayesian t tests for accepting and
rejecting the null hypothesis. Psychon Bull Rev 16:225–237.

Ruthruff E, Allen PA, Lien M-C, Grabbe J (2008) Visual word recognition without central
attention: evidence for greater automaticity with greater reading ability. Psychon Bull
Rev 15:337–343.

Ruz M, Nobre AC (2008) Attention modulates initial stages of visual word processing. J Cogn
Neurosci 20:1727–1736.

Stevens WD, Kravitz DJ, Peng CS, Tessler MH, Martin A (2017) Privileged Functional Connectivity
between the Visual Word Form Area and the Language System. J Neurosci 37:5288–
5297.

Stigliani A, Weiner KS, Grill-Spector K (2015) Temporal Processing Capacity in High-Level Visual
Cortex Is Domain Specific. J Neurosci 35:12412–12424.

Striem-Amit E, Cohen L, Dehaene S, Amedi A (2012) Reading with sounds: sensory substitution
selectively activates the visual word form area in the blind. Neuron 76:640–652.

Stroop JR (1935) Studies of interference in serial verbal reactions. J Exp Psychol 18:643–662.

Taylor JSH, Rastle K, Davis MH (2013) "Can cognitive models explain brain activation during work
and pseudoword reading: A meta-analysis of 36 neuroimaging studies": Correction to
Taylor, Rastle, and Davis (2012). Psychol Bull 139:900.

Vidal C, Content A, Chetail F (2017) BACS: The Brussels Artificial Character Sets for studies in
cognitive psychology and neuroscience. Behav Res Methods 49:2093–2112.

Vildavski VY, Lo Verde L, Blumberg G, Parsey J, Norcia AM (2022) PseudoSloan: A perimetric-
complexity and area-controlled font for vision and reading research. J Vis 22:7.

Vinckier F, Dehaene S, Jobert A, Dubus JP, Sigman M, Cohen L (2007) Hierarchical coding of
letter strings in the ventral stream: dissecting the inner organization of the visual word-
form system. Neuron 55:143–156.

Virtanen P et al. (2020) SciPy 1.0: fundamental algorithms for scientific computing in Python.
Nat Methods 17:261–272.

Vogel AC, Miezin FM, Petersen SE, Schlaggar BL (2012) The putative visual word form area is
functionally connected to the dorsal attention network. Cereb Cortex 22:537–549.

White AL, Kay KN, Tang KA, Yeatman JD (2023) Engaging in word recognition elicits highly
specific modulations in visual cortex. Curr Biol 33:1308-1320.e5.

White AL, Palmer J, Boynton GM, Yeatman JD (2019) Parallel spatial channels converge at a
bottleneck in anterior word-selective cortex. Proc Natl Acad Sci U S A 116:10087–10096.

Woolnough O, Donos C, Curtis A, Rollo PS, Roccaforte ZJ, Dehaene S, Fischer-Baum S, Tandon N
(2022) A Spatiotemporal Map of Reading Aloud. J Neurosci 42:5438–5450.

Woolnough O, Donos C, Murphy E, Rollo PS, Roccaforte ZJ, Dehaene S, Tandon N (2023)
Spatiotemporally distributed frontotemporal networks for sentence reading. Proc Natl
Acad Sci U S A 120:e2300252120.

Woolnough O, Donos C, Rollo PS, Forseth KJ, Lakretz Y, Crone NE, Fischer-Baum S, Dehaene S,
Tandon N (2021) Spatiotemporal dynamics of orthographic and lexical processing in the
ventral visual pathway. Nat Hum Behav 5:389–398.

Woolnough O, Forseth KJ, Rollo PS, Tandon N (2019) Uncovering the functional anatomy of the
human insula during speech. Elife 8 Available at: http://dx.doi.org/10.7554/eLife.53086.

Yarkoni T, Barch DM, Gray JR, Conturo TE, Braver TS (2009) BOLD correlates of trial-by-trial
reaction time variability in gray and white matter: a multi-study fMRI analysis. PLoS One
4:e4257.

Yeatman JD, Dougherty RF, Ben-Shachar M, Wandell BA (2012) Development of white matter
and reading skills. Proc Natl Acad Sci U S A 109:E3045-53.

Yeatman JD, White AL (2021) Reading: The Confluence of Vision and Language. Annu Rev Vis Sci
7:487–517.

Extended Data

Real Words					Pseudowords				
fall	fair	heat	talk	bulb	onll	kide	tiet	cais	spon
dung	feet	game	stop	jeep	soad	evey	iraw	onod	mext
warm	rate	wide	send	role	sood	veen	geal	alst	nean
team	cold	pads	wait	maze	pust	reen	gope	woud	sain
wall	bats	nets	bald	owed	cich	sont	rown	fole	nage
dual	leak	hush	feat	fake	edea	hing	rops	gomy	fize
cart	lose	word	baby	fish	hane	yeal	onte	mowl	firl
muck	news	lily	ness	hell	tris	sloy	wist	mank	alat
flap	paid	cans	flee	fast	bour	tand	yext	wiky	diew
foam	pens	moss	lead	none	surn	nost	whis	lato	weat
ants	near	nude	kept	bows	nery	samb	gove	fook	dasy
size	bees	crap	lack	safe	coar	doel	tuch	hece	jure
oath	sort	cops	butt	lost	halp	sind	woth	lood	inlk
line	main	ache	menu	show	eved	moad	aung	higs	fike
bunk	lone	pans	view	free	moen	pich	houd	alar	juch
flat	peer	mint	grow	dark	welp	ment	wele	pase	sani
poor	club	chub	axis	drum	youd	jook	frew	ovew	dake
lung	wind	cord	boot	note	toat	lely	hais	inen	purd
hind	lads	seat	melt	lady	vero	mang	derm	leat	lind
edge	care	step	wife	norm	cass	yeen	neny	vass	aror
move	lied	deed	coin	lent	sork	toul	dook	juru	tind
hail	pier	cosy	aids	oats	mear	dowl	mard	wien	grur
mass	bust	male	fuse	lush	lery	pign	mune	foor	gool
oral	liar	area	lain	dame	hery	basy	wesk	sond	egen
plan	race	gaps	jars	walk	ween	clus	cerm	knto	wany
jets	type	gone	stay	arid	bory	blso	palk	rean	tery
junk	lids	hair	hour	oils	rith	golt	goor	werr	parm
fire	thin	fare	ways	sent	frme	tast	hony	mave	elsk
lord	comb	huge	shot	milk	heah	lesk	milm	wene	blus
gets	hope	tree	bred	cult	feld	iney	huss	oner	fing
bass	loss	jerk	seem	call	lons	liet	traw	cont	dopy
test	buys	rock	east	dash	fack	soll	havy	lome	arom
ones	isle	wild	mine	desk	unth	yoom	muro	kner	lish
coup	hymn	dear	ally	rise	eang	como	helf	roso	masy
dose	horn	pass	grip	duly	woen	sath	sest	shem	wead
jazz	caps	aide	west	ball	cand	thil	timp	lelf	eart
born	neck	fine	sign	thus	thow	eath	mish	bere	foon
hall	dead	foot	loaf	bolt	loat	rell	sids	dath	boin
bump	town	died	haul	late	rata	tood	terl	dort	foss
boys	gale	live	loop	dump	lous	clat	darl	tere	pook
dole	drab	limb	heir	chew	fift	vech	tirl	loof	werk
army	bake	skin	gasp	fade	nore	riss	toto	frow	loor
knot	deaf	fads	haze	bank	bood	beey	gaid	lact	fith
nuns	gold	bony	rest	pain	seef	frop	bext	shus	gide
huts	cane	doom	duck	meat	thad	hent	onch	hasy	thip

 Extended Data Figure 1-1: All words and pseudowords used in the experiment.

LH

RH

Word Form Broca's

Language Broca's

VWFA-2

VWFA-1

OTS-Limbs

STS

MFUS-Text

FFA-2

FFA-1

V4

sub-091

sub-118

sub-143

sub-145

1147

LH

RH

sub-146

sub-147

sub-150

sub-151

1148

LH   RH

sub-152

sub-154

sub-159

sub-160

1149

LH   **RH**

sub-161

sub-163

sub-164

sub-166

1150
1151   **Extended Data Figure 1-2:** Regions of interest in individual subject anatomical surface space.
1152
1153

ROI	Stimulus	Task	Mean ΔPSC	SEM	95% CI	t-value	BF	p-value
VWFA1	Text	LD - FC	0.15	0.02	0.09, 0.2	6.05	44.95	<0.001
		LD - SC	0.12	0.02	0.07, 0.17	5.77	34.29	<0.001
		SC - FC	0.03	0.02	0.02, 0.08	1.47	1	0.147
	False Fonts	LD - FC	-0.06	0.03	-0.14, 0.01	2.03	2	0.07
		LD - SC	-0.09	0.03	-0.16, -0.02	3.31	6.35	0.004
		SC - FC	0.03	0.03	-0.04, 0.1	1.04	1.08	0.3
VWFA2	Text	LD - FC	0.12	0.02	0.07, 0.16	6.5	527.66	<0.001
		LD - SC	0.1	0.02	0.05, 0.14	5.67	1603.96	<0.001
		SC - FC	0.02	0.02	0.01, 0.07	1.35	0.84	0.18
	False Fonts	LD - FC	-0.07	0.02	-0.12, 0.01	2.71	8.79	0.01
		LD - SC	-0.09	0.02	-0.15, -0.03	3.88	32.28	0.001
		SC - FC	0.03	0.02	-0.03, 0.08	1.05	0.57	0.3

1154 **Extended Data Table 2-1:** Statistics for stimulus and task differences for VWFA-1 and -2, reported separately.
1155

1156
1157 **Extended Data Figure 2-1:** Individual participants' mean beta responses in the left VWFA (VWFA-1 and VWFA-2
1158 together) for **A)** text (real words and pseudowords) and **B)** false fonts. In each panel, each participant contributes
1159 two points that are connected by a vertical line. The horizontal position of each point represents that subject's
1160 VWFA response during the fixation task. The vertical position of the darker points represents the response during
1161 the lexical decision task, and the vertical position of the lighter points represents the stimulus color task.
1162

ROI	Stimulus	Task	Mean ΔPSC	SEM	95% CI	t-value	BF	p-value
FFA1	Text	LD - FC	-0.02	0.01	-0.04, 0.02	1.15	0.44	0.26
		LD - SC	-0.03	0.01	-0.06, 0.001	2.38	1.86	0.06
		SC - FC	0.01	0.01	-0.01, 0.04	1.13	0.86	0.26
	False Fonts	LD - FC	-0.005	0.02	-0.05, 0.04	0.31	0.29	0.76
		LD - SC	0.009	0.02	-0.03, 0.05	0.54	0.28	0.76
		SC - FC	-0.01	0.02	-0.05, 0.03	0.86	0.32	0.76
FFA2	Text	LD - FC	0.025	0.02	-0.01, 0.06	1.65	0.6	0.72
		LD - SC	0.014	0.02	-0.02, 0.05	0.95	0.34	0.72
		SC - FC	0.012	0.02	-0.02, 0.05	0.81	0.7	0.72
	False Fonts	LD - FC	-0.03	0.02	-0.08, 0.02	1.48	1.42	0.72
		LD - SC	-0.04	0.02	-0.09, 0.01	2.07	1.55	0.72
		SC - FC	0.01	0.02	0.04, 0.06	0.55	0.39	0.86
OTS-Limbs	Text	LD - FC	0.02	0.01	-0.01, 0.04	1.45	0.5	0.72
		LD - SC	0.02	0.01	-0.005, 0.043	1.96	0.67	0.72
		SC - FC	-0.003	0.01	-0.03, 0.02	0.28	0.29	0.91
	False Fonts	LD - FC	-0.014	0.01	-0.05, 0.02	0.95	0.34	0.72
		LD - SC	-0.033	0.01	-0.07, -0.001	2.52	1.25	0.48
		SC - FC	0.019	0.01	-0.013, 0.05	1.46	0.62	0.72
STS	Text	LD - FC	0.009	0.007	-0.009, 0.03	1.2	0.5	0.72
		LD - SC	0.01	0.007	-0.008, 0.03	1.35	0.7	0.72
		SC - FC	-0.001	0.007	-0.02, 0.02	0.15	0.27	0.96
	False Fonts	LD - FC	0.01	0.01	-0.01. 0.04	1.11	0.41	0.4
		LD - SC	-0.002	0.01	-0.03, 0.02	0.16	0.26	0.87
		SC - FC	0.01	0.01	-0.1, 0.04	1.28	0.67	0.4
V4	Text	LD - FC	-0.01	0.009	-0.04, 0.01	1.38	0.52	0.81
		LD - SC	-0.009	0.009	-0.03, 0.14	0.95	0.33	0.81
		SC - FC	-0.004	0.009	-0.03, 0.02	0.42	0.29	0.89
	False Fonts	LD - FC	0	0.01	-0.03, 0.03	0.03	0.26	0.98
		LD - SC	-0.001	0.01	-0.03, 0.03	0.05	0.26	0.98
		SC - FC	0.001	0.01	-0.03, 0.03	0.08	0.26	0.98
MFUS-Text	Text	LD - FC	0.07	0.02	0.02, 0.12	3.68	1.99	0.07
		LD - SC	0.06	0.02	0.01, 0.11	3.29	2.14	0.07
		SC - FC	0.01	0.02	-0.04, 0.06	0.54	0.32	0.86
	False Fonts	LD - FC	-0.04	0.03	-0.11, 0.02	1.6	0.53	0.72
		LD - SC	-0.06	0.03	-0.12, 0.01	2.1	1.44	0.72
		SC - FC	0.012	0.03	-0.05, 0.08	0.47	0.28	0.89

Extended Data Figure 3-1: Statistics for task effects on responses to each stimulus type for all control ROIs in the left hemisphere. P-values are corrected for multiple comparisons using FDR correction. All p-values were corrected for post-hoc tests within ROIs, and then concatenated and corrected for multiple comparisons across ROIs. Adjusted P-values are reported in the final column of the table.

Extended Data Figure 3-2: Mean BOLD percent signal change as a function of stimulus and task types in control ROIs in the right hemisphere. Error bars represent standard error of the mean.