1 **Compact Programmable Control of Protein Secretion in Mammalian Cells**
2

3 Authors: Alexander E. Vlahos[1], Connor C. Call[1], Samarth E. Kadaba[1], Siqi Guo[1,2], Xiaojing J. Gao[1,3]*
4
5 1: Department of Chemical Engineering, Stanford University, Stanford, CA, 94305, USA
6 2: The Chinese Undergraduate Visiting Research (UGVR) Program, Stanford, CA, 94305, USA
7 3: Neurosciences Interdepartmental Program, Stanford University, Stanford, CA, 94305, USA
8
9 *Corresponding Author = xjgao@stanford.edu

10
Abstract:

Synthetic biology currently holds immense potential to engineer the spatiotemporal control of intercellular
signals for biomedicine. Programming behaviors using protein-based circuits has advantages over
traditional gene circuits such as compact delivery and direct interactions with signaling proteins. Previously,
we described a generalizable platform called RELEASE to enable the control of intercellular signaling
through the proteolytic removal of ER-retention motifs compatible with pre-existing protease-based circuits.
However, these tools lacked the ability to reliably program complex expression profiles and required
numerous proteases, limiting delivery options. Here, we harness the recruitment and antagonistic behavior
of endogenous 14-3-3 proteins to create RELEASE-NOT to turn off protein secretion in response to
protease activity. By combining RELEASE and RELEASE-NOT, we establish a suite of protein-level
processing and output modules called Compact RELEASE (compRELEASE). This innovation enables
functions such as logic processing and analog signal filtering using a single input protease. Furthermore,
we demonstrate the compactness of the post-translational design by using polycistronic single transcripts
to engineer cells to control protein secretion via lentiviral integration and leverage mRNA delivery to
selectively express cell surface proteins only in engineered cells harboring inducible proteases.
CompRELEASE enables complex control of protein secretion and enhances the potential of synthetic
protein circuits for therapeutic applications, while minimizing the overall genetic payload.

28
29
30 **Keywords:** intercellular communication, synthetic biology, proteases, 14-3-3 proteins, protein circuits,
31 mRNA delivery

Introduction:

The field of synthetic biology is uniquely suited to develop biomolecular circuits to engineer cells[1–4] for various biomedical applications, such as improving cancer immunotherapies[5,6] and cell therapies[7]. These engineered circuits can enable cells to respond to combinatorial environmental inputs[8,9], interrogate natural systems[10], and produce controlled therapeutic responses[11]. In particular, an emerging strategy for programming cellular function in mammalian cells is to use orthogonal proteases to create synthetic protein circuits[12,13]. Protein-based circuits can implement synthetic signal-processing, featuring advantages such as fast operation, compact design, and robust context-independent performance compared to traditional transcriptional circuits[12–14]. Previously, we[15] and others[16] described generalizable platforms ("RELEASE" and "membER/lumER", respectively), which interfaced with pre-existing protease-based circuits[12,13] and enabled the control of intercellular signals through proteolytic removal of ER-retention motifs (**Fig. 1a-c**). However, to program complex expression patterns such as logic operations or quantitative processing, multiple orthogonal proteases were required[12,13], which may hinder their use in viral vectors with limited packaging capacities[17]. Therefore, to fully unlock the potential of these synthetic protein circuits for biomedicine, there is a need to harness additional post-translational modifications to develop synthetic protein components that condense genetic payloads while retaining or even enhancing circuit functionality.

Post-translational modifications of proteins represent a cornerstone in cellular regulation, orchestrating a diverse array of functions crucial for cellular homeostasis and responsiveness[18–20]. Among these, the interplay between post-translational modifications and protein-protein interactions, particularly with 14-3-3 scaffolding proteins, stands as a pivotal nexus of cellular control[21,22]. The 14-3-3 protein family is expressed in all eukaryotes and are adept at recognizing phosphorylated motifs on target proteins[23]. When 14-3-3 proteins bind to phosphorylated motifs, they can mediate diverse cellular outcomes, including sequestration from specific cellular compartments, stabilization of protein complexes, or modulation of enzymatic activity[21,24]. Of particular interest is their ability to block interactions between the COPI retrograde transport machinery and ER-retention motifs to control the surface expression of membrane proteins[25–30] (**Fig. 1d**). Although surface expression is mediated by phosphorylation-dependent recruitment of 14-3-3 proteins, this specific protein-protein interaction may be exploitable when using pre-existing synthetic protease-based circuits[12,13,31].

Given the size limitation of viral vectors for genetic delivery[17], we sought to expand the RELEASE platform and enable compact programmable control of protein secretion. Specifically, we used a small 14-3-3 recruiting peptide to manipulate protein-protein interactions with endogenous 14-3-3 scaffolding proteins to create RELEASE-NOT, an approach to turn off protein secretion in response to protease activity. With the ability to control the activation or repression of protein secretion in response to protease activity, we created a suite of RELEASE variants known as Compact RELEASE (compRELEASE). The compRELEASE suite achieved functional completeness of all Boolean logic gates and enabled analog signal filtering without requiring additional processing proteases[12,13]. We also focused on the compact delivery of synthetic protein circuits, and optimized polycistronic constructs under the control of a single promoter to create engineered cells with inducible control of protein secretion. Finally, to highlight the utility of post-translational circuits, we leveraged the mRNA delivery of RELEASE to selectively express cell surface proteins only in engineered cells harboring inducible proteases. This study builds on the RELEASE platform to unlock its potential as a protein-level processing and output module to enable circuit-level control of protein secretion, while minimizing the overall genetic payload.

Materials and Methods:

Plasmid generation:

81 All plasmids were constructed using general practices. Backbones were linearized via restriction digestion,
82 and inserts were generated using PCR, or purchased from Twist Biosciences. A complete list of plasmids
83 and their respective amounts used for each experiment can be found in **Supplementary Table 1**.
84 Furthermore, all new plasmids used in this study will be deposited with annotations to Addgene
85 (https://www.addgene.org/Xiaojing_Gao/).
86
Tissue culture: Human Embryonic Kidney (HEK) 293 cells (ATCC, CRL-1573) and HEK293T-LentiX
(Takara Biosciences) were cultured in standard culture conditions (37ºC and 5% CO2) in Dulbecco's
Modified Eagle Media (DMEM), supplemented with 10% Fetal Bovine Serum (FBS, ThermoFisher; catalog#
FB12999102), 1X Pen/Strep (Genesee; catalog# 25-512), 1X non-essential amino acids (Genesee;
catalog# 25-536), and 1X sodium pyruvate (Santa Cruz Biotechnology; catalog# sc-286966). APRE-19
cells (ATCC, CRL-2302) were cultured in standard culture conditions in DMEM/F-12 (1:1) GlutaMAX media
(ThermoFisher; catalog# 10565018), supplemented with 10% FBS, and 1X PenStrep. K562 cells (ATCC,
CCL-243) were cultured in RPMI-1640 (Sigma-Aldrich; catalog# R8758) media supplemented with 10%
FBS, and 1X PenStrep. HEK293 cells were used for all transient transfection experiments and K562 cells
were used for mRNA transfection experiments. HEK293T cells were used to produce lentivirus, described
below. Cells tested negative for mycoplasma.
98
99 **Transient transfection:** HEK293 cells were cultured in 96-well tissue-culture treated plates under standard
100 culture conditions. At 70-90% confluency, cells were transiently transfected with plasmids using the
101 jetOPTIMUS® DNA transfection Reagent (Polyplus transfection; catalog# 117-15), as per manufacturer's
102 instructions.
103
Lentiviral transduction: To generate lentivirus, HEK293T-LentiX cells (Takara Biosciences) were
transfected with 750 ng of an equimolar mixture of the three third-generation packaging plasmids (pMD2.G,
pRSV-Rev, pMDLg/pRRE) and 750 ng of donor plasmids using 200 uL of jetOPTIMUS® DNA transfection
Reagent (Polyplus transfection, catalog# 117-15). pMD2.G (Addgene plasmid #12259,
https://www.addgene.org/12259/), pRSV-Rev (Addgene plasmid #12253,
https://www.addgene.org/12253/), and pMDLg/pRRE (Addgene plasmid # 12251,
https://www.addgene.org/12251/) were gifts from Didier Trono. After 2 - 3 days of incubation, the lentivirus
was harvested and filtered through a 0.45 micron filter (Millipore) to remove any cellular debris. The
harvested virus was precipitated using the Lentivirus Precipitation Solution (Alstem; catalog# VC100) and
centrifuged at 1,500 x g for 30 minutes to remove any residual media. The concentrated virus was then
resuspended in supplemented DMEM, aliquoted and frozen at -80C for later use.
115
116 Lentivirus was titrated on HEK293 or APRE-19 cells by serial dilution in supplemented DMEM or DMEM/F-
117 12, respectively. 72 hours after incubation, the percentage of mCherry-positive cells were quantified using
118 flow cytometry. The cells were selected using 1000 ng/mL of puromycin (ThermoFisher Scientific; catalog#
119 J61278-MB) for one week until >90% of cells were mCherry-positive before being used for *in vitro* or *in vivo*
120 experiments.
121
K562 inducible reporter cell line: To generate the reporter cell line, K562 were electroporated in Amaxa
solution (Lonza Nucleofector 2b, setting T0-16) with 1µg of reporter donor (pEF1a-p450-rapalog-split-
TEVP-T2A-mCherry) and 1 µg of SuperPiggyBac transposase. After 7 days, the cells were treated with
1000 ng/mL of puromycin for two weeks to ensure that the construct was stably integrated. A monoclonal
population was generated through serial dilution. The expression of the fluorescent reporter was measured
by microscopy and flow cytometry. The SuperPiggyBac transposase plasmid was a gift from the Bassik
lab.
129

mRNA Synthesis and Transfection: The IgG-RELEASE construct was cloned into a backbone containing optimized 5' and 3' UTRs and a T7 promoter. The plasmid backbone was a gift from the Elowitz lab. Using a reverse primer containing a 200 bp polyA sequence, we added a Poly-A tail to the DNA template via PCR. In vitro transcribed mRNA was produced and purified by GeneScript Inc. based on the DNA template, with Cap1 and modified bases (100% N1-methyl-pseudouridine).

Briefly, $2.5x10^5$ wildtype K562 cells and $2.5x10^5$ reporter K562 cells were mixed for a total of $5.0x10^5$ cells per well and transfected with various amounts of the purified mRNA using the TransIT®-mRNA Transfection Kit (Mirus Bio, catalog# MIR2250). To induce the reporter cell lines, Rapalog AP21967 (also known as A/C heterodimerizer, purchased from Takara Biosciences; catalog# 635056) was administered at the time of transfection. All experiments were induced with 100 nM of rapalog, unless otherwise stated.

Flow cytometry and data analysis: Following 24 hours after mRNA transfection, K562 cells were pelleted at 300 x g for 7 minutes and reconstituted in PBS supplemented with 5% Bovine Serum Albumin (BSA, company name). Following two wash steps, the cells were incubated with an anti-human IgG antibody conjugated to Alexa Fluor-488 (Southern Biotech Catalog# 2015-30; 1/750 dilution in PBS + 2% BSA) for 1 hour. Following incubation, the cells were washed twice in PBS + 2% BSA and then filtered through a 40 µm sieve. Cells were analyzed by flow cytometry (BioRad ZE5 Cell Analyzer) and Everest Software (version 3.1). FlowJo (version 10.9.0) was used to process the flow cytometry data.

For analysis, cells were gated for cells that were expressing the IgG-RELEASE-T2A-BFP construct (BFP+). The median fluorescence intensity (MFI) of AF488 was calculated for wildtype (mCherry-) and reporter (mCherry+) cells, and then compared between uninduced and induced conditions. For each condition, an IgG isotype control (Southern Biotech Catalog# 0109-30) to correct the MFI of AF488 for non-specific staining.

Measuring protein secretion: Protein secretion was measured by a Secreted Embryonic Alkaline Phosphatase (SEAP) assay, as previously described[15]. Briefly, two days after transient transfection or induction with rapalog, the supernatant was collected and heat-inactivated at 70C for 45 minutes. After heat-inactivation, 10 - 40 uL of supernatant was mixed with dH_2O to a final volume of 80 uL. This mixture was then mixed with with 100 uL of 2X SEAP Buffer (20 mM homoarginine (ThermoFisher catalog# H27387), 1 mM MgCl2, and 21% (v/v) dioethanolamine (ThermoFisher, catalog# A13389)) and 20 µL of the p-nitrophenyl phosphate (PNPP, Acros Organics catalog# MFCD00066288) substrate (120 mM). Samples were measured via kinetic measurements (1 reading/minute) for a total of 30 minutes at 405 nm using a SpectraMax iD3 spectrophotometer (Molecular Devices) with the Softmax pro software (version 7.0.2)

Secreted NanoLuc® (secNLuc) was measured by taking the supernatant from samples that were previously transiently transfected or induced with rapalog. Briefly, 10 uL of supernatant was mixed with 45 µL of dH2O and then mixed with 50 uL of the Nano-Glo® Luciferase Assay system (Promega; catalog# N1110), per the manufacturer's instructions. Samples were measured using a SpectraMax iD3 spectrophotometer (Molecular Devices) with the Softmax pro software (version 7.0.2).

Statistical Analysis:
Values are reported as the average of four biological replicates, and representative of at least two independent biological experiments. For experiments comparing two groups, a Student's T-test was used to determine statistical significance, following confirmation that equal variance could be assumed (F-test). If equal variance could not be assumed, then a Welch's correction was used. For experiments with more than two groups, a one-way ANOVA with a post hoc Tukey's test was used for multiple comparisons among

179 the different means. Data was considered statistically significant when the p-value was less than 0.05. Data
180 presented are average +/- SEM, unless otherwise stated. All statistical analysis was performed using Prism
181 9.0 (GraphPad).
182
183 **Results:**
184
185 **Using additional protein motifs to modulate secretion with RELEASE:**
186
The original RELEASE platform used the dilysine ER-retention motif (**Fig. 1b**) to retain tagged proteins of
interest via retrograde transport, mediated through interactions with the COPI complex[32–35] (**Fig. 1a**). Only
after proteolytic removal of the retention motif would the protein of interest be transported through the
conventional secretory pathway, processed into its secretory form via furin (**Fig. 1c**), and secreted.
However, the dilysine motif (**Fig. 1b**, -KKXX-COOH) only functioned when placed at the C-terminus[34,36] and
limited the potential of RELEASE to be used in polycistronic constructs separated by self-cleaving 2A
peptides[14,37]. Indeed, since 2A peptides leave a small peptide scar at the C-terminus, when transiently
transfecting a polyprotein encoding Secreted Alkaline Embryonic Phosphatase (SEAP) fused to RELEASE
and the mNeon reporter protein (**Supplementary Fig. 1a**), we observed protein secretion with and without
co-expression of TEVP (**Supplementary Figure 1b**). Therefore, we identified additional ER-retention
protein motifs such as the diarginine motif (-RXR-, **Fig. 1b**)[38] that were not limited to the C-terminus of
membrane proteins and created updated RELEASE constructs (**Fig. 1e – left panel**). Using these new
RELEASE constructs, we transiently transfected HEK293 cells (**Fig. 1e – right panel**) and observed a
significant increase in SEAP secretion when co-expressing the cognate protease relative to when the
protease was absent (**Fig. 1e – left panel**). As expected, these new RELEASE variants retained their ER-
retention capabilities when encoded as the first protein in a polypeptide construct separated via a T2A
peptide (**Supplementary Fig. 1c**). For the remainder of this study, we used RELEASE variants with the
RXR retention motif unless otherwise stated. Furthermore, this new design unlocked the ability to generate
fusion proteins with RELEASE to potentially impart new behaviors.
206
To test the inhibitory effect of 14-3-3 proteins on the ER-retention activity of RELEASE to facilitate protein
secretion (**Fig.1d**)[27,39], we fused the 14-3-3ζ protein to the C-terminus of RELEASE (**Supplementary Fig.
2a**). As expected, the fusion of 14-3-3ζ resulted in a significant increase in protein secretion, compared to
RELEASE alone (**Supplementary Fig. 2b**). This fusion highlighted the potential of using 14-3-3 proteins
to block the ER-retention capabilities of RELEASE, however since our focus was to keep constructs as
small as possible, we sought an alternative strategy focused on the recruitment of endogenous 14-3-3
proteins to drive protein secretion.
214
Previously, the R18 peptide has been described to bind the conserved amphipathic ligand groove of 14-3-
3 proteins[40] via the WLDLE motif[41], and we hypothesized that its fusion to the C-terminus of RELEASE
would be sufficient to recruit 14-3-3 proteins and drive protein secretion (**Fig. 1f – left panel**). To evaluate
this, we created two RELEASE constructs: one with the R18 peptide and the other containing a mutant
variant lacking the two key acidic residues (referred to as R18 mutant) critical for mediating binding with
14-3-3 proteins[40]. The addition of R18 was sufficient in blocking the retention capabilities of RELEASE and
resulted in a significant increase in SEAP secretion relative to the RELEASE construct with the R18 mutant
peptide (**Fig. 1f**). This result shows that the ER-retention capabilities of RELEASE can be inhibited through
the addition of the R18 peptide at the C-terminus to drive protein secretion.
224
225 **Negative regulation of protein secretion and functional completeness:**
226

An important capability of synthetic protein circuits is the ability to integrate multiple inputs[12–14,31]. The ability
of negative regulation (i.e., NOT logic) is essential for implementing all possible Boolean logic operations,
in addition to the previous RELEASE-enabled compact implementation of AND/OR[15]. We attempted to use
CHOMP to create a two-protease circuit to turn off secretion in response to the expression of the first
protease (**Supplementary Fig.** 3a, TEVP, orange pac-man), however, protein secretion did not completely
return to baseline even at high levels of the repressing protease (**Supplementary Fig. 3b**). Instead of
relying on two proteases to impart negative regulation with CHOMP, we asked if we could program NOT
logic directly into the RELEASE platform.

By placing a protease cut site (such as TEVP) between the ER-retention motif and R18 peptide we created
RELEASE-NOT (**Fig. 2a**). RELEASE-NOT relies on the recruitment of 14-3-3 proteins to enable constitutive
secretion of a tagged protein of interest, until proteolytic removal of the R18 peptide results in restoration
of the ER-retention activity (**Fig. 2a**). Co-expression of TEVP with a TEVP-inducible RELEASE-NOT
significantly reduced protein secretion relative to when protease was not expressed (**Fig. 2b**). Like
RELEASE, we validated the modularity of RELEASE-NOT by switching the cytosolic protease cut site to
respond to other proteases such as the hepatitis C virus protease (HCVP) (**Supplementary Fig. 4**). Since
protein secretion with RELEASE-NOT is dependent on recruitment of 14-3-3 proteins, we overexpressed
exogenous 14-3-3ζ protein and observed a significant increase in SEAP secretion (**Supplementary Fig.
5b**). However, exogenous expression of 14-3-3ζ proteins reduced the dynamic range by increasing
baseline secretion (**Supplementary Fig. 5c**).

In addition to R18, we wanted to assess the compatibility of RELEASE-NOT with phosphorylation-
dependent binding motifs that recruit 14-3-3 proteins, such as the SWTY motif[42]. The SWTY motif is
phosphorylated by kinases within the PI3K/Akt pathway, which can be activated by the presence of FBS[29].
As expected, RELEASE-NOT using the SWTY motif, resulted in SEAP secretion in the absence of protease
and a significant decrease when co-expressing the cognate protease (**Supplementary Fig. 6a**). To validate
that protein secretion was due to phosphorylation-dependent recruitment of endogenous 14-3-3 proteins,
we created a variant where the phospho-responsive threonine residue was mutated to alanine (i.e., SWTY
→ SWAY). The RELEASE-NOT using SWAY resulted in a complete abrogation of SEAP secretion
(**Supplementary Fig.** 6b), suggesting phosphorylation was important for recruiting 14-3-3 proteins to inhibit
ER-retention and drive SEAP secretion. Although the SWTY motif was suitable for imparting NOT logic,
similar to the dilysine motif (**Supplementary Fig.** 1a), SWTY functions only at the C-terminus[42], therefore
making it incompatible when encoded by polycistronic constructs separated by self-cleaving 2A peptides.
In comparison, RELEASE-NOT with the R18 peptides preserved negative regulation of protein secretion
when encoded in polycistronic constructs (**Supplementary Fig. 7**).

Now that we could implement OR, AND, and NOT logic, we created a suite of RELEASE variants known
as Compact RELEASE (compRELEASE) for the different Boolean Logic gates and validated their behavior
through co-transfection with and without different combinations of input proteases (**Fig. 2c**). An important
characteristic of these variants is that the R18 peptide cannot inhibit the ER-retention capabilities of the N-
terminal p450 motif since it confers ER retention by directly inserting into the ER membrane[43,44]. For
example, to secrete SEAP under the control of RELEASE-NIMPLY, only the N-terminal p450 motif must be
removed since the C-terminal retention motif is being inhibited by the R18 peptide (**Supplementary Fig.
8a**). Any other combination of binary inputs resulted in either the p450 motif remaining intact, or the removal
of the R18 peptide and restoration of the C-terminal ER-retention motif activity (**Supplementary Fig. 8b**).
Most of the Boolean gates (5 of 8) were implemented using single RELEASE constructs, but some gates,
such as XNOR were created by co-expressing RELEASE variants with NOR and AND logic (**Fig. 2c**). The
compRELEASE suite highlights the modularity of the RELEASE design to create variants with predictable
logic operation functions without requiring additional proteases[12,13,45].

276
Quantitative processing using compRELEASE:

278
In addition to implementing Boolean logic, it would be advantageous to use the compRELEASE suite to
enable more core circuit functions, such as analog signal filtering[46]. Considering some intercellular signals
such as cytokines and growth factors have pleiotropic activity based on their concentrations[47–49], the ability
to selectively respond to specific concentrations ranges of inputs would be critical for enabling future
biomedical therapies. Proteolytic activation and repression of protein secretion within the compRELEASE
suite are dependent on cleavage efficiencies of the protease cut sites, which can be manipulated by
changing the canonical amino acid sequence (**Fig. 3a**). For example, the canonical TEVP cut site has a
serine amino acid (S) at the P1' position (**Fig. 3a – bolded residue**), and mutations to this residue to
alternative amino acids such tyrosine (Y) or leucine (L) have been reported to reduce cleavage efficiency[50].
Indeed, when using modified protease cut sites harboring these mutations, we observed a shift in the
activation thresholds for both RELEASE (**Fig. 3b**) and RELEASE-NOT (**Fig. 3c**).

290
Now that we could reliably manipulate the cleavage efficiencies of our constructs, we sought to achieve
analog signal processing by modifying the RELEASE-NIMPLY and RELEASE-IMPLY variants so that both
of their cut sites were cleaved by the same protease, but with different cleavage efficiencies. In addition,
these two variants were selected because they were single constructs that contained both activating and
repression arms for protein secretion (**Fig. 2c**). To implement band-pass filtering of protein secretion (**Fig.**
3d), RELEASE-NIMPLY was modified to contain two TEVP cut sites with the activating arm of protein
secretion to be more efficiently cleaved compared to the repression arm (**Fig. 3e**). This ensured that protein
secretion would first increase and then decrease, to increasing amounts of TEVP (**Fig. 3f**). We implemented
a similar strategy using the RELEASE-IMPLY variant to enable band-stop filtering (**Fig. 3g**), except that the
repressing arm of protein secretion was more efficiently cleaved than the activating arm. In this design,
protein secretion would first decrease and then increase, to increasing amounts of TEVP (**Fig. 3i**). These
two variants enabled quantitative processing using the compRELEASE suite to improve the compactness
of the circuit and while reducing the number of required proteases previously shown to be required to
achieve the same behavior[12].

305
Delivering RELEASE:

307
Unlike traditional gene regulation circuits that often require multiple integration events to create functional
circuits[51,52], synthetic protein circuits function at the post-translational level and can be encoded within a
single polycistronic construct. However, manipulating the relative concentrations of individual protein
components in single transcript designs is not trivial and may require additional optimization. To
demonstrate the utility of using synthetic protein circuits to engineer new behaviors into cells, we designed
a polycistronic construct encoding a puromycin-resistant cassette, a mCherry reporter protein, a rapalog-
inducible split HCVP and a secNLuc-RELEASE all under the control of the EF1α promoter (**Fig. 4a**). Using
lentiviral transduction, we created a stable HEK293 cell line with rapalog-inducible control of secNluc
secretion (**Fig, 4b**).

317
The initial single-transcript design encoded a split HCVP protease into the cytoplasm to activate secNluc
secretion in response to rapalog (**Fig. 4c – left panel**), however the dynamic range was poor (2.26-fold
increase). We have previously shown that sub-cellular localization of intermediate protease components
can modulate the activation of RELEASE[15], so we localized one half of the split protease to the ER using
the signal anchor sequence of cytochrome p450[43,44]. This fusion was expected to improve RELEASE
activation (**Fig. 4c – middle panel**), and we observed an increase in secNluc secretion following induction
with rapalog. However, this topology resulted in an increased baseline and did not significantly improve the

dynamic range (1.96-fold increase). Therefore, our next approach was designed to manipulate the cleavage
efficiency of the HCVP cut site, reducing the baseline of secNluc secretion, and improving the dynamic
range of the circuit. We have previously shown that modifying the flanking residues of the HCVP cut site
from their native sequences to GS-linkers significantly reduced the cleavage efficiency[15]. Indeed, using a
single transcript design with an ER-localized split HCVP and a less efficient HCVP-inducible RELEASE
construct resulted in the highest dynamic range (6.14-fold increase) due decreasing baseline secretion,
with comparable activation levels as the original design (**Fig. 4c – right panel**).
332
In addition to stable integration, synthetic protein circuits have the potential to be delivered as mRNA
transcripts without any risk of insertional mutagenesis[53]. Since RELEASE requires expression of the
cognate protease to control protein secretion, we hypothesized that we could leverage this property to
selectively express protein payloads in engineered cells, without activating RELEASE in wildtype cells (**Fig.**
4d). To validate this selectivity, we created an mRNA transcript encoding a human IgG-RELEASE and the
BFP reporter separated by a T2A peptide (**Fig. 4d**). Previously, human IgG has been used as a synthetic
surface marker[54], and we transfected this mRNA into a 50/50 mixture of wildtype (mCherry⁻) and engineered
(mCherry⁺) K562 cells. The engineered K562 cells contained a rapalog-inducible TEVP, and 24 hours after
transfection we assessed the surface expression of IgG using flow cytometry (**Fig. 4e**). From the mixture
of cells, only engineered K562 cells (mCherry⁺) showed a significant increase in the surface expression of
IgG when induced with rapalog (**Fig. 4f**). In comparison wildtype cells (mCherry⁻) did not show any
significant induction and comparable surface expression with the uninduced population of engineered cells
(**Fig. 4f**). Since mRNA expression is transient, the IgG retained in the ER of wildtype cells should eventually
degrade via the proteasomal degradation pathway. This work highlights the potential of using mRNA to
deliver RELEASE-tagged protein payloads that were selectivity expressed only when delivered into
engineered cells.
349
350   **Discussion:**
351
Here, we introduced the compRELEASE suite, a collection of engineered proteins with expanded input-
processing capabilities, enabling the precise control over intercellular signals in mammalian cells. The
compRELEASE suite was engineered by harnessing the complementary functions of RELEASE and
RELEASE-NOT to combine them in various configurations introduce novel behaviors. By functioning as
both the processing and output module, the compRELEASE suite allowed for the integration of multiple
inputs (**Fig. 2c**) and the implementation of analog-signal filtering (**Fig. 3**) to control protein secretion.
Notably, this functionality was achieved without the need of additional processing proteases[12,13]. The
reduction in the number of proteases is important as it streamlines the entire circuit, reduces the genetic
payload, and frees up proteases for use in designing synthetic receptors[55,56]. However, it should be noted
that the compactness enabled by compRELEASE is specifically applicable for protein secretion (**Fig. 2 and**
3) and the surface expression of proteins. Furthermore, given that the programmable features of
compRELEASE are independent of specific tagged protein of interest, we hypothesized that these variants
would uphold the plug-and-play adaptability of outputs demonstrated with the original RELEASE platform[15].
365
Since 14-3-3 proteins are ubiquitously expressed in all eukaryotic cells[23], we expect compRELEASE to
maintain its functionality across various cell types. Nonetheless, it is important to acknowledge that
differential expression of 14-3-3 proteins[23] in specific cells and tissues may influence the extent of protein
secretion, particularly when employing logic gates (**Fig. 2c**) or quantitative processing (**Fig. 3i**) dependent
on RELEASE-NOT (**Fig. 2a**). Notably, the reversible process of 14-3-3 recruitment and binding to the R18
peptide could be enhanced by overexpressing the 14-3-3ζ isoform to increase protein secretion using
RELEASE-NOT (**Supplementary Fig. 5c**). While this study primarily emphasized the use of the R18
peptide to modulate protein secretion (**Fig. 1f and 2a**), it is important to note that RELEASE-NOT also

demonstrated compatibility with phosphorylation-dependent binding motifs for recruiting 14-3-3 proteins
(**Supplementary Fig. 6)**. Through harnessing the interactions between phosphorylated peptides and 14-3-
3 proteins[57], there exists an opportunity to develop a class of phospho-responsive components to modulate
protein secretion and further expand the programming capabilities of compRELEASE. With over 200 distinct
human proteins documented to interact with 14-3-3 proteins[58], the potential to monitor shifts in vital cellular
processes such as cell cycle progression, glucose metabolism and apoptosis[21,59] may hold significant
implications for advancing therapeutics applications, such as cell transplantation and cancer
immunotherapies in the future.
382
The obstacle of delivery remains a significant challenge in the integration of synthetic protein circuits. While
these circuits can be encoded within a single polycistronic construct, it's imperative to fine-tune individual
components to ensure proper circuit functionality. Instead of relying on large and cumbersome Internal
Ribosomal Entry Sites (IRES) to regulate expression levels[60], we capitalized on the inherent modularity and
composability of engineered proteases and RELEASE to optimize our circuit's dynamic range at the post-
translational level (**Fig. 4c**). Although not explicitly tested here, fine-tuning the relative expression levels of
individual proteins has been found to be contingent on the positioning of individual genetic elements and
the selection of specific 2A peptides[61]. The ability to optimize the relative expression levels at the genetic
levels, in parallel with their respective activities at the protein level, holds the potential to enhance the
tuneability of synthetic protein circuits.
393
The examples presented thus far have employed a single compRELEASE construct to govern a single
output. However, the potential for more complex circuit functions is substantial, as multiple variants
introducing different logic (**Fig. 2c**) or quantitative processing (**Fig. 3e, f**) can be integrated. For instance,
by stacking RELEASE and RELEASE-NOT inducible variants to the same protease, one can establish
single-input multi-output behavior driven by the presence or absence of a given protease input. Introducing
these behaviors within engineered cells and co-delivering them with therapeutic cells to modulate the
dynamic immune response may be a suitable approach for improving the long-term engraftment without
the need of systemic immunosuppression. Moreover, the ability to exert on-demand control of outputs via
mRNA (**Fig. 4f**), ensures that only engineered cells equipped with the synthetic protease circuit can activate
and regulate their expression. The inherent compactness, delivery potential, and composability of synthetic
protein circuits enhanced by compRELEASE will be an invaluable tool for implementing programmable cell
therapies that control intercellular signals for biomedicine.
406

Acknowledgements: We would like to thank the Bassik lab and Elowitz lab for kindly sharing plasmids
used in this work. This work was funded by NIH (4R00EB027723-02, X.J.G.), the International Human
Frontier Science Program Organization (LT000221/2021-L, A.E.V.), National Science Foundation graduate
research fellowships program (DGE-2146755, C.C.C.), Cellular and Molecular Biology Training Grant (NIH
5 T32 GM007276, C.C.C.).

Author Contributions: A.E.V. and X.J.G. conceived and directed the study. A.E.V, N.E., C.C.C., S.K.,
and S.Q. performed the experiments. A.E.V. analyzed the data for the manuscript. A.E.V., C.C.C., and
X.J.G. wrote the manuscript. All authors provided feedback on the manuscript.

Figures:

417
Figure 1 – Harnessing protein motifs to control protein secretion with RELEASE: a) Co-translated
secreted proteins are normally transported through the conventional secretory pathway and secreted into
the extracellular space. The Retained Endoplasmic Cleavable Secretion (**RELEASE**) platform harnesses
b) native ER retention motifs (purple diamond) to retain tagged proteins of interest in the ER. Only through
the activation or expression of a protease such as TEVP (orange partial circle), the ER-retention motif will
be removed, and the protein of interest will be transported through the conventional secretory pathway.
Upon reaching the trans-golgi apparatus, the tagged protein of interest will be solubilized through cleavage
from the **c)** furin endoprotease and then secreted. **d)** Phosphorylation-dependent recruitment of
endogenous 14-3-3 scaffolding proteins to control the surface expression of membrane proteins. **e)**
Additional ER-retention motifs such as the diarginine (-RXR-) motif are compatible with RELEASE. Co-
expression of proteases such as TEVP (orange partial circle), or HCVP (yellow partial circle) with the
respective RELEASE constructs significantly increased SEAP secretion. **f)** The fusion of the 14-3-3
antagonist (R18 peptide) to the C-terminus of RELEASE inhibited ER retention activity and resulted in the

secretion of SEAP. In comparison, a fused R18 mutant which had two acidic residues required for 14-3-3 recruitment mutated into lysines (D12K and E14K) could not overcome the retention capabilities of RELEASE, and SEAP was not secreted. Each dot represents a biological replicate. Mean values were calculated from four replications (**d** and **f**). **The** error bars represent ± SEM. The results are representative of at least two independent experiments; significance was tested using an unpaired two-tailed Student's *t*-test between the two indicated conditions for each experiment. ****$p < 0.0001$.

Figure 2 – Functional completeness using Compact RELEASE: a) The RELEASE platform was modified to enable negative regulation of protein secretion (RELEASE-NOT) in response to protease activity. Without protease activity, the tagged protein of interest in secreted due to the antagonistic activity of the R18 peptide on the ER-retention motif. Proteolytic removal of the R18 peptide restores the ER-retention activity, resulting in the tagged protein of interest to be retained (**NOT** logic). **b)** With RELEASE-NOT, the co-expression of proteases such as TEVP resulted in a significant decrease in SEAP secretion. **c)** Due to the modularity of engineered proteins, RELEASE variants were engineered to enable binary logic gates. This collection of RELEASE variants is referred to as the Compact RELEASE (compRELEASE) suite. For each indicated gate, TEVP and HCVP served as binary inputs, which are either included or excluded in transfections. SEAP secretion served as an output. The design of each compRELEASE variant is shown above each respective graph. Each dot represents a biological replicate. Mean values were calculated from four replications (**c-d**). **The** error bars represent ± SEM. The results are representative of at least two independent experiments; significance was tested using an unpaired two-tailed Student's t-test between the two indicated conditions for each experiment. For experiments with multiple conditions, a one-way ANOVA with a Tukey's post-hoc comparison test was used to assess significance. ***$p < 0.001$. ****$p < 0.0001$.

454
Figure 3 – Quantitative processing using the Compact RELEASE suite: a) RELEASE and RELEASE-
NOT control the activation or repression of protein secretion in response to protease activity, respectively.
The cleavage efficiencies of RELEASE and RELEASE-NOT to TEVP were modified by creating constructs
using protease cut sites with different residues at the P1 position (bolded letter). The inverted triangle
indicates the position of proteolytic cleavage. **b)** Input-output curve of three RELEASE constructs (positive
regulation) with modified protease cut sites titrated with TEVP plasmid. Changing the TEVP cut site from
the canonical sequence reduced the cleavage efficiency and increased the activation threshold required
for protein secretion. **c)** The same changes in TEVP cut site sequences affected the repression thresholds
of proteins tagged with RELEASE-NOT. The data is fitted to a Hill equation with a variable slope
represented by the respective lines. **d)** Expected input-output curve of band-pass activation. **e)** To achieve
band-pass activation of protein secretion, a NIMPLY compRELEASE variant was modified to contain two
TEVP cut sites with different cutting efficiencies (K_{cat}). Band-pass filter activation requires the first TEVP
cut site (indicated as cut site 1) to have a higher K_{cat} than the second TEVP cut site (indicated as cut site
2). This ensures that at low concentrations of TEVP, the first cut site will be cleaved, and the tagged protein
of interest will be secreted. As the concentration of TEVP increases, the second cut site will be cleaved,
restoring the ER-retention activity and the protein of interest will be retained. **f)** Input-output curve of the
RELEASE band-pass construct titrated with TEVP plasmid. **g)** Expected input-output curve of band-stop
filter activation. **h)** To achieve band-stop activation of protein secretion, an IMPLY compRELEASE variant
designed with two TEVP cut sites with different cutting efficiencies was created. Under basal conditions
without TEVP, the tagged protein of interest will be secreted since the ER-retention motif is inhibited by the
R18 peptide. As the concentration of TEVP increases, the distal cut site will be preferentially cleaved
(indicated as cut site 1), restoring ER-retention activity and reducing SEAP secretion. If the concentrations
of TEVP continue to increase, then the proximal cut site will be cleaved (indicated as cut site 2) and the
tagged protein of interest will be secreted. **i)** Input-output curve of the RELEASE band-stop construct titrated
with TEVP plasmid. Each dot represents a biological replicate. Mean values were calculated from four

replications (**b-c, f and i**). The error bars represent ± SEM. The results are representative of at least two
independent experiments; significance was tested using an unpaired two-tailed Student's *t*-test between
the two indicated conditions for each experiment. For experiments with multiple conditions, a one-way
ANOVA with a Tukey's post-hoc comparison test was used to assess significance. ***$p < 0.001$. ****$p < 0.0001$.
484

485
Figure 4 – Delivery of RELEASE: a) Schematic of the stable integration of a polycistronic gene encoding
all the protein components required for inducible protein secretion via lentiviral transduction. **b)** Engineered
cells contain a split HCVP protease expressed in the cytoplasm to control the secretion of Nluc following
induction with rapalog. **c)** Input-output curves for engineered cells where secreted Nluc functions as the
output and rapalog concentration serves as the input. The curve represents a Hill equation with a variable
slope, and the grey box represents the top and bottom plateaus of the fit. To increase Nluc secretion, one
half of the split HCVP was localized to the ER via the p450-motif, however this design resulted in an
increase in background secretion and a reduction in the dynamic range. To reduce the background and
increase the dynamic range, the cleavage efficiency of the HCVP cut site of RELEASE was reduced by
modifying the flanking residues, as previously described[15]. **d)** mRNA delivery of a synthetic surface marker
(IgG) tagged with RELEASE and BFP into wildtype and engineered K562 cells. K562 cells were engineered
to stably express a rapalog-inducible TEVP protease and the mCherry reporter protein. **e**) Experimental
setup of RNA-delivered IgG-RELEASE into a mixture of wildtype and engineered K562 cells with and
without rapalog. Following 24 hours, the cells were harvested, stained for the surface expression of IgG

and processed through flow cytometry. **f)** Only engineered K562 cells containing the inducible split TEVP
showed a significant increase in the surface expression of IgG. Wildtype K562 cells did not show any
significant differences. Each dot represents a biological replicate. Mean values were calculated from four
replicates (**c** and **f**). The error bars represent ± SEM. The results are representative of at least two
independent experiments; significance was tested using a one-way ANOVA with a Tukey's post-hoc
comparison test was used to assess significance. ****$p < 0.0001$.

Supplementary Figures:

507
Supplementary Figure 1 – ER retention motifs in polyprotein constructs: a) Schematic of polyprotein
construct containing SEAP-RELEASE under the control of the dilysine retention motif (KKXX-COOH) and
the mNeon reporter protein separated via a T2A self-cleaving peptide. **b)** The retention capabilities of
RELEASE were negatively affected, and SEAP secretion was measured even in the absence of TEVP.
There was a significant increase in SEAP secretion when expressing TEVP, however the dynamic range
was lower than previously reported[15]. **c)** Schematic of polyprotein construct containing SEAP under the
control of the diarginine retention motif (-RXR-) and the mNeon reporter protein separated via a T2A self-
cleaving peptide. **d)** The retention capabilities of these RELEASE variant using the diarginine retention
motif were intact and we observed a significant increase in SEAP secretion when co-expressing TEVP.
Each dot represents a biological replicate. Mean values were calculated from four biological replicates (**b**,
d). The error bars represent ± SEM. The results are representative of at least two independent experiments;
significance was tested using an unpaired two-tailed Student's t-test between the two indicated conditions.
**$p < 0.01$, ****$p < 0.0001$.

521
<u>Supplementary Figure 2 – Fusing 14-3-3ζ to block ER-retention activity of RELEASE:</u> a) Schematic
of the standard RELEASE construct and a RELEASE-construct fused to the 14-3-3ζ protein. **b)** In the
absence of protease, there was a significant increase in SEAP secretion when transiently transfecting the
RELEASE-14-3-3 fusion compared to the standard RELEASE alone. Each dot represents a biological
replicate. Mean values were calculated from four biological replicates (**b**, **d**). The error bars represent ±
SEM. The results are representative of at least two independent experiments; significance was tested using
an unpaired two-tailed Student's *t*-test between the two indicated conditions. **p < 0.01, ****p < 0.0001.

529
Supplementary Figure 3 – Negative regulation of protein secretion using CHOMP: a) Schematic of
CHOMP circuit for negative regulation of protein secretion. A HCVP-inducible RELEASE construct controls
the secretion of SEAP. The HCVP protease can be repressed by TEVP, by cleaving the co-peptide of
HCVP, which will significantly reduce its activity. **b)** SEAP secretion was dependent on the co-expression
of HCVP. Increasing the amount of TEVP significantly reduced SEAP secretion, however negative
regulation of SEAP secretion plateaued and could not reach basal levels. Each dot represents a biological
replicate. Mean values were calculated from four replicates (**b**). The error bars represent ± SEM. The results
are representative of at least two independent experiments; significance was tested using a one-way
ANOVA with a Tukey's post-hoc comparison test was used to assess significance. ****$p < 0.0001$.

539
Supplementary Figure 4 – HCVP-inducible RELEASE-NOT: a) Schematic of HCVP-inducible
RELEASE-NOT. When co-expressed with HCVP, SEAP secretion decreased relative to when HCVP was
absent. Each dot represents a biological replicate. Mean values were calculated from four biological
replicates +/- SEM. The results are representative of at least two independent experiments; significance
was tested using an unpaired two-tailed Student's t-test between the two indicated conditions for each
experiment. **** = $p < 0.0001$.

546
Supplementary Figure 5 – Exogenous expression of 14-3-3ζ to boost activity of RELEASE NOT: a)
Mechanism of action for RELEASE-NOT to control protein secretion in the absence of protease. **b)**
Schematic of adding exogenous 14-3-3 scaffolding proteins to boost the secretion of RELEASE-NOT in the
absence of protease. **c)** Increasing the amount of exogenous 14-3-3ζ resulted in a significant increase in
SEAP secretion with and without the co-expression of HCVP. **d)** Although we observed an increase in
SEAP secretion as increasing the amount of exogenous 14-3-3ζ, the fold change between the ON and OFF
state decreased. 14-3-3ζ also increased SEAP secretion in the presence of HCVP, resulting in a reduction
in the overall fold change as 14-3-3ζ increased. Each dot represents a biological replicate. Mean values
were calculated from four replicates (**b-c**). The error bars represent ± SEM. The results are representative
of at least two independent experiments; significance was tested using a one-way ANOVA with a Tukey's
post-hoc comparison test was used to assess significance. ***$p < 0.001$, ****$p < 0.0001$.

558
Supplementary Figure 6 – Phosphorylation-dependent recruitment of 14-3-3 proteins to program
protein secretion: a) Schematic of RELEASE-NOT using the SWTY motif to enable phosphorylation-
dependent recruitment of 14-3-3 proteins to block the retention activity of RXR. **b)** Schematic of RELEASE-
NOT where the phospho-responsive threonine residue is mutated to alanine (SWTY-COOH → SWAY-
COOH). **c)** When co-expressing TEVP we observed a significant decrease in SEAP secretion only when
using the RELEASE-NOT construct with the SWTY motif. RELEASE-NOT with the SWAY motif did not
result in any secretion of SEAP. Each dot represents a biological replicate. Mean values were calculated
from four replicates. The error bars represent ± SEM. The results are representative of at least two
independent experiments; significance was tested using a one-way ANOVA with a Tukey's post-hoc
comparison test was used to assess significance. ****$p < 0.0001$.

569
Supplementary Figure 7 – R18 function is compatible when encoded by polycistronic constructs:
a) RELEASE-NOT encoded by a polycistronic construct separated by self-cleaving T2A peptides is compatible with negative regulation of protein secretion. Each dot represents a biological replicate. Mean values were calculated from four biological replicates +/- SEM. The results are representative of at least two independent experiments; significance was tested using an unpaired two-tailed Student's t-test between the two indicated conditions for each experiment. **** = $p < 0.0001$.

Retained   Retained   Secreted   Retained

Supplementary Figure 8 – Schematic of RELEASE-NIMPLY: a) In the absence of any protease, the tagged protein of interest is retained due to the ER-retention capabilities of the N-terminal signal anchor of cytochrome p450. **b)** When TEVP is co-expressed, the R18 peptide is removed restoring the retention capabilities of the RXR ER-retention motif and the protein of interest is retained. **c)** When HCVP is co-expressed, the N-terminal ER-retention motif is removed, and the C-terminal retention motif is being inhibited by the recruitment of 14-3-3 proteins via the intact R18 peptide. This results in the tagged protein of interest being secreted. **d**) Co-expression of HCVP and TEVP resulted in the tagged protein of interest being retained. Although the N-terminal ER-retention motif is removed by HCVP, the activity of the C-terminal ER-retention motif is restored by removal of the R18 peptide.

References:

1. Roybal, K. T. & Lim, W. A. Synthetic immunology: hacking immune cells to expand their therapeutic capabilities. *Annu. Rev. Immunol.* **35,** 229–253 (2017).
2. Lim, W. A. Designing customized cell signalling circuits. *Nat. Rev. Mol. Cell Biol.* **11,** 393–403 (2010).
3. Dueber, J. E., Yeh, B. J., Chak, K. & Lim, W. A. Reprogramming control of an allosteric signaling switch through modular recombination. *Science* **301,** 1904–1908 (2003).
4. Elowitz, M. & Lim, W. A. Build life to understand it. *Nature* **468,** 889–890 (2010).
5. Roybal, K. T. *et al.* Precision Tumor Recognition by T Cells With Combinatorial Antigen-Sensing Circuits. *Cell* **164,** 770–779 (2016).
6. Allen, G. M. *et al.* Synthetic cytokine circuits that drive T cells into immune-excluded tumors. *Science* **378,** eaba1624 (2022).
7. Ye, H. *et al.* Self-adjusting synthetic gene circuit for correcting insulin resistance. *Nat. Biomed. Eng.* **1,** 0005 (2017).
8. Cho, J. H. *et al.* Engineering advanced logic and distributed computing in human CAR immune cells. *Nat. Commun.* **12,** 792 (2021).
9. Hernandez-Lopez, R. A. *et al.* T cell circuits that sense antigen density with an ultrasensitive threshold. *Science* **371,** 1166–1171 (2021).
10. Bashor, C. J., Horwitz, A. A., Peisajovich, S. G. & Lim, W. A. Rewiring cells: synthetic biology as a tool to interrogate the organizational principles of living systems. *Annu. Rev. Biophys.* **39,** 515–537 (2010).
11. Morsut, L. *et al.* Engineering customized cell sensing and response behaviors using synthetic notch receptors. *Cell* **164,** 780–791 (2016).
12. Gao, X. J., Chong, L. S., Kim, M. S. & Elowitz, M. B. Programmable protein circuits in living cells. *Science* **361,** 1252–1258 (2018).
13. Fink, T. *et al.* Design of fast proteolysis-based signaling and logic circuits in mammalian cells. *Nat. Chem. Biol.* **15,** 115–122 (2019).
14. Chen, Z. & Elowitz, M. B. Programmable protein circuit design. *Cell* **184,** 2284–2301 (2021).
15. Vlahos, A. E. *et al.* Protease-controlled secretion and display of intercellular signals. *Nat. Commun.* **13,** 912 (2022).
16. Praznik, A. *et al.* Regulation of protein secretion through chemical regulation of endoplasmic reticulum retention signal cleavage. *Nat. Commun.* **13,** 1323 (2022).
17. Ghosh, S., Brown, A. M., Jenkins, C. & Campbell, K. Viral vector systems for gene therapy: A comprehensive literature review of progress and biosafety challenges. *Applied Biosafety* **25,** 7–18 (2020).
18. Patwardhan, A., Cheng, N. & Trejo, J. Post-Translational Modifications of G Protein-Coupled Receptors Control Cellular Signaling Dynamics in Space and Time. *Pharmacol. Rev.* **73,** 120–151 (2021).
19. Ardito, F., Giuliani, M., Perrone, D., Troiano, G. & Lo Muzio, L. The crucial role of protein phosphorylation in cell signaling and its use as targeted therapy (Review). *Int. J. Mol. Med.* **40,** 271–280 (2017).
20. Proud, C. G. Phosphorylation and signal transduction pathways in translational control. *Cold Spring Harb. Perspect. Biol.* **11,** (2019).
21. Pennington, K. L., Chan, T. Y., Torres, M. P. & Andersen, J. L. The dynamic and stress-adaptive signaling hub of 14-3-3: emerging mechanisms of regulation and context-dependent protein-protein interactions. *Oncogene* **37,** 5587–5604 (2018).
22. Luo, D. *et al.* Interacting domains of P14-3-3 and actin involved in protein-protein interactions of living cells. *Arch. Microbiol.* **193,** 651–663 (2011).
23. Wang, W. & Shakes, D. C. Molecular evolution of the 14-3-3 protein family. *J. Mol. Evol.* **43,** 384–398 (1996).
24. Fan, X. *et al.* 14-3-3 Proteins Are on the Crossroads of Cancer, Aging, and Age-Related

638  Neurodegenerative Disease. *Int. J. Mol. Sci.* **20,** (2019).

639 25. Heusser, K. *et al.* Scavenging of 14-3-3 proteins reveals their involvement in the cell-surface
640  transport of ATP-sensitive K+ channels. *J. Cell Sci.* **119,** 4353–4363 (2006).

641 26. Rajan, S. *et al.* Interaction with 14-3-3 proteins promotes functional expression of the potassium
642  channels TASK-1 and TASK-3. *J. Physiol. (Lond.)* **545,** 13–26 (2002).

643 27. Mrowiec, T. & Schwappach, B. 14-3-3 proteins in membrane protein transport. *Biol. Chem.* **387,**
644  1227–1236 (2006).

645 28. Coblitz, B. *et al.* C-terminal recognition by 14-3-3 proteins for surface expression of membrane
646  receptors. *J. Biol. Chem.* **280,** 36263–36272 (2005).

647 29. Chung, J.-J. *et al.* PI3K/Akt signalling-mediated protein surface expression sensed by 14-3-3
648  interacting motif. *FEBS J.* **276,** 5547–5558 (2009).

649 30. Rong, J. *et al.* 14-3-3 protein interacts with Huntingtin-associated protein 1 and regulates its
650  trafficking. *J. Biol. Chem.* **282,** 4748–4756 (2007).

651 31. Fink, T. & Jerala, R. Designed protease-based signaling networks. *Curr. Opin. Chem. Biol.* **68,**
652  102146 (2022).

653 32. Girod, A. *et al.* Evidence for a COP-I-independent transport route from the Golgi complex to the
654  endoplasmic reticulum. *Nat. Cell Biol.* **1,** 423–430 (1999).

655 33. Cole, N. B., Ellenberg, J., Song, J., DiEuliis, D. & Lippincott-Schwartz, J. Retrograde transport of
656  Golgi-localized proteins to the ER. *J. Cell Biol.* **140,** 1–15 (1998).

657 34. Jackson, L. P. *et al.* Molecular basis for recognition of dilysine trafficking motifs by COPI. *Dev. Cell*
658  **23,** 1255–1262 (2012).

659 35. Shiwarski, D. J., Crilly, S. E., Dates, A. & Puthenveedu, M. A. Dual RXR motifs regulate nerve
660  growth factor-mediated intracellular retention of the delta opioid receptor. *Mol. Biol. Cell* **30,** 680–
661  690 (2019).

662 36. Jackson, M. R., Nilsson, T. & Peterson, P. A. Identification of a consensus motif for retention of
663  transmembrane proteins in the endoplasmic reticulum. *EMBO J.* **9,** 3153–3162 (1990).

664 37. Szymczak, A. L. *et al.* Correction of multi-gene deficiency in vivo using a single "self-cleaving" 2A
665  peptide-based retroviral vector. *Nat. Biotechnol.* **22,** 589–594 (2004).

666 38. Shikano, S. & Li, M. Membrane receptor trafficking: evidence of proximal and distal zones
667  conferred by two independent endoplasmic reticulum localization signals. *Proc. Natl. Acad. Sci.*
668  *USA* **100,** 5783–5788 (2003).

669 39. Yuan, H., Michelsen, K. & Schwappach, B. 14-3-3 dimers probe the assembly status of multimeric
670  membrane proteins. *Curr. Biol.* **13,** 638–646 (2003).

671 40. Wang, B. *et al.* Isolation of high-affinity peptide antagonists of 14-3-3 proteins by phage display.
672  *Biochemistry* **38,** 12499–12504 (1999).

673 41. Petosa, C. *et al.* 14-3-3zeta binds a phosphorylated Raf peptide and an unphosphorylated peptide via
674  its conserved amphipathic groove. *J. Biol. Chem.* **273,** 16305–16310 (1998).

675 42. Shikano, S., Coblitz, B., Sun, H. & Li, M. Genetic isolation of transport signals directing cell surface
676  expression. *Nat. Cell Biol.* **7,** 985–992 (2005).

677 43. Park, J. W., Reed, J. R., Brignac-Huber, L. M. & Backes, W. L. Cytochrome P450 system proteins
678  reside in different regions of the endoplasmic reticulum. *Biochem. J.* **464,** 241–249 (2014).

679 44. Szczesna-Skorupa, E. & Kemper, B. Endoplasmic reticulum retention determinants in the
680  transmembrane and linker domains of cytochrome P450 2C1. *J. Biol. Chem.* **275,** 19409–19415
681  (2000).

682 45. Holt, B. A. & Kwong, G. A. Protease circuits for processing biological information. *Nat. Commun.*
683  **11,** 5021 (2020).

684 46. Hart, Y. *et al.* Paradoxical signaling by a secreted molecule leads to homeostasis of cell levels. *Cell*
685  **158,** 1022–1032 (2014).

686 47. McFarlane, A., Pohler, E. & Moraga, I. Molecular and cellular factors determining the functional
687  pleiotropy of cytokines. *FEBS J.* **290,** 2525–2552 (2023).

688 48. Martinez-Sanchez, M. E., Huerta, L., Alvarez-Buylla, E. R. & Villarreal Luján, C. Role of cytokine

combinations on CD4+ T cell differentiation, partial polarization, and plasticity: continuous network
modeling approach. *Front. Physiol.* **9,** 877 (2018).
49. Biernacka, A., Dobaczewski, M. & Frangogiannis, N. G. TGF-β signaling in fibrosis. *Growth*
Factors **29,** 196–202 (2011).
50. Kapust, R. B., Tözsér, J., Copeland, T. D. & Waugh, D. S. The P1' specificity of tobacco etch virus
protease. *Biochem. Biophys. Res. Commun.* **294,** 949–955 (2002).
51. Xie, M. *et al.* β-cell-mimetic designer cells provide closed-loop glycemic control. *Science* **354,**
1296–1301 (2016).
52. Xie, M. & Fussenegger, M. Designing cell function: assembly of synthetic gene circuits for cell
biology applications. *Nat. Rev. Mol. Cell Biol.* **19,** 507–525 (2018).
53. Kowalski, P. S., Rudra, A., Miao, L. & Anderson, D. G. Delivering the Messenger: Advances in
Technologies for Therapeutic mRNA Delivery. *Mol. Ther.* **27,** 710–728 (2019).
54. Tycko, J. *et al.* High-Throughput Discovery and Characterization of Human Transcriptional
Effectors. *Cell* **183,** 2020–2035.e16 (2020).
55. Daringer, N. M., Dudek, R. M., Schwarz, K. A. & Leonard, J. N. Modular extracellular sensor
architecture for engineering mammalian cell-based devices. *ACS Synth. Biol.* **3,** 892–902 (2014).
56. Kipniss, N. H. *et al.* Engineering cell sensing and responses using a GPCR-coupled CRISPR-Cas
system. *Nat. Commun.* **8,** 2212 (2017).
57. Ballone, A., Centorrino, F. & Ottmann, C. 14-3-3: A Case Study in PPI Modulation. *Molecules* **23,**
(2018).
58. Pozuelo Rubio, M. *et al.* 14-3-3-affinity purification of over 200 human phosphoproteins reveals
new links to regulation of cellular metabolism, proliferation and trafficking. *Biochem. J.* **379,** 395–
408 (2004).
59. Diallo, K., Oppong, A. K. & Lim, G. E. Can 14-3-3 proteins serve as therapeutic targets for the
treatment of metabolic diseases? *Pharmacol. Res.* **139,** 199–206 (2019).
60. Koh, E. Y. C. *et al.* An internal ribosome entry site (IRES) mutant library for tuning expression level
of multiple genes in mammalian cells. *PLoS One* **8,** e82100 (2013).
61. Liu, Z. *et al.* Systematic comparison of 2A peptides for cloning multi-genes in a polycistronic vector.
Sci. Rep. **7,** 2193 (2017).
718