1 **Sexual dimorphism of insular cortex function in persistent alcohol drinking despite**
2 **aversion in mice**
3

4 Claudia Fornari[1], Carmen Guerrero-Márquez[1], Praneeth Namburi[2], Yoni Couderc[1], Céline
5 Nicolas[1*] and Anna Beyeler[1*]
6
Affiliations:
[1] University of Bordeaux, INSERM, Neurocentre Magendie, U1215, 146 rue Léo Saignat, 33000,
Bordeaux, France.
[2] Institute for Medical Engineering & Science, Massachusetts Institute of Technology, 77
Massachusetts Ave., Cambridge, MA, 02139, USA.
*The authors scientifically contributed equally to this work

Send correspondence to anna.beyeler@inserm.fr or celine.nicolas@u-bordeaux.fr
Short Title: Sex differences in persistent alcohol drinking
Keywords: sex differences, binge drinking, persistent drinking, insular cortex, chemogenetic,
calcium fiber photometry

17
18 **ABSTRACT**:
19

BACKGROUND: One major hallmark of alcohol use disorder (AUD) is the persistence of alcohol
drinking despite negative consequences. Among the indicators of AUD vulnerability, binge
drinking is a strong risk factor. Although the lifetime prevalence of binge and AUD has been
historically higher in men than women, this gap dramatically narrowed in the last decade.
Additionally, sex differences in AUD and binge drinking have been shown in clinical and preclinical
studies, respectively. The insular cortex plays an important role in AUD, and the anterior (aIC)
and posterior (pIC) divisions have dimorphic functions. However, the contributions of the aIC and
pIC sections in alcohol binge drinking and alcohol persistent drinking despite aversion, as well as
the sexual dimorphism of these contributions, remained to be uncovered.
29
30 METHODS: First, by combining the drinking in the dark model with chemogenetics, we studied
31 the causal role of aIC and pIC excitatory neurons in binge and persistent ethanol drinking in
32 C57BL6/J male (n=49) and female (n=49) mice. Second, using calcium fiber photometry, we
33 investigated pIC neuronal activity in both sexes (male n=14, female n=11) during both binge and
34 persistent ethanol drinking.
35
RESULTS: We identified a higher binge and persistent ethanol consumption in females compared
to males. Chemogenetic inhibition of aIC glutamatergic neurons reduced bitter solutions intake
independently of the solvent (ethanol or water), in both sexes. In contrast, inhibition of pIC
glutamatergic neurons exclusively reduced persistent ethanol drinking in female mice. Finally,
using fiber photometry recordings, we uncovered that pIC glutamatergic neuron activity was
selectivity increased during ethanol persistent drinking in female mice.
42
43 CONCLUSIONS: These findings suggest a sex-dependent function of the pIC in persistent
44 ethanol drinking, providing a starting point in our understanding of the insular cortex function in
45 the neurobiology of AUD in both sexes.

46
47
48

INTRODUCTION

Alcohol use disorder (AUD) is a chronic relapsing disorder defined by the persistence of excessive alcohol consumption despite physical and psychological negative consequences (1), which are a major impediment to AUD treatment (2,3). AUD is a major public health burden as 6% of the world population is affected by alcohol morbidity or mortality (4). Among the indicators of vulnerability to AUD, binge drinking, an episodic and excessive pattern of alcohol consumption leading to the intoxication threshold of 80 mg/dL, has been identified as a strong risk factor (3,5–7). Indeed, a longitudinal study showed that 43% of adolescents (13 to 18 years old) that binge-drink alcohol developed AUD at the age of 21, whereas only 7% of non-bingers developed AUD at the same age (8). In preclinical studies, binge-drinking is modeled using the drinking in the dark (DID) procedure, where mice have repeated access to alcohol for a limited time during the dark phase of the circadian rhythm (active period) which allows to reach blood ethanol concentration (BEC) of 80 mg/dL as observed in humans (8–10). Interestingly, this model can be adapted to study the persistence of alcohol drinking despite aversive outcomes, for example by adulterating ethanol with quinine, a bitter and aversive substance to mice (11–14).

Historically, men have been more likely than women to drink alcohol and exhibit pathological drinking behaviors. In 2020, AUD lifetime prevalence was still higher in men than women, reaching respectively 36% and 23% (15). However, a recent longitudinal study showed that the increase rate of AUD over a decade was drastically higher in women than men, with respective rates of 84% and 35% (16). These epidemiological data highlight that the sex difference of AUD prevalence is narrowing. Furthermore, women transition faster to AUD after regular or chronic alcohol consumption (17,18) and are subsequently more likely to develop alcohol-related diseases (e.g. cardiovascular and hepatic diseases) (19–21). Interestingly, preclinical studies revealed sex differences in alcohol intake, with female rodents drinking more than males across different models of intake, including binge drinking and drinking despite taste aversion (12,22–

29). This higher propensity to alcohol drinking in females is proposed to be, at least in part, due to their lowest sensitivity to aversive properties of alcohol, as female rodents are more resistant to ethanol conditioned taste aversion (30,31).

The insular cortex (or insula), is strongly involved in drug addiction (32–34), including AUD (35–37). Indeed, human functional imaging studies have demonstrated that alcohol cues trigger greater activity responses in the insula, in alcohol dependent subjects (38,39). Furthermore, insula white matter volume is linked to binge drinking frequency in adolescents (40). Indeed, a higher white matter volume in the left insula was associated with stronger drinking motivations, and therefore a greater frequency of binge drinking. Anatomically, the insula is subdivided into the anterior (aIC) and posterior (pIC) sections, which are proposed to have antagonistic functions in a wide range of behaviors (41–43). In AUD patients, structural studies report a reduction of the aIC volume and gray matter density compared to healthy controls, which is correlated with higher compulsive drinking measures (44–47). In addition, aIC functional connectivity with several brain regions (e.g. hippocampus, medial orbitofrontal regions) is higher in patients with AUD compared to social drinkers (48) and healthy controls (49). Intriguingly, less information is known about the role of pIC in AUD. Indeed, only two recent studies reported a decrease of pIC gray matter volumes in AUD patients (49,50) and a higher resting state of the pIC relative to healthy controls (49), suggesting that the pIC plays a role in AUD as well. Preclinical studies confirmed the role of both aIC and pIC in ethanol drinking behaviors. Indeed, chemogenetic activation or inhibition of aIC neurons respectively decrease and increase alcohol intake in several behavioral paradigms such as self-administration and intermittent 2-bottle choice (51–53). In contrast, pharmacological inactivation of pIC neurons decreases alcohol self-administration (54), suggesting divergent functions of aIC and pIC in alcohol drinking. However, how sex as a biological variable influence those divergent contributions remained untested.

Recent findings point towards structural sexual dimorphism of the insula, with a larger volume in

men than women in physiological conditions, whereas in AUD long-term abstinent patients this

ratio is inverted (47). In support of these clinical findings, preclinical studies highlighted sex

differences of insula functions in alcohol drinking behaviors. For example, in alcohol binge

drinking mice, optogenetic stimulation of aIC terminals in the dorso-lateral striatum induced larger

excitatory post-synaptic currents and smaller AMPA/NMDA current ratio recorded in medium

spiny neurons compared to water drinking mice (55). Interestingly, this synaptic plasticity was

observed in males but not female mice (55). Additionally, in mice, short-term ethanol exposure

increases the excitability of pIC projection neurons targeting the bed nucleus of the stria terminalis

neurons in females, but not males (56). Although the literature pinpoints the insular cortex as a

key player in alcohol-related behaviors, the sexual dimorphism of aIC and pIC function in binge

and/or persistent alcohol intake despite quinine aversion remains to be uncovered.

Using an adapted DID model, we repeatedly demonstrated a higher binge and persistent ethanol

consumption despite aversion in female compared to male mice. Combining the DID protocol with

chemogenetics in a within-subject experimental design, we showed that inhibition of aIC

glutamatergic neurons reduced bitter solutions intake independently of the solvent (e.g. ethanol

or water), in both sexes. In contrast, pIC glutamatergic neuron inhibition exclusively reduced

persistent ethanol drinking in female mice. Then, using *in vivo* calcium fiber photometry

recordings, we identified that pIC excitatory neurons are activated during drinking of all liquids

tested, when we did not consider sex as a variable. However, when comparing males and

females, we uncovered that pIC glutamatergic neuron activity was selectivity increased in female

mice during ethanol persistent drinking.

Altogether, we identified a sex-dependent function of the pIC in persistent ethanol drinking,

providing a starting point in our understanding of the insular cortex function in the neurobiology of

AUD in both sexes.

METHODS AND MATERIALS

For additional details on the methods, see the Supplement.

Stereotactic surgeries

Chemogenetic viral injection. The adeno-associated viruses encoding the inhibitory hM4Di receptor coupled to mCherry under the control of CaMKII promoter (AAV$_9$-CaMKIIa-hM4D(Gi)-mCherry, Addgene), to target glutamatergic neurons, or the control viral vector (AAV9/2-mCaMKII-mCherry-WPRE, ETH Zürich) were injected bilaterally (200-250 nL, 1nL/sec) in the aIC (antero-posterior +1.7mm; medial-lateral ±3.1mm; dorso-ventral -3.5mm from the bregma) or in the pIC (antero-posterior -0.35mm; medial-lateral ±4.0mm; dorso-ventral -4.2mm from the bregma).

Calcium fiber photometry viral injection and fiber implantation. A viral vector coding for the calcium sensor GCaMP6f (AAV$_9$-CaMKII-GCaMP6f-WPRE-SV40, Addgene) was injected unilaterally (250 nl, 1nl/sec) in the right pIC. An optic fiber (400µm diameter, 0.48 NA, >90% efficiency) inserted in a ceramic ferrule was implanted 50µm above the virus injection site.

Drinking in the dark procedure (DID)

The DID protocol is based on the circadian rhythm where animals have access to different liquids (e.g. ethanol) during the nocturnal period (8,9). Two and a half hours after the onset of the dark phase, the water bottle in the home cage was replaced by different liquids for 2 hours per day for 4 consecutive days, followed by 3 days with water access only. This 7-day cycle is repeated for 4 weeks (**Figure 2C**), starting with tap water (cycle 0), then ethanol (20% v/v in tap water, cycles 1 and 2) to study binge drinking behavior, and ethanol adulterated with quinine (500µM of quinine, cycle 3) to study persistent drinking behavior despite aversion. Finally, water or water adulterated with quinine (500µM quinine) consumption was measured during a single 2-hour session and served as control. Daily liquid consumption was obtained by weighting the bottle before and after the 2-hour session, and mice were weighed the first day of each cycle.

[Blood ethanol concentration measurement.](#) To measure blood ethanol concentrations (BEC),

blood sampling was performed by incision of the lateral tail vein 30 minutes or 2 hours after the

onset of the ethanol session. The samples were immediately centrifuged at 10,000 rpm for 10

minutes at 4°C. The BEC in mg/dL was obtained from the serum using the Analox GL5 analyzer.

We excluded 3 males and 3 females from the BEC analysis after 2-hour ethanol intake because

the blood sample did not contain enough serum.

Chemogenetic inhibition of aIC or pIC excitatory neurons during binge and persistent

ethanol drinking

The same DID procedure as described above was used. Additionally, a subgroup of animals

underwent three more days of water+quinine followed by a last cycle of ethanol+quinine to control

the resumption of persistent ethanol drinking. The day before the chemogenetic manipulation an

intraperitoneal (i.p.) injection of vehicle (NaCl, 0.1 mL/kg) was performed 30 minutes before the

drinking session for habituation. The causal role of aIC or pIC glutamatergic neurons on drinking

was tested after injection of CNO (i.p. 3 mg/kg in NaCl, Tocris) or vehicle 30 minutes before the

last session of binge (day 18), persistent ethanol drinking (day 25) and a single session of water

or water+quinine (day 29). To note, mice tested for chemogenetic inhibition of pIC on

water+quinine drinking (day 29) were CNO naive.

Coding properties of pIC excitatory neurons during binge and persistent ethanol drinking

The same DID procedure as described before was used, except that no initial water cycle was

performed. To study the coding properties of pIC neurons during binge and persistent ethanol

drinking, the calcium signal and the drinking behavior (e.g. licking) has to be synchronized. Thus,

after two ethanol binge cycles in their home cage, mice (n=14 males and n=11 females)

underwent three cycles of ethanol drinking (cycle 3 to 5) in multifunctional boxes equipped with

lickometers (**Figure S2A**). Then, after one cycle of ethanol+quinine drinking in the home cages

(cycle 6), mice underwent a cycle of ethanol+quinine drinking in the multifunctional boxes (cycle

7). Finally, a water and water+quinine photometry recording session was performed in these

boxes as a control. To note 3 male and 2 female ethanol naive mice were included in water

recording analysis and 4 male and 2 female ethanol naive mice were included in water+quinine

recording analysis.

Fiber photometry recordings. The recordings were performed with Neurophotometrics fiber

photometry system (FP3002 V2), on the last day of the last ethanol and ethanol+quinine cycles,

and during a single water and water+quinine session (**Figure S2A**). The recordings were

performed using a 4-branch patch cord (400μm diameter, NA=0.48, 1 branch per animal). Then,

the drinking sprout was filled with the appropriate solution, and the recording lasted 1 hour as

previously done (33). Mice were kept in the boxes for an additional hour for a total of a 2-hour

session. The drinking sprout was automatically refilled to give *ad-libitum* access to the mice. The

optimal wavelength for GCaMP6f excitation is 470 nm, while the isosbestic wavelength is 415 nm.

To minimize the photobleaching effect of the recording, the light intensity at the tip of the patch

cord was adjusted between 80 and 100 μW for both channels.

Calcium fiber photometry data analysis. Photometry recordings were analyzed using custom

Python scripts which can be downloaded at:

https://github.com/praneethnamburi/photometry_collab_ab. The timestamps for lick events

were extracted and classified in bouts. A bout event is defined as a lick or a group of licks with an

inter lick interval (ILI) less than or equal to 10 seconds, for the peri-event analysis. The first 10

minutes of signals were discarded. Then, GCaMP and isosbestic signals were de-trended using

the airPLS algorithm (57), and the detrended isosbestic signal was regressed from the detrended

GCaMP signal. The resulting signal was bandpass filtered with cutoff frequencies of 0.2 and 6 Hz

using a finite impulse response filter implemented in the scipy package. Finally, a time window of

-10 to +10 s was defined for the peri-event analysis of the resulting signal aligned to the first lick

of each bout. Signal from each window was z-normalized by subtracting the mean of the baseline

and dividing the result by the standard deviation of the baseline, where the baseline signal was

defined as the signal from 10 s to 8 s before a lick bout onset. Z-normalized calcium signals were

averaged for each male and female mouse during the reference (-8 to -5 sec) and licking window

(0 to +3 sec) to describe the signal changes.

RESULTS

Sexual dimorphism of ethanol binge drinking: validation of the drinking in the dark model

After 2 ethanol cycles (**Figure 1A**), the average of ethanol consumption over 2-hour access was

higher in females than males (**Figure 1B,** unpaired Student t-test, t=3.864, p=0.0007) whereas

the BEC levels were similar between sexes (**Figure 1C**), with 23% of males and 15% of females

reaching the binge intoxication threshold of 80 mg/dL. The literature suggests that mice drink the

majority of the total ethanol intake at the beginning of the DID sessions (10). Thus, we measured

the proportion of ethanol consumed during the first 30-minutes of the 2-hour session and we

showed that males consumed 67% and females 84% of the total ethanol intake during this period

(**Figure 1D**). Moreover, in an additional session, ethanol intake and BEC were measured after 30

minutes of ethanol access and revealed a higher ethanol intake (**Figure 1E**, unpaired Student t-

test, t=3.679, p=0.0009) and BEC (**Figure 1F**, unpaired Mann-Whitney, U=71, p=0.0318) in

females than males with a respective proportion of 75% and 44% of mice reaching the intoxication

threshold of 80 mg/dL which validates the model.

Chemogenetic inhibition of aIC and pIC glutamatergic neurons during binge and persistent

ethanol drinking

At the behavioral level, for all liquids during the sessions without chemogenetic manipulation,

animals injected into aIC or pIC showed similar results, so we grouped them together (**Figure 3A-**

H). Across the sessions, both male and female mice had similar levels of water intake as well as

average water intake (**Figure 3A, B**). In contrast, female mice consumed a larger amount of

ethanol across the sessions (**Figure 3C**, 2-way RM-ANOVA, sex factor $F_{1,97}=94.91$, p<0.0001

and session factor $F_{5.192, 503.7}=4.629$, p=0.0003 without interaction $F_{6, 582}=0.5145$, p=0.7976) and

on average than males (**Figure 3D**, unpaired Student t-test, t=9.742, p<0.0001) although the

absolute volume of ethanol solution was similar between sexes (**Figure S1A**). Similarly, when

ethanol was adulterated with quinine, female mice had a higher consumption than males across

the sessions (**Figure 3E**, 2-way RM-ANOVA, sex factor $F_{1, 96}=94.71$, p<0.0001 and session factor

$F_{1.880, 180.5}=4.931$, p=0.0095 without interaction $F_{2, 192}=1.233$, p=0.2938), on average (**Figure 3F**,

unpaired Student t-test, t=9.732, p<0.0001) as well as on absolute volume (**Figure S1B**, unpaired

Student t-test, t=4.472, p<0.0001). Both ethanol and ethanol adulterated with quinine drinking

were independent of the hormonal cycle in female mice (**Figure S1C, D**). To test the strength of

persistent ethanol drinking despite quinine aversion, a subgroup of mice had access to

water+quinine for 3 sessions, followed by 4 sessions of ethanol+quinine access. Both males and

females show an increase of ethanol+quinine intake after exposure to 3 days of quinine diluted in

water (**Figure 3G**, 2-way RM-ANOVA, sex factor $F_{1, 35}=11.00$, p=0.0021, session factor

$F_{6, 210}=17.35$, p<0.0001, with an interaction $F_{6, 210}=3.316$, p=0.0039). Interestingly, in both sexes,

the average of quinine solution consumed was larger when quinine was mixed to ethanol, than

when quinine was diluted in water (**Figure 3H**, 2-way RM-ANOVA, sex factor $F_{1, 46}=17.32$,

p=0.0001, liquid factor $F_{1, 46}=52.47$, p<0.0001, with a sex x liquid interaction $F_{1, 46}=8.905$,

p=0.0045). These results validate the aversiveness of quinine and that ethanol reintroduction to

quinine is sufficient to resume ethanol persistent drinking in both sexes.

To study the causal role of aIC and pIC neurons in binge and persistent ethanol drinking, we used

a chemogenetic approach (**Figure 2A-C**), that we validated using *in-vivo* electrophysiology

(**Figure S1E, F**, paired Wilcoxon test, p=0.023), to inhibit aIC or pIC glutamatergic neurons by

expressing hM4Di, under the CaMKIIα promoter, before the mice underwent the behavioral

protocol. Then, we investigated the effect of chemogenetic inhibition of aIC or pIC excitatory

neurons on binge and persistent ethanol intake. Chemogenetic inhibition of aIC or pIC

glutamatergic neurons (**Figure S1M, N**) do not influence ethanol drinking in both sexes (**Figure**

4A, 2-way ANOVA, sex factor $F_{1, 42}=33.65$, $p<0.0001$, virus factor $F_{1, 42}=0.9755$, $p=0.3290$, with

no sex x virus interaction $F_{1, 42}=1.340$, $p=0.2535$, **Figure 4F**, 2-way ANOVA, sex factor

$F_{1, 26}=27.71$, $p<0.0001$, virus factor $F_{1, 26}=0.0001289$, $p=0.9910$, without sex x virus interaction

$F_{1, 26}=0.2937$, $p=0.5925$). In contrast, the inhibition of aIC excitatory neurons reduced

ethanol+quinine drinking independently of the sex (**Figure 4B**, 2-way ANOVA, sex factor

$F_{1, 41}=18.71$, $p<0.0001$, virus factor $F_{1, 41}=8.624$, $p=0.0054$, with no sex x virus interaction

$F_{1, 41}=2.201$, $p=0.1456$), whereas the inhibition of pIC excitatory neurons reduced ethanol+quinine

intake exclusively in female mice (**Figure 4G**, 2-way ANOVA, sex factor $F_{1, 26}=42.01$, $p<0.0001$,

virus factor $F_{1, 26}=14.04$, $p=0.0009$, with a sex x virus interaction $F_{1, 26}=7.055$, $p=0.0133$). To

control whether aIC and pIC glutamatergic neurons inhibition was specific to ethanol persistent

drinking, we tested the effect of the same chemogenetic manipulation on water and water+quinine

intake. The inhibition of aIC or pIC glutamatergic neurons did not change water consumption nor

in males neither in females (**Figure 4C, H**). In contrast, inhibition of aIC neurons decreased

water+quinine intake independently of the sex (**Figure 4D**, 2-way ANOVA, sex factor $F_{1, 19}=1.481$,

$p=0.2386$, virus factor $F_{1, 19}=10.80$, $p=0.0039$, with no sex x virus interaction $F_{1, 19}=1.296$,

$p=0.2691$), while, we observed a strong trend of pIC inhibition to reduce water+quinine intake

mainly driven by the male group (**Figure 4L**, 2-way ANOVA, sex factor $F_{1, 19}=2.677$, $p=0.1183$,

virus factor $F_{1, 19}=4.230$, $p=0.0537$, with no sex x virus interaction $F_{1, 19}=2.999$, $p=0.0995$). Finally,

to rule out any behavioral confound due to CNO treatment, the locomotion was measured in

control mice that received CNO or vehicle injection, and in mice expressing the inhibitory viral

vector in aIC or pIC that received CNO injection. Locomotion was similar between the three

groups (**Figure 4E, J**). Finally, CNO injection in control animals did not alter ethanol,

ethanol+quinine and water intake compared to control animals injected with vehicle (**Figure S1G-**

L).

Together, our results demonstrate a divergent function of aIC and pIC glutamatergic neurons on

drinking behaviors with the aIC neurons supporting bitter solution drinking in both sexes and the

pIC neurons underpinning persistent ethanol drinking only in females.

Coding properties of pIC glutamatergic neurons during binge and persistent ethanol

drinking

Using fiber photometry with the genetically encoded calcium sensor GCaMP6f

(**Figure 5A, B, S2F**), we recorded pIC glutamatergic neurons activity during licking behaviors

over 1 hour (**Figure 5C**, **S2**). Behaviorally, the average of ethanol intake across all drinking

sessions, excluding the recording session, was similar between males and females (**Figure 5D**),

whereas the average of ethanol+quinine intake was higher in females (**Figure 5E,** unpaired

Student t-test, t=4.023, p=0.0024). Both sexes did the same average number of bouts for ethanol,

ethanol+quinine, water and water+quinine during the recording sessions (**Figure S2B-E**). At the

neural activity level, calcium signal in pIC neurons was increased during ethanol licking compared

to the reference independently of the sex (**Figure 5F, G**, 2-way RM-ANOVA, sex factor

$F_{1, 10}$=1.593, p=0.2355, period factor $F_{1, 10}$=7.708, p=0.0196, with no sex x period interaction

$F_{1, 10}$=0.3726, p=0.5552). Furthermore, in females, ethanol+quinine licking increased the calcium

signal in pIC neurons compared to the reference. Moreover, pIC neurons activity during

ethanol+quinine licking was higher in female compared to male mice (**Figure 5H, I**, 2-way RM-

ANOVA, sex factor $F_{1, 10}$=8.368, p=0.0160, period factor $F_{1, 10}$=12.98, p=0.0048, with a

sex x period interaction $F_{1, 10}$=5.026, p=0.0489). To control the specificity of the increase of pIC

neuronal activity in female mice in response to persistent ethanol drinking, we performed fiber

photometry recording during water and water+quinine licking. Independently of the sex, the

calcium signal in pIC neurons increased during water licking compared to the reference (**Figure**

5J, K, 2-way RM-ANOVA, sex factor $F_{1, 12}$=1.081, p=0.3189, period factor $F_{1, 12}$=12.51, p=0.0041,

without interaction $F_{1, 12}$=3.316, p=0.0936). Similarly, independently of the sex, pIC neurons

exhibited an increase of calcium signal during water+quinine licking compared to the reference

(**Figure 5L, M,** 2-way RM-ANOVA, sex factor $F_{1, 15}$=1.708, p=0.2109, period factor $F_{1, 15}$=8.119,

p=0.0122, without interaction $F_{1, 15}$=0.4080, p=0.5326)**.**

DISCUSSION

The goal of our study was to elucidate whether the aIC and pIC have sexually dimorphic function

on ethanol binge drinking and persistent ethanol drinking despite bitter aversion in mice. First, we

observed higher binge and persistent ethanol intake in females compared to males (**Figure 3C-**

F). Second, we demonstrated that, independently of sex, chemogenetic inhibition of aIC

glutamatergic neurons reduced the intake of quinine adulterated solutions (e.g. ethanol and water,

Figure 4B, D). In contrast, chemogenetic inhibition of pIC glutamatergic neurons reduced

exclusively persistent ethanol drinking in females but not males (**Figure 4G**). Finally, pIC neuronal

activity increases during ethanol, water and water+quinine drinking, independently of sex,

(**Figure 5F, G, J-M**). However, pIC neuronal activity responses to persistent ethanol intake,

specifically in females but not in males (**Figure 5H, I**). All together, these findings suggest a causal

role of aIC glutamatergic neurons in bitter liquid drinking, and a sex-dependent function of pIC

excitatory neurons in persistent ethanol drinking.

Sex-dependent behavior on binge and persistent ethanol drinking

Our results repeatedly showed a higher binge and persistent ethanol intake per body mass in

females compared to males, independently of the estrous cycle, which is consistent and extend

previous literature highlighting sex differences in drug addiction (58,59). Indeed, multiple

preclinical studies also reported a higher binge (12,26,56) and persistent (12,27,60) ethanol intake

per body mass in females, than in males. Importantly, to rule out any confound on sex-dependent

aversion resistance for quinine, we adulterated the ethanol solution with 500 µM quinine, a

concentration for which both male and female mice have similar aversive threshold (61).

Therefore, the sex differences observed in the consumption of ethanol adulterated with quinine is

independent of the aversion sensitivity but driven toward the ethanol intake.

Bitter taste processing in aIC excitatory neurons

We demonstrated that inhibition of aIC glutamatergic neurons reduced the intake of solutions

adulterated with quinine (e.g. ethanol or water) but did not alter binge ethanol intake. Our results

support and extend Haaranen and al. study, which demonstrated that chemogenetic inhibition of

aIC neurons had no effect on alcohol intake in alcohol-preferring male rats in a two-bottle choice

model (53). However, it contrasts with previous literature demonstrating that chemogenetic

manipulation of aIC neurons modulates alcohol binge drinking, although the results are divergent.

Indeed, while silencing aIC neurons increased operant responses for alcohol (51), it decreased

alcohol intake in a two-bottle choice paradigm (62). Altogether, these results suggest that aIC

function in ethanol binge drinking might be dependent on the animal model used. Moreover, while

we specifically manipulated aIC glutamatergic neurons, the studies cited previously manipulated

all aIC mature neurons independently of their phenotype which could explain some discrepancy

in the results observed.

On the other hand, we demonstrated that aIC excitatory neurons play a role in bitter solution

intake as shown by the reduction of ethanol and water adulterated with quinine consumption

following aIC glutamatergic inhibition. These results extend previous findings demonstrating a

reduction of ethanol+quinine intake in male mice compared to controls after inhibition of aIC

neurons, as well as its projections to locus coeruleus or nucleus accumbens core (62,63). In

addition, Chen and Lasek reported an increase expression of the neuronal activity marker *cFos*

in the aIC after ethanol+quinine intake in male mice compared to water, ethanol and quinine

drinking (64). Overall, these observations confirm the implication of aIC in quinine-adulterated

ethanol drinking in males, and our study extends this finding to female mice. In regard to the role

of aIC in water+quinine intake, the literature is more divergent. On one hand, inhibition or lesion

of the aIC respectively had no effect (65,66) or decreased (67) sensitivity to quinine, leading to

an increase of the intake. On the other hand, activation of aIC neurons and its projection to

basolateral amygdala also decreased quinine sensitivity (41,68). These findings contrast with our

results where aIC excitatory inhibition reduced quinine intake. In our study, we chose not to water

deprive our animals in order to observe behavioral responses as close as possible to physiological

conditions. In contrast, in previously cited studies, animals were water deprived before quinine

exposure which could have induced a physiological stress altering aIC response to quinine.

Overall, these results suggest a potential role of aIC in bitter solution intake, however further

investigations are necessary to sharpen our understanding of aIC neurons and its projections in

aversive solutions consumption in males and females.

Uncovering sex-dependent pIC function in persistent ethanol intake

We demonstrated that inhibition of pIC glutamatergic neurons reduces persistent ethanol drinking

only in females without altering ethanol, water, or water adulterated with quinine drinking. To our

knowledge, these results are the first demonstration of a sex-dependent function of pIC neurons

on persistent ethanol drinking. Using the two-bottle choice paradigm, a recent study showed a

higher expression of the neuronal activity marker *cFos* in the pIC of males compared to females

after ethanol+quinine intake (69). Importantly, in this study ethanol was adulterated with 100 µM

of quinine. Previous reports suggests that females are not sensitive to ethanol adulterated with

quinine at this concentration whereas males are (27) which could explain the discrepancy with

our results where we used a concentration of quinine at 500 µM. Furthermore, using an operant

alcohol drinking model in male rats, pharmacological inactivation of pIC neurons reduces alcohol

intake (54) while in our study, chemogenetic inhibition of pIC excitatory neurons did not change

ethanol drinking nor in males neither in females. Several methodological aspects can, at least in

part, support this discrepancy. First, operant and voluntary alcohol drinking could recruit different

neuronal circuits, second, the methods used for the neuronal inhibition (e.g. pharmacology vs

chemogenetics) and the animal model used (e.g. rat vs mouse) were different.

To rule out the possibility that pIC glutamatergic inhibition was due to non-selective effects of

CNO, we performed multiple control experiments. Indeed, it has been shown that CNO can bind

to non-DREADD targets, and does not cross the blood brain barrier. However, clozapine, a CNO

metabolite, is reaching the brain and can also have multiple pharmacological targets which can

induce several physiological and behavioral effects (70,71). In our control experiments, we found

that CNO injection in mice expressing the control viral vector in pIC, had no effects on either liquid

consumption (e.g. ethanol, ethanol+quinine or water) or locomotion. Furthermore, CNO injection

in mice injecting with the viral vector carrying the gene coding for hM4Di also had no effect on

locomotion. These results indicate first, that CNO does not have non-specific effects neither on

drinking nor on locomotion and secondly, that the inhibition of pIC excitatory neurons did not affect

locomotion. Overall, these results confirm that the behavioral effect of CNO on ethanol+quinine

drinking is due to selective inhibition of pIC excitatory neurons.

Functional sex differences in pIC coding properties of persistent ethanol drinking

The chemogenetic manipulations demonstrated the necessity of pIC excitatory neurons to change

persistent ethanol drinking in females, but these artificial manipulations do not reflect how pIC

excitatory neurons encode this behavior. Thus, we performed *in vivo* fiber photometry recordings

of pIC calcium signal and identified an increased activity of pIC glutamatergic neurons during

ethanol licking, as well as water and water+quinine licking, independently of the sex. These results

consolidate recent findings demonstrating that pIC excitatory neurons respond to water intake

(72) which suggests that these neurons encode drinking behaviors in both males and females.

Furthermore, pIC neuronal activity increased during water+quinine drinking, which is consistent

with previous studies reporting that pIC neurons encode bitter taste, independently of the sex

(43,73). In contrast, we decipher a sex-dependent pIC neuronal coding property in persistent

ethanol drinking with an increase of pIC excitatory neurons activity exclusively in females during ethanol adulterated with quinine drinking. Our results strengthen a recent study demonstrating that short ethanol exposure increases intrinsic excitability of pIC-BNST projection in female mice only (56) highlighting pIC glutamatergic neurons and their projections as one of the key regions supporting sex-dependent functions in alcohol-related behaviors. One limitation of our study is the manipulation and the recording of pIC excitatory neurons regardless of their outputs. The pIC is a highly wired region sending dense projections to several regions (43,74), including the central amygdala, a brain region highly involved in alcohol-related behaviors and supporting sex-differences in alcohol intake (75–77). Therefore, future investigations will be crucial to understand the sex-dependent function of pIC neurons in alcohol-related behaviors at the circuit level.

Conclusions

To conclude, our study demonstrated that aIC and pIC have different roles in drinking behaviors. While aIC excitatory neuron inhibition decreases bitter-tastant solutions intake independently of mice sex, pIC glutamatergic neurons inhibition exclusively reduces persistent ethanol intake despite aversion in female mice. In line with this result, pIC glutamatergic neurons activity increased specifically in females during ethanol consumption despite bitter aversion. Our results provide a starting point to further characterize the role of pIC circuits in sex-dependent addictive-related behaviors.

ACKNOWLEDGEMENTS

The authors acknowledge the entire Beyeler Lab. We thank Harold Haun and Irene Lorrai for their helpful feedback to set up the drinking in the dark model. We thank Guillaume Ferreira for interesting discussions. We thank Sara Laumond, Julie Tessaire and the technical staff of the animal housing facility of the Neurocentre Magendie (INSERM U1215) for their invaluable support.

FUNDING

We acknowledge the support of the Région Nouvelle-Aquitaine, the 'Avenir program' of the French institute of health (INSERM), the Fondation NRJ-Institut de France, the 'Agence nationale pour la recherche' (ANR), and of the 'Fondation Schlumberger pour l'éducation et la recherche' (FSER) to AB. We are grateful for the support of the Fondation pour la recherche médicale (FRM) to AB and CN (ARF201909009147) as well as the 'ATIP program' (CNRS) to CN.

AUTHORSHIP CONTRIBUTIONS

CF, CN and AB designed the study. CF, CGM, YC, CN and AB carried out the experiments and performed data analysis. PN wrote the custom Python code for fiber photometry analysis. CF, CN and AB wrote the manuscript. All authors critically reviewed the content and approved the final version before submission.

DISCLOSURE

The authors declare that they do not have any competing interests or conflicts of interest (financial or non-financial) related to the material presented in this manuscript.

REFERENCES

1. Crocq M-A, Guelfi J-D (2015): *DSM-5: manuel diagnostique et statistique des troubles mentaux*, 5e éd. Issy-les-Moulineaux: Elsevier Masson.

2. Moos RH, Moos BS (2007): Treated and Untreated Alcohol-Use Disorders: Course and Predictors of Remission and Relapse. *Eval Rev* 31: 564–584.

3. Koob GF, Volkow ND (2010): Neurocircuitry of Addiction. *Neuropsychopharmacology* 35: 217–238.

4. Organisation mondiale de la santé (Ed.) (2018): *Global Status Report on Alcohol and Health 2018*. Geneva: World health organization.

5. Drinking Levels Defined | National Institute on Alcohol Abuse and Alcoholism (NIAAA) (n.d.):
Retrieved September 21, 2023, from https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-
consumption/moderate-binge-drinking

6. Gowin JL, Sloan ME, Stangl BL, Vatsalya V, Ramchandani VA (2017): Vulnerability for
Alcohol Use Disorder and Rate of Alcohol Consumption. *Am J Psychiatry* 174: 1094–1101.

7. Tavolacci M-P, Berthon Q, Cerasuolo D, Dechelotte P, Ladner J, Baguet A (2019): Does
binge drinking between the age of 18 and 25 years predict alcohol dependence in adulthood? A
retrospective case–control study in France. *BMJ Open* 9: e026375.

8. Rhodes WA, Singleton E, McMillan TB, Perrino CS (2005): Does knowledge of college
drinking policy influence student binge drinking? *J Am Coll Health J ACH* 54: 45–49.

9. Thiele TE, Navarro M (2014): "Drinking in the dark" (DID) procedures: a model of binge-like
ethanol drinking in non-dependent mice. *Alcohol Fayettev N* 48: 235–241.

10. Wilcox MV, Cuzon Carlson VC, Sherazee N, Sprow GM, Bock R, Thiele TE, *et al.* (2014):
Repeated binge-like ethanol drinking alters ethanol drinking patterns and depresses striatal
GABAergic transmission. *Neuropsychopharmacol Off Publ Am Coll Neuropsychopharmacol* 39:
579–594.

11. Siciliano CA, Noamany H, Chang C-J, Brown AR, Chen X, Leible D, *et al.* (2019): A cortical-
brainstem circuit predicts and governs compulsive alcohol drinking. *Science* 366: 1008–1012.

12. Sneddon EA, White RD, Radke AK (2019): Sex Differences in Binge-Like and Aversion-
Resistant Alcohol Drinking in C57BL/6J Mice. *Alcohol Clin Exp Res* 43: 243–249.

13. Bauer MR, McVey MM, Boehm SL (2021): Three Weeks of Binge Alcohol Drinking
Generates Increased Alcohol Front-Loading and Robust Compulsive-Like Alcohol Drinking in
Male and Female C57BL/6J Mice. *Alcohol Clin Exp Res* 45: 650–660.

14. Schuh KM, Sneddon EA, Nader AM, Muench MA, Radke AK (2022): Orbitofrontal cortex
subregion inhibition during binge-like and aversion-resistant alcohol drinking. *Alcohol* 99: 1–8.

15. Slade T, Chapman C, Swift W, Keyes K, Tonks Z, Teesson M (2016): Birth cohort trends in
the global epidemiology of alcohol use and alcohol-related harms in men and women:
systematic review and metaregression. *BMJ Open* 6: e011827.

16. Grant BF, Chou SP, Saha TD, Pickering RP, Kerridge BT, Ruan WJ, *et al.* (2017):
Prevalence of 12-Month Alcohol Use, High-Risk Drinking, and DSM-IV Alcohol Use Disorder in
the United States, 2001-2002 to 2012-2013: Results From the National Epidemiologic Survey
on Alcohol and Related Conditions. *JAMA Psychiatry* 74: 911–923.

17. Ehlers CL, Gizer IR, Vieten C, Gilder A, Gilder DA, Stouffer GM, *et al.* (2010): Age at regular
drinking, clinical course, and heritability of alcohol dependence in the San Francisco Family
Study: a gender analysis. *Am J Addict Am Acad Psychiatr Alcohol Addict* 19: 101–110.

18. Diehl A, Croissant B, Batra A, Mundle G, Nakovics H, Mann K (2007): Alcoholism in women:
is it different in onset and outcome compared to men? *Eur Arch Psychiatry Clin Neurosci* 257:
344–351.

19. Erol A, Karpyak VM (2015): Sex and gender-related differences in alcohol use and its
consequences: Contemporary knowledge and future research considerations. *Drug Alcohol*
Depend 156: 1–13.

20. Eagon PK (2010): Alcoholic liver injury: Influence of gender and hormones. *World J*
Gastroenterol WJG 16: 1377–1384.

21. Green CA, Perrin NA, Polen MR (2004): Gender differences in the relationships between
multiple measures of alcohol consumption and physical and mental health. *Alcohol Clin Exp*
Res 28: 754–764.

22. Strong MN, Yoneyama N, Fretwell AM, Snelling C, Tanchuck MA, Finn DA (2010): "Binge"
drinking experience in adolescent mice shows sex differences and elevated ethanol intake in
adulthood. *Horm Behav* 58: 82–90.

23. Loi B, Colombo G, Maccioni P, Carai MAM, Franconi F, Gessa GL (2014): High alcohol
intake in female Sardinian alcohol-preferring rats. *Alcohol Fayettev N* 48: 345–351.

24. Priddy BM, Carmack SA, Thomas LC, Vendruscolo JCM, Koob GF, Vendruscolo LF (2017):
Sex, strain, and estrous cycle influences on alcohol drinking in rats. *Pharmacol Biochem Behav*
152: 61–67.

25. Satta R, Hilderbrand ER, Lasek AW (2018): Ovarian Hormones Contribute to High Levels of
Binge-Like Drinking by Female Mice. *Alcohol Clin Exp Res* 42: 286–294.

26. Evans O, Rodríguez-Borillo O, Font L, Currie PJ, Pastor R (2020): Alcohol Binge Drinking
and Anxiety-Like Behavior in Socialized Versus Isolated C57BL/6J Mice. *Alcohol Clin Exp Res*
44: 244–254.

27. Fulenwider HD, Nennig SE, Price ME, Hafeez H, Schank JR (2019): Sex Differences in
Aversion-Resistant Ethanol Intake in Mice. *Alcohol Alcohol* 54: 345–352.

28. Chuong V, Farokhnia M, Khom S, Pince CL, Elvig SK, Vlkolinsky R, *et al.* (n.d.): The
glucagon-like peptide-1 (GLP-1) analogue semaglutide reduces alcohol drinking and modulates
central GABA neurotransmission. *JCI Insight* 8: e170671.

29. Gage GA, Muench MA, Jee C, Kearns DN, Chen H, Tunstall BJ (2023): Intermittent-access
operant alcohol self-administration promotes binge-like drinking and drinking despite negative
consequences in male and female heterogeneous stock rats. *Neuropharmacology* 235: 109564.

30. Schramm-Sapyta NL, Francis R, MacDonald A, Keistler C, O'Neill L, Kuhn CM (2014):
Effect of sex on ethanol consumption and conditioned taste aversion in adolescent and adult
rats. *Psychopharmacology (Berl)* 231: 1831–1839.

31. Cailhol S, Mormède P (2001): Sex and Strain Differences in Ethanol Drinking: Effects of
Gonadectomy. *Alcohol Clin Exp Res* 25: 594–599.

32. Venniro M, Caprioli D, Zhang M, Whitaker LR, Zhang S, Warren BL, *et al.* (2017): The
anterior insular cortex→central amygdala glutamatergic pathway is critical to relapse after
contingency management. *Neuron* 96: 414-427.e8.

33. Ghareh H, Alonso-Lozares I, Schetters D, Herman RJ, Heistek TS, Van Mourik Y, *et al.*
(n.d.): Role of anterior insula cortex in context-induced relapse of nicotine-seeking. *eLife* 11:
e75609.

34. Nicolas C, Tauber C, Lepelletier F-X, Chalon S, Belujon P, Galineau L, Solinas M (2017):
Longitudinal Changes in Brain Metabolic Activity after Withdrawal from Escalation of Cocaine
Self-Administration. *Neuropsychopharmacology* 42: 1981–1990.

35. Campbell EJ, Flanagan JPM, Walker LC, Hill MKRI, Marchant NJ, Lawrence AJ (2019):
Anterior Insular Cortex is Critical for the Propensity to Relapse Following Punishment-Imposed
Abstinence of Alcohol Seeking. *J Neurosci* 39: 1077–1087.

36. Campbell EJ, Lawrence AJ (2021): It's more than just interoception: The insular cortex
involvement in alcohol use disorder. *J Neurochem* 157: 1644–1651.

37. Pina MM, Pati D, Neira S, Taxier LR, Stanhope CM, Mahoney AA, *et al.* (2023): Insula
Dynorphin and Kappa Opioid Receptor Systems Regulate Alcohol Drinking in a Sex-Specific
Manner in Mice. *J Neurosci* 43: 5158–5171.

38. Myrick H, Anton RF, Li X, Henderson S, Drobes D, Voronin K, George MS (2004):
Differential Brain Activity in Alcoholics and Social Drinkers to Alcohol Cues: Relationship to
Craving. *Neuropsychopharmacology* 29: 393–402.

39. Ihssen N, Cox WM, Wiggett A, Fadardi JS, Linden DEJ (2011): Differentiating Heavy from
Light Drinkers by Neural Responses to Visual Alcohol Cues and Other Motivational Stimuli.
Cereb Cortex 21: 1408–1415.

40. Chung T, Clark DB (2014): Insula White Matter Volume Linked to Binge Drinking Frequency
Through Enhancement Motives in Treated Adolescents. *Alcohol Clin Exp Res* 38: 1932–1940.

41. Peng Y, Gillis-Smith S, Jin H, Tränkner D, Ryba NJP, Zuker CS (2015): Sweet and bitter
taste in the brain of awake behaving animals. *Nature* 527: 512–515.

42. Méndez-Ruette M, Linsambarth S, Moraga-Amaro R, Quintana-Donoso D, Méndez L,
Tamburini G, *et al.* (2019): The Role of the Rodent Insula in Anxiety. *Front Physiol* 10: 330.

43. Nicolas C, Ju A, Wu Y, Eldirdiri H, Delcasso S, Couderc Y, *et al.* (2023): Linking emotional
valence and anxiety in a mouse insula-amygdala circuit. *Nat Commun* 14: 5073.

44. Grodin EN, Cortes CR, Spagnolo PA, Momenan R (2017): Structural deficits in salience
network regions are associated with increased impulsivity and compulsivity in alcohol
dependence. *Drug Alcohol Depend* 179: 100–108.

45. Senatorov VV, Damadzic R, Mann CL, Schwandt ML, George DT, Hommer DW, *et al.*
(2015): Reduced anterior insula, enlarged amygdala in alcoholism and associated depleted von
Economo neurons. *Brain* 138: 69–79.

46. Makris N, Oscar-Berman M, Jaffin SK, Hodge SM, Kennedy DN, Caviness VS, *et al.* (2008):
Decreased Volume of the Brain Reward System in Alcoholism. *Biol Psychiatry* 64: 192–202.

47. Thompson BL, Maleki N, Kelly JF, Sy KTL, Oscar-Berman M (2021): Brain, behavioral,
affective, and sex correlates of recovery from alcohol use disorders. *Alcohol Clin Exp Res* 45:
1578–1595.

48. Halcomb ME, Chumin EJ, Goñi J, Dzemidzic M, Yoder KK (2019): Aberrations of anterior
insular cortex functional connectivity in nontreatment-seeking alcoholics. *Psychiatry Res*
Neuroimaging 284: 21–28.

49. Manuweera T, Kisner MA, Almira E, Momenan R (2022): Alcohol use disorder-associated
structural and functional characteristics of the insula. *J Neurosci Res* 100: 2077–2089.

50. Galandra C, Basso G, Manera M, Crespi C, Giorgi I, Vittadini G, *et al.* (2018): Salience
network structural integrity predicts executive impairment in alcohol use disorders. *Sci Rep* 8:
14481.

51. Jaramillo AA, Randall PA, Stewart S, Fortino B, Van Voorhies K, Besheer J (2018):
Functional role for cortical-striatal circuitry in modulating alcohol self-administration.
Neuropharmacology 130: 42–53.

52. Haaranen M, Schäfer A, Järvi V, Hyytiä P (2020): Chemogenetic Stimulation and Silencing
of the Insula, Amygdala, Nucleus Accumbens, and Their Connections Differentially Modulate
Alcohol Drinking in Rats. *Front Behav Neurosci* 14: 580849.

53. Haaranen M, Scuppa G, Tambalo S, Järvi V, Bertozzi SM, Armirotti A, *et al.* (2020): Anterior
insula stimulation suppresses appetitive behavior while inducing forebrain activation in alcohol-
preferring rats. *Transl Psychiatry* 10: 150.

54. Pushparaj A, Le Foll B (2015): Involvement of the caudal granular insular cortex in alcohol
self-administration in rats. *Behav Brain Res* 293: 203–207.

55. Haggerty DL, Munoz B, Pennington T, Viana Di Prisco G, Grecco GG, Atwood BK (2022):
The role of anterior insular cortex inputs to dorsolateral striatum in binge alcohol drinking. *eLife*
11: e77411.

56. Marino RAM, Girven KS, Figueiredo A, Navarrete J, Doty C, Sparta DR (2021): Binge
ethanol drinking associated with sex-dependent plasticity of neurons in the insula that project to
the bed nucleus of the stria terminalis. *Neuropharmacology* 196: 108695.

57. Zhang Z-M, Chen S, Liang Y-Z (2010): Baseline correction using adaptive iteratively
reweighted penalized least squares. *The Analyst* 135: 1138.

58. Nicolas C, Russell TI, Pierce AF, Maldera S, Holley A, You Z-B, *et al.* (2019): Incubation of
Cocaine Craving After Intermittent-Access Self-administration: Sex Differences and Estrous
Cycle. *Biol Psychiatry* 85: 915–924.

59. Nicolas C, Zlebnik NE, Farokhnia M, Leggio L, Ikemoto S, Shaham Y (2022): Sex
Differences in Opioid and Psychostimulant Craving and Relapse: A Critical Review ((M. Nader,
editor)). *Pharmacol Rev* 74: 119–140.

60. Martins de Carvalho L, Chen H, Sutter M, Lasek AW (2023): Sexually dimorphic role for
insular perineuronal nets in aversion-resistant alcohol consumption. *Front Psychiatry* 14:
1122423.

61. Sneddon EA, Ramsey OR, Thomas A, Radke AK (2020): Increased Responding for Alcohol
and Resistance to Aversion in Female Mice. *Alcohol Clin Exp Res* 44: 1400–1409.

62. De Oliveira Sergio T, Lei K, Kwok C, Ghotra S, Wegner SA, Walsh M, *et al.* (2021): The role
of anterior insula–brainstem projections and alpha-1 noradrenergic receptors for compulsion-like
and alcohol-only drinking. *Neuropsychopharmacology* 46: 1918–1926.

63. Seif T, Chang S-J, Simms JA, Gibb SL, Dadgar J, Chen BT, *et al.* (2013): Cortical activation
of accumbens hyperpolarization-active NMDARs mediates aversion-resistant alcohol intake. *Nat
Neurosci* 16: 1094–1100.

64. Chen H, Lasek AW (2020): Perineuronal nets in the insula regulate aversion-resistant
alcohol drinking. *Addict Biol* 25: e12821.

65. Wu Y, Chen C, Chen M, Qian K, Lv X, Wang H, *et al.* (2020): The anterior insular cortex
unilaterally controls feeding in response to aversive visceral stimuli in mice. *Nat Commun* 11:
640.

66. Hashimoto K, Spector AC (2014): Extensive Lesions in the Gustatory Cortex in the Rat Do
Not Disrupt the Retention of a Presurgically Conditioned Taste Aversion and Do Not Impair
Unconditioned Concentration-Dependent Licking of Sucrose and Quinine. *Chem Senses* 39:
57–71.

67. Bales MB, Schier LA, Blonde GD, Spector AC (2015): Extensive Gustatory Cortex Lesions
Significantly Impair Taste Sensitivity to KCl and Quinine but Not to Sucrose in Rats ((S. A.
Simon, editor)). *PLOS ONE* 10: e0143419.

68. Wang L, Gillis-Smith S, Peng Y, Zhang J, Chen X, Salzman CD, *et al.* (2018): The coding of
valence and identity in the mammalian taste system. *Nature* 558: 127–131.

69. Arnold ME, Butts AN, Erlenbach TR, Amico KN, Schank JR (2023): Sex differences in
neuronal activation during aversion-resistant alcohol consumption. *Alcohol Clin Exp Res* 47:
240–250.

70. Gomez JL, Bonaventura J, Lesniak W, Mathews WB, Sysa-Shah P, Rodriguez LA, *et al.*
(2017): Chemogenetics revealed: DREADD occupancy and activation via converted clozapine.
Science 357: 503–507.

71. Manvich DF, Webster KA, Foster SL, Farrell MS, Ritchie JC, Porter JH, Weinshenker D
(2018): The DREADD agonist clozapine N-oxide (CNO) is reverse-metabolized to clozapine and
produces clozapine-like interoceptive stimulus effects in rats and mice. *Sci Rep* 8: 3840.

72. Zhao Z, Soria-Gómez E, Varilh M, Covelo A, Julio-Kalajzić F, Cannich A, *et al.* (2020): A
Novel Cortical Mechanism for Top-Down Control of Water Intake. *Curr Biol* 30: 4789-4798.e4.

73. Chen X, Gabitto M, Peng Y, Ryba NJP, Zuker CS (2011): A Gustotopic Map of Taste
Qualities in the Mammalian Brain. *Science* 333: 1262–1266.

74. Gogolla N (2017): The insular cortex. *Curr Biol CB* 27: R580–R586.

75. Tunstall BJ, Kirson D, Zallar LJ, McConnell SA, Vendruscolo JCM, Ho CP, *et al.* (2019):
Oxytocin blocks enhanced motivation for alcohol in alcohol dependence and blocks alcohol
effects on GABAergic transmission in the central amygdala. *PLoS Biol* 17: e2006421.

76. de Guglielmo G, Crawford E, Kim S, Vendruscolo LF, Hope BT, Brennan M, *et al.* (2016):
Recruitment of a Neuronal Ensemble in the Central Nucleus of the Amygdala Is Required for
Alcohol Dependence. *J Neurosci* 36: 9446–9453.

77. Agoglia AE, Zhu M, Quadir SG, Bluitt MN, Douglass E, Hanback T, *et al.* (2022): Sex-
specific plasticity in CRF regulation of inhibitory control in central amygdala CRF1 neurons after
chronic voluntary alcohol drinking. *Addict Biol* 27: e13067.

Figure 1. Validation of the drinking in the dark model. **(A)** Experimental timeline of binge ethanol exposure according to the drinking in the dark procedure. Blood was sampled after 2 hours of session at day 18 and after 30 minutes of session at day 22. **(B, C)** Average of ethanol intake (**B**) and blood ethanol concentration (**C**) after a 2 hours of ethanol binge drinking (day 18) in male and female mice. The red dashed line represents the intoxication threshold of 80 mg/dL of ethanol in blood. **(D)** Proportion of ethanol intake during the first 30 minutes and last 90 minutes of a 2-hour session. **(E, F)** Average of ethanol intake (**E**) and blood ethanol concentration (**F**) after 30 minutes of session (day 22) in male and female mice. Data are shown as mean ± SEM. *p<0.05 ***p<0.001.

Figure 2. Experimental approach for chemogenetic manipulation of anterior (aIC) and posterior

(pIC) insular cortex neurons during drinking in the dark procedure. **(A)** Experimental design to

inhibit aIC and pIC excitatory neurons. A viral vector carrying the gene coding for the inhibitory

receptor hM4Di fused to the fluorescent reporter mCherry, or the control virus, containing only the

gene for mCherry, was bilaterally injected in the aIC or pIC. **(B)** Behavioral timeline during the

drinking in the dark procedure. **(C)** Representative images of the chemogenetic viral vector

expression in the aIC (left) and the pIC (right) neurons. The mouse brain atlas delineation has

been overlaid to the image, with the borders of the aIC and pIC in bold.

Figure 3. Sexual dimorphism of binge and persistent ethanol drinking. **(A, B)** Kinetic (**A**) and average (**B**) of water intake during cycle 0 in male and female mice. **(C, D)** Kinetic (**C**) and average (**D**) of ethanol intake (binge drinking) during cycle 1 and 2 in male and female mice. **(E, F)** Kinetic (**E**) and average (**F**) of ethanol+quinine intake (persistent drinking) during cycle 3 in male and female mice. **(G)** Kinetic of water+quinine (water+Q) and ethanol+quinine (EtOH+Q) intake in males and females. # p<0.05 compared to day 39. **(H)** Comparison of water and ethanol adulterated with quinine intake during the last session exposure in male and female mice. Data are shown as mean ± SEM. ***p<0.001 represent significant t-test and main effect of 2-way ANOVA. ##p<0.01 ###p<0.001 represent significant Bonferroni post-hoc test.

Figure 4. Chemogenetic inhibition of anterior (aIC) or posterior (pIC) insular cortex excitatory neurons during binge and persistent ethanol intake. **(A)** Average of ethanol intake during chemogenetic manipulation of the aIC glutamateric neurons at day 18 in males and females. **(B)** Average of ethanol+quinine intake during chemogenetic manipulation of the aIC at day 25, in males and females. **(C)** Average of water intake during chemogenetic manipulation of the aIC at day 29, in males and females. **(D)** Average of water+quinine intake during chemogenetic manipulation of the aIC at day 29, in male and female mice. **(E)** Average of total distance traveled in an open field over 15 minutes. **(F)** Average of ethanol intake during chemogenetic manipulation of the pIC neurons at day 18, in males and females. **(G)** Average of ethanol+quinine intake during chemogenetic manipulation of the pIC neurons at day 25, in males and females. **(H)** Average of water intake during chemogenetic manipulation of the pIC neurons at day 29, in males and females. **(I)** Average of water+quinine intake during chemogenetic manipulation of the pIC neurons day 29, in an independent group of male and female mice. **(J)** Average of total distance traveled in an open field over 15 minutes. Data are shown as mean ± SEM. **p<0.01 ***p<0.001

Figure 5. Coding properties of posterior insular cortex (pIC) excitatory neurons during binge and

persistent ethanol drinking. **(A)** Viral strategy to record calcium changes in pIC excitatory neurons.

A viral vector carrying the gene coding for the calcium sensor GCaMP6f was injected unilaterally

in the pIC. **(B)** Representative image of GCaMP6f expression in pIC excitatory neurons. The

mouse brain atlas delineation has been overlaid to the image, with the borders of the pIC in bold.

(C) Housing conditions during photometry recordings to synchronize calcium signal and licking

behavior. **(D)** Average of ethanol intake during all binge-drinking sessions except the recording

day (day 32) in male and female mice. **(E)** Average of ethanol+quinine intake during all persistent

drinking sessions except the recording day (day 46) in male and female mice. **(F)** Peri-ethanol

licking time course of the calcium signal in pIC glutamatergic neurons in males and females. **(G)**

Average of calcium signal in pIC glutamatergic neurons during reference and ethanol licking, in

both sexes. **(H)** Peri-ethanol+quinine licking time course of the calcium signal in pIC glutamatergic

neurons in male and female mice. **(I)** Average calcium signal in pIC glutamatergic neurons during

reference and ethanol+quinine licking, in both sexes. **(J)** Peri-water licking time course of the

calcium signal recorded from pIC glutamatergic neurons in males and females. **(K)** Average of

calcium signal from pIC glutamatergic neurons during reference and water licking in both sexes.

(L) Peri-water+quinine licking time course of the calcium signal from pIC glutamatergic neurons,

in male and female mice. **(M)** Average of calcium signal from pIC glutamatergic neurons during

reference and water+quinine licking, in both sexes. Data are shown as mean $\pm$ SEM. *$p < 0.05$

**$p < 0.01$ represent significant t-test and main effect of 2-way ANOVA. ##$p < 0.01$ represents

significant Bonferroni post-hoc test.

A experimental timeline

binge drinking

2 hr 30 min
days 8 ↓ ↓ 22

☐ 2h EtOH (20%) ↓ blood sampling

housing conditions

ethanol 20% ♂ or ♀

B after 2 hours of ethanol intake

EtOH intake (g/kg)

***

13 13

♂ ♀

C BEC (mg/dL)

23% 15%

13 13

♂ ♀

D proportion of ethanol intake after 2 hours

♂ (n=16)

last 90 min 33%

first 30 min 67%

♀ (n=16)

last 90 min 16%

first 30 min 84%

E after 30 minutes of ethanol

EtOH intake (g/kg)

***

16 16

♂ ♀

F BEC (mg/dL)

44% 75%

16 16

♂ ♀

Sexual dimorphism of binge and persistent ethanol intake

Chemogenetic inhibition of excitatory neurons in anterior insula (aIC)

A. binge drinking — EtOH intake (g/kg); control, hM4Di; ♂ 14, 14; ♀ 12, 11

B. persistent drinking — EtOH+Q intake (g/kg); control, hM4Di; *** ; ♂ 14, 14; ♀ 12, 10

C. water — Water intake (mL/kg); control, hM4Di; ♂ 14, 14; ♀ 12, 10

D. water+quinine — Water+Q intake (mL/kg); control, hM4Di; ** ; ♂ 5, 5; ♀ 7, 7

E. locomotion control — Total distance in OFT (m); control vehicle, control CNO, hM4Di CNO; ♂ 6, 7; ♀ 6, 6, 10

Chemogenetic inhibition of excitatory neurons in posterior insula (pIC)

F. binge drinking — EtOH intake (g/kg); control, hM4Di; ♂ 7; ♀ 7, 8

G. persistent drinking — EtOH+Q intake (g/kg); control, hM4Di; ** ; # ; ♂ 7; ♀ 7, 8

H. water — Water intake (mL/kg); control, hM4Di; ♂ 7; ♀ 6, 8

I. water+quinine — Water+Q intake (mL/kg); control, hM4Di; 0.053 ; ♂ 6; ♀ 6, 6

J. locomotion control — Total distance in OFT (m); control vehicle, control CNO, hM4Di CNO; ♂ 4, 3; ♀ 5, 5, 6

A Stereotaxic surgeries

AAV9-CaMKII-GCaMP6f

optic fiber

posterior insula

GCaMP Calm
M13

B Viral expression

Bregma -0.47 mm   GCaMP6f DAPI

600 μm   50 μm

C Liquid consumption

recording conditions

Calcium signal

Licks z-score

10 s

♂ or ♀

D
EtOH intake (g/kg)
7   5
♂   ♀

E
EtOH+Q intake (g/kg)
**
7   5
♂   ♀

Calcium signal recordings of posterior insula glutamatergic neurons during drinking

binge drinking

F
Calcium signal (z-score)
reference window
Licking onset
Time (s)
reference window   licking window   male n=7   female n=5

G
Calcium signal (z-score)
7   5
♂♀ reference
♂♀ licking *

persistent drinking

H
Calcium signal (z-score)
reference window
Licking onset
Time (s)
reference window   licking window   male n=7   female n=5

I
Calcium signal (z-score)
##
##
7   5
♂♀ reference
♂♀ licking *

water drinking

J
Calcium signal (z-score)
reference window
Licking onset
Time (s)
reference window   licking window   male n=9   female n=5

K
Calcium signal (z-score)
9   5
♂♀ reference **
♂♀ licking **

water+quinine drinking

L
Calcium signal (z-score)
reference window
Licking onset
Time (s)
reference window   licking window   male n=10   female n=7

M
Calcium signal (z-score)
10   7
♂♀ reference
♂♀ licking *