Sex differences in insular cortex function in persistent alcohol drinking despite aversion in mice

Claudia Fornari[1][2], Daria Ricci[1], Yoni Couderc[1], Carmen Guerrero-Márquez[1], Praneeth Namburi[3], Camille Penet[1], Céline Nicolas[1][2]* and Anna Beyeler[1]*

Affiliations:
[1] University of Bordeaux, INSERM 1215, Neurocentre Magendie, 146 rue Léo Saignat, 33000, Bordeaux, France.
[2] University of Bordeaux, CNRS UMR 5287, Institut de Neurosciences Cognitives et Intégratives d'Aquitaine, 2 rue Dr Hoffmann Martinot, 33000, Bordeaux, France.
[3] Institute for Medical Engineering & Science, Massachusetts Institute of Technology, 77 Massachusetts Ave., Cambridge, MA, 02139, USA.
*The authors scientifically contributed equally to this work
Send correspondence to anna.beyeler@inserm.fr +33 557 574 072 or celine.nicolas@u-bordeaux.fr +33 661787799

Running Title: Sex differences in persistent alcohol drinking
Keywords: sex differences, persistent drinking, insular cortex, binge-drinking, chemogenetic, calcium fiber photometry

ABSTRACT

One major hallmark of alcohol use disorder (AUD) is the persistence of drinking despite negative consequences. Among indicators of AUD vulnerability, binge drinking has been identified as one of the strongest risk factors. Although the lifetime prevalence of both binge drinking and AUD has historically been higher in men than women, this gap has dramatically narrowed in the last decade. Additionally, sex differences in AUD and binge drinking have been found in clinical and preclinical studies. At the neurobiological level, the insular cortex plays an important role in AUD, with the anterior (aIC) and posterior (pIC) divisions supporting different functions. However, the contributions of the aIC and pIC sections in sexual dimorphism of alcohol binge drinking and the persistence of alcohol drinking despite aversion remain to be uncovered. Using the drinking in the dark model in mice, we validated that female mice have a higher binge ethanol intake compared to males. To evaluate persistent ethanol consumption despite aversion, we supplemented ethanol with the bitter compound quinine, and found a higher persistent drinking in females compared to males. Using fiber photometry recordings, we revealed that aIC activity was increased during binge and persistent ethanol consumption independently of sex, whereas pIC glutamatergic neuron activity was higher during persistent ethanol drinking, specifically in female mice. Using chemogenetics, we revealed that inhibition of aIC glutamatergic neurons reduced intake of bitter solutions independently of the solvent (ethanol or water) in both sexes. In addition, inhibition of pIC glutamatergic neurons exclusively reduced persistent ethanol drinking in females, while decreasing quinine consumption only in males. These findings suggest a sex-dependent function of the pIC in the persistence of ethanol consumption, providing a starting point in understanding sex-specific functions of the insular cortex in the neurobiology of AUD.

INTRODUCTION

Alcohol use disorder (AUD) is a chronic relapsing disorder defined by the persistence of excessive alcohol consumption despite physical and psychological negative consequences (1), which is a considerable impediment to AUD treatment (2,3). AUD is a major public health burden, as 6% of the world population is affected by alcohol morbidity or mortality (4). Among the indicators of vulnerability to AUD, binge drinking has been identified as a strong risk factor (3,5–7). Binge drinking is a recreational pattern of alcohol use characterized by episodic and excessive consumption that raises blood alcohol concentration to 80 mg/dL or higher, typically within about two hours. A longitudinal study showed that 43% of adolescents (13 to 18 years old) who binge-drink alcohol developed AUD at the age of 21, whereas only 7% of non-bingers developed AUD at the same age (8). In preclinical studies, binge drinking is modeled using the drinking in the dark (DID) procedure, where mice have repeated access to alcohol for a limited time during the dark phase of the circadian rhythm (active period), which allows to reach blood ethanol concentration (BEC) of 80 mg/dL as observed in humans (9–11). One core symptom of drug addiction is the persistence of drug use despite negative consequences. This specific aspect can be modeled in animals with different approaches. Among them, the DID procedure can be adapted to study the persistence of alcohol drinking despite aversive outcomes by adulterating ethanol with quinine (e.g., 100-500µM), a bitter and aversive substance for mice (12–15).

Historically, men have been reported as more likely than women to drink alcohol and exhibit pathological drinking behaviors. In 2020, AUD lifetime prevalence was still lower in women than men, reaching respectively 23% and 36% (16). However, a recent longitudinal study showed that the increase rate of AUD over a decade was drastically higher in women than men, with respective rates of 84% and 35% (17). These epidemiological data highlight that the sex difference in AUD prevalence is narrowing. Additionally, women transition faster to AUD after regular or chronic alcohol consumption (18,19) and are subsequently more likely to develop alcohol-related diseases (e.g. cardiovascular and hepatic diseases, 19–21). Interestingly, preclinical studies revealed sex differences in alcohol intake as well (23), with female rodents drinking more than males across different models of intake, including binge drinking and drinking despite taste aversion (13,24–31). This higher propensity to alcohol drinking in females is proposed to be, at least in part, due to their lowest sensitivity to aversive properties of alcohol, as female rodents are more resistant to ethanol-conditioned taste aversion (32–34).

The insular cortex (or insula) is strongly involved in drug addiction (35–37), including AUD (38–43). Indeed, human functional imaging studies in alcohol-dependent subjects have demonstrated that alcohol cues trigger greater activity responses in the insula compared to control neutral cues (44,45). Furthermore, insula white matter volume is correlated to binge drinking frequency in adolescents (46). Anatomically, the insula is subdivided into the anterior (aIC) and posterior (pIC) sections, which are proposed to have antagonistic functions in a wide range of behaviors (47–49). In AUD patients, structural studies report a reduction of the aIC volume and gray matter density compared to healthy controls, which is correlated with higher compulsive drinking measures (50–53). In addition, aIC functional connectivity with several brain regions (e.g., hippocampus, medial orbitofrontal regions) is higher in patients with AUD compared to social drinkers (54) and healthy controls (55). Intriguingly, less information is known about the role of pIC in AUD. Indeed, only two recent studies reported a decrease in pIC gray matter volumes in AUD patients (55,56) and a higher resting state activity of the pIC relative to healthy controls (55), suggesting the pIC also plays a role in AUD. Preclinical studies confirmed the contribution of both aIC and pIC in ethanol-drinking behaviors. Indeed, chemogenetic activation or inhibition of aIC neurons respectively decreased and increased alcohol intake in several behavioral paradigms such as self-administration and intermittent 2-bottle choice (41,57,58). In contrast, pharmacological inactivation of pIC neurons decreased alcohol self-administration (59), suggesting divergent functions of aIC

and pIC in alcohol drinking. However, how sex as a biological variable influences those divergent contributions remained untested.

Recent findings point towards structural sexual dimorphism of the insula, with a larger volume in men than women in physiological conditions (60), whereas in AUD long-term abstinent patients, this ratio is inverted in both right aIC and pIC (53). In support of these clinical findings, preclinical studies highlighted sex differences in insula functions in alcohol-drinking behaviors. For example, in alcohol binge-drinking mice, optogenetic stimulation of aIC terminals in the dorsolateral striatum induced larger excitatory post-synaptic currents and a smaller ratio of AMPA/NMDA currents recorded in medium spiny neurons compared to water-drinking mice (61). Interestingly, this synaptic plasticity was observed in male but not in female mice (61). Additionally, in mice, short-term ethanol exposure increases the excitability of pIC projection neurons targeting the bed nucleus of the stria terminalis neurons in females, but not males (62). Although the literature pinpoints the insular cortex as a key player in alcohol-related behaviors, sex differences in aIC and pIC function in binge and/or persistent alcohol intake despite quinine aversion remain to be uncovered.

Using the DID model, we repeatedly demonstrated a higher binge and aversion-resistant ethanol consumption in female compared to male mice. Combining the DID protocol with *in vivo* calcium fiber photometry recordings, we identified that aIC excitatory neurons are activated during the consumption of ethanol, ethanol adulterated with quinine, and water independently of sex. In the pIC, excitatory neurons are activated during binge ethanol drinking in both sexes, and during ethanol+quinine and water intake in females only. Importantly, sex differences during drinking behavior were identified only during ethanol+quinine consumption, with pIC neurons showing higher response in females than males. Then, using chemogenetics, we showed that inhibition of aIC glutamatergic neurons reduced bitter solution intake independently of solvent (e.g., ethanol or water) and sex. In contrast, inhibition of pIC glutamatergic neurons decreased the intake of quinine only in males and reduced persistent ethanol drinking exclusively in female mice.

Altogether, we identified a sex-dependent function of the pIC in persistent ethanol drinking, providing key elements to the characterization of insular cortex function in AUD in a sex-specific manner.

METHODS AND MATERIALS
For additional details on the methods, see the Supplementary information.

Stereotactic surgeries
Chemogenetic viral injection. The adeno-associated viruses encoding the inhibitory hM4Di receptor coupled to mCherry under the control of CaMKII promoter (AAV9-CaMKIIa-hM4D(Gi)-mCherry, titer >$1.0e^{+13}$ vg/mL, Addgene, Watertown, Massachusetts, USA), to target glutamatergic neurons, or the control viral vector (AAV9/2-mCaMKII-mCherry-WPRE, titer $6.3e^{+12}$ vg/mL, ETH Zürich, Switzerland) were injected bilaterally (200-250 nL, 1 nL/sec) in the aIC (anteroposterior +1.7 mm; mediolateral ±3.1 mm; dorsoventral -3.5 mm from the bregma) or the pIC (anteroposterior -0.35 mm; mediolateral ±4.0 mm; dorsoventral -4.2 mm from the bregma).

Calcium fiber photometry viral injection and fiber implantation. A viral vector coding for the calcium sensor GCaMP6f (AAV9-CaMKII-GCaMP6f-WPRE-SV40, titer $2.5e^{+12}$ vg/mL, Addgene) was injected unilaterally (250 nL, 1 nL/sec) in the right aIC or pIC. An optic fiber (400 μm diameter, 0.48 NA, >90% efficiency) inserted in a ceramic ferrule was implanted 50 μm above the virus injection site.

Drinking in the dark procedure (DID)
The DID protocol is based on the circadian rhythm where animals have access to different liquids (e.g., ethanol) during the nocturnal period (9,10). Two and a half hours after the onset of the dark

phase, the water bottle in the home cage was replaced by different liquids for 2 hours per day for 4 consecutive days, followed by 3 days with water access only. This 7-day cycle is repeated for 4 weeks, starting with tap water (cycle 0), then ethanol (20% v/v in tap water, cycles 1 and 2) to study binge drinking behavior, and ethanol adulterated with quinine (500 µM of quinine, cycle 3) to study persistent drinking despite aversion. Finally, water or water adulterated with quinine (500 µM quinine) consumption was measured during a single 2-hour session and served as a control. Additionally, a subgroup of animals underwent three more days of water+quinine, followed by a last cycle of ethanol+quinine to control the resumption of persistent ethanol drinking (**Fig. 2A**). Daily liquid consumption was obtained by weighing the bottle before and after the 2-hour session, and mice were weighed on the first day of each cycle. Food was available *ad libitum* during drinking sessions performed in the home cages, and until the beginning of the sessions in multifunctional boxes for fiber photometry.

Blood ethanol concentration measurement. To measure blood ethanol concentrations (BEC), blood sampling was performed by incision of the lateral tail vein 30 minutes or 2 hours after the onset of the ethanol session. The samples were immediately centrifuged at 10,000 rpm for 10 minutes at 4°C. The BEC in mg/dL was obtained from the serum using the Analox GL5 analyzer. This experiment has been performed twice, and consistent results were observed between experiments. We excluded 3 males and 3 females from the BEC analysis after a 2-hour ethanol intake because the blood sample did not contain enough serum.

Coding properties of pIC excitatory neurons during binge and persistent ethanol drinking
The same DID procedure as described before was used, except that no initial water cycle was performed. To study the coding properties of aIC and pIC neurons during binge and persistent ethanol drinking, the calcium signal and the drinking behavior (e.g., licking) have to be synchronized. Thus, after two ethanol binge cycles in their home cage, mice (n=30 males and n=31 females) underwent three cycles of ethanol drinking (cycles 3 to 5) in multifunctional boxes equipped with lickometers (**Fig. S2A**). Then, after one cycle of ethanol+quinine drinking in the home cages (cycle 6), mice underwent a cycle of ethanol+quinine drinking in the multifunctional boxes (cycle 7). Finally, a water and water+quinine photometry recording session was performed in these boxes as a control. The analysis of pIC neuronal activity during water and water+quinine consumption includes 3 males and 2 females, and 4 males and 2 females, respectively, who were ethanol-naive. The pIC recording experiment has been performed six times, aIC recording experiment has been repeated twice (**Table S2**). Replicated experiments showed consistent results.

Fiber photometry recordings. The recordings were performed with Neurophotometrics fiber photometry system (FP3002 V2, San Diego, California, USA), on the last day of the last ethanol and ethanol+quinine cycles, and during a single water and water+quinine session. The recordings were performed using a 4-branch patch cord (400 µm diameter, 0.48 NA, 1 branch per animal, Doric Lenses). Then, the drinking sprout was filled with the appropriate solution, and the recording lasted 1 hour as previously done (36). Mice were kept in the boxes for an additional hour for a total of a 2-hour session. The drinking sprout was automatically refilled to give *ad-libitum* access to the mice. The optimal wavelength for GCaMP6f excitation is 470 nm, while the isosbestic wavelength is 415 nm. To minimize the photobleaching effect of the recording, the light intensity at the tip of the patch cord was adjusted between 80 and 100 µW for both channels. The sampling rate was settled at 30 Hz for photometry recordings.

Calcium fiber photometry data analysis. Photometry recordings were analyzed using custom Python scripts https://github.com/praneethnamburi/Photometry_and_Licks_BeyelerLab. The timestamps for lick events were extracted and classified in bouts. To minimize potential confounds in the neuronal activity analyzed due to differences in the behavior across mice (e.g., number of licking bouts), the analysis of aIC and pIC neuronal activity was first performed on single licks and then on the combination of single licks and bouts (**Table S4, S5**), with a maximum of 5 licks/bouts

per animal for both analyses. A bout is defined as a minimum of two licks with an inter-lick interval (ILI) less than or equal to 10 seconds for the peri-event analysis. The first 10 minutes of the test mice did not have access to liquid solutions, thus the signals were discarded from the analysis. Then, GCaMP and isosbestic signals were de-trended using the airPLS algorithm (63), and the detrended isosbestic signal was regressed from the detrended GCaMP signal. The resulting signal was bandpass filtered with cutoff frequencies of 0.2 and 6 Hz using a finite impulse response filter implemented in the scipy package. Finally, a time window of -10 to +10 s was defined for the peri-event analysis of the resulting signal aligned to each lick (time 0 s). We controlled that no event happened during the -10 to 0 s. The signal from each window was z-normalized by subtracting the mean of the baseline and dividing the result by the standard deviation of the baseline, where the baseline signal was defined as the signal from -10 s to -8 s before a lick onset. Z-normalized calcium signals were averaged for each male and female mouse during the reference (-8 to -5 s), the pre-lick (-3 to 0 s), the post-lick (0 to +3 s) and the post-ingestive (+6 to +9 s) windows to describe the signal changes.

Code availability. Photometry recordings were analyzed using custom Python scripts, which can be downloaded at: https://github.com/praneethnamburi/Photometry_and_Licks_BeyelerLab, no restrictions apply.

Chemogenetic inhibition of aIC or pIC excitatory neurons during binge and persistent ethanol drinking

The same DID procedure as described above was used. The day before the chemogenetic manipulation an intraperitoneal (i.p.) injection of vehicle (NaCl, 0.1 mL/kg) was performed 30 minutes before the drinking session for habituation. The causal role of aIC or pIC glutamatergic neurons on drinking was tested after injection of CNO (i.p. 3 mg/kg in NaCl, Tocris, Bristol, United-Kingdom) in hM4Di mice 30 minutes before the last session of binge (day 18), persistent ethanol drinking (day 25) and a single session of water or water+quinine (day 29). To control for potential non-specific effects of CNO, mice expressing the control viral vector received either CNO or vehicle injections. As no behavioral differences were observed between the control-CNO and control-vehicle groups, these groups were pooled into a single control group. To note, some of the mice tested for chemogenetic inhibition of pIC on water+quinine drinking (day 29) were CNO naive. This experiment has been performed two times for the inhibition of pIC neurons and two times for the aIC inhibition (**Table S3**), with consistent results between experiment.

RESULTS

Behavioral characterization of binge ethanol consumption using the drinking in the dark protocol in male and female mice

After two 4-day cycles of ethanol consumption (**Fig. 1A**), the average 2-hour ethanol intake reported to the animal bodyweight, was higher in females than males (**Fig. 1B,** unpaired Student t-test, t=3.870, p=0.0007), whereas the blood ethanol concentration (BEC) was similar between sexes (**Fig. 1C**). After two hours of ethanol consumption, similar percentages of males (23%) and females (15%) reached the binge intoxication threshold of 80 mg/dL (Fischer's exact test, p>0.9999). As previous studies suggest that mice drink most of the ethanol at the beginning of the DID sessions (11), we measured the proportion of ethanol consumed during the first 30 minutes of the 2-hour session. The proportion of ethanol consumed during the first 30 minutes was higher than the proportion drunk during the remaining 90 minutes for both males (67% vs 33%) and females (84% vs 16%), and was higher in females than males (84% vs 67%), supporting a pattern of binge consumption in both sexes, even stronger in females (**Fig. 1D,** 2-way ANOVA, sex factor $F_{1, 60}=0.000$, p>0.999, ethanol proportion $F_{1, 60}=132.2$, p<0.0001, with an interaction $F_{1, 60}=14.16$, p=0.0004). Moreover, in an additional session, both ethanol intake and BEC were measured after

30 minutes of ethanol access, and were found to be higher in females compared to males (**Fig. 1E**, unpaired Student t-test, t=3.668, p=0.0009, and **Fig. 1F**, unpaired Mann-Whitney, U=71, p=0.0318 respectively). At this short time point, 75% of female and 37.5% of male mice reached the intoxication threshold of 80 mg/dL (Fisher's exact test, p=0.0732).

Fig. 1 Validation of the drinking in the dark model. A Experimental timeline of binge ethanol exposure according to the drinking in the dark procedure. Blood was sampled after 2 hours of session on day 18 and after 30 minutes of session on day 22. **B, C** Average of ethanol intake (B) and blood ethanol concentration (C) after 2 hours of ethanol binge drinking (day 18) in male (n=13) and female (n=13) mice. The red dashed line represents the intoxication threshold of 80 mg/dL. **D** Proportion of ethanol intake during the first 30 minutes and last 90 minutes of a 2-hour session in male (n=16) and female (n=16) mice. **E, F** Average of ethanol intake (E) and blood ethanol concentration (F) after 30 minutes of session (day 22) in male (n=16) and female (n=16) mice. Data are shown as mean ± SEM. *p<0.05 **p<0.01 ***p<0.001 represent significant Student t-test or Mann-Whitney test.

Sex differences in ethanol binge drinking during the drinking in the dark protocol

First, we analyzed drinking behavior across all sessions excluding the days of the chemogenetic test and independently of the viral vector injection site (e.g. aIC or pIC, **Fig. 2A**). Altogether, female mice had a higher level of water (**Fig. 2B,** 2-way RM-ANOVA, sex factor $F_{1, 109}$=7.208, p=0.0084, session factor $F_{3, 327}$=3.342, p=0.0195, without interaction $F_{3, 327}$=0.1697, p=0.9168), ethanol (**Fig. 2D,** 2-way RM-ANOVA, sex factor $F_{1, 109}$=116.7, p<0.0001, session factor $F_{5.331, 581.7}$=0.0027, p=0.0027, without interaction $F_{6, 654}$=0.9494, p=0.4589), and ethanol+quinine (**Fig. 2F**, 2-way RM-ANOVA, sex factor $F_{1, 108}$=103.4, p<0.0001, session factor $F_{2, 216}$=1.951, p=0.1446, without interaction $F_{2, 216}$=1.951, p=0.1446) intake compared to males, across the sessions. This higher liquid intake in females was also observed on the average intake for the three solutions (**Fig. 2C,** unpaired Student t-test, t=2.685, p=0.0084, **Fig. 2E,** unpaired Student t-test, t=10.80, p<0.0001**, Fig. 2G,** unpaired Student t-test, t=10.17, p<0.0001). However, it is important to note that the difference between females and males is larger for ethanol as well as ethanol+quinine, compared to water (effect size Cohen's d=0.04 for water 0.30 for ethanol and 0.25 for ethanol+quinine). Water, ethanol, and ethanol+quinine intake were higher in females than in males when expressed both in absolute volume (mL) and body weight-normalized volume (mL/kg, **Supplementary Fig. S1A-F**). To further analyze the link between binge ethanol drinking and persistent ethanol consumption, we correlated the average ethanol intake with the average ethanol+quinine intake. For male mice, we observed a positive correlation between binge drinking and persistent drinking (**Supplementary Fig. S1G**), and a strong tendency for a positive correlation in females (**Supplementary Fig. S1H**).

Finally, consumption of both ethanol and ethanol adulterated with quinine was independent of the hormonal cycle in female mice (**Supplementary Fig. S1I-K**). To test the strength of persistent ethanol drinking despite quinine aversion, a subgroup of mice that had undergone binge and persistent drinking had access to water+quinine only for 3 sessions, followed by 4 sessions of ethanol+quinine access. Interestingly, both males and females consume more ethanol+quinine

than water+quinine (**Fig. 2H**, 2-way RM-ANOVA, sex factor $F_{1, 46}=20.64$, $p<0.0001$, session factor $F_{6, 276}=23.41$, $p<0.0001$, with an interaction $F_{6, 276}=3.450$, $p=0.0027$). This observation was validated on average, as for both sexes, consumption of ethanol+quinine was larger than water+quinine (**Fig. 2I**, 2-way RM-ANOVA, sex factor $F_{1, 46}=17.85$, $p=0.0001$, liquid factor $F_{1, 46}=85.86$, $p<0.0001$, with an interaction $F_{1, 46}=9.787$, $p=0.0030$). These results validate the appetitive property of ethanol, as its addition to quinine is sufficient to increase quinine drinking in both sexes. Therefore, we named the consumption of ethanol+quinine persistent drinking despite aversion.

Fig. 2 Sex differences in binge and persistent ethanol drinking. A Behavioral timeline during the drinking in the dark procedure. **B, C** Kinetic (B) and average (C) of water intake during cycle 0 in male (n=55) and female (n=56). **D, E** Kinetic (D) and average (E) of ethanol intake during cycles 1 and 2 in male (n=55) and female (n=56). **F, G** Kinetic (F) and average (G) of ethanol+quinine intake during cycle 3 in male (n=55) and female (n=55). **H** Kinetic of water+quinine (water+Q) and ethanol+quinine (ethanol+Q) intake in males (n=22) and females (n=26). **I** Comparison of water+quinine and ethanol+quinine intake during the last session of exposure in male (n=22) and female (n=26) mice. Data are shown as mean ± SEM. **p<0.01, ***p<0.001 represents a significant Student t-test. #p<0.05 ##p<0.01 ###p<0.001 represents a significant Bonferroni post-hoc test.

Neural activity of right aIC and pIC glutamatergic neurons during binge and persistent ethanol drinking

We used fiber photometry with the genetically encoded calcium sensor GCaMP6f (**Fig. 3A, S2A**), to record aIC (**Fig. 3B, S2B**) and pIC (**Fig.3C, S2K**) glutamatergic neuron activity during single lick of different solutions over 1-hour sessions (**Fig. 3D**). Behaviorally, the average ethanol and ethanol+quinine intake across all drinking sessions, excluding the recording session, was higher in females than males (**Fig. 3E,** unpaired Mann-Whitney test, U=145, p=0.0143, **Fig. 3F,** unpaired Student t-test, t=2.357, p=0.0233). For animals with neuronal recordings in the right aIC, a similar average number of single licks or events (i.e., combination of single licks and licking bouts) were analyzed across ethanol, ethanol+quinine, water, and water+quinine conditions (**Supplementary Fig. S2C-J**). Comparable results were obtained in animals recorded in the right pIC (**Supplementary Fig. S2L-O, R, S**), except for the water condition, in which males exhibited a higher number of single licks and events compared to females (**Supplementary Fig. S2P**, unpaired Student's t-test, t=2.382, p=0.0309; **Supplementary Fig. S2Q**, unpaired Mann-Whitney test, U=119, p=0.03).

At the neural level, we segmented the signal recorded in the insular cortex of the right hemisphere in four 3-seconds time windows centered around licking onset: reference (-8 to -5 s), pre-lick (-3 to 0 s), post-lick (0 to +3 s), and post-ingestive (+6 to +9 s, **Fig. 3**). When centered around ethanol consumption, the activity of aIC neurons was different across the four windows, independently of sex (**Fig. 3G, H**, 2-way RM-ANOVA, sex factor $F_{1, 11}$=0.2616, p=0.6191, period factor $F_{3, 33}$=7.374, p=0.0007, without interaction $F_{3, 33}$=0.3421, p=0.7950). In contrast, the activity of pIC neurons was increased during ethanol post-lick compared to the reference window, in both males and females, and was higher in males during the post-ingestive window, compare to the male reference and to the female post-ingestive windows (**Fig. 3I**, 2-way RM-ANOVA, sex factor $F_{1, 15}$=2.916, p=0.1083, period factor $F_{3, 45}$=9.974, p<0.0001, with an interaction $F_{3, 45}$=2.875, p=0.0465).

Regarding ethanol+quinine drinking, aIC neurons activity was different across periods, independently of sex (**Fig. 3J, K,** 2-way RM-ANOVA, sex factor $F_{1, 12}$=0.1030, p=0.7538, period factor $F_{3, 36}$=4.757, p=0.0068, without interaction $F_{3, 36}$=0.7980, p=0.5031). Interestingly, pIC neuronal activity around ethanol+quinine licking varied only in female mice. First, we observed higher pIC neuron activation during the pre-lick period compared to the reference, indicating an anticipatory activation of these neurons. Neuronal activity was further increased during the post-lick period compared to pre-lick, indicating that the peak activity response is occurring triggered by ethanol+quinine licking. Notably, pIC neural activity was higher in females than in males, specifically during this ethanol+quinine post-lick period, when licking occurred (**Fig. 3L**, 2-way RM-ANOVA, sex factor $F_{1, 13}$=3.447, p=0.0862, period factor $F_{3, 39}$=11.75, p<0.0001, with an interaction $F_{3, 39}$=3.717, p=0.0192).

Additionally, we performed fiber photometry recordings during water and water+quinine licking. The activity of aIC neurons was different across periods, independently of sex (**Fig. 3M, N**, 2-way RM-ANOVA, sex factor $F_{1, 15}$=0.09366, p=0.7638, period factor $F_{3, 45}$=3.956, p=0.0138, without interaction $F_{3, 45}$=0.1703, p=0.9159), whereas pIC neuron activity was increased in females during post-lick and post-ingestive periods relative to the reference, and was higher in females than males in the post-ingestive period (**Fig. 3O,** 2-way RM-ANOVA, sex factor $F_{1, 15}$=1.101, p=0.3107, period factor $F_{3, 45}$=6.288, p=0.0012, with an interaction $F_{3, 45}$=2.814, p=0.0498). For water+quinine licking, the activity of aIC neuron was similar across periods, while it was different for pIC neurons, independently of sex (**Fig. 3P-R**, 2-way RM-ANOVA for pIC, sex factor $F_{1, 20}$=0.9419, p=0.3434, period factor $F_{3, 60}$=2.962, p=0.0392, without interaction $F_{3, 60}$=0.8110, p=0.4928). Similar patterns of neuronal activity were observed in the aIC when analyzing the first 5 events (i.e., combination of single licks and licking bouts, **Supplementary Fig. S3A-B, S3D-E, S3G-H, S3J- K**). In the pIC, comparable neuronal activity was observed when analyzing the first 5 events for all liquids (**Supplementary Fig. S3F, S3I, S3L**), except for ethanol where pIC activity was different across periods, without specific effects (**Supplementary Fig. S3C,** 2-way RM-ANOVA, sex factor $F_{1, 16}$=0.6862, p=0.4196, period factor $F_{3, 48}$=13.06, p<0.0001, without interaction $F_{3, 48}$=0.5825, p=0.6294).

Furthermore, we assessed whether consumption history calculated as the average ethanol or ethanol+quinine intake across all sessions except the recording session, correlated with aIC and pIC neuronal calcium signals during consumption. Activity of aIC neurons did not correlate with intake history in either sex (**Supplementary Fig. S4A-D**). Similarly, pIC activity showed no correlation with ethanol intake history in males or females (**Supplementary Fig. S4E, F**), nor with ethanol+quinine intake in males (**Supplementary Fig. S4G**). However, females exhibited a trend toward a positive correlation between ethanol+quinine intake history and pIC activity (**Supplementary Fig. S4H**, Pearson's R^2=0.46, p=0.07).

Fig. 3 Coding properties of anterior (aIC) and posterior (pIC) insular cortex excitatory neurons during single licks of binge and persistent ethanol consumption. A Viral strategy to record calcium changes in aIC or pIC excitatory neurons. A viral vector carrying the gene coding for the calcium sensor GCaMP6f was injected unilaterally in the aIC or the pIC. **B, C** Representative image of GCaMP6f expression in aIC (B) or pIC (C) excitatory neurons. The mouse brain atlas delineation has been overlaid to the image, with the borders of the aIC or pIC in bold. **D** Conditions during photometry recordings to synchronize calcium signal and licking behavior. **E** Average ethanol intake during all binge-drinking sessions except the recording day in male (n=21) and female (n=24) mice. **F** Average ethanol+quinine intake during all persistent drinking sessions except the recording day in male (n=21) and female (n=22) mice. **G** Peri-ethanol licks analysis of the calcium signal in the aIC (up) and pIC (bottom) during single ethanol licks in male and female mice. **H, I** Average of calcium signal in the aIC (H) or pIC (I) during ethanol licking for the reference, pre-lick, post-lick, and post-ingestive periods in male (n=8 for aIC and n=7 for pIC) and female (n=5 for aIC and n=10 for pIC) mice. **J** Peri-ethanol+quinine licks analysis of the calcium signal in the aIC (up) and pIC (bottom) single ethanol+quinine licks in male and female mice. **K, L** Average of calcium signal in the aIC (K) or pIC (L) during ethanol+quinine licking for the reference, pre-lick, post-lick, and post-ingestive periods in male (n=7 for aIC and n=7 for pIC) and female (n=7 for aIC and n=8 for pIC) mice. **M** Peri-water licks analysis of the calcium signal in the aIC (up) and pIC (bottom) during single water licks in male and female mice. **N, O** Average of calcium signal in the aIC (N) or pIC (O) during water licking for the reference, pre-lick, post-lick, and post-ingestive periods in male (n=9 for aIC and n=10 for pIC) and female (n=8 for aIC and n=7 for pIC) mice. **P** Peri-water+quinine licks analysis of the calcium signal in the aIC (up) and pIC (bottom) during single water+quinine licks in male and female mice. **Q, R** Average of calcium signal in the aIC (Q) or pIC (R) during water+quinine licking for the reference, pre-lick, post-lick, and post-ingestive periods in male (n=6 for aIC and n=12 for pIC) and female (n=7 for aIC and n=10 for pIC) mice. Data are shown as mean ± SEM. *p<0.05 **p<0.01 ***p<0.001 represent significant Student t-test, or Mann-Whitney test, or a main effect of the 2-way RM ANOVA. #p<0.05, ##p<0.01, ¤p<0.05, ¤¤p<0.01, ¤¤¤p<0.01 represent significant Bonferroni post-hoc tests.

Chemogenetic inhibition of aIC and pIC glutamatergic neurons during binge and persistent ethanol drinking

To study the causal role of the aIC and pIC excitatory neurons in binge ethanol drinking and persistent ethanol drinking (ethanol adulterated with quinine), we used a chemogenetic approach (**Fig. 4A-C**) to inhibit aIC or pIC glutamatergic neurons by expressing hM4Di, under the control of the CaMKIIα promoter. Beforehand, we tested the viral approach using *in-vivo* electrophysiology and confirmed a decrease in the average firing rate from 30 to 55 minutes after CNO injection compared to baseline (**Supplementary Fig. S4A, B**, paired Wilcoxon test, p=0.0232). Chemogenetic inhibition of aIC glutamatergic neurons (**Supplementary Fig. S5C**) did not influence ethanol drinking in both sexes (**Fig. 4D**). The same neuronal manipulation reduced ethanol+quinine drinking independently of sex (**Fig. 4E**, 2-way ANOVA, sex factor $F_{1,48}$=15.78, p=0.0002, virus factor $F_{1,48}$=6.180, p=0.0165, interaction $F_{1,48}$=3.151, p=0.0822). Moreover, inhibition of aIC neurons did not change water consumption in males or females (**Fig. 4F**), whereas it decreased water+quinine drinking independently of the sex (**Fig. 4G**, 2-way ANOVA, sex factor $F_{1,25}$=1.541, p=0.2259, virus factor $F_{1,25}$=14.72, p=0.0008, interaction $F_{1,25}$=3.839, p=0.0613).

Similarly, to the aIC, inhibition of pIC excitatory neurons (**Supplementary Fig. S5D**) did not influence ethanol drinking in both sexes (**Fig. 4H**). However, pIC inhibition reduced ethanol+quinine intake exclusively in female mice (**Fig. 4I**, 2-way ANOVA, sex factor $F_{1,31}$=58.57, p<0.0001, virus factor $F_{1,31}$=14.24, p=0.0007, with an interaction $F_{1,31}$=5.263, p=0.0287). As for aIC inhibition, pIC silencing also did not alter water consumption in males nor females (**Fig. 4J**). However, in contrary to aIC, pIC silencing altered water+quinine intake in male mice (**Fig. 4K**, 2-way ANOVA, sex factor $F_{1,24}$=6.284, p=0.0194, virus factor $F_{1,24}$=7.063, p=0.0138, with an interaction $F_{1,24}$=4.582, p=0.0427).

To rule out any behavioral confound due to the treatment, CNO or vehicle injection was performed in control mice. CNO injection in control animals did not alter ethanol, ethanol+quinine or water intake compared to control animals injected with vehicle (**Supplementary Fig. S5E-G, I-K**). Finally, we found that locomotion was similar between control mice injected with vehicle or CNO, and mice expressing the inhibitory DREADDs injected with CNO (**Supplementary Fig. S5H, L**). Together, our results demonstrate a sex-dependent function of the antero-posterior insula axis on drinking behaviors. The aIC neurons are recruited for bitter solution drinking independently of sex, whereas the pIC sustains persistent ethanol drinking in females, and water+quinine drinking in males.

Fig. 4 Chemogenetic inhibition of anterior (aIC) or posterior (pIC) insular cortex excitatory neurons during binge and persistent ethanol intake. A Experimental design to inhibit aIC and pIC excitatory neurons. A viral vector carrying the gene coding for the inhibitory receptor hM4Di fused to the fluorescent reporter mCherry, or the control virus, containing only the gene for mCherry, was bilaterally injected in the aIC or pIC. **B**, **C** Representative images of the chemogenetic viral vector expression in aIC (B) and pIC (C) neurons. The mouse brain atlas delineation has been overlaid on the image, with the borders of the aIC and pIC in bold. **D** Average ethanol intake during chemogenetic manipulation of aIC glutamatergic neurons (day 18) in male (control n=16, hM4Di n=11) and female (control n=14, hM4Di n=12) mice. **E** Average of ethanol+quinine intake during chemogenetic manipulation of aIC neurons (day 25), in male (control n=16, hM4Di n=11) and female (control n=14, hM4Di n=11) mice. **F** Average water intake during chemogenetic manipulation of aIC neurons (day 29), in male (control n=16, hM4Di n=11) and female (control n=14, hM4Di n=11) mice. **G** Average of water+quinine intake during chemogenetic manipulation of aIC neurons (day 29), in male (n=6 for both groups) and female (control n=9, hM4Di n=8) mice. **H** Average ethanol intake during chemogenetic manipulation of pIC neurons (day 18), in male (control n=8, hM4Di n=9) and female (control n=8, hM4Di n=10) mice. **I** Average of ethanol+quinine intake during chemogenetic manipulation of pIC neurons (day 25), in male (control n=8, hM4Di n=9) and female (control n=8, hM4Di n=10) mice. **J** Average of water intake during chemogenetic manipulation of pIC neurons (day 29), in male (control n=8, hM4Di n=9) and female (control n=8, hM4Di n=10) mice. **K** Average of water+quinine intake during chemogenetic manipulation of pIC neurons (day 29), in an independent group of males (control n=7, hM4Di n=6) and female (control n=7, hM4Di n=8) mice. Data are shown as mean ± SEM. *p<0.05 ***p<0.001 represents the significant main effect of the 2-way ANOVA. ###p<0.001 represents a significant Bonferroni post-hoc test.

DISCUSSION

The goal of our study was to elucidate whether the aIC and pIC exhibit sex-dependent functions in ethanol binge drinking and ethanol persistent drinking despite bitter aversion in mice. First, we observed a higher binge and persistent ethanol intake in females compared to males (**Fig. 2E, G**). Second, at the neuronal activity level, we showed that aIC neuronal activity was increased around ethanol, ethanol+quinine, and water across all periods, independently of sex (**Fig. 3H, K, N**), but not around water+quinine consumption (**Fig. 3Q**). In addition, right pIC neuronal activity increased in both sexes during ethanol drinking (**Fig. 3I**), and only in females during ethanol+quinine and water consumption (**Fig. 3L, O**). However, sex differences during licking behavior were observed only for ethanol+quinine drinking, where pIC neural activity was higher in females than in males. For water+quinine licking, we observed increased activity of pIC neurons during water+quinine licking across periods, independently of sex (**Fig. 3R**). Additionally, we demonstrated that chemogenetic inhibition of aIC glutamatergic neurons reduced the intake of quinine-adulterated solutions (e.g., ethanol and water, **Fig. 4E, G**) regardless of sex, whereas chemogenetic inhibition of pIC glutamatergic neurons reduced persistent ethanol drinking exclusively in females and quinine drinking in males (**Fig. 4I, K**). Altogether, these findings suggest a topographical and sex-dependent divergent function of the insular cortex in bitter liquid drinking, highlighting the pIC glutamatergic neurons as a substrate of persistent ethanol drinking in female mice.

Sex-dependent behavior on binge and persistent ethanol drinking
Our results repeatedly showed a higher binge and persistent ethanol intake in females compared to males, independently of the estrous cycle (**Supplementary Fig. S1J, K**), which is consistent with and extends previous literature highlighting sex differences in drug addiction (64,65). Indeed, multiple preclinical studies also reported a higher binge (13,28,62) and persistent (13,29,66) ethanol intake per body mass in females compared to males. Importantly, to rule out any confound on sex-dependent aversion resistance for quinine, we adulterated the ethanol solution with 500 μM quinine, a concentration for which both male and female mice have similar aversive thresholds (67). Therefore, the sex difference observed in the consumption of ethanol adulterated with quinine is independent of the aversion sensitivity but driven by the motivation to consume ethanol. We further demonstrated that females exhibited higher (BEC) than males after 30 minutes of alcohol consumption (**Fig. 1F**). This difference may reflect sex-dependent variations in ethanol metabolism and absorption. Indeed, Robinson et al. (68) reported that, following intragastric administration of ethanol (1 g/kg), BEC peaked more rapidly in female than male rats, indicating faster gastric absorption in females. In addition, several studies have shown that females metabolize ethanol more rapidly than males (69,70), which supports our results showing that, after 2 hours of ethanol intake, a lower percentage of females reaches the BEC intoxication threshold of 80 mg/dL than males (**Fig. 1C**).
A key advantage of the DID model is that it rapidly induces high levels of ethanol drinking and intoxication in mice without food or water deprivation, similar to what is observed during human binge alcohol drinking. The protocol can also be adapted to model persistent drinking despite aversion by adding quinine to ethanol. This within-animal approach allows investigation of the transition from episodic binge drinking to persistent alcohol consumption. While quinine adulteration is widely used to mimic persistent ethanol drinking despite aversive consequences, it does not capture the broader social, legal, and health negative consequences associated with AUD in humans, nor its slow, multifactorial development. We acknowledge this limit in our study, and for this reason, we chose to refer to "persistent" rather than "compulsive" ethanol intake.

Coding properties of right aIC glutamatergic neurons during binge and persistent ethanol intake
Our results revealed that right aIC excitatory neurons are activated during ethanol and ethanol+quinine consumption across periods independently of sex (**Fig. 3H, K**). A recent study

demonstrated that the projection from the aIC to the dorsolateral striatum encodes binge ethanol consumption in males but not females (61). In contrast, our recordings measured the global glutamatergic neuronal population in the aIC, which may account for the differing results. These findings suggest that sex-dependent functions of the aIC during binge drinking may be mediated by distinct neural projections. For persistent ethanol consumption, few studies have investigated the aIC neuronal activity after ethanol+quinine drinking. Chen and Lasek reported an increased expression of the neuronal activity marker *cFos* in the aIC after ethanol+quinine intake compared to water, ethanol and quinine drinking in male mice (71). Additionally, a recent *in vivo* electrophysiological study in male rats demonstrated that aIC neurons with increased firing at ethanol drinking onset also enhanced their activity during ethanol+quinine consumption, demonstrating that aIC neurons in males code both binge and persistent ethanol drinking (72). However, to our knowledge, no study has examined the dynamics of aIC activation in females during persistent alcohol consumption. Thus, our results corroborate previous findings and confirm the role of the right aIC in persistent consumption in males, and extend this to females.

Finally, our results showed increased response of right aIC glutamatergic neurons around water consumption (**Fig. 3N**), but not water+quinine intake (**Fig. 3Q**). For water intake, this is in line with and extends the findings of a recent study highlighting that aIC neurons were activated after water licking in male mice (73). For water+quinine consumption, Chen and colleagues revealed aIC activation after quinine tasting (71). However, these findings were obtained using a two-bottle choice paradigm, in which quinine was presented alongside with water, allowing animals to choose between the two solutions. This procedural difference likely accounts for the discrepancy with our results, as the availability of a competing non-aversive solution engages specific choice and value-comparison circuits, which include the aIC, that is not recruited in our single-solution DID paradigm.

Function of aIC excitatory neurons in bitter taste processing
Our results revealed that aIC neuron inhibition does not alter binge ethanol intake in both sexes. These results support and extend Haaranen et al. study, which showed that chemogenetic inhibition of aIC neurons did not affect alcohol intake in alcohol-preferring male rats in a two-bottle choice model (58). However, it contrasts with previous literature demonstrating that chemogenetic manipulation of aIC neurons modulates alcohol binge drinking, although the results are divergent. Indeed, while silencing aIC neurons increased operant responses for alcohol (41), it decreased alcohol intake in a two-bottle choice paradigm (74). Altogether, these results suggest that aIC function in ethanol binge drinking might be dependent on the animal model used. Moreover, while we specifically manipulated aIC glutamatergic neurons, the studies cited previously manipulated all aIC mature neurons independently of their phenotype, which could explain some discrepancy in the results observed.

We also demonstrated that inhibition of aIC glutamatergic neurons reduced the intake of solutions adulterated with quinine (e.g., ethanol or water) independently of sex. These results confirm and extend previous experiments performed in male rats and showing a reduction of ethanol+quinine intake after inhibition of aIC neurons, as well as inhibition of aIC projections to locus coeruleus or nucleus accumbens core (74,75). Thus, our results bring additional and complementary information to these studies.

Regarding the role of aIC in water+quinine intake, the literature is divergent and, to our knowledge, only addresses this question in males so far. On one hand, optogenetic inhibition or lesion of the aIC respectively had no effect (76,77) or decreased (78) sensitivity to quinine, leading to an increase in intake. On the other hand, activation of aIC neurons and their projection to the basolateral amygdala also decreased quinine sensitivity (47,79). As in our study aIC excitatory neuron inhibition had no effect on water+quinine intake, our findings are consistent with two studies (76,77) and contrast with three others (42, 28, 69). Moreover, we chose to not water-deprive our animals to observe more ethological behavioral responses. In contrast, in the other

studies, animals were water-deprived before quinine exposure, which could likely have induced a physiological stress altering aIC response to quinine. In addition, Staszko et al., have shown that previous taste experience with water adulterated with quinine decreases the number of active cells and neuronal response in the aIC (80). Thus, in our study, the absence of modulation of water+quinine intake by aIC neuron inhibition could be explained by pre-exposure to quinine before chemogenetic manipulation. Finally, *cFos*+ neuron distribution in aIC after quinine exposition significantly varies across aIC subareas and the median-lateral axis (81), whereas in our study we targeted the entire aIC. Thus, further investigations are necessary to sharpen our understanding of aIC neurons in aversive solution consumption in both sexes.

Function of the insular cortex in processing information from the peripheral nervous system
The insular cortex is a key brain region integrating taste perception and post-ingestive effects of drug intake. For example, Naqvi and Bechara (82), proposed that the insular cortex integrates the conscious pleasure generated by the bodily effect of drug intake and the pain following withdrawal. In the context of AUD, alcohol taste activates gustatory pathways, including the insula (83), and a recent study demonstrated a positive correlation between aversive taste reactivity to orally delivered 20% EtOH and pIC cFos+ cells in rats (84). In addition, fMRI studies revealed that the insular cortex is activated after gastric distention in rodents (85,86), and that the pIC in particular receives interoceptive visceral information (87). However, to our knowledge, no studies have investigated how peripheral information after alcohol ingestion is integrated by the insula to regulate subsequent alcohol consumption in rodents. Additional studies are necessary to understand how the integration of gustatory and post-ingestive signals by the insula shapes alcohol consumption in both sexes.

Functional sex differences in right pIC coding properties of persistent ethanol drinking
To decipher how pIC excitatory neurons encode binge drinking and persistent ethanol drinking, we performed *in vivo* fiber photometry recordings of the right pIC. We observed increased activity of right pIC glutamatergic neurons in response to ethanol in both sexes (**Fig. 3I**). In contrast, activity increased only in females for ethanol+quinine and water (**Fig. 3L, O**). Importantly, sex differences during the licking period itself are detected exclusively during ethanol+quinine drinking with females showing higher right pIC neuron activity than males (**Fig. 3L**). In contrast, during ethanol and water consumption, sex differences emerged after the licking behavior, within the post-ingestive period (+6 to +9 s). Finally, during water+quinine drinking, pIC neuronal activity was different across periods, independently of sex (**Fig. 3R**).
Our results showed a sex-dependent functional dynamic of right pIC neuronal activity in persistent ethanol drinking (**Fig. 3L**). In physiological conditions, Lezzi et al. showed that pIC neurons in females are less excitable than in males (89). In contrast, a recent study demonstrated that short ethanol exposure increases the intrinsic excitability of pIC neurons projecting to the bed nucleus of the stria terminalis, in female mice only (62). This study, strengthened by our present results, suggests that ethanol consumption modulates pIC neuronal excitability. Altogether, it highlights pIC glutamatergic neurons and their projections as a key neural population supporting sex-dependent phenotypes in alcohol-related behaviors. Moreover, we revealed sex differences in pIC neuronal activity in the post-ingestive period after ethanol and water consumption, which was higher in males for ethanol, and higher in female for water (**Fig. 3I, O**). These results may reflect a sex-specific processing of post-ingestive interoceptive signals by the pIC, rather than differences strictly related to the consummatory behavior. Finally, right pIC neuronal activity increased during water+quinine drinking, which is consistent with previous studies reporting that pIC neurons encode bitter taste, independently of the sex (49,88).
Importantly, some literature suggests a lateralization of insular cortex functions, including water and alcohol drinking. Indeed, several neuroimaging studies report lateralized right-sided posterior insula atrophy in AUD patients (56,90,91). For this reason, we focused fiber photometry recordings

on the right insula. However, unilateral recordings remain one of the limitations of our study. Understanding the lateralization of insular cortex function in alcohol consumption is an important question that will require further investigation.

<u>Uncovering sex-dependent pIC function in persistent ethanol intake</u>
Fiber photometry recordings reflect the functional dynamics of pIC glutamatergic neurons in response to ethanol intake, however, it does not provide information on its causal role on persistent ethanol intake. To address this question, we used chemogenetic manipulations to demonstrate the necessity of pIC excitatory neurons to change persistent ethanol drinking in females. We demonstrated sex differences in the function of pIC glutamatergic neurons in drinking solutions adulterated with bitter taste. To our knowledge, these results are the first demonstration of a sex-dependent function of pIC neurons on persistent ethanol drinking. Indeed, in males, inhibition of pIC glutamatergic neurons reduced only water adulterated with quinine intake (**Fig. 4K**), while in females, the same neuronal manipulation decreased exclusively persistent ethanol drinking (**Fig. 4I**). Previous literature highlighted the role of the insular cortex in aversive taste processing (88,92). While the function in bitter tastant has been first dedicated to the pIC (47), multiple studies revealed that the distribution of neurons activated by quinine is not strictly specific to the pIC and can be found in the entire insular cortex (73–76). However, so far, no study has investigated the causal role of the insular cortex in the anteroposterior axis for bitter-tastant solution drinking. Therefore, our results provide new evidence on how the insular cortex influences bitter-solutions drinking in a sex-dependent manner. Using the two-bottle choice paradigm, a recent study showed a higher expression of the neuronal activity marker *cFos* in the pIC of males compared to females after ethanol+quinine intake (93). Importantly, in this study ethanol was adulterated with 100 µM of quinine. A previous report suggests that females are not sensitive to ethanol adulterated with quinine at this concentration whereas males are (29) which could explain the discrepancy with our results where we used a concentration of quinine at 500 µM. Furthermore, using an operant alcohol drinking model in male rats, pharmacological inactivation of pIC neurons reduces alcohol intake (59) while in our study, chemogenetic inhibition of pIC excitatory neurons did not change ethanol drinking nor in males neither in females. Several methodological aspects can, at least in part, support this discrepancy. First, operant and voluntary alcohol drinking could recruit different neuronal circuits, second, the methods used for the neuronal inhibition (e.g. pharmacology vs chemogenetics) and the animal model used (e.g. rat vs mouse) were different.
To rule out the possibility that pIC glutamatergic inhibition was due to the non-selective effects of CNO, we performed multiple control experiments. Indeed, it has been shown that CNO can bind to non-DREADD targets, and does not cross the blood-brain barrier. However, clozapine, a CNO metabolite, is reaching the brain and can also have multiple pharmacological targets that can induce several physiological and behavioral effects (94,95). In our control experiments, we found that CNO injection in mice expressing the control viral vector in pIC had no effects on either liquid consumption (e.g., ethanol, ethanol+quinine or water) or locomotion. Furthermore, CNO injection in mice injected with the viral vector carrying the gene coding for hM4Di also did not affect locomotion. These results indicate that CNO does not have non-specific effects either on drinking or locomotion and secondly, that the inhibition of pIC excitatory neurons did not affect locomotion. Overall, these results confirm that the behavioral effect of CNO on ethanol+quinine drinking in females and water+quinine intake in males is due to selective inhibition of pIC excitatory neurons. Finally, in our study the manipulation and the recording of pIC excitatory neurons was performed regardless of their outputs. However, the pIC is a highly wired region sending dense projections to several regions (49,96), including the central amygdala, a brain region highly involved in alcohol-related behaviors and supporting sex differences in alcohol intake (97–99). Therefore, future investigations will be crucial to understand the sex-dependent function of pIC neurons in alcohol-related behaviors at the circuit level.

<u>Theoretical circuit mechanism within the pIC underlying sex differences in persistent ethanol drinking in mice</u>

Persistent ethanol drinking typically reflects a behavior in which reward and aversion counterbalance each other, driving the pursuit of the reward (e.g., ethanol) despite the associated aversion (e.g., quinine). In our study, female mice drink more ethanol+quinine than males (**Fig. 2G**), indicating that the balance between both modalities is shifted toward the reward. At the neuronal level, we demonstrated that this sex difference engages pIC glutamatergic neurons (**Fig. 3L, 4I**). Interestingly, the literature reports that pIC neurons can mediate both reward and aversive behavior in a projection-specific manner. Indeed, optogenetic activation of pIC to the central amygdala (CeA) induces avoidance behavior and learning in male and female mice (100), showing that this pathway drives aversive-like behaviors in both sexes. In contrast, using a nosepoke-triggered photostimulation paradigm or a real-time place preference task, Girven and colleagues showed that photostimulation of pIC to the bed nucleus of the stria termianlis (BNST) terminals is rewarding in both male and female mice (101), demonstrating that this pIC output mediates reward-related behaviors. Thus, in our experimental context, both pIC-CeA and pIC-BNST projections may be, respectively, engaged by the aversive and rewarding components of ethanol+quinine. Moreover, ethanol drinking selectively enhances the excitatory properties of pIC to BNST neurons in females but not males (62). Consequently, ethanol drinking may potentiate the reward-associated pIC-BNST pathway in females, shifting circuit engagement toward reward over the pIC-CeA aversion circuit, thus promoting higher persistent ethanol drinking in females than males. Finally, this mechanism may additionally reflect modulation through local GABAergic microcircuitry in the pIC (**Supplementary Fig. S6**).

<u>Translation relevance</u>

Human studies have primarily examined neurobiological alterations of the aIC in AUD men and women, reporting reduced gray matter density compared to healthy controls (50–53). In addition, AUD men and women had reduced aIC volume compared to healthy controls (50), which is associated with higher compulsivity and impulsivity. In our study, we showed that aIC encodes and is required for persistent ethanol consumption (**Fig. 3K, 4E**) in both sexes, which is in line with the human literature supporting that the anatomy and function of the aIC are altered in AUD patients of both sexes. In contrast to aIC, less information is known about the role of the pIC in AUD patients. Only two studies reported decreased gray matter volume in AUD men and women (56) and elevated resting-state activity in AUD women (55), making it difficult to draw strong conclusions regarding its potentially sexually dimorphic role in AUD. In our study, we identify sex differences in pIC function during persistent ethanol intake in mice, laying the groundwork for future translational investigations of this subregion in men and women suffering from AUD.

<u>Conclusions</u>

To conclude, our study demonstrated that aIC and pIC play different roles in drinking behaviors. While right aIC glutamatergic neurons encode binge and persistent ethanol drinking independently of sex, right pIC glutamatergic neurons are active during persistent ethanol consumption exclusively in females. In line with these results, inhibition of posterior insular cortex (pIC) excitatory neurons selectively reduced intake of ethanol adulterated with quinine in female mice, indicating a sex-specific role of pIC glutamatergic neurons in mediating persistent ethanol consumption despite aversion. In contrast, inhibition of aIC glutamatergic neurons reduced bitter solution drinking independently of sex. Altogether, our results provide a starting point to characterize further the role of aIC and pIC circuits in sex-dependent addictive-related behaviors.

ACKNOWLEDGEMENTS

The authors acknowledge the entire Beyeler Lab. We thank Harold Haun and Irene Lorrai for their helpful feedback on setting up the drinking in the dark model. We thank Guillaume Ferreira for the

interesting discussions. We thank Sara Laumond, Julie Tessaire, Nathalie Aubailly, and the technical staff of the animal housing facility of the Neurocentre Magendie (INSERM U1215) for their invaluable support.

We acknowledge the support of the Région Nouvelle-Aquitaine, the 'Avenir program' of the French Institute of Health (INSERM), the 'Fondation NRJ-Institut de France', the 'Agence Nationale pour la Recherche' (ANR), the 'Fondation Schlumberger pour l'Education et la Recherche' (FSER), and the 'Fondation Bettencourt-Schueller' to AB. We are grateful for the support of the 'Fondation pour la Recherche Médicale' (FRM) to AB, CN (ARF201909009147) and CF (FDT202404018152), as well as the 'ATIP program' (CNRS) to CN. Finally, this research was supported by the French Addictions Fund (Fonds de lutte contre les addictions) under the 2024 call for proposals on Addictive Behaviors and Drugs (CAD): Prevention, Mechanisms, Identification, and Support, jointly led by IReSP and INCa (CAD-V124-014) to CN.

CONFLICT OF INTEREST

The authors declare that they do not have any competing interests or conflicts of interest (financial or non-financial) related to the material presented in this manuscript.

REFERENCES

1. Crocq MA, Guelfi JD. DSM-5: manuel diagnostique et statistique des troubles mentaux. 5e éd. Issy-les-Moulineaux: Elsevier Masson; 2015.
2. Moos RH, Moos BS. Treated and untreated alcohol-use disorders: course and predictors of remission and relapse. Eval Rev. 2007 Dec;31(6):564–84.
3. Koob GF, Volkow ND. Neurocircuitry of Addiction. Neuropsychopharmacology [Internet]. 2010 Jan [cited 2023 Sep 21];35(1):217–38. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2805560/
4. Organisation mondiale de la santé, editor. Global status report on alcohol and health 2018. Geneva: World health organization; 2018.
5. NIAAA. Drinking Levels Defined | National Institute on Alcohol Abuse and Alcoholism [Internet]. 2023 [cited 2023 Sep 21]. Available from: https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/moderate-binge-drinking
6. Gowin JL, Sloan ME, Stangl BL, Vatsalya V, Ramchandani VA. Vulnerability for Alcohol Use Disorder and Rate of Alcohol Consumption. Am J Psychiatry [Internet]. 2017 Nov [cited 2023 Sep 21];174(11):1094–101. Available from: http://ajp.psychiatryonline.org/doi/10.1176/appi.ajp.2017.16101180
7. Tavolacci MP, Berthon Q, Cerasuolo D, Dechelotte P, Ladner J, Baguet A. Does binge drinking between the age of 18 and 25 years predict alcohol dependence in adulthood? A retrospective case–control study in France. BMJ Open [Internet]. 2019 May [cited 2023 Sep 21];9(5):e026375. Available from: https://bmjopen.bmj.com/lookup/doi/10.1136/bmjopen-2018-026375
8. Hill KG, White HR, Chung IJ, Hawkins JD, Catalano RF. Early Adult Outcomes of Adolescent Binge Drinking: Person- and Variable-Centered Analyses of Binge Drinking Trajectories. Alcohol Clin Exp Res. 2000 Jun;24(6):892–901.
9. Rhodes WA, Singleton E, McMillan TB, Perrino CS. Does knowledge of college drinking policy influence student binge drinking? J Am Coll Health J ACH. 2005 Aug;54(1):45–9.
10. Thiele TE, Navarro M. "Drinking in the dark" (DID) procedures: a model of binge-like ethanol drinking in non-dependent mice. Alcohol Fayettev N. 2014 May;48(3):235–41.
11. Wilcox MV, Cuzon Carlson VC, Sherazee N, Sprow GM, Bock R, Thiele TE, et al. Repeated binge-like ethanol drinking alters ethanol drinking patterns and depresses striatal GABAergic transmission. Neuropsychopharmacol Off Publ Am Coll Neuropsychopharmacol. 2014 Feb;39(3):579–94.
12. Siciliano CA, Noamany H, Chang CJ, Brown AR, Chen X, Leible D, et al. A cortical-brainstem circuit predicts and governs compulsive alcohol drinking. Science. 2019 Nov 22;366(6468):1008–12.
13. Sneddon EA, White RD, Radke AK. Sex Differences in Binge-Like and Aversion-Resistant Alcohol Drinking in C57BL/6J Mice. Alcohol Clin Exp Res. 2019 Feb;43(2):243–9.

14. Bauer MR, McVey MM, Boehm SL. Three Weeks of Binge Alcohol Drinking Generates Increased Alcohol Front-Loading and Robust Compulsive-Like Alcohol Drinking in Male and Female C57BL/6J Mice. Alcohol Clin Exp Res. 2021 Mar;45(3):650–60.
15. Schuh KM, Sneddon EA, Nader AM, Muench MA, Radke AK. Orbitofrontal cortex subregion inhibition during binge-like and aversion-resistant alcohol drinking. Alcohol. 2022 Mar;99:1–8.
16. Slade T, Chapman C, Swift W, Keyes K, Tonks Z, Teesson M. Birth cohort trends in the global epidemiology of alcohol use and alcohol-related harms in men and women: systematic review and metaregression. BMJ Open. 2016 Oct;6(10):e011827.
17. Grant BF, Chou SP, Saha TD, Pickering RP, Kerridge BT, Ruan WJ, et al. Prevalence of 12-Month Alcohol Use, High-Risk Drinking, and DSM-IV Alcohol Use Disorder in the United States, 2001-2002 to 2012-2013: Results From the National Epidemiologic Survey on Alcohol and Related Conditions. JAMA Psychiatry. 2017 Sep 1;74(9):911–23.
18. Ehlers CL, Gizer IR, Vieten C, Gilder A, Gilder DA, Stouffer GM, et al. Age at regular drinking, clinical course, and heritability of alcohol dependence in the San Francisco Family Study: a gender analysis. Am J Addict Am Acad Psychiatr Alcohol Addict. 2010 Mar 1;19(2):101–10.
19. Diehl A, Croissant B, Batra A, Mundle G, Nakovics H, Mann K. Alcoholism in women: is it different in onset and outcome compared to men? Eur Arch Psychiatry Clin Neurosci. 2007 Sep;257(6):344–51.
20. Erol A, Karpyak VM. Sex and gender-related differences in alcohol use and its consequences: Contemporary knowledge and future research considerations. Drug Alcohol Depend. 2015 Nov 1;156:1–13.
21. Eagon PK. Alcoholic liver injury: Influence of gender and hormones. World J Gastroenterol WJG. 2010 Mar 21;16(11):1377–84.
22. Green CA, Perrin NA, Polen MR. Gender differences in the relationships between multiple measures of alcohol consumption and physical and mental health. Alcohol Clin Exp Res. 2004 May;28(5):754–64.
23. Zachry JE, Johnson AR, Calipari ES. Sex Differences in Value-Based Decision Making Underlie Substance Use Disorders in Females. Alcohol Alcohol Oxf Oxfs [Internet]. 2019 Jul [cited 2024 Mar 11];54(4):339–41. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6671483/
24. Strong MN, Yoneyama N, Fretwell AM, Snelling C, Tanchuck MA, Finn DA. "Binge" drinking experience in adolescent mice shows sex differences and elevated ethanol intake in adulthood. Horm Behav. 2010 Jun;58(1):82–90.
25. Loi B, Colombo G, Maccioni P, Carai MAM, Franconi F, Gessa GL. High alcohol intake in female Sardinian alcohol-preferring rats. Alcohol Fayettev N. 2014 Jun;48(4):345–51.
26. Priddy BM, Carmack SA, Thomas LC, Vendruscolo JCM, Koob GF, Vendruscolo LF. Sex, strain, and estrous cycle influences on alcohol drinking in rats. Pharmacol Biochem Behav. 2017 Jan;152:61–7.
27. Satta R, Hilderbrand ER, Lasek AW. Ovarian Hormones Contribute to High Levels of Binge-Like Drinking by Female Mice. Alcohol Clin Exp Res. 2018 Feb;42(2):286–94.
28. Evans O, Rodríguez-Borillo O, Font L, Currie PJ, Pastor R. Alcohol Binge Drinking and Anxiety-Like Behavior in Socialized Versus Isolated C57BL/6J Mice. Alcohol Clin Exp Res. 2020 Jan;44(1):244–54.
29. Fulenwider HD, Nennig SE, Price ME, Hafeez H, Schank JR. Sex Differences in Aversion-Resistant Ethanol Intake in Mice. Alcohol Alcohol. 2019 Jul 1;54(4):345–52.
30. Chuong V, Farokhnia M, Khom S, Pince CL, Elvig SK, Vlkolinsky R, et al. The glucagon-like peptide-1 (GLP-1) analogue semaglutide reduces alcohol drinking and modulates central GABA neurotransmission. JCI Insight. 8(12):e170671.
31. Gage GA, Muench MA, Jee C, Kearns DN, Chen H, Tunstall BJ. Intermittent-access operant alcohol self-administration promotes binge-like drinking and drinking despite negative consequences in male and female heterogeneous stock rats. Neuropharmacology. 2023 Sep;235:109564.
32. Schramm-Sapyta NL, Francis R, MacDonald A, Keistler C, O'Neill L, Kuhn CM. Effect of sex on ethanol consumption and conditioned taste aversion in adolescent and adult rats. Psychopharmacology (Berl). 2014 Apr;231(8):1831–9.
33. Cailhol S, Mormède P. Sex and Strain Differences in Ethanol Drinking: Effects of Gonadectomy. Alcohol Clin Exp Res. 2001;25(4):594–9.
34. Gereau GB, Torruella-Suárez ML, Sizer SE, Xia M, Zhou D, Wykoff LA, et al. GABA release from central amygdala neurotensin neurons differentially modulates ethanol consumption in male and female mice. Neuropsychopharmacology [Internet]. 2024 Jun [cited 2024 Sep 12];49(7):1151–61. Available from: https://www.nature.com/articles/s41386-024-01830-5

35. Venniro M, Caprioli D, Zhang M, Whitaker LR, Zhang S, Warren BL, et al. The anterior insular cortex→central amygdala glutamatergic pathway is critical to relapse after contingency management. Neuron. 2017 Oct 11;96(2):414-427.e8.

36. Ghareh H, Alonso-Lozares I, Schetters D, Herman RJ, Heistek TS, Van Mourik Y, et al. Role of anterior insula cortex in context-induced relapse of nicotine-seeking. eLife. 11:e75609.

37. Nicolas C, Tauber C, Lepelletier FX, Chalon S, Belujon P, Galineau L, et al. Longitudinal Changes in Brain Metabolic Activity after Withdrawal from Escalation of Cocaine Self-Administration. Neuropsychopharmacology. 2017 Sep;42(10):1981–90.

38. Campbell EJ, Flanagan JPM, Walker LC, Hill MKRI, Marchant NJ, Lawrence AJ. Anterior Insular Cortex is Critical for the Propensity to Relapse Following Punishment-Imposed Abstinence of Alcohol Seeking. J Neurosci [Internet]. 2019 Feb 6 [cited 2023 Sep 21];39(6):1077–87. Available from: https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.1596-18.2018

39. Campbell EJ, Lawrence AJ. It's more than just interoception: The insular cortex involvement in alcohol use disorder. J Neurochem [Internet]. 2021 Jun [cited 2023 Sep 21];157(5):1644–51. Available from: https://onlinelibrary.wiley.com/doi/10.1111/jnc.15310

40. Pina MM, Pati D, Neira S, Taxier LR, Stanhope CM, Mahoney AA, et al. Insula Dynorphin and Kappa Opioid Receptor Systems Regulate Alcohol Drinking in a Sex-Specific Manner in Mice. J Neurosci [Internet]. 2023 Jul 12 [cited 2023 Sep 21];43(28):5158–71. Available from: https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.0406-22.2023

41. Jaramillo AA, Randall PA, Stewart S, Fortino B, Van Voorhies K, Besheer J. Functional role for cortical-striatal circuitry in modulating alcohol self-administration. Neuropharmacology. 2018 Mar 1;130:42–53.

42. Jaramillo AA, Agan VE, Makhijani VH, Pedroza S, McElligott ZA, Besheer J. Functional role for suppression of the insular–striatal circuit in modulating interoceptive effects of alcohol. Addict Biol [Internet]. 2018 Sep [cited 2024 Sep 12];23(5):1020–31. Available from: https://onlinelibrary.wiley.com/doi/10.1111/adb.12551

43. Jaramillo AA, Van Voorhies K, Randall PA, Besheer J. Silencing the insular-striatal circuit decreases alcohol self-administration and increases sensitivity to alcohol. Behav Brain Res [Internet]. 2018 Aug [cited 2024 Sep 12];348:74–81. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0166432818302250

44. Myrick H, Anton RF, Li X, Henderson S, Drobes D, Voronin K, et al. Differential Brain Activity in Alcoholics and Social Drinkers to Alcohol Cues: Relationship to Craving. Neuropsychopharmacology. 2004 Feb;29(2):393–402.

45. Ihssen N, Cox WM, Wiggett A, Fadardi JS, Linden DEJ. Differentiating Heavy from Light Drinkers by Neural Responses to Visual Alcohol Cues and Other Motivational Stimuli. Cereb Cortex. 2011 Jun 1;21(6):1408–15.

46. Chung T, Clark DB. Insula White Matter Volume Linked to Binge Drinking Frequency Through Enhancement Motives in Treated Adolescents. Alcohol Clin Exp Res. 2014 Jul;38(7):1932–40.

47. Peng Y, Gillis-Smith S, Jin H, Tränkner D, Ryba NJP, Zuker CS. Sweet and bitter taste in the brain of awake behaving animals. Nature. 2015 Nov 26;527(7579):512–5.

48. Méndez-Ruette M, Linsambarth S, Moraga-Amaro R, Quintana-Donoso D, Méndez L, Tamburini G, et al. The Role of the Rodent Insula in Anxiety. Front Physiol. 2019 Mar 29;10:330.

49. Nicolas C, Ju A, Wu Y, Eldirdiri H, Delcasso S, Couderc Y, et al. Linking emotional valence and anxiety in a mouse insula-amygdala circuit. Nat Commun. 2023 Aug 21;14(1):5073.

50. Grodin EN, Cortes CR, Spagnolo PA, Momenan R. Structural deficits in salience network regions are associated with increased impulsivity and compulsivity in alcohol dependence. Drug Alcohol Depend. 2017 Oct;179:100–8.

51. Senatorov VV, Damadzic R, Mann CL, Schwandt ML, George DT, Hommer DW, et al. Reduced anterior insula, enlarged amygdala in alcoholism and associated depleted von Economo neurons. Brain. 2015 Jan;138(1):69–79.

52. Makris N, Oscar-Berman M, Jaffin SK, Hodge SM, Kennedy DN, Caviness VS, et al. Decreased Volume of the Brain Reward System in Alcoholism. Biol Psychiatry. 2008 Aug;64(3):192–202.

53. Thompson BL, Maleki N, Kelly JF, Sy KTL, Oscar-Berman M. Brain, behavioral, affective, and sex correlates of recovery from alcohol use disorders. Alcohol Clin Exp Res. 2021 Aug;45(8):1578–95.

54. Halcomb ME, Chumin EJ, Goñi J, Dzemidzic M, Yoder KK. Aberrations of anterior insular cortex functional connectivity in nontreatment-seeking alcoholics. Psychiatry Res Neuroimaging. 2019 Feb;284:21–8.

55. Manuweera T, Kisner MA, Almira E, Momenan R. Alcohol use disorder-associated structural and functional characteristics of the insula. J Neurosci Res. 2022;100(11):2077–89.

56. Galandra C, Basso G, Manera M, Crespi C, Giorgi I, Vittadini G, et al. Salience network structural integrity predicts executive impairment in alcohol use disorders. Sci Rep. 2018 Sep 27;8(1):14481.

57. Haaranen M, Schäfer A, Järvi V, Hyytiä P. Chemogenetic Stimulation and Silencing of the Insula, Amygdala, Nucleus Accumbens, and Their Connections Differentially Modulate Alcohol Drinking in Rats. Front Behav Neurosci. 2020 Nov 4;14:580849.

58. Haaranen M, Scuppa G, Tambalo S, Järvi V, Bertozzi SM, Armirotti A, et al. Anterior insula stimulation suppresses appetitive behavior while inducing forebrain activation in alcohol-preferring rats. Transl Psychiatry. 2020 Dec;10(1):150.

59. Pushparaj A, Le Foll B. Involvement of the caudal granular insular cortex in alcohol self-administration in rats. Behav Brain Res. 2015 Oct 15;293:203–7.

60. ÖZ F, ACER N, KATAYIFÇI N, AYTAÇ G, KARAALİ K, SİNDEL M. The role of lateralisation and sex on insular cortex: 3D volumetric analysis. Turk J Med Sci [Internet]. 2021 Jun 28 [cited 2025 Aug 1];51(3):1240–8. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8283486/

61. Haggerty DL, Munoz B, Pennington T, Viana Di Prisco G, Grecco GG, Atwood BK. The role of anterior insular cortex inputs to dorsolateral striatum in binge alcohol drinking. eLife. 2022 Sep 13;11:e77411.

62. Marino RAM, Girven KS, Figueiredo A, Navarrete J, Doty C, Sparta DR. Binge ethanol drinking associated with sex-dependent plasticity of neurons in the insula that project to the bed nucleus of the stria terminalis. Neuropharmacology. 2021 Sep 15;196:108695.

63. Zhang ZM, Chen S, Liang YZ. Baseline correction using adaptive iteratively reweighted penalized least squares. The Analyst. 2010;135(5):1138.

64. Nicolas C, Russell TI, Pierce AF, Maldera S, Holley A, You ZB, et al. Incubation of Cocaine Craving After Intermittent-Access Self-administration: Sex Differences and Estrous Cycle. Biol Psychiatry. 2019 Jun;85(11):915–24.

65. Nicolas C, Zlebnik NE, Farokhnia M, Leggio L, Ikemoto S, Shaham Y. Sex Differences in Opioid and Psychostimulant Craving and Relapse: A Critical Review. Nader M, editor. Pharmacol Rev. 2022 Jan;74(1):119–40.

66. Martins de Carvalho L, Chen H, Sutter M, Lasek AW. Sexually dimorphic role for insular perineuronal nets in aversion-resistant alcohol consumption. Front Psychiatry. 2023 Feb 28;14:1122423.

67. Sneddon EA, Ramsey OR, Thomas A, Radke AK. Increased Responding for Alcohol and Resistance to Aversion in Female Mice. Alcohol Clin Exp Res. 2020 Jul;44(7):1400–9.

68. Robinson DL, Brunner LJ, Gonzales RA. Effect of Gender and Estrous Cycle on the Pharmacokinetics of Ethanol in the Rat Brain. Alcohol Clin Exp Res [Internet]. 2002 Feb [cited 2025 Aug 1];26(2):165–72. Available from: https://onlinelibrary.wiley.com/doi/10.1111/j.1530-0277.2002.tb02521.x

69. Crippens D, White ML, George MA, Jaworski JN, Brunner LJ, Lancaster FE, et al. Gender Differences in Blood Levels, But Not Brain Levels, of Ethanol in Rats. Alcohol Clin Exp Res [Internet]. 1999 Mar [cited 2025 Aug 1];23(3):414–20. Available from: https://onlinelibrary.wiley.com/doi/10.1111/j.1530-0277.1999.tb04131.x

70. Kishimoto R, Ogishi Y, Ueda M, Matsusaki M, Amako K, Goda K, et al. Gender-related differences in mouse hepatic ethanol metabolism. J Nutr Sci Vitaminol (Tokyo). 2002 Jun;48(3):216–24.

71. Chen H, Lasek AW. Perineuronal nets in the insula regulate aversion-resistant alcohol drinking. Addict Biol. 2020 Nov;25(6):e12821.

72. Starski P, Morningstar MD, Katner SN, Frasier RM, De Oliveira Sergio T, Wean S, et al. Neural Activity in the Anterior Insula at Drinking Onset and Licking Relates to Compulsion-Like Alcohol Consumption. J Neurosci [Internet]. 2024 Feb 28 [cited 2025 Jul 31];44(9):e1490232023. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10904088/

73. Zhao Z, Covelo A, Couderc Y, Mitra A, Varilh M, Wu Y, et al. Cannabinoids regulate an insula circuit controlling water intake. Curr Biol [Internet]. 2024 May 6 [cited 2025 Aug 1];34(9):1918-1929.e5. Available from: https://www.cell.com/current-biology/abstract/S0960-9822(24)00393-2

74. De Oliveira Sergio T, Lei K, Kwok C, Ghotra S, Wegner SA, Walsh M, et al. The role of anterior insula–brainstem projections and alpha-1 noradrenergic receptors for compulsion-like and alcohol-only drinking. Neuropsychopharmacology. 2021 Oct;46(11):1918–26.

75. Seif T, Chang SJ, Simms JA, Gibb SL, Dadgar J, Chen BT, et al. Cortical activation of accumbens hyperpolarization-active NMDARs mediates aversion-resistant alcohol intake. Nat Neurosci. 2013 Aug;16(8):1094–100.

76. Wu Y, Chen C, Chen M, Qian K, Lv X, Wang H, et al. The anterior insular cortex unilaterally controls feeding in response to aversive visceral stimuli in mice. Nat Commun. 2020 Jan 31;11(1):640.

77. Hashimoto K, Spector AC. Extensive Lesions in the Gustatory Cortex in the Rat Do Not Disrupt the Retention of a Presurgically Conditioned Taste Aversion and Do Not Impair Unconditioned Concentration-Dependent Licking of Sucrose and Quinine. Chem Senses. 2014 Jan;39(1):57–71.

78. Bales MB, Schier LA, Blonde GD, Spector AC. Extensive Gustatory Cortex Lesions Significantly Impair Taste Sensitivity to KCl and Quinine but Not to Sucrose in Rats. Simon SA, editor. PLOS ONE. 2015 Nov 23;10(11):e0143419.

79. Wang L, Gillis-Smith S, Peng Y, Zhang J, Chen X, Salzman CD, et al. The coding of valence and identity in the mammalian taste system. Nature. 2018 Jun;558(7708):127–31.

80. Staszko SM, Boughter JD, Fletcher ML. The impact of familiarity on cortical taste coding. Curr Biol CB [Internet]. 2022 Nov 21 [cited 2024 Mar 8];32(22):4914-4924.e4. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9691541/

81. King MS. Distribution of Fos-immunoreactive neurons in the gustatory cortex elicited by intra-oral infusion of taste solutions in conscious rats. Brain Res [Internet]. 2018 Mar [cited 2024 Mar 8];1683:67–77. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0006899318300325

82. Naqvi NH, Bechara A. The insula and drug addiction: an interoceptive view of pleasure, urges, and decision-making. Brain Struct Funct. 2010 Jun;214(5–6):435–50.

83. Brasser SM, Castro N, Feretic B. Alcohol Sensory Processing and its Relevance for Ingestion. Physiol Behav [Internet]. 2015 Sep 1 [cited 2025 Aug 1];148:65–70. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4388769/

84. Mukherjee A, Paladino M, McSain S, Gilles-Thomas E, Lichte D, Camadine R, et al. Escalation of alcohol intake is associated with regionally decreased insular cortex activity but not associated with changes in taste quality [Internet]. 2023 [cited 2025 Aug 1]. Available from: http://biorxiv.org/lookup/doi/10.1101/2022.10.06.511140

85. Min DK, Tuor UI, Chelikani PK. Gastric distention induced functional magnetic resonance signal changes in the rodent brain. Neuroscience [Internet]. 2011 Apr [cited 2025 Aug 1];179:151–8. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0306452211001035

86. Roelofs TJM, Luijendijk MCM, van der Toorn A, Camps G, Smeets PAM, Dijkhuizen RM, et al. Good taste or gut feeling? A new method in rats shows oro-sensory stimulation and gastric distention generate distinct and overlapping brain activation patterns. Int J Eat Disord. 2021 Jul;54(7):1116–26.

87. (Bud) Craig A. Interoception: the sense of the physiological condition of the body. Curr Opin Neurobiol [Internet]. 2003 Aug [cited 2025 Aug 1];13(4):500–5. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0959438803000904

88. Chen X, Gabitto M, Peng Y, Ryba NJP, Zuker CS. A Gustotopic Map of Taste Qualities in the Mammalian Brain. Science. 2011 Sep 2;333(6047):1262–6.

89. Iezzi D, Cáceres-Rodríguez A, Strauss B, Chavis P, Manzoni OJ. Sexual differences in neuronal and synaptic properties across subregions of the mouse insular cortex. BioRxiv Prepr Serv Biol. 2023 Oct 20;2023.10.18.562844.

90. Billaux P, Maurage P, Cabé N, Laniepce A, Segobin S, Pitel A. Insular volumetry in severe alcohol use disorder and Korsakoff's syndrome through an anatomical parcellation: Let us go back to basics. Addict Biol [Internet]. 2023 Oct [cited 2025 Aug 1];28(10):e13324. Available from: https://onlinelibrary.wiley.com/doi/10.1111/adb.13324

91. Li L, Yu H, Liu Y, Meng Y jing, Li X jing, Zhang C, et al. Lower regional grey matter in alcohol use disorders: evidence from a voxel-based meta-analysis. BMC Psychiatry [Internet]. 2021 May 11 [cited 2025 Aug 1];21:247. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8111920/

92. Gehrlach DA, Dolensek N, Klein AS, Roy Chowdhury R, Matthys A, Junghänel M, et al. Aversive state processing in the posterior insular cortex. Nat Neurosci [Internet]. 2019 Sep [cited 2024 Jan 24];22(9):1424–37. Available from: https://www.nature.com/articles/s41593-019-0469-1

93. Arnold ME, Butts AN, Erlenbach TR, Amico KN, Schank JR. Sex differences in neuronal activation during aversion-resistant alcohol consumption. Alcohol Clin Exp Res. 2023;47(2):240–50.

94. Gomez JL, Bonaventura J, Lesniak W, Mathews WB, Sysa-Shah P, Rodriguez LA, et al. Chemogenetics revealed: DREADD occupancy and activation via converted clozapine. Science. 2017 Aug 4;357(6350):503–7.

95. Manvich DF, Webster KA, Foster SL, Farrell MS, Ritchie JC, Porter JH, et al. The DREADD agonist clozapine N-oxide (CNO) is reverse-metabolized to clozapine and produces clozapine-like interoceptive stimulus effects in rats and mice. Sci Rep. 2018 Mar 1;8(1):3840.

96. Gogolla N. The insular cortex. Curr Biol CB. 2017 Jun 19;27(12):R580–6.

97. Tunstall BJ, Kirson D, Zallar LJ, McConnell SA, Vendruscolo JCM, Ho CP, et al. Oxytocin blocks enhanced motivation for alcohol in alcohol dependence and blocks alcohol effects on GABAergic transmission in the central amygdala. PLoS Biol. 2019 Apr 16;17(4):e2006421.

98. de Guglielmo G, Crawford E, Kim S, Vendruscolo LF, Hope BT, Brennan M, et al. Recruitment of a Neuronal Ensemble in the Central Nucleus of the Amygdala Is Required for Alcohol Dependence. J Neurosci. 2016 Sep 7;36(36):9446–53.

99. Agoglia AE, Zhu M, Quadir SG, Bluitt MN, Douglass E, Hanback T, et al. Sex-specific plasticity in CRF regulation of inhibitory control in central amygdala CRF1 neurons after chronic voluntary alcohol drinking. Addict Biol. 2022 Jan;27(1):e13067.

100. Schiff HC, Bouhuis AL, Yu K, Penzo MA, Li H, He M, et al. An Insula–Central Amygdala Circuit for Guiding Tastant-Reinforced Choice Behavior. J Neurosci. 2018 Feb 7;38(6):1418–29.

101. Girven KS, Aroni S, Navarrete J, Marino RAM, McKeon PN, Cheer JF, et al. Glutamatergic input from the insula to the ventral bed nucleus of the stria terminalis controls reward-related behavior. Addict Biol. 2021 May;26(3):e12961.