1 Title:

Evolution of hind limb morphology of Titanosauriformes (Dinosauria, Sauropoda) via 3D Geometric Morphometrics reveals wide-gauge posture as an exaptation for gigantism.

By ADRIÁN PÁRAMO[1,2*], P. MOCHO[2,3,4,5], F. ESCASO[2], F. ORTEGA[2].

1. Centro de Interpretación Paleontológica de La Rioja, Calle Mayor 10, Igea, Spain; e-mails: paramoblazquez@gmail.com.

2. Grupo de Biología Evolutiva, Facultad de Ciencias, Universidad Nacional de Educación a Distancia (UNED), Avda. Esparta s/n, 28232. Las Rozas. Madrid, Spain; e-mails: f.escaso@ccia.uned.es, fortega@ccia.uned.es.

3. Instituto Dom Luiz, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal; e-mails: pdmocho@fc.ul.pt

4. Departamento de Geologia, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal

5 Dinosaur Institute, Natural History Museum of Los Angeles County, 900 Exposition Blvd, 90007 Los Angeles, CA, USA.

*Corresponding author.

Abstract

The sauropod hind limb is the main support that allowed their gigantic body masses and a wide range of dynamic stability adaptations. It is closely related with the position of the centre of masses of their multi-toned barrel-shaped bodies, and experiences one of the most noticeable posture changes along macronarian evolution. Deeply-branched macronaria achieve increasingly arched hind limbs in what it is known as wide-gauge posture. However, it is not clear if this evolutionary trend is related to the evolutionary cascade toward gigantism despite some titanosaurian were the largest terrestrial vertebrates ever-existed. We tested through means of 3D geometric morphometrics for evolutionary changes in hind limb morphology along Macronaria phylogenetic tree. The macronarian hind limb does become progressively more arched toward deeply-branched Saltasauridae. However, there is morphological convergence between different subclades. Wide-gauge posture does not correlate with changes in body size deeper in the macronarian evolutionary tree, acting as exaptation to gigantism. Despite some titanosaurian subclades becoming some of the largest vertebrates, there is a phyletic body size decrease in Macronaria.

Main Text

1. Introduction

Sauropod dinosaurs evolved a distinct body plan that allowed them to reach some of the largest body masses of terrestrial vertebrate (e.g., Bonaparte and Coria, 1993; Carballido et al., 2017; Lacovara et al., 2014) by several orders of magnitude compared to other vertebrates, including other megaherbivore dinosaurs. Sauropods were dominant herbivorous dinosaurs throughout much of the Mesozoic (Sander, 2013; Sander et al., 2011). Several morphological features can be related to the evolutionary cascade that allowed the acquisition of their colossal sizes, like the vertebral pneumaticity related to avian-like air sacs, high metabolic rates, cranial morphology, feeding mechanisms (among several others, see Sander 2013). There are also several appendicular features related with the development of columnar limbs (Bates et al., 2016; Lefebvre et al., 2022; Salgado et al., 1997; Sander, 2013; Ullmann et al., 2017). The mechanical stability of the columnar limbs allowed them to support their multi-toned body masses.

Previous works based on sauropod anatomical description, systematics, traditional morphometrics, biomechanics and GMM suggest that there is an important evolutionary trend in the appendicular skeleton toward the acquisition of a stable limb posture known as wide-gauge (Klinkhamer et al., 2019; Lefebvre et al., 2022; Salgado et al., 1997; Sander, 2013; Ullmann et al., 2017; Upchurch et al., 2004; Voegele et al., 2021; Wilson and Carrano, 1999; Wilson, 2002). This feature appears along deeply-branched Neosauropoda which exhibit a progressively arched limb posture (Salgado et al., 1997; Upchurch et al., 2004; Voegele et al., 2021; Wilson and Carrano, 1999; Wilson, 2002). The wide stance may have enhanced lateral stability during locomotion allowing them to adapt to a range of new topographic and ecological niches (Mannion and Upchurch, 2010; Ullmann et al., 2017). However, the widening of the titanosauriformes body and the acquisition of the wide-gauge stance is still poorly understood. Although the arched limbs and wider postures appeared in the largest sauropods among Titanosauriformes (Bates et al., 2016; Henderson, 2006; Ullmann et al., 2017; Wilson and Carrano, 1999) these features were acquired independently between small and large titanosauriforms of different subclades (Henderson 2006; Bates et al. 016). Limb posture may have been related with achieving greater ranges of ecological niches though biomechanical stability rather than body size increases alone (Bates et al., 2016; Henderson, 2006; Ullmann et al., 2017). The analysis of the hind limb through 2D and 3D geometric morphometrics (GMM) allow us to analyze complex morphological changes along the titanosauriformes phylogeny (e.g., Lefebvre et al., 2022, Ullmann et al., 2017). Here we will use 3D-digitized and reconstructed titanosauriformes hind limbs (Table 1, see also Fig. 1) to test whether there is a relationship between wider hind limb posture (as the hind limbs are the main weight support of sauropod body, even after wide-gauge posture acquisition; see Bates et al., 2016) and potential morphological convergence between different titanosauriformes subclades regardless of their body size (as proxied by hind limb size). The analysis presented here can be seen as an expansion of Lefebvre et al. (2022) study on sauropod limb evolution as we analyse a sample comprised mostly of Late Cretaceous Lithostrotian sauropods (their study included a broad diversity of sauropodomorph taxa including several titanosaurs).

2. Results

87   2.1. Titanosauriformes Morphospace Occupation
88

A Principal Component Analysis was performed to generate an occupation morphospace,
obtaining a total of 16 shape PCs (after Anderson's χ test). The first six PCs accounted for
78.3% of the cumulative morphological variation (Table 2). The non-parametric Kruskal-Wallis
test shows that no single shape variable reports significant differences among sauropod
subclades (Table 3 and 4). Here we comment on the results of the first three shape PCs (>50%
of the cumulative variance) due to space limitations, but a full description and visualization of
the complete PCA and phylomorphospace projections can be found in Appendix S3.

PC1 (summarizing 29.92% of the total variance; Fig. 2a-b) is associated with characters
describing the orientation and compression of the femoral shaft, the length of the distal femoral
condyles, the orientation of the fourth trochanter, and the length and width of the zeugopod
bones (Fig. 2c). Non-titanosaurian macronarians occupy negative values, whereas non-
lithostrotian titanosaurs are distributed between weakly negative (e.g., *Bonatitan*) and positive
PC1 values (Fig. 4b). Lithostrotian titanosaurs occupy a wide range of values, with
Aeolosaurus, the specimens of Colossosauria and *Lirainosaurus* cluster at negative PC1 values
(however, *Ampelosaurus* occupy negative PC1 values approaching zero; Fig. 2b). Most of the
non-saltasaurid, non-colossosaur, non-lirainosauridae and non-aeolosaurini lithostrotians are
broadly distributed in the morphospace between negative and positive values, with *Bonitasaura*
occupies the farthest negative PC1 values and overlapping with Colossosauria and
Lirainosaurinae (Fig. 2b). The highest positive PC1 values are occupied by saltasaurids (i.e.,
Saltasaurus and *Neuquensaurus*) (Fig. 2b). There is a trend from non-titanosaur macronarians at
negative values to the titanosaur node at weakly negative values (Fig. 2b). In this PC1, the
Titanosauria node splits near zero score, with *Lohuecotitan* occupies weakly positive PC1
values near zero score [in recent phylogenetic analyses, *Lohuecotitan* has been recovered as a
member of Lithostrotia (Díez Díaz et al., 2020; Navarro et al., 2022)], but other deeply nested
titanosaurs occupy positive scores in PC1 (Fig. 2b). The trend toward positive values follows
with several other deeply nested lithostrotian, however both Colossosauria and the two members
of Liranosaurinae fall into negative PC1 values (Lirainosaurinae was not recovered as a
monophyletic group in our current topology; Fig. 2b). There are no significant differences
between the different sauropod subclades in this PC (Appendix S4).

PC2 (summarizing 14.83% of the variance; Fig. 3a-b) is associated with characters describing
the orientation of the femoral shaft; the length and orientation of the femur proximal end; the
length, width and orientation of the distal femoral condyles; the orientation of the tibial shaft;
the length and orientation of the cnemial crest; the length, width and orientation of the tibial
distal end; the orientation of the tibial ascending process; the orientation of the fibular shaft and
fibular anterior crest (Fig. 3c). Both the early branching macronarian *Oceanotitan* -a possible
member of Somphospondyli (Mocho et al., 2019) - and the euhelopodid somphospondyli
Euhelopus occupy positive and negative values near zero score, whereas the early-diverging
somphospondylan *Ligabuesaurus* is distributed at slightly more positive PC2 values than
Oceanotitan (Fig. 3b). However, there is no clear pattern among the titanosaur subclade
morphospace (Fig. 3a) or the phylomorphospace (Fig. 3b) among progressively deeply
branching Titanosauriformes. The Saltasaurids and *Aeolosaurus* occupy positive PC2 scores
(Fig. 3b) showing some overlap with Colossosauria at weakly positive values of PC2 (Fig. 3b).

Lirainosaurines and colossosaurians are distributed at weakly negative to positive values in PC2
(Fig. 3b). However, non-saltasaurid, non-aeolosaurini, non-colossosaur, non-lithostrotian
titanosaurs are broadly distributed along the morphospace, most of them between negative
values (i.e. *Dreadnoughtus*) and positive PC2 values (i.e *Jainosaurus*, *Antarctosaurus*) (Fig. 3a-
b). It is remarkable the hind limb occupation of the titanosaur *Dreadnoughtus* at strongly
negative PC2 values (Fig. 3a-b). The Mann-Whitney's U test found no significant differences in
the PC2 in any of the pairwise comparisons (Appendix S4).

Finally, PC3 (summarizing 11.61% of the variance; Fig. 3a-b, 4a-b) is related to characters
describing the orientation of the femur shaft and head; the length of the femoral lateral bulge;
the location and orientation of the fourth trochanter; the orientation of the distal femoral
condyles; the length and orientation of the tibial cnemial crest; the length of the tibial distal end;
the length of the tibial ascending process; and the length and morphology of the fibula (Fig. 4c).
Oceanotitan and *Euhelopus* were plotted toward progressively more positive values (Fig. 4a)
and the node was estimated at slightly more negative values of PC3 than *Euhelopus* (Fig. 4b).
Ligabuesaurus occupies positive values of PC3 closer to zero than early branching
titanosauriformes (Fig. 4a). In the phylomorphospace, there is a trend from positive values to
negative values of PC3 (Fig. 4b) from more basally branching titanosauriforms to more deeply
nested ones. Non-lithostrotian titanosaurs occupy negative PC3 values but the non-saltasaurid,
non-lirainosaurid and non-colossosaurian lithostrotians trends towards positive PC3 values (Fig.
4a-b), whereas Lirainosauridae, Colossosauria and Saltasauridae occupy negative values of PC3
(Fig. 4a-b). Colossosauria, Saltasauridae and *Lirainosaurus*, the most deeply-branched
representative of Lirainosauridae according to our phylogenetic hypothesis, were plotted toward
increasingly more negative PC3values (Fig. 4a-b). *Magyarosaurus* is the only lithostrotian
occupying high positive values, clearly separated from all other sauropods (Fig. 4a-b). The pair-
wise Mann-Whitney's U test found no significant differences between sauropod subclades along
this PC (Appendix S4).

2.2. Size distribution
There is a trend in hind limb centroid size distribution from large, non-titanosaurian
macronarians in the Early Cretaceous (e.g., *Euhelopus*) to small, deeply nested lithostrotian
titanosaurs in the Late Cretaceous (Fig. 5). This trend coincides with the morphospace
occupation recovers by PC1 (early-branching and larger non-titanosaurian macronarians among
negative PC1 values and progressively more deeply nested and smaller titanosaurs to positive
PC1 values (Fig. 5). The smallest lithostrotians are concentrated at deeply branching nodes with
the Ibero-Armorican lirainosaurines, *Bonatitan reigi*, *Magyarosaurus* spp. and members of
Saltasaurinae (Fig. 5).

However, the RMA models found no significant correlation between the shape variables and the
log-transformed centroid size (Table 6, see Fig. 6 and Appendix S3). The PC1 model (r^2 =
0.105, p-value= 0.204; Fig. 6a) found negative allometry but no significant correlation and the
percentage of variance explained by the hind limb size differences was small. Almost all of the
RMA found a negative relationship, except for several sub-sampled RMA models like the PC2
against log-transformed centroid size for the sample of lithostrotian titanosaurs only (Fig. 6b).
None of the RMA models for the sub-samples found significant correlation (see Appendix S3).

2.3. Phylogenetic trends
Pagel's lambda (λ) estimation shows a significative phylogenetic signal in log-transformed hind
limb centroid size (λ = 0.982), and PC1 (λ = 0.715), PC3 (λ = 0.760), PC5 (λ = 0.778) and PC6

(λ = 0.697) suggesting a significant trend in the evolution of Titanosauriformes (Table 7). We
estimated ancestral characters using log-transformed hind limb centroid size and those shape
variables that exhibit significant signal over the evolution of Titanosauriformes and tested for a
directionality or trend (Fig. 7). The resulting tests recover significant trends toward a decrease in
hind limb size across all Titanosauriformes subclades, positive PC1 values up to
Somphospondyli Titanosauriformes, and negative PC3 values across all Titanosauriformes
subclades (electronic Appendix S4, table S4.8, S4.9, S4.10).

3. Discussion
190

3.1. Titanosauriformes hind limb morphological convergence
Analysis of the shape variables extracted by PCA on the procrustes coordinates of the sauropod
taxa exhibits a large overlapping between the different sauropod subclades (e.g., Fig. 2-4) across
all the resulting shape PCs. Both the non-parametric tests on the hind limb shape variables and
the size, and the phylogenetic ANOVA accounting for the time-calibrated supertree topology
suggest the lack of enough and significant morphological differences between the different
sauropod subclades studied in this analysis. Based on the lack of significant phylogenetic
differences and the presence of morphological similarities, the evolutionary pattern of the
titanosaurian hind limb may be explained by convergent evolution, consistent with previous
analyses (Lefebvre et al., 2022; Páramo, 2020; Páramo et al., 2020; Ullmann et al., 2017).

Considering the analysed sample, the acquisition of a wide-gauge locomotion results as the
main source of morphological variability in titanosaur sauropods (Fig. 2, 5) having a
significative phylogenetic signal. The trend toward a more arched limb posture persists in the
more deeply nested titanosaurs. *Oceanotitan dantasi*, a possible representative of
Somphospondyli Macronaria, exhibits a columnar hind limb characterized by having no
deflection of the femoral head with respect to the tibial condyle (though some titanosauriformes
exhibits this feature, see Royo-Torres, 2009), and a straight and long, lateromedially-narrow
zeugopod with few anterior rotations of the fibula (Fig. 2). Whereas the more deeply branching
titanosauriforms exhibit the typical titanosaurian hind limb configuration having a more arched
posture with lateromedially more robust femora, increased medial or proximo-medial deflection
of the femoral head Distally, the hind limb exhibits a slight rotation of the femoral distal ends,
and increasingly lateromedially robust zeugopods. The robust zeugopod elements being the only
resemblance to *O. dantasi* as our outgroup (e.g., Fig. 2). The highest positive PC1 values
correlate with several hyper-robust taxa of different titanosaur subclades that exhibit
lateromedially and anteroposteriorly wide stylopods and zeugopods. In these taxa, the zeugopod
bones are extremely shorter proximodistally, extremely arched with a predominance of tibiae
characterized by short cnemial crests but somewhat rotated interlocking with anteriorly
deflected and robust fibulae as in the saltasaurine *Saltasaurus loricatus*, the lithostrotian
Magyarosaurus spp. and the non-lithostrotian titanosaur *Dreadnoughtus schrani* (based on the
super-tree: Fig. 2,5). The acquisition of this particular morphology is correlated to development
of gigantism within Titanosauria (Carrano, 2005; Lefebvre et al., 2022; Ullmann et al., 2017), at
least when analysing early branching members of Lithostrotia. However, specimens occupying
higher PC1 values exhibit the most hyper-robust and arched hind limb including representatives
of the smallest lithostrotians and the largest non-lithostrotian titanosaur studied (Fig. 5). The
analyses of evolutionary trends presented here reveal that the trend toward titanosauriform
gigantism is shifting toward adaptation to dwarfism in some lithostrotian titanosaurs like
Magyarosaurus and *Neuquensaurus* (see discussion on hind limb size variability below). In
general, the results obtained here confirm the previously proposed trend toward the acquisition

229 of robust and arched hind limbs in Titanosauria (Fig. 7). Nevertheless, once the hind limb
230 mechanical stability is acquired the increasingly arched and robust morphologies established
231 within Titanosauria cannot be fully related to an increase in body mass (Fig. 6a) and are better
232 explained as convergence between different subclades (Figs. 2 and 6, see evolutionary trend
233 breakdown in Table 7). Saltasaurine lithostrotians are characterized by this type of extreme
234 morphology, with hyper-robust limb bones. Even large saltasaurid sauropods like
235 *Opisthocoelicaudia skarzyinski* exhibit this type of hind limb (Borsuk-Bialynicka, 1977).
236 However, this morphology is not exclusive of saltasaurids, thereby other titanosaurs exhibit the
237 homoplastic hyper-robust and arched hind limb (i.e., *Dreadnoughtus* and *Magyarosaurus*). It is
238 probable that this morphological trend was hard-coded in the sauropod bauplan and, after the
239 stable posture was acquired, was still present but relatable to a significant variability of
240 biomechanical adaptations (Voegele et al., 2021) as our analyses suggest. Despite
241 biomechanical diversity associated with hind limb morphology is still unclear, several studies
242 point out that morphological differences in the fore limb elongation in sauropod may be related
243 to different feeding niche capabilities (Bates et al., 2016; Upchurch et al., 2021; Vidal et al.,
244 2020; Voegele et al., 2020), including discussion on possible bipedal/tripodal rearing abilities
245 that are much more developed, particularly in sauropods with this kind of hind limb (Upchurch
246 et al., 2021). Interestingly, *Dreadnoughtus* exhibits this hyper-robust hind limb with subequal
247 autopodial lengths and a wedged sacrum (Vidal et al., 2020; Voegele et al., 2021), an anteriorly
248 placed body Center of Masses (CoM; Bates et al., 2016) and possibly high browsing feeding
249 capabilities (e.g., Vidal et al., 2020). While the small saltasaurids in this study exhibit short
250 necks, stout bodies and a posteriorly placed CoM (Bates et al., 2016) the pneumaticity of some
251 saltasaurid tails (Upchurch et al., 2021; Zurriaguz and Cerda, 2017) may suggest the possibility
252 of a more anterior location of the CoM. It is possible that the acquisition of a hyper-robust and
253 arched hind limb morphology is related with an improved skeletal support in high browsing and
254 extremely large sauropods like *Dreadnoughtus*, which in combination with their size and neck
255 dorsiflexion capabilities allows them to feed in a high niche stratification, independent of the
256 acquisition of rearing capabilities.

257 Although PC2 does not provide phylogenetic signal over the titanosaur evolution, it reveals
258 significant differences among smaller titanosaurian taxa with robust hind limbs and especially
259 hyper-robust zeugopods (Fig. 3) like *Magyarosaurus* and the saltasaurines *Neuquensaurus* and
260 *Saltasaurus* that can be related with different roles in the necessary mechanical stability for
261 rearing capabilities. The titanosaur *Dreadnoughtus* exhibits a slightly arched hind limb posture
262 with a robust zeugopod having a medially deflected cnemial crest of the tibia and a sigmoidal
263 fibula that is slightly posteriorly deflected. In this taxon, the fibula, although sigmoidal,
264 articulates in a mostly straight anatomical position with the particularly robust tibia. On the
265 other hand, *Magyarosaurus*, *Neuquensaurus* and *Saltasaurus* occupy increasingly positive PC2
266 values (Fig. 3). *Magyarosaurus* exhibit a short zeugopod in which the tibia is slightly laterally
267 rotated, the cnemial crest is laterally projected and the fibula is slightly sigmoidal having its
268 proximal third with an anterior projection and a medially deflected anterolateral crest.
269 Saltasaurines exhibit an extreme condition characterized by short and robust zeugopods.
270 Members of this clade show laterally rotated tibiae with posteriorly deflected distal ends, and
271 sigmoidal fibulae that articulate in an oblique position with the anteriorly projected proximal
272 third and anteriorly placed lateral crest. Also, this anatomical fibular configuration produces a
273 more anterior displacement of the distal attachment of the *M. iliofibularis* and a probable
274 distribution of the stress, which previous myological studies suggest that there is no evident
275 body size/phylogenetic pattern (Voegele et al., 2021). The differences among robust hind limb
276 titanosaurian morphologies may suggest that its convergence is due to other sources of variation
277 (i.e., different biomechanical adaptations). It is also important to note that all the lithostrotian
278 titanosaurs that exhibit a slightly rotated fibula, whether sigmoid or not, and extremely arched

hind limb morphology independent of its size, exhibit femora with reduced fibular epicondyles
(after Otero et al., 2020).

Previous studies also pointed to the distinct morphology of the members of Colossosauria
sampled in this analysis, especially in its extremely elongate and gracile zeugopod (González
Riga et al., 2018; Páramo et al., 2020). Here we found a distinctive plotting for the hind limb of
Muyelensaurus and *Mendozasaurus*, often overlapping with *Aeolosaurus,* as in PC1 and PC3
morphospaces (Figs. 2-4). It is especially relevant that these specimens occupy distinctive PC1
and PC3 scores, which exhibit a signal over titanosaur evolution (corresponding to a specific
lineage within Colossosauria). In addition, the phylomorphospaces indicate a shift away from
the main early-branching lithostrotian morphospace toward a straighter, elongated hind limb
convergent with non-titanosaurian macronarians (Fig 2a-b, Fig. 4a-b). In the case of
Aeolosaurus. the phylomorphospace produces a long branched and a wide morphospace
occupation among titanosaurs (Figs. 2-4) as the titanosaurian node occupies an entirely new area
of the morphospace. Representatives of this group exhibit extremely different morphologies,
with *Aeolosaurus* and *Antarctosaurus* often much more similar to the early branching non-
titanosaurian titanosauriformes than to other titanosaurs (e.g., Fig. 2b, 4b). Colossosauria
includes several of the largest titanosaurs ever known (Carballido et al., 2017; González Riga et
al., 2019) as well as some of the smallest lithostrotian titanosaurs of South America that are
placed within Rinconsauria, which have been recently recovered as an early branching lineage
of Colossosauria (Carballido et al., 2017; González Riga et al., 2019). Additionally, some recent
phylogenetic hypotheses recovered the aeolosaurini lithostrotians as deeply branching
rincosaurs as well (Carballido et al., 2017; Silva Junior et al., 2022). Whether the appendicular
skeleton of Aeolosaurini and Colossosauria exhibits morphological convergence between these
two different lithostrotian subclades or they represent the same morphological trend within the
same lineage of specialization is still unknown. This depends on whether the latter phylogenetic
hypotheses including Aeolosaurini within Colossosauria are accepted as valid. It is important to
note that despite the small to medium size of *Muyelensaurus* and *Aeolosaurus*, *Mendonzasaurus*
can be considered as large sauropod and exhibit morphological differences to other large
sauropods from our sample (i.e., the titanosaurs *Dreadnoughtus* and *Jainosaurus* or the non-
titanosaurian somphospondylan *Ligabuesaurus*), similar to previous analyses of separate
specimens (Páramo et al., 2020). *Antarctosaurus* seems to be the only large titanosaur that
shares a similar morphospace with the members of Colossosauria analysed here (i.e., PC1-PC3,
Figs. 2-4). The inclusion of hind limbs elements of other members of this clade (i.e.,
Patagotitan, *Epachthosaurus*) in future analyses might shed light about this morphological
pattern, as the zeugopod elements of several of these taxa resemble those of other non-
colossosaur titanosaurs (e.g., *Dreadnoughtus* and *Patagotitan*; see also *Argentinosaurus* in
Páramo et al. 2020). Also, *Aeolosaurus* exhibits a different morphology than other post-early
Coniacian lithostrotians titanosaurs producing a widening of the occupied morphospace (Fig.
7b-c).

Despite these small differences in the hind limb morphology in key features that may be related
to the giant titanosaurian body size, our analyses indicate that there is a great deal of
morphological convergence between the different titanosaurian subclades (e.g., Table 2-3).
Also, no significant differences were found in any of the shape variables, either in the pairwise
subclade comparisons (Appendix S4), or in the shape PCs that indicate slightly less variance
than expected by evolution under Brownian motion (i.e., shape PC1, PC3 in this text, Table 5;
shape PC5-6 in Appendix S4).

3.2. Hind limb size evolution
Our results with the current time-calibrated supertree topology indicate that there is a trend in
the evolution of titanosaurs towards the acquisition of smaller sizes (Fig. 5 and 7, Tables 5-6).
In the lights of our current sample, once many of the hind limb features relatable to the
acquisition of wide-gauge posture are acquired, there is a phylogenetical trend, close to pure
Brownian motion model, toward an overall drecreasing size (log-transformed hind limb centroid
size $\lambda = 0.982$), which is consistent with previous results that indicated that lithostrotians (or
even all macronarian sauropods) may not follow the Cope's rule (Carrano, 2006; de Souza and
Santucci, 2014). However, this could be due to our current lithostrotian sample as some of the
Saltasaurinae or closely related sister taxa (i.e., *Opisthocoelicauda* and *Alamosaurus*, which are
usually considered members of a more inclusive clade, Saltasauridae) are also large
lithostrotians, especially *Alamosaurus*. However, it is important to note that we also found
several key traits in our shape variables (most importantly summarized in PC1 and PC3, 41.53%
of the total sample variance between them; Table 7) that are usually related to the acquisition of
gigantism and that exhibit significant signal about titanosaur evolution. When we tested for a
significant correlation between these traits and the log-transformed hind limb centroid size, no
significant correlation was found between any of the shape variables and size (Table 4, Fig. 6).

Within Titanosauria, the hind limb arched morphology (increasing wide-gauge posture) does
not correspond to the significant trend toward a size decreasing observed in this sauropod clade
(see Fig. 7, Table 6-7). Large titanosaurs show very different hind limb morphologies. Some
lithostrotians exhibit plesiomorphic columnar, slightly arched hind limbs with elongated or even
gracile zeugopod elements (*Antarctosaurus*, *Aeolosaurus* or the extremely slender hind limb of
the colossosaurs *Medonzasaurus* and *Muyelensaurus*) (Fig. 2), whereas other large titanosaurs
(i.e. *Dreadnoughtus)* exhibit an extremely robust hind limb with slightly reduced zeugopods
that may be expected of the typical trend toward the acquisition of an arched morphology in
Titanosauriformes (Fig. 2a). In this context, it is remarkable that most of the lithostrotians that
exhibit extremely arched hind limbs, robust elements and reduced zeugopod elements are
increasingly smaller, like *Saltasaurus* or *Magyarosaurus* (Fig. 2, see discussion on
morphological convergence above). Our test found that the trend toward this type of hind limb
breaks toward titanosaurian sauropods, with more variable morphologies, large overlapping of
morphospaces whereas the hind limb morphology decreases (Fig. 7, Table 6-7). Most
importantly, PC3 exhibit some of the morphological features that are classically related to wide-
gauge arching morphology and our analyses found a significant trend toward increasingly
arched hind limbs (Table 8). However, the post-Cenomanian lithostrotians exhibit a large
morphological variability, with several taxa exhibiting plesiomorphic columnar femora (i.e.,
Ampelosaurus, *Aeolosaurus*, *Magyarosaurus*) and slightly straight zeugopodial elements, with
anteriorly expanded tibial proximal ends, and straight non-rotated fibula expanded to anterior
view (Fig. 4,7; see discussion on morphological convergence above). Among these
lithostrotians, *Ampelosaurus* and *Aeolosaurus* are medium-sized sauropods, whereas
Magyarosaurus exhibits both plesiomorphic hind limb morphology and the smallest size.
Similarly, the large size variability in the post-early Coniacian cannot be related with the
morphological features that are classically related with the acquisition of the sauropod
gigantism (Fig. 7) as discussed before. Herein, titanosaurs that produce large PC1-PC3 changes
at this peak are both moderately-to-large in size inducing a displacement of the morphospace: (i)
toward the plesiomorphic hind limb morphology, plotted toward negative PC1 and positive PC3
values (i.e., *Aeolosaurus*) or (ii) toward a morphology intermediate between the main PC1-PC3
robust arched typical titanosaurian hind limb morphology (i.e., *Jainosaurus*) (Fig. 7).

Here it is important to point out that when comparing our results with other lithostrotian
titanosaurs not included in the current analysis, similar morphological variability not relatable to
hind limb size increases (and therefore body size) is observed. The arched morphology with

extremely robust zeugopodial elements exhibited in saltasaurid sauropods is also found in the
dwarf non-lithostrotian titanosaur *Diamantinasaurus* (Poropat et al., 2021, 2015). Its tibia shares
similarities with those of *Dreadnougthus* (anteroposteriorly and lateromedially wide proximal
end, extremely short and laterally-projected cnemial crest and lateromedially expanded distal
end) (Poropat et al., 2015; Ullmann and Lacovara, 2016) whereas the fibula is similar to the
slightly straight fibula of *Magyarosaurus* having a proximally anterior deflection (Poropat et al.,
2015; APB direct observation on *Magyarosaurus* sp. specimens). Despite slight differences in
the proportion of the zeugopodial elements the hind limb morphology of *Diamantinasaurus* is
similar to those of other small taxa. However, *Rapetosaurus* exhibits a completely different hind
limb configuration that is more similar to that observed in *Lirainosaurus*. Thereby, whereas the
fibula is slightly straight having an anteriorly expanded proximal third (Curry Rogers, 2009) as
in *Magyarosaurus* and *Diamantinasaurus*, both the tibia and fibula are extremely lateromedially
compressed as in *Lirainosaurus* or *Muyelensaurus* (APB pers. obs.). Our results indicate that
small titanosaurs exhibit similar morphological variability.

Among large titanosaurs, *Elaltitan* exhibits a (virtually restored) robust femur, a slightly robust
and plesiomorphic straight tibia but with a posteriorly rotated distal end, and an extremely
sigmoidal and anteriorly projected fibula like those of members of Saltasaurinae but slightly
lateromedially narrow compared to the latter (Páramo 2020; Páramo et al., 2020). Despite some
differences, other robust large lithostrotians (i.e., *Dreadnoughtus*) exhibit similar hind limb
morphology. However, other large lithostrotians as *Argentinosaurus* exhibits a plesiomorphic
fibula (Páramo 2020; Páramo et al., 2020). The hind limb of *Patagotitan* exhibits a less arched
posture, with a slightly straighter femur whereas the fibula is robust, sigmoidal and has an
anterior expansion of the proximal third (Otero et al., 2020). The stylopodium is not as robust
and arched as in *D. schrani* but it may still exhibit the slightly plesiomorphic titanosaurian
straight hind limb, which is a shift toward robust morphology in the earlier branching
Colossosauria as *Mendozasaurus neguyelap* (Figs. 2 and 4) but still lacking the extreme medial
deflection of the femoral head seen in our PC1 positive and PC3 negative values.
Petrobrasaurus and *Narambuenatitan* are both large titanosaurs that exhibit similar medial
deflection of the femoral head to *Dreadnougthus*, and in the case of *Narambuenatitan*, even
more so than *Saltasaurus* and *Neuquensaurus* (Páramo et al., 2020). However, the tibia of
Petrobrasaurus is extremely lateromedially narrow as in *Mendozasaurus* (Páramo et al., 2020;
APB pers. obs.) instead of the typical robust tibiae of the extremely arched hind limb found in
our results (see Fig. 2). Only *Uberabatitan* exhibits a clear morphology like the hind limb
morphology found in *Dreadnougthus* and our positive PC1 scores. Another large titanosaur that
exhibits a lateromedially narrow proximal tibial end is *Ruyangosaurus* (Lu et al., 2009). Despite
it its fragmentary, the femur is straight (with a rounded shaft) and has a tibia with lateromedially
narrow proximal end similar to that observed in *Ligabuesaurus* and the members of
Colossosauria (which are closely positioned in our analyses) (Figs. 2 and 4). Other more deeply
branching lithostrotian that exhibit a hind limb that closely resemble the early-branching
Colossosauria or exhibit a plesiomorphic tibia as in *Abditosaurus* (Vila et al., 2022). In this
taxon, the femur is anteroposteriorly-narrow with highly eccentric shaft, like other Ibero-
Armorican lithostrotians (e.g., *Ampelosaurus*). *Abditosaurus* also preserves an extremely
lateromedially narrow tibial proximal end (Vila et al., 2022). Only the fibula is sigmoid and
anteriorly projected, but lateromedially narrow with an expanded anteromedially crest, which is
larger anteroposteriorly than in *Lohuecotitan*, *Lirainosaurus* (Vila et al., 2022) and probably
like those of *Magyarosaurus* being the single character that resembles our PC1 results.

Moreover, we must consider that our age estimations for the nodes of our supertree are
extremely conservative and are based on the topology after the reduced tips of our sample (see
Table 1). Titanosaur sauropods appeared unambiguously during the Early Cretaceous (D'Emic,
2012; Mannion et al., 2019b; Poropat et al., 2016) as well as lithostrotian titanosaurs appeared

early after Valanginian-Hauterivian (Mannion et al., 2019b; Poropat et al., 2016). Considering
several recent phylogenetic hypotheses, the colossosaurian node may have branched in the early
Albian (Gorscak et al., 2023; Sallam et al., 2018; Vila et al., 2022). Saltasauridae have also
often been estimated at the transit between the Early Cretaceous and Late Cretaceous, with
Jiangshanosaurus considered as an early-branching saltasaurid (Poropat et al., 2016). However,
the recent redescription of *Jiangshanosaurus* material has shed light on its phylogenetic
affinities and it may be an euhelopodid or at least not as a deeply branched titanosaur (Mannion
et al., 2019a). Saltasaurid lithostrotians are nevertheless traced back to the late Albian in recent
studies (Sallam et al. 2018; Vila et al., 2022). The trends observed in our study may be
accentuated with still a general body size decreasing among lithostrotian sauropods (Fig 9a).
However, a series of convergences in the appendicular skeleton seem to evolve independently of
body size (e.g., Fig. 7b), (e.g., the acquisition of the plesiomorphic columnar titanosaur hind
limb among medium to large lithostrotians). Some recent phylogenetic hypothesis shows
uncertain phylogenetic affinities for some of the sauropods studied, particularly for some of the
lithostrotian taxa. This is the case for the Ophistocoelicaudiinae affinities that have been
suggested for some members of Lirainosauridae and *Lohuecotitan*, which is also proposed a
subclade within Saltasauridae (Vila et al., 2022) as Opisthocoelicaudiinae. *Atsinganosaurus* an
Ibero-Armorican lithostrotian was recently recovered member of Lirainosaurinae (Díez Díaz et
al., 2020, 2018) or as a lognkosaurian colossosaurs (Gorscak and O'Connor, 2019; Vila et al.,
2022). *Atsinganosaurus* is a small lithostrotian with a hind limb morphology similar to that of
Lirainosurus (Díez Díaz et al., 2018). The extremely gracile limbs of *Atsinganosaurus* resemble
those of the small Rinconsauria lithostrotian (González Riga et al., 2019; this study) or even the
large *Mendozasaurus*. However, its affinities to Colossosauria will still indicate a convergence
between small Opisthocoelicaudiinae and lognkosaurian colossosaurs according to the
phylogenetic hypothesis of Vila et al. (2022). This phylogenetic hypothesis still indicates that
the large-sized titanosaurs *Dreadnoughtus* and *Alamosaurus* (robust medially-bevelled femur
and sigmoid fibula; see Lehman and Coulson, 2002; Wick and Lehman, 2014) exhibit a
morphologically convergence in the hind limb with the small saltasaurines as *Neuquensaurus*
and *Saltasaurus*.

It seems that these morphological similarities are due to other biomechanical aspects after the
acquisition of the arched hind limb, as well as to other morphological features related to the
wide-gauge posture of the appendicular skeleton. It is possible that minor differences that do not
show significant phylogenetic signal and recover by other shape-PCs (Appendix S4), are key
features for other adaptations that are also important such as the trade-off between speed and
rearing stability, which may have shaped limb morphology in the Titanosauriformes,
particularly in the zeugopods (e.g., Upchurch et al., 2021).

3.3. Caveats of this study
Our analyses include a wide range of titanosaurs from most of the proposed subclades.
However, many hind limb elements are fragmentary and required virtual restoration according
to Páramo et al., (2020, 2022). Traditional studies propose excluding incomplete specimens
from the sample. However, in this case, several of our taxa from an already small sample (n =
17) could not be included in this study because they do not have complete specimens to
calculate the mean shape for each hind limb element type (e.g., *Mendozasaurus neguyelap* with
several tibia and fibula specimens, *Oceanotitan dantasi* with only the fragmentary elements of
its left hind limb). The exclusion of potentially informative areas or taxa may hinder
palaeobiological studies (Brown et al., 2012) and landmarks estimation may be a more
informative procedure (Arbour and Brown, 2014; Brown et al., 2012). Also, it may be
interesting to include several taxa that have been examined in previous studies (e.g., Páramo

2020), but many of these taxa lack one or more hind limb element types. In this case, we did not
choose to estimate the entire morphology of an element type, because we lack the necessary and
more powerful tools to do so, such as partial least squares estimation methods (Torres-Tamayo
et al., 2020)(e.g., Torres-Tamayo *et al.* 2020).

Our reconstructions of the titanosaur hind limbs can also bias our study. The lack of the
astragalus in most of the titanosaurs studied (whether due to the lack of available 3D-scanned
specimens or the lack of a preserved astragalus) hinder the estimation of the position of the
zeugopodial elements. In this study, it is used the most conservative assumption in the overall
position of the distal part of the hind limb as the fibula could be positioned even more distally,
without reaching proximally the femur and interlocked with the tibia in several specimens (e.g.,
Neuquensaurus australis, specimen MCS-5-25/26, APB. pers. obs.). This is especially relevant
for the robust and arched hind limb with extremely robust zeugopodia, as *Dreadnoughtus*
schrani, *Neuquensaurus* spp., and *Saltasaurus loricatus*, but see also *Uberabatitan riberoi*
(Salgado and Carvalho, 2008).

The sample is also small and we did not opt to include several advanced statistical hypothesis
tests like phylogenetic convergence (Stayton, 2015) because we did not meet the requirements.
Instead, we choose a conservative set of tests to discuss the true morphological convergence
over titanosaurian hind limb evolution with a mix of phylogenetic and non-phylogenetic
methods. We chose Butler & Gowsami (2008) change frequencies analysis because, although its
statistical properties are less well known than those of independent contrast analyses, it is less
sensitive to topological imprecision and somewhat independent of branch length differences
(Butler and Goswami, 2008; de Souza and Santucci, 2014). Following this reasoning, we also
opted not to estimate independent phylogenetic contrasts that may be sensitive to large
differences in branch length after our lithostrotian-biased sample to test for trait correlation
between shape-PCs against log-transformed centroid size contrary to previous studies (Bates et
al., 2016). Instead, we opted for the traditional use of test for correlation between tree tips
(specimens) shape-PCs against log-transformed centroid sizes via RMA and without
incorporating the phylogenetic tree topology of our current time-calibrated supertree.

4. Conclusions

Our results suggest that the main features related to the acquisition of an arched hind limb
posture (presence of lateromedially wide femora with robust zeugopods) are typical of more
deeply nested titanosaurs such as saltasaurines (Carrano, 2005; Lefebvre et al., 2022; Sander et
al., 2011; Ullmann et al., 2017; Vila et al., 2022; Wilson and Carrano, 1999)(Wilson & Carrano
1999; Carrano 2005; Sander *et al.* 2011; Ullmann *et al.* 2017; Lefebvre *et al.* 2022; Vila *et al.*
2022) does exhibit significant phylogenetic signal about titanosaur evolution (Table 5, Fig. 7).
The arched morphology usually related with wide-gauge posture is an exaptation initially
related to increasing body size. However, once fully acquired in Titanosauria, these features are
no longer related to increasing body size, as increases in the arched posture and the development
of hyper-robust zeugopods are features shared by several of both the smallest and largest taxa
from different titanosaurian lineages (Fig. 6, Table 4). Also, there is an evolutionary trend
toward decreasing titanosaurian hind limb size (and body size; Fig. 7), based on the size of the
hind limb centroid. This trend is congruent with previous studies focused on the size evolution
of the Titanosauriformes (de Souza and Santucci 2014), despite an increase in arched
morphology and robustness of the hind limb in deeply nested titanosaur subclades.

The lack of correlation between several of these morphological traits with body size may also be
related to wide morphological variation within Lithostrotia (Figs. 2-4). It can be noted that,
along with a trend toward increasingly arched hind limb and reduced zeugopods with

increasingly rotated fibulae, there is also a large morphological convergence between the different titanosaurian subclades, as indicated by both our non-phylogenetic and phylogenetic analyses (Table 2-3). Many of the morphological changes without significant signal on titanosaur evolution is related to the morphology and anatomical position of zeugopod bones. Large morphological convergences unrelated to body size may be a response to different morphofunctional adaptations. Our results show several differences in the morphology of zeugopod elements without signal on the titanosaurian evolution, especially those regarding with the fibula (rotation, sigmoidal morphology, anterior or posterior displacement of the proximal end that affects the anteroposterior position of the lateral bulge). These morphological differences are also related with its relative position and articulation with the tibia and correlate with the expression of femoral features like the relative development of the posterior epicondyle of the fibular condyle. These features translate in changes in the length and morphology of the hind limb distal musculature that previous studies indicate that exhibit no clear evolutionary trend within titanosaurs. Also, our results show that specimens placed in different subclades share similar morphologies across the shape-PCs that exhibit no phylogenetical signal on titanosaur evolution. The observed changes in the zeugopod morphology may be related with different morphofunctional and ecomorphological adaptations and the convergences in hind limb morphology of titanosaurs may explain biomechanical similarities. This may explain the differences observed between small and medium to large titanosaurs with either arched or columnar hind limb (i.e. PC1 similarities) across those shape-PCs that lack significant signal on titanosaur evolution. However, to test the hypothesis of differences in biomechanical adaptation (i.e., movement speed or differences in feeding niche specialization) further analysis with additional parts of the skeleton must be included.

5. Acknowledgments

This work is funded by Ministry of Science and Innovation of Spain project PID2019-111488RB-I00 and the Junta de Comunidades de Castilla-La Mancha projects (SBPLY/19/180801/000044 and SBPLY/21/180801/000045). Access to fossil vertebrate collections in Argentina was made possible by Spanish Ministry of Economy and Competitiveness grant EEBB-I-16-11875. We want to thank BP. Kear and Uppsala's University Natural History Museum staff for facilitating access to their collections. Access to Uppsala's University Museum of Natural History paleovertebrate collections was possible thanks to ERG-2020 EAVP research grant. Digitizing of macronarian sauropod *Oceanotitan dantasi* holotype was possible thanks to ERG-2020 EAVP research grant. Author PM was funded by FCT/MCTES for one CEEC Individual contract (CEECIND/00726/2017) and a Haaga Family Postdoctoral Fellow (2016-2018) in the NHMLA (CA, USA). We want to thank S. Chapman and Natural History Museum of London (UK) staff for facilitating access to their collections. Access to NHM (UK) was made possible thanks to the European Union Council Synthesys program GB-TAF-6153. We want to also thank the alumni of the Faculty of Arts from the UCM (Madrid, Spain) and preparation staff for providing the sample for this study. We also want to thank many others for access to specimens under their care, including S. Langreo from the Museo de Paleontología de Castilla-La Mancha (Cuenca, Spain); J.A. Ramírez de la Peciña and C. Corral from the MCNA (Vitoria, Spain); R. Coria from the MCF (Plaza Huincul, Argentina); L. Filippi from the MRS (Rincón de los Sauces, Argentina); M. Reguero from the MLP (La Plata, Argentina); A. Otero from CONICET (La Plata, Argentina); S. Devincenzi from the IANIGLA (Mendoza, Argentina); and B. González Riga from the UNCUYO (Mendoza, Argentina); C. Muñoz from the MPCA (Cipolletti, Argentina); I. Cerda from the CONICET (Cipolletti, Argentina); P. Ortiz from the PVL (Tucumán, Argentina); A.G. Kramarz and M. Ezcurra from the MACN (Buenos Aires, Argentina) and J. Le Loeuff from the Musée des Dinosauries d'Espéraza (Espéraza, France).

6. Bibliography

Arbour JH, Brown CM. 2014. Incomplete specimens in geometric morphometric analyses.
Methods Ecol Evol **5**:16–26. doi:10.1111/2041-210X.12128
Bates KT, Mannion PD, Falkingham PL, Brusatte SL, Hutchinson JR, Otero A, Sellers WI,
Sullivan C, Stevens KA, Allen V. 2016. Temporal and phylogenetic evolution of the
sauropod dinosaur body plan. *Royal Society Open Science* **3**:150636.
doi:10.1098/rsos.150636
Bininda-Emonds ORP. 2004. The evolution of supertrees. *Trends in ecology & evolution*
19:315–22. doi:10.1016/j.tree.2004.03.015
Bonaparte JF, Coria RA. 1993. Un nuevo y gigantesco saurópodo Titanosaurio de la Formación
Río Limay (Albiano-Cenomaniano) de la provincia del Neuquén, Argentina.
Ameghiniana **30**:271–282.
Bonnan MF. 2007. Linear and Geometric Morphometric Analysis of Long Bone Scaling
Patterns in Jurassic Neosauropod Dinosaurs: Their Functional and Paleobiological
Implications. *The Anatomical Record: Advances in Integrative Anatomy and
Evolutionary Biology* **290**:1089–1111. doi:10.1002/ar.20578
Bonnan MF, Wilhite RD, Masters SL, Yates AM, Gardner CK, Aguiar A. 2013. What Lies
Beneath: Sub-Articular Long Bone Shape Scaling in Eutherian Mammals and
Saurischian Dinosaurs Suggests Different Locomotor Adaptations for Gigantism. *PLoS
ONE* **8**:e75216. doi:10.1371/journal.pone.0075216
Borsuk-Bialynicka M. 1977. A New Camarasaurid Sauropod *Opisthocoelicaudia skarzynskii*
gen. n., sp. n. from the Upper Cretaceous of Mongolia. *Paleontologia Polonica* **37**:5–
64.
Brown CM, Arbour JH, Jackson DA. 2012. Testing of the effect of missing data estimation and
distribution in morphometric multivariate data analyses. *Syst Biol* **61**:941–954.
doi:10.1093/sysbio/sys047
Butler RJ, Goswami A. 2008. Body size evolution in Mesozoic birds: little evidence for Cope's
rule. *Journal of Evolutionary Biology* **21**:1673–1682. doi:10.1111/j.1420-
9101.2008.01594.x
Carballido JL, Pol D, Otero A, Cerda IA, Salgado JL, Garrido AC, Ramezani J, Cúneo NR,
Krause JM. 2017. A new giant titanosaur sheds light on body mass evolution among
sauropod dinosaurs. *Proceedings of the Royal Society B: Biological Sciences*
284:20171219. doi:10.1098/rspb.2017.1219
Carrano MT. 2006. Body-Size Evolution in the Dinosauria In: Carrano MT, Gaudin TJ, Blob
RW, Wible J, editors. Amniote Paleobiology: Perspectives on the Evolution of
Mammals, Birds, and Reptiles. pp. 225–268.
Carrano MT. 2005. The Evolution of Sauropod Locomotion In: Curry Rogers KA, Wilson JA,
editors. The Sauropods: Evolution and Paleobiology. University of California Press. pp.
229–250.
Curry Rogers K. 2009. The postcranial osteology of *Rapetosaurus krausei* (Sauropoda:
Titanosauria) from the Late Cretaceous of Madagascar. *Journal of Vertebrate
Paleontology* **29**:1046–1086. doi:10.1671/039.029.0432
de Souza LM, Santucci RM. 2014. Body size evolution in Titanosauriformes (Sauropoda,
Macronaria). *Journal of Evolutionary Biology* **27**:2001–2012. doi:10.1111/jeb.12456
D'Emic MD. 2012. The early evolution of titanosauriform sauropod dinosaurs. *Zoological
Journal of the Linnean Society* **166**:624–671. doi:10.1111/j.1096-3642.2012.00853.x
Díez Díaz V, Garcia G, Pereda Suberbiola X, Jentgen-Ceschino B, Stein K, Godefroit P,
Valentin X. 2020. A new titanosaur (Dinosauria: Sauropoda) from the Upper
Cretaceous of Velaux-La-Bastide Neuve (southern France). *Historical Biology* **00**:1–20.
doi:10.1080/08912963.2020.1841184

Díez Díaz V, Garcia G, Pereda-Suberbiola X, Jentgen-Ceschino B, Stein K, Godefroit P,
Valentin X. 2018. The titanosaurian dinosaur *Atsinganosaurus velauciensis*
(Sauropoda) from the Upper Cretaceous of southern France: New material,
phylogenetic affinities, and palaeobiogeographical implications. *Cretaceous Research*
91:429–456. doi:10.1016/j.cretres.2018.06.015
González Riga BJ, Lammana MC, Otero A, Leonardo D. Ortiz D, Kellner AW, Ibiricu LM.
2019. An overview of the appendicular skeletal anatomy of South American
titanosaurian sauropods, with definition of a newly recognized clade. *Anais da*
Academia Brasileira de Ciências **91**:e20180374. doi:https://doi.org/10.1590/0001-
3765201920180374
González Riga BJ, Mannion PD, Poropat SF, Ortiz David LD, Coria JP. 2018. Osteology of the
Late Cretaceous Argentinean sauropod dinosaur *Mendozasaurus neguyelap*:
implications for basal titanosaur relationships. *Zoological Journal of the Linnean*
Society 1–46. doi:10.1093/zoolinnean/zlx103/4816851
Gorscak E, Lamanna M, Schwarz D, Díez Díaz V, Salem B, Sallam H, Wiechmann M. 2023. A
new titanosaurian (Dinosauria: Sauropoda) from the Upper Cretaceous (Campanian)
Quseir Formation of the Kharga Oasis, Egypt. *Journal of Vertebrate Paleontology*.
doi:10.1080/02724634.2023.2199810
Gorscak E, O'Connor PM. 2019. A new African Titanosaurian Sauropod Dinosaur from the
middle Cretaceous Galula Formation (Mtuka Member), Rukwa Rift Basin,
Southwestern Tanzania. *PLOS ONE* **14**:e0211412. doi:10.1371/journal.pone.0211412
Gunz P, Mitteroecker P, Bookstein FL. 2005. Semilandmarks in three dimensions In: Slice DE,
editor. Modern Morphometrics in Physical Anthropology. KLUWER ACADEMIC
PUBL. pp. 73–98.
Henderson DM. 2006. Burly gaits: centers of mass, stability, and the trackways of sauropod
dinosaurs. *Journal of Vertebrate Paleontology* **26**:907–921. doi:10.1671/0272-
4634(2006)26
Holliday CM, Ridgely RC, Sedlmayr JC, Witmer LM. 2010. Cartilaginous epiphyses in extant
archosaurs and their implications for reconstructing limb function in dinosaurs. *PloS*
one **5**:1–16. doi:10.1371/journal.pone.0013120
Klinkhamer AJ, Mallison H, Poropat SF, Sloan T, Wroe S. 2019. Comparative Three-
Dimensional Moment Arm Analysis of the Sauropod Forelimb: Implications for the
Transition to a Wide-Gauge Stance in Titanosaurs. *The Anatomical Record* **302**:794–
817. doi:10.1002/ar.23977
Lacovara KJ, Lamanna MC, Ibiricu LM, Poole JC, Schroeter ER, Ullmann PV, Voegele KK,
Boles ZM, Carter AM, Fowler EK, Egerton VM, Moyer AE, Coughenour CL, Schein
JP, Harris JD, Martínez RD, NOVAS FE. 2014. A Gigantic, Exceptionally Complete
Titanosaurian Sauropod Dinosaur from Southern Patagonia, Argentina. *Sci Rep* **4**.
doi:10.1038/srep06196
Lefebvre R, Houssaye A, Mallison H, Cornette R, Allain R. 2022. A path to gigantism: Three-
dimensional study of the sauropodomorph limb long bone shape variation in the context
of the emergence of the sauropod bauplan. *Journal of Anatomy*. doi:10.1111/joa.13646
Legendre P. 2018. lmodel2: Model II Regression.
Lehman TM, Coulson AB. 2002. A Juvenile Specimen of the Sauropod Dinosaur Alamosaurus
sanjuanensis from the Upper Cretaceous of Big Bend National Park, Texas. *Journal of*
Paleontology **76**:156–172. doi:10.1666/0022-3360(2002)076<0156:AJSOTS>2.0.CO;2
Lu JC, Xu L, Jia SH, Zhang XL, Zhang JM, Yang LL, You HL, Ji Q. 2009. A new gigantic
sauropod dinosaur from the Cretaceous of Ruyang, Henan, China. *Geological Bulletin*
of China **28**:1–10. doi:10.1111/j.1095-8312.1965.tb00944.x
Mallison H. 2011. Digitizing Methods for Paleontology: Applications, Benefits and Limitations
In: Elewa A, editor. Computational Paleontology. Berling: Springer. pp. 7–43.
Mannion PD, Upchurch P. 2010. A quantitative analysis of environmental associations in
sauropod dinosaurs. *Paleobiology* **36**:253–282. doi:10.1666/08085.1

Mannion PD, Upchurch P, Jin X, Zheng W. 2019a. New information on the Cretaceous
sauropod dinosaurs of Zhejiang Province, China: Impact on Laurasian titanosauriform
phylogeny and biogeography. *Royal Society Open Science* **6**. doi:10.1098/rsos.191057
Mannion PD, Upchurch P, Schwarz D, Wings O. 2019b. Taxonomic affinities of the putative
titanosaurs from the Late Jurassic Tendaguru Formation of Tanzania: phylogenetic and
biogeographic implications for eusauropod dinosaur evolution. *Zoological Journal of*
the Linnean Society **185**:784–909. doi:10.1093/zoolinnean/zly068
Mazzetta GV, Christiansen P, Fariña R a. 2004. Giants and Bizarres: Body Size of Some
Southern South American Cretaceous Dinosaurs. *Historical Biology* **16**:71–83.
doi:10.1080/08912960410001715132
Mocho P, Royo-Torres R, Ortega F. 2019. A new macronarian sauropod from the Upper
Jurassic   of   Portugal.   *Journal   of   Vertebrate   Paleontology*   **39**:1–23.
doi:10.1080/02724634.2019.1578782
Navarro BA, Ghilardi AM, Aureliano T, Díaz VD, Bandeira KLN, Cattaruzzi AGS, Iori FV,
Martine AM, Carvalho AB, Anelli LE, Fernandes MA, Zaher H. 2022. A new nanoid
titanosaur (Dinosauria, Sauropoda) from the Upper Cretaceous of Brazil. *Ameghiniana*
59:317–354. doi:10.5710/AMGH.25.08.2022.3477
Otero A, Carballido JL, Moreno AP. 2020. The Appendicular Osteology of *Patagotitan*
Mayorum (Dinosauria, Sauropoda). *Journal of Vertebrate Paleontology* **40**:1–19.
doi:10.1080/02724634.2020.1793158
Páramo A. 2020. Estudio del esqueleto apendicular de los titanosaurios Ibero-Armoricanos
mediante morfometría geométrica: estado del arte y perspectivas futurasI Congreso
Iberoamericano de Morfometría Geométrica.
Páramo A, Mocho P, Ortega F. 2020. Three-dimensional analysis of the titanosaurian limb
skeleton: implications for systematic analysis. *Journal of Iberian Geology* **46**:369–402.
doi:10.1007/s41513-020-00139-8
Poropat SF, Kundrát M, Mannion PD, Upchurch P, Tischler TR, Elliott DA. 2021. Second
specimen of the Late Cretaceous Australian sauropod dinosaur Diamantinasaurus
matildae provides new anatomical information on the skull and neck of early
titanosaurs.   *Zoological   Journal   of   the   Linnean   Society*   **192**:610–674.
doi:10.1093/zoolinnean/zlaa173
Poropat SF, Mannion PD, Upchurch P, Hocknull SA, Kear BP, Kundrát M, Tischler TR, Sloan
T, Sinapius GHK, Elliott JA, Elliott DA. 2016. New Australian sauropods shed light on
Cretaceous   dinosaur   palaeobiogeography.   *Scientific   Reports*   **6**:1–12.
doi:10.1038/srep34467
Poropat SF, Upchurch P, Mannion PD, Hocknull SA, Kear BP, Sloan T, Sinapius GHK, Elliott
DA. 2015. Revision of the sauropod dinosaur *Diamantinasaurus matildae* Hocknull et
al. 2009 from the mid-Cretaceous of Australia: Implications for Gondwanan
titanosauriform   dispersal.   *Gondwana   Research*   **27**:995–1033.
doi:10.1016/j.gr.2014.03.014
R Core Team. 2022. R: A Language and Environment for Statistical Computing. Vienna,
Austria: R Foundation for Statistical Computing.
Revell LJ. 2012. phytools: An R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* **3**:217–223.
Royo-Torres R. 2009. El saurópodo de Peñarroya de Tastavins. *Monograpfías Turonlenses* **6**:1–
548.
Salgado JL, Carvalho IDS. 2008. *Uberabatitan ribeiroi*, a new titanosaur from the Marília
Formation (Bauru Group, Upper Cretaceous), Minas Gerais, Brazil. *Palaeontology*
51:881–901. doi:10.1111/j.1475-4983.2008.00781.x
Salgado L, Coria RA, Calvo JO. 1997. Evolution of titanosaurid sauropods: Phytogenetic
analysis based on the postcranial evidence. *Ameghiniana* **34**:3–32.
Sallam HM, Gorscak E, O'Connor PM, El-Dawoudi IA, El-Sayed S, Saber S, Kora MA, Sertich
JJW, Seiffert ER, Lamanna MC. 2018. New Egyptian sauropod reveals Late Cretaceous
dinosaur dispersal between Europe and Africa. *Nature Ecology & Evolution* **2**:445–451.
doi:10.1038/s41559-017-0455-5

Sander PM. 2013. An Evolutionary Cascade Model for Sauropod Dinosaur Gigantism -
Overview, Update and Tests. *PLOS ONE* **8**:e78573. doi:10.1371/journal.pone.0078573
Sander PM, Christian A, Clauss M, Fechner R, Gee CT, Griebeler E-M, Gunga H-C, Hummel J,
Mallison H, Perry SF, Preuschoft H, Rauhut OWM, Remes K, Tütken T, Wings O,
Witzel U. 2011. Biology of the sauropod dinosaurs: the evolution of gigantism.
Biological Reviews **86**:117–155. doi:10.1111/j.1469-185X.2010.00137.x
Schlager S. 2017. Morpho and Rvcg – Shape Analysis in R In: Zheng G, Li S, Szekely G,
editors. Statistical Shape and Deformation Analysis. Elsevier. pp. 217–256.
doi:10.1016/B978-0-12-810493-4.00011-0
Schliep K, Potts AJ, Morrison DA, Grimm GW. 2017. Intertwining phylogenetic trees and
networks. *Methods in Ecology and Evolution* **8**:1212–1220.
Schliep KP. 2011. phangorn: phylogenetic analysis in R. *Bioinformatics* **27**:592–593.
doi:10.1093/bioinformatics/btq706
Schwarz D, Wings O, Meyer CA. 2007. Super sizing the giants: first cartilage preservation at a
sauropod dinosaur limb joint. *Journal of the Geological Society* **164**:61–65.
doi:10.1144/0016-76492006-019
Silva Junior JCG, Martinelli AG, Iori FV, Marinho TS, Hechenleitner EM, Langer MC. 2022.
Reassessment of Aeolosaurus maximus, a titanosaur dinosaur from the Late Cretaceous
of   Southeastern   Brazil.   *Historical   Biology*   **34**:403–411.
doi:10.1080/08912963.2021.1920016
Stayton C. 2015. The definition, recognition, and interpretation of convergent evolution, and
two new measures for quantifying and assessing the significance of convergence.
Evolution **69**. doi:10.1111/evo.12729
Torres-Tamayo N, Schlager S, García-Martínez D, Sanchis-Gimeno JA, Nalla S, Ogihara N,
Oishi M, Martelli S, Bastir M. 2020. Three-dimensional geometric morphometrics of
thorax-pelvis covariation and its potential for predicting the thorax morphology: A case
study on Kebara 2 Neandertal. *Journal of Human Evolution* **147**:102854.
doi:10.1016/j.jhevol.2020.102854
Ullmann PV, Bonnan MF, Lacovara KJ. 2017. Characterizing the Evolution of Wide-Gauge
Features in Stylopodial Limb Elements of Titanosauriform Sauropods via Geometric
Morphometrics. *The Anatomical Record* **300**:1618–1635. doi:10.1002/ar.23607
Ullmann PV, Lacovara KJ. 2016. Appendicular osteology of *Dreadnoughtus schrani*, a giant
titanosaurian (Sauropoda, Titanosauria) from the Upper Cretaceous of Patagonia,
Argentina.   *Journal   of   Vertebrate   Paleontology*   **36**.
doi:10.1080/02724634.2016.1225303
Upchurch P, Barrett PM, Dodson P. 2004. Sauropoda In: Weishampel DB, Dodson P, Osmolska
H, editors. The Dinosauria. Berkeley and Los Angeles: University of California Press.
pp. 259–322.
Upchurch P, Mannion PD, Xu X, Barrett PM. 2021. Re-assessment of the Late Jurassic
eusauropod dinosaur *Hudiesaurus sinojapanorum* Dong, 1997, from the Turpan Basin,
China, and the evolution of hyper-robust antebrachia in sauropods. *Journal of*
Vertebrate Paleontology **41**. doi:10.1080/02724634.2021.1994414
Vidal D, Mocho P, Aberasturi A, Sanz JL, Ortega F. 2020. High browsing skeletal adaptations
in Spinophorosaurus reveal an evolutionary innovation in sauropod dinosaurs. *Sci Rep*
10:1–10. doi:10.1038/s41598-020-63439-0
Vila B, Sellés A, Moreno-Azanza M, Razzolini NL, Gil-Delgado A, Canudo JI, Galobart À.
2022. A titanosaurian sauropod with Gondwanan affinities in the latest Cretaceous of
Europe. *Nature Ecology & Evolution*. doi:10.1038/s41559-021-01651-5
Voegele KK, Bonnan MF, Siegler S, Langel CR, Lacovara KJ. 2022. Constraining
Morphologies of Soft Tissues in Extinct Vertebrates Using Multibody Dynamic
Simulations: A Case Study on Articular Cartilage of the Sauropod Dreadnoughtus.
Frontiers in Earth Science **10**.
Voegele KK, Ullmann PV, Lamanna MC, Lacovara KJ. 2021. Myological reconstruction of the
pelvic girdle and hind limb of the giant titanosaurian sauropod dinosaur *Dreadnoughtus*
schrani. *Journal of Anatomy* **238**:576–597. doi:10.1111/joa.13334

Voegele KK, Ullmann PV, Lamanna MC, Lacovara KJ. 2020. Appendicular myological
reconstruction of the forelimb of the giant titanosaurian sauropod dinosaur
Dreadnoughtus schrani. *Journal of Anatomy* 1–22. doi:10.1111/joa.13176
Wick SL, Lehman TM. 2014. A complete titanosaur femur from West Texas with comments
regarding hindlimb posture. *Cretaceous Research* **49**:39–44.
doi:10.1016/j.cretres.2014.02.003
Wiley DF, Amenta N, Alcantara DA, Ghosh D, Kil YJ, Delson E, Harcourt-Smith W, Rohlf FJ,
St John K, Hamann B. 2005. Evolutionary morphingIEEE Visualization, 2005. VIS 05.
Presented at the IEEE Visualization, 2005. VIS 05. pp. 431–438.
doi:10.1109/VISUAL.2005.1532826
Wilson JA, Carrano MT. 1999. Titanosaurs and the origin of "wide-gauge" trackways; a
biomechanical and systematic perspective on sauropod locomotion. *Paleobiology*
25:252–267.
Wilson JeffreyA. 2002. Sauropod dinosaur phylogeny: critique and cladistic analysis.
Zoological Journal of the Linnean Society **136**:215–275. doi:10.1046/j.1096-
3642.2002.00029.x
Zurriaguz VL, Cerda IA. 2017. Caudal pneumaticity in derived titanosaurs (Dinosauria:
Sauropoda). *Cretaceous Research* **73**:14–24. doi:10.1016/j.cretres.2017.01.001

808 7. Methods

810 7.1. 3D Geometric Morphometrics

To analyse the morphology 17 macronarian sauropods hind limbs (Table 1) were 3D-digitized
using 3D Geometric Morphometrics tool-kit (3D-GMM; Gunz et al., 2005 e.g., Páramo et al.,
2020). 3D digitizing process were based on the methodology proposed in previous analyses
(i.e., Mallison, 2011; Páramo et al., 2020) and 3D-GMM analyses were conducted in R
statistical software v4.1.3 (R Core Team, 2022).

To reconstruct each taxa hind limb, we firstly virtually-restored the 3D reconstructions of the
digitized femora, tibiae and fibulae and calculated the grand mean shape for each sampled taxon
using previous datasets (Páramo 2020; Páramo et al., 2020). The grand mean specimens for
each taxon were mounted on an estimated anatomical position (accounting for the lack of
astragalus on our sample to guide the distal zeugopod articulation). A comprehensive
description of the methodology can be accessed in Appendix 2.

Each element of the hind limb is separated by an articular cap much larger than in other known
extant archosaurs (Bonnan et al., 2013; Holliday et al., 2010; Schwarz et al., 2007; Voegele et
al., 2022). The femur, and the hind limb overall, may exhibit the best correlation between
cartilage cap and bone morphology constrained by its role as support under most of the stress of
the body mass (Bonnan et,al., 2013; Voegele et al., 2022). Despite cartilaginous cap thickness is
not constant and may vary among taxa, we set up our model hind limbs with a similar constant
space between the femur and the zeugopod bones in all the sampled taxa. The anatomical
mounts include an additional space of 2% of the element length between stylopod and zeugopod
(following Voegele et al., 2022).

A total of 28 landmarks and 12 semilandmark curves were placed on the hind limb bones partly
based on previous studies (Lefebvre et al., 2022; Páramo et al., 2020; Table 2, Fig. 2, Appendix
1) using IDAV Landmark Editor software (Wiley et al., 2005) (dataset with the landmark and
semilandmark curves coordinates can be accessed in the Appendix S1. Semilandmarks were
then slid in R using package "Morpho" (Schlager, 2017)(Schlager 2017) following Gunz et al.,

(2005). To remove size differences, spatial position and orientation, the resulting landmark and
semilandmark configurations were superimposed via Generalized Procrustes Analysis (GPA)
using the "procSym" function in *Morpho* package. Morphological variance was analysed with
Principal Component Analysis (PCA; results in Table 3) saving the expected number of
Principal Components (PCs) which summarize a significant amount of variance after an
Anderson Chi's test (see Bonnan, 2007).

Evolutionary trend analyses were accomplished after the estimation of a consensus tree
topology using the MRP-supertree methodology (Bininda-Emonds, 2004) with the *phangorn*
package (Schliep et al., 2017; Schliep, 2011). For supertree construction we compiled several of
the more recent phylogenetic hypotheses that includes all the available sampled data (resulting
supertree can be accessed in Appendix 1; trimmed supertree in Fig. 2). The resulting
phylogenetic relationships were projected onto the shape-PCA to visualize the
phylomorphospace. To analyse true morphological convergences between titanosaur sub-clades
we tested for morphological differences in the hind limb skeleton using the shape variables
(PCs) without the phylogenetic relationships involved using: (i) Mann Whitney U's test and
Kruskal-Wallis non-parametric tests accounting for the uneven distribution of the group (sub-
clade) samples and (ii) phylogenetic ANOVA using the time-calibrated supertree topology with
the *phytools* R package (Revell, 2012).

7.2. Hind limb size distribution and phylogenetic signal
Sauropod body mass was proxied by hind limb centroid size collected from the GPA. Body
mass can be calculated preferably using both humeral and femoral measurements (Mazzetta et
al., 2004) or whole-body volumetric estimation (Bates et al., 2016). However, as the hind limb
is the main sauropod body mass support, it can be better used as a "conservative-minimal"
approach to its body mass, with larger hind limb corresponding to giant titanosaur taxa. We
tested for allometric relationships between the sauropod hind limb shape variables (PCs) and the
centroid size as proxy to body mass via Reduced Major Axis (RMA) regression using *lmodel2*
R package (Legendre, 2018)(Legendre 2018).

We used the time-calibrated supertree topology and our shape variables (PCs) and hind limb
centroid size to generate the Pagel's lambda (λ) with "phytools" R package and test for
phylogenetic signal of these traits. All hypotheses of statistical correlations, dissimilarity tests
and phylogenetic signal tests were accepted as significant using an alpha level of 0.05 (a
comprehensive report of the results and a copy of the R code and packages used can be accessed
in Appendix S4 and S5 respectively). Once identified those PCs that exhibit significant
phylogenetical signal, as well as the log-transformed centroid size, we estimated their ancestral
characters (ACEs) using maximum-likelihood and a simple Brownian evolutionary model
similar to the one assumed for estimation of the Pagel's lambda. We used the ACEs to observe
trends in the evolution of titanosaur hind limb size and morphology (based on our shape-PCs).
We evaluated the differences in body size (proxied by log-transformed hind limb centroid size
ACEs) between terminal taxa and internal nodes and between internal nodes of
Titanosauriformes, Somphospondyli, Titanosauria and Lithostrotia, as well as their subclades.
The sum of changes, mean change, median change, positive, negative and the total amount of
changes were evaluated for each of the above clades following Butler and Goswami (2008). We
used a $\chi2$ godness-of-fit test to evaluate whether body size increasing or decreasing, and shape-
PCs occur at the same frequency (50%-50% null hypothesis) or have a positive or negative
tendency over titanosaurian evolution.

Figure 1. Sample of several 3D reconstruction of macronarian hind limbs used in this study. (a)
Oceanotitan dantasi; (b) *Ligabuesarusu leanzai*; (c) *Lohuecotitan pandafilandi*; (d) *Saltasaurus*
loricatus. All taxa in anterior and posterior views.

Figure 2. PCA results on the GPA aligned landmark and semilandmark curves of the hind limb.
(a) PC1-PC2 biplot. (b) PC1-PC2 phylomorphospace with projected phylogenetic tree. (c)
Representation of the shape change along PC1, blue are negative scores, red are positive scores.
Percentage of variance of each PC in brackets under corresponding axis. *Ae – Aeolosaurus, Amp*
– Ampelosaurus, An – Antarctosaurus, Bo – Bonatitan, Bon – Bonitasaura, Dre –
Dreadnoughuts, Eu – Euhelopus, Ja – Jainosaurus, Li – Lirainosaurus, Lig – Ligabuesaurus,
Lo – Lohuecotitan, Ma – Magyarosaurus, Me – Mendozasaurus, Mu – Muyelensaurus, Ne –
Neuquensaurus, Sa – Saltasaurus.

Figure 3. PCA results for the GPA aligned landmark and semilandmark curves of the hind
limbs. (a) PC2-PC3 biplot. (b) PC2-PC3 phylomorphospace with projected phylogenetic tree.
(c) Representation of the shape change along PC2, blue are negative scores, red are positive
scores. Percentage of variance of each PC in brackets under corresponding axis. *Ae –*
Aeolosaurus, Amp – Ampelosaurus, An – Antarctosaurus, Bo – Bonatitan, Bon – Bonitasaura,
Dre – Dreadnoughuts, Eu – Euhelopus, Ja – Jainosaurus, Li – Lirainosaurus, Lig –
Ligabuesaurus, Lo – Lohuecotitan, Ma – Magyarosaurus, Me – Mendozasaurus, Mu –
Muyelensaurus, Ne – Neuquensaurus, Sa – Saltasaurus.

Figure 4. PCA results for the GPA aligned landmark and semilandmark curves of the hind
limbs. (a) PC1-PC3 biplot. (b) PC1-PC3 phylomorphospace with projected phylogenetic tree.
(c) Representation of the shape change along PC3, blue are negative scores, red are positive
scores. Percentage of variance of each PC in brackets under corresponding axis. *Ae –*
Aeolosaurus, Amp – Ampelosaurus, An – Antarctosaurus, Bo – Bonatitan, Bon – Bonitasaura,
Dre – Dreadnoughuts, Eu – Euhelopus, Ja – Jainosaurus, Li – Lirainosaurus, Lig –
Ligabuesaurus, Lo – Lohuecotitan, Ma – Magyarosaurus, Me – Mendozasaurus, Mu –
Muyelensaurus, Ne – Neuquensaurus, Sa – Saltasaurus.

Figure 5. Time calibrated supertree with PC1-PC2 results and hind limb centroid size for each
sauropod.

Figure 6. RMA results of the first three shape PCs against the logarithm of the hind limb
centroid size. (a) PC1 against log-Centroid size, all taxa RMA in dark red: intercept = 0.221,
slope = -0.102, r^2 = 0.105, p = 0.204; Titanosauria only partial RMA in dashed green: intercept
= 0.203, slope = -0.092, r^2 = 0.118, p = 0.229; Lithostrotia only partial RMA in dotted orange:
intercept = 0.246, slope = -0.120, r^2 = 0.319, p = 0.07; (b) PC2 against log-Centroid size, all
taxa RMA in dark red: intercept = 0.155, slope = -0.072, r^2 = 0.054, p = 0.371; Titanosauria
only partial RMA in dashed green: intercept = 0.158, slope = -0.075, r^2 = 0.117, p = 0.232;

Lithostrotia only partial RMA in dotted orange: intercept = -0.127, slope = 0.066, r^2 = 0, p =
0.952; (c) PC3 against log-Centroid size (Csize), all taxa RMA in dark red: intercept = 0.137,
slope = -0.064, r^2 = 0.055, p = 0.363; Titanosauria only partial RMA in dashed green: intercept
= 0.110, slope = -0.055, r^2 = 0.236, p = 0.078; Lithostrotia only partial RMA in dotted orange:
intercept = 0.140, slope = -0.073, r^2 = 0.313, p = 0.074. *Ae – Aeolosaurus, Amp –*
Ampelosaurus, An – Antarctosaurus, Bo – Bonatitan, Bon – Bonitasaura, Dre – Dreadnoughtus,
Eu – Euhelopus, Ja – Jainosaurus, Li – Lirainosaurus, Lig – Ligabuesaurus, Lo – Lohuecotitan,
Ma – Magyarosaurus, Me – Mendozasaurus, Mu – Muyelensaurus, Ne – Neuquensaurus, Sa –
Saltasaurus.

Figure 7. Evolution of log-transformed hind limb centroid size, shape PC1 and PC3 according
to our sample and time-calibrated supertree topology. Ba – Barremian, Be – Berriasian, Ce –
Cenomanian, Co – Conacian, H – Hauterivian, J – Jurassic, Late-K – Late Cretaceous, Ti –
Tithonian, Tu – Turonian, Va – Valanginian. my – million years from present, sc – subclade
Colossosauria + *Bonitasaura*.

TABLE CAPTIONS
944

Table 1. Specimen sample used in this study. Proximodistal hind limb length measured in mm.

Table 2. PCA results over GPA aligned coordinates. Variance explained by each shape PC.

Table 3. Kruskal-Wallis test on shape PCA variables between the most inclusive subclades
analysed.

Table 4. Phylogenetic ANOVA test on shape PCA variables between the most inclusive
subclades studied.

Table 5. RMA models of the shape PCs against log-transformed Centroid size. CI – confidence
interval.

Table 6. Estimated phylogenetic signal via Pagel's lambda. *p* value of log-likelihood ratio test
after 1000 simulations. * Indicates significant relationships for an alpha of 0.05. log-Csize –
Log-transformed hind limb centroid size.

Table 7. Results of pairwise ancestor-descendant comparisons for log-transformed centroid size
in macronarian sauropods in our time-calibrated supertree (17 terminal taxa). n = ingroup
internal nodes + terminal taxa. *= accepted as significant with alpha <0.05. º= Lithostrotia +
Antarctosaurus.

Table 8. Results of ancestor-descendant pairwise comparisons for shape PC1 in macronarian
sauropods in our time-calibrated supertree (17 terminal taxa). n = ingroup internal nodes +
terminal taxa. *= accepted as significant with alpha <0.05. º= Lithostrotia + *Antarctosaurus*.

Table 9. Results of ancestor-descendant pairwise comparisons for shape PC3 in macronarian
sauropods in our time-calibrated supertree (17 terminal taxa). n = ingroup internal nodes +
terminal taxa. *= accepted s significant with alpha <0.05. º= Lithostrotia + *Antarctosaurus*.

TABLES

Table 1. Specimen sample used in this study. Proximodistal hind limb length measured in mm.		
	Clade	Hind limb_length (mm)
Aeolosaurus	Aeolosaurini	1839
Ampelosaurus	Lirainosauridae	1411
Antarctosaurus	Titanosauria	2351
Bonatitan	Lithostrotia	963
Bonitasaura	Lithostrotia	1987
Dreadnoughtus	Titanosauria	3184
Euhelopus	Euhelopodidae	1683
Jainosaurus	Titanosauria	2188
Ligabuesaurus	Somphospondyli	2866
Lirainosaurus	Lirainosauridae	1074
Lohuecotitan	Lithostrotia	1120
Magyarosaurus	Lithostrotia	750
Mendozasaurus	Colossosauria	2334
Muyelensaurus	Colossosauria	1450
Neuquensaurus	Saltasauridae	1200
Oceanotitan	Macronaria	1892
Saltasaurus	Saltasauridae	1230

Table 2. PCA results over GPA aligned coordinates. Variance explained by each shape PC.		
	Explained variance (%)	Cumulative variance (%)
PC1	29.92	29.92
PC2	14.83	44.75
PC3	11.61	56.36
PC4	8.83	65.19
PC5	7.02	72.21
PC6	6.09	78.3
PC7	4.18	82.48
PC8	3.75	86.23
PC9	3.12	89.35
PC10	2.19	91.53
PC11	2.15	93.68
PC12	1.95	95.64
PC13	1.56	97.19

PC14	1.18	98.38
PC15	0.92	99.29
PC16	0.71	100

Table 3. Kruskal-Wallis test on shape PCA variables between the most inclusive subclades analysed.

	Chi-sq	p-value	p-adjusted
PC1	11.954	0.153	1
PC2	5.592	0.693	1
PC3	7.886	0.445	1
PC4	9.647	0.291	1
PC5	5.17	0.739	1
PC6	7.618	0.472	1
PC7	8.915	0.35	1
PC8	8.941	0.347	1
PC9	10.206	0.251	1
PC10	10.422	0.237	1
PC11	4.768	0.782	1
PC12	5.66	0.685	1
PC13	8.886	0.352	1
PC14	9.856	0.275	1
PC15	10.248	0.248	1
PC16	4.578	0.802	1

Table 4. Phylogenetic ANOVA test on shape PCA variables between the most inclusive subclades studied.

		Df	SS	MS	R^2	F	Z	Pr(>F)
PC1	~Clade	8	0.073	0.009	0.466	0.872	-0.105	0.531
	Residuals	8	0.083	0.010	0.533	-	-	-
	Total	16	0.157	-	-	-	-	-
PC2	~Clade	8	0.001	1.50E+04	0.685	0.218	0.573	0.313
	Residuals1	8	5,52E+04	6.90E+09	0.314	-	-	-
	Total1	16	0.002	-	-	-	-	-
PC3	~Clade	8	1.54E+04	1.93E+09	0.285	0.399	-0.925	0.82
	Residuals	8	3.86E+04	4,83E+08	0.714	-	-	-
	Total	16	5.40E+04	-	-	-	-	-
PC4	~Clade	8	1.49E+04	1.87E+09	0.3534	0.546	-0.496	0.658
	Residuals	8	2.73E+04	3,41E+09	0.646	-	-	-
	Total	16	4.22E+04	-	-	-	-	-
PC5	~Clade	8	3.77E+04	4,72E+09	0.556	125.362	0.121	0.452
	Residuals	8	3.01E+04	3,77E+09	0.443	-	-	-
	Total	16	6.79E+04	-	-	-	-	-
PC6	~Clade	8	3.24E+04	4,06E+09	0.590	144.158	0.590	0.264
	Residuals	8	2.25E+04	2,82E+09	0.409	-	-	-
	Total	16	5.50E+04	-	-	-	-	-
	~Clade	8	1.63E+04	2,05E+09	0.434	0.768	-0.315	0.625
	Residuals	8	2.13E+04	2,66E+09	0.565	-	-	-
	Total	16	3.76E+04	-	-	-	-	-

Table 5. RMA models of the shape PCs against log-transformed Centroid size. CI – confidence interval.

	intercept	slope	CI 2.5% Slope	CI 97.5% Slope	r^2	P
PC1	0.221	-0.102	-0.168	-0.062	0.105	0.204
PC2	0.155	-0.072	-0.12	-0.043	0.054	0.371
PC3	0.137	-0.064	-0.106	-0.038	0.055	0.363
PC4	0.12	-0.055	-0.093	-0.033	0.026	0.534
PC5	-0.107	0.049	0.03	0.082	0.079	0.275
PC6	-0.1	0.046	0.028	0.076	0.086	0.254

Table 6. Estimated phylogenetic signal via Pagel's lambda. *p* value of log-likelihood ratio test after 1000 simulations. * Indicates significant relationships for an alpha of 0.05. log-Csize – Log-transformed hind limb centroid size.

	Lambda	p
log-Csize	0.982	0.003*
PC1	0.715	0.000*
PC2	0	1
PC3	0.76	0.002*
PC4	0	1
PC5	0.778	0.031*
PC6	0.697	0.01*

Table 7. Results of pairwise ancestor-descendant comparisons for log-transformed centroid size in macronarian sauropods in our time-calibrated supertree (17 terminal taxa). n = ingroup internal nodes + terminal taxa. *= accepted as significant with alpha <0.05. º= Lithostrotia + *Antarctosaurus*.

Clade	Mean	Sum	Skew	Median	n	Positive changes	Negative changes	χ^2	p
Titanosauriformes	-0.127	-3.819	-0.223	-0.032	30	12	18	1.2	0.273
Somphospondyli	-0.366	-10.252	-0.148	-0.256	28	4	24	14.286	0.000*
Titanosauria	-0.288	-7.488	-0.088	-0.182	26	5	21	9.846	0.002*
Lithostrotiaº	-0.283	-6.22	-0.193	-0.307	22	5	17	6.545	0.011*

Table 8. Results of ancestor-descendant pairwise comparisons for shape PC1 in macronarian sauropods in our time-calibrated supertree (17 terminal taxa). n = ingroup internal nodes + terminal taxa. *= accepted as significant with alpha <0.05. º= Lithostrotia + *Antarctosaurus*.

Clade	Mean	Sum	Skew	Median	n	Positive changes	Negative changes	χ^2	p
Titanosauriformes	0.051	1.518	0.98	0.043	30	29	1	26.133	0.000*
Somphospondyli	0.01	0.53	1.15	0.012	2	20	7	6.259	0.012

9	8	8		8				*
Titanosauria	0.011	0.291	1.076	0.003	26	14	10	0.667	0.414
Lithostrotiaº	0.011	0.24	1.079	0.002	22	11	10	0.048	0.827

Table 9. Results of ancestor-descendant pairwise comparisons for shape PC3 in macronarian sauropods in our time-calibrated supertree (17 terminal taxa). n = ingroup internal nodes + terminal taxa. *= accepted s significant with alpha <0.05. º= Lithostrotia + *Antarctosaurus*.								
Clade	Mean	Sum	Skew	Median	n	Positive changes	Negative changes	χ^2	p
Titanosauriformes	-0.035	-1.047	1.697	-0.038	30	2	28	22.533	0.000*
Somphospondyli	-0.023	-0.639	2.174	-0.024	28	2	26	20.571	0.000*
Titanosauria	-0.008	-0.21	2.483	-0.009	26	3	23	15.385	0.000*
Lithostrotiaº	-0.003	-0.074	2.323	-0.004	22	4	17	8.048	0.005*

300mm

Clade
- Aeolosaurini
- Colossosauria
- Euhelopodidae
- Lirainosauridae
- Lithostrotia
- Macronaria
- Saltasauridae
- Somphospondyli
- Titanosauria

A

PC3 (11.61%)

Ma
Eu
Oc
Lig
Ae
Amp   *Bo*
Dre   *Ne*   *Sa*
Mu   *An*
Lo   *Bon*
Ja   *Me*
Li

PC2 (14.83%)

B

PC3 (11.61%)

PC2 (14.83%)

Clade

⊡	Aeolosaurini	■	Lirainosauridae	▼	Saltasauridae
▲	Colossosauria	◆	Lithostrotia	⊞	Somphospondyli
⊠	Euhelopodidae	●	Macronaria	⊕	Titanosauria

C

PC2
− ⟷ +

A

B

Clade

⊠ Aeolosaurini	■ Lirainosauridae	▼ Saltasauridae
▲ Colossosauria	◆ Lithostrotia	⊞ Somphospondyli
⊠ Euhelopodidae	● Macronaria	⊕ Titanosauria

C

PC3

− +