β-catenin inhibition disrupts the homeostasis of osteogenic/adipogenic differentiation leading to the development of glucocorticoid-induced osteonecrosis of femoral head

Chenjie Xia[1,2*], Huihui Xu[1*], Liang Fang[1], Jiali Chen[1], Wenhua Yuan[1], Danqing Fu[3], Xucheng Wang[1], Bangjian He[4], Luwei Xiao[1], Chengliang Wu[1], Peijian Tong[4], Di Chen[5†], Pinger Wang[1†], Hongting Jin[1†]

[1] Institute of Orthopadics and Traumatology, The First Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou, Zhejiang 310053, China
[2] Department of Orthopedic Surgery, Lihuili Hospital Affiliated to Ningbo University, Ningbo, Zhejiang 315041, China
[3] School of Basic Medical Sciences, Zhejiang Chinese Medical University, Hangzhou, Zhejiang 310053, China
[4] Department of Orthopedic Surgery, the First Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou, Zhejiang 310003, China
[5] Faculty of Pharmaceutical Sciences, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen 518055, China

*These authors have an equal contribution to this work.
†Corresponding author:
Di Chen
Faculty of Pharmaceutical Sciences
Shenzhen Institute of Advanced Technology
Chinese Academy of Sciences
1068, Xueyuan Road
Shenzhen 518055, China
Email: di.chen@siat.ac.cn

Ping-er Wang
Institute of Orthopaedics and Traumatology
The First Affiliated Hospital of Zhejiang Chinese Medical University
548, Binwen Road
Hangzhou, Zhejiang Province 310053, China
Email: pingerwang@zcmu.edu.cn

Hongting Jin
Institute of Orthopaedics and Traumatology
The First Affiliated Hospital of Zhejiang Chinese Medical University
548, Binwen Road
Hangzhou, Zhejiang Province 310053, China.
Email: hongtingjin@zcmu.edu.cn

Abstract

Glucocorticoid-induced osteonecrosis of the femoral head (GONFH) is a common refractory joint disease characterized by bone damage and the collapse of femoral head structure. However, the exact pathological mechanisms of GONFH remain unknown. Here, we observed abnormal osteogenesis and adipogenesis associated with decreased β-catenin in the necrotic femoral head of GONFH patients. In vivo and in vitro studies further revealed that glucocorticoid exposure disrupted osteogenic/adipogenic differentiation bone marrow stromal cells (BMSCs) by inhibiting β-catenin signaling in glucocorticoid-induced GONFH rats. Col2$^+$ lineage largely contributes to BMSCs, and was found an osteogenic commitment in the femoral head through 9 months of lineage trace. Specific deletion of *β-catenin* in Col2$^+$ cells shifted their commitment from osteoblasts to adipocytes, leading to a full spectrum of disease phenotype of GONFH in adult mice. Overall, we uncover that β-catenin inhibition disrupting the homeostasis of osteogenic/adipogenic differentiation contribute to the development of GONFH, and identify an ideal genetic modified mouse model of GONFH.

Keywords: Glucocorticoid; Osteonecrosis of the femoral head; β-catenin; Osteogenic/adipogenic differentiation; Col2$^+$ lineage

Introduction

Glucocorticoids, potent immunity regulators, are widely used in the treatment of various autoimmune diseases such as systemic lupus erythematosus (SLE), nephrotic syndrome (NS) and rheumatoid arthritis (RA)*(Yang et al., 2021)*. However, excessive usage of glucocorticoids has been reported to cause severe adverse effects, especially femoral head osteonecrosis in the joint*(Zaidi et al., 2010; Tao et al., 2017)*. As a destructive joint disease, glucocorticoid-induced osteonecrosis of the femoral head (GONFH) disables about 100,000 Chinese and more than 20,000 Americans annually*(Petek et al., 2019; Zhao et al., 2020)*, bringing a huge financial burden on the society. Currently, there is no ideal medical treatment for GONFH due to its unclear pathological mechanisms. Most patients eventually have to undergo a joint replacement surgery*(Scaglione et al., 2015)*. Therefore, in-depth study of GONFH pathogenesis is urgently needed.

Previous *ex-vivo* studies on human necrotic femoral heads revealed prominent pathological

features of GONFH including fat droplet clusters, trabecular bone loss, empty lacunae of

osteocyte at the early stage and extra subchondral bone destruction, structure collapse at the late

stage*(Qin et al., 2015; Wang et al., 2018a)*. Glucocorticoid-induced animal models are commonly

used to study GONFH pathogenesis, which mimic the early necrotic changes of GONFH but

without structure collapse*(Zheng et al., 2018)*. Several hypotheses, such as damaged blood

supply*(Sun et al., 2021; Wang et al., 2018b)*, abnormal lipid metabolism*(Lavernia et al., 1999;*

Chang et al., 1993) and bone cell apoptosis*(Chen et al., 2022; Zalavras et al., 2003)* have been

proposed to explain the occurrence and development of GONFH. However, the therapies

developed according to the guideline established based on these theories have not been successful

in prevention of GONFH progression. Clinical evidence about decreased replication and

osteogenic differentiation capacity of bone marrow mesenchymal stromal cells (BMSCs)*(Lee et*

al., 2006; Wang et al., 2008; Houdek et al., 2016) and the efficacy of BMSCs

transplantation*(Wang et al., 2019a; Tabatabaee et al., 2015; Gangji and Hauzeur, 2005)* in

GONFH patients indicate that GONFH might be a BMSCs-related disease. BMSCs contain

mesenchymal progenitor cells which give rise to osteoblasts and adipocytes, and glucocorticoids

have been shown to regulate osteogenic/adipogenic differentiation of BMSCs *in vitro(Yin et al.,*

2006; Han et al., 2019). Various treatments targeting on promoting osteogenic differentiation of

BMSCs alleviate early necrotic phenotype in glucocorticoid-induced GONFH animal

models*(Chen et al., 2020; Zhu et al., 2020; Feng et al., 2022)*. Thus, we hypothesize that

imbalanced osteogenic/adipogenic differentiation of BMSCs plays a dominant role in GONFH

pathogenesis. Recent lineage tracing and single-cell RNA sequencing studies revealed that

BMSCs are a group of heterogeneous multipotent cells*(Gao et al., 2021)*, and skeletal-derived

mesenchymal progenitor cells including collagen II (Col2)[+] lineage and Osterix (Osx)[+] lineage

compose a large proportion of BMSCs*(Ono et al., 2014; Liu et al., 2013)*. However, which

sub-population of BMSCs significantly contributes to the GONFH pathogenesis remains

unknown.

BMSCs differentiation is a complex process and is regulated by multiple signaling pathways.

Among them, canonical Wnt/β-catenin signaling functions as a switch in determining

osteogenic/adipogenic differentiation of BMSCs*(Song et al., 2012)*. When β-catenin is

accumulated in the nucleus, interacting with TCF/Lef transcription factors to activate downstream target genes including Runt-related transcription factor 2 (Runx2) and Osterix*(Yuan et al., 2016)*. On the contrary, inhibition of β-catenin induces expressions of CCAAT/enhancer binding protein alpha (C/EBPα) and peroxisome proliferator activated receptor gamma (PPAR-γ) for adipogenesis*(Takada et al., 2009; Xu et al., 2016)*. Our previous study showed a significant decrease of β-catenin in the glucocorticoid-induced GONFH rat model*(Zhang et al., 2019)*. Other groups also reported an involvement of β-catenin signaling in GONFH development*(Zhang et al., 2021; Chen et al., 2019; Zhao et al., 2021; Yu et al., 2016; Zhang et al., 2015)*.

Here, abnormal osteogenesis and adipogenesis with decreased β-catenin signaling were observed in the necrotic femoral heads of GONFH patients and glucocorticoid-induced GONFH rats. Activation of β-catenin signaling effectively alleviated the necrotic changes in GONFH rats by restoring glucocorticoid exposure-induced imbalanced osteogenic/adipogenic differentiation of BMSCs. Interestingly, specific deletion of *β-catenin* in Col2[+] cells, but not Osx[+] cells led to a full spectrum of disease phenotype of GONFH in mice even including subchondral bone destruction and femoral head collapse. Our present study provided novel molecular mechanisms of GONFH pathogenesis, and an ideal genetic modified mouse model for GONFH study.

Results

Abnormal osteogenesis and adipogenesis with decreased β-catenin in necrotic femoral head of GONFH patients

To determine the pathogenesis of GONFH, we harvested the surgical specimens from GONFH patients with femoral head necrosis (n=15) and samples from trauma patients without femoral head necrosis (n=10). Compared to the non-necrotic femoral head, we observed subchondral bone destruction (Figure 1A, black asterisk), liquefied necrotic foci (Figure 1A, black arrow), deformed and collapsed outline in the necrotic femoral head. μCT analysis further showed sparse and cracked trabeculae in the collapsed region (Figure 1B, red box) and liquefied necrotic region (Figure 1B, yellow box) compared to the corresponding regions (Figure 1B, white boxes) in the non-necrotic femoral heads, which were further confirmed by the decreased BV/TV, Tb.N and Tb.Th and increased Tb.Sp (Figure 1C). The representative images of ABH staining (Figure 1D) revealed sparse trabeculae, massive accumulated fat droplets (Figure 1D, black triangles), numerous empty lacunae of osteocytes (Figure 1D, black arrows) and extensive subchondral bone

destruction (Figure 1D, yellow arrows) in the necrotic femoral head. Histomorphological quantitative analyses showed decreased trabecular bone area (Figure 1E), increased fat droplet area (Figure 1E) and empty lacunae rate (Figure 1F) in the necrotic femoral heads. TUNEL staining revealed more apoptotic osteocytes in the necrotic femoral head (Figure 1G). We then detected the expression of osteogenic (Runx2, Osterix) and adipogenic (FABP4, PPAR-γ, CEBP/α) marker genes. The results of IHC staining showed a down-regulation of Runx2 and an up-regulation of FABP4 in the necrotic regions (Figure 1H). Western blot analysis showed increased expressions of PPAR-γ and CEBP/α and a decreased expression of Osterix in the human necrotic femoral head tissues (Figure 1I). These findings indicate an involvement of abnormal osteogenesis and adipogenesis during GONFH pathogenesis. β-catenin can regulate osteogenesis and adipogenesis downstream target genes (Runx2, Osterix, FABP4, PPAR-γ and CEBP/α) expressions(Tencerova and Kassem, 2016). The significant decrease of β-catenin in the necrotic femoral heads (Figure 1G, 1H) indicate an important role of β-catenin signaling in GONFH pathogenesis.

Inhibition of β-catenin signialing leads to abnormal osteogenesis and adipogenesis in glucocorticoid-induced GONFH rats

To further analyze the role of β-catenin in GONFH pathogenesis, a glucocorticoid-induced GONFH rat model was established by continuous MPS induction, as previously described(Zheng et al., 2018). Gross anatomy analysis showed a local melanocratic region on the femoral head surface of GONFH rats without any structure deformity or collapse (Figure 2A, black arrow). No color change was observed on the femoral head surface from the Wnt agonist 1 treated rats. μCT images and quantitative analysis of bone microstructure showed severe trabecular bone loss in the femoral head of GONFH rats, especially in the subchondral region (Figure 2A, 2B). Systemic injection of Wnt agonist 1 effectively increased bone mass (Figure 2A), and alleviated decreased BV/TV, Tb.N and Tb.Th and increased Tb.Sp in GONFH rats (Figure 2B). ABH staining and histomorphological quantitative analysis revealed that the GONFH rats presented sparse trabeculae, massive fat accumulation (Figure 2C, black arrow) and numerous empty lacunae of osteocytes (Figure 2C, black arrows) in the necrotic region of femoral heads (Figure 2C-2E). However, no subchondral bone destruction, femoral head collapse and deformity was occurred in this GONFH rat model (Figure 2C). TUNEL staining reveled increased apoptotic osteocytes in the

femoral heads of GONFH rats (Figure 2F). ALP is an osteoblast marker reflecting the capacity of

osteogenesis. FABP4 is essential for lipid formation and metabolism. The decreased β-catenin and

ALP expressions and increased FABP4 expression confirmed abnormal osteogenesis and

adipogenesis during rat GONFH development (Figure 2G, 2H), consistent with the findings in

human GONFH femoral heads. Interestingly, activation of β-catenin by systematically injecting

Wnt agonist 1 for 6 weeks attenuated the early necrotic changes (Figure 2C-2E) and osteocyte

apoptosis (Figure 2F), and restored ALP and FABP4 expressions in the femoral head of GONFH

rats (Figure 2G, 2H). Overall, these data indicate that β-catenin inhibition-induced abnormal

osteogenesis and adipogenesis contributes to GONFH pathogenesis.

β-catenin directs dexamethasone-induced osteogenic/adipogenic differentiation of rat BMSCs

We further investigated cellular mechanisms involved in the necrotic femeral head of

glucocorticoid-induced GONFH rats. Primary bone marrow cells were extracted from the

proximal femur of 4-week-old SD rats and cultivated to P3. Flow cytometry analysis identified

that they were BMSCs, with $CD90^+$, $CD29^+$, $CD45^-$ and $CD11b^-$ characteristics (Figure 3A). ALP

staining showed that the osteogenic differentiation of BMSCs were significantly inhibited after

exposing to 10^2 nM or higher concentration of Dex for 7 days (Figure 3B). Oil red O staining

showed a progressive adipogenic differentiation of BMSCs with the increased concentration of

Dex (Figure 3B). These findings indicated that increased adipogenic differentiation and decreased

osteogenic differentiation of BMSCs caused the abnormal osteogenesis and adipogenesis, finally

leading to the femoral head necrosis in GONFH rats. The decreased expressions of β-catenin,

Runx2, ALP and increased expressions of PPAR-γ, CEBP/α further revealed that Dex exposure

inhibited β-catenin signaling, and redirected BMSC differentiation from osteoblasts to adipocytes

(Figure 3C, 3D). However, this Dex-induced abnormal differentiation of BMSCs were restored by

SKL2001 through activation of β-catenin singling (Figure 3E-3G). Taking together, these findings

suggest that β-catenin inhibition redirects the direction of BMSC differentiation, contributing to

the GONFH development.

Loss-of-function of β-catenin in $Col2^+$, but not Osx^+, -expressing cells leads to a GONFH-like phenotype in femoral head

Skeletal progenitor cells is a major source of BMSCs, and both $Col2^+$ lineage and Osx^+

lineage can trans-differentiate into BMSCs *(Gao et al., 2021; Ono et al., 2014)*. To determine

which subset of BMSCs contributes more to GONFH pathogenesis, we generated *β-catenin*^{Col2ER}

mice and *β-catenin*^{OsxER} mice, in which *β-catenin* gene was specifically deleted in Col2$^+$ cells and

Osx$^+$ cells, respectively. For mapping their cell fate in the femoral head, we induced *Tomato*^{Col2ER}

mice and *Tomato*^{OsxER} mice at 2-weeks old and analyzed at the age of 1 month. The fluorescent

images showed that Col2$^+$ cells were evidently expressed in the chondrocytes of articular cartilage,

secondary ossification center and growth plate (Figure 4A, white arrowheads), osteoblasts (Figure

4A, green arrowheads), osteocytes (Figure 4A, yellow arrowheads) and stromal cells (Figure 4A,

red arrowheads) underneath the growth plate, while Osx$^+$ cells were identified only in osteoblasts,

osteocytes and stromal cells underneath the growth plate (Figure 4A). We then trace Col2$^+$ lineage

for 9 months and found that Col2$^+$ chondrocytes in secondary ossification center and growth plate

were replaced by Col2$^+$ osteoblasts, osteocytes and stromal cells with age (Figure 4B), and these

Col2$^+$ cells continuously produced osteoblasts and osteocytes in the femoral head (Figure 4B).

These findings indicated the self-renew ability and osteogenic commitment of Col2$^+$ cells. We also

induced *Tomato*^{Col2ER} mice at the age of 1 month, but identified a poor *Cre*-recombinase efficiency

24 hours and 3 months later (Figure 4C). IHC assay showed a significant decrease of β-catenin in

the femoral head of 3-month-old *β-catenin*^{Col2ER} mice and *β-catenin*^{OsxER} mice (Figure 4D, 4E),

indicating successful establishments of these *β-catenin* conditional knockout mouse models.

Then, we analyzed morphologic changes of the femoral head from 3-months old

β-catenin^{Col2ER} mice and *β-catenin*^{OsxER} mice. No change of appearance was observed on the

femoral head both in *β-catenin*^{Col2ER} mice and *β-catenin*^{OsxER} mice compared to *Cre*-negative

littermates (Figure 5A, 6A). µCT images and quantitative analysis of decreased BV/TV, Tb.N,

Tb.Th and increased Tb.Sp indicated that both *β-catenin*^{Col2ER} mice and *β-catenin*^{OsxER} mice

presented severe bone loss in the femoral heads (Figure 5B, 6B). ABH staining and

histomorphological analysis further revealed that bone trabeculae were extensively replaced by fat

droplet clusters in the femoral head (especially in the subchondral region) of *β-catenin*^{Col2ER} mice

(Figure 5C, 5D), similar to the necrotic femoral heads in GONFH patients and

glucocorticoid-induced rat models. However, no fat droplet accumulation but only sparse and thin

bone trabeculae were observed in the femoral head of *β-catenin*^{OsxER} mice (Figure 6C, 6D).

Compared to *Cre*-negative littermates, the number of empty lacunae was increased in the femoral

head in *β-catenin*^{Col2ER} mice (Figure 5E), but no changed in *β-catenin*^{OsxER} mice (Figure 6E).

TUNEL staining reveled more apoptotic osteocytes in *β-catenin*^{Col2ER} mice compared to

Cre-negative littermates (Figure 5F). All these findings indicate that loss-of-function of β-catenin

in Col2$^+$ cells, but not in Osx$^+$ cells, led to an GONFH-like phenotype in femoral head.

Furthermore, decreased Runx2, ALP and increased PPAR-γ, CEBP/α expressions in

β-catenin^{Col2ER} mice (Figure 5G) revealed that this GONFH-like phenotype was caused by

imbalance of osteogenic/adipogenic differentiation of Col2$^+$ cells.

Older *β-catenin*^{Col2ER} mice display subchondral bone destruction and collapse tendency in

the femoral heads

Femoral head collapse occurs at the end-stage of GONFH and represents a poor joint

function*(Hines et al., 2021)*. To determine the occurrence of femoral head collapse, we further

analyzed 6-month-old *β-catenin*^{Col2ER} mice. Gross appearance showed that older *β-catenin*^{Col2ER}

mice lost smooth and regular outline on the femoral heads compared to *Cre*-negative littermates

and 3-month-old *β-catenin*^{Col2ER} mice (Figure 7A). μCT images showed severe bone loss (Figure

7B, 7C), subchondral bone destruction (Figure 7B, yellow arrows), local collapse (Figure 7B, red

arrows) and integral deformity (Figure 7B, red dotted lines) in the femoral heads of older

β-catenin^{Col2ER} mice. μCT quantitative analysis confirmed a significant increase of subchondral

bone defect area on the femoral head surface in older *β-catenin*^{Col2ER} mice (Figure 7D). Besides of

these early necrotic changes including sparse bone trabeculae, accumulated fat droplets (Figure 7E,

black triangles) and increased empty lacunae of osteoctye (Figure 7E, black arrows, Figure 7F),

ABH staining also showed subchondral bone destruction (Figure 7E, yellow arrows), femoral head

collapse or deformity (Figure 7E, red arrows and red dotted lines) in older *β-catenin*^{Col2ER} mice.

Compared to *Cre*-negative littermates, the older *β-catenin*^{Col2ER} mice presented a decrease of

loading-bearing stiffness, indicating a poor biomechanical support of the femoral heads (Figure

7G).

Wnt agonist 1 could not alleviate GONFH-like phenotype in *β-catenin*^{Col2ER} mice

As the therapeutic effects of Wnt agonist 1 on rat GONFH, we further determined whether it

could alleviate the GONFH-like phenotype in *β-catenin*^{Col2ER} mice. After induction with TM,

β-catenin^{Col2ER} mice were immediately injected with Wnt agonist 1 three times once week until

they were analyzed at the age of 3 months. However, μCT images and quantitative analysis of

BV/TV, Tb.N, Tb.Th and Tb.Sp showed severe bone loss were not restored in *β-catenin*[Col2ER] mice

by treating with Wnt agonist 1 (Figure 8A, 8B). Accumulated fat droplets, sparse trabeculae and

empty lacunae were still existed in the femoral heads of Wnt agonist 1 treated *β-catenin*[Col2ER] mice

(Figure 8C-8E). IHC staining showed no change of decreased ALP and increased FABP4 in the

femoral heads of *β-catenin*[Col2ER] mice after Wnt agonist 1 treatment (Figure 8F, 8G). These

findings indicate that Wnt agonist 1 could not restore the imbalanced osteogenic/adipogenic

differentiation of Col2[+] cells in *β-catenin*[Col2ER] mice.

Discussion

Despite it is well known that glucocorticoids are the certain etiology of GONFH, its

pathogenesis remains largely unknown. Current evidence including reduced osteogenic

differentiation ability of BMSCs from GONFH patients and clinical therapeutical effects of

BMSCs transplantation indicates that GONFH could be resulted from abnormal differentiation of

BMSCs*(Houdek et al., 2016; Wang et al., 2019a; Tabatabaee et al., 2015; Gangji and Hauzeur,*

2005). Glucocorticoid-induced animal models are commonly used for GONFH research, and

several previous studies have revealed that direct BMSCs modifications with miRNA*(Cao et al.,*

2021; Gu et al., 2016), lncRNA*(Wang et al., 2018c; Xu et al., 2022)* and circRNA*(Chen et al.,*

2020; Xiang et al., 2020) or microenvironment interventions by injecting platelet-rich plasma*(Xu*

et al., 2021; Wang et al., 2022), exosome*(Li et al., 2020; Fang et al., 2019)* and growth

factors*(Guzman et al., 2021; Rackwitz et al., 2012)* could promote osteogenic differentiation of

BMSCs and exhibit anti-GONFH effects in animals. In the present studies, we found severe

trabecular bone loss and massive fat droplets accumulation in the necrotic femoral heads of

GONFH patients and glucocorticoid-induced GONFH rats, confirming increased adipogenesis and

decreased osteogenesis in GONFH pathogenesis. Dex inhibited osteogenic differentiation of rat

BMSCs and promoted their adipogenic differentiation, revealing that this abnormal osteogenesis

and adipogenesis in GONFH rats were caused by the imbalance in osteogenic/adipogenic

differentiation of BMSCs. As β-catenin plays an important role in BMSCs differentiation, we then

analyzed its role in GONFH pathogenesis. The results showed a significant inhibition of β-catenin

signaling in the necrotic femoral heads of GONFH patients and glucocorticoid-induced GONFH

rats, while Wnt agonist 1 attenuated accumulated fat droplets and sparse trabeculae in GONFH

rats by activation of β-catenin signaling. The subsequent cellular experiments revealed that

β-catenin activation restored the imbalance of osteogenic/adipogenic differentiation of BMSCs caused by long-term exposure of Dex. Overall, these findings indicate that β-catenin inhibition-induced imbalanced osteogenic/adipogenic differentiation of BMSCs contributed to GONFH.

BMSCs are a group of heterogeneous cells, and their majorities are derived from skeletal progenitors*(Gao et al., 2021)*. It has reported that both Col2$^+$ progenitors and Osx$^+$ progenitors provide a large proportion of BMSCs*(Ono et al., 2014)*. Adult transgenic mice with conditional deletion of *β-catenin* in Col2$^+$ cells, but not Osx$^+$ cells, displayed a GONFH-like phenotype. Extensive trabecular bone were replaced by fat droplets in the femoral head (especially in the subchondral bone region) of *β-catenin^{Col2ER}* mice compared to *β-catenin^{OsxER}* mice. Cell fate mapping revealed a apparent difference in the distributions of Osx$^+$ cells and Col2$^+$ cells in the femoral head when mice induced at 2-week-old. Besides of the similar parts with Osx$^+$ cells (osteoblasts, osteocytes and stromal cells underneath growth plate), Col2$^+$ cells also largely expressed in the chondrocytes of articular cartilage, secondary ossification center and growth plate. Whether these Col2$^+$ chondrocytes in secondary ossification center and growth plate caused this GONFH-like phenotype, as it has been reported that the osteogenic commitment of these Col2$^+$ chondrocytes*(Shu et al., 2021; Li et al., 2022)*. At the age of 3 month, Col2$^+$ chondrocytes in the secondary ossification center and growth plate were almost replaced by Col2$^+$ osteoblasts, osteocytes and stromal cells, while GONFH-like phenotype including fat droplets accumulation and sparse bone trabeculae could be still observed in 6-month-old β-catenin^{Col2ER} mice, indicating that not only Col2$^+$ chondrocytes but all Col2$^+$ cells in different cellular morphology were involved in this GONFH-like phenotype. Nine months of lineage trace further revealed that Col2$^+$ cells had self-renew ability and could continuously produced osteoblasts. The decreased ALP, Runx2 and increased CEBP/α and PPAR-γ in *β-catenin^{Col2ER}* mice revealed that deletion of *β-catenin* shifted the commitment of Col2$^+$ cells from osteoblasts to adipocytes, which should be responsible for this GONFH-like phenotype.

Femoral head collapse is the key feature of late stage GONFH and is disastrous to hip joint*(Wang et al., 2019b)*. Although it has been well recognized that poor mechanical supports caused structural collapse of necrotic femoral head*(Fan et al., 2011)*, its detailed pathogenesis is poorly understood due to lack of ideal animal models. Glucocorticoid-induced bipedal and

quadrupedal animal models only exhibit the early necrotic changes of GONFH*(Xu et al., 2018)*,

without structure collapse of femoral head in MPS-induced rats. Other models using chemical or

physical approaches are all failed to mimic full-range osteonecrosis of GONFH in humans*(Wang*

et al., 2021). Importantly, we found that femoral head collapse occurred spontaneously in older

β-catenin^{Col2ER} mice. Besides the early necrotic changes including trabecular bone loss, empty

lacunae and fat droplets accumulation, extensive subchondral bone destruction were identified in

6-month-old *β-catenin*^{Col2ER} mice. Previous studies have revealed that strong trabeculae and

integrated subchondral bone are critical to maintain femoral head morphology*(Kawano et al.,*

2020; Motomura et al., 2011). A significant decrease of loading-bearing stiffness confirmed a poor

biomechanical supports in 6-month-old *β-catenin*^{Col2ER} mice. Therefore, it could be concluded that

β-catenin inhibition-induced imbalanced osteogenic/adipogenic differentiation of Col2^{+} cells

caused a poor biomechanical supports (manifesting with sparse trabeculae and subchondral bone

destruction), eventually leading to femoral head collapse in *β-catenin*^{Col2ER} mice.

Osteocytes are the bone cells with longest life, up to decades within their mineralized

environment*(Dallas et al., 2013; Bonewald, 2011)*. However, numerous empty lacunae of

osteocyte were observed in the necrotic femoral head of GONFH patients and

glucocorticoid-induced rats. Glucocorticoids are known to induce cell apoptosis*(Weinstein, 2012)*,

and TUNEL staining confirmed increased osteocyte apoptosis in the human and rat necrotic

femoral heads, leaving behinds these empty lacunae. The importance of β-catenin in the regulation

of apoptosis is well established in skeletal system*(Jähn et al., 2012; Kitase et al., 2010)*.

Consistently, the expression of β-catenin was significantly decreased in the human and rat necrotic

femoral heads. Moreover, we found that genetic deletion of β-catenin in Col2^{+} cells spontaneously

caused osteocyte apoptosis in mice and activation of β-catenin by systematically injecting Wnt

agonist 1 alleviated osteocyte apoptosis in glucocorticoid-induced rats, which indicate that

β-catenin inhibition-induced osteocyte apoptosis contributed to empty lacunae of GONFH.

In summary, our studies revealed that β-catenin inhibition induces the imbalance in

osteogenic/adipogenic differentiation of BMSCs and plays a crucial role in GONFH pathogenesis.

We also provided a *Col2*-specific *β-catenin* knockout mouse model, for the first time, which

mimics full spectrum of osteonecrosis phenotype of GONFH.

Materials and methods

Human specimen collection

Fifteen necrotic femoral head samples were collected from GONFH patients (ARCO grades III-IV) who received arthroplasty surgery. Ten non-necrotic femoral head samples were acquired from femoral neck fracture patients. All experiments with human specimens were approved by the Ethics Committee of the First Affiliated Hospital of Zhejiang Chinese Medical University (2018KL-005).

Animals and interventions

Thirty 2-month-old male Sprague-Dawley (SD) rats were purchased from the Animal Laboratory Center of Zhejiang Chinese Medical University (Zhejiang, China, SCXK 2014-0004), and randomly divided into the sham group, the GONFH group and the treatment group (n = 10 per group). We adopted Zheng's methods to establish a glucocorticoid-induced GONFH rat model*(Zheng et al., 2018)*. Firstly, rats in the GONFH group and the treatment group received once injection of lipopolysaccharide (0.2 mg/kg body weight, Sigma, USA) by tail vein, followed by three continuous intraperitoneal injections of 100 mg/kg body methylprednisolone (MPS, Pfizer, USA). At the subsequent 6 weeks, MPS were intraperitoneally injected at the dosage of 40 mg/kg body weight three times every week. Rats in the treatment group were intravenously injected with Wnt agonist 1 (5 mg/kg body weight, Selleck, USA) at synchronous frequency with MPS.

Col2-CreER^{T2} mice, *β-catenin$^{flox/flox}$* mice, *Osterix-CreER^{T2}* mice and *Rosa26^{tdTomato}* mice were obtained from Jackson Laboratory. To map cell fate of Col2$^+$ cells and Osx$^+$ cells in the femoral head, *Col2-CreER^{T2};Rosa26^{tdTomato}* (*Tomato^{Col2ER}*) mice and *Osterix-CreER^{T2};Rosa26^{tdTomato}* (*Tomato^{OsxER}*) mice induced by intraperitoneally injecting with tamoxifen (TM, 1 mg/10g body weight/day, diluted in corn oil) for 5 consecutive days at the age of 2 or 4 weeks, and traced as long as 9-month-old. To further investigate the role of β-catenin in GONFH pathogenesis, *Col2-CreER^{T2};β-catenin$^{flox/flox}$* (*β-catenin^{Col2ER}*) mice and *Osterix-CreER^{T2};β-catenin$^{flox/flox}$* (*β-catenin^{OsxER}*) mice were generated and induced with 5 continuous injection of TM. After induction, Wnt agonist 1 (5 mg/kg body weight) was intravenously injected into *β-catenin^{Col2ER}* mice three times every week until they were sacrificed at the age of 3 months. All animal experiments were approved by the Animal Ethics Committee of Zhejiang Chinese Medical University (No. 20190401-10).

Stereomicroscopy observation and biomechanical testing

Femoral head tissues harvested from human patients, glucocorticoid-induced GONFH rats, $β$-$catenin^{Col2ER}$ mice and $β$-$catenin^{OsxER}$ mice were viewed under a stereomicroscope (Model C-DSD230, Nikon, Japan) to assess their appearance characteristics. We further detected the loading-bearing stiffness of femoral head samples using a biomechanical testing machine (EnduraTec TestBenchTM system, Minnetonka, MN, USA) with an axial compression speed of 0.5 mm/min.

μCT analysis

Femoral head samples were scanned with a micro-computed tomography (μCT,Skyscan 1176，Bruker μCT, Kontich, Belgium) at a resolution of 9 μm. Three-dimensional (3D) structure was reconstructed using NRecon Software v1.6 as described previously*(Xia et al., 2019)*. The region of interest was selected for morphometric analysis including bone volume fraction (BV/TV, %), average trabecular thickness (Tb.Th, mm), average trabecular number (Tb.N, 1/mm), average trabecular separation (Tb.Sp, mm) and subchondral bone defect area (%).

Histology, immunohistochemistry and histomorphometry

Femoral head tissues were processed into 3-μm-thick paraffin section or 10-μm-thick frozen section as described previously*(Xia et al., 2020a)*. The paraffin sections were stained with Alcian Blue Hematoxylin (ABH) / Orange G for histological analysis. The trabecular bone area (%), fat droplet area (%) and empty lacuna rate (%) were measured using OsteoMetrics software (Decatur, GA). The frozen sections were performed with 4',6-diamidino-2-phenylindole (DAPI) staining and hematoxylin and eosin (H&E) staining for evaluating *Cre*-recombination efficiency.

Immunohistochemistry (IHC) assay were performed on the paraffin sections according to the previously established procedures*(Xia et al., 2020b)*. Briefly, sections were immersed into 0.01 M citrate buffer (Solarbio, Beijing, China) at 60℃ for 4 h to repair antigen. After washing with phosphate buffer (PBS) three times, the sections were incubated with primary antibodies including β-catenin (diluted 1:500, abcam, ab32572, UK), Runx2 (1:200, Abcam, ab76956, UK), alkaline phosphatase (ALP, 1:200, ARIGO, ARG57422, CN), osteocalcin (OCN; diluted 1:200, Takara, UK), PPAR-γ (1:300, Abcam,ab19481, UK), CEBP/α (1:400, Huabio, EM1710-20, CA) and fatty acid-binding protein (FABP4, diluted 1:200, abcam, ab92501, UK) overnight at 4℃. After incubation with secondary antibodies for 30 min at room temperature, tissue sections were stained

with diaminobenzidine (DAB) solution to detect positive staining and hematoxylin for counterstaining. Mean density (IOD/Area) as the parameter of immunohistochemical quantitative analysis was evaluated using Image-Pro Plus software (Media Cybernetics, Silver Spring, USA).

Rat BMSCs isolation and identification

BMSCs isolation were conducted as previously described*(Zhu et al., 2020)*. Briefly, bone marrow cells were isolated from the proximal femur of 4-week-old SD rats. After centrifugation at 200g for 10 min, cells were resuspended with α-MEM (Gibco, Grand Island, NY) containing 10% fetal bovine serum (FBS). By changing the medium 48 h later, no-adherent cells were removed and the primary BMSCs were harvested. At 100% confluence, primary cultures were detached using trypsin and subcultured at 8×10^3 cells/cm^2. A total of 50 μL third passage (P3) BMSCs suspension (4×10^6/mL) were mixed with 5 μL of rat CD90-APC, rat anti-CD29-BV421, rat anti-CD45-FITC or rat CD11b-PE (BD Pharmingen, Franklin Lakes, NJ, USA) in each flow cytometry tube. After incubation at room temperature for 30 min followed by twice washes of a standing buffer, the cells were resuspend by 500 μL staining buffer for flow cytometry analysis.

Rat BMSCs intervention and staining

To investigate the role of glucocorticoids on BMSCs differentiation, P3 BMSCs were treated with gradient concentrations of dexamethasone (Dex, 0, 10, 10^2, 10^3, 10^4, 10^5 nM) with or without SKL2001 (a agonist of β-catenin) for 72 h. For oil-red O staining, BMSCs were treated with adipogenic medium (α-MEM containing 10% FBS, 0.25 mM methylisobutylxanthine, 5 μg/mL insulin, and 50 μM indomethacin) for 72 h followed by 5 μg/mL insulin alone for an additional 48 h. After fixed in 4% paraformaldehyde for 10 min, BMSCs were stained in 60% saturated oil-red O solution for 5 min. For ALP staining, BMSCs were treated with osteogenic medium (α-MEM containing 10% FBS, 50 μg/mL ascorbic acid and 5 mM β-glycerophosphate) for 14 d, fixed in 4% paraformaldehyde for 10 min and stained with ALP staining kit (Thermo Fisher Scientific, Rockford, IL, USA) for 15 min.

Western blot analysis

According to the appearance and CT images, two piece (about $2*2*2$ cm^3) of necrotic tissues were cut from each human necrotic femoral head, and the corresponding region from human fractured femoral head. These necrotic and non-necrotic tissues were mixed and ground into powder after processed by liquid nitrogen, respectively. Tissue powder and rat P3 BMSCs were

lysed by protein extraction reagent, and the liquid supernatant containing total protein was

obtained through centrifuging at 12000 g for 20 min. Then, proteins were separated by SDS-PAGE

and transferred onto nitrocellulose membranes. The membrane were incubated overnight with

primary antibodies including β-catenin, Runx2, Osterix, ALP, PPAR-γ and CEBP/α at 1:1000

dilution. Next, the membranes were incubated with appropriate horseradish peroxidase-conjugated

secondary antibodies at 37 ℃ for 1 h. The immune complexes were detected using a

chemiluminescence kit and visualized via the ChemiDoc XRS Gel documentation system (Bio-rad,

CA). The gray intensity was measured by the software of Quantity One (Bio-Rad, CA).

Statistical analysis

All data were presented as mean ± standard deviation (SD). Statistical analyses including

unpaired Student's *t*-test (two groups) and one-way ANOVA followed by Tukey's test (multiple

groups) were performed with the software of SPSS 22.0 software. *$P<0.05$, **$P<0.01$ was

considered statistically significant.

Data Availability Statement
Raw data are mainly attached in the current study. Any other data involving this study are available from the corresponding author upon request.

Acknowledgements
This work has been supported by Chinese National Natural Science Foundation (grants nos. 81873325, 82074469, 82104885), Natural Science Foundation of Zhejiang Province (grant no. Q22H271191), Ningbo medical key discipline (No. 2022-B01).

Author Contributions
Di Chen, Chenjie Xia and Hongting Jin conceived this study. Chenjie Xia wrote the paper. Di Chen and Hongting Jin revised the paper. Chenjie Xia, Huihui Xu, Ping-er Wang, Liang Fang, Jiali Chen, Wenhua Yuan, Danqing Fu, Xucheng Wang performed the experiments and analyzed the data. Bangjian He, Luwei Xiao, Peijian Tong and Chengliang Wu collected the human samples.

References
Bonewald LF. 2011. The amazing osteocyte. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research* **26**:229-238. doi: 10.1002/jbmr.320, PMID: 21254230
Cao Y, Jiang C, Wang X, Wang H, Yan Z, Yuan H. 2021. Reciprocal effect of microRNA-224 on osteogenesis and

adipogenesis in steroid-induced osteonecrosis of the femoral head. *Bone* **145**:115844. doi: 10.1016/j.bone.2021.115844, PMID: 33453444

Chang CC, Greenspan A, Gershwin ME. 1993. Osteonecrosis: current perspectives on pathogenesis and treatment. *Seminars in arthritis and rheumatism* **23**:47-69. doi: 10.1016/s0049-0172(05)80026-5, PMID: 8235665

Chen CY, Rao SS, Yue T, Tan YJ, Yin H, Chen LJ, Luo MJ, Wang Z, Wang YY, Hong CG, Qian YX, He ZH, Liu JH, Yang F, Huang FY, Tang SY, Xie H. 2022. Glucocorticoid-induced loss of beneficial gut bacterial extracellular vesicles is associated with the pathogenesis of osteonecrosis. *Science advances* **8**:eabg8335. doi: 10.1126/sciadv.abg8335, PMID: 35417243

Chen G, Wang Q, Li Z, Yang Q, Liu Y, Du Z, Zhang G, Song Y. 2020. Circular RNA CDR1as promotes adipogenic and suppresses osteogenic differentiation of BMSCs in steroid-induced osteonecrosis of the femoral head. *Bone* **133**:115258. doi: 10.1016/j.bone.2020.115258, PMID: 32018039

Chen XJ, Shen YS, He MC, Yang F, Yang P, Pang FX, He W, Cao YM, Wei QS. 2019. Polydatin promotes the osteogenic differentiation of human bone mesenchymal stem cells by activating the BMP2-Wnt/β-catenin signaling pathway. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* **112**:108746. doi: 10.1016/j.biopha.2019.108746, PMID: 30970530

Dallas SL, Prideaux M, Bonewald LF. 2013. The osteocyte: an endocrine cell ... and more. *Endocrine reviews* **34**:658-690. doi: 10.1210/er.2012-1026, PMID: 23612223

Fan M, Peng J, Qin L, Lu S. 2011. Experimental animal models of osteonecrosis. *Rheumatology international* **31**:983-994. doi: 10.1007/s00296-011-1819-9, PMID: 21340568

Fang S, Li Y, Chen P. 2019. Osteogenic effect of bone marrow mesenchymal stem cell-derived exosomes on steroid-induced osteonecrosis of the femoral head. *Drug design, development and therapy* **13**:45-55. doi: 10.2147/DDDT.S178698, PMID: 30587927

Feng X, Xiang Q, Jia J, Guo T, Liao Z, Yang S, Cai X, Liu X. 2022. CircHGF suppressed cell proliferation and osteogenic differentiation of BMSCs in ONFH via inhibiting miR-25-3p binding to SMAD7. *Molecular therapy. Nucleic acids* **28**:99-113. doi: 10.1016/j.omtn.2022.02.017, PMID: 35402073

Gangji V, Hauzeur JP. 2005. Treatment of osteonecrosis of the femoral head with implantation of autologous bone-marrow cells. Surgical technique. *The Journal of bone and joint surgery. American volume* **87 Suppl 1**:106-112. doi: 10.2106/JBJS.D.02662, PMID: 15743852

Gao Q, Wang L, Wang S, Huang B, Jing Y, Su J. 2021. Bone Marrow Mesenchymal Stromal Cells: Identification, Classification, and Differentiation. *Frontiers in cell and developmental biology* **9**:787118. doi: 10.3389/fcell.2021.787118, PMID: 35047499

Gu C, Xu Y, Zhang S, Guan H, Song S, Wang X, Wang Y, Li Y, Zhao G. 2016. miR-27a attenuates adipogenesis and promotes osteogenesis in steroid-induced rat BMSCs by targeting PPARγ and GREM1. *Scientific reports* **6**:38491. doi: 10.1038/srep38491, PMID: 27910957

Guzman RA, Maruyama M, Moeinzadeh S, Lui E, Zhang N, Storaci HW, Tam K, Huang EE, Utsunomiya T, Rhee C, Gao Q, Yao Z, Yang YP, Goodman SB. 2021. The effect of genetically modified platelet-derived growth factor-BB over-expressing mesenchymal stromal cells during core decompression for steroid-associated osteonecrosis of the femoral head in rabbits. *Stem cell research & therapy* **12**:503. doi: 10.1186/s13287-021-02572-7, PMID: 34526115

Han L, Wang B, Wang R, Gong S, Chen G, Xu W. 2019. The shift in the balance between osteoblastogenesis and adipogenesis of mesenchymal stem cells mediated by glucocorticoid receptor. *Stem cell research & therapy* **10**:377. doi: 10.1186/s13287-019-1498-0, PMID: 31805987

Hines JT, Jo WL, Cui Q, Mont MA, Koo KH, Cheng EY, Goodman SB, Ha YC, Hernigou P, Jones LC, Kim SY, Sakai T, Sugano N, Yamamoto T, Lee MS, Zhao D, Drescher W, Kim TY, Lee YK, Yoon BH, Baek SH,

Ando W, Kim HS, Park JW. 2021. Osteonecrosis of the Femoral Head: an Updated Review of ARCO on Pathogenesis, Staging and Treatment. *Journal of Korean medical science* **36**:e177. doi: 10.3346/jkms.2021.36.e177, PMID: 34155839

Houdek MT, Wyles CC, Packard BD, Terzic A, Behfar A, Sierra RJ. 2016. Decreased Osteogenic Activity of Mesenchymal Stem Cells in Patients With Corticosteroid-Induced Osteonecrosis of the Femoral Head. *The Journal of arthroplasty* **31**:893-898. doi: 10.1016/j.arth.2015.08.017, PMID: 26404846

Jähn K, Lara-Castillo N, Brotto L, Mo CL, Johnson ML, Brotto M, Bonewald LF. 2012. Skeletal muscle secreted factors prevent glucocorticoid-induced osteocyte apoptosis through activation of β-catenin. *European cells & materials* **24**:197-209; discussion 209-210. doi: 10.22203/ecm.v024a14, PMID: 22972510

Kawano K, Motomura G, Ikemura S, Yamaguchi R, Baba S, Xu M, Nakashima Y. 2020. Differences in the microarchitectural features of the lateral collapsed lesion between osteonecrosis and subchondral insufficiency fracture of the femoral head. *Bone* **141**:115585. doi: 10.1016/j.bone.2020.115585, PMID: 32795680

Kitase Y, Barragan L, Qing H, Kondoh S, Jiang JX, Johnson ML, Bonewald LF. 2010. Mechanical induction of PGE2 in osteocytes blocks glucocorticoid-induced apoptosis through both the β-catenin and PKA pathways. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research* **25**:2657-2668. doi: 10.1002/jbmr.168, PMID: 20578217

Lavernia CJ, Sierra RJ, Grieco FR. 1999. Osteonecrosis of the femoral head. *The Journal of the American Academy of Orthopaedic Surgeons* **7**:250-261. doi: 10.5435/00124635-199907000-00005, PMID: 10434079

Lee JS, Lee JS, Roh HL, Kim CH, Jung JS, Suh KT. 2006. Alterations in the differentiation ability of mesenchymal stem cells in patients with nontraumatic osteonecrosis of the femoral head: comparative analysis according to the risk factor. *Journal of orthopaedic research : official publication of the Orthopaedic Research Society* **24**:604-609. doi: 10.1002/jor.20078, PMID: 16514658

Li L, Wang Y, Yu X, Bao Y, An L, Wei X, Yu W, Liu B, Li J, Yang J, Xia Y, Liu G, Cao F, Zhang X, Zhao D. 2020. Bone marrow mesenchymal stem cell-derived exosomes promote plasminogen activator inhibitor 1 expression in vascular cells in the local microenvironment during rabbit osteonecrosis of the femoral head. *Stem cell research & therapy* **11**:480. doi: 10.1186/s13287-020-01991-2, PMID: 33176873

Li X, Yang S, Yuan G, Jing D, Qin L, Zhao H, Yang S. 2022. Type II collagen-positive progenitors are important stem cells in controlling skeletal development and vascular formation. *Bone research* **10**:46. doi: 10.1038/s41413-022-00214-z, PMID: 35739091

Liu Y, Strecker S, Wang L, Kronenberg MS, Wang W, Rowe DW, Maye P. 2013. Osterix-cre labeled progenitor cells contribute to the formation and maintenance of the bone marrow stroma. *PloS one* **8**:e71318. doi: 10.1371/journal.pone.0071318, PMID: 23951132

Motomura G, Yamamoto T, Yamaguchi R, Ikemura S, Nakashima Y, Mawatari T, Iwamoto Y. 2011. Morphological analysis of collapsed regions in osteonecrosis of the femoral head. *The Journal of bone and joint surgery. British volume* **93**:184-187. doi: 10.1302/0301-620X.93B225476, PMID: 21282756

Ono N, Ono W, Nagasawa T, Kronenberg HM. 2014. A subset of chondrogenic cells provides early mesenchymal progenitors in growing bones. *Nature cell biology* **16**:1157-1167. doi: 10.1038/ncb3067, PMID: 25419849

Petek D, Hannouche D, Suva D. 2019. Osteonecrosis of the femoral head: pathophysiology and current concepts of treatment. *EFORT open reviews* **4**:85-97. doi: 10.1302/2058-5241.4.180036, PMID: 30993010

Qin L, Yao D, Zheng L, Liu WC, Liu Z, Lei M, Huang L, Xie X, Wang X, Chen Y, Yao X, Peng J, Gong H, Griffith JF, Huang Y, Zheng Y, Feng JQ, Liu Y, Chen S, Xiao D, Wang D, Xiong J, Pei D, Zhang P, Pan X, Wang X, Lee KM, Cheng CY. 2015. Phytomolecule icaritin incorporated PLGA/TCP scaffold for

steroid-associated osteonecrosis: Proof-of-concept for prevention of hip joint collapse in bipedal emus and mechanistic study in quadrupedal rabbits. *Biomaterials* **59**:125-143. doi: 10.1016/j.biomaterials.2015.04.038, PMID: 25968462

Rackwitz L, Eden L, Reppenhagen S, Reichert JC, Jakob F, Walles H, Pullig O, Tuan RS, Rudert M, Nöth U. 2012. Stem cell- and growth factor-based regenerative therapies for avascular necrosis of the femoral head. *Stem cell research & therapy* **3**:7. doi: 10.1186/scrt98, PMID: 22356811

Scaglione M, Fabbri L, Celli F, Casella F, Guido G. 2015. Hip replacement in femoral head osteonecrosis: current concepts. *Clinical cases in mineral and bone metabolism : the official journal of the Italian Society of Osteoporosis, Mineral Metabolism, and Skeletal Diseases* **12**:51-54. doi: 10.11138/ccmbm/2015.12.3s.051, PMID: 27134633

Shu HS, Liu YL, Tang XT, Zhang XS, Zhou B, Zou W, Zhou BO. 2021. Tracing the skeletal progenitor transition during postnatal bone formation. *Cell stem cell* **28**:2122-2136.e3. doi: 10.1016/j.stem.2021.08.010, PMID: 34499868

Song L, Liu M, Ono N, Bringhurst FR, Kronenberg HM, Guo J. 2012. Loss of wnt/β -catenin signaling causes cell fate shift of preosteoblasts from osteoblasts to adipocytes. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research* **27**:2344-2358. doi: 10.1002/jbmr.1694, PMID: 22729939

Sun H, Zhang W, Yang N, Xue Y, Wang T, Wang H, Zheng K, Wang Y, Zhu F, Yang H, Xu W, Xu Y, Geng D. 2021. Activation of cannabinoid receptor 2 alleviates glucocorticoid-induced osteonecrosis of femoral head with osteogenesis and maintenance of blood supply. *Cell death & disease* **12**:1035. doi: 10.1038/s41419-021-04313-3, PMID: 34718335

Tabatabaee RM, Saberi S, Parvizi J, Mortazavi SM, Farzan M. 2015. Combining Concentrated Autologous Bone Marrow Stem Cells Injection With Core Decompression Improves Outcome for Patients with Early-Stage Osteonecrosis of the Femoral Head: A Comparative Study. *The Journal of arthroplasty* **30**:11-15. doi: 10.1016/j.arth.2015.06.022, PMID: 26143238

Takada I, Kouzmenko AP, Kato S. 2009. Wnt and PPARgamma signaling in osteoblastogenesis and adipogenesis. *Nature reviews. Rheumatology* **5**:442-447. doi: 10.1038/nrrheum.2009.137, PMID: 19581903

Tao SC, Yuan T, Rui BY, Zhu ZZ, Guo SC, Zhang CQ. 2017. Exosomes derived from human platelet-rich plasma prevent apoptosis induced by glucocorticoid-associated endoplasmic reticulum stress in rat osteonecrosis of the femoral head via the Akt/Bad/Bcl-2 signal pathway. *Theranostics* **7**:733-750. doi: 10.7150/thno.17450, PMID: 28255363

Tencerova M, Kassem M. 2016. The Bone Marrow-Derived Stromal Cells: Commitment and Regulation of Adipogenesis. *Frontiers in endocrinology* **7**:127. doi: 10.3389/fendo.2016.00127, PMID: 27708616

Wang A, Ren M, Wang J. 2018b. The pathogenesis of steroid-induced osteonecrosis of the femoral head: A systematic review of the literature. *Gene* **671**:103-109. doi: 10.1016/j.gene.2018.05.091, PMID: 29859289

Wang BL, Sun W, Shi ZC, Lou JN, Zhang NF, Shi SH, Guo WS, Cheng LM, Ye LY, Zhang WJ, Li ZR. 2008. Decreased proliferation of mesenchymal stem cells in corticosteroid-induced osteonecrosis of femoral head. *Orthopedics* **31**:444. doi: 10.3928/01477447-20080501-33, PMID: 19292322

Wang C, Meng H, Wang Y, Zhao B, Zhao C, Sun W, Zhu Y, Han B, Yuan X, Liu R, Wang X, Wang A, Guo Q, Peng J, Lu S. 2018a. Analysis of early stage osteonecrosis of the human femoral head and the mechanism of femoral head collapse. *International journal of biological sciences* **14**:156-164. doi: 10.7150/ijbs.18334, PMID: 29483834

Wang C, Sun W, Ling S, Wang Y, Wang X, Meng H, Li Y, Yuan X, Li J, Liu R, Zhao D, Lu Q, Wang A, Guo Q, Lu S, Tian H, Li Y, Peng J. 2019b. AAV-Anti-miR-214 Prevents Collapse of the Femoral Head in

Osteonecrosis by Regulating Osteoblast and Osteoclast Activities. *Molecular therapy. Nucleic acids* **18**:841-850. doi: 10.1016/j.omtn.2019.09.030, PMID: 31739209

Wang C, Wu Z, Li X, Shi L, Xie Q, Liu D, Guo Z, Zheng W. 2021. An animal model of early-stage femoral head osteonecrosis induced by cryo-insult in small tailed Han sheep. *Journal of orthopaedic translation* **26**:84-91. doi: 10.1016/j.jot.2020.06.004, PMID: 33437627

Wang Q, Yang Q, Chen G, Du Z, Ren M, Wang A, Zhao H, Li Z, Zhang G, Song Y. 2018c. LncRNA expression profiling of BMSCs in osteonecrosis of the femoral head associated with increased adipogenic and decreased osteogenic differentiation. *Scientific reports* **8**:9127. doi: 10.1038/s41598-018-27501-2, PMID: 29904151

Wang Y, Luan S, Yuan Z, Lin C, Fan S, Wang S, Ma C, Wu S. 2022. Therapeutic effect of platelet-rich plasma on glucocorticoid-induced rat bone marrow mesenchymal stem cells in vitro. *BMC musculoskeletal disorders* **23**:151. doi: 10.1186/s12891-022-05094-2, PMID: 35168574

Wang Z, Sun QM, Zhang FQ, Zhang QL, Wang LG, Wang WJ. 2019a. Core decompression combined with autologous bone marrow stem cells versus core decompression alone for patients with osteonecrosis of the femoral head: A meta-analysis. *International journal of surgery (London, England)* **69**:23-31. doi: 10.1016/j.ijsu.2019.06.016, PMID: 31301432

Weinstein RS. 2012. Glucocorticoid-induced osteonecrosis. *Endocrine* **41**:183-190. doi: 10.1007/s12020-011-9580-0, PMID: 22169965

Xia C, Ge Q, Fang L, Yu H, Zou Z, Zhang P, Lv S, Tong P, Xiao L, Chen D, Wang PE, Jin H. 2020a. TGF-β/Smad2 signalling regulates enchondral bone formation of Gli1+ periosteal cells during fracture healing. *Cell proliferation* **53**:e12904. doi: 10.1111/cpr.12904, PMID: 32997394

Xia C, Wang P, Fang L, Ge Q, Zou Z, Dong R, Zhang P, Shi Z, Xu R, Zhang L, Luo C, Ying J, Xiao L, Shen J, Chen D, Tong P, Jin H. 2019. Activation of β-catenin in Col2-expressing chondrocytes leads to osteoarthritis-like defects in hip joint. *Journal of cellular physiology* **234**:18535-18543. doi: 10.1002/jcp.28491, PMID: 30912140

Xia C, Zou Z, Fang L, Ge Q, Zhang P, Xu H, Xu R, Shi Z, Lin H, Ding X, Xiao L, Tong P, Wang PE, Jin H. 2020b. Bushenhuoxue formula promotes osteogenic differentiation of growth plate chondrocytes through β-catenin-dependent manner during osteoporosis. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* **127**:110170. doi: 10.1016/j.biopha.2020.110170, PMID: 32334373

Xiang S, Li Z, Weng X. 2020. Changed cellular functions and aberrantly expressed miRNAs and circRNAs in bone marrow stem cells in osteonecrosis of the femoral head. *International journal of molecular medicine* **45**:805-815. doi: 10.3892/ijmm.2020.4455, PMID: 31922208

Xu C, Wang J, Zhu T, Shen Y, Tang X, Fang L, Xu Y. 2016. Cross-Talking Between PPAR and WNT Signaling and its Regulation in Mesenchymal Stem Cell Differentiation. *Current stem cell research & therapy* **11**:247-254. doi: 10.2174/1574888x10666150723145707, PMID: 26201865

Xu HH, Li SM, Fang L, Xia CJ, Zhang P, Xu R, Shi ZY, Zou Z, Ge QW, Wang P, Tong PJ, Jin HT. 2021. Platelet-rich plasma promotes bone formation, restrains adipogenesis and accelerates vascularization to relieve steroids-induced osteonecrosis of the femoral head. *Platelets* **32**:950-959. doi: 10.1080/09537104.2020.1810221, PMID: 32835568

Xu J, Gong H, Lu S, Deasey MJ, Cui Q. 2018. Animal models of steroid-induced osteonecrosis of the femoral head-a comprehensive research review up to 2018. *International orthopaedics* **42**:1729-1737. doi: 10.1007/s00264-018-3956-1, PMID: 29705870

Xu Y, Jiang Y, Wang Y, Jia B, Gao S, Yu H, Zhang H, Lv C, Li H, Li T. 2022. LINC00473-modified bone marrow mesenchymal stem cells incorporated thermosensitive PLGA hydrogel transplantation for steroid-induced

osteonecrosis of femoral head: A detailed mechanistic study and validity evaluation. *Bioengineering & translational medicine* **7**:e10275. doi: 10.1002/btm2.10275, PMID: 35600648

Yang N, Sun H, Xue Y, Zhang W, Wang H, Tao H, Liang X, Li M, Xu Y, Chen L, Zhang L, Huang L, Geng D. 2021. Inhibition of MAGL activates the Keap1/Nrf2 pathway to attenuate glucocorticoid-induced osteonecrosis of the femoral head. *Clinical and translational medicine* **11**:e447. doi: 10.1002/ctm2.447, PMID: 34185425

Yin L, Li YB, Wang YS. 2006. Dexamethasone-induced adipogenesis in primary marrow stromal cell cultures: mechanism of steroid-induced osteonecrosis. *Chinese medical journal* **119**:581-588. PMID: 16620700

Yu Z, Fan L, Li J, Ge Z, Dang X, Wang K. 2016. Lithium prevents rat steroid-related osteonecrosis of the femoral head by β-catenin activation. *Endocrine* **52**:380-390. doi: 10.1007/s12020-015-0747-y, PMID: 26458922

Yuan Z, Li Q, Luo S, Liu Z, Luo D, Zhang B, Zhang D, Rao P, Xiao J. 2016. PPARγ and Wnt Signaling in Adipogenic and Osteogenic Differentiation of Mesenchymal Stem Cells. *Current stem cell research & therapy* **11**:216-225. doi: 10.2174/1574888x10666150519093429, PMID: 25986621

Zaidi M, Sun L, Robinson LJ, Tourkova IL, Liu L, Wang Y, Zhu LL, Liu X, Li J, Peng Y, Yang G, Shi X, Levine A, Iqbal J, Yaroslavskiy BB, Isales C, Blair HC. 2010. ACTH protects against glucocorticoid-induced osteonecrosis of bone. *Proceedings of the National Academy of Sciences of the United States of America* **107**:8782-8787. doi: 10.1073/pnas.0912176107, PMID: 20421485

Zalavras C, Shah S, Birnbaum MJ, Frenkel B. 2003. Role of apoptosis in glucocorticoid-induced osteoporosis and osteonecrosis. *Critical reviews in eukaryotic gene expression* **13**:221-235. PMID: 14696969

Zhang C, Zou YL, Ma J, Dang XQ, Wang KZ. 2015. Apoptosis associated with Wnt/β-catenin pathway leads to steroid-induced avascular necrosis of femoral head. *BMC musculoskeletal disorders* **16**:132. doi: 10.1186/s12891-015-0606-2, PMID: 26037065

Zhang P, Xu H, Wang P, Dong R, Xia C, Shi Z, Xu R, Fang L, Zou Z, Ge Q, Tong P, Jin H. 2019. Yougui pills exert osteoprotective effects on rabbit steroid-related osteonecrosis of the femoral head by activating β-catenin. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* **120**:109520. doi: 10.1016/j.biopha.2019.109520, PMID: 31629251

Zhang XY, Li HN, Chen F, Chen YP, Chai Y, Liao JZ, Gan B, Chen DP, Li S, Liu YQ. 2021. Icariin regulates miR-23a-3p-mediated osteogenic differentiation of BMSCs via BMP-2/Smad5/Runx2 and WNT/β-catenin pathways in osteonecrosis of the femoral head. *Saudi pharmaceutical journal : SPJ : the official publication of the Saudi Pharmaceutical Society* **29**:1405-1415. doi: 10.1016/j.jsps.2021.10.009, PMID: 35002378

Zhao D, Zhang F, Wang B, Liu B, Li L, Kim SY, Goodman SB, Hernigou P, Cui Q, Lineaweaver WC, Xu J, Drescher WR, Qin L. 2020. Guidelines for clinical diagnosis and treatment of osteonecrosis of the femoral head in adults (2019 version). *Journal of orthopaedic translation* **21**:100-110. doi: 10.1016/j.jot.2019.12.004, PMID: 32309135

Zhao H, Yeersheng R, Xia Y, Kang P, Wang W. 2021. Hypoxia Enhanced Bone Regeneration Through the HIF-1α/β-Catenin Pathway in Femoral Head Osteonecrosis. *The American journal of the medical sciences* **362**:78-91. doi: 10.1016/j.amjms.2021.03.005, PMID: 33727018

Zheng LZ, Wang JL, Kong L, Huang L, Tian L, Pang QQ, Wang XL, Qin L. 2018. Steroid-associated osteonecrosis animal model in rats. *Journal of orthopaedic translation* **13**:13-24. doi: 10.1016/j.jot.2018.01.003, PMID: 29662787

Zhu W, Guo M, Yang W, Tang M, Chen T, Gan D, Zhang D, Ding X, Zhao A, Zhao P, Yan W, Zhang J. 2020. CD41-deficient exosomes from non-traumatic femoral head necrosis tissues impair osteogenic differentiation and migration of mesenchymal stem cells. *Cell death & disease* **11**:293. doi:

List of Figure Captions

Figure 1. Abnormal osteogenesis and adipogenesis with decreased β-catenin signaling in the necrotic femoral heads of GONFH patients. Necrotic (n=15) and Non-necrotic (n=10) femoral head samples were obtained from the patients with GONFH or femoral neck fracture, respectively. (A) Gross anatomy analysis of human necrotic and non-necrotic femoral head samples. Black asterisks: subchondral collapsed region. Black arrows: liquefied necrotic region. (B) μCT images of human necrotic and non-necrotic femoral heads. Red and Yellow boxed area: 3D images of subchondral collapsed region and liquefied necrotic region, respectively. White boxed areas: 3D images of corresponding regions in the non-necrotic femoral heads. (C) Quantitative analysis of

BV/TV, Tb.N, Tb.Th and Tb.Sp on the necrotic regions. (D) ABH staining of human necrotic and non-necrotic femoral heads. a,c: high magnification images of bone marrow; b,d: high magnification images of bone trabeculae; Yellow arrows: subchondral bone destruction; black arrows: empty lacunae of osteocyte; black triangles: fat droplet. (E) Histomorphological quantitative analysis of trabecular bone area and fat droplet area. (F) Histomorphological quantitative analysis of empty lacunae rate. (G) TUNEL staining of osteocytes in human necrotic and non-necrotic femoral heads. (H) IHC staining of Runx2 and FABP4 expressions in human necrotic and non-necrotic femoral heads. (I) Western blot of β-catenin, Osterix, PPAR-γ and CEBP/α in human necrotic and non-necrotic femoral heads.

Figure 2. Systemic injection of Wnt agonist 1 alleviates abnormal osteogenesis and adipogenesis in rat GONFH femoral heads. The rat model of GONFH was established by a co-induction of lipopolysaccharide and methylprednisolone (MPS). The femoral head samples were harvested after GONFH rats were intravenously injected with Wnt agonist 1 for 6 weeks. (A) Gross anatomy analysis and µCT images of femoral heads in sham, model and Wnt agonist 1 treated groups. (B) µCT quantitative analysis of BV/TV, Tb.N, Tb.Th and Tb.Sp in each group. (C) ABH staining of

femoral heads in each group. a-c, high magnification images of the representative region; black arrows: empty lacunae of osteocyte; black triangles: fat droplet. (D) Histomorphological quantitative analysis of trabecular bone area and fat droplet area in each group. (E) Histomorphological quantitative analysis of empty lacunae rate in each group. (F) TUNEL staining of osteocytes in the femoral heads of GONFH rats. (G, H) IHC staining and quantitative analysis of β-catenin, ALP and FABP4 in rat femoral heads of each group.

Figure 3. Activation of β-catenin redirected Dex-induced imbalanced osteogenic/adipogenic differentiation of BMSCs. Primary BMSCs were extracted from the proximal femur of 4-week-old SD rats and cultivated to passage 3 for subsequent experiments. (A) Flow cytometry analyzing the

surface markers of rat BMSCs. (B) ALP staining and oil red O staining of BMSCs at the increasing concentrations of dexamethasone (Dex). (C, D) Western blot and quantitative analysis of β-catenin, Runx2, ALP, PPAR-γ and CEBP/α in BMSCs at the increasing concentrations of Dex. (E) ALP staining and oil red O staining of rat BMSCs at the condition with or without 10^4 nM Dex and 20 μM SKL2001 (an agonist of β-catenin). (F, G) Western blot and quantitative analysis of β-catenin, Runx2, ALP, PPAR-γ and CEBP/α in rat BMSCs at the condition with or without 10^4 nM Dex and 20 μM SKL2001.

Figure 4. Fate mapping of Col2$^+$ cells and Osx$^+$ cells in the femoral head. *Tomato^{Col2ER}* mice and *Tomato^{OsxER}* mice were continuously received 5 dose of TM injections (1 mg/10 g body weight) at the age of 2 weeks for fate mapping analysis. (A) Distributions of Col2$^+$ and Osx$^+$ cells in the femoral head of 1-month-old *Tomato^{Col2ER}* mice and *Tomato^{OsxER}* mice. White arrowheads: chondrocyte; Green arrowheads: osteoblast; Yellow arrowheads: osteocyte; Red arrowheads: bone marrow stromal cell; GP: growth plate; SOC: second ossification center; BM: bone marrow. (B) Lineage trace of Col2$^+$ cells in the femoral head for 9 months. a: high magnification image of bone marrow region. (C) Poor *Cre*-recombinase efficiency in the femoral head of *Tomato^{Col2ER}* mice with TM induction at the age of 1 month. (D, E) Femoral heads were harvested from 3-month-old

β-catenin^{Col2ER} mice and β-catenin^{OsxER} mice to detect expression of β-catenin. IHC staining and quantitative analysis of β-catenin in the femoral heads of 3-month-old β-catenin^{Col2ER} mice and β-catenin^{OsxER} mice.

Figure 5. Deletion of *β-catenin* in Col2[+] cells leads to a GNOFH-like phenotype. Femoral heads were harvested from 3-month-old *β-catenin^{Col2ER}* mice and *Cre*-negative littermates with 5 continuous dosage of TM injections (1 mg/10 g body weight) at the age of 2 weeks. (A) Gross anatomy analysis of femoral heads in *β-catenin^{Col2ER}* mice and *Cre*-negative littermates. (B) Representative μCT images and quantitative analysis of BV/TV, Tb. N, Tb. Th and Tb. Sp in the femoral heads of *β-catenin^{Col2ER}* mice. (C) ABH staining of femoral heads in *β-catenin^{Col2ER}* mice. a-d: high magnification images of representative subchondral bone region; Black triangle arrowheads: fat droplet; Black arrows: empty lacunae of osteocyte. (D) Histomorphological quantitative analysis of trabecular bone area and fat droplet area in the femoral heads of *β-catenin^{Col2ER}* mice. (E) Histomorphological quantitative analysis of empty lacunae rate in the femoral heads of *β-catenin^{Col2ER}* mice. (F) TUNEL staining of osteocytes in the femoral heads of *β-catenin^{Col2ER}* mice. (G) IHC staining and quantitative analysis of Runx2, ALP, PPAR-γ and CEBP/α in the femoral heads of *β-catenin^{Col2ER}* mice.

Figure 6. Loss-of-function of *β-catenin* in Osx[+] cells causes bone loss in the femoral head. Femoral heads were harvested from 3-month-old *β-catenin^{OsxER}* mice and *Cre*-negative littermates with 5 continuous dosage of TM injections (1 mg/10 g body weight) at the age of 2 weeks. (A) Gross anatomy analysis of femoral heads in *β-catenin^{OsxER}* mice and *Cre*-negative littermates. (B) μCT images and quantitative analysis of BV/TV, Tb. N, Tb. Th and Tb. Sp in the femoral heads of *β-catenin^{OsxER}* mice. (C) ABH staining of femoral heads in *β-catenin^{OsxER}* mice. (D) Histomorphological quantitative analysis of trabecular bone area and fat droplet area in the femoral heads of *β-catenin^{OsxER}* mice. (E) Histomorphological quantitative analysis of empty lacunae rate in the femoral heads of *β-catenin^{OsxER}* mice.

Figure 7. Older *β-catenin*^Col2ER^ mice display subchondral bone destruction and collapse tendency in femoral heads. Femoral heads were harvested from 6-month-old *β-catenin*^Col2ER^ mice and *Cre*-negative littermates with 5 continuous dosage of TM injections (1 mg/10 g body weight) at the age of 2 weeks. (A) Gross anatomy analysis of femoral heads in 6-month-old *β-catenin*^Col2ER^ mice. (B) Representative μCT images of femoral heads in 6-month-old *β-catenin*^Col2ER^ mice. Red dotted lines: integral deformity. Red arrows: local collapse. Yellow arrows: subchondral bone destruction. (C) Quantitative analysis of BV/TV in the femoral heads in 6-month-old *β-catenin*^Col2ER^ mice. (D) Quantitative analysis of subchondral bone defect area on the femoral head surface in older *β-catenin*^Col2ER^ mice. (E) ABH staining of femoral heads in 6-month-old *β-catenin*^Col2ER^ mice. a-c: high magnification images of representative subchondral bone region. Red dotted lines: integral deformity. Red arrows: local collapse. Black arrows: empty lacunae of osteoctye. Black triangles: fat droplet. (F) Histomorphological quantitative analysis of empty lacunae rate. (G) Loading-bearing stiffness of the femoral heads in 6-month-old *β-catenin*^Col2ER^ mice.

Figure 8. Systemic injection of Wnt agonist 1 can not alleviate the GONFH-like phenotype in *β-catenin^{Col2ER}* mice. *β-catenin^{Col2ER}* mice with 5 continuous dosage of TM injections (1 mg/10 g body weight) were treated with Wnt agonist 1 three times once week until sacrifice at the age of 3 months. (A) Representative μCT images of femoral heads in Wnt agonist 1 treated *β-catenin^{Col2ER}* mice. (B) Quantitative analysis of BV/TV, Tb.N, Tb.Th and Tb.Sp in the femoral heads of Wnt agonist 1 treated *β-catenin^{Col2ER}* mice. (C) ABH staining of the femoral heads in Wnt agonist 1 treated *β-catenin^{Col2ER}* mice. a-b: high magnification images of subchondral bone region. Black triangles: fat droplet. (D) Histomorphological quantitative analysis of trabecular bone area and fat droplet area in the femoral heads of Wnt agonist 1 treated *β-catenin^{Col2ER}* mice. (E) Histomorphological quantitative analysis of empty lacunae rate in the femoral heads of Wnt agonist 1 treated *β-catenin^{Col2ER}* mice. (F, G) IHC staining and quantitative analysis of ALP and FABP4 in the femoral heads of Wnt agonist 1 treated *β-catenin^{Col2ER}* mice.