Title

Auditory evoked-potential abnormalities in a mouse model of 22q11.2 Deletion Syndrome and their interactions with hearing impairment

Authors

Chen Lu[1] and Jennifer F. Linden[1,2]

1 Ear Institute, University College London, London, WC1X 8EE, UK

2 Department of Neuroscience, Physiology, & Pharmacology, University College London, London WC1E 6BT, UK

Corresponding Author

Jennifer F. Linden (j.linden@ucl.ac.uk)

Short title

AEP abnormalities in a 22q11.2DS mouse model

Keywords

22q11.2DS, *Df1/+* mouse, hearing loss, auditory evoked potential

Abstract

Background

The 22q11.2 chromosomal microdeletion is one of the strongest known genetic risk factors for schizophrenia; ~30% of carriers develop schizophrenia in adulthood. Up to 60% of 22q11.2 deletion carriers also have mild to moderate hearing impairment, primarily from chronic middle-ear inflammation. The *Df1*/+ mouse model of the 22q11.2 deletion replicates the large inter-individual variation in hearing ability observed among deletion carriers. Here we used the *Df1/+* mouse model of the 22q11.2 deletion to investigate how genetic risk for schizophrenia and experience of hearing impairment interact to affect auditory brain function.

Methods

We measured peripheral hearing sensitivity and cortical auditory evoked potentials (AEPs) in *Df1/+* mice and their WT littermates, exploiting large inter-individual variation in hearing ability among *Df1/+* mice to distinguish cortical effects of genetic background from those of experience with hearing impairment. We quantified gain and adaptation of central auditory responses to repeated tone presentations by comparing brainstem and cortical tone-evoked potentials and by analysing the growth of cortical AEPs as tone intensity level or inter-tone interval duration increased.

Results

Central auditory gain (ratio of tone-evoked cortical and brainstem activity) was elevated in *Df1/+* mice with hearing impairment. Growth of cortical AEPs with increasing sound level was abnormally large in *Df1/+* mice regardless of hearing impairment. Finally, growth of cortical AEPs with increasing inter-tone interval was abnormal only for *Df1/+* mice without hearing impairment.

Conclusions

Auditory brain abnormalities in 22q11.2DS are a function of both genotype and hearing phenotype.

Introduction

22q11.2 Deletion Syndrome (22q11.2DS) has been extensively utilised as a model to study the aetiology of schizophrenia. It is the most frequent chromosomal microdeletion syndrome (1) and one of the strongest cytogenetic risk factors for schizophrenia (2). Nearly 30% of 22q11.2 deletion carriers develop schizophrenia during their lifetime, and 1% of schizophrenia patients are estimated to have the 22q11.2 deletion (3–5). Importantly, identified schizophrenia symptoms are indistinguishable between schizophrenia patients with the 22q11.2 deletion and those with idiopathic schizophrenia (4, 6), so insights gained from studying brain abnormalities in 22q11.2DS might have more general application to understanding schizophrenia-related brain abnormalities. Mouse models of 22q11.2 deletion are also well established, enabling researchers to investigate, at the mechanistic level, what brain abnormalities arise from this genetic risk factor for schizophrenia and how experiential risk factors might affect their development (7).

One experiential risk factor for schizophrenia that may be particularly relevant for 22q11.2DS patients is hearing impairment. Up to 60% of 22q11.2 deletion carriers have mild to moderate hearing impairment, primarily arising from chronic middle ear inflammation (8, 9). Hearing impairment is a recognized experiential risk factor for schizophrenia in the general population (10), but its possible contribution to development of schizophrenia in 22q11.2DS patients has not been studied. Surprisingly, the influence of hearing impairment on auditory evoked-potential (AEP) abnormalities in 22q11.2DS patients has not been systematically examined either, even though AEP abnormalities are well-documented in 22q11.2DS and schizophrenia patients (11–15).

Here we investigated whether AEP abnormalities in the *Df1/+* mouse model of 22q11.2DS depend only upon genotype or are affected by hearing phenotype. *Df1/+* mice have a 1.2 Mb microdeletion homologous to the minimal 1.5 Mb deletion in 22q11.2DS (16) and replicate multiple physiological abnormalities observed in 22q11.2DS patients (17–20), including high inter-individual variation in hearing sensitivity. Like 22q11.2DS patients, *Df1/+* mice are susceptible to chronic middle-ear problems, and up to 60% have mild to moderate hearing impairment in one or both ears (21, 22). Moreover, previous work has suggested that auditory brain abnormalities in *Df1/+* mice correlate with hearing impairment (23). Density of parvalbumin-immunoreactive inhibitory interneurons in the auditory cortex was abnormally low in *Df1/+* mice with hearing impairment, and gain of click-evoked cortical AEPs was abnormally high. No significant differences in AEPs were observed between *Df1/+* mice with normal hearing and their WT littermates; however, conclusions were limited by the use of a broadband click stimulus at a single sound level and repetition rate.

In the current study, we first conducted a comprehensive review of methods for AEP measurement in mice, which revealed striking diversity in approaches. Adopting one of the simplest and best-validated

strategies, we then investigated AEP abnormalities in *Df1/+* mice in detail, using tones to rule out frequency-dependent confounds that constrain interpretation of click-evoked AEPs. In particular, we studied how AEP abnormalities in *Df1/+* mice with and without hearing impairment were affected by varying tone intensity level and inter-tone interval duration. Our data reveal that some AEP abnormalities in *Df1/+* mice are independent of hearing impairment and others depend on degree of hearing impairment. The results point to specific AEP measures that could be reliable biomarkers for auditory brain abnormalities in 22q11.2DS patients, robust to individual differences in degree of hearing impairment.

Methods and Materials

Animals

Experiments were conducted in 29 *Df1/+* mice (18 females, 11 males) and 22 WT littermates (8 females, 14 males). *Df1/+* mice were 8.6 to 13.3 weeks old with a mean age of 10.3 ± 1.4 weeks, and WT mice were 8.6 to 13.3 weeks old with a mean age of 10.2 ± 1.3 weeks. All experiments were performed in accordance with a Home Office project licence approved under the United Kingdom Animal Scientific Procedures Act of 1986.

Auditory stimuli

Click-evoked auditory brainstem responses (ABRs): stimuli were 50 μs monophasic clicks ranging in sound level from 20 to 90 dB SPL in 5 dB steps, repeated 500 times at each sound level with an inter-onset interval of 50 ms.

Fixed-intensity auditory evoked potentials (AEPs) and tone-evoked ABR: stimuli were 5 ms 16 kHz tones at 80 dB SPL, repeated 1000 times at an inter-tone interval (ITI) of 300 ms.

Level-dependent AEP growth functions: stimuli were 5 ms 16 kHz tones at 70, 80, 90, or 100 dB SPL, repeated 1000 times for each sound level with an ITI of 300 ms.

Time-dependent AEP growth functions: stimuli were 5 ms 16 kHz tones at 80 dB SPL, repeated 1000 times at ITIs of 200, 250, 300, 350 or 450 ms.

See Supplementary Information for further details.

Recording procedures

Recording procedures were as described previously (23). Briefly, ketamine/medetomidine-anesthetized mice were oriented with the tested ear directed toward the speaker. The opposite ear was blocked with an earplug during the recording to ensure monaural stimulation. Auditory brainstem responses (ABRs)

were collected from electrodes at the bulla of the tested ear and the vertex. Auditory evoked potentials (AEP) were recorded single-ended from the electrode over the contralateral auditory cortex relative to the stimulated ear.

Analysis

ABR signals were filtered with a 100-3000 Hz band-pass filter before subtraction (vertex minus bulla electrode) to obtain the differential ABR signal. The hearing threshold was defined as the lowest sound intensity level with a characteristic ABR wave deflection at least twice as large as the time-dependent standard error in the mean ABR waveform. The baseline and peak of ABR wave 1 were manually labeled on averaged waveforms to measure ABR wave 1 amplitude.

Three key deflections of the AEP waveform (P1, N1, and P2) were manually selected from the averaged waveform after removing heartbeat noise (24). P1 is the highest deflection between 15 to 30 ms post stimulus onset; N1 is the lowest deflection between 25 to 60 ms post stimulus onset; and P2 is the highest deflection between 60 to 120 ms post stimulus onset.

The data were statistically analyzed and graphically visualized using Python 3. Non-parametric statistical methods were adopted (Mann-Whitney U rank test, Kruskal-Wallis H test with post-hoc Dunn's test, and Spearman's correlation test). All significance tests were conducted two-tailed with α = 0.05, and exact p-values are stated where significant but greater than 0.001.

See Supplementary Information for more details on animal breeding and maintenance, acoustic stimulation, experimental procedures, and data processing.

Results

Comprehensive literature review reveals a need for standardized approaches to auditory-evoked potential measurement in mice

Procedures for auditory evoked-potential (AEP) recording are relatively well established for human studies, but there is no standardized AEP recording paradigm for mice. To find out how AEP measurement in mice has been performed in the past, we reviewed 115 papers that were obtained from Web of Science using search terms 'auditory evoked potentials', 'event-related potentials', 'mouse', 'rodent'. Papers that only covered auditory brainstem responses (ABR, also sometimes phased as 'early AEP') were excluded. There was surprisingly little consistency between approaches used by unaffiliated labs. After consolidating the search results to eliminate papers using repeated methods, we still had 50 publications describing different approaches to AEP recording in mice. Electrode locations varied from study to study; some used single or symmetric electrode recording from one brain region and others recorded from multiple brain areas simultaneously (Figure 1A). Clustering all positive electrode locations

across studies, we found that in the majority of previous studies, positive electrodes had been placed above auditory cortex, hippocampus and/or frontal areas (Figure 1B), while reference electrodes were typically located either at the occipital bone or over the frontal/olfactory lobe.

In the absence of a standardized approach to mouse AEP recording, we adopted the simplest strategy compatible with future measurement of intracranial auditory cortical activity correlating with the AEP. Specifically, we placed a single subdermal positive electrode above the auditory cortex contralateral to the monaurally stimulated ear, and a reference electrode over the frontal/olfactory lobe (25–28). Our definitions of the AEP wave deflections P1, N1 and P2 are consistent with definitions used in those studies (29). We note also that the same electrode positioning with reversed polarity was used in one of the most detailed characterizations of subcutaneous cortical AEPs in mice to date (30). Aside from the reversal of polarity, our AEP waveforms were consistent with those recorded by Postal et al. 2022, and our P1, N1 and P2 correspond to N23, P35 and N61 from their study.

We used this simple and well-validated approach to AEP measurement in mice to quantify AEP abnormalities in the *Df1/+* mouse model of 22q11.2DS, for comparison with AEP abnormalities observed in 22q11.2DS patients (12). We focused particularly on a question that has been largely overlooked in previous studies of 22q11.2DS patients and 22q11.2DS model mice: are AEP abnormalities in *Df1/+* mice related to hearing impairment, which is a common comorbidity of 22q11.2DS?

High inter-individual and inter-ear variation in hearing sensitivity in *Df1/+* mice necessitates ear-by-ear analysis of effects of hearing impairment

We first confirmed previous findings that *Df1/+* mice replicate the high inter-individual and inter-ear variation in hearing sensitivity also observed in 22q11.2DS patients (9, 21, 23). We quantified peripheral hearing sensitivity of each ear in each mouse using the click-evoked ABR threshold. ABRs were measured with subdermal electrodes (Figure 2A) during presentations of click stimuli varying in sound level from 20dB to 90dB SPL. The hearing threshold of an ear was defined as the lowest click intensity level that elicited at least one clear characteristic ABR waveform deflection (Figure 2B).

Hearing thresholds of *Df1/+* ears were bimodally distributed (Figure 2C), as demonstrated previously (21, 23). On average, hearing thresholds for *Df1/+* ears were significantly elevated compared to those from their WT littermates (*Df1/+*, 54.02 ± 14.41 dB SPL from 56 ears, WT, 35.23 ± 4.64 dB SPL from 44 ears; Wilcoxon rank-sum test, p < 0.001), but many *Df1/+* ears had WT-like hearing thresholds. Here, we defined an ear as exhibiting hearing impairment if its hearing threshold was more than 2.5 standard deviations above the mean hearing threshold for WT ears (47 dB SPL). Then we grouped *Df1/+* ears into "normal hearing" (*Df1/+* NH, 23 ears) and "hearing impaired" (*Df1/+* HI, 33 ears).

We adopted individual ears rather than mice as the unit for analysis because there was no significant correlation between the hearing thresholds of the two ears in *Df1/+* mice (Spearman's rank correlation, ρ = 0.31, p = 0.12). Among all *Df1/+* mice tested in both ears, 26% (7/27) had normal hearing in both ears, 33% (9/27) had unilateral hearing impairment (6 in right ear, 3 in left ear), and 41% (11/27) had bilateral hearing impairment (Figure 2D).

Tone-evoked auditory cortical potentials and brainstem responses confirm increased central auditory gain in *Df1/+* mice with hearing impairment

We recorded tone-evoked AEPs simultaneously with ABRs to test the hypothesis that central auditory gain is elevated in *Df1/+* mice with hearing impairment. A previous study had concluded that *Df1/+* mice with hearing impairment exhibited increased central auditory gain, based on the observation that the ratio of click-evoked AEP to ABR magnitude was abnormally elevated for *Df1/+* ears with hearing impairment (23). This conclusion is consistent with a substantial literature demonstrating that hearing impairment increases central auditory excitability (31). However, it is possible that alterations in click-evoked AEP/ABR ratio with hearing impairment could arise simply from the different frequency sensitivities of the ABR versus AEP generators (for example, if hearing impairment primarily affected high sound frequencies, which drive the ABR more strongly than the AEP), not from changes in central auditory gain.

To rule out this possible alternative explanation, we investigated the relationship between hearing impairment and the AEP/ABR ratio measure of central auditory gain using 80 dB SPL, 16 kHz tones instead of clicks, and confirmed that elevated central auditory gain is evident in tone-evoked brain activity in *Df1/+* mice. For this analysis, clicks were used only for categorizing *Df1/+* ears as normal hearing or hearing impaired based on click-evoked ABR threshold, not for analysis of central auditory gain. As expected given the click-evoked ABR thresholds, the amplitude of ABR wave 1 evoked by a 80 dB SPL, 16 kHz tone was much smaller for *Df1/+* HI ears than for *Df1/+* NH or WT ears (Figure 3A and C; Kruskal-Wallis test, $p < 0.001$, post-hoc tests, $p_{Df1/+ \text{ NH-}Df1/+ \text{ HI}} < 0.001$, $p_{Df1/+ \text{ NH-WT}} = 0.98$, $p_{Df1/+ \text{ HI-WT}} < 0.001$). However, there was no corresponding drop in the amplitude of cortical AEP waveforms evoked at *Df1/+* HI ears compared to WT ears, and tone stimulation of *Df1/+* NH ears produced AEPs with *larger* P1-N1 amplitude than for WT ears (Figure 3B, D and E; Kruskal-Wallis test, P1-N1, $p = 0.0015$, post-hoc tests, $p_{Df1/+ \text{ NH-}Df1/+ \text{ HI}} = 0.017$, $p_{Df1/+ \text{ NH-WT}} = 0.017$, $p_{Df1/+ \text{ HI-WT}} = 0.79$; N1-P2, $p = 0.034$, post-hoc tests, $p_{Df1/+ \text{ NH-}Df1/+ \text{ HI}} = 0.20$, $p_{Df1/+ \text{ NH-WT}} = 0.20$, $p_{Df1/+ \text{ HI-WT}} = 0.83$). The gain of tone-evoked auditory brain activity was elevated in *Df1/+* mice, not only during stimulation of *Df1/+* HI ears but also (for the P1-N1 complex) during stimulation of *Df1/+* NH ears (Figure 3F and G; Kruskal-Wallis tests, P1-N1: $p < 0.001$, post-hoc tests, $p_{Df1/+ \text{ NH-}Df1/+ \text{ HI}} = 0.083$, $p_{Df1/+ \text{ NH-WT}} = 0.043$, $p_{Df1/+ \text{ HI-WT}} < 0.001$; N1-P2, $p < 0.001$, post-hoc tests, $p_{Df1/+ \text{ NH-}Df1/+ \text{ HI}} = 0.037$, $p_{Df1/+ \text{ NH-WT}} = 0.11$, $p_{Df1/+ \text{ HI-WT}} < 0.001$).

Further analysis demonstrated that central auditory gain in *Df1/+* mice was positively correlated with the degree of hearing impairment in the stimulated ear. We examined the relationship between click-evoked hearing thresholds and tone-evoked ABR wave 1 amplitude, AEP P1-N1 or N1-P2 amplitude, or AEP/ABR ratio measures of central auditory gain in *Df1/+* mice. Unsurprisingly, the amplitude of tone-evoked ABR wave 1 was negatively correlated with click-evoked hearing threshold in the stimulated ear (Figure 3G; Spearman's rank correlation test, ρ = -0.68, p < 0.001). Amplitudes of tone-evoked P1-N1 and N1-P2 complexes measured over the contralateral auditory cortex were also negatively correlated with click-evoked hearing thresholds (Figure 3H-I; P1-N1 complex: ρ = -0.45, p = 0.0015; N1-P2 complex: ρ = -0.31, p = 0.034). However, there were robust positive correlations between tone-evoked central auditory gain measures (AEP/ABR ratios) and click-evoked hearing thresholds, for both the P1-N1 and N1-P2 complex (Figure K-L; P1-N1 complex AEP/ABR: ρ = 0.33, p = 0.024; N1-P2 complex AEP/ABR: ρ = 0.43, p = 0.0024).

Interestingly, we also found that latencies of late tone-evoked AEP waves were longer in *Df1/+* than WT mice, regardless of hearing impairment (Supplementary Figure 1). Tone stimulation of *Df1/+* ears with or without hearing impairment evoked N1 and P2 waves with significantly longer latencies than those evoked in WT mice, while P1 wave latencies were not significantly different (Supplementary Figure 1A-C). There was no significant correlation between AEP wave latencies in *Df1/+* mice and the click-evoked hearing threshold in the stimulated ear for any of the three AEP waves (Supplementary Figure 1D-F). Thus, tone-evoked N1 and P2 wave latencies were significantly longer for *Df1/+* than WT mice, even when the stimulated *Df1/+* ears had normal hearing thresholds.

Higher level-dependent AEP growth in *Df1/+* mice, with or without hearing impairment

Next, we tested another stimulus paradigm that might reveal AEP abnormalities in *Df1/+* mice: a tone stimulus varying in sound intensity level (16kHz, 70 to 100 dB SPL). A previous study using a similar stimulus paradigm reported steeper level-dependent AEP growth in the *Df(h22q11)/+* mouse model of 22q11.2DS than in WT mice (25). Here, we compared level-dependent AEP (LDAEP) growth functions not only between *Df1/+* mice and their WT littermates, but also among *Df1/+* mice, analyzing effects of hearing impairment in the stimulated ear.

We found that AEP wave amplitudes grew more steeply with increasing sound level in *Df1/+* mice than WT mice, regardless of whether the stimulated *Df1/+* ear was normal hearing or hearing impaired (Figure 4). This effect was clearly evident both in example AEP waveform recordings and in group averages (Figure 4A-B). We calculated the AEP amplitude as a linear function of sound level and used the function slope to represent the level-dependence of the AEP (Figure 4C and E). For both the P1-N1 and N1-P2 complexes, the LDAEP amplitude change was larger in *Df1/+* mice than in their WT littermates (Figure 4D and F; P1-N1: Kruskal-Wallis test, p < 0.001, post-hoc tests, $p_{Df1/+ NH-Df1/+ HI}$ = 0.092, $p_{Df1/+ NH-WT}$ < 0.001,

p$_{Df1/+ \text{ HI-WT}}$ = 0.040; N1-P2: Kruskal-Wallis test, p = 0.0020, post-hoc tests, p$_{Df1/+ \text{ NH-}Df1/+ \text{ HI}}$ = 0.75, p$_{Df1/+ \text{ NH-WT}}$ = 0.022, p$_{Df1/+ \text{ HI-WT}}$ = 0.0042). There were no significant differences in LDAEP amplitude change for data obtained during stimulation of *Df1/+* NH versus *Df1/+* HI ears, suggesting that level-dependent AEP amplitude change could be a biomarker for central auditory abnormalities in *Df1/+* mice that are robust to hearing impairment.

Level-dependent changes in AEP wave latencies were less robust as a potential biomarker for abnormalities in *Df1/+* mice (Supplementary Figure 2). AEP P1 and N1 latencies decreased more steeply with increasing tone intensity level in *Df1/+* mice compared to their WT littermates, but effects were not consistently significant for the two waves across *Df1/+* NH and *Df1/+* HI ears. For P2, there were no significant differences between the groups in the relationship between wave latency and sound level.

Higher inter-stimulus-interval time-dependent AEP growth in *Df1/+* mice with normal hearing

A recent study of auditory processing endophenotypes in 22q11.2DS patients showed that tone-evoked AEP amplitudes were more strongly dependent on inter-tone interval (ITI) in patients without psychotic symptoms than in either patients with psychotic symptoms or neurotypical controls (12). We wondered if similar abnormalities might be evident in *Df1/+* mice, and if so, how these abnormalities might relate to hearing impairment. We measured the sensitivity of the AEP to changes in the temporal structure of the stimulus (time-dependent AEP, TDAEP) using 16kHz, 80 dB SPL tones presented at varying ITIs.

At first we used ITIs varying from 200 ms to 350 ms (Supplementary Figure 3), and observed that AEP amplitude increased with ITI to a plateau in many cases. The turning point for the plateau in WT and *Df1/+* HI data was between 300 and 350 ms, while the turning point for *Df1/+* NH data was not clear and could have been at longer ITIs. Therefore, we extended the range of ITIs to 200-450 ms in further experiments.

We found that TDAEP amplitude change was significantly larger for *Df1/+* NH recordings than for either WT or *Df1/+* HI recordings (Figure 5). To quantify TDAEP amplitude change, we calculated the best-fit linear function relating AEP amplitude to the logarithm of ITI duration (Figure 5C and E; note logarithmic scale of x-axis). We used the slope of this function as our measure of TDAEP amplitude change (Figure 5C-F). For both the P1-N1 and N1-P2 complexes, TDAEP amplitude change was abnormally large in *Df1/+* NH recordings, relative to either WT data or *Df1/+* HI data (Figure 5D and F; P1-N1: Kruskal-Wallis test, p < 0.001, post-hoc tests, p$_{Df1/+ \text{ NH-}Df1/+ \text{ HI}}$ = 0.060, p$_{Df1/+ \text{ NH-WT}}$ < 0.001, p$_{Df1/+ \text{ HI-WT}}$ = 0.069; N1-P2: p = 0.0010, post-hoc tests, p$_{Df1/+ \text{ NH-}Df1/+ \text{ HI}}$ = 0.10, p$_{Df1/+ \text{ NH-WT}}$ < 0.001, p$_{Df1/+ \text{ HI-WT}}$ = 0.10). This result indicates that central auditory adaptation to repeated tones differs between *Df1/+* NH and WT mice, and suggests that hearing impairment in *Df1/+* HI mice may either mask or counteract this abnormality.

We also examined the relationship between AEP wave latency and ITI, comparing *Df1/+* NH, *Df1/+* HI, and WT data. For the P1, N1, and P2 waves, latency decreased similarly with increasing ITIs in all groups (Supplementary Figure 4). There were no significant group differences in the relationship between AEP wave latency and ITI for P1 or P2, and for N1, a weak group difference was not robust to post-hoc testing.

Discussion

Our results reveal multiple abnormalities in cortical auditory evoked potentials (AEPs) in the *Df1/+* mouse model of 22q11.2DS and demonstrate that some of these abnormalities are strongly influenced by mild to moderate hearing impairment, a common co-morbidity of 22q11.2DS. Using one of the best-validated of the surprisingly diverse and unstandardized approaches to mouse AEP measurement in the literature, we measured tone-evoked cortical AEPs along with ABRs in both *Df1/+* mice and their WT littermates. We quantified hearing thresholds ear-by-ear in every mouse using click-evoked ABR thresholds, so that we could exploit the high inter-individual and inter-ear variation in hearing sensitivity typical of *Df1/+* mice (and 22q11.2DS patients) to probe the relationship between AEP abnormalities and hearing impairment. We found 3 key AEP abnormalities in *Df1/+* mice, which exhibited differing relationships to hearing impairment. Tone-evoked central auditory gain (AEP amplitude normalized by ABR amplitude) was elevated primarily in *Df1/+* mice with hearing impairment, and strongly correlated with the level of hearing impairment in the stimulated ear. Another measure of central auditory excitability --- growth of cortical AEP amplitude with increasing sound level --- was robustly elevated in all *Df1/+* mice, even when hearing was normal in the stimulated ear. Finally, growth of AEP amplitude with increasing inter-tone interval, a measure of recovery from repetition suppression, was abnormal in *Df1/+* mice only when hearing was normal in the tested ear. In summary, these results show that *Df1/+* mice have abnormalities in central auditory gain (AEP/ABR ratio) that are exacerbated by hearing impairment; abnormalities in central auditory excitability (level-dependent AEP growth) that are robust to hearing impairment; and abnormalities in auditory adaptation (inter-stimulus-interval time-dependent AEP growth) that are either counteracted or masked by hearing impairment.

Implications of results

Two points need to be kept in mind when considering the implications of our results.

First: the unusual nature of hearing impairment in *Df1/+* mice (and 22q11.2DS patients) means that ear-by-ear analysis of hearing sensitivity is essential, and monaural auditory stimulation for AEP measurement is advisable. Like 22q11.2DS patients, *Df1/+* mice are susceptible to chronic middle ear inflammation that can cause mild to moderate hearing impairment in both ears or in only one ear (21, 23). Hearing impairment in our cohort of *Df1/+* mice was frequently unilateral, and there was minimal correlation in hearing thresholds between the two ears. Therefore, we focused on obtaining AEP

recordings from each hemisphere in every animal during monaural stimulation of the contralateral ear. (Due the anatomy and physiology of the auditory system, monaural stimulation elicits stronger cortical AEPs contralateral than ipsilateral to the stimulated ear; see for example Postal et al. 2022 (30)). We found no evidence that AEPs measured during monaural stimulation depended on the presence or absence or hearing impairment in the unstimulated ear, justifying our approach. However, it is likely that AEPs measured during binaural stimulation would depend on hearing thresholds in both ears, increasing the variability and complexity of experimental results. Therefore, one implication of this work in *Df1/+* mice is that AEP abnormalities in 22q11.2DS patients might best be identified using monaural auditory stimulation paradigms, rather than the usual binaural stimulation paradigms (e.g., (11–13, 32)).

Second: high inter-individual and inter-ear variation in hearing sensitivity among *Df1/+* mice (and 22q11.2DS patients) complicates interpretation of AEP abnormalities, even with monaural stimulation. As in most AEP studies of 22q11.2DS patients (11–13, 32), we presented exactly the same set of auditory stimuli to each subject (and each ear). We did not adjust the sound level of the stimuli to be constant relative to the hearing threshold of the stimulated ear, because this alternative approach would have necessitated the use of extremely loud stimuli for stimulation of ears with moderate hearing impairment, undermining our ability to ensure monaural stimulation using an earplug in the opposite ear. However, the use of fixed-intensity stimuli meant that inevitably, weaker auditory nerve signals were evoked in ears with hearing impairment than in ears with normal hearing. The AEP/ABR ratio measure of central auditory gain attempts to normalize for differences in auditory nerve activity during stimulation of *Df1/+* HI versus *Df1/+* NH or WT ears, making comparisons across all three groups more valid. However, differences in auditory nerve activity are not taken into account by measures of level-dependent AEP growth (LDAEP) and inter-stimulus-interval time-dependent AEP growth (TDAEP), complicating comparisons between hearing-impaired and normal-hearing data groups. Therefore, the most compelling abnormalities in LDAEP and TDAEP are those revealed in *Df1/+* NH versus WT comparisons: higher LDAEP growth, and higher TDAEP growth.

Biomarkers for central auditory processing abnormalities in 22q11.2DS

Interestingly, similar LDEAP and TDAEP abnormalities have been mentioned previously in studies of 22q11.2DS models or patients. Didriksen et al. (2017) reported higher LDAEP growth in the *Df(h22q11)/+* mouse model of 22q11.2DS than in WT littermates. Notably, hearing thresholds were assessed ear-by-ear using ABR in a separate cohort of these mice and were not significantly different between *Df(h22q11)/+* and WT animals, although unusually high for WT mice (25). Francisco et al. (2020) (12) found higher TDAEP growth in 22q11.2DS patients without psychotic symptoms compared to neurotypical age-matched controls (although no significant differences were observed between patients with psychotic symptoms and controls). Participants were included in that study only if they passed an audiometric test to ensure that hearing thresholds were no more than 25 dB above normal in both ears.

Together with our own data, these two studies suggest that both LDAEP growth and TDAEP growth are abnormally elevated in 22q11.2DS, at least in the absence of hearing impairment. Our work further demonstrates that elevated LDAEP growth may be a particularly useful biomarker for central auditory processing abnormalities in 22q11.2DS, because this AEP abnormality is robust to hearing impairment in the stimulated ear.

What mechanisms might underlie the AEP abnormalities we observed in *Df1/+* mice? From a systems-level perspective, increased central auditory gain and LDAEP growth suggest increased central auditory excitability, and increased TDAEP growth suggests altered adaptation. Hearing impairment is already known to cause increased excitability and decreased inhibitory synaptic transmission in the auditory cortex (31, 33–36). Moreover, previous work has found that parvalbumin-expressing (PV) inhibitory interneurons, which play a critical role in auditory cortical gain control (37), are abnormally sparse in the auditory cortex of *Df1/+* mice --- and this PV interneuron deficit correlates with degree of hearing impairment (23). More generally, deficits in GABAergic signaling have been reported in the prefrontal cortex and hippocampus in other mouse models of 22q11.2DS (38, 39), and human 22q11.2DS cerebral cortical organoids have been found to display high resting membrane potential and increased spontaneous firing (40). Therefore, it is possible that both the 22q11.2 deletion and hearing impairment alter auditory cortical excitability and adaptation, and that auditory brain abnormalities in 22q11.2DS arise from interactions between genetic vulnerability and experience of hearing impairment.

Conclusions

We found multiple abnormalities in cortical auditory evoked potentials (AEPs) in the *Df1/+* mouse model of 22q11.2DS, some of which were strongly influenced by hearing impairment, a common co-morbidity of 22q11.2DS. *Df1/+* mice exhibited 3 key AEP abnormalities: (1) increased central auditory gain, which was especially pronounced for stimulation of ears with hearing impairment; (2) increased growth of AEP amplitude with sound level, regardless of the presence or absence of hearing impairment in the stimulated ear; and (3) increased growth of AEP amplitude with inter-stimulus-interval time duration when hearing was normal in the stimulated ear. Our results indicate that level-dependent AEP growth may be especially useful as a biomarker for auditory brain abnormalities in 22q11.2DS, because this measure was robust to hearing impairment in the stimulated ear. Moreover, the results suggest that both the 22q11.2 deletion and experience of hearing impairment drive changes in central auditory function that produce AEP abnormalities.

Acknowledgements

We thank Dr. Alain de Cheveigne for advising on data analysis. We thank Dr. Pedro J. Gonçalves for discussion on the comparison between experimental data and modelling data. We would also like to

acknowledge Ashviniy Thamilmaran for literature data collection, and infrastructure support from the NIHR UCLH BRC Hearing Health Theme. This research was funded by the UCL Institute for Mental Health (Small Grant 2022 to JFL) and the UK Medical Research Council (MR/P006221/1 to JFL).

Disclosures

The authors report no conflicts of interest.

Other information

Confirmation of references for Figure 1, for article proof: #1 (41); #2 (42); #3 (43); #4 (44); #5 (45); #6 (46); #7 (47); #8 (48); #9 (49); #10 (50); #11 (51); #12 (52); #13 (53); #14 (54); #15 (27); #16 (26); #17 (25); #18 (55); #19 (56); #20 (28); #21 (57); #22 (30); #23 (58); #24 (59); #25 (60); #26 (61); #27 (62); #28 (63); #29 (64); #30 (65); #31 (66); #32 (67); #33 (68); #34 (69); #35 (70); #36 (71); #37 (72); #38 (73); #39 (74); #40 (75); #41 (76); #42 (77); #43 (78); #44 (79); #45 (80); #46 (81); #47 (82); #48 (83); #49 (84); #50 (85)

Reference

1. McDonald-McGinn DM, Sullivan KE, Marino B, Philip N, Swillen A, Vorstman JAS, *et al.* (2015): 22q11.2 deletion syndrome. *Nat. Rev. Dis. Primers*. 1: 15071.

2. Singh T, Poterba T, Curtis D, Akil H, Al Eissa M, Barchas JD, *et al.* (2022): Rare coding variants in ten genes confer substantial risk for schizophrenia. *Nature.* 604(7906): 509–16.

3. Schneider M, Debbané M, Bassett AS, Chow EWC, Fung WLA, van den Bree M, *et al.* (2014): Psychiatric disorders from childhood to adulthood in 22q11.2 deletion syndrome: results from the international consortium on brain and behavior in 22q11.2 deletion syndrome. *Am. J. Psychiatry*. 171(6): 627–39.

4. Monks S, Niarchou M, Davies AR, Walters JTR, Williams N, Owen MJ, *et al.* (2014): Further evidence for high rates of schizophrenia in 22q11.2 deletion syndrome. *Schizophr. Res.* 153(1–3): 231–36.

5. Bassett AS, Costain G, Fung WLA, Russell KJ, Pierce L, Kapadia R, *et al.* (2010): Clinically detectable copy number variations in a canadian catchment population of schizophrenia. *J. Psychiatr. Res.* 44(15): 1005–9.

6. Bassett AS, Chow EWC, AbdelMalik P, Gheorghiu M, Husted J, Weksberg R (2003): The schizophrenia phenotype in 22q11 deletion syndrome. *Am. J. Psychiatry*. 160(9): 1580–86.

7. Meechan DW, Maynard TM, Tucker ES, Fernandez A, Karpinski BA, Rothblat LA, *et al.* (2015): Modeling a model: mouse genetics, 22q11.2 deletion syndrome, and disorders of cortical circuit development. *Prog. Neurobiol.* 130: 1–28.

8. Van Eynde C, Swillen A, Lambeens E, Verhaert N, Desloovere C, Luts H, *et al.* (2016): Prevalence and nature of hearing loss in 22q11.2 deletion syndrome. *J. Speech Lang. Hear. Res.* 59(3): 583–89.

9. Verheij E, Derks LSM, Stegeman I, Thomeer HGXM (2017): Prevalence of hearing loss and clinical otologic manifestations in patients with 22q11.2 deletion syndrome: a literature review. *Clin. Otolaryngol.* 42(6): 1319–28.

10. Linszen MMJ, Brouwer RM, Heringa SM, Sommer IE (2016): Increased risk of psychosis in patients with hearing impairment: review and meta-analyses. *Neurosci. Biobehav. Rev.* 62: 1–20.

11. Larsen KM, Pellegrino G, Birknow MR, Kjær TN, Baaré WFC, Didriksen M, *et al.* (2018): 22q11.2 deletion syndrome is associated with impaired auditory steady-state gamma response. *Schizophr. Bull.* 44(2): 388–97.

12. Francisco AA, Foxe JJ, Horsthuis DJ, DeMaio D, Molholm S (2020): Assessing auditory processing endophenotypes associated with schizophrenia in individuals with 22q11.2 deletion syndrome. *Transl. Psychiatry*. 10(1): 85.

13. Cantonas L-M, Mancini V, Rihs TA, Rochas V, Schneider M, Eliez S, *et al.* (2021): Abnormal auditory processing and underlying structural changes in 22q11.2 deletion syndrome. *Schizophr. Bull.* 47(1): 189–96.

14. Cantonas L-M, Tomescu MI, Biria M, Jan RK, Schneider M, Eliez S, *et al.* (2019): Abnormal development of early auditory processing in 22q11.2 deletion syndrome. *Transl. Psychiatry*. 9(1): 138.

15. Larsen KM, Mørup M, Birknow MR, Fischer E, Olsen L, Didriksen M, *et al.* (2019): Individuals with 22q11.2 deletion syndrome show intact prediction but reduced adaptation in responses to repeated sounds: evidence from bayesian mapping. *Neuroimage Clin.* 22: 101721.

16. Lindsay EA, Botta A, Jurecic V, Carattini-Rivera S, Cheah YC, Rosenblatt HM, *et al.* (1999): Congenital heart disease in mice deficient for the digeorge syndrome region. *Nature.* 401(6751): 379–83.

17. Paylor R, Lindsay E (2006): Mouse models of 22q11 deletion syndrome. *Biol. Psychiatry*. 59(12): 1172–79.

18. Paylor R, McIlwain KL, McAninch R, Nellis A, Yuva-Paylor LA, Baldini A, *et al.* (2001): Mice deleted for the digeorge/velocardiofacial syndrome region show abnormal sensorimotor gating and learning and memory impairments. *Hum. Mol. Genet.* 10(23): 2645–50.

19. Sivagnanasundaram S, Fletcher D, Hubank M, Illingworth E, Skuse D, Scambler P (2007): Differential gene expression in the hippocampus of the df1/+ mice: a model for 22q11.2 deletion syndrome and schizophrenia. *Brain Res.* 1139: 48–59.

20. Sumitomo A, Horike K, Hirai K, Butcher N, Boot E, Sakurai T, *et al.* (2018): A mouse model of 22q11.2 deletions: molecular and behavioral signatures of parkinson's disease and schizophrenia. *Sci. Adv.* 4(8): eaar6637.

21. Fuchs JC, Zinnamon FA, Taylor RR, Ivins S, Scambler PJ, Forge A, *et al.* (2013): Hearing loss in a mouse model of 22q11.2 deletion syndrome. *PLoS ONE*. 8(11): e80104.

22. Fuchs JC, Linden JF, Baldini A, Tucker AS (2015): A defect in early myogenesis causes otitis media in two mouse models of 22q11.2 deletion syndrome. *Hum. Mol. Genet.* 24(7): 1869–82.

23. Zinnamon FA, Harrison FG, Wenas SS, Liu Q, Wang KH, Linden JF (2022): Increased central auditory gain

and decreased parvalbumin-positive cortical interneuron density in the df1/+ mouse model of schizophrenia correlate with hearing impairment. *Biological Psychiatry Global Open Science*

24.   de Cheveigné A (2016): Sparse time artifact removal. *J. Neurosci. Methods*. 262: 14–20.

25.   Didriksen M, Fejgin K, Nilsson SRO, Birknow MR, Grayton HM, Larsen PH, *et al.* (2017): Persistent gating deficit and increased sensitivity to nmda receptor antagonism after puberty in a new mouse model of the human 22q11.2 microdeletion syndrome: a study in male mice. *J. Psychiatry Neurosci.* 42(1): 48–58.

26.   Lovelace JW, Wen TH, Reinhard S, Hsu MS, Sidhu H, Ethell IM, *et al.* (2016): Matrix metalloproteinase-9 deletion rescues auditory evoked potential habituation deficit in a mouse model of fragile x syndrome. *Neurobiol. Dis.* 89: 126–35.

27.   Aedo C, Terreros G, León A, Delano PH (2016): The corticofugal effects of auditory cortex microstimulation on auditory nerve and superior olivary complex responses are mediated via alpha-9 nicotinic receptor subunit. *PLoS ONE*. 11(5): e0155991.

28.   Khani A, Lanz F, Loquet G, Schaller K, Michel C, Quairiaux C (2019): Large-scale networks for auditory sensory gating in the awake mouse. *eNeuro*. 6(5):

29.   Maxwell CR, Liang Y, Weightman BD, Kanes SJ, Abel T, Gur RE, *et al.* (2004): Effects of chronic olanzapine and haloperidol differ on the mouse n1 auditory evoked potential. *Neuropsychopharmacology*. 29(4): 739–46.

30.   Postal O, Bakay W, Dupont T, Buck A, Élodie Daoud, Petit C, *et al.* (2022): Characterizing subcutaneous cortical auditory evoked potentials in mice. *Hear. Res.* 422: 108566.

31.   Persic D, Thomas ME, Pelekanos V, Ryugo DK, Takesian AE, Krumbholz K, *et al.* (2020): Regulation of auditory plasticity during critical periods and following hearing loss. *Hear. Res.* 397: 107976.

32.   Rihs TA, Tomescu MI, Britz J, Rochas V, Custo A, Schneider M, *et al.* (2013): Altered auditory processing in frontal and left temporal cortex in 22q11.2 deletion syndrome: a group at high genetic risk for schizophrenia. *Psychiatry Res.* 212(2): 141–49.

33.   Sarro EC, Kotak VC, Sanes DH, Aoki C (2008): Hearing loss alters the subcellular distribution of presynaptic gad and postsynaptic gabaa receptors in the auditory cortex. *Cereb. Cortex*. 18(12): 2855–67.

34.   Kotak VC, Takesian AE, Sanes DH (2008): Hearing loss prevents the maturation of gabaergic transmission in the auditory cortex. *Cereb. Cortex*. 18(9): 2098–2108.

35.   Resnik J, Polley DB (2021): Cochlear neural degeneration disrupts hearing in background noise by increasing auditory cortex internal noise. *Neuron*. 109(6): 984-996.e4.

36.   Chambers AR, Resnik J, Yuan Y, Whitton JP, Edge AS, Liberman MC, *et al.* (2016): Central gain restores auditory processing following near-complete cochlear denervation. *Neuron*. 89(4): 867–79.

37.   Gothner T, Gonçalves PJ, Sahani M, Linden JF, Hildebrandt KJ (2021): Sustained activation of pv+ interneurons in core auditory cortex enables robust divisive gain control for complex and naturalistic stimuli. *Cereb. Cortex*. 31(5): 2364–81.

38.   Al-Absi A-R, Qvist P, Okujeni S, Khan AR, Glerup S, Sanchez C, *et al.* (2020): Layers ii/iii of prefrontal cortex in df(h22q11)/+ mouse model of the 22q11.2 deletion display loss of parvalbumin interneurons and modulation of neuronal morphology and excitability. *Mol. Neurobiol.*

39.   Al-Absi A-R, Thambiappa SK, Khan AR, Glerup S, Sanchez C, Landau AM, *et al.* (2022): Df(h22q11)/+ mouse model exhibits reduced binding levels of gabaa receptors and structural and functional dysregulation in the inhibitory and excitatory networks of hippocampus. *Mol. Cell. Neurosci.* 122: 103769.

40.   Khan TA, Revah O, Gordon A, Yoon S-J, Krawisz AK, Goold C, *et al.* (2020): Neuronal defects in a human cellular model of 22q11.2 deletion syndrome. *Nat. Med.* 26(12): 1888–98.

41.   Welch TM, Church MW, Shucard DW (1985): A method for chronically recording brain-stem and cortical auditory evoked potentials from unanesthetized mice. *Electroencephalogr. Clin. Neurophysiol.* 60(1): 78–83.

42.   Frenkel M, Sherman GF, Bashan KA, Galaburda AM, LoTurco JJ (2000): Neocortical ectopias are associated with attenuated neurophysiological responses to rapidly changing auditory stimuli. *Neuroreport*. 11(3): 575–79.

43.   Tang J, Wotjak CT, Wagner S, Williams G, Schachner M, Dityatev A (2001): Potentiated amygdaloid auditory-evoked potentials and freezing behavior after fear conditioning in mice. *Brain Res.* 919(2): 232–41.

44.   Simosky JK, Stevens KE, Kem WR, Freedman R (2001): Intragastric dmxb-a, an alpha7 nicotinic agonist, improves deficient sensory inhibition in dba/2 mice. *Biol. Psychiatry*. 50(7): 493–500.

45.   Siegel SJ, Connolly P, Liang Y, Lenox RH, Gur RE, Bilker WB, *et al.* (2003): Effects of strain, novelty, and nmda blockade on auditory-evoked potentials in mice. *Neuropsychopharmacology*. 28(4): 675–82.

46.   Süer C, Dolu N, Ozesmi C (2004): The effect of immobilization stress on sensory gating in mice. *Int. J. Neurosci.* 114(1): 55–65.

47.   Kojima N, Sakamoto T, Endo S, Niki H (2005): Impairment of conditioned freezing to tone, but not to context, in fyn-transgenic mice: relationship to nmda receptor subunit 2b function. *Eur. J. Neurosci.* 21(5): 1359–69.

48.   Amann LC, Gandal MJ, Halene TB, Ehrlichman RS, White SL, McCarren HS, *et al.* (2010): Mouse behavioral endophenotypes for schizophrenia. *Brain Res. Bull.* 83(3–4): 147–61.

49.   Kyuhou S, Gemba H (2007): Fast cortical oscillation after thalamic degeneration: pivotal role of nmda receptor. *Biochem. Biophys. Res. Commun.* 356(1): 187–92.

50.   Liao W, Gandal MJ, Ehrlichman RS, Siegel SJ, Carlson GC (2012): MeCP2+/- mouse model of rtt reproduces auditory phenotypes associated with rett syndrome and replicate select eeg endophenotypes of autism spectrum disorder. *Neurobiol. Dis.* 46(1): 88–92.

51.   Moessnang C, Habel U, Schneider F, Siegel SJ (2012): The electrophysiological signature of motivational salience in mice and implications for schizophrenia. *Neuropsychopharmacology*. 37(13): 2846–54.

52.   De Rojas JO, Saunders JA, Luminais C, Hamilton RH, Siegel SJ (2013): Electroencephalographic changes following direct current deep brain stimulation of auditory cortex: a new model for investigating neuromodulation. *Neurosurgery*. 72(2): 267–75; discussion 275.

53.   Goffin D, Brodkin ES, Blendy JA, Siegel SJ, Zhou Z (2014): Cellular origins of auditory event-related potential deficits in rett syndrome. *Nat. Neurosci.* 17(6): 804–6.

54.   Fejgin K, Nielsen J, Birknow MR, Bastlund JF, Nielsen V, Lauridsen JB, *et al.* (2014): A mouse model that recapitulates cardinal features of the 15q13.3 microdeletion syndrome including schizophrenia- and epilepsy-related alterations. *Biol. Psychiatry*. 76(2): 128–37.

55.   Jones NC, Hudson M, Foreman J, Rind G, Hill R, Manning EE, *et al.* (2018): Brain-derived neurotrophic factor haploinsufficiency impairs high-frequency cortical oscillations in mice. *Eur. J. Neurosci.* 48(8): 2816–25.

56.   Jin N, Lipponen A, Koivisto H, Gurevicius K, Tanila H (2018): Increased cortical beta power and spike-wave discharges in middle-aged app/ps1 mice. *Neurobiol. Aging*. 71: 127–41.

57.   Aguilar DD, Radzik LK, Schiffino FL, Folorunso OO, Zielinski MR, Coyle JT, *et al.* (2021): Altered neural oscillations and behavior in a genetic mouse model of nmda receptor hypofunction. *Sci. Rep.* 11(1): 9031.

58.   Umbricht D, Vyssotky D, Latanov A, Nitsch R, Brambilla R, D'Adamo P, *et al.* (2004): Midlatency auditory event-related potentials in mice: comparison to midlatency auditory erps in humans. *Brain Res.* 1019(1–2): 189–200.

59.   Black MD, Stevens RJ, Rogacki N, Featherstone RE, Senyah Y, Giardino O, *et al.* (2011): AVE1625, a cannabinoid cb1 receptor antagonist, as a co-treatment with antipsychotics for schizophrenia: improvement in cognitive function and reduction of antipsychotic-side effects in rodents. *Psychopharmacology (Berl)*. 215(1): 149–63.

60.   Schuelert N, Dorner-Ciossek C, Brendel M, Rosenbrock H (2018): A comprehensive analysis of auditory event-related potentials and network oscillations in an nmda receptor antagonist mouse model using a novel wireless recording technology. *Physiol. Rep.* 6(16): e13782.

61.   Aleksandrov AA, Knyazeva VM, Volnova AB, Dmitrieva ES, Polyakova NV, Gainetdinov RR (2019): Trace amine-associated receptor 1 agonist modulates mismatch negativity-like responses in mice. *Front. Pharmacol.* 10: 470.

62.   O'Reilly JA (2019): Double-epoch subtraction reveals long-latency mismatch response in urethane-anaesthetized mice. *J. Neurosci. Methods*. 326: 108375.

63.   Singh Y, Leinonen H, Fazaludeen F, Jaronen M, Guest D, Buckley N, *et al.* (2019): Loss of cln5 leads to altered gad1 expression and deficits in interneuron development in mice. *Hum. Mol. Genet.* 28(19): 3309–22.

64.   Pan W, Lyu K, Zhang H, Li C, Chen P, Ying M, *et al.* (2020): Attenuation of auditory mismatch negativity in serotonin transporter knockout mice with anxiety-related behaviors. *Behav. Brain Res.* 379: 112387.

65.   Aleksandrov AA, Dmitrieva ES, Knyazeva VM, Simon YA, Polyakova NV, Stankevich LN, *et al.* (2022): Sensory gating in taar1 knockout mice. *J. Evol. Biochem. Physiol.* 58(4): 979–85.

66.   Hudson MR, Foreman J, Rind G, Manning EE, Jones NC, van den Buuse M (2022): Differential effects of chronic methamphetamine treatment on high-frequency oscillations and responses to acute methamphetamine and nmda receptor blockade in conscious mice. *Brain Sci.* 12(11):

67.   Li Y, Wang X, Chen J, Li Z, Yang P, Qin L (2020): Aberrant auditory steady-state response of awake mice

induced by chronic interferon-α treatment. *Front. Pharmacol.* 11: 584425.

68.   Ehlers CL, Somes C (2002): Long latency event-related potentials in mice: effects of stimulus characteristics and strain. *Brain Res.* 957(1): 117–28.

69.   Wang J, Ikonen S, Gurevicius K, Van Groen T, Tanila H (2003): Altered auditory-evoked potentials in mice carrying mutated human amyloid precursor protein and presenilin-1 transgenes. *Neuroscience.* 116(2): 511–17.

70.   Tang J, Wagner S, Schachner M, Dityatev A, Wotjak CT (2003): Potentiation of amygdaloid and hippocampal auditory-evoked potentials in a discriminatory fear-conditioning task in mice as a function of tone pattern and context. *Eur. J. Neurosci.* 18(3): 639–50.

71.   Gurevicius K, Gureviciene I, Valjakka A, Schachner M, Tanila H (2004): Enhanced cortical and hippocampal neuronal excitability in mice deficient in the extracellular matrix glycoprotein tenascin-r. *Mol. Cell. Neurosci.* 25(3): 515–23.

72.   Radek RJ, Miner HM, Bratcher NA, Decker MW, Gopalakrishnan M, Bitner RS (2006): Alpha4beta2 nicotinic receptor stimulation contributes to the effects of nicotine in the dba/2 mouse model of sensory gating. *Psychopharmacology (Berl).* 187(1): 47–55.

73.   Nagy D, Tingley FD, Stoiljkovic M, Hajós M (2015): Application of neurophysiological biomarkers for huntington's disease: evaluating a phosphodiesterase 9a inhibitor. *Exp. Neurol.* 263: 122–31.

74.   Wen TH, Lovelace JW, Ethell IM, Binder DK, Razak KA (2019): Developmental changes in eeg phenotypes in a mouse model of fragile x syndrome. *Neuroscience.* 398: 126–43.

75.   Kim B, Shin J, Kim Y, Choi JH (2020): Destruction of erp responses to deviance in an auditory oddball paradigm in amyloid infusion mice with memory deficits. *PLoS ONE.* 15(3): e0230277.

76.   Dong H-W, Erickson K, Lee JR, Merritt J, Fu C, Neul JL (2020): Detection of neurophysiological features in female r255x mecp2 mutation mice. *Neurobiol. Dis.* 145: 105083.

77.   Rumschlag JA, Razak KA (2021): Age-related changes in event related potentials, steady state responses and temporal processing in the auditory cortex of mice with severe or mild hearing loss. *Hear. Res.* 412: 108380.

78.   Gurevicius K, Lipponen A, Tanila H (2013): Increased cortical and thalamic excitability in freely moving appswe/ps1de9 mice modeling epileptic activity associated with alzheimer's disease. *Cereb. Cortex.* 23(5): 1148–58.

79.   Carreño-Muñoz MI, Chattopadhyaya B, Agbogba K, Côté V, Wang S, Lévesque M, *et al.* (2022): Sensory processing dysregulations as reliable translational biomarkers in syngap1 haploinsufficiency. *Brain.* 145(2): 754–69.

80.   Kat R, Kas MJH (2022): Largely unaffected auditory and visual sensory processing phenotypes in the evoked potentials of fmr1 ko2 mice. *Eur. J. Neurosci.* 56(8): 5260–73.

81.   Bickel S, Lipp H-P, Umbricht D (2008): Early auditory sensory processing deficits in mouse mutants with reduced nmda receptor function. *Neuropsychopharmacology.* 33(7): 1680–89.

82.   Shahriari Y, Krusienski D, Dadi YS, Seo M, Shin H-S, Choi JH (2016): Impaired auditory evoked potentials and oscillations in frontal and auditory cortex of a schizophrenia mouse model. *World J. Biol. Psychiatry.* 17(6): 439–48.

83.   Wang X, Pinto-Duarte A, Behrens MM, Zhou X, Sejnowski TJ (2015): Characterization of spatio-temporal epidural event-related potentials for mouse models of psychiatric disorders. *Sci. Rep.* 5: 14964.

84.   Ahnaou A, Moechars D, Raeymaekers L, Biermans R, Manyakov NV, Bottelbergs A, *et al.* (2017): Emergence of early alterations in network oscillations and functional connectivity in a tau seeding mouse model of alzheimer's disease pathology. *Sci. Rep.* 7(1): 14189.

85.   Marton TF, Seifikar H, Luongo FJ, Lee AT, Sohal VS (2018): Roles of prefrontal cortex and mediodorsal thalamus in task engagement and behavioral flexibility. *J. Neurosci.* 38(10): 2569–78.

Figures

Figure 1

A

Figure 1. Comprehensive review of electrode placement strategies used for measurement of auditory evoked potentials (AEPs) in the mouse. (A) Electrode positions (red cross, positive electrode; gray circle, negative electrode) used for mouse AEP measurements in 50 publications representative of the diversity of approaches in 115 relevant studies identified through a literature search. (B) Summary diagram illustrating range of positive electrode positions used in previous studies. Positive electrodes were most frequently positioned over the auditory cortex (purple), frontal cortex (green), or hippocampus (orange).

Figure 2

Figure 2. Experimental setup and ear-by-ear quantification of peripheral hearing sensitivity. (A) Auditory brainstem responses (ABRs) and auditory evoked potentials (AEPs) were measured with subdermal electrodes in anesthetized mice. (B) Hearing sensitivity of each ear was quantified by identifying the click-evoked ABR threshold, i.e. the minimum click intensity eliciting a detectable ABR waveform. Three examples are shown, with click-evoked ABR thresholds indicated by arrowheads: left (WT) 35 dB SPL, middle (*Df1/+*) 35 dB SPL, right (*Df1/+*) 70 dB SPL. Dashed vertical line indicates click onset. (C) Distributions of click-evoked ABR thresholds in WT ears (blue circles) and *Df1/+* ears (purple crosses). Ears with hearing loss were defined as those with click-evoked ABR thresholds more than 2.5 standard deviations above the mean WT threshold (i.e., above 47 dB SPL; dashed line). (D) Comparison of click-evoked ABR thresholds in the right and left ears; conventions as in C.

Figure 3

Figure 3. Central auditory gain (AEP/ABR amplitude ratio) was elevated in *Df1/+* mice for 16 kHz tone stimulus, and positively correlated with click-evoked ABR threshold. (A-B) Averaged tone-evoked ABR waveforms were measured between tested ear bulla and vertex (A) and AEP waveforms were measured from the contralateral hemisphere (B) during monaural stimulation of WT NH, *Df1/+* NH, and *Df1/+* HI ears with an 80 dB SPL, 16 kHz tone. Dashed vertical line indicates stimulus onset. (C) Tone-evoked amplitude of ABR wave 1 was significantly reduced for *Df1/+* HI ears versus *Df1/+* NH or WT NH ears. (D-E) However, cortical AEP waves measured contralateral to the stimulated ear showed a different pattern. Amplitude of the AEP P1-N1 complex was largest for stimulation of *Df1/+* NH ears (D); a similar trend was evident for the AEP N1-P2 complex (E). (F-G) Central gain (tone-evoked contralateral AEP amplitude normalized by tone-evoked ABR wave 1 amplitude) was abnormally elevated for the P1-N1 complex during stimulation of *Df1/+* NH ears (F), and for both the P1-N1 and N1-P2 complexes during stimulation of *Df1/+* HI ears (F,G). (H-J) Amplitude of ipsilateral ABR wave 1 (H), contralateral AEP P1-N1 complex (I), or contralateral AEP N1-P2 complex (J) evoked by a 80 dB SPL, 16 kHz tone, versus click-evoked ABR threshold for the stimulated ear. (H) Negative correlation with hearing threshold in the stimulated ear was weaker for tone-evoked AEP than ABR. (K-L)

Central gain of tone-evoked responses was positively correlated with click-evoked ABR threshold for both the P1-N1 complex (K) and N1-P2 complex (L). Each animal typically contributed two data points, one for each ear, with NH or HI status determined separately for each ear based on click-evoked ABR thresholds. Hearing sensitivity in the contralateral ear is indicated by the darkness of symbols: darker symbols indicate cases in which the contralateral ear was hearing impaired. Plot conventions: blue circles, WT; yellow cross, *Df1/+* NH with normal hearing in contralateral ear; brown cross, *Df1/+* NH with hearing impairment in contralateral ear; red cross, *Df1/+* HI with normal hearing in contralateral ear; dark red cross, *Df1/+* HI with hearing impairment in contralateral ear; gray solid line, 2D least-mean-squares best-fit line to the *Df1/+* data; asterisks, significance threshold for Dunn's post-hoc tests (C-G) or Spearman's rank correlation tests (H-L) (*$p < 0.05$, **$p < 0.01$, ***$p < 0.005$).

Figure 4

Figure 4. Growth of AEP amplitude with increasing tone intensity was steeper in *Df1/+* mice than WT mice, even for stimulation of *Df1/+* ears with normal hearing. (A-B) Example AEP waveforms (A) and average AEP waveforms (B) evoked by a 16 kHz tone presented at different sound intensity levels.

(C,E) AEP amplitudes for the tone-evoked P1-N1 complex (C) and N1-P2 complex (E) grew more steeply with increasing sound level for *Df1/+* mice than WT mice, regardless of the hearing status for the stimulated *Df1/+* ear. Thin lines represent recordings from contralateral hemispheres of individual ears; thick line with error bars shows mean ± SEM. (D,F) For both the P1-N1 complex (D) and the N1-P2 complex (F), slopes of level-dependent AEP growth functions were higher for stimulation of either *Df1/+* NH or *Df1/+* HI ears than for WT NH ears. Plot conventions as in Figure 2 H-L.

Figure 5. Growth of AEP amplitude with increasing inter-tone interval (ITI) was steeper for stimulation of *Df1/+* NH ears than for WT ears. (A-B) Example AEP waveforms (A) and average AEP waveforms (B) evoked by 80 dB SPL, 16 kHz tones presented at different ITIs. (C,E) AEP amplitudes for

the tone-evoked P1-N1 complex (C) and N1-P2 complex (E) grew more steeply with increasing ITI in *Df1/+* mice than WT mice, particularly at longer ITIs. Note the apparent plateau in AEP amplitude at 300-450 ms ITIs in WT data, and continued growth with ITI in *Df1/+* data, especially for stimulation of *Df1/+* ears with normal hearing. (D,F) For both the P1-N1 complex (D) and the N1-P2 complex (F), slopes of ITI-dependent AEP growth functions were higher for stimulation of *Df1/+* NH ears than WT ears. A similar trend, but no significant difference, was observed for comparison of ITI-dependent AEP growth-function slopes for stimulation of *Df1/+* HI ears versus WT ears; note however that interpretation of this latter result is complicated by the fact that the fixed-intensity tone stimulus was closer to hearing threshold for *Df1/+* HI ears than WT (NH) ears. Plot conventions as Figure 4.

Supplementary Information

Methods and Materials

Animal breeding and maintenance

Df1/+ mice were developed on a C57BL/6J background (1) and had been maintained on a C57BL/6J background for well over 25 generations, through pairing of *Df1/+* males either with WT females from the colony or (at least yearly) with newly acquired C57BL/6J females from Charles River UK. Mice were raised in standard cages and mouse housing facilities, on a standard 12 h-light/12 h-dark cycle.

Acoustic stimulation

Auditory stimuli were generated at a sample rate of 195,312.5 Hz using a digital signal processor (Tucker Davis Technologies, TDT RX6), attenuated as needed (TDT PA5), amplified (TDT SA1), and presented using a free-field speaker (TDT FF1) positioned 10 cm from the ear directed toward the speaker. Speaker output was calibrated to within 1.5 dB of target values before each set of experiments using a G.R.A.S. free-field 1/4" microphone, placed at the location of the ear to be tested.

Experimental procedures

Animals were anesthetized with intraperitoneally injected ketamine (70-100 mg/kg) and medetomidine (0.24-0.50 mg/kg) for recording, and were also given subcutaneous doses of carprofen (4.8-8.3 mg/kg) for analgesia and atropine (0.10-0.16 mg/kg) to minimize respiratory secretions. Supplementary doses of ketamine or ketamine/medetomidine mixture were administered if a toe-pinch response or whisker twitching was observed. Body temperature was maintained at 37-38°C using a homeothermic blanket (Harvard Apparatus). Each mouse was placed on an elevated platform and oriented with the tested ear directed toward the speaker. The opposite ear was blocked with an earplug during the recording to ensure monaural stimulation.

Subdermal electrodes (Rochester Medical) were inserted at the vertex; the bulla of the tested ear; the ipsilateral and contralateral auditory cortex relative to the tested ear; and the olfactory bulb (ground electrode). Data was acquired at a 24,414 Hz sample rate (TDT RX5) using a low-impedance headstage and signal amplifier (TDT RA4LI and RA16SD, 20x gain overall, 2.2 Hz - 7.5 kHz filtering) along with a custom low-pass filter designed to remove attenuation switching transients (100 kHz cutoff). Stimulus presentation and data acquisition was controlled using software from TDT (Brainware) and custom software written in MATLAB.

Data processing

To check the quality of AEP recordings, distributions of root mean square (RMS) values of the 1000 trials in every AEP recording session were checked. No clear outlier trials that were discrete from the rest of RMS distribution were identified for any of the recordings; therefore, we didn't exclude any trials from the analysis. Three key deflections of the AEP waveform (P1, N1, and P2) were manually selected from the averaged waveform of all 1000 trials after removing heartbeat noise (2). Heartbeats were identified as large differences between the detrended waveform and a detrended, smoothed waveform, and eliminated by subtracting these differences from the detrended waveform.

Across all recordings from a stimulated ear and the contralateral auditory cortex, 4 recordings were excluded because no clear P1, N1 and P2 could be identified, and 2 were excluded because the mice died from possible anaesthetics overdose during recording.

References

1.   Lindsay EA, Botta A, Jurecic V, Carattini-Rivera S, Cheah YC, Rosenblatt HM, et al. Congenital heart disease in mice deficient for the DiGeorge syndrome region. Nature. 1999 Sep 23;401(6751):379–83.

2.   de Cheveigné A. Sparse time artifact removal. J Neurosci Methods. 2016 Mar 15;262:14–20.

Supplementary Figures

Supplementary Figure 1

Supplementary Figure 1. Latencies of AEP waves N1 and P2 evoked by a fixed-intensity tone were abnormally elevated in *Df1/+* mice and uncorrelated with click-evoked ABR threshold in the stimulated ear. (A-C) A fixed-intensity tone (80 dB SPL, 16 kHz) evoked longer latency N1 and P2 waves in *Df1/+* mice than WT mice, regardless of whether the stimulated *Df1/+* ear had a normal or impaired hearing threshold (B,C; N1: *Df1/+* NH 39.96 ± 5.05 ms, *Df1/+* HI 42.98 ± 6.82 ms, WT 36.39 ± 6.04 ms, p < 0.001, post-hoc tests, $p_{Df1/+ NH-Df1/+ HI}$ = 0.23, $p_{Df1/+ NH-WT}$ = 0.020, $p_{Df1/+ HI-WT}$ < 0.001; P2:, *Df1/+* NH 98.64 ± 13.75 ms, *Df1/+* HI 97.25 ± 13.28 ms, WT 88.74 ± 15.09 ms, p = 0.015, post-hoc tests, $p_{Df1/+ NH-Df1/+ HI}$ = 0.70, $p_{Df1/+ NH-WT}$ = 0.042, $p_{Df1/+ HI-WT}$ = 0.046). There were no significant differences between WT and *Df1/+* data for latency of the P1 wave (A; P1: *Df1/+* NH 24.35 ± 3.31 ms, *Df1/+* HI 24.66 ± 3.05 ms, WT 23.24 ± 2.33 ms, Kruskal-Wallis test, p = 0.057). (D-F) No significant correlations between the latency of AEP waves evoked by an 80 dB SPL, 16 kHz tone and the click-evoked ABR threshold of the stimulated *Df1/+* ear (Spearman's rank correlation tests, P1, N1 and P2 all p > 0.05). Plot conventions as in Figure 3.

Supplementary Figure 2

Supplementary Figure 2. Decrease of AEP wave latency with increasing tone intensity was slightly more pronounced in *Df1/+* mice than WT mice. (A,C,E) Latency for the P1 (A) and N1 (C) AEP waves evoked by a 16 kHz tone dropped slightly with increasing sound level, while latency for the P2 wave (E) was stable across groups (P1: Kruskal-Wallis test, p = 0.0017, post-hoc tests, $p_{Df1/+ NH-Df1/+ HI}$ = 0.18, $p_{Df1/+ NH-WT}$ = 0.0019, $p_{Df1/+ HI-WT}$ = 0.059; N1: Kruskal-Wallis test, p < 0.001, post-hoc tests, $p_{Df1/+ NH-Df1/+ HI}$ = 0.090, $p_{Df1/+ NH-WT}$ = 0.072, $p_{Df1/+ HI-WT}$ < 0.001; P2: Kruskal-Wallis test, p = 0.61). (B,D,F) Slopes of level-dependent latency growth functions for P1 (B) and N1 (D) were slightly steeper in Df1/+ ears, but the growth function for P2 didn't show significant between-group differences (F). Plot conventions as in Figure 4.

Supplementary Figure 3

Supplementary Figure 3. Plateau in growth of AEP wave amplitude at 300-350 ms inter-tone intervals (ITIs) for WT but not *Df1/+* mice. For both P1-N1 and N1-P2 complexes, AEP amplitude showed a plateau at longer ITIs (300-350 ms) in WT mice. In contrast, AEP wave amplitude kept increasing over this ITI interval in *Df1/+* mice, for stimulation of both *Df1/+* NH and *Df1/+* HI ears. Solid lines represent AEP recordings contralateral to the stimulated ears; thick line with error bars shows mean ± SEM.

Supplementary Figure 4

Supplementary Figure 4. No differences between *Df1/+* and WT mice in the relationship between AEP wave latency and ITI for a fixed-intensity tone. (A,C,E) Latencies of the P1 (A), N1 (C), and P2 (E) waves evoked by an 80 dB SPL, 16 kHz tone dropped with increasing ITI. Note that although AEP wave latencies were slightly elevated overall for both *Df1/+* NH and *Df1/+* HI data compared to WT data (see also Supplementary Figure 1), the relationship between wave latency and ITI was generally consistent across the three groups. Slopes of the ITI-dependent latency growth functions were not significantly different between groups for P1 or P2, and were different for N1 only overall, not in post-hoc tests (Kruskal-Wallis test, P1: p = 0.11; N1: p = 0.040, post-hoc tests, $p_{Df1/+ NH-Df1/+ HI}$ = 0.051, $p_{Df1/+ NH-WT}$ = 0.39, $p_{Df1/+ HI-WT}$ = 0.10; P2: p = 0.086). Plot conventions as in Figure 4.