Grove-level analysis of titer and prevalence of "*Candidatus* Liberibacter asiaticus" and

Wolbachia in *Diaphorina citri,* vector of citrus Huanglongbing

Marina Mann[1], Luke Thompson[2], Lynn Johnson[3], Michelle Heck[1,2,*]

[1]Plant Pathology and Plant-Microbe Biology Section, School of Integrative Plant Science, Cornell

University, Ithaca, NY, 14853, USA

[2]Emerging Pests and Pathogens Research Unit, Robert W. Holley Center, USDA Agricultural

Research Service, Ithaca, NY, 14853, USA

[3] Cornell Statistical Consulting Unit, Cornell University, Ithaca, NY 14853, USA

* Corresponding author

Michelle Heck, michelle.heck@usda.gov

Abstract

Huanglongbing (HLB, or citrus greening disease) affects all citrus varieties world-wide. In

the USA, Asia, and South America the causal agent is "*Candidatus* Liberibacter asiaticus" (*C*Las),

a phloem-limited, uncultured, alphaproteobacterium. The hemipteran insect vector, *Diaphorina*

citri (Asian citrus psyllid) acquires and transmits *C*Las in a circulative, propagative manner. In

addition to *C*Las, *D. citri* hosts multiple symbiotic bacteria including *Wolbachia* (wDi)*. In D. citri*,

wDi has been sequenced and studied but specific roles in *D. citri* biology are unknown. Using well

established quantitative PCR methods we measured *C*Las titer in *D. citri* collected from four

groves in central Florida with distinct HLB management strategies and tested whether *C*Las and

wDi titer were correlated in a sub-set of these insects. Grove site had the largest effect on *C*Las

titer. Sex had no effect on *C*Las titer, while higher wDi titer was correlated with *C*Las-uninfected

insects. Our results suggest that more directed follow-up research is necessary and important to

clarify whether field management tactics influence *C*Las titer in *D. citri* and to better understand

gene-by-environment interactions among *D. citri*, wDi and *C*Las. Now that millions of trees in

Florida have been treated with injectable formulations of oxytetracycline, which is likely to

decrease bacterial populations in *D. citri*, this study may represent the last biologically meaningful

snapshot of grove-level vector-pathogen ecology in the state during the HLB epidemic.

Introduction

Huanglongbing (HLB, also known as citrus greening disease) affects all citrus varieties

world-wide, a total of 61 countries as of December 2022) (Ghosh et al. 2023) and after decades

of targeted research for solutions and over $1 billion US dollars spent in research, HLB remains

the most devastating disease of citrus with no cure. It is caused by an unculturable, intracellular,

phloem-limited bacteria in the genus "*Candidatus* Liberibacter". The predominant species

causing HLB is "*Ca.* L. asiaticus" (*C*Las). Two other species, "*Ca.* L. africanus" and "*Ca.* L.

americanus" have also been described in multiple African countries and South-Central American

countries, respectively. *C*Las is spread from tree to tree by the insect vector *Diaphorina citri*

(Asian citrus psyllid), a hemipteran in the family Psyllidae which feeds exclusively on the phloem

of citrus trees as well as periwinkle and species of *Murraya*. *C*Las is transmitted by *D. citri* in an

intracellular, circulative, propagative manner (Ammar et al. 2016; Inoue et al. 2009; Higgins et al.

2022). While *C*Las can also spread via grafting infected citrus budwood, vector-borne dispersal is

the predominant and least controllable method causing new citrus infections.

Management of the vector requires an understanding of basic epidemiology and vector

biology in the field. In the United States, strict testing regimens are followed in Texas and

California to maintain quarantine areas for the vector and pathogen. But the state of Florida has

nearly succumbed to HLB. Despite the frequency of psyllid testing and a known increase in

insecticide resistance by *D. citri* (Chen et al. 2022; Vázquez-García et al. 2013; Naeem et al. 2019;

Rao et al. 2014; Pardo et al. 2018; Venkatesan et al. 2022), relatively few papers have been

published on population-level dynamics of *C*Las-*D. citri* infections in the field. A few studies have

looked at *D. citri*-*C*Las titer and prevalence in different situations, including specific field sites,

month to month changes, year to year trends, and differences between countries (Bové et al.

1993; Wulff et al. 2020; Capoor et al. 1974; Coy and Stelinski 2015; Hall 2018; Manjunath et al.

2008; Razi et al. 2014; Tiwari et al. 2010; Sétamou et al. 2020; Ukuda-Hosokawa et al. 2015;

Walter et al. 2012). These studies found high levels of variation in *C*Las titer across both the

individual and population level, but the latest of these was conducted in 2017. Genetic studies

have also shown that the variation in *C*Las acquisition, transmission and titer levels is stable and

heritable, consistent with these phenotypes being regulated by *D. citri* genes and proteins

(Ramsey et al. 2015; Mann et al. 2018; Ammar et al. 2018).

In addition to *C*Las, *D. citri* hosts multiple symbiotic bacteria which it maintains because

they provide various benefits (such as nutritional supplements and defensive polyketides)

(Hosseinzadeh et al. 2019b; Nakabachi et al. 2013). *Wolbachia* (wDi) is one such endosymbiont

found in *D. citri*, as well as over 50% of all insects. At least two genetically distinct populations of

wDi have been shown to infect *D. citri* in Florida (Chu et al. 2019). Understanding how wDi

interacts with both *D. citri* and *C*Las may be important to the future management of psyllid

population (Chu et al. 2019; Jain et al. 2017). In *D. citri*, wDi has been sequenced (Neupane et al.

2022; Saha et al. 2012; Shippy et al. 2019) and studied but its specific role in *D. citri* is still unclear.

Studies suggest wDi is 100% present in *D. citri* individuals (Hosseinzadeh et al. 2019b; Guidolin

and Cônsoli 2013; Hoffmann et al. 2014) and is found infecting many organs (Hosseinzadeh et al.

2019b; Ren et al. 2018) of *D. citri*. The bacteriome, the endosymbiotic organ where the other

psyllid endosymbiont bacteria reside, maintains lower titers as compared to other organs

(Hosseinzadeh et al. 2019b).

The goal of this study was to compare *C*Las titer of psyllids collected from four groves in

Florida in 2018 with high disease pressure to better understand the natural variation that exists

in *C*Las titer across different populations of *D. citri* and to test whether factors such as sex or wDi

titer influence *C*Las titer.

Methods

<u>Sample collection</u>

Four unique citrus groves in central Florida, near lake Okeechobee, were sampled for

adult *Diaphorina citri* during the first week of October, 2018 (see **Figure S1**). Individual *D. citri*

were collected into 70% ethanol using mouth aspiration. At least 1200 individual *D. citri* were

collected from each field. Samples were then air dried and randomly aliquoted into groups of 100

and promptly flash frozen in liquid nitrogen and stored at -80 °C until DNA extractions.

<u>DNA extraction and quantification</u>

DNA extractions were performed in three batches on a total of 1062 samples, randomly

selected from each field site in varying amounts, using the Qiagen DNeasy Blood and Tissue 96

extraction kit. This kit allowed 96 samples to be extracted simultaneously with a filter-based

method. The methods advertised with the kit were followed with minor adjustments as follows.

First, individual *D. citri* were sexed under a dissecting scope over ice, then moved via paint brush

to the 96-well racks of sample tubes supplied by the Qiagen kit. Three, 3.2mm steel beads were

added to each tube and the samples were capped using the rubber tube caps provided,

throughout the first and last steps of the protocol. Samples were disrupted for 2 minutes at 1000

rpm using a mechanical homogenization with a SPEX Sample Prep MiniG 1600 (Thermo Fisher).

The samples were kept on ice before and after sample disruption. All centrifugation steps were

performed at the maximum 4000 rpm (RCF of 3434 x g) available when using the Hermle Z466 3L

swing-out rotor with square microplate buckets and inserts to accommodate the stacked column

and flow-through plates. Samples were lysed for 45 min at 56 ℃ in a water bath and mixed once

midway. Following lysis, 4 ul RNase A per sample was added to purify the genomic DNA, at which

point, instead of tube caps, plastic PCR plate covers were used to cover the 96-well racks. Samples

were eluted once using 50 ul AE buffer then stored at -20 ℃ until DNA quantification and purity

checks, which were performed using a Nanodrop 2000.

<u>Quantitative PCR</u>

Quantitative polymerase chain reaction (qPCR) was performed on all 1062 samples using

TaqMan hydrolysis probes, testing for the presence and relative titer quantification of the

"*Candidatus* Liberibacter asiaticus" (*C*Las) 16S rRNA gene. The 16S rRNA forward primer (5'-

TCGAGCGCGTATGCAATACG-3'), reverse primer (5'-GCGTTATCCCGTAGAAAAAGGTAG-3') and

probe (5'-FAM-AGACGGGTG/ZEN/AGTAACGCG-3') were used, as previously established (Wenbin

Li, 2006). The probe includes a fluorescent dye (5' 6-FAM) and a quencher (ZEN, 3' Iowa Black

FQ). Standard thermal conditions were used during the qPCR. Specifically, 95 °C for 10 min (1

cycle) and 95 °C for 15 sec and 60 °C for 60 sec (40 cycles total). Each qPCR 384-well plate included

a standard curve dilution series (a serial dilution from 10^9 down to 10^3 of the synthetic plasmid

pUC57 containing a CLas 16S rDNA gene fragment). The resulting y-intercept was 38.303, R^2 =

0.964, slope = -3.095, and efficiency = 110.452%. Each qPCR plate also contained positive and

negative controls run with two technical replicates each. A subset of samples from the first batch

were subjected to additional qPCR using SYBR Green reagents to access wDi presence-absence

and estimate titer. The wDi *ftsZ* gene (ftsZ-81) (A. S. Guidolin 2013) forward primer

(5'AGCAGCCAGAGAAGCAAGAG3') and reverse primer (5'TACGTCGCACACCTTCAAAA3'), and the

Wingless (*WgL*) (MyLo L. Thao 2000) forward primer (GCTCTCAAAGATCGGTTTGACGG) and

reverse (GCTGCCACGAACGTTACCTTC) primers were run in parallel on the same 386 well plate.

All samples on every qPCR plate were run in duplicate (2 technical replicates) then averaged to

create the reported final quantitative cycle (Cq) values. The maximum number of qPCR cycles is

40, where Cq = 40 means the primers did not create a product that could be amplified so the

sample is non-infected or titers are below the limit of detection, while Cq < 40 suggests copies of

the target gene are present, and the insect is infected. CLas cell counts were calculated from Cq

using the standard curve and the QuantStudio6 software. Each qPCR well contained 20 ng of DNA

from a single *D. citri.* The QuantStudio6 software associated with the qPCR machine used the

standard curve dilution values and resulting best-fit linear equation and efficiency metrics to

internally calculate the *C*Las cell equivalent to the number of copies of the *C*Las rDNA sequence

in a single *D. citri*. The values were exported from the software as an excel document. *C*Las cell

equivalents for technical replicates of each sample were averaged (note, Cq = 40 yielded

undetermined values, which were transformed to zeros for calculations of cell equivalents), then

all samples from each field were averaged. Field mean *C*Las cell equivalent was found by taking

the difference.

Statistical Methods

A linear mixed-effects model (LMM) was used to analyze the *C*Las Cq data collected on

the individual insects. The model contained fixed effects of sex and grove and a random effect of

DNA extraction batch. Fixed effects were tested using F tests, and post-hoc pairwise comparisons

were performed using Tukey's HSD. Model assumptions of normality and homogeneous variance

were assessed visually using the model residuals. *P*-values < 0.05 were considered statistically

significant. The mean absolute titer of 918 individuals was calculated from a standard curve, then

the mean absolute titer was calculated for each field site. For each pairwise field comparison, the

difference in absolute mean was taken by subtracting the second from the first (**Table 1**). A subset

of the samples from batch number one, representing all four fields, was additionally tested for

the dual effect of wDi (Cq values normalized by *Wingless* Cq) and grove on *C*Las Cq. R packages

(R Core Team 2022) were required for the analysis of both *C*Las and wDi, including *ggplot2*

(Wickham 2-16)*, ggpubr* (Kassambara 2020), *emmeans* (Lenth 2023), and *lmertest* (Kuznetsova

et al. 2017)*.* To test for interactions between *C*Las and wDi in the insects, wDi Cq and *C*Las Cq

were converted into binary values as follows: for wDi, a value of 0 was assigned for Cq values

below the median, and a value of 1 was assigned for values above median and for *C*Las, a value

of 0 was assigned for Cq values less than 40, and a value of 1 for undermined Cq values or Cq=40).

The association between wDi and *C*Las titer was assessed using a Wilcoxon test, which clarified

whether wDi Cq above and below the median was different in CLas (+) and (-) insects, and a Chi-

squared test in R to test whether the wDi Cq distribution around the median differs from the

distributions of *C*Las Cq below or at Cq=40.

Results

1. Each grove had a distinct management strategy.

Groves were assigned names corresponding to their town of origin to protect the identity

of the participating growers. Psyllid infestations were found by scouting for flush points with

nymph honeydew deposits. A total of 450 adult *D. citri* were used from the Port St. Lucie grove.

This grove was one of two organic groves we sampled and did not utilize any pesticides,

herbicides or other chemical inputs at the time of sampling. It maintained a high *D. citri*

population as well as the highest observed levels of in-field entomopathogenic fungi (Higgins et

al. 2022) and other insects (including ladybeetles, ants, caterpillars and wasps). Multiple citrus

species as well as *Murraya spp.* grew within the grove. The majority of *D. citri* samples were

taken from *Citrus sinensis* (Valencia orange). Trees ranged in age from recently re-planted to well

established, bearing trees. We used a total of 164 *D. citri* adults from *C. sinensis* in the second

grove site, referred to as Pioneer. This grove was also organically managed and under a regimen

of periodic nitrogen, phosphorous and potassium (NPK) supplementation. Trees were mature

and bearing and well-spaced out. A total of 176 samples were used from an experimental plot

within a third grove, Whidden Corner. This grove was composed of mostly small, juvenile,

recently planted *C. sinensis* trees undergoing heavy pesticide treatment for *D. citri* control, with

applications occurring every three weeks. A total of 272 adult *D. citri* were used from a fourth

citrus grove, referred to as Montura. These insects were sampled from *C. sinensis,* and this grove

also experienced insecticide sprays every three weeks. The trees were bearing, tall, and thick

with leaves. In all groves, rootstock variety was unknown, and the majority of *D. citri* were

collected from flush.

2. Grove had the largest effect on *C*Las Cq.

There was a significant difference in *C*Las Cq among grove sites after controlling for the

effects of sex and DNA extraction batch (F = 212.408, df = 3, *P* < 0.001). The effect of DNA

extraction batch was included in the model to rule out a potential bias, and we found no

between-batch variability (var < 0.001) and low, or no standard deviation (SD < 0.001), suggesting

high accuracy of technical replications of DNA extraction.

The four populations of *D. citri* fell into three broad categories based on *C*Las titer and

secondarily, infection rates. After accounting for effects of batch and sex, Pioneer represents the

"low titer" population and has the lowest overall titer (bacterial load per individual *D. citri*) with

mean Cq = 39.0 and standard error (SE) = 0.32. Montura has the highest titer (mean Cq = 30.4,

SE = 0.28) and represents the "high titer" population, while Port St. Lucie (mean Cq = 32.1, SE =

0.30) and Whidden Corner (mean Cq = 31.6, SE = 0.30) report similar bacterial titers and make up

the third population of "medium titer" individuals. When absolute titer of *C*Las is calculated from

Cq and a standard curve, the three groups are still present. Means were calculated using only

samples with Cq<40, and Montura had the highest titer (29,630 *C*Las cell equivalents), Pioneer

the lowest titer (41 *C*Las cell equivalents), while Whidden Corner and Port St. Lucie fell in the

middle (8,912 and 11,270 *C*Las cell equivalents, respectively).

The percent infection rate is calculated using the proportion of insects with Cq values

lower than 40, as Cq = 40 are considered non-infected with *C*Las. While the three groups above

are determined by *C*Las titer, their infection rates do not necessarily match the trend. The "low

titer" population represented by Pioneer did also have the lowest *C*Las infection rate (26.82 %).

However, the "medium titer" populations represented by Port St. Lucie (97.11 %) and Whidden

Corner (96.02 %) recorded the highest percent infection rates. Finally, the "high titer" population

represented by Montura maintained the second-lowest percent infection rate (94.85 %). The

three populations as determined by mean Cq can be seen in **Figure 1** and pair-wise results

between grove populations (**Table 1**) additionally confirm visual differences between groves.

Figure 1. Box plot of *C*Las Cq distribution by citrus grove collection site and colored by sex.

Box plot center lines are the median, box top and bottom edges are the first and third quartile

respectively, and outliers are points that extend past the tails. NA samples (grey) are those that

have a *C*Las Cq value but no recorded sex. M = male (blue), F = female (red).

Table 1: Statistical results from each pair-wise field comparison using both mean *C*Las Cq and

absolute titer of *C*Las (genome copies of *C*Las in 20 ng of *D. citri* DNA).

	*C*Las cell equivalent	Quantification Cycle (Cq) of *C*Las[3]				
Field site	**Mean difference[1]**	**Mean difference[2]**	**SE**	**t.ratio**	**df**	***P***
Montura vs. Pioneer	29,588	-8.870	0.371	-23.910	910	< 0.001
Montura vs. Port St. Lucie	18,360	-1.711	0.285	-5.995	763	< 0.001
Montura vs. Whidden Corner	20,717	-1.139	0.369	-3.086	850	0.011
Pioneer vs. Port St. Lucie	-11,228	7.159	0.360	19.906	476	< 0.001
Pioneer vs. Whidden Corner	-8,870	7.731	0.411	18.806	924	< 0.001
Port St. Lucie vs. Whidden Corner	2,357	0.573	0.362	1.580	305	0.391

[1]The mean absolute *C*Las titer, measured as the number of genome copies per 20ng of *D. citri*
DNA, of all samples with Cq < 40 per field site, was subtracted from each other to yield this
difference in mean absolute *C*Las titer by field. "Montura vs. Pioneer" means the mean *C*Las cell
equivalent of the Pioneer psyllids was subtracted from the mean *C*Las cell equivalent of Montura
psyllids, in this case, yielding a positive value showing Montura psyllids have a higher bacterial
load (total number of *C*Las genome copies) than psyllids from the Pioneer grove site.
[2]The difference in adjusted mean Cq value between two field sites. For example, the mean
difference of -8.870 shows that Montura has 8.87 Cq values more (meaning higher titer) than
Pioneer.
[3]The Linear Mixed-effect model transforms the data and accounts for co-variates. DF = degrees
of freedom are calculated using the Kenward-Rodger method, SE = Standard Error, Mean
difference = estimated relative difference in means, t.ratio = test statistic for testing the pairwise
contrasts, *P*-values are adjusted using the Tukey method. *C*Las = "*Candidatus* Liberibacter
asiaticus", Cq = quantification cycle in a quantitative polymerase chain reaction (qPCR).
230

3. The sex of *D. citri* and color morph had no effect on *C*Las Cq means.

Sex of each *D. citri* individual was recorded. Of the 929 samples with recorded sex, the

overall ratio of males to females was nearly equal (Male: n=454, 48.86%; Female: n=475, 51.13%)

as was the ratio of males to females in each grove (Montura: Female=41.54%, n=113,

Male=46.32%, n=126; Pioneer: Female=42.68%, n=70, Male=39.02%, n=64; Port St. Lucie:

Female=50.00%, n=225, Male=42.22%, n=190; Whidden Corner: Female=38.07%, n=67,

Male=42.05%, n=74) (see **Table S1A** for further breakdown of male-female results).

The effect of sex on *C*Las Cq was tested and included in our model as *C*Las infection has

been shown to affect *D. citri* behavior differently by sex. There was no statistically significant

effect of sex on *C*Las Cq after controlling for grove and DNA extraction batch (F = 0.482, df =1, *P*

= 0.487). This can also be seen in **Figure 1**, as the blue and red colors show a random distribution

of sex by Cq value for each of the four groves.

In laboratory experiments, the color morph of *D. citri* was shown to be associated with

*C*Las titers (Hosseinzadeh et al. 2019a). While it was technically challenging to accurately assign

the color morphology to each specimen due to the insects being stored in ethanol prior to data

collection, no obvious associations between *C*Las titer and color morph emerged in these insects

(**Table S1B**). The lack of association between color morphology and *C*Las infection status has

previously been reported in the analysis of other grove-captured, Florida psyllid populations (Hall

et al. 2018).

4. *Wolbachia* (wDi) has higher titer in uninfected *D. citri.*

A subset of *D. citri* samples from the first DNA extraction batch were tested for wDi titer

(Total = 176: Port St, Lucie = 46, Pioneer = 42, Montura = 41, Whidden Corner = 47). WDi Cq was

compared to *C*Las Cq using a linear regression. The regression showed that there was no

correlation between *C*Las and wDi Cq in these *D. citri* from Florida citrus groves ($R^2 = 0.0005$)

(Figure S2). Additionally, wDi Cq was normalized against the *D. citri Wingless* gene (**Figure S3).** A

linear model assessed the relative effects of both grove and normalized wDi Cq on *C*Las Cq, and

also found that wDi was not significantly correlated with *C*Las Cq (F = 0.646, df = 1, *P* = 0.426) and

that the source grove maintains the strongest effect on *C*Las titer (F = 22.881, df = 3, *P* < 0.001,

Table S2).

In the regression analysis, we noticed that there was a large number of wDi samples with

low Cq values when *C*Las Cq = 40 (*C*Las uninfected samples, **Figure S2)**. We tested whether wDi

titer is significantly different depending on the presence or absence of *C*Las (present when Cq <

40, absent when Cq = 40), and found that when *C*Las Cq = 40 in a sample, wDi Cq was more likely

to be below the median wDi Cq value (Wilcoxon, n=119, *P* = 0.017). A Chi^2 test confirmed that

the proportion of wDi with Cq above and below the median differs significantly from expected

when compared to the *C*Las Cq distribution based on infection or no infection, X^2(1, n=119)

=6.086, *P*=0.013. *C*Las-uninfected *D. citri* tend to have higher wDi titers as compared to *C*Las-

infected individuals.

Discussion

Natural variation in vector competency plays a role in disease epidemiology. Vector

competency is the potential of a vector to transmit a pathogen. In *D. citri*, vector competency is

apparently in part regulated by *D. citri* genes and proteins functioning at distinct tissue barriers

which *C*Las must overcome prior to inoculation into a new host tree (Ammar et al. 2018; Mann

et al. 2018; Kruse et al. 2017, 2018; Hosseinzadeh and Heck 2023). *D. citri* acquires and transmits

*C*Las in an intracellular, circulative, propagative manner. Vector competency in our study was

measured through infection rate and *C*Las titer for each population of vectors studied. Studies

have shown that approximation of *C*Las titer in individual insects can predict vector capacity.

There is a strong positive correlation between *C*Las titer in the psyllid and transmission efficiency

(Ammar et al. 2018; Hosseinzadeh et al. 2019a). Broadly, our study shows that there are

population-level differences in *D. citri* vector competency that are likely maintained through a

combination of gene by environment interactions.

During circulative propagative transmission, *C*Las must cross the gut epithelial cell layer

to enter the body cavity of the psyllid. Next, *C*Las must circulate throughout the body and

replicate in tissues, especially the salivary glands. Finally, *C*Las must be inoculated into healthy

citrus plants while feeding. The efficiency of the first and second steps can be approximated by

using qPCR to detect *C*Las titer in adults that have acquired *C*Las, while the third step can only be

measured by careful transmission studies. Also, the development of vector-competent adults

requires *C*Las acquisition to occur during the psyllid's nymphal stages (Inoue et al. 2009; Ammar

et al. 2016; Igwe et al. 2022). This study approximates the grove-wide vector capacity of four *D.*

citri populations through measurements of *C*Las titer in individual insects. A key assumption of

this study is that the majority of trees in all citrus-growing counties of Florida are infected with

*C*Las, and thus, all adult psyllids that are infected with *C*Las likely acquired as nymphs and are

competent transmitters. However, our results suggest these competent transmitters may still fall

into categories of low to high vector capacity, as determined by mean *C*Las titer of the population.

Other factors may influence *D. citri* vector competency and the variation we observed in

our study. Prior work has shown that host switching influences heritable variation in *D. citri* vector

competency (Ramsey et al. 2022). While grove-collected sampling makes it difficult to control or

understand all variables that may influence *C*Las titer in psyllids, all four *D. citri* populations were

sampled from the same citrus variety (*Citrus sinensis*), within the same week in October, 2018,

so it represents a snapshot in time of the potential of *D. citri* in these regions to acquire and

transmit. Additionally, the role that symbionts like wDi might play in mediating vector-pathogen

interactions is unclear, so the effect of wDi on *C*Las titer from these four fields was also tested.

Our results show that *D. citri* has higher wDi titer in healthy individuals, while field site has the

strongest effect on *C*Las titer, and sex does not have any effect. The field effect could be related

to the heritability of *D. citri* genes involved in vector competency, the management regimen, or

transient access to and feeding on non-citrus host plants such as *Murraya spp.* in the grove agro-

ecosystem. More work is required to fully explore these hypotheses.

Abandoned groves have also been shown to maintain high *D. citri* populations that travel

frequently between managed and abandoned areas (Sétamou et al. 2022; Boina et al. 2009) with

equally high *C*Las titers and infection rates as alive and managed groves (Hall 2018). Prior to 2010,

Tiwari et al (Tiwari et al. 2010) studied the *D. citri-C*Las infection rate and found 1.9 % in

abandoned and 1.2 % in managed grove populations while the mean *C*Las Cq value was 32.52 in

abandoned and 25.95 in managed groves. Over 10 years after Tiwari et al collected their data,

the infection rates found by our study in the least infected, but still managed, grove are at least

1.5x higher incidence while our other three groves have at least 50x higher infection rates. This

is undoubtedly due to the vastly increased number of infected trees in Florida over the past ten

years. A study by Pacheco et al (De Souza Pacheco et al. 2020) shows that when *D. citri* are raised

on *C*Las-infected citrus for four consecutive generations in a greenhouse setting, the percent-

infection and the titer in individuals increased with each generation from 42% to 100% infection

incidence, and 47.9 to 5,390 *C*Las cells per 100 ng of DNA. These results suggest that interruptions

in vector generations (such as through regular insecticide applications or during seasons of low

flush) in the field may help keep *C*Las titer and incidence lower in vector populations. The increase

in infection incidence seen in Florida groves despite various management innovations over the

last decade is a testament to both the effect of continuous populations of *D. citri* replicating on

infected citrus and reference to the accumulating impact HLB has wrought on the Florida citrus

industry.

Vector competency of *D. citri* populations and individuals varies widely in the field as we

have seen through this study and others (Capoor et al. 1974; Coy and Stelinski 2015; Hall 2018;

Ukuda-Hosokawa et al. 2015; Ammar et al. 2018; Sétamou et al. 2016; Xu et al. 1988), in addition

to laboratory-kept colonies (Inoue et al. 2009; Hosseinzadeh et al. 2019b; De Souza Pacheco et

al. 2020). In agreement with our results, Chu et al (Chu et al. 2016) show that three wild

populations of *D. citri* vary in their *C*Las titer and *C*Las infection incidence, and that like our study,

population (grove site) has the strongest effect on all their models. While grove is highly

correlated in the current study with *C*Las titer, the base differences may still come down to

molecular or genetic differences between the psyllid populations, *C*Las strains, other

endosymbionts, and unknown environmental factors. Methods to track genetic changes in *C*Las

between populations of *D. citri* show that greenhouse and field *D. citri* maintain diverging strains

of *C*Las (Higgins et al. 2022), and that microbiota vary with *D. citri* collection method. Research

has shown that the ability of *D. citri* to acquire and transmit *C*Las may be genetically linked

(Ammar et al. 2018). Studies that separate acquisition and transmission (Ammar et al. 2016;

Ukuda-Hosokawa et al. 2015; Ammar et al. 2018; Xu et al. 1988) agree that these are likely two

separate traits, but that they are correlated, such that higher acquisition rates often lead to more

efficient transmission (Ammar et al. 2018). The current study used *C*Las titer as a measure of

acquisition in adult *D. citri*, (which may also include effects of internal *C*Las replication) but the

high numbers of infected individuals (infection rate) also suggests transmission of *C*Las is

occurring in the field and that adult psyllids acquired as nymphs from infected trees. Now that

direct injection of the antibiotic oxytetracycline is becoming a wide-spread HLB management

tactic in Florida, understanding the impact of OTC on *C*Las titer in psyllids will be important to

understand if the use of OTC will indirectly lower vector competency in psyllid populations.

The effect of wDi titer on *C*Las in *D. citri* is still uncertain, and there was no effect of sex

on wDi or *C*Las observed in this study. However, in a related psyllid, *Cacopsylla pyri* (pear psyllid)

studied by Serbina et al (Serbina et al. 2022), wDi titer was reduced in males. Several lab studies

investigated interactions among wDi, *D. citri* and *C*Las. Hosseinzadeh et al. (Hosseinzadeh et al.

2019b) found that wDi titer varied between sexes, and *D. citri* had higher titer in males that were

infected with *C*Las than uninfected males, though wDi did not significantly differ between *C*Las

infected and uninfected populations as a whole. Others found that wDi titer was significantly

higher in *C*Las-negative *D. citri*, except when controlling for population effect (Chu et al. 2019;

Ren et al. 2018; Chu et al. 2016). Fagen et al (Fagen 2012) showed the opposite trend, that *C*Las

has a negative relationship with other *D. citri* obligate endosymbionts, but a positive relationship

to wDi regarding titer. Our results, using insects collected from citrus groves instead of

greenhouse and chamber-raised populations, contradict some of these previous lab-based

results. We saw 100% wDi infection rate among our individuals tested, and the titer of wDi

tended to be higher in *C*Las-uninfected *D. citri* individuals. These contradictions underscore the

importance to not over-generalize the results from lab studies and the importance of field-based

inquiries on vector-microbe interactions.

Conclusion

This snapshot assessment of vector populations showed grove-specific differences in *C*Las

infection. Specifically, infection rate and titer of *C*Las within individual *D. citri* varied by grove site,

exhibiting three patterns of vector competency. However, no conclusions could be drawn on the

components of field management that influenced *C*Las titer in *D. citri.* Additionally, while there

was no impact of sex on *C*Las titer, wDi titers were higher in *C*Las-uninfected individuals. These

results contradict some studies using lab-reared psyllid colonies. Unknown environmental factors

may contribute to titer levels in insects in the grove and caution is necessary when interpreting

results from lab-reared insects for field-based applications. The data suggest that future studies

should prioritize understanding the effect of grove management practices, including the use of

non-citrus plants that are hosts of *D. citri* in the grove, on the dynamics of *D. citri* infection by

*C*Las. Additionally, understanding the contributions that both environment and psyllid genes –

the G x E interaction – have on vector competency will help make all future genetic, genomic and

field-based studies of HLB more informed when developing control strategies or other programs

aimed blocking *C*Las transmission by *D. citri*.

Acknowledgements

We thank Southern Gardens Citrus, specifically, Tim Eyrich and Shannon Leahy – and Fort Pierce

USDA ARS researcher Dr. David Hall and technician Kathy Moulton, for their help, organization,

and permission to collect live psyllids from grove sites in Florida, USA.

Financial Support

Funding for this research was graciously provided by NIFA-AFRI Predoctoral Fellowship award #

2021-67011-35143 for Marina Mann, USDA ARS CRIS project 8062-22410-007-000-D, and USDA

NIFA grants 2020-70029-33176 and 2022-70029-38503.

Competing Interests

The authors declare that they have no competing interests.

References

Ammar, E.-D., Hall, D. G., Hosseinzadeh, S., and Heck, M. 2018. The quest for a non-vector
psyllid: Natural variation in acquisition and transmission of the huanglongbing pathogen
'Candidatus Liberibacter asiaticus' by Asian citrus psyllid isofemale lines. PLoS ONE.
13:e0195804.
398

399 Ammar, E.-D., Ramos, J. E., Hall, D. G., Dawson, W. O., and Shatters, R. G. 2016. Acquisition,
400 Replication and Inoculation of Candidatus Liberibacter asiaticus following Various Acquisition
401 Periods on Huanglongbing-Infected Citrus by Nymphs and Adults of the Asian Citrus Psyllid.
402 PLoS ONE. 11:e0159594.
403

Boina, D. R., Meyer, W. L., Onagbola, E. O., and Stelinski, L. L. 2009. Quantifying Dispersal of
Diaphorina citri (Hemiptera: Psyllidae) by Immunomarking and Potential Impact of Unmanaged
Groves on Commercial Citrus Management. Environ Entomol. 38:1250–1258.
Bové, J. M., Garnier, M., Ahlawat, Y. S., Chakraborty, N. K., and Varma, A. 1993. Detection of the
Asian Strains of the Greening BLO by DNA-DNA Hybridization in Indian Orchard Trees and
Malaysian Diaphorina citri Psyllids. International Organization of Citrus Virologists Conference
Proceedings (1957-2010). 12 Available at: https://escholarship.org/uc/item/0ng306x8
[Accessed July 19, 2023].
412

413 Capoor, S. P., Rao, D. G., and Viswanath, S. M. 1974. Greening Disease of Citrus in the Deccan
414 Trap Country and its Relationship with the Vector, *Diaphorina citri* Kuwayama. International
415 Organization of Citrus Virologists Conference Proceedings (1957-2010). 6 Available at:
416 https://escholarship.org/uc/item/6rm6x1tw [Accessed July 19, 2023].
417

418 Chen, Q., Li, Z., Liu, S., Chi, Y., Jia, D., and Wei, T. 2022. Infection and distribution of *Candidatus*
419 Liberibacter asiaticus in citrus plants and psyllid vectors at the cellular level. Microbial
420 Biotechnology. 15:1221–1234.
421

422 Chu, C., Hoffmann, M., Braswell, W. E., and Pelz-Stelinski, K. S. 2019. Genetic variation and
423 potential coinfection of *Wolbachia* among widespread Asian citrus psyllid (*Diaphorina citri*
424 Kuwayama) populations. Insect Science. 26:671–682.
425

426 Chu, C.-C., Gill, T. A., Hoffmann, M., and Pelz-Stelinski, K. S. 2016. Inter-Population Variability of
427 Endosymbiont Densities in the Asian Citrus Psyllid (*Diaphorina citri* Kuwayama). Microb Ecol.
428 71:999–1007.
429

430 Coy, M. R., and Stelinski, L. L. 2015. Great Variability in the Infection Rate of ' *Candidatus*
431 Liberibacter Asiaticus' in Field Populations of *Diaphorina citri* (Hemiptera: Liviidae) in Florida.
432 Florida Entomologist. 98:356–357.
433

434 De Souza Pacheco, I., Manzano Galdeano, D., Spotti Lopes, J. R., and Machado, M. A. 2020.
435 Development on Infected Citrus over Generations Increases Vector Infection by '*Candidatus*
436 Liberibacter Asiaticus in *Diaphorina citri*.' Insects. 11:469.
437

438 Fagen, J. R. 2012. Characterization of the Relative Abundance of the Citrus Pathogen *Ca*.
439 Liberibacter Asiaticus in the Microbiome of Its Insect Vector, *Diaphorina citri*, using High
440 Throughput 16S rRNA Sequencing. TOMICROJ. 6:29–33.
441

442 Ghosh, D., Kokane, S., Savita, B. K., Kumar, P., Sharma, A. K., Ozcan, A., et al. 2023.
443 Huanglongbing Pandemic: Current Challenges and Emerging Management Strategies. Plants.
444 12:160.
445

446 Guidolin, A. S., and Cônsoli, F. L. 2013. Molecular Characterization of *Wolbachia* Strains
447 Associated with the Invasive Asian Citrus Psyllid *Diaphorina citri* in Brazil. Microb Ecol. 65:475–
448 486.
449

450 Hall, D. G. 2018. Incidence of " *Candidatus* Liberibacter asiaticus" in a Florida population of
451 Asian citrus psyllid. J Appl Entomol. 142:97–103.

Higgins, S. A., Mann, M., and Heck, M. 2022. Strain Tracking of ' *Candidatus* Liberibacter asiaticus', the Citrus Greening Pathogen, by High-Resolution Microbiome Analysis of Asian Citrus Psyllids. Phytopathology®. 112:2273–2287.

Hoffmann, M., Coy, M. R., Gibbard, H. N. K., and Pelz-Stelinski, K. S. 2014. *Wolbachia* Infection Density in Populations of the Asian Citrus Psyllid (Hemiptera: Liviidae). Environ Entomol. 43:1215–1222.

Hosseinzadeh, S., and Heck, M. 2023. Variations on a theme: factors regulating interaction between *Diaphorina citri* and "*Candidatus* Liberibacter asiaticus" vector and pathogen of citrus huanglongbing. Current Opinion in Insect Science. 56:101025.

Hosseinzadeh, S., Ramsey, J., Mann, M., Bennett, L., Hunter, W. B., Shams-Bakhsh, M., et al. 2019a. Color morphology of *Diaphorina citri* influences interactions with its bacterial endosymbionts and '*Candidatus* Liberibacter asiaticus' ed. Janice L. Bossart. PLoS ONE. 14:e0216599.

Hosseinzadeh, S., Shams-Bakhsh, M., Mann, M., Fattah-Hosseini, S., Bagheri, A., Mehrabadi, M., et al. 2019b. Distribution and Variation of Bacterial Endosymbiont and "*Candidatus* Liberibacter asiaticus" Titer in the Huanglongbing Insect Vector, *Diaphorina citri* Kuwayama. Microb Ecol. 78:206–222.

Igwe, D. O., Higgins, S. A., and Heck, M. 2022. An Excised Leaf Assay to Measure Acquisition of ' *Candidatus* Liberibacter asiaticus' by Psyllids Associated with Citrus Huanglongbing Disease. Phytopathology®. 112:69–75.

Inoue, H., Ohnishi, J., Ito, T., Tomimura, K., Miyata, S., Iwanami, T., et al. 2009. Enhanced proliferation and efficient transmission of *Candidatus* Liberibacter asiaticus by adult *Diaphorina citri* after acquisition feeding in the nymphal stage. Annals of Applied Biology. 155:29–36.
Jain, M., Fleites, L. A., and Gabriel, D. W. 2017. A Small *Wolbachia* Protein Directly Represses Phage Lytic Cycle Genes in " *Candidatus* Liberibacter asiaticus" within Psyllids ed. Katherine McMahon. mSphere. 2:e00171-17.

Kassambara, A. 2020. ggpubr: "ggplot2" based publication ready plots. (R package version 0.4.0). Available at: https://CRAN.R-project.org/package=ggpubr.

Kruse, A., Fattah-Hosseini, S., Saha, S., Johnson, R., Warwick, E., Sturgeon, K., et al. 2017. Combining 'omics and microscopy to visualize interactions between the Asian citrus psyllid

491 vector and the Huanglongbing pathogen *Candidatus* Liberibacter asiaticus in the insect gut ed.
492 Hanu R Pappu. PLoS ONE. 12:e0179531.
493

494 Kruse, A., Ramsey, J. S., Johnson, R., Hall, D. G., MacCoss, M. J., and Heck, M. 2018. *Candidatus*
495 Liberibacter asiaticus Minimally Alters Expression of Immunity and Metabolism Proteins in
496 Hemolymph of *Diaphorina citri* , the Insect Vector of Huanglongbing. J. Proteome Res. 17:2995–
497 3011.
498

499 Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B. 2017. **lmerTest** Package: Tests in
500 Linear Mixed Effects Models. J. Stat. Soft. 82 Available at: http://www.jstatsoft.org/v82/i13/
501 [Accessed August 24, 2023].
502

503 Lenth, R. 2023. emmeans: Estimated Marginal Means, aka Least-Squares Means. Available at:
504 https://CRAN.R-project.org/package=emmeans.
505

506 Manjunath, K. L., Halbert, S. E., Ramadugu, C., Webb, S., and Lee, R. F. 2008. Detection of '
507 *Candidatus* Liberibacter asiaticus' in *Diaphorina citri* and Its Importance in the Management of
508 Citrus Huanglongbing in Florida. Phytopathology®. 98:387–396.
509

510 Mann, M., Fattah-Hosseini, S., Ammar, E.-D., Stange, R., Warrick, E., Sturgeon, K., et al. 2018.
511 *Diaphorina citri* Nymphs Are Resistant to Morphological Changes Induced by "*Candidatus*
512 Liberibacter asiaticus" in Midgut Epithelial Cells ed. Marvin Whiteley. Infect Immun. 86:e00889-
513 17.
514

515 Naeem, A., Afzal, M. B. S., Freed, S., Hafeez, F., Zaka, S. M., Ali, Q., et al. 2019. First report of
516 thiamethoxam resistance selection, cross resistance to various insecticides and realized
517 heritability in Asian citrus psyllid *Diaphorina citri* from Pakistan. Crop Protection. 121:11–17.
518

519 Nakabachi, A., Ueoka, R., Oshima, K., Teta, R., Mangoni, A., Gurgui, M., et al. 2013. Defensive
520 Bacteriome Symbiont with a Drastically Reduced Genome. Current Biology. 23:1478–1484.
521 Neupane, S., Bonilla, S. I., Manalo, A. M., and Pelz-Stelinski, K. S. 2022. Complete de novo
522 assembly of *Wolbachia* endosymbiont of *Diaphorina citri* Kuwayama (Hemiptera: Liviidae) using
523 long-read genome sequencing. Sci Rep. 12:125.
524

525 Pardo, S., Martínez, A. M., Figueroa, J. I., Chavarrieta, J. M., Viñuela, E., Rebollar-Alviter, Á., et
526 al. 2018. Insecticide resistance of adults and nymphs of Asian citrus psyllid populations from
527 Apatzingán Valley, Mexico: Resistance of *D. citri* to bifenthrin, malathion and chlorpyrifos. Pest.
528 Manag. Sci. 74:135–140.
529

530   R Core Team. 2022. R: A language and environment for statistical computing. Available at:
531   https://www-R-project.org/.
532

533   Ramsey, J. S., Ammar, E.-D., Mahoney, J. E., Rivera, K., Johnson, R., Igwe, D. O., et al. 2022. Host
534   Plant Adaptation Drives Changes in *Diaphorina citri* Proteome Regulation, Proteoform
535   Expression, and Transmission of ' *Candidatus* Liberibacter asiaticus', the Citrus Greening
536   Pathogen. Phytopathology®. 112:101–115.
537

538   Ramsey, J. S., Johnson, R. S., Hoki, J. S., Kruse, A., Mahoney, J., Hilf, M. E., et al. 2015. Metabolic
539   Interplay between the Asian Citrus Psyllid and Its *Profftella* Symbiont: An Achilles' Heel of the
540   Citrus Greening Insect Vector ed. Murad Ghanim. PLoS ONE. 10:e0140826.
541

542   Rao, C. N., Shivankar, V. J., Sandnya, D., David, K. J., and Dhengre, V. N. 2014. Insecticide
543   resistance in field populations of Asian Citrus psyllid, *Diaphorina citri* Kuwayama (Hemiptera:
544   Psyllidae). Pesticide Research Journal. 26:42–47.
545

546   Razi, M. F., Keremane, M. L., Ramadugu, C., Roose, M., Khan, I. A., and Lee, R. F. 2014.
547   Detection of Citrus Huanglongbing-Associated ' *Candidatus* Liberibacter asiaticus' in Citrus and
548   *Diaphorina citri* in Pakistan, Seasonal Variability, and Implications for Disease Management.
549   Phytopathology®. 104:257–268.
550

551   Ren, S.-L., Li, Y.-H., Ou, D., Guo, Y.-J., Qureshi, J. A., Stansly, P. A., et al. 2018. Localization and
552   dynamics of *Wolbachia* infection in Asian citrus psyllid *Diaphorina citri* , the insect vector of the
553   causal pathogens of Huanglongbing. MicrobiologyOpen. 7:e00561.
554

555   Saha, S., Hunter, W. B., Reese, J., Morgan, J. K., Marutani-Hert, M., Huang, H., et al. 2012.
556   Survey of Endosymbionts in the *Diaphorina citri* Metagenome and Assembly of a *Wolbachia*
557   wDi Draft Genome ed. Dan Zilberstein. PLoS ONE. 7:e50067.
558

559   Serbina, Š. L., Gajski, D., Malenovský, I., Corretto, E., Schuler, H., and Dittmer, J. 2022.
560   *Wolbachia* infection dynamics in a natural population of the pear psyllid *Cacopsylla pyri*
561   (Hemiptera: Psylloidea) across its seasonal generations. Sci Rep. 12:16502.
562

563   Sétamou, M., Alabi, O. J., Kunta, M., Dale, J., and Da Graça, J. V. 2020. Distribution of
564   *Candidatus* Liberibacter asiaticus in Citrus and the Asian Citrus Psyllid in Texas Over a Decade.
565   Plant Disease. 104:1118–1126.
566

567 Sétamou, M., Alabi, O. J., Kunta, M., Jifon, J. L., and Da Graça, J. V. 2016. Enhanced Acquisition
568 Rates of ' *Candidatus* Liberibacter asiaticus' by the Asian Citrus Psyllid (Hemiptera: Liviidae) in
569 the Presence of Vegetative Flush Growth in Citrus. J Econ Entomol. 109:1973–1978.
570

571 Sétamou, M., Tarshis Moreno, A., and Patt, J. M. 2022. Source or Sink? The Role of Residential
572 Host Plants in Asian Citrus Psyllid Infestation of Commercial Citrus Groves ed. Arash Rashed.
573 Journal of Economic Entomology. 115:438–445.
574

575 Shippy, T. D., Hosmani, P. S., Flores-Gonzalez, M., Mann, M., Miller, S., Weirauch, M. T., et al.
576 2019. *Diaci v3.0: Chromosome-level assembly,* de novo *transcriptome and manual annotation of*
577 Diaphorina citri, *insect vector of Huanglongbing*. Genomics. Available at:
578 http://biorxiv.org/lookup/doi/10.1101/869685 [Accessed August 24, 2023].
579

580 Tiwari, S., Lewis-Rosenblum, H., Pelz-Stelinski, K., and Stelinski, L. L. 2010. Incidence of
581 *Candidatus* Liberibacter asiaticus Infection in Abandoned Citrus Occurring in Proximity to
582 Commercially Managed Groves. jnl. econ. entom. 103:1972–1978.
583

584 Ukuda-Hosokawa, R., Sadoyama, Y., Kishaba, M., Kuriwada, T., Anbutsu, H., and Fukatsu, T.
585 2015. Infection Density Dynamics of the Citrus Greening Bacterium "*Candidatus* Liberibacter
586 asiaticus" in Field Populations of the Psyllid *Diaphorina citri* and Its Relevance to the Efficiency
587 of Pathogen Transmission to Citrus Plants ed. H. Goodrich-Blair. Appl Environ Microbiol.
588 81:3728–3736.
589

590 Vázquez-García, M., Velázquez-Monreal, J., Medina-Urrutia, V. M., Cruz-Vargas, C. D. J.,
591 Sandoval-Salazar, M., Virgen-Calleros, G., et al. 2013. Insecticide Resistance in Adult *Diaphorina*
592 *citri* Kuwayama from Lime Orchards in Central West Mexico. Southwestern Entomologist.
593 38:579–596.
594

595 Venkatesan, T., Chethan, B. R., and Mani, M. 2022. Insecticide Resistance and Its Management
596 in the Insect Pests of Horticultural Crops. In *Trends in Horticultural Entomology*, ed. M. Mani.
597 Singapore: Springer Nature Singapore, p. 455–490. Available at:
598 https://link.springer.com/10.1007/978-981-19-0343-4_14 [Accessed August 24, 2023].
599

600 Walter, A. J., Hall, D. G., and Duan, Y. P. 2012. Low Incidence of ' *Candidatus* Liberibacter
601 asiaticus' in *Murraya paniculata* and Associated *Diaphorina citri*. Plant Disease. 96:827–832.
602

603 Wickham, H. 2-16. ggplot2: Elegant graphics for data analysis.
604

Wulff, N., Daniel, B., Sassi, R., Moreira, A., Bassanezi, R., Sala, I., et al. 2020. Incidence of
Diaphorina citri Carrying *Candidatus* Liberibacter asiaticus in Brazil's Citrus Belt. Insects. 11:672.
Xu, C. F., Xia, Y. H., Li, K. B., and Ke, C. 1988. Further Study of the Transmission of Citrus
Huanglungbin by a Psyllid, *Diaphorina citri* Kuwayama. International Organization of Citrus
Virologists Conference Proceedings (1957-2010). 10 Available at:
https://escholarship.org/uc/item/0w42q0r7 [Accessed July 19, 2023].
611

3   **Figure S1:** Map of four grove sites in the central Florida region. A) Port St. Lucie, B) Pioneer, C)

4   Whidden Corner, D) Montura.

Figure S2: Linear regression of 176 samples with paired *C*Las-*Wolbachia* Cq values, showing no

correlation (R^2=0.00055) in *D. citri*. Grey area is 95% confidence, while the blue line is the best fit line

which follows the equation stated. Note that many of the *Wolbachia* samples with Cq < 20 Cq have *C*Las

Cq = 40 (uninfected).

Figure S3: Normalization of *Wolbachia* wDi Cq values using a psyllid gene internal standard. In addition to measuring the Cq of wDi (A), wDi Cq can be normalized by dividing with *Wingless* Cq (B), which is the *D. citri* endogenous control gene. Paired *C*Las Cq values are also available for each sample (C), and a standard curve was used while measuring *C*Las Cq instead of *Wingless.* W*ingless (WgL)* Cq alone can be seen in (D). The goal of running the reference gene is to see consistent Cq values, despite variation in the variable of interest (in this case, testing for wDi Cq. Two 386-well qPCR plates were run (First: Pioneer and Port St. Lucie, Second: Montura and Whidden Corner samples). Statistical results are from pair-wise T-tests.

30   **Table S1:** Data for sex and color of individual *Diaphorina citri* from four citrus groves. Pioneer is

31   a consistent outlier due to a higher proportion of *C*Las-uninfected (Cq = 40) samples relative to

32   other groves, yielding a higher mean Cq as well.

A	*Total	Sex	% of Total	% of Sex w/Cq=40	**Mean Cq
Montura	239	Female	47.28	4.42	30.14
		Male	52.71	3.17	29.94
Pioneer	134	Female	52.24	65.71	37.33
		Male	47.76	82.81	37.18
Port St. Lucie	415	Female	54.22	4.89	31.73
		Male	45.78	5.79	31.65
Whidden Corner	141	Female	47.52	0.00	32.01
		Male	52.48	2.70	30.89

B	*Total	Color	% of Total	% of Color w/Cq=40	**Mean Cq
Montura	239	Blue	17.99	4.65	31.75
		Orange	26.36	15.87	29.27
		White	55.65	51.13	29.45
Pioneer	134	Blue	32.09	69.77	37.54
		Orange	47.01	79.37	37.31
		White	20.90	67.86	36.86
Port St. Lucie	415	Blue	20.24	4.76	32.23
		Orange	40.96	6.47	31.51
		White	38.80	4.35	31.60
Whidden Corner	141	Blue	13.48	0.00	31.49
		Orange	45.39	1.56	31.81
		White	41.13	1.72	30.99

35   *NA samples are excluded from total counts and from all percent and mean calculations.

36   **mean Cq uses samples with Cq<40, excludes samples with Cq=40.

37 **Table S2**: Statistical results and post-hoc tests from the linear model used to test the effect of

38 grove on *C*Las Cq while controlling for normalized *Wolbachia* Cq.

Pair-wise grove (field) comparisons controlling for effect of *Wolbachia* *lm*(CLas.Cq ~ Field + Norm_Wolb, data=dataset)					
Contrast	**Est**	**SE**	**t.ratio**	**df**	**P**
Montura vs. Pioneer	-7.264	0.996	-7.296	114	<0.0001
Montura vs. Port St. Lucie	-0.278	0.943	-0.295	114	0.9911
Montura vs. Whidden Corner	-2.329	0.943	-2.471	114	0.0701
Pioneer vs. Port St. Lucie	6.986	0.961	7.271	114	<0.0001
Pioneer vs. Whidden Corner	4.935	1.028	4.800	114	<0.0001
Port St. Lucie vs. Whidden Corner	-2.052	0.974	-2.107	114	0.1572

*The estimate (Est) value of -7.264 (for example) shows that Montura has on average about 7

points smaller Cq value than Pioneer, which can be understood as a multi-fold higher mean *C*Las

titer.

Table S2 Legend: DF = degrees of freedom, SE = Standard Error, Est = estimate, t.ratio = test

statistic for testing the pairwise contrasts, *P* < 0.0001 is adjusted using the Tukey method. *C*Las =

"*Candidatus* Liberibacter asiaticus", Norm.Wolb = *Wolbachia* titer normalized by dividing

Wolbachia Cq by *Wingless* Cq, Cq = quantification cycle in a quantitative polymerase chain

reaction (qPCR).