Grove-level analysis of titer and prevalence of *"Candidatus* Liberibacter asiaticus" and

Wolbachia in *Diaphorina citri,* vector of citrus Huanglongbing

Marina Mann[1], Luke Thompson[2], Lynn Johnson[3], Michelle Heck[1,2,*]

[1]Plant Pathology and Plant-Microbe Biology Section, School of Integrative Plant Science, Cornell

University, Ithaca, NY, 14853, USA

[2]Emerging Pests and Pathogens Research Unit, Robert W. Holley Center, USDA Agricultural

Research Service, Ithaca, NY, 14853, USA

[3]Cornell Statistical Consulting Unit, Cornell University, Ithaca, NY 14853, USA

* Corresponding author

Michelle Heck, michelle.heck@usda.gov

Abstract

Huanglongbing (HLB, or citrus greening disease) affects all citrus varieties world-wide. In

the USA, Asia, and South America the causal agent is *"Candidatus* Liberibacter asiaticus" (*C*Las),

a phloem-limited, uncultured, alphaproteobacterium. The hemipteran insect vector, *Diaphorina*

citri (Asian citrus psyllid) acquires and transmits *C*Las in a circulative, propagative manner. In

addition to *C*Las, *D. citri* hosts multiple symbiotic bacterial species including *Wolbachia* (wDi). In

D. citri, wDi has been sequenced and studied but specific roles in *D. citri* biology are unknown.

Using well established quantitative PCR methods we measured *C*Las titer in *D. citri* collected

from four groves in central Florida with distinct HLB management strategies and tested

whether *C*Las and wDi titer were correlated in a sub-set of these insects. Grove site had the

largest effect on *C*Las titer. Sex had no effect on *C*Las titer, while higher wDi titer was correlated

with non-infected insects. Our results suggest that more directed follow-up research is

necessary and important to clarify whether field management tactics influence *C*Las titer in *D.*

citri and to better understand gene-by-environment interactions among *D. citri*, wDi and *C*Las.

Now that millions of trees in Florida have been treated with injectable formulations of

oxytetracycline, which is likely to decrease bacterial populations in *D. citri*, this study may

represent the last biologically meaningful snapshot of grove-level vector-pathogen ecology in

the state during the HLB epidemic.

Introduction

Huanglongbing (HLB, also known as citrus greening disease) affects all citrus varieties

world-wide, a total of 61 countries as of December 2022) (Ghosh et al. 2023) and after decades

of targeted research for solutions and over $1 billion US dollars spent in research, HLB remains

the most devastating disease of citrus with no cure. It is caused by an unculturable,

intracellular, phloem-limited bacteria in the genus "*Candidatus* Liberibacter". The predominant

species causing HLB is "*Ca.* L. asiaticus" (*C*Las). Two other species, "*Ca.* L. africanus" and "*Ca.* L.

americanus" have also been described in multiple African countries and South-Central

American countries, respectively. *C*Las is spread from tree to tree by the insect vector

Diaphorina citri (Asian citrus psyllid), a hemipteran in the family Psyllidae which feeds

exclusively on the phloem of citrus trees as well as species of *Murraya*. *C*Las is transmitted by *D.*

citri in an intracellular, circulative, propagative manner (Ammar et al. 2016; Inoue et al. 2009;

Higgins et al. 2022). While *C*Las can also spread via grafting infected citrus budwood, vector-

borne dispersal is the predominant and least controllable method causing new citrus infections.

Lastly, *C*Las can infect periwinkle via dodder, but periwinkle is not a host for *D. citri* (Garnier and

Bove 1983).

Management of the vector requires an understanding of basic epidemiology and vector

biology in the field. In the United States, strict testing regimens are followed in Texas and

California to maintain quarantine areas for the vector and pathogen. But HLB is now endemic in

Florida. Despite the frequency of psyllid testing and a known increase in insecticide resistance

by *D. citri* (Chen et al. 2022; Vázquez-García et al. 2013; Naeem et al. 2019; Rao et al. 2014;

Pardo et al. 2018; Venkatesan et al. 2022), relatively few papers have been published on

population-level dynamics of *C*Las-*D. citri* infections in the field. A few studies have looked at *D.*

citri-*C*Las titer and prevalence in different situations, including specific field sites, month to

month changes, year to year trends, and differences between countries (Bové et al. 1993; Wulff

et al. 2020; Capoor et al. 1974; Coy and Stelinski 2015; Hall 2018; Manjunath et al. 2008; Razi et

al. 2014; Tiwari et al. 2010; Sétamou et al. 2020; Ukuda-Hosokawa et al. 2015; Walter et al.

2012). These studies reported high levels of variation in *C*Las titer across both the individual and

population level. Genetic studies have also shown that psyllid populations vary in their *C*Las

acquisition, transmission and titer phenotypes and that this variation is stable and heritable

over many generations, consistent with these phenotypes being regulated by *D. citri* genes and

proteins (Ramsey et al. 2015; Mann et al. 2018; Ammar et al. 2018).

In addition to *C*Las, *D. citri* hosts multiple symbiotic bacteria which it maintains because

they provide various benefits (such as nutritional supplements and defensive polyketides)

(Hosseinzadeh et al. 2019b; Nakabachi et al. 2013). *Wolbachia* (wDi) is one such endosymbiont

found in *D. citri*, as well as over 50% of all insects. At least two genetically distinct populations

of wDi have been shown to infect *D. citri* in Florida (Chu et al. 2019). Understanding how wDi

interacts with both *D. citri* and *C*Las may be important to the future management of psyllid

population (Chu et al. 2019; Jain et al. 2017). In *D. citri*, wDi has been sequenced (Neupane et

al. 2022; Saha et al. 2012; Shippy et al. 2019) and studied but its specific role in *D. citri* is still

unclear. Studies suggest wDi is 100% present in *D. citri* individuals (Hosseinzadeh et al. 2019b;

Guidolin and Cônsoli 2013; Hoffmann et al. 2014) and is found infecting many organs

(Hosseinzadeh et al. 2019b; Ren et al. 2018) of *D. citri*. The bacteriome, the endosymbiotic

organ where the other psyllid endosymbiont bacteria reside, maintains lower titers as

compared to other organs (Hosseinzadeh et al. 2019b).

The goal of this study was to compare *C*Las titer of psyllids collected from four groves in

Florida in 2018 with high disease pressure to better understand the natural variation that exists

in *C*Las titer across different populations of *D. citri* and to test whether factors such as sex or

wDi titer influence *C*Las titer.

Methods

<u>Sample collection</u>

Four citrus groves in central Florida, near lake Okeechobee, were sampled for adult

Diaphorina citri during the first week of October, 2018 (see **Figure S1**). Individual *D. citri* were

collected into 70% ethanol using mouth aspiration. At least 1200 individual *D. citri* were

collected from each field. Samples were air dried and randomly aliquoted into groups of 100,

promptly flash frozen in liquid nitrogen and stored at -80 °C until DNA extractions.

<u>DNA extraction and quantification</u>

DNA extractions were performed in three batches on a total of 1062 samples, randomly

selected from each field site in varying amounts, using the Qiagen DNeasy Blood and Tissue 96

extraction kit. This kit allowed 96 samples to be extracted simultaneously with a filter-based

method. The methods advertised with the kit were followed with minor adjustments as follows.

First, individual *D. citri* were sexed under a dissecting scope over ice, then moved via paint

brush to the 96-well racks of sample tubes supplied by the Qiagen kit. Three, 3.2mm steel

beads were added to each tube and the samples were capped using the rubber tube caps

provided, throughout the first and last steps of the protocol. Samples were disrupted for 2

minutes at 1000 rpm using a mechanical homogenization with a SPEX Sample Prep MiniG 1600

(Thermo Fisher). The samples were kept on ice before and after sample disruption. All

centrifugation steps were performed at the maximum 4000 rpm (RCF of 3434 x g) available

when using the Hermle Z466 3L swing-out rotor with square microplate buckets and inserts to

accommodate the stacked column and flow-through plates. Samples were lysed for 45 min at

56 °C in a water bath and mixed once midway. Following lysis, 4 ul RNase A per sample was

added to purify the genomic DNA, at which point, instead of tube caps, plastic PCR plate covers

were used to cover the 96-well racks. Samples were eluted once using 50 ul AE buffer then

stored at -20 °C until DNA quantification and purity checks, which were performed using a

Nanodrop 2000.

 Quantitative PCR

Quantitative polymerase chain reaction (qPCR) was performed on all 1062 samples

using TaqMan hydrolysis probes, testing for the presence and relative titer quantification of the

"*Candidatus* Liberibacter asiaticus" (*C*Las) 16S rRNA gene. The 16S rRNA forward primer (5′-

TCGAGCGCGTATGCAATACG-3′), reverse primer (5′-GCGTTATCCCGTAGAAAAAGGTAG-3′) and

probe (5′-FAM-AGACGGGTG/ZEN/AGTAACGCG-3′) were used, as previously established

(Wenbin Li, 2006). The probe includes a fluorescent dye (5′ 6-FAM) and a quencher (ZEN, 3′

Iowa Black FQ). Standard thermal conditions were used during the qPCR. Specifically, 95 °C for

10 min (1 cycle) and 95 °C for 15 sec and 60 °C for 60 sec (40 cycles total). Each qPCR 384-well

plate included a standard curve dilution series (a serial dilution from 10^9 down to 10^3 of the

synthetic plasmid pUC57 containing a *C*Las 16S rDNA gene fragment). The resulting y-intercept

was 38.303, R^2 = 0.964, slope = -3.095, and efficiency = 110.452%. Each qPCR plate also

contained positive and negative controls run with two technical replicates each. A subset of

samples from the first batch were subjected to additional qPCR using SYBR Green reagents to

access wDi presence-absence and estimate titer. The wDi *ftsZ* gene (ftsZ-81) (A. S. Guidolin

2013) forward primer (5'AGCAGCCAGAGAAGCAAGAG3') and reverse primer

(5'TACGTCGCACACCTTCAAAA3'), and the *Wingless* (*WgL*) (MyLo L. Thao 2000) forward primer

(GCTCTCAAAGATCGGTTTGACGG) and reverse (GCTGCCACGAACGTTACCTTC) primers were run

in parallel on the same 386 well plate. All samples on every qPCR plate were run in duplicate (2

technical replicates) then averaged to create the reported final quantitative cycle (Cq) values.

The maximum number of qPCR cycles is 40, where Cq = 40 means the primers did not create a

product that could be amplified so the sample is non-infected or titers are below the limit of

detection, while Cq < 40 suggests copies of the target gene are present, and the insect is

infected. *C*Las cell counts were calculated from Cq using the standard curve and the

QuantStudio6 software. Each qPCR well contained 20 ng of DNA from a single *D. citri.* The

QuantStudio6 software associated with the qPCR machine used the standard curve dilution

values and resulting best-fit linear equation and efficiency metrics to internally calculate the

*C*Las cell equivalent to the number of copies of the *C*Las rDNA sequence in a single *D. citri*. The

values were exported from the software as an excel document. *C*Las cell equivalents for

technical replicates of each sample were averaged (note, Cq = 40 yielded undetermined values,

which were transformed to zeros for calculations of cell equivalents), then all samples from

each field were averaged. Field mean *C*Las cell equivalent was found by taking the difference.

<u>Statistical Methods</u>

A linear mixed-effects model (LMM) was used to analyze the *C*Las Cq data collected on

the individual insects... Two models were compared to determine whether grove makes a

significant difference as a random effect. Model 1 included both DNA extraction batch and

grove as random effects, and sex as a fixed effect. Model 2 was reduced, containing only batch

as a random effect and sex as a fixed effect, with grove absent. An ANOVA was used to test the

difference between the two models. Model assumptions of normality and homogeneous

variance were assessed visually using the model residuals. *P*-values < 0.05 were considered

statistically significant. The mean absolute titer of 918 individuals was calculated from a

standard curve, then the mean absolute titer was calculated for each field site. For each

pairwise field comparison, non-zero titers were used to calculate field-specific means, the

means subtracted to find the difference and the absolute value was reported to highlight the

magnitude of the difference the second from the first (**Figure 2**). A subset of the samples from

batch number one, representing all four fields, was additionally tested for the dual effect of wDi

(Cq values normalized by *Wingless* Cq) and grove on *C*Las Cq. R packages (R Core Team 2022)

were required for the analysis of both *C*Las and wDi, including *ggplot2* (Wickham 2-16)*, ggpubr*

(Kassambara 2020), and *lmertest* (Kuznetsova et al. 2017). To test for interactions between *C*Las

and wDi in the insects, wDi Cq and *C*Las Cq were converted into binary values as follows: for

wDi, a value of 0 was assigned for Cq values below the median, and a value of 1 was assigned

for values above median and for *C*Las, a value of 0 was assigned for Cq values less than 40, and

a value of 1 for undermined Cq values or Cq=40).   The association between wDi and *C*Las titer

was assessed using a Wilcoxon test, which clarified whether wDi Cq above and below the

median was different in *C*Las (+) and (-) insects, and a Chi-squared test in R to test whether the

wDi Cq distribution around the median differs from the distributions of *C*Las Cq below or at

Cq=40.

Results

1. Each grove had a distinct characteristics, including tree age, type and management

169   strategies.

170   Groves were assigned names corresponding to their town of origin to protect the

identity of the participating growers. Psyllid infestations were found by scouting for flush points

with nymph honeydew deposits. A total of 450 adult *D. citri* were used from the Port St. Lucie

grove. This grove was one of two organically managed groves we sampled from, however, Port

St. Lucie did not utilize any pesticides, herbicides or other chemical inputs to our knowledge, at

the time of sampling. It maintained a high *D. citri* population as well as the highest observed

levels of in-field entomopathogenic fungi (Higgins et al. 2022) and other insects (including

ladybeetles, ants, caterpillars and wasps). Multiple citrus species as well as *Murraya spp.* grew

within the grove. The majority of *D. citri* samples were taken from *Citrus sinensis* (Valencia

orange). Trees ranged in age from recently re-planted to well established, bearing trees. We

used a total of 164 *D. citri* adults from *C. sinensis* in the second grove site, referred to as

Pioneer. This grove was also organically managed, and while organic pesticides were used, only

a regimen of periodic nitrogen, phosphorous and potassium (NPK) supplementation was

described in particular during sample collection, and any other treatments utilized are

unknown. Trees were mature and bearing and well-spaced out. A total of 176 samples were

used from an experimental plot within a third grove, Whidden Corner. This grove was

composed of mostly small, juvenile, recently planted *C. sinensis* trees undergoing heavy

pesticide treatment for *D. citri* control, with applications occurring every three weeks. A total of

272 adult *D. citri* were used from a fourth citrus grove, referred to as Montura. These insects

were sampled from *C. sinensis,* and this grove also experienced insecticide sprays every three

weeks. The trees were bearing, tall, and thick with leaves. In all groves, rootstock type was

unknown, and the majority of *D. citri* were collected from flush.

2. Grove had the largest effect on *C*Las Cq.

The random effect of grove site significantly outweighed any effect of sex or DNA

extraction batch on *C*Las Cq (Chi^2 = 462.01 , df = 2, *P* < 0.001). The effect of DNA extraction

batch was included in the model to rule out a potential bias, and we found no between-batch

variability (var. = 0.010, SD = 0.010) suggesting high accuracy of technical replications of DNA

extraction.

The four populations of *D. citri* fell into three broad categories based on *C*Las titer and

secondarily, infection rates. After accounting for effects of batch and sex, Pioneer represents

the "low titer" population and has the lowest overall titer (bacterial load per individual *D. citri*)

with mean Cq = 39.0 and standard error (SE) = 0.32. Montura has the highest titer (mean Cq =

30.4, SE = 0.28) and represents the "high titer" population, while Port St. Lucie (mean Cq = 32.1,

SE = 0.30) and Whidden Corner (mean Cq = 31.6, SE = 0.30) report similar bacterial titers and

make up the third population of "medium titer" individuals. When absolute titer of *C*Las is

calculated from Cq and a standard curve, the three groups are still present. Means were

calculated using only samples with Cq<40, and Montura had the highest titer (29,630 *C*Las cell

equivalents), Pioneer the lowest titer (41 *C*Las cell equivalents), while Whidden Corner and Port

St. Lucie fell in the middle (8,912 and 11,270 *C*Las cell equivalents, respectively).

The percent infection rate is calculated using the proportion of insects with Cq values

lower than 40, as Cq = 40 are considered non-infected with *C*Las. While the three groups above

are determined by *C*Las titer, their infection rates do not necessarily match the trend. The "low

titer" population represented by Pioneer did also have the lowest *C*Las infection rate (26.82 %).

However, the "medium titer" populations represented by Port St. Lucie (97.11 %) and Whidden

Corner (96.02 %) recorded the highest percent infection rates. Finally, the "high titer"

population represented by Montura maintained the second-lowest percent infection rate

(94.85 %). The three populations as determined by mean Cq can be seen in **Figure 1** and pair-

wise results between grove populations (**Figure 2**) additionally confirm visual differences

between groves.

Figure 1. Box plot of *C*Las Cq distribution by citrus grove collection site and colored by sex.

Box plot center lines are the median, box top and bottom edges are the first and third quartile

respectively, and outliers are points that extend past the tails. NA samples (grey) are those that

have a *C*Las Cq value but no recorded sex. M = male (blue), F = female (red).

Mann et al 2023

Absolute CLas Cell Equivalent

	Port St. Lucie	Pioneer	Whidden Corner	Montura
Montura	18,360	29,588	20,717	**29,630**
Whidden Corner	2,357	8,870	**8,912**	
Pioneer	11,288	**41**		
Port St. Lucie	**11,270**			

Quantification Cycle (Cq) of CLas

	Port St. Lucie	Pioneer	Whidden Corner	Montura
Montura	1.7	8.6	1.2	**30.4**
Whidden Corner	0.5	7.4	**31.6**	
Pioneer	6.9	**39**		
Port St. Lucie	**32.1**			

Figure 2: Pair-wise field comparisons using both mean CLas Cq and titer of CLas (genome copies of CLas in 20 ng of *D. citri* DNA). Field-specific means were computed using only non-zero absolute titer values, or Cq < 40 values and the absolute values are reported.

3. The sex of *D. citri* and color morph had no effect on CLas Cq means.

Sex of each *D. citri* individual was recorded. Of the 929 samples with recorded sex, the overall ratio of males to females was nearly equal (Male: n=454, 48.86%; Female: n=475, 51.13%) as was the ratio of males to females in each grove (Montura: Female=41.54%, n=113, Male=46.32%, n=126; Pioneer: Female=42.68%, n=70, Male=39.02%, n=64; Port St. Lucie: Female=50.00%, n=225, Male=42.22%, n=190; Whidden Corner: Female=38.07%, n=67, Male=42.05%, n=74) (see **Table S1A** for further breakdown of male-female results).

The effect of sex on CLas Cq was tested and included in our model as CLas infection has been shown to affect *D. citri* behavior differently by sex. There was no effect of sex on CLas Cq

after controlling for grove and DNA extraction batch (F = 0.489, df =1, *P* = 0.484). This can also

be seen in **Figure 1**, as the blue and red colors show a random distribution of sex by Cq value

for each of the four groves.

In laboratory experiments, the color morph of *D. citri* was shown to be associated with

*C*Las titers (Hosseinzadeh et al. 2019a). While it was technically challenging to accurately assign

the color morphology to each specimen due to the insects being stored in ethanol prior to data

collection, no obvious associations between *C*Las titer and color morph emerged in these

insects (**Table S1B**). The lack of association between color morphology and *C*Las infection status

has previously been reported in the analysis of other grove-captured, Florida psyllid

populations (Hall et al. 2018).

4.   *Wolbachia* (wDi) has higher titer in non-infected *D. citri.*

A subset of *D. citri* samples from the initial DNA extraction were tested for wDi titer. The

total number of samples tested was 176 (Port St, Lucie = 46, Pioneer = 42, Montura = 41,

Whidden Corner = 47). A linear regression compared wDi Cq to *C*Las Cq , revealing no

correlation between *C*Las and wDi Cq among *D. citri* from these Florida citrus groves (R^2 =

0.0005) **(Figure S2)**. Additionally, wDi Cq was normalized against the *D. citri Wingless* gene

(Figure S3). A linear model assessed the relative effects of both grove and normalized wDi Cq

on *C*Las Cq. The analysis indicated that wDi was not significantly correlated with *C*Las Cq (F =

0.646, df = 1, *P* = 0.426). Moreover, the source grove had the most substantial effect on *C*Las

titer (F = 22.881, df = 3, *P* < 0.001, **Table S2**).

During the regression analysis, many wDi samples exhibited low Cq values when *C*Las Cq

= 40 (indicating non-infected samples, see **Figure S2)**. We investigated if wDi titer differed

significantly based on CLas presence (defined as Cq < 40) or absence (Cq = 40 Our analysis

revealed that when CLas is absent, wDi Cq was more likely to be below the median wDi Cq

value (Wilcoxon, n=119, P = 0.017). Furthermore, a Chi^2 test confirmed a significant difference

in the proportion of wDi samples with Cq above and below the median CLas Cq distribution,

when that distribution is divided by infection or no infection (X^2(1, n = 119) = 6.086, P = 0.013).

This exploratory testing appears to show that non-infected $D. citri$ tend to have higher wDi

titers compared to CLas-infected individuals.

Discussion

Natural variation in vector competency plays a role in disease epidemiology. Vector

competency is the potential of a vector to transmit a pathogen. In $D. citri$, vector competency is

in part regulated by $D. citri$ genes and proteins functioning at distinct tissue barriers which CLas

must overcome prior to inoculation into a new host tree (Ammar et al. 2018; Mann et al. 2018;

Kruse et al. 2017, 2018; Hosseinzadeh and Heck 2023). The infection of the host tree and citrus

variety has also been shown to play a role in modulating vector capacity (Ramsey et al. 2022;

Chin et al. 2020; Lombardi et al. 2024; Coletta-Filho et al. 2014). $D. citri$ acquires and transmits

CLas in an intracellular, circulative, propagative manner. Vector competency in our study was

measured through infection rate and CLas titer for each individual in each population studied.

Studies have shown that approximation of CLas titer in individual insects can predict vector

capacity. There is a strong positive correlation between CLas titer in the psyllid and

transmission efficiency (Ammar et al. 2018; Hosseinzadeh et al. 2019a). Broadly, our study

shows that there are population-level differences in *D. citri* vector competency that are likely

maintained through a combination of gene by environment interactions.

During circulative propagative transmission, *C*Las must cross the gut epithelial cell layer

to enter the body cavity of the psyllid and to be considered "acquired". Next, *C*Las must

circulate throughout the body and replicate in tissues, especially the salivary glands, where it

has been shown to reach high relative titers. From the salivary glands, transmission is possible

and can be inoculated into healthy citrus plants while feeding. The efficiency of the first and

second steps can be approximated by using qPCR to detect *C*Las titer in adults that have

acquired *C*Las, while the third step can only be measured by careful transmission studies. The

development of vector-competent adults requires *C*Las acquisition to occur during the psyllid's

nymphal stages (Inoue et al. 2009; Ammar et al. 2016; Igwe et al. 2022). This study

approximates the grove-wide vector capacity of four *D. citri* populations through

measurements of *C*Las titer in individual adult insects. A key assumption of this study is that the

majority of trees in all citrus groves of Florida are infected with *C*Las and thus, all adult psyllids

will have fed on *C*Las-infected citrus as nymphs and are competent transmitters as adults.

However, even if this assumption did not hold for every individual in our dataset, multiple

studies on vector acquisition rates in the lab have recorded <100% acquisition, despite feeding

on known infected material (Pelz-Stelinski et al. 2010; De Souza Pacheco et al. 2020). Samples

that do test positive for *C*Las are competent transmitters and fall into categories of low to high

vector capacity, as determined by their mean *C*Las titer (Ammar et al. 2018).

Many factors may influence *D. citri* vector competency and the variation we observed in

our study. Prior work has shown that host switching influences variation in *D. citri* vector

competency (Ramsey et al. 2022; Lombardi et al. 2024). While grove-collected sampling makes

it difficult to control or understand all variables that may influence CLas titer in psyllids, all four

D. citri populations were sampled from the same citrus variety (*Citrus sinensis*), within the same

week in October, 2018, so it represents a snapshot in time of the potential of *D. citri* in these

regions to acquire and transmit. Additionally, the role that symbionts like wDi might play in

mediating vector-pathogen interactions is unclear, so the effect of wDi on CLas titer from these

four fields was also tested. Our results show that *D. citri* has higher wDi titer in CLas-negative

individuals, while field site has the strongest effect on CLas titer, and sex does not have any

effect. The field effect could be related to the heritability of *D. citri* genes involved in vector

competency, the management regimen, or transient access to and feeding on non-citrus host

plants such as *Murraya spp.* in the grove agro-ecosystem. More work is required to fully explore

these hypotheses.

Abandoned groves have also been shown to maintain high *D. citri* populations that

travel frequently between managed and abandoned areas (Sétamou et al. 2022; Boina et al.

2009) with equally high CLas titers and infection rates as alive and managed groves (Hall 2018).

Prior to 2010, Tiwari et al. (Tiwari et al. 2010) studied the *D. citri-C*Las infection rate and found

1.9 % in abandoned and 1.2 % in managed grove populations while the mean CLas Cq value was

32.52 in abandoned and 25.95 in managed groves. Over 10 years after Tiwari et al. collected

their data, the infection rates found by our study in the least infected, but still managed, grove

are at least 1.5x higher incidence while our other three groves have at least 50x higher infection

rates. This is undoubtedly due to the vastly increased number of both managed and abandoned

infected trees in Florida over the past ten years. A study by De Souza Pacheco et al. (De Souza

Pacheco et al. 2020) shows that when *D. citri* are raised on *C*Las-infected citrus for four

consecutive generations in a greenhouse setting, the percent-infection and the titer in

individuals increased with each generation from 42% to 100% infection incidence, and 47.9 to

5,390 *C*Las cells per 100 ng of DNA. These results suggest that interruptions in vector

generations (such as through regular insecticide applications or during seasons of low flush) in

the field may help keep *C*Las titer and incidence lower in vector populations. The increase in

infection incidence seen in Florida groves despite various management innovations over the

last decade is a testament to both the effect of continuous populations of *D. citri* replicating on

infected citrus and reference to the accumulating impact HLB has wrought on the Florida citrus

industry.

Vector competency of *D. citri* populations and individuals varies widely in the field as we

have seen through this study and others (Capoor et al. 1974; Coy and Stelinski 2015; Hall 2018;

Ukuda-Hosokawa et al. 2015; Ammar et al. 2018; Sétamou et al. 2016; Xu et al. 1988), in

addition to laboratory-kept colonies (Inoue et al. 2009; Hosseinzadeh et al. 2019b; De Souza

Pacheco et al. 2020). In agreement with our results, Chu et al (Chu et al. 2016) show that three

wild populations of *D. citri* (including a total of 123 insects collected in 2015) vary in their *C*Las

titer and *C*Las infection incidence, and that like our study, population (grove site) has the

strongest effect on all their models. Coy and Stelinski collected their samples in 2014 from six

locations and showed large vector capacity variation ranging from 35-100%, but they collected

less than 300 samples in total (Coy and Stelinski 2015). Hall published results from two studies

with multiple time points and similar numbers of samples as the current study, but only from

one location, and again showed strong variability (Hall 2018). This study not only adds to the

literature on in-field vector variability, but it fills a gap in the literature by studying multiple

locations with over 150 samples from each location, in 2018 – the most recent timepoint for a

study like this, available to our knowledge in the literature. One study focusing on

overwintering survival rates assessed psyllids in northern Florida from 2017-2018, but their

measurements of *C*Las titer were based on sticky card captures and did not record any psyllids

from commercial groves (Martini et al. 2020). Finally, a recent paper, published in 2023 by

Ebert et al., surveyed multiple commercial sites in Florida. However, their samples were

collected between 2008 and 2012, (Ebert et al. 2023) making our study the latest published

record of commercial psyllid vector capacity variability in Florida and perhaps the last one

before the wide-spread adoption of injectable oxytetracycline (OTC) for HLB management in

Florida. Now that direct injection of the OTC is becoming a wide-spread HLB management tactic

in Florida, understanding the impact of OTC on *C*Las titer in psyllids will be important to

understand if the use of OTC will indirectly lower vector competency in psyllid populations.

While grove is highly correlated in the current study with *C*Las titer, the base differences

may still come down to molecular or genetic differences between the psyllid populations, *C*Las

strains, other endosymbionts, and unknown environmental factors. The ability of *D. citri* to

acquire and transmit *C*Las may be genetically linked (Ammar et al. 2018) and studies that

separate acquisition and transmission (Ammar et al. 2016; Ukuda-Hosokawa et al. 2015;

Ammar et al. 2018; Xu et al. 1988) agree that these are likely two separate traits, but that they

are correlated, such that higher acquisition rates often lead to more efficient transmission

(Ammar et al. 2018). The current study used *C*Las titer as a measure of acquisition in adult *D.*

citri, (which may also include effects of internal *C*Las replication) but the high numbers of

infected individuals (infection rate) also suggests transmission of CLas is occurring in the field

and that adult psyllids acquired as nymphs from infected trees.

The relationship between wDi titer on CLas in *D. citri* requires further investigation.

However, in a related psyllid, *Cacopsylla pyri* (pear psyllid) studied by Serbina et al (Serbina et

al. 2022), wDi titer was reduced in males. Several lab studies investigated interactions among

wDi, *D. citri* and CLas. Hosseinzadeh et al. (Hosseinzadeh et al. 2019b) found that wDi titer

varied between sexes, and *D. citri* had higher titer in males that were infected with CLas than

non-infected males, though wDi did not significantly differ between CLas infected and non-

infected populations as a whole. Others found that wDi titer was significantly higher in CLas-

negative *D. citri*, except when controlling for population effect (Chu et al. 2019; Ren et al. 2018;

Chu et al. 2016). Fagen et al (Fagen 2012) showed the opposite trend, that CLas has a negative

relationship with other *D. citri* obligate endosymbionts, but a positive relationship to wDi

regarding titer. Our results, using insects collected from citrus groves instead of greenhouse

and chamber-raised populations, contradict some of these previous lab-based results. We saw

100% wDi infection rate among our individuals tested, and the titer of wDi tended to be higher

in CLas non-infected *D. citri* individuals. These contradictions underscore the importance to not

over-generalize the results from lab studies and the importance of field-based inquiries on

vector-microbe interactions.

Conclusion

This snapshot assessment of vector populations showed grove-specific differences in

CLas infection. Specifically, infection rate and titer of CLas within individual *D. citri* varied by

grove site, exhibiting three patterns of vector competency. However, no conclusions could be

drawn on the components of grove site that influenced CLas titer in *D. citri.* Additionally, while

there was no impact of sex on CLas titer, wDi titers were higher in non-infected individuals.

These results contradict some studies using lab-reared psyllid colonies. Unknown

environmental factors may contribute to titer levels in insects in the grove and we stress that

caution is necessary when applying and interpreting results from lab-reared insects to field-

based studies and applications. The data suggest that future studies should prioritize

understanding the effect of grove management practices, including the potential benefits of

using non-citrus plants that are hosts of *D. citri* or of its natural enemies in the grove.

Additionally, understanding the contributions that both environment and psyllid genes – the G

x E interaction – have on vector competency will help make all future genetic, genomic and

field-based studies of HLB more informed when developing control strategies or other

programs aimed at blocking CLas transmission by *D. citri.*

Acknowledgements

We thank Tim Eyrich and Shannon Leahy (Southern Gardens), Dr. David Hall and Kathy Moulton

(USDA Agricultural Research Service, retired), for their help, organization, and outreach to the

growers which enabled us to collect live psyllids from grove sites in Florida, USA.

Financial Support

Funding for this research was graciously provided by NIFA-AFRI Predoctoral Fellowship award #

2021-67011-35143 for Marina Mann, USDA ARS CRIS project 8062-22410-007-000-D, and USDA

NIFA grants 2020-70029-33176 and 2022-70029-38503.

Competing Interests

The authors declare that they have no competing interests.

References

Ammar, E.-D., Hall, D. G., Hosseinzadeh, S., and Heck, M. 2018. The quest for a non-vector psyllid:
Natural variation in acquisition and transmission of the huanglongbing pathogen '*Candidatus*
Liberibacter asiaticus' by Asian citrus psyllid isofemale lines. PLoS ONE. 13:e0195804.
421

422   Ammar, E.-D., Ramos, J. E., Hall, D. G., Dawson, W. O., and Shatters, R. G. 2016. Acquisition, Replication
423   and Inoculation of *Candidatus* Liberibacter asiaticus following Various Acquisition Periods on
424   Huanglongbing-Infected Citrus by Nymphs and Adults of the Asian Citrus Psyllid. PLoS ONE.
425   11:e0159594.
426

427   Boina, D. R., Meyer, W. L., Onagbola, E. O., and Stelinski, L. L. 2009. Quantifying Dispersal of *Diaphorina*
428   *citri* (Hemiptera: Psyllidae) by Immunomarking and Potential Impact of Unmanaged Groves on
429   Commercial Citrus Management. Environ Entomol. 38:1250–1258.
430

431   Bové, J. M., Garnier, M., Ahlawat, Y. S., Chakraborty, N. K., and Varma, A. 1993. Detection of the Asian
432   Strains of the Greening BLO by DNA-DNA Hybridization in Indian Orchard Trees and Malaysian
433   Diaphorina citri Psyllids. International Organization of Citrus Virologists Conference Proceedings (1957-
434   2010). 12 Available at: https://escholarship.org/uc/item/0ng306x8 [Accessed July 19, 2023].
435

436   Capoor, S. P., Rao, D. G., and Viswanath, S. M. 1974. Greening Disease of Citrus in the Deccan Trap
437   Country and its Relationship with the Vector, *Diaphorina citri* Kuwayama. International Organization of
438   Citrus Virologists Conference Proceedings (1957-2010). 6 Available at:
439   https://escholarship.org/uc/item/6rm6x1tw [Accessed July 19, 2023].
440

441   Chen, Q., Li, Z., Liu, S., Chi, Y., Jia, D., and Wei, T. 2022. Infection and distribution of *Candidatus*
442   Liberibacter asiaticus in citrus plants and psyllid vectors at the cellular level. Microbial Biotechnology.
443   15:1221–1234.
444

445   Chin, E. L., Ramsey, J. S., Mishchuk, D. O., Saha, S., Foster, E., Chavez, J. D., et al. 2020. Longitudinal
446   Transcriptomic, Proteomic, and Metabolomic Analyses of *Citrus sinensis* (L.) Osbeck Graft-Inoculated
447   with " *Candidatus* Liberibacter asiaticus." J. Proteome Res. 19:719–732.
448

449   Chu, C., Hoffmann, M., Braswell, W. E., and Pelz-Stelinski, K. S. 2019. Genetic variation and potential
450   coinfection of *Wolbachia* among widespread Asian citrus psyllid (*Diaphorina citri* Kuwayama)
451   populations. Insect Science. 26:671–682.
452

453 Chu, C.-C., Gill, T. A., Hoffmann, M., and Pelz-Stelinski, K. S. 2016. Inter-Population Variability of
454 Endosymbiont Densities in the Asian Citrus Psyllid (*Diaphorina citri* Kuwayama). Microb Ecol. 71:999–
455 1007.
456

457 Coletta-Filho, H. D., Daugherty, M. P., Ferreira, C., and Lopes, J. R. S. 2014. Temporal Progression of
458 '*Candidatus* Liberibacter asiaticus' Infection in Citrus and Acquisition Efficiency by *Diaphorina citri*.
459 Phytopathology®. 104:416–421.
460

461 Coy, M. R., and Stelinski, L. L. 2015. Great Variability in the Infection Rate of '*Candidatus* Liberibacter
462 Asiaticus' in Field Populations of *Diaphorina citri* (Hemiptera: Liviidae) in Florida. Florida Entomologist.
463 98:356–357.
464

465 De Souza Pacheco, I., Manzano Galdeano, D., Spotti Lopes, J. R., and Machado, M. A. 2020. Development
466 on Infected Citrus over Generations Increases Vector Infection by '*Candidatus* Liberibacter Asiaticus in
467 *Diaphorina citri*.' Insects. 11:469.
468

469 Ebert, T. A., Shawer, D., Brlansky, R. H., and Rogers, M. E. 2023. Seasonal Patterns in the Frequency of
470 Candidatus Liberibacter Asiaticus in Populations of *Diaphorina citri* (Hemiptera: Psyllidae) in Florida.
471 Insects. 14:756.
472

473 Fagen, J. R. 2012. Characterization of the Relative Abundance of the Citrus Pathogen *Ca.* Liberibacter
474 Asiaticus in the Microbiome of Its Insect Vector, *Diaphorina citri*, using High Throughput 16S rRNA
475 Sequencing. TOMICROJ. 6:29–33.
476

477 Garnier, M., and Bove, J. 1983. Transmission of the organism associatied with Citrus Greening disease
478 from sweet orange to periwinkle by dodder. Phytopathology. 73:1358–1363.
479

480 Ghosh, D., Kokane, S., Savita, B. K., Kumar, P., Sharma, A. K., Ozcan, A., et al. 2023. Huanglongbing
481 Pandemic: Current Challenges and Emerging Management Strategies. Plants. 12:160.
482

483 Guidolin, A. S., and Cônsoli, F. L. 2013. Molecular Characterization of *Wolbachia* Strains Associated with
484 the Invasive Asian Citrus Psyllid *Diaphorina citri* in Brazil. Microb Ecol. 65:475–486.
485

486 Hall, D. G. 2018. Incidence of "*Candidatus* Liberibacter asiaticus" in a Florida population of Asian citrus
487 psyllid. J Appl Entomol. 142:97–103.
488

489 Higgins, S. A., Mann, M., and Heck, M. 2022. Strain Tracking of '*Candidatus* Liberibacter asiaticus', the
490 Citrus Greening Pathogen, by High-Resolution Microbiome Analysis of Asian Citrus Psyllids.
491 Phytopathology®. 112:2273–2287.
492

493 Hoffmann, M., Coy, M. R., Gibbard, H. N. K., and Pelz-Stelinski, K. S. 2014. *Wolbachia* Infection Density in
494 Populations of the Asian Citrus Psyllid (Hemiptera: Liviidae). Environ Entomol. 43:1215–1222.
495

496 Hosseinzadeh, S., and Heck, M. 2023. Variations on a theme: factors regulating interaction between
497 *Diaphorina citri* and "*Candidatus* Liberibacter asiaticus" vector and pathogen of citrus huanglongbing.
498 Current Opinion in Insect Science. 56:101025.
499

500 Hosseinzadeh, S., Ramsey, J., Mann, M., Bennett, L., Hunter, W. B., Shams-Bakhsh, M., et al. 2019a.
501 Color morphology of *Diaphorina citri* influences interactions with its bacterial endosymbionts and
502 '*Candidatus* Liberibacter asiaticus' ed. Janice L. Bossart. PLoS ONE. 14:e0216599.
503

504 Hosseinzadeh, S., Shams-Bakhsh, M., Mann, M., Fattah-Hosseini, S., Bagheri, A., Mehrabadi, M., et al.
505 2019b. Distribution and Variation of Bacterial Endosymbiont and "*Candidatus* Liberibacter asiaticus"
506 Titer in the Huanglongbing Insect Vector, *Diaphorina citri* Kuwayama. Microb Ecol. 78:206–222.
507

508 Igwe, D. O., Higgins, S. A., and Heck, M. 2022. An Excised Leaf Assay to Measure Acquisition of '
509 *Candidatus* Liberibacter asiaticus' by Psyllids Associated with Citrus Huanglongbing Disease.
510 Phytopathology®. 112:69–75.
511

512 Inoue, H., Ohnishi, J., Ito, T., Tomimura, K., Miyata, S., Iwanami, T., et al. 2009. Enhanced proliferation
513 and efficient transmission of *Candidatus* Liberibacter asiaticus by adult *Diaphorina citri* after acquisition
514 feeding in the nymphal stage. Annals of Applied Biology. 155:29–36.
515

516 Jain, M., Fleites, L. A., and Gabriel, D. W. 2017. A Small *Wolbachia* Protein Directly Represses Phage Lytic
517 Cycle Genes in "*Candidatus* Liberibacter asiaticus" within Psyllids ed. Katherine McMahon. mSphere.
518 2:e00171-17.
519

520 Kassambara, A. 2020. ggpubr: "ggplot2" based publication ready plots. (R package version 0.4.0).
521 Available at: https://CRAN.R-project.org/package=ggpubr.
522

523 Kruse, A., Fattah-Hosseini, S., Saha, S., Johnson, R., Warwick, E., Sturgeon, K., et al. 2017. Combining
524 'omics and microscopy to visualize interactions between the Asian citrus psyllid vector and the
525 Huanglongbing pathogen *Candidatus* Liberibacter asiaticus in the insect gut ed. PLoS ONE. 12:e0179531.
526

527 Kruse, A., Ramsey, J. S., Johnson, R., Hall, D. G., MacCoss, M. J., and Heck, M. 2018. *Candidatus*
528 Liberibacter asiaticus Minimally Alters Expression of Immunity and Metabolism Proteins in Hemolymph
529 of *Diaphorina citri* , the Insect Vector of Huanglongbing. J. Proteome Res. 17:2995–3011.
530

531 Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B. 2017. **lmerTest** Package: Tests in Linear Mixed
532 Effects Models. J. Stat. Soft. 82 Available at: http://www.jstatsoft.org/v82/i13/ [Accessed August 24,
533 2023].

Lombardi, R. L., Ramsey, J. S., Mahoney, J. E., MacCoss, M. J., Heck, M. L., and Slupsky, C. M. 2024. Longitudinal Transcriptomic, Proteomic, and Metabolomic Response of *Citrus sinensis* to *Diaphorina citri* Inoculation of *Candidatus* Liberibacter asiaticus. J. Proteome Res. :acs.jproteome.3c00485.

Manjunath, K. L., Halbert, S. E., Ramadugu, C., Webb, S., and Lee, R. F. 2008. Detection of '*Candidatus* Liberibacter asiaticus' in *Diaphorina citri* and Its Importance in the Management of Citrus Huanglongbing in Florida. Phytopathology®. 98:387–396.

Mann, M., Fattah-Hosseini, S., Ammar, E.-D., Stange, R., Warrick, E., Sturgeon, K., et al. 2018. *Diaphorina citri* Nymphs Are Resistant to Morphological Changes Induced by "*Candidatus* Liberibacter asiaticus" in Midgut Epithelial Cells ed. Marvin Whiteley. Infect Immun. 86:e00889-17.

Martini, X., Malfa, K., Stelinski, L. L., Iriarte, F. B., and Paret, M. L. 2020. Distribution, Phenology, and Overwintering Survival of Asian Citrus Psyllid (Hemiptera: Liviidae), in Urban and Grove Habitats in North Florida ed. Blake Bextine. Journal of Economic Entomology. 113:1080–1087.

Naeem, A., Afzal, M. B. S., Freed, S., Hafeez, F., Zaka, S. M., Ali, Q., et al. 2019. First report of thiamethoxam resistance selection, cross resistance to various insecticides and realized heritability in Asian citrus psyllid *Diaphorina citri* from Pakistan. Crop Protection. 121:11–17.

Nakabachi, A., Ueoka, R., Oshima, K., Teta, R., Mangoni, A., Gurgui, M., et al. 2013. Defensive Bacteriome Symbiont with a Drastically Reduced Genome. Current Biology. 23:1478–1484.

Neupane, S., Bonilla, S. I., Manalo, A. M., and Pelz-Stelinski, K. S. 2022. Complete de novo assembly of *Wolbachia* endosymbiont of *Diaphorina citri* Kuwayama (Hemiptera: Liviidae) using long-read genome sequencing. Sci Rep. 12:125.

Pardo, S., Martínez, A. M., Figueroa, J. I., Chavarrieta, J. M., Viñuela, E., Rebollar-Alviter, Á., et al. 2018. Insecticide resistance of adults and nymphs of Asian citrus psyllid populations from Apatzingán Valley, Mexico: Resistance of *D. citri* to bifenthrin, malathion and chlorpyrifos. Pest. Manag. Sci. 74:135–140.

Pelz-Stelinski, K. S., Brlansky, R. H., Ebert, T. A., and Rogers, M. E. 2010. Transmission Parameters for *Candidatus* Liberibacter asiaticus by Asian Citrus Psyllid (Hemiptera: Psyllidae). J. Econ. Entom. 103:1531–1541.

R Core Team. 2022. R: A language and environment for statistical computing. Available at: https://www-R-project.org/.

Ramsey, J. S., Ammar, E.-D., Mahoney, J. E., Rivera, K., Johnson, R., Igwe, D. O., et al. 2022. Host Plant
Adaptation Drives Changes in *Diaphorina citri* Proteome Regulation, Proteoform Expression, and
Transmission of '*Candidatus* Liberibacter asiaticus', the Citrus Greening Pathogen. Phytopathology®.
112:101–115.

Ramsey, J. S., Johnson, R. S., Hoki, J. S., Kruse, A., Mahoney, J., Hilf, M. E., et al. 2015. Metabolic Interplay
between the Asian Citrus Psyllid and Its *Profftella* Symbiont: An Achilles' Heel of the Citrus Greening
Insect Vector ed. Murad Ghanim. PLoS ONE. 10:e0140826.

Rao, C. N., Shivankar, V. J., Sandnya, D., David, K. J., and Dhengre, V. N. 2014. Insecticide resistance in
field populations of Asian Citrus psyllid, *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae). Pesticide
Research Journal. 26:42–47.

Razi, M. F., Keremane, M. L., Ramadugu, C., Roose, M., Khan, I. A., and Lee, R. F. 2014. Detection of
Citrus Huanglongbing-Associated '*Candidatus* Liberibacter asiaticus' in Citrus and *Diaphorina citri* in
Pakistan, Seasonal Variability, and Implications for Disease Management. Phytopathology®. 104:257–
268.

Ren, S.-L., Li, Y.-H., Ou, D., Guo, Y.-J., Qureshi, J. A., Stansly, P. A., et al. 2018. Localization and dynamics
of *Wolbachia* infection in Asian citrus psyllid *Diaphorina citri* , the insect vector of the causal pathogens
of Huanglongbing. MicrobiologyOpen. 7:e00561.

Saha, S., Hunter, W. B., Reese, J., Morgan, J. K., Marutani-Hert, M., Huang, H., et al. 2012. Survey of
Endosymbionts in the *Diaphorina citri* Metagenome and Assembly of a *Wolbachia* wDi Draft Genome ed.
Dan Zilberstein. PLoS ONE. 7:e50067.

Serbina, Š. L., Gajski, D., Malenovský, I., Corretto, E., Schuler, H., and Dittmer, J. 2022. *Wolbachia*
infection dynamics in a natural population of the pear psyllid *Cacopsylla pyri* (Hemiptera: Psylloidea)
across its seasonal generations. Sci Rep. 12:16502.

Sétamou, M., Alabi, O. J., Kunta, M., Dale, J., and Da Graça, J. V. 2020. Distribution of *Candidatus*
Liberibacter asiaticus in Citrus and the Asian Citrus Psyllid in Texas Over a Decade. Plant Disease.
104:1118–1126.

Sétamou, M., Alabi, O. J., Kunta, M., Jifon, J. L., and Da Graça, J. V. 2016. Enhanced Acquisition Rates of '
Candidatus Liberibacter asiaticus' by the Asian Citrus Psyllid (Hemiptera: Liviidae) in the Presence of
Vegetative Flush Growth in Citrus. J Econ Entomol. 109:1973–1978.

Sétamou, M., Tarshis Moreno, A., and Patt, J. M. 2022. Source or Sink? The Role of Residential Host
Plants in Asian Citrus Psyllid Infestation of Commercial Citrus Groves ed. Arash Rashed. Journal of
Economic Entomology. 115:438–445.

Shippy, T. D., Hosmani, P. S., Flores-Gonzalez, M., Mann, M., Miller, S., Weirauch, M. T., et al. 2019. *Diaci*
v3.0: Chromosome-level assembly, de novo *transcriptome and manual annotation of* Diaphorina citri,
insect vector of Huanglongbing. Genomics. Available at: http://biorxiv.org/lookup/doi/10.1101/869685
[Accessed August 24, 2023].
619

620   Tiwari, S., Lewis-Rosenblum, H., Pelz-Stelinski, K., and Stelinski, L. L. 2010. Incidence of *Candidatus*
621   Liberibacter asiaticus Infection in Abandoned Citrus Occurring in Proximity to Commercially Managed
622   Groves. jnl. econ. entom. 103:1972–1978.
623

624   Ukuda-Hosokawa, R., Sadoyama, Y., Kishaba, M., Kuriwada, T., Anbutsu, H., and Fukatsu, T. 2015.
625   Infection Density Dynamics of the Citrus Greening Bacterium "*Candidatus* Liberibacter asiaticus" in Field
626   Populations of the Psyllid *Diaphorina citri* and Its Relevance to the Efficiency of Pathogen Transmission
627   to Citrus Plants ed. H. Goodrich-Blair. Appl Environ Microbiol. 81:3728–3736.
628

629   Vázquez-García, M., Velázquez-Monreal, J., Medina-Urrutia, V. M., Cruz-Vargas, C. D. J., Sandoval-
630   Salazar, M., Virgen-Calleros, G., et al. 2013. Insecticide Resistance in Adult *Diaphorina citri* Kuwayama
631   from Lime Orchards in Central West Mexico. Southwestern Entomologist. 38:579–596.
632

633   Venkatesan, T., Chethan, B. R., and Mani, M. 2022. Insecticide Resistance and Its Management in the
634   Insect Pests of Horticultural Crops. In *Trends in Horticultural Entomology*, ed. M. Mani. Singapore:
635   Springer Nature Singapore, p. 455–490. Available at: https://link.springer.com/10.1007/978-981-19-
636   0343-4_14 [Accessed August 24, 2023].
637

638   Walter, A. J., Hall, D. G., and Duan, Y. P. 2012. Low Incidence of ' *Candidatus* Liberibacter asiaticus' in
639   *Murraya paniculata* and Associated *Diaphorina citri*. Plant Disease. 96:827–832.
640

641   Wickham, H. 2-16. ggplot2: Elegant graphics for data analysis.
642

643   Wulff, N., Daniel, B., Sassi, R., Moreira, A., Bassanezi, R., Sala, I., et al. 2020. Incidence of *Diaphorina citri*
644   Carrying *Candidatus* Liberibacter asiaticus in Brazil's Citrus Belt. Insects. 11:672.
645

646   Xu, C. F., Xia, Y. H., Li, K. B., and Ke, C. 1988. Further Study of the Transmission of Citrus Huanglungbin by
647   a Psyllid, *Diaphorina citri* Kuwayama. International Organization of Citrus Virologists Conference
648   Proceedings (1957-2010). 10 Available at: https://escholarship.org/uc/item/0w42q0r7 [Accessed July 19,
649   2023].
650