1 **Thymine DNA glycosylase combines sliding, hopping, and nucleosome interactions to**
2 **efficiently search for 5-formylcytosine.**
3

Brittani L. Schnable[1,2], Matthew A. Schaich[2,3], Vera Roginskaya[2,3], Liam P. Leary[2,3], Tyler M.
Weaver[4], Bret D. Freudenthal[4], Alexander C. Drohat[5], and Bennett Van Houten[1,2,3]

[1]Molecular Biophysics and Structural Biology Graduate Program, University of Pittsburg, PA
15260, USA
[2]UPMC Hillman Cancer Center, University of Pittsburgh, Pittsburgh, PA 15213, USA
[3]Department of Pharmacology and Chemical Biology, School of Medicine, University of
Pittsburgh, Pittsburgh, PA 15213, USA
[4]Department of Biochemistry and Molecular Biology, Department of Cancer Biology, University
of Kansas Medical Center, Kansas City, KS 66160, USA
[5]Department of Biochemistry and Molecular Biology, University of Maryland School of Medicine,
Baltimore, Maryland 21201, USA.

Corresponding author
Bennett Van Houten, PhD
5117 Centre Ave
Hillman Cancer Center
Research Pavilion, Suite 2.6
Pittsburgh, PA 15213
Phone: 412-623-7762
Fax: 412-623-7761
Email: vanhoutenb@upmc.edu

Abstract

Base excision repair is the main pathway involved in active DNA demethylation. 5-

formylctyosine and 5-carboxylcytosine, two oxidized moieties of methylated cytosine, are

recognized and removed by thymine DNA glycosylase (TDG) to generate an abasic site. Using

single molecule fluorescence experiments, we studied TDG in the presence and absence of 5-

formylctyosine. TDG exhibits multiple modes of linear diffusion, including hopping and sliding, in

search of a lesion. We probed TDG active site variants and truncated N-terminus revealing how

these variants alter the lesion search and recognition mechanism of TDG. On DNA containing

an undamaged nucleosome, TDG was found to either bypass, colocalize with, or encounter but

not bypass the nucleosome. However, truncating the N-terminus reduced the number of

interactions with the nucleosome. Our findings provide unprecedented mechanistic insights into

how TDG searches for DNA lesions in chromatin.

Introduction

DNA can be damaged by a variety of endogenous and exogenous sources and these potentially mutagenic bases are removed via base excision repair (BER)[1]. In humans, BER is initiated by one of eleven damage specific DNA glycosylases that recognize a lesion and cleave the glycosidic bond to leave an abasic site. The repair is completed by the actions of several proteins that restore the correct base pair. Thymine DNA glycosylase (TDG) is a monofunctional glycosylase that initiates BER by recognizing a specific repertoire of lesions, including 5-methylcytosine (5mC) that has been deaminated to form a G:T mismatch or 5mC that has been oxidized by the ten eleven translocation (TET) family of enzymes into 5-formylcytosine (5fC) and 5-carboxylcytosine (5caC)[2-5]. Interestingly, TDG is the only known mammalian DNA glycosylase that creates an embryonic lethal phenotype in mice upon depletion: presumably due to the significant role that TDG plays in active demethylation, in addition to the action on environmental DNA damage[6,7].

While TDG is known to preferentially remove T from G:T mismatches in the context of a CpG, other substrates do not show this sequence context dependence [8,9]. One such lesion is 5fC, which has an abundance of 6×10^4 per human genome, and the removal of 5fC was similar in the context of a CpG and outside a CpG[10-12]. Previous studies have shown that N140 of TDG is essential for the catalytic activity on 5fC because of interactions with the water nucleophile. The N140A variant greatly reduces the catalytic activity 16,000-fold[5]. It has been shown that R275 plays an essential role in catalysis by filling in the void in the DNA that is generated once the substrate nucleobase is flipped into the active site[13]. In addition to TDG specificity for G:T mismatches and 5fC, TDG is able to bind to DNA nonspecifically with an affinity of 293 ± 64 nM and has a footprint of 10 base pairs[14,15]. While the roles for these key active site amino acids in catalysis is well-established, their role in mediating TDG searching for DNA lesions in the presence of excess unmodified DNA is unknown.

A major unexplored consideration for any glycosylase search process in a cell is the

presence of chromatin. DNA packaged into nucleosomes would interrupt a DNA sliding

mechanism and inhibit the accessibility of specific sites in the absence of nucleosome sliding

and/or remodeling. Among DNA glycosylases this phenomenon is particularly relevant to TDG,

as the methylated DNA where its substrates are formed are associated with higher degree of

chromatinization[16]. TDG has also been shown to directly interact with nucleosome core

particles, however kinetic studies also demonstrated TDG activity was only reduced ~two-fold

for uracil substrates present on a nucleosome core particle[17,18].

Here, we utilized single-molecule analysis of DNA-binding proteins from nuclear extracts

(SMADNE)[19] to directly observe the search process of fluorescently labeled TDG in real-time on

DNA containing 5fC substrates or nucleosome core particles. By observing TDG scanning the

DNA as it searches for 5fC, we determined that TDG searches for damaged sites using a

combination of sliding and hopping and that TDG diffusivity was highly dependent on DNA

tension. This combined search mechanism was also observed on DNA containing NCPs. In

some instances, TDG hopped over nucleosomes to continue searching the DNA, but in other

cases it approached the NCP but did not bypass. We have also found that specific amino acid

side chains that are essential for catalytic activity also impact the dwell time and the ability of

TDG to diffuse on DNA. Additionally, although ΔN-term TDG (residues 82-410) is catalytically

active, we demonstrate that the N-terminus plays a significant role in binding and searching

unmodified DNA as well as playing a key role in engaging undamaged NCPs.

Results

TDG binds DNA specifically and nonspecifically.

TDG-HaloTag was overexpressed in U2OS cells, and the nuclear extracts were

quantified using western blots and SDS gels to determine a 60-fold difference in overexpressed

fluorescently tagged WT TDG to endogenous TDG (Fig. S2). To determine the binding behavior

of TDG on DNA containing a lesion of interest, we initially performed the single-molecule

imaging with TDG-HaloTag-JF635 and DNA containing 5fC generated by nick translation (Fig.

1a, c). Analysis of the resulting kymographs revealed complex binding dynamics, including non-

motile (12.6%) and motile TDG binding events (i.e., exhibited 1D diffusion, 87.4%). Further co-

localization analysis of red TDG-HaloTag-JF635 with the blue 5fC fiducial marker in the DNA

(fluorescein dUTP) revealed the stationary events were primarily TDG bound to 5fC sites (Fig.

1d). The motile events primarily represent TDG binding to the DNA nonspecifically. Of note,

diffusion events that are less than 0.3 seconds are not detected. Fitting the motile events to a

cumulative residence time distribution plot (CRTD) with a one-phase exponential decay, a

binding lifetime of 7.5 ± 0.3 seconds was obtained (Fig. 1d-e, Table 1). Only motile events with a

binding lifetime of 7.9 ± 0.1 sec were observed in additional single-molecule imaging

experiments with TDG-HaloTag-JF635 and unmodified DNA, supporting that the stationary

events reflect TDG bound to a lesion (Fig. 1f-g, Table 1). In contrast to the motile events, the

CRTD plot of stationary events that co-localized with the 5fC fiducial marker were fit with a two-

phase exponential decay with binding lifetimes of 14.7 ± 1.7 and 101.1 ± 7.1 seconds (Fig. 1d-e,

Table 1). We hypothesize that these longer TDG lifetimes represent binding to and cleaving the

5fC. This is consistent with previous kinetic experiments that indicate it takes ~ 68 secs (k_{max} of

0.61 min[-1]) to cleave 5fC at 22°C, plus the time for TDG to dissociate from the generated abasic

site at[2]. Notably, the shorter 14.7 ± 1.7 second binding events, that co-localized with the 5fC

fiducial marker, exclusively occur after a long-lived event. These shorter events likely represent

TDG binding to an abasic site generated by its catalytic function or a nicked abasic site

processed by apurinic/apyrimidinic endonuclease 1 (APE1) in the extracts[19].

Since the nick translated DNA contains multiple sites with a region of 5fC, we generated

a 12 kb DNA substrate that contains a single center-positioned 5fC to more easily observe if

lifetime differs over time, indicating turnover. We then performed the correlative optical

tweezers-fluorescence microscopy (CTFM) experiment using TDG-HaloTag-JF635 with the

single 5fC DNA. A CRTD plot of stationary events that co-localized with the 5fC were fit with a

two-phase exponential decay with binding lifetimes of 0.67± 0.06 sec and 41.8 ± 12.2 sec (Fig.

1d-e,h-i, Table 1), consistent with the short- and long-lived binding lifetimes observed on the

DNA substrate containing 10 5fC sites (Fig. 1 e). These data further indicate that long-lived

events represent TDG binding to the substrate and the short-lived events are mostly likely TDG

rebinding to the catalytic product abasic site (Fig. 1h-i).

TDG exhibits mixed modes of linear diffusion during its search for DNA damage.

After observing TDG bind and move on DNA nonspecifically, we sought to determine the

mode of linear diffusion TDG uses on DNA. Linear diffusion occurs through sliding and hopping

mechanisms that have distinct behaviors. Sliding is when a protein tracks the DNA helix without

dissociating, leading to coupled translational and rotational diffusion[20,21]. Alternatively, hopping is

when a protein micro-dissociates as it translocates along the DNA, leading to uncoupled

translational and rotational diffusions. Notably, the sliding and hopping mechanisms can be

distinguished by determining whether the diffusion coefficient of TDG changes as a function of

salt concentration: diffusion via sliding is unaffected by salt concentration, whereas diffusion for

hopping is sensitive to salt concentration[22-24]. To further differentiate whether TDG uses a

sliding, hopping, or multimodal 1D diffusion, we performed additional single-molecule imaging

experiments with TDG-HaloTag-JF635 and unmodified DNA at varying ionic concentrations (75

to 150 mM NaCl). We observed an increase in the diffusion from 1.6×10^{-2} $\mu m^2/s$ at 75 mM NaCl,

to 2.8×10^{-2} $\mu m^2/s$ at 100 mM NaCl, and 3.1×10^{-2} $\mu m^2/s$ at 150 mM NaCl (Fig. 2a, Table 1,

Supplemental Fig. 5a-d). The theoretical limit of diffusion of TDG-HaloTag sliding along the DNA

was calculated to be 2.56×10^{-1} $\mu m^2/s$ (see Supplemental Note). Any rates that exceed this

theoretical limit of diffusion would reflect hopping events. While there was an increase in the

diffusion with increasing ionic strength and a widening of the distribution of diffusivities, none of

the values were above the theoretical limit of diffusion to unambiguously signify hopping. Since

the crystal structures (PDBs 2RBA and 4Z7B) show TDG interacts with both strands of the helix

when bound to a lesion, we labeled and mixed TDG-HaloTag with two different colored labels,

JF635 (red) and JF552 (green) to directly visualize hopping. Two types of behavior were

observed: (1) JF635-TDG and JF552-TDG collide, in which the two orthogonally labeled TDG

proteins did not bypass one another (62.5%), and (2) JF635-TDG and JF552-TDG molecules

bypass one another (37.5%) (Fig. 2b). The only way it is possible to observe both of these

events is for TDG to have the ability to not only slide on the DNA but also to hop.

TDG catalytic variants reveal R275 is essential for base detection.

N140 is essential for activating the water molecule for catalysis to occur and R275 fills in

the hole in the duplex that is generated when the base is flipped into the active site. The

difference in expression of each variant compared to endogenous TDG is 72-fold for N140A, 45-

fold for R275A, and 21-fold for R275L (Fig. S2). Mutating to R275A and R275L decreases the

excision of T from G:T pairs by 8-fold and 30-fold, respectively, and also reduces the binding

affinity of either variant to G:T containing substrates 3-fold[13]. It is unknown if either variant

affects the ability of TDG to search for DNA lesions[3,13,25]. To investigate the role of these key

amino acids on the lesion search and recognition mechanism of TDG, we performed additional

single-molecule imaging with TDG-HaloTag-N140A, R275L, and R275A variants and DNA

containing regions of 5fC. Consistent with the WT TDG, we observed motile and non-motile

events for all three variants tested, which represent nonspecific and 5fC binding, respectively.

The N140A nonspecific events (motile) bound with a lifetime of 15.9 ± 0.23 seconds (84.8%)

and the specific events (non-motile) bound with a lifetime of 52.9 ± 1.1 seconds (15.2%) (Fig.

3a, Supplemental Figs. 1d and 6a-b). This shorter lifetime for specific events (compared to WT)

and the absence of short-lived rebinding events, further suggesting that wild type TDG is able to

cleave the 5fC moiety and rebind to the abasic site product. The R275A nonspecific events

(motile) bound with a lifetime of 1.07 ± 0.02 seconds (92.5%) and the specific events (non-

motile) bound with a lifetime of 10.0 ± 1.1 seconds (7.5%) (Fig. 3b, Supplemental Figs. 1e and

6c-d). The R275L nonspecific events (motile) bound with a lifetime of 0.7 ± 0.02 seconds

(88.7%) and the specific events (non-motile) bound with a lifetime of 1.7 ± 0.06 seconds (11.3%)

(Fig. 3c, Supplemental Figs. 1e and 6e-f). N140A, R275A, and R275L bound to unmodified DNA

with lifetimes of 7.7 ± 0.24, 2.8 ± 0.13, and 1.8 ± 0.02 seconds, respectively (Fig. 3a-c, Table 1),

with the R275A and R275L lifetimes shorter than the lifetime of TDG-HaloTag-JF635 binding to

unmodified DNA. Together these data reveal that disrupting key active site residues disrupts the

ability of TDG to bind DNA as well as demonstrating a reduced binding lifetime at sites of 5fC, in

agreement with previous bulk biochemical studies[13,26].

Since we observed the binding lifetimes of the TDG variants to differ from wild type TDG,

we examined the diffusion of these variants by mean squared displacement (MSD) analysis.

MSD analysis measures the displacement of the particle over a given time window, over the

course of the event length to determine the diffusivity and α, the anomalous diffusion coefficient.

N140A had a diffusion coefficient of 2.2×10^{-2} μm^2/s, which was not significantly different from

wild type (Fig. 3d). This indicates that N140 does not significantly affect the movement of TDG

on unmodified DNA, but the decreased binding lifetime leads to less DNA being sampled before

dissociation. The diffusion coefficient for R275A (1.3×10^{-1} μm^2/s) and R275L (8.7×10^{-2} μm^2/s)

were significantly faster than both the wild type and N140A variant (Fig. 3a, Table 1), suggesting

that R275 is more important for searching unmodified DNA than N140.

We can estimate the residence time per base pair by utilizing the stepping rate (see

Supplemental Note). The base pair residence time for wild type was estimated to be 2.1 ± 1.6

µs. For N140A, the base pair residence time was estimated to be 2.6 ± 2.2 µs, which is similar

to the time for wild type TDG (2.1 ± 1.6 µs). In contrast, the R275A and R275L TDG variants had

shorter residence times of 0.5 ± 0.3 µs and 0.7 ± 0.5 µs, respectively (Supplemental Table 1).

This may indicate that the R275A/L variants are no longer able to flip out and sample base

pairs, which agrees with previous biochemical studies that indicated R275 plays a role in

promoting or stabilizing the flipped nucleotide[13]. While the faster diffusion observed for the

R275A and R275L TDG variants increases the amount of DNA sampled during each binding

event, the rapid search may be less effective if TDG is more likely to miss the lesion or modified

base, similar to behavior observed with other DNA glycosylases wedge residue variants[27].

The N-terminus of TDG facilitates its movement on DNA.

Previous work identified the first 81 amino acids are not essential for the catalytic activity

of TDG[28]. However, we hypothesized that the intrinsically disordered N-terminus may play a key

role for TDG search of unmodified DNA (Supplemental Fig. 1a, c). To investigate the role of the

TDG N-terminal region on lesion search and lesion recognition mechanism, we performed

additional single-molecule imaging experiments with TDG-HaloTag containing a truncation of

the N-terminal 81 residues (ΔN-term-TDG-HaloTag) and DNA containing regions of 5fC

modification. The ΔN-term-TDG-HaloTag was expressed 50-fold higher compared to

endogenous TDG (Fig S2). We observed similar behavior with ΔN-term-TDG-HaloTag as full-

length TDG-HaloTag, where long-lived, stationary events were followed by shorter lived

stationary events (Fig. 4a). Interestingly, the nonspecific lifetime of ΔN-term-TDG-HaloTag was

1.2 ± 0.02 seconds which is three-fold shorter than full-length TDG-HaloTag (Fig. 4b). This

reduced lifetime on nonspecific DNA supports the hypothesis that the N-terminus of TDG is

important for searching the DNA.

A previous AFM study determined that TDG bends the DNA ~30° when searching for a

lesion, whereas the DNA is bent ~70° when TDG is bound to the lesion[29]. We hypothesized

based on the published AFM study[29] that the necessity of DNA bending by TDG may result in

shorter lifetimes at higher DNA tensions because TDG cannot fully engage with the extended

DNA, but our data showed this is not the case. In our prior experiments we used a tension of 10

pN. By increasing the tension from 5 pN to 40 pN with full-length TDG, we observed similar

lifetimes of 6.49 ± 0.02, 6.45 ± 0.06, and 6.25 ± 0.20 seconds at 5, 10 and 20 pN, respectively

(Fig. 4c, Table 2). At 30 pN and 40 pN, we observed lifetimes that were significantly ~two-fold

higher than at lower tensions (Fig. 4c, Table 2). For ΔN-term TDG, we also observed a two-fold

increase in the binding lifetimes with increasing tension. Additionally, ΔN-term TDG stays bound

to the unmodified DNA with a 3-5-fold shorter lifetime than full-length at the respective tensions

(Table 2). Shorter lifetimes for ΔN-term compared to full-length further supports that idea that

the N-terminus is essential for prolonged interaction with nonspecific DNA. MSD analysis

indicated a decrease in diffusivity with increasing DNA tension for both full-length and ΔN-term

TDG, showing an ~ 5-fold decrease from 5 to 40 pN (Fig. 4e-f, Table 2). Longer lifetimes and a

decrease in diffusivity at higher tensions could indicate that TDG is better at interrogating the

DNA for the base modification. Furthermore, these data indicate that bending of the DNA is not

essential for TDG to bind productively to DNA. This idea is further supported by an increase in

the residence time per base pair (Supplemental Note, Supplemental Table 2). The anomalous

diffusion exponent, α, provides further insight into the movement of a particle, with α = 1 is

random diffusion, α < 1 sub-diffusive (constrained motion), and α > 1 is super diffusive. For full-

length WT TDG a decrease in the ~~anomalous diffusion exponent, α,~~ was observed but we did

not see a significant decrease in α for ΔN-term TDG (Fig. 4e-f, Table 2). The TDG variants,

N140A, R275L, and R275A, were also investigated at 40 pN to further investigate base

interrogation investigate base interrogation at the idea of base interrogation. TDG N140A had a

similar behavior to WT TDG: a decrease in both alpha and diffusivity were observed with an

increase in tension (Fig. 4g). At high tension, the diffusivity of TDG R275L significantly

decreased compared to low tension, but no significant difference was observed for TDG R275A.

(Fig. 4h-i), supporting the idea that the size of the side chain aids in flipping in the base. These

data indicate that DNA tension, perhaps through increasing the distance between stacked bases

(~5% increase in distance at 40 pN compared to 10 pN), provides better interaction of the N-

terminal domain to increase the affinity to DNA increasing the dwell time and slowing the rate of

diffusion to a mode of constrained motion.

N-terminus of TDG mediates nucleosome interactions.

We sought to understand how chromatin structure impacts the TDG search mechanism

by reconstituting an undamaged nucleosome using a 601 sequence[30]and ligating the

nucleosome into 6.2 kb biotinylated handles using the DNA tethering kit from LUMICKS

(Supplemental Fig. 7). We then performed single-imaging experiments using TDG-HaloTag and

the Cy3 labeled mononucleosome-containing DNA substrate. We observed three different

behaviors when TDG encountered the nucleosome: (1) TDG approaches from one side and

hops over the nucleosome (bypass) (Fig. 5a), (2) TDG collides with but does not move past the

nucleosome (no bypass) (Fig. 5b), and (3) TDG colocalizes with the nucleosome after the initial

encounter (Fig. 5c). Further analysis revealed that TDG bypasses the nucleosome during 33.3%

of encounters, did not bypass the nucleosome during 39.2% of encounters, and colocalized with

the nucleosome during 27.5% of the encounters. CRTD analysis fit to a one-phase exponential

indicated that TDG bound to nucleosomes with a lifetime of 16.5 ± 1.9 seconds (Fig. 5e), which

is 2-fold longer than non-damaged DNA, and 4.5-fold shorter than binding to 5fC.

Since we observed the N-terminus is important for movement on DNA, we also sought to

see if the N-terminus plays a role in binding or bypassing nucleosomes. To address this, we

performed additional single-imaging experiments using TDG-HaloTag-ΔN-term and the

mononucleosome-containing DNA substrate. These experiments revealed a significant

decrease in the frequency of ΔN-term colocalized with the nucleosome (8.3%), an increase in

the number of events that approached but did not bypass the nucleosome (61.1%), and saw a

similar frequency for bypass (30.6%) (Fig. 5e). This decrease in TDG colocalization with the

nucleosome indicates that the N-terminus plays an important role in nucleosome binding.

Discussion

In this study, we utilized single-molecule methods to determine how TDG searches for

DNA damage as well as uncovering how key active site residues, domains, and the presence of

nucleosomes alter this search process. Similar to other DNA glycosylases, TDG forms

moderately stable interactions with unmodified DNA to optimize its search[31,32]. Despite the

ongoing 1D search, in our single molecule regime TDG primarily binds its modified sites directly

out of solution in a 3D diffusion mechanism, with a limited number of events that bind

unmodified DNA first and then slide into the lesion. Of note, sliding interactions of less than 100

base pairs are not discernable on our system so these events that appear as 3D may have

some component of 1D diffusion not observed, as was observed for human uracil DNA

glycosylase hUNG2[33,34]. Furthermore, we found that TDG bound specific lesions sites 10-fold

longer than unmodified DNA. Since TDG is able to bind a wide array of modified bases,

including 5caC and G:T mismatch, it is plausible that a similar behavior would be observed:

TDG will bind to the modified base and become nonmotile, with rebinding to the generate abasic

site. While it is currently not known how long TDG will bind to different lesions, it is likely that

TDG will bind to a specific base modification for a time that is directly correlated to the

biochemically determined cleavage rate followed by subsequent rebinding to an abasic site.

Examining TDG under different ionic conditions, as well as two color experiment, demonstrated

that TDG is able to both slide and hop along the DNA, which appears to be common amongst

bacterial and mammalian glycosylases[33,35,36]. We estimated TDG's search length on λ DNA

utilizing the observed diffusion constant in $\frac{bp^2}{sec}$ and the nonspecific lifetime of 7.9 sec (See

Supplemental Note) to be 5225 bp or ~10% of λ DNA per encounter.

The complex lesion search mechanism used by TDG has several potential benefits

during interrogation of the genome for DNA lesions. Due to TDG's high affinity for nondamaged

DNA, the ability to switch between sliding and hopping may also be advantageous in situations

where TDG encounters other factors on the DNA, including other transcription factors or

nucleosomes (Fig. 6). With a sliding-exclusive method, full dissociation would need to occur to

sample the DNA after the block. However, hopping would allow TDG to circumvent the block

and continue scanning on the other side of an obstacle. This use of dual linear diffusion modes

leads to the question of whether certain cellular conditions favor TDG hopping versus sliding on

DNA. For example, if TDG has a sequence preference, it may adopt a conformation at these

preferred sequences where tight binding leads to direct sliding as TDG further investigates that

region of DNA. While it is currently unclear whether TDG has sequence preference, it is

possible that sequences enriched in CG dinucleotides could result in initiating sliding behaviors.

This type of increased binding to particular sequences would be highly advantages in regions

with a heavier burden of DNA modification either due to (1) an enrichment of methylated CG

dinucleotides or (2) TET enzymes localizing to methylated CpG rich islands to oxidize 5-methlyC

groups for their removal by TDG and subsequent transcriptional activation[37]. It has been

previously determined the glycosylase OGG1 slides and hops along the DNA in search of a

lesion[31,35], whereas hUNG2, a uracil DNA glycosylase family member, performs 3D diffusion,

short range sliding, and hopping to find a lesion[33]. Thus, TDG exhibiting multiple modes of

diffusion is in agreement with the behavior observed for other mammalian glycosylases and

appears to be essential to efficiently scan the DNA and the ability to switch between modes may

be dependent upon sequence or other cellular factors (Fig. 6a).

Our single molecule work also revealed the active site variants N140A, R275A, and

R275L change the behavior of TDG on DNA. N140A had a shorter dwell time on 5fC but did not

exhibit a faster diffusion than wild type. This contrasts other previous studies with the

glycosylase OGG1, where the catalytically dead glycosylase bound much longer than that of the

WT[19,38]. On the other hand, R275A/L variants both exhibited a shorter dwell time as well as a

faster diffusion. One explanation for this behavior is that R275 probes the DNA sequence during

the sliding, enabling efficient detection of DNA damage but also slowing the speed of sliding

interactions. Further, it is possible that the lack of the positive charge causes TDG to have lower

binding affinity and thus shorter dwell times and faster rates of diffusion. A previous study on

bacterial glycosylases Fpg, Nei, and Nth has also shown that if a residue that interrogates the

DNA is changed to alanine, faster diffusivity was observed, indicating that efficient recognition of

a base is dependent on a single active site residue[27]. Together, these findings provide an in-

depth description of how key catalytic residues of TDG contribute to its search for DNA damage

and provides insight into how its mixed mode of hopping and sliding searches contribute to

efficient damage detection and repair.

Deleting the disordered N-terminus of TDG, consisting of the first 81 amino acids, did not

reduce the ability to bind to 5fC. However, the ability of ΔN-term TDG to search nonspecific DNA

was significantly reduced, due to the three-fold shorter lifetime of ΔN-term TDG bound to DNA

nonspecifically. While the tension on DNA at any given point in the cell is not well known, it is

known that nucleosomes begin to unwrap at 5 pN[39], hairpins begin to pull apart 10-15 pN[40,41],

G-quadruplexes begin to unfold 15-20 pN[42], RNA polymerase has been shown to exert a force

of 25-30 pN on DNA[43] and double strand DNA begins to unwind at ~50 pN[44,45]. With increasing

tension, the λ DNA may begin to have regions that may potentially allow for TDG to interrogate

the DNA more easily perhaps due to widening of the base steps and less stacking interactions,

allowing better interrogation of the DNA at higher tensions. The observed decrease in α,

indicating more constrained motion on DNA, for full-length but not ΔN-term TDG suggests that

the disordered N-terminus may play a role in allowing TDG to thoroughly interrogate the DNA for

a lesion. The N-terminus may also play a role in TDG switching between hopping and sliding. If

TDG needs to form for tighter interactions with the DNA in order to slide, the N-terminus, which

contains several positively charged amino acid residues, may adopt a confirmation that limits

hoping to better probe an area. Similarly, a study has shown that the N-terminus of hUNG2 has

an increase in contact with DNA during molecular crowding which results in an efficient sliding

and prevents hUNG2 from dissociating and diffusing into bulk solution[46].

We have shown that not only is TDG able to hop over a nucleosome, but it also interacts

with a nucleosome with an apparent dwell time of 16.5 ± 1.9 seconds. Deleting the N-terminus

significantly reduces the binding frequency to nucleosomes (Fig. 6b). This interaction of TDG

with nucleosomes may act as a scaffold to help facilitate removal of oxidized cytosine in

heterochromatic regions. Alternatively, perhaps TDG associates with nucleosomes prior to the

arrival of TET enzymes, and thus would increase the efficiency of oxidative demethylation by

already being in the correct location prior to the initial 5mC oxidation step in heterochromatic

regions. Efficient active oxidative demethylation is vital for life and of interest for potential

therapeutic applications[47], and this work gives key mechanistic insight on this process. TDG

interacting with a nucleosome could also alter nucleosome dynamics and may interfere with fork

progression and DNA replication, and thus might explain its degradation during S-phase in

mammalian cells.

Taken together, our studies offer new insight into how key active site residues and

domains of TDG contribute to its efficient search for DNA damage, and how chromatin can act

either as a roadblock or a scaffold during the search process. Of note, these studies with

nucleosome core particles represent a unique look at how base excision repair enzyme interact

with nucleosomes that do not have free DNA ends present. Future studies will need to be

completed to determine the impact nucleosome unwrapping, chromatin remodelers, and how

post translational modifications of TDG will impact its ability to efficiently search for its

substrates.

Online Methods

DNA substrate preparation

Nick translation for confocal imaging. Lambda DNA was biotinylated as described previously[19].

Briefly, DNA was purchased from New England Biotechnologies and treated with Klenow

fragment polymerase (NEB) and a dNTP mix containing biotinylated dCTP. This results in one

side of the DNA containing four biotins and the other containing six biotins.

Next, 1 μg of the biotinylated lambda DNA is treated with Nt.BspQI (NEB) for 1 hour at 50°C

then heat inactivated for 20 minutes at 80°C, which will generate single strand breaks at 10

different sites along the DNA by cutting 3' of the recognition site 5'-GCTCTTCN-3'. To

incorporate 5fC at the 10 sites, 800 ng nicked DNA was incubated with a dNTP mix containing

d5fCTP, fluorescein labeled dUTP, dATP, and dGTP were incubated and 10 units of DNA

polymerase 1 (NEB) for 6 minutes at 37°C. Reaction was inactivated by heating at 75°C for 20

minutes and allowed to slowly cool to room temperature. To seal the nicks, a reaction containing

800 ng of the nick translated DNA, 8 mM ATP and 10 units of T4 ligase (NEB) were incubated

overnight at 16°C. Reaction was heat inactivated for 10 minutes at 65°C and allowed to cool to

room temperature slowly. Of the 10 d5fCTP sites, only eight d5fCTP sites can be resolved as

two of the sites are within 400 bp of each other and one site is near the end of λ DNA and thus

too close to the bead (Supplemental Fig. 4a-c).

Arms ligation for confocal imaging. Using the LUMICKS DNA tethering kit and protocol, a

sequence of interest is ligated into two handles that are each 6.3 kb in length to generate a

substrate that is 12.6 kb in total. For the defined 5fC, the following oligonucleotide sequences

were ordered from Trilink and IDT, respectively:

Top strand (5fC): 5'-phosphate-*CAAC* ACC AGT CCA TCG CTC A**5fC**G TAC AGA GCT G–3'

Bottom strand: 5'- phosphate – *ACCA* CAG CTC TGT ACG TGA GCG ATG GAC TGG T-3'

Annealing reactions were heated at 95°C for 5 minutes in buffer containing 10 mM Tris-HCl (pH

8.0) and 100 mM KCl, and the heat block was turned off and reactions then cooled slowly to

room temperature. In a 10 μL reaction, 25 nM of the duplex, 2.5 μL of each handle, 1 μL of 10X

ligase buffer, and 0.5 μL of ligase were mixed and incubated at 16°C overnight. Reaction was

heat inactivated at 65°C for 10 minutes and allowed to slowly cool to room temperature.

Substrate was stored at 4°C and protected from light.

Undamaged nucleosome were reconstituted on a 601 sequence as described

previously[30]. Briefly, Human histone H3 C96S C110A, H2A K119C, H2B, and H4 were ordered

from the Histone Source at Colorado State University. Incubate H2A K119C and H2B in 2

mg/mL guanidinium buffer at room temperature for 2 hours. Mix equimolar amounts of H2A and

H2B and dialyze against high salt refolding buffer a total of 3 times, and at least 8 hours for

each exchange at 4°C. Repeat the same process for H3 C96S C110A and H4 to refold the

tetramer. Purify the H2A K119C/H2B dimer and the H3 C96S C110A/H4 tetramer over Superdex

200 column. Incubate the H2A K119C/H2B dimer with 0.7mM TCEP for 20 minutes at 4°C. Cy3-

maleimide dye was added to the H2A K119C in a 2:1 molar ratio and incubated at room

temperature for 1 hour while rocking. Reaction was quenched with 10 mM DTT, the dimer was

purified over a Superdex S200 column, and frozen in 50% glycerol. After confirming the

stoichiometry with SDS gel, add equal volume of 100% glycerol to store the H2A K119C/H2B

dimer and the H3 C96S C110A/H4 tetramer at -20°C.

To reconstitute the nucleosome on DNA, the following ultramer sequences were ordered from

IDT:

Top Strand: 5'-phosphate-*CAAC* TGA GAC CAT GTA CCC AGT TCG AAT CGG ATG TAT ATA

TCT GAC ACG TGC CTG GAG ACT AGG GAG TAA TCC CCT TGG CGG TTA AAA CGC GGG

GGA CAG CGC GTA CGT GCG TTT AAG CGG TGC TAG AGC TGT CTA CGA CCA ATT GAG

CGG CCT CGG CAC CGG GAT TCT CGA TAA CTC AGC AAT AGT GGG TCT CA – 3'

Bottom strand: 5'- phosphate -*ACCA* TGA GAC CCA CTA TTG CTG AGT TAT CGA GAA TCC

CGG TGC CGA GGC CGC TCA ATT GGT CGT AGA CAG CTC TAG CAC CGC TTA AAC GCA

CGT ACG CGC TGT CCC CCG CGT TTT AAC CGC CAA GGG GAT TAC TCC CTA GTC TCC

AGG CAC GTG TCA GAT ATA TAC ATC CGA TTC GAA CTG GGT ACA TGG TCT CA – 3'

The annealed DNA, H2A K119C/H2B dimer, and the H3 C96S C110A/H4 tetramer are mixed in

a 1:2:1 molar ratio and equilibrate in dialysis tubing against high salt buffer for 30 minutes. To

remove the high salt buffer, a series of dialysis steps occur to transition from 1.5 M NaCl to

0.125 M NaCl. The reconstituted nucleosome is concentrated, and heat shocked at 55°C for 30

minutes, prior to spinning over a 10-40% sucrose gradient for 40 hours at 125,000 xG at 4°C.

Fractions containing reconstituted nucleosomes are combined, buffered exchanged into TE

buffer, and concentrated to ~1 µM and stored at 4°C. To ligate into the LUMICKS handles, 25

nM of the reconstitute nucleosome, 2.5 µL of each handle, 1 µL of 10X ligase buffer, and 0.5 µL

of ligase were mixed and incubated at 16°C overnight. Substrate was stored at 4°C and

protected from light.

Plasmid Constructs

Wild type TDG-HaloTag and TDG-HaloTag variants were expressed from plasmids constructed

utilizing Gene Universal Inc with the pHTN HaloTag CMV-neo vector (Promega). (Supplemental

Figs. 1a-e)

Nuclear Extract

Nuclear extracts were prepared as described previously[19]. Briefly, U2OS cells were cultured in

5% CO2 in Dulbecco's modified Eagle's medium (DMEM) supplemented with 4.5 g/L glucose,

10% fetal bovine serum (Gibco), and 5% penicillin/streptavidin (Life Technologies). To obtain

transient overexpression of TDG-HaloTag, cells were transfected with 4 µg of plasmid per 4

million cells using the lipofectamine 2000 reagent and protocol (Thermo Fisher Cat# 11668019).

Nuclear extraction was performed 24 hours after transfection utilizing nuclear extraction kit from

Abcam (ab113474). Following the protocol from the kit, 10 µL single-use aliquots were made,

flash-frozen and, stored at -80°C. Extracts were diluted in buffer containing 20 mM HEPES pH

7.5, 100 mM NaCl, 0.2 mM EDTA, 0.1 mg/ml BSA, 1 mM DTT, 1 mM Trolox, and 5% glycerol.

Nuclear Extract Quantification

SDS gel. A total of 2.5 µg of each nuclear extract was loaded in duplicate on a 4-12% Bis Tris

gels (Invitrogen NP0323BOX) with a dye free sample buffer (4X buffer: 40% glycerol, 200 mM

Tris-Cl (pH 6.8), and 8% SDS). The gel was imaged using a laser scanner for Cy5 (Typhoon,

Amersham). Gels were then stained with Coomassie blue and imaged. The total intensity for

each lane was measured and the lanes were averaged together in order to normalize for

loading. Finally, each of the variant extracts were normalized to the wild type TDG in order to

look at expression levels and the level of free dye. (Supplemental Figure 2a-c)

Western Blotting. Different amounts of purified proteins and nuclear extracts were denatured at

95°C for 10 minutes. Equal volumes were loaded on a 4-20% tris-glycine polyacrylamide gel

(Invitrogen XP04202BOX). Proteins were transferred onto a polyvinylidene difluoride membrane

and blocked in 20% nonfat dry milk in PBST (1x phosphate-buffered saline containing 0.1%

Tween 20) for 1 hr at room temperature. Membranes were probed with TDG primary antibody

(1:500 Sigma HPA052263) overnight at 4°C. Membranes were washed 3 times for 14 min in

PSBT and incubated with peroxidase conjugated rabbit secondary antibody for 1 h at room

temperature. Membranes were washed again 3 times for 15 minutes before developing using

SuperSignal West Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific #34095).

(Supplemental Figure 2d),

Confocal Imaging in the LUMICKS C-trap

Imaging. Fluorophores and dyes utilized were excited with the laser closest to their maximum

intensity. Fluorescein was excited with a 488 nm laser and emission collected with a 500–550

nm band pass filter, Cy3 and HaloTag-JF552 were excited with a 561 nm laser and the emission

was collected with a 575–625 nm band pass filter, and HaloTag-JF635 was excited with a 638

nm laser and emission collected with a 650– 750 nm band pass filter. All data were collected

with a 1.2 NA 60x water emersion objective and photons measured with single-photon

avalanche photodiode detectors. The laser power was set to 5% and kymographs were

collected continuously with 0.1 msec exposure for each pixel of size 100 nm, which resulted in

approximately 30 frames/sec for λ DNA. All data was collected with a minimum of two different

preparations of nuclear extracts across multiple days.

Single-molecule experiments were then performed with a LUMICKS C-trap using a 5-

channel flow chamber. For experiments using λ or nick translated DNA, ~4 μm streptavidin

coated beads and experiments using the LUMICKS handles, ~1.7 μm streptavidin coated beads

were used. The beads were captured using the dual optical tweezers and then transferred to a

second chamber containing the biotinylated DNA and the biotinylated DNA molecule tethered

between the streptavidin coated beads. Once the DNA molecule is tethered, it is transferred to a

final chamber containing a 1:10 dilution of the nuclear extract with TDG-HaloTag-JF635. To

determine the concentration of the protein in the flow cell, background photon intensities were

determined to generate the standard curve of purified HaloTag labeled with JF635 or JF552.

The background intensities of the protein from the nuclear extract were measured and the curve

was utilized to determine protein concertation (Supplemental Figure 3a-b).

Analysis. Kymographs were analyzed with custom software from LUMICKS, Pylake. The

kymographs were exported as .h5 files from the C-trap and viewed using the utility "C-Trap .h5

Visualization GUI (2020)" by John Watters (harbor.lumicks.com). Line tracking was performed

using a custom script from LUMICKS based on a Gaussian fit over the line to determine its

positions[48]. Only events that associated and dissociated during the course of a 5-to-10-minute

kymograph were tracked. Events are plotted on cumulative residence time distribution (CRTD)

plot in order to determine the binding lifetime. Specific binding event curves are fit with a two-

phase exponential decay, for catalytically active WT protein. One-phase exponential decays

were sued for the variants, which have little or no catalytic activity, and nonspecific binding

events for the variants and WT. The two-phase fit for WT represents a second rebinding event to

the catalytic product abasic site with a shorter dwell time. The goodness of the fit for the CRTD

curves range from R^2 = 0.96 – 0.99. Mean squared displacement (MSD) analysis was

performed using a custom script from LUMICKS based on the following equation:

497
$$MSD\,(n\Delta t) = \frac{1}{N-n}\sum_{i=1}^{N-n}(x_{i+n} - x_i)^2 \qquad\qquad (1)$$

498   Where N is the total number of frames in the phase, n is the number of frames at a given time

499   step, Δt is the time increment of one frame, and x_i is the particle position in the ith frame[27]. The

500   diffusion coefficient (D) was determined by fitting a linear model of one-dimensional diffusion to

501   the MSD plots:

502
$$MSD\,(n\Delta t) = 2D(n\Delta t)^{\alpha} + y \qquad\qquad (2)$$

where α is the anomalous diffusion coefficient and y is a constant (y-intercept). Fittings resulting

in R^2 < 0.8 or using <10% of the MSD plot were not considered.

Data Availability

All data are available from the corresponding author upon reasonable request.

Code Availability

Python scripts used to calculate lifetime and MSD analysis have been deposited at

https://harbor.lumicks.com/scripts.

Contributions

Acknowledgements

We greatly appreciate the discussions with the lab members during the course of these

experiments. This work was supported by NIH R35ES031638 (BVH), T32GM088119 (BLS), the

Hillman Postdoctoral Fellowship for Innovative Cancer Research and F32ES034982 (MAS),

2P30CA047904 to the UPMC Hillman Cancer Center, the major equipment grant

S10OD032158-01A1 (BVH), R35-GM136225 (ACD), R35GM128652 (BDF), and F32GM140718

(TMW).

Competing interests

None declared.

Figure Legends

Figure 1: TDG binds DNA specifically and nonspecifically.

(A) Cartoon schematic showing how 5fC was incorporated into nick translated λ DNA. (B)

Cartoon schematic showing 28 base pair duplex DNA containing a single 5fC (orange) ligated

into 6 kb LUMICKS handle kit, with handle 2 containing ATTO 488 (blue). (C) A diagram

depicting the order of reagents, which are under laminar flow, are captured in the flowcell. (D) A

cartoon depiction of the DNA substrate used for TDG binding, with 5fC sites shown in blue, and

an example kymograph with TDG binding shown in red. Specific event indicated with grey

asterisk and nonspecific event indicated with teal asterisk. (E) Cumulative Resident Time

Distribution (CRTD) analysis fit to a two-phase decay of TDG binding DNA containing 5fC

specifically (n = 70) and nonspecifically (n = 487). (F) A cartoon depiction of the unmodified DNA

substrate with an example kymograph of TDG binding and moving. (G) CRTD analysis fit to a

one-phase decay of TDG binding unmodified λ DNA (n = 155). (H) An example kymograph of

TDG binding to a single 5fC. (I) CRTD analysis fit to a two-phase decay of TDG binding to 5fC

(n = 28). Created with BioRender.com.

Figure 2: TDG exhibits linear diffusion on DNA.

(A) Scatter plot of the diffusion coefficient ($\log_{10}D$) calculated for TDG with increasing ionic

strength on unmodified λ. Dashed line, D_{lim}, theoretical limit to free diffusion for TDG-HaloTag.

****$p < 0.0001$ by two-way ANOVA. (B) An example kymograph of the two-color TDG

experiment. Separate TDG-HaloTag extracts were labeled with either JF635 (red) or JF552

(green) and mixed at a 1:1 ratio. White arrows indicate bypass events and yellow arrows

indicate collision events.

Figure 3: TDG catalytic variants indicate R275 is essential for base detection.

Experiments conducted with unmodified or nick translated DNA containing 5fC. (A) CRTD

analysis fit to a one-phase decay of N140A TDG binding to unmodified (n = 163), specifically to

5fC (n = 30) and nonspecifically (n = 168). (B) CRTD analysis fit to a one-phase decay of R275A

TDG binding to unmodified (n = 110), specifically to 5fC (n = 24) and nonspecifically (n = 296).

(C) CRTD analysis fit to a one-phase decay of R275L TDG binding to unmodified (n = 269),

specifically to 5fC (n = 42) and nonspecifically (n = 330). (D) Scatter plot of the diffusion

coefficient ($\log_{10}D$) calculated for each variant TDG on unmodified λ. Dashed line, D_{lim},

theoretical limit to free diffusion for TDG-HaloTag. ****$p < 0.0001$ by two-way ANOVA.

Figure 4: The N-terminus of TDG is significant for its movement on DNA. Tension alters

diffusivity and α for ΔN-term and full-length TDG.

(a) A cartoon depiction of the DNA substrate with 5fC sites shown in blue, and a representative

kymograph with TDG binding shown in red. Break in kymograph to show location of fiducial

markers. (b) CRTD analysis fit to a two-phase decay of TDG binding DNA containing 5fC

specifically (n = 32). CRTD analysis fit to a one-phase decay of TDG binding nonspecifically (n =

243). (c) CRTD analysis fit to a one-phase decay of full-length TDG binding to unmodified DNA

at 5 pN (n = 172), 10 pN (n = 27), 20 pN (n = 53), 30 pN (n = 169), and 40 pN (n = 102). (d)

CRTD analysis fit to a one-phase decay of ΔN-term TDG binding to unmodified DNA at 5 pN (n

= 113), 10 pN (n = 201), 20 pN (n = 197), 30 pN (n = 293), and 40 pN (n = 203) (e) Plot of

diffusion coefficients (D) versus alpha (α) for full-length for 5 pN (n = 50), 10 pN (n = 41), 20 pN

(n = 41), 30 pN (n = 57), and 40 pN (n = 33). (f) Plot of diffusion coefficients (D) versus alpha (α)

for ΔN-term for 5 pN (n = 30), 10 pN (n = 36), 20 pN (n = 24), 30 pN (n = 22), and 40 pN (n =

35). (g) Plot of diffusion coefficients (D) versus alpha (α) for N140A for 10 pN (n = 33) and 40 pN

(n = 34). (h) Plot of diffusion coefficients (D) versus alpha for 10 pN (n = 19) and 40 pN (n = 27).

(α) for R275L. (i) Plot of diffusion coefficients (D) versus alpha (α) for R275A for 10 pN (n = 14)

and 40 pN (n = 19).

Figure 5: N-terminus of TDG is important for interacting with nucleosomes.

A cartoon depiction of the DNA substrate with Cy3 labeled nucleosome. Representative

kymograph with full-length TDG (shown in red) (a) bypassing, (b) does not bypass, or (c)

colocalizes with the nucleosome shown in green. (d) Representative kymograph of ΔN-term

TDG (red) colocalizing with nucleosome (green). (e) CRTD analysis fit to a one-phase decay of

TDG interacting with the NCP (n = 28). (f) Stacked bar graph showing the fraction of bypass

(green), no bypass (grey), and colocalized (teal) events for full-length and ΔN-term. shows there

is p = 0.0011 by X^2.

Figure 6: Working model.

TDG uses multiple modes of linear diffusion to efficiently search for 5fC. (a) WT TDG (gray)

fused to HaloTag (red) scans dsDNA with sliding and hopping interactions, with a lifetime of 7.9

± 0.06 sec and diffusivity of 0.028 ± 0.02 µm2/sec. Upon 5fC engagement (orange), TDG binds

with a lifetime of 72.9 ± 5.4 sec. These interactions are dependent on active site residues N140,

R275, and the N-terminal domain. (b) When TDG encounters a nucleosome core particle, it can

bypass, not bypass, or colocalize with the NCP (16.5 ± 1.9 sec). Importantly the N-terminal

domain of TDG stabilizes the interaction with the NCP, resulting in a four-fold increase in

colocalization frequency. Created with BioRender.com.

Table 1. Binding lifetimes and diffusivity

Protein	DNA	Lifetime(s) and Percentages[b]	Average binding lifetime (τ_{avg})[b]	Diffusivity on undamaged DNA ($\mu m^2/s$)[c]
Wild Type	Unmodified	7.9 ± 0.06	72.9 ± 5.4	0.028 ± 0.02
	5fC, specific[a]	101.1 ± 7.1 (67.3 ± 3%)		
		14.7 ± 1.7 (32.7 ± 3%)		
	5fC, nonspecific	7.5 ± 0.3		
	Undamaged NCP	16.5 ± 1.9		
N140A	Unmodified	7.7 ± 0.24		0.022 ± 0.02
	5fC, specific	52.6 ± 1.1		
	5fC, nonspecific	15.9 ± 0.23		
R275A	Unmodified	2.8 ± 0.13		0.130 ± 0.09
	5fC, specific	10.0 ± 1.1		
	5fC, nonspecific	1.07 ± 0.02		
R275L	Unmodified	1.8 ± 0.02		0.087 ± 0.07
	5fC, specific	1.7 ± 0.06		
	5fC, nonspecific	0.7 ± 0.02		
ΔN term	5fC, specific	24.55 ± 3.57	10.2 ± 1.6	
		1.11 ± 18		
	5fC, nonspecific	1.2 ± 0.02		

592 [a] Nick translated DNA
593 [b] The tau is the best fit value ± the standard error of the fit to the observed data
594 [c] The diffusivity is the mean ± the standard error of the mean
595

601 **Table 2. Binding lifetimes and diffusivities dependent upon tension**

Protein	Tension (pN)	Lifetime(s)[a]	Diffusivity ($\mu m^2/s$)[b]	α[c]
Wild Type	5	6.49 ± 0.04	0.034 ± 0.019	1.07 ± 0.22
	10	6.45 ± 0.06	0.032 ± 0.027	1.05 ± 0.24
	20	6.25 ± 0.20	0.026 ± 0.016	1.11 ± 0.27
	30	10.31 ± 0.11	0.008 ± 0.006	0.83 ± 0.32
	40	8.20 ± 0.27	0.007 ± 0.006	0.72 ± 0.35
ΔN term	5	0.99 ± 0.02	0.049 ± 0.067	0.99 ± 0.25
	10	1.39 ± 0.02	0.050 ± 0.049	1.04 ± 0.35
	20	1.58 ± 0.02	0.047 ± 0.052	0.98 ± 0.40
	30	2.03 ± 0.02	0.020 ± 0.015	1.19 ± 0.46
	40	2.03 ± 0.02	0.012 ± 0.009	0.99 ± 0.40
N140A	10	5.9 ± 0.32	0.025 ± 0.017	1.05 ± 0.29
	40	5.52 ± 0.12	0.007 ± 0.005	0.85 ± 0.32
R275A	10	0.7 ± 0.03	0.239 ± 0.179	0.90 ± 0.43
	40	0.73 ± 0.04	0.277 ± 0.403	1.01 ± 0.29
R275L	10	1.4 ± 0.37	0.143 ± 0.095	1.07 ± 0.34
	40	0.66 ± 0.02	0.037 ± 0.031	1.05 ± 0.45

602 [a] The tau is the best fit value ± the standard error of the fit to the observed data
603 [b] The diffusivity is the mean ± the standard error of the mean
604 [c] The alpha is the mean ± the standard error of the mean
605

References

1.   Whitaker, A.M., Schaich, M.A., Smith, M.R., Flynn, T.S. & Freudenthal, B.D. Base
excision repair of oxidative DNA damage: from mechanism to disease. *Front Biosci*
(Landmark Ed) **22**, 1493-1522 (2017).

2.   Maiti, A. & Drohat, A.C. Thymine DNA glycosylase can rapidly excise 5-formylcytosine
and 5-carboxylcytosine: potential implications for active demethylation of CpG sites. *J*
Biol Chem **286**, 35334-35338 (2011).

3.   Maiti, A., Michelson, A.Z., Armwood, C.J., Lee, J.K. & Drohat, A.C. Divergent
mechanisms for enzymatic excision of 5-formylcytosine and 5-carboxylcytosine from
DNA. *J Am Chem Soc* **135**, 15813-22 (2013).

4.   Pidugu, L.S., Dai, Q., Malik, S.S., Pozharski, E. & Drohat, A.C. Excision of 5-
Carboxylcytosine by Thymine DNA Glycosylase. *J Am Chem Soc* **141**, 18851-18861
(2019).

5.   Pidugu, L.S. et al. Structural Basis for Excision of 5-Formylcytosine by Thymine DNA
Glycosylase. *Biochemistry* **55**, 6205-6208 (2016).

6.   Cortázar, D. et al. Embryonic lethal phenotype reveals a function of TDG in maintaining
epigenetic stability. *Nature* **470**, 419-23 (2011).

7.   Cortellino, S. et al. Thymine DNA glycosylase is essential for active DNA demethylation
by linked deamination-base excision repair. *Cell* **146**, 67-79 (2011).

8.   Bennett, M.T. et al. Specificity of human thymine DNA glycosylase depends on N-
glycosidic bond stability. *J Am Chem Soc* **128**, 12510-9 (2006).

9.   Morgan, M.T., Bennett, M.T. & Drohat, A.C. Excision of 5-halogenated uracils by human
thymine DNA glycosylase. Robust activity for DNA contexts other than CpG. *J Biol Chem*
282, 27578-86 (2007).

10.   Ito, S. et al. Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-
carboxylcytosine. *Science* **333**, 1300-3 (2011).

11.   Kellinger, M.W. et al. 5-formylcytosine and 5-carboxylcytosine reduce the rate and
substrate specificity of RNA polymerase II transcription. *Nat Struct Mol Biol* **19**, 831-3
(2012).

12.   DeNizio, J.E. et al. TET-TDG Active DNA Demethylation at CpG and Non-CpG Sites. *J*
Mol Biol **433**, 166877 (2021).

13.   Maiti, A., Morgan, M.T. & Drohat, A.C. Role of two strictly conserved residues in
nucleotide flipping and N-glycosylic bond cleavage by human thymine DNA glycosylase.
J Biol Chem **284**, 36680-36688 (2009).

14.   Morgan, M.T., Maiti, A., Fitzgerald, M.E. & Drohat, A.C. Stoichiometry and affinity for
thymine DNA glycosylase binding to specific and nonspecific DNA. *Nucleic Acids Res*
39, 2319-29 (2011).

15.   Maiti, A., Morgan, M.T., Pozharski, E. & Drohat, A.C. Crystal structure of human thymine
DNA glycosylase bound to DNA elucidates sequence-specific mismatch recognition.
Proc Natl Acad Sci U S A **105**, 8890-5 (2008).

16.   Hashimshony, T., Zhang, J., Keshet, I., Bustin, M. & Cedar, H. The role of DNA
methylation in setting up chromatin structure during development. *Nat Genet* **34**, 187-92
(2003).

17.   Tarantino, M.E., Dow, B.J., Drohat, A.C. & Delaney, S. Nucleosomes and the three
glycosylases: High, medium, and low levels of excision by the uracil DNA glycosylase
superfamily. *DNA Repair (Amst)* **72**, 56-63 (2018).

18.   Deckard, C.E. & Sczepanski, J.T. Reversible chromatin condensation by the DNA repair
and demethylation factor thymine DNA glycosylase. *Nucleic Acids Res* **49**, 2450-2459
(2021).

19.   Schaich, M.A. et al. Single-molecule analysis of DNA-binding proteins from nuclear
656   extracts (SMADNE). *Nucleic Acids Res* **51**, e39 (2023).

657   20.   Bigman, L.S., Greenblatt, H.M. & Levy, Y. What Are the Molecular Requirements for
658   Protein Sliding along DNA? *J Phys Chem B* **125**, 3119-3131 (2021).

659   21.   Bigman, L.S. & Levy, Y. Protein Diffusion Along Protein and DNA Lattices: Role of
660   Electrostatics and Disordered Regions. *Annu Rev Biophys* **52**, 463-486 (2023).

661   22.   Bagchi, B., Blainey, P.C. & Xie, X.S. Diffusion constant of a nonspecifically bound protein
662   undergoing curvilinear motion along DNA. *J Phys Chem B* **112**, 6282-4 (2008).

663   23.   Berg, O.G., Winter, R.B. & von Hippel, P.H. Diffusion-driven mechanisms of protein
664   translocation on nucleic acids. 1. Models and theory. *Biochemistry* **20**, 6929-48 (1981).

665   24.   Tafvizi, A., Huang, F., Fersht, A.R., Mirny, L.A. & van Oijen, A.M. A single-molecule
666   characterization of p53 search on DNA. *Proc Natl Acad Sci U S A* **108**, 563-8 (2011).

667   25.   Dow, B.J., Malik, S.S. & Drohat, A.C. Defining the Role of Nucleotide Flipping in Enzyme
668   Specificity Using (19)F NMR. *J Am Chem Soc* **141**, 4952-4962 (2019).

669   26.   Coey, C.T. & Drohat, A.C. Defining the impact of sumoylation on substrate binding and
670   catalysis by thymine DNA glycosylase. *Nucleic Acids Res* **46**, 5159-5170 (2018).

671   27.   Nelson, S.R., Dunn, A.R., Kathe, S.D., Warshaw, D.M. & Wallace, S.S. Two glycosylase
672   families diffusively scan DNA using a wedge residue to probe for and identify oxidatively
673   damaged bases. *Proc Natl Acad Sci U S A* **111**, E2091-9 (2014).

674   28.   Coey, C.T. et al. Structural basis of damage recognition by thymine DNA glycosylase:
675   Key roles for N-terminal residues. *Nucleic Acids Res* **44**, 10248-10258 (2016).

676   29.   Buechner, C.N., Maiti, A., Drohat, A.C. & Tessmer, I. Lesion search and recognition by
677   thymine DNA glycosylase revealed by single molecule imaging. *Nucleic Acids Res* **43**,
678   2716-29 (2015).

679   30.   Ryan, B.J., Weaver, T.M., Spencer, J.J. & Freudenthal, B.D. Generation of Recombinant
680   Nucleosomes Containing Site-Specific DNA Damage. *Methods Mol Biol* **2701**, 55-76
681   (2023).

682   31.   Rowland, M.M., Schonhoft, J.D., McKibbin, P.L., David, S.S. & Stivers, J.T. Microscopic
683   mechanism of DNA damage searching by hOGG1. *Nucleic Acids Res* **42**, 9295-303
684   (2014).

685   32.   Dunn, A.R., Kad, N.M., Nelson, S.R., Warshaw, D.M. & Wallace, S.S. Single Qdot-
686   labeled glycosylase molecules use a wedge amino acid to probe for lesions while
687   scanning along DNA. *Nucleic Acids Res* **39**, 7487-98 (2011).

688   33.   Porecha, R.H. & Stivers, J.T. Uracil DNA glycosylase uses DNA hopping and short-range
689   sliding to trap extrahelical uracils. *Proc Natl Acad Sci U S A* **105**, 10791-6 (2008).

690   34.   Schonhoft, J.D. & Stivers, J.T. Timing facilitated site transfer of an enzyme on DNA. *Nat
691   Chem Biol* **8**, 205-10 (2012).

692   35.   Blainey, P.C., van Oijen, A.M., Banerjee, A., Verdine, G.L. & Xie, X.S. A base-excision
693   DNA-repair protein finds intrahelical lesion bases by fast sliding in contact with DNA.
694   *Proc Natl Acad Sci U S A* **103**, 5752-7 (2006).

695   36.   Friedman, J.I. & Stivers, J.T. Detection of damaged DNA bases by DNA glycosylase
696   enzymes. *Biochemistry* **49**, 4957-67 (2010).

697   37.   Williams, K., Christensen, J. & Helin, K. DNA methylation: TET proteins-guardians of
698   CpG islands? *EMBO Rep* **13**, 28-35 (2011).

699   38.   Schaich, M.A., Weaver, T.M., Roginskaya, V., Freudenthal, B.D. & Van Houten, B.
700   Single-molecule analysis of purified proteins and nuclear extracts: Insights from 8-
701   oxoguanine glycosylase 1. *DNA Repair (Amst)* **134**, 103625 (2024).

702   39.   Díaz-Celis, C. et al. Assignment of structural transitions during mechanical unwrapping
703   of nucleosomes and their disassembly products. *Proc Natl Acad Sci U S A* **119**,
704   e2206513119 (2022).

40.   Bercy, M. & Bockelmann, U. Hairpins under tension: RNA versus DNA. *Nucleic Acids*
Res **43**, 9928-36 (2015).
41.   Liu, Y., Galior, K., Ma, V.P. & Salaita, K. Molecular Tension Probes for Imaging Forces at
the Cell Surface. *Acc Chem Res* **50**, 2915-2924 (2017).
42.   Mitra, J. et al. Extreme mechanical diversity of human telomeric DNA revealed by
fluorescence-force spectroscopy. *Proc Natl Acad Sci U S A* **116**, 8350-8359 (2019).
43.   Wang, H.Y., Elston, T., Mogilner, A. & Oster, G. Force generation in RNA polymerase.
Biophys J **74**, 1186-202 (1998).
44.   Rouzina, I. & Bloomfield, V.A. Force-induced melting of the DNA double helix 1.
Thermodynamic analysis. *Biophys J* **80**, 882-93 (2001).
45.   Williams, M.C., Rouzina, I. & Bloomfield, V.A. Thermodynamics of DNA interactions from
single molecule stretching experiments. *Acc Chem Res* **35**, 159-66 (2002).
46.   Rodriguez, G. et al. Disordered N-Terminal Domain of Human Uracil DNA Glycosylase
(hUNG2) Enhances DNA Translocation. *ACS Chem Biol* **12**, 2260-2263 (2017).
47.   Nuñez, J.K. et al. Genome-wide programmable transcriptional memory by CRISPR-
based epigenome editing. *Cell* **184**, 2503-2519.e17 (2021).
48.   Mangeol, P., Prevo, B. & Peterman, E.J. KymographClear and KymographDirect: two
tools for the automated quantitative analysis of molecular and cellular dynamics using
kymographs. *Mol Biol Cell* **27**, 1948-57 (2016).

Figure 1

Figure 2

Figure 3

Figure 5

a

Mixed 1D diffusion: sliding and hopping

5fC engagement

b

No bypass (39.2%)

Bypass (33.0%)

Colocalize (27.5%)

Figure 6