TITLE: Impairment of the glial phagolysosomal system drives prion-like propagation in a *Drosophila* model of Huntington's disease

ABBREVIATED TITLE: Phagocytic glial defects drive huntingtin proteotoxicity

AUTHORS and AFFILIATIONS: Graham H. Davis[1,2,3], Aprem Zaya[3], and Margaret M. Panning Pearce[1,2,3,*]

[1]Rowan University, Department of Biological and Biomedical Sciences, Glassboro, NJ 08028, [2]Saint Joseph's University, Department of Biology, Philadelphia, PA 19131, and [3]University of the Sciences, Department of Biological Sciences, Philadelphia, PA 19104

*Correspondence should be addressed to Maggie Panning Pearce at pearcem@rowan.edu.

Number of pages: 48

Number of Figures and Tables: 18 Figures and 4 Tables

Abstract Word Count: 238

Introduction Word Count: 612

Discussion Word Count: 1342

CONFLICT OF INTEREST STATEMENT: The authors declare no competing financial interests.

ACKNOWLEDGMENTS: We would like to thank Sabrina Abbruzzese, Ryan Buist, Georgiana Moore, and David Tomlinson for experimental assistance, and all former and current members of the Pearce Lab for their support and many fruitful discussions about this project. This work was supported by funding from University of the Sciences, Rowan University, the W.W. Smith Charitable Trusts, and the National Institutes of Health (awards NS128847 and AG063295).

ABSTRACT

Protein misfolding, aggregation, and spread through the brain are primary drivers of neurodegenerative diseases pathogenesis. Phagocytic glia are responsible for regulating the load of pathogenic protein aggregates in the brain, but emerging evidence suggests that glia may also act as vectors for aggregate spread. Accumulation of protein aggregates could compromise the ability of glia to eliminate toxic materials from the brain by disrupting efficient degradation in the phagolysosomal system. A better understanding of phagocytic glial cell deficiencies in the disease state could help to identify novel therapeutic targets for multiple neurological disorders. Here, we report that mutant huntingtin (mHTT) aggregates impair glial responsiveness to injury and capacity to degrade neuronal debris in male and female adult *Drosophila* expressing the gene that causes Huntington's disease (HD). mHTT aggregate formation in neurons impairs engulfment and clearance of injured axons and causes accumulation of phagolysosomes in glia. Neuronal mHTT expression induces upregulation of key innate immunity and phagocytic genes, some of which were found to regulate mHTT aggregate burden in the brain. Finally, a forward genetic screen revealed Rab10 as a novel component of Draper-dependent phagocytosis that regulates mHTT aggregate transmission from neurons to glia. These data suggest that glial phagocytic defects enable engulfed mHTT aggregates to evade lysosomal degradation and acquire prion-like characteristics. Together, our findings reveal new mechanisms that enhance our understanding of the beneficial and potentially harmful effects of phagocytic glia in HD and potentially other neurodegenerative diseases.

SIGNIFICANCE STATEMENT

Deposition of amyloid aggregates is strongly associated with neurodegenerative disease progression and neuronal cell loss. Many studies point to glial cells as dynamic mediators of disease, capable of phagocytosing toxic materials, but also promoting chronic inflammation and proteopathic aggregate spread. Thus, glia have emerged as promising therapeutic targets for disease intervention. Here, we demonstrate in a *Drosophila* model of Huntington's disease that neuronal mHTT aggregates interfere with glial phagocytic engulfment, phagolysosomal processing, and innate immunity transcriptional responses. We also identify Rab10 as a novel modifier of prion-like transmission of mHTT aggregates. Our findings add to a growing narrative of glia as double-edged players in neurodegenerative diseases.

INTRODUCTION

Neuron-glia crosstalk is critical for maintaining homeostasis in the central nervous system (CNS), and disruption of these intercellular interactions is increasingly recognized as a central component of many neurological disorders, including neurodegenerative diseases. Glia perform immune surveillance functions in the CNS and respond to neuronal injury by altering gene expression (Magaki et al., 2018) and clearing damaged cells (Raiders et al., 2021; Zheng and Tuszynski, 2023). These glial responses may initially be neuroprotective, but prolonged glial reactivity propels the neurodegenerative disease state, for example, by driving premature loss of living neurons or functional synapses (Neniskyte et al., 2011; Hong et al., 2016; Dejanovic et al., 2022) and inducing neuroinflammation (Liddelow et al., 2017). Expanding our understanding of how glia influence neuron function and survival could reveal promising new therapeutic strategies for neurodegenerative diseases.

A pathological hallmark of most neurodegenerative diseases is the accumulation of misfolded proteins into intra- or extracellular amyloid aggregates in vulnerable regions of the CNS (Knowles et al., 2014). Protein aggregates form due to age-associated decline in cellular protein folding capacity (Santra et al., 2019; Stein et al., 2022) and overwhelming of degradative pathways, including the ubiquitin-proteasome system, autophagy, and phagocytosis (Aman et al., 2021; Duong et al., 2021; Wodrich et al., 2022). As professional phagocytes of the brain, microglia and astrocytes clear damaged and dysfunctional cells (Paolicelli et al., 2011; Wakida et al., 2018; Herzog et al., 2019; Lee et al., 2021) and other pathogenic material, such as protein aggregates (Liu et al., 2017; Choi et al., 2020). Defective glial clearance of debris may be a driving force underlying neurodegenerative disease, highlighted by a growing list of genetic risk variants associated with phagocytic and endolysosomal pathways (Podleśny-Drabiniok et al., 2020). Endolysosomal impairment promotes protein aggregate accumulation; in turn, aggregates can drive further endolysosomal dysfunction, including deacidification and membrane permeabilization of intracellular vesicles (Krasemann et al., 2017; Heckmann et al., 2019; Burbidge et al., 2022; Lee et al., 2022). Aggregates that escape degradation may gain the ability

to spread and seed soluble proteins in a prion-like manner (Jucker and Walker, 2018; Monaco and

Fraldi, 2020).

Mechanisms by which pathogenic aggregates dysregulate endolysosomal processing remain

largely unknown. Intracellular membrane fusion events that regulate endo/phagosome maturation are

catalyzed by Rab GTPases (Ng and Tang, 2008; Chan et al., 2011; Langemeyer et al., 2018), enzymes

that cycle between active (GTP-bound) and inactive (GDP-bound) states to organize endomembranes

into distinct functional domains (Hall, 1990; Chan et al., 2011). Rab dysfunction is implicated in several

neurodegenerative diseases—e.g., upregulation of Rab4, Rab5, Rab7, and Rab27 has been observed

in Alzheimer's disease (AD) (Ginsberg et al., 2011), and several Rabs are known substrates of leucine

rich repeat kinase 2 (LRRK2), a genetic risk factor in familial Parkinson's disease (PD) (Jeong et al.,

2018). Intriguingly, spread of α-synuclein between cultured neuronal and enteroendocrine cells is

mediated by the LRRK2 substrate Rab35 (Bae et al., 2018; Rodrigues et al., 2022), suggesting that

Rab-dependent processes may contribute to formation and propagation of pathogenic aggregate

seeds.

Here, we investigated the impact of protein aggregates generated in neurons on phagocytic glial

cell functions in a *Drosophila* model of Huntington's disease (HD). We report that neuronal expression

of aggregation-prone mutant huntingtin (mHTT) protein reduces the ability of glia to clear axonal debris

and to mount phagocytic and innate immunity transcriptional responses following acute nerve injury.

We also observed mHTT-induced changes to numbers of glial lysosomes and Rab+ vesicles in

uninjured brains, and identify Rab10 as a novel modifier of prion-like spreading of mHTT aggregates in

adult fly brains. Together, these studies shed light on mechanisms by which phagocytic glia respond to

and are impaired by accumulation of pathogenic aggregates in neurons.

MATERIALS AND METHODS

Fly husbandry

Fly stocks and crosses were raised on standard cornmeal/molasses media on a 12 hr light/12 hr dark cycle at 25°C, unless otherwise noted. No sex-specific differences were observed in any experiments, so both males and females were utilized in this study. Transgenic or mutant flies were either generated for this study (described below), purchased from Bloomington Drosophila Stock Center, or kindly provided by collaborators. Genotypes and sources of all fly stocks used in this study are listed in Table 1, and complete genotypes of flies used in each figure are listed in Table 2.

Acute antennal nerve injury was performed by bilateral removal of the second and third antennal segments from anesthetized adult flies (MacDonald et al., 2006). For quantitative PCR analyses, maxillary palps were removed in addition to third antennal segments to sever all olfactory receptor neuron (ORN) axons. Axotomized flies were incubated for the indicated times on standard media prior to dissection and processing for imaging.

Cloning and Drosophila transgenesis

pUASTattB(HTT$_{ex1}$Q25-V5) and pUASTattB(HTT$_{ex1}$Q91-V5) plasmids were cloned by PCR amplification of HTT exon 1 (HTT$_{ex1}$) cDNA using a reverse primer that inserted an in-frame, C-terminal V5 epitope tag (GKPIPNPLLGLDST) and ligation into the pQUASTattB vector backbone via XhoI and XbaI restriction sites. pQUASTattB(HTT$_{ex1}$Q25-GFP) and pQUASTattB(HTT$_{ex1}$Q91-GFP) were generated by replacing the mCherry sequence in pQUASTattB(HTT$_{ex1}$Q25-mCherry) and pQUASTattB(HTT$_{ex1}$Q91-mCherry) plasmids previously generated by our lab (Pearce et al., 2015) with an in-frame, C-terminal GFP sequence via Gibson Assembly (New England Biolabs, Inc. Ipswich, MA). pUASTattB(pHluorin-HTT$_{ex1}$Q25-tdTomato) and pUASTattB(pHluorin-HTT$_{ex1}$Q91-tdTomato) plasmids were generated by PCR amplification of HTT$_{ex1}$ cDNAs and subcloning to replace the CD4 sequence in the pUASTattB(MApHS) plasmid (Han et al., 2014) via In-Fusion cloning (Takara Bio USA, Inc., Mountain View, CA). pUASTattB(mCherry-Galectin-3) and pUASTattB(mCherry-Galectin-8) were cloned by PCR amplification of human LGals3 and LGals8 cDNAs from the pHAGE-mKeima-LGALS3

and pHAGE-FLAG-APEX2-LGALS8 plasmids (Addgene plasmids #175780 and #175758, Watertown,

MA) (Eapen et al., 2021) and insertion downstream of an in-frame mCherry sequence in the pUASTattB

vector backbone via Gibson Assembly.

Plasmids were microinjected into *w-* embryos with the su(Hw)attP8, attP24, VK19, or VK27

φC31 attP integration sites at BestGene, Inc (Chino Hills, CA). Table 1 lists detailed genotype

information for all transgenic flies generated in this study.

Drosophila brain dissection and sample preparation

Adult fly brains were dissected in ice-cold phosphate-buffered saline (PBS) containing either no

detergent or 0.03% (PBS/0.03T), 0.1% (PBS/0.1T), or 0.3% (PBS/0.3T) Triton X-100. Dissected brains

were fixed in 4% paraformaldehyde (PFA) in the dark at room temperature (RT) for 20 minutes. For

imaging of GFP or mCherry fluorescence signals, brains were washed 7X in PBS/0.03T buffer before

incubation in Slowfade Gold Antifade Mountant (Invitrogen, Carlsbad, CA). When imaging pHluorin-

tagged and other pH-sensitive constructs, such as GFP-LAMP1, brains were washed 7X in PBS for at

least 50 minutes at RT before incubation in Slowfade. For immunostaining, brains were washed 7X in

PBS/0.3T after fixation, blocked in PBS/0.3T containing 5% normal goat serum (Lampire Biological

Laboratories, Pipersville, PA) for 30 min at RT, incubated in primary antibodies diluted in blocking

solution for 24-48 hours at 4°C, washed 7X in PBS/0.3T, and then incubated in secondary antibodies

diluted in blocking solution for 16-20 hours at 4°C. After a final set of 7X PBS/0.3T washes, dissected

brains were incubated in Slowfade. For Magic Red and LysoTracker staining, brains were dissected in

PBS and incubated in 1:1,000 LysoTracker Red DND-99 (Invitrogen, Carlsbad, CA) or 1:1,250 Magic

Red (ImmunoChemistry, Davis, CA) diluted in PBS for 20 minutes at RT. Brains were washed 5X in

PBS for 15 min total, fixed in 4% PFA in PBS/0.1T for 20 minutes at RT, washed 6X in PBS/0.1T for 30

min total, and incubated in Slowfade. Following incubation in Slowfade for 1-24 hours at 4°C in the

dark, all brains were bridge-mounted in Slowfade on glass microscopy slides under #1.5 cover glass,

and edges were sealed using clear nail polish.

Primary antibodies used in this study include: chicken anti-GFP (RRID: AB_300798; 1:1,000;

Abcam, Cambridge, UK), chicken anti-GFP (RRID: AB_2534023; 1:500; Thermo Fisher Scientific Inc.,

Waltham, MA), rat anti-N-cadherin (clone DN-Ex #8; RRID: AB_528121; 1:75; Developmental Studies

Hybridoma Bank, Iowa City, IA), mouse anti-Repo (clone 8D12; RRID: AB_528448; 1:25;

Developmental Studies Hybridoma Bank, Iowa City, IA), mouse anti-V5 (RRID: AB_2556564; 1:125;

Thermo Fisher Scientific Inc., Waltham, MA), and rabbit anti-Draper (1:500; a kind gift from Marc

Freeman, Vollum Institute, Portland, OR). Secondary antibodies include: AlexaFluor 405 goat anti-

rabbit (RRID: AB_221605; 1:250), FITC-conjugated donkey anti-chicken (RRID: AB_2340356; 1:250;

Jackson Immuno Research Labs, West Grove, PA), AlexaFluor 568 goat anti-mouse (RRID:

AB_2534072; 1:250), AlexaFluor 647 goat anti-rabbit (RRID: AB_2535812; 1:250), and AlexaFluor 647

goat anti-rat IgGs (RRID: AB_141778; 1:250) (Invitrogen, Carlsbad, CA).

Image acquisition

All microscopy data were collected on a Leica SP8 laser-scanning confocal system equipped

with 405 nm, 488 nm, 561 nm, and 633 nm lasers and 40X 1.3 NA or 63X 1.4 NA oil objective lenses.

Leica LAS-X software was used to establish optimal settings during each microscopy session and to

collect optical z-slices of whole-mounted brain samples with Nyquist-optimized sampling criteria.

Optical zoom was used to magnify and establish a region of interest (ROI) in each sample. For images

showing a single glomerulus, confocal slices were collected to generate ~73 x 73 x 20 µm (*xyz*) stacks,

with z-axis boundaries established using fluorescent signal in DA1 or VA1lm ORN axons. For images of

a single antennal lobe, confocal slices were collected to generate ~117 x 117 x 26 µm (*xyz*) stacks,

which were located using HTT_{ex1} fluorescence in ORN axons.

Post-imaging analysis

Raw confocal data were analyzed in 2D using ImageJ/FIJI (RRID:SCR_002285; NIH, Bethesda,

MD) or in 3D using Imaris image analysis software (RRID:SCR_007370; Bitplane, Zürich, Switzerland).

Methods used for image segmentation and semi-automated quantification of fluorescent signals were

previously described (Donnelly et al., 2020). Briefly, raw confocal data were cropped to establish an

ROI for further analysis and displayed as a 2D sum intensity projection (ImageJ) or a 3D volume

(Imaris). Background fluorescence was subtracted from raw confocal images. mCD8-GFP and HTT_{ex1}-

mCherry fluorescent signals was segmented using the 'Surfaces' function in Imaris (surface detail =

0.25 μm, background subtraction = 0.75 μm). Using the 'split touching objects' option, seed point

diameter was set to 0.85 μm. pHluorin- and tdTomato-labeled VA1lm ORN axons were quantified in

central 30 x 30 pixels, 50-slice ROIs from sum intensity projections of each VA1lm glomerulus.

pHluorin- and tdTomato-labeled Or83b+ ORN axons or MBN soma were quantified from central 100 x

100 pixels, 75-slice ROIs of each antennal lobe or mushroom body calyx, and background intensity was

subtracted from sum intensity projections. Quantification of Toll-6[MIMICGFP] fluorescence was performed

using the 'Spots' function in Imaris (xy diameter = 0.5 μm, z-diameter = 1.0 μm), and glial nuclei-

associated GFP-Jra signal was quantified using the "Surfaces" function to identify Repo+/GFP+ nuclei

(surface detail = 0.2 μm, background subtraction = 1.6 μm, number of voxels >10).

171   mCherry-tagged $mHTT_{ex1}$ and GFP-tagged $wtHTT_{ex1}$ aggregates were identified and quantified

as previously reported (Donnelly et al., 2020). Briefly, mCherry+ surfaces were segmented and

measured in Imaris (surface detail = 0.2 μm, background subtraction = 0.45 μm, seed point diameter =

0.85 μm). Seeded $wtHTT_{ex1}$ aggregates were identified as $mHTT_{ex1}$-mCherry+ objects that overlapped

with $wtHTT_{ex1}$-GFP signal. Intracellular vesicles were identified using the 'Surfaces' algorithm in Imaris

(surface detail = 0.2 μm, background subtraction = 0.4 μm, volume >0.001 μm^3) to segment fluorescent

signals associated with lysosomes (MR+, LTR+, LAMP1-GFP+, GFP-LAMP1+, mCherry-Galectin-3+,

or mCherry-Galectin-8+) or phagosomes (YRab+ or YFP-Rab+). Intracellular vesicles associated with

$mHTT_{ex1}$ aggregates were identified by filtering for lysosomal or phagosomal surfaces within 0.2 μm of

$mHTT_{ex1}$ objects using the "Shortest Distance" calculation in Imaris.

qPCR

183   Transgenic *Drosophila* were flash frozen in liquid nitrogen, and heads were collected using a

microsieve with a 230 nm filter to separate bodies from heads and 170 nm filter to separate heads from

appendages. Total RNA was extracted from isolated fly heads using the Zymo Direct-Zol RNA miniprep

kit (Zymo Research, Irvine, CA). Extracted RNA was quantified on a Nanodrop 2000 (Thermo Fisher

Scientific), and equal quantities of each sample were subjected to cDNA synthesis using the High-

Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Waltham, MA). qPCR was performed

on a T100 Thermal Cycler with a CFX384 Real-Time System (Bio-Rad Laboratories, Hercules, CA)

using 10 ng of input RNA per replicate and TB Green Premix Ex Taq II (Takara Bio, Kusatsu, Japan).

Sequences of all qPCR primers used in this study are listed in Table 3.

Experimental design and statistical analyses

All quantified data were organized and analyzed in Prism 9 software (Graphpad, San Diego,

CA). Power analyses to determine appropriate sample sizes for each experiment were calculated using

a Sample Size calculator available at ClinCalc.com ($\alpha = 0.05$, $\beta = 0.2$). All quantifications are graphed

as mean ± standard error of the mean (s.e.m). A single glomerulus or antennal lobe represented one

biological replicate. Statistical comparisons were performed using the following tests where appropriate:

unpaired, two-tailed *t*-test when comparing two samples and ANOVA followed by *post hoc* multiple

comparison tests when comparing ≥3 samples. Detailed statistical information for each experiment,

including sample sizes (*n*), means, and test results are summarized in Table 4.

RESULTS

Expression of mHTT exon 1 fragments in neurons inhibits phagocytic clearance of axonal debris

HD is a monogenic neurodegenerative disease caused by expansion of a CAG repeat region in exon 1 of the *huntingtin (HTT/IT15)* gene beyond a pathogenic threshold of 37 repeats, leading to production of mHTT proteins containing expanded N-terminal polyglutamine (polyQ≥37) tracts (Scherzinger et al., 1999). mHTT proteins are prone to misfolding and accumulate into insoluble aggregates (Wanker, 2000), whereas wild-type HTT (wtHTT) proteins containing polyQ≤36 tracts remain soluble unless seeded by a pre-formed HTT aggregate (Preisinger et al., 1999; Chen et al., 2001). To recapitulate these molecular features of HD, we generated transgenic flies that express N-terminal exon 1 fragments of human mHTT (mHTT$_{ex1}$) containing a polyQ91 tract or wtHTT (wtHTT$_{ex1}$) with a polyQ25 tract via the GAL4-UAS (Brand and Perrimon, N., 1993) or QF-QUAS binary expression systems (Potter and Luo, 2011). As we have previously reported (Pearce et al., 2015; Donnelly et al., 2020), fluorescent protein fusions of mHTT$_{ex1}$ and wtHTT$_{ex1}$ appeared punctate (i.e., aggregated or insoluble) or diffuse (i.e., soluble), respectively, in axon termini of the DA1 type of olfactory receptor neurons (ORNs), which synapse in the DA1 glomerulus of the antennal lobe in the central brain of adult flies (Couto et al., 2005) (Fig. 1A and B).

To monitor the capacity of glia to maintain CNS homeostasis in the presence of mHTT$_{ex1}$ aggregates, we performed a series of experiments that quantified glial responses to acute injury in the adult fly CNS. Surgical removal of the second and third antennal segments initiates Wallerian degeneration of ORN axons, inducing a robust phagocytic glial response that involves upregulation of the scavenger receptor, Draper, and clearance of axonal debris within 7 days (MacDonald et al., 2006). To determine if neuronal mHTT$_{ex1}$ expression affects the ability of glial cells to efficiently clear axonal debris, we co-expressed mCherry-tagged HTT$_{ex1}$ transgenes with membrane-targeted GFP (mCD8-GFP) in DA1 ORNs and measured GFP and mCherry fluorescence intensities following antennal nerve axotomy (Fig. 1A-D). Interestingly, mHTT$_{ex1}$ expression was associated with reduced steady-state mCD8-GFP levels in DA1 ORN axons in 2 and 4 week-old flies (Fig. 1E), likely due to neurotoxicity caused by accumulation of mHTT$_{ex1}$ aggregates over time. Quantification of DA1 ORN axons remaining

after antennal nerve injury revealed that clearance of axonal debris was reduced in flies co-expressing

neuronal mHTT$_{ex1}$ compared with controls expressing wtHTT$_{ex1}$ (Fig. 1C). This effect could be observed

as early as 1 day post-injury, suggesting that mHTT$_{ex1}$ causes defects in both clearance and

engulfment of axonal debris. Delayed clearance of axonal debris was exacerbated in older (14 and 28

day-old) mHTT$_{ex1}$-expressing flies (Fig. 1C), possibly related to a decline in Draper activity during

normal aging (Purice et al., 2016) and/or enhanced glial dysfunction compounded by mHTT$_{ex1}$

aggregate accumulation. Quantification of mCherry fluorescence indicated that clearance of axonal

mHTT$_{ex1}$ was also slowed compared to wtHTT$_{ex1}$ (Fig. 1D), suggesting that glia are deficient in

degrading both axonal debris and neuronal mHTT$_{ex1}$ aggregates.

We also tested whether formation of mHTT$_{ex1}$ aggregates in glial cells impacts ORN axonal

debris clearance following acute injury. Restricting expression of mHTT$_{ex1}$ to glia resulted in

appearance of heterogeneously-sized mHTT$_{ex1}$ aggregates throughout the brain (Fig. 2A). Glial

mHTT$_{ex1}$ aggregates also slowed injury-induced clearance of mCD8-GFP-labeled axons compared with

wtHTT$_{ex1}$-expressing controls (Fig. 2B-D), though glial mHTT$_{ex1}$ expression did not affect DA1 ORN

axon abundance in 1 day-old flies (Fig. 2E). Together, these findings indicate that mHTT$_{ex1}$ aggregates

originating in either neurons or glia slow efficient clearance of injured ORN axons by phagocytic glia.

Neuronal mHTT aggregates impair nascent phagosome formation

Phagocytosis occurs via 4 major steps: (1) extension of phagocyte membranes toward

extracellular "find me" cues, (2) recognition of "eat me" signals by scavenger receptors, (3) cytoskeletal

and plasma membrane reorganization to surround and internalize extracellular material, and (4)

maturation of nascent phagosomes through sequential endomembrane fusions that culminate at the

lysosome (Vieira et al., 2002). We have previously reported that the conserved scavenger receptor,

Draper/Ced-1/MEGF10 (MacDonald et al., 2006; Evans et al., 2015; Iram et al., 2016), regulates

engulfment, clearance, and intercellular spreading of mHTT$_{ex1}$ aggregates originating in ORN axons

(Pearce et al., 2015; Donnelly et al., 2020). To determine whether delayed clearance of injured axons

containing mHTT$_{ex1}$ aggregates (Fig. 1C and 2D) could result from defective Draper-dependent

engulfment, we co-expressed V5-tagged mHTT$_{ex1}$ or wtHTT$_{ex1}$ with the ratiometric membrane-

associated pH sensor (MApHS), consisting of the transmembrane domain of CD4 flanked by N-terminal

ecliptic pHluorin and C-terminal tdTomato (Han et al., 2014), in the VA1lm type of ORNs. Ecliptic

pHluorin is brightest at pH 7.5 and dims as pH drops, quenching at pH <6.0 (Miesenböck et al., 1998),

whereas tdTomato fluorescence is pH resistant. Thus, internalization of MApHS-labeled ORN axonal

debris into a rapidly-acidified nascent phagosome can be monitored by calculating pHluorin:tdTomato

fluorescence intensity ratios (Han et al., 2014). Indeed, pHluorin:tdTomato ratios in VA1lm axons were

decreased at 14 and 25 hours post-axotomy (Fig. 3A-C), and this effect was lost when *draper* was

deleted from glia using the tissue-specific CRISPR/Cas9-TriM method (Fig. 3C and E-G) (Poe et al.,

2019), demonstrating that this construct accurately reports nascent phagosome acidification following

engulfment. Notably, the decrease in pHluorin:tdTomato ratio following injury was less pronounced in

VA1lm ORN axons co-expressing mHTT$_{ex1}$ compared with wtHTT$_{ex1}$-expressing controls (Fig. 3A-C),

suggesting that mHTT$_{ex1}$ aggregates impair nascent phagosome formation following engulfment.

Neuronal mHTT accumulates in low pH intracellular compartments

271   mHTT$_{ex1}$ expression was associated with a slight but significant decrease in steady-state

pHluorin:tdTomato ratios in MApHS-labeled axons compared with wtHTT$_{ex1}$-expressing controls (Fig.

3D), suggesting that even in the absence of acute injury, mHTT$_{ex1}$ signals for ORN axon engulfment. To

test this, we generated transgenic flies that express mHTT$_{ex1}$ or wtHTT$_{ex1}$ fused to N-terminal pHluorin

and C-terminal tdTomato fluorescent proteins, herein referred to as mHTT-associated pH sensor

(mHApHS) or wtHTT-associated pH sensor (wtHApHS). Accumulation of HTT$_{ex1}$ protein in low pH

cellular compartments would cause a decrease in the ratio of pHluorin:tdTomato fluorescence for these

constructs. wtHApHS and mHApHS transgenes were expressed in either Or83b+ ORNs, which

encompass 70-80% of all adult ORNs (Fig. 4A-B) (Larsson et al., 2004), or mushroom body neurons

(MBNs; Fig. 4D-E), which are downstream in the fly olfactory circuit and innervate the learning and

memory center of the fly CNS (McGuire et al., 2001). In both ORN axons and MBN soma,

pHluorin:tdTomato ratios associated with mHApHS were significantly decreased compared with

wtHApHS controls (Fig. 4C and F), suggesting that $mHTT_{ex1}$ proteins accumulate in acidified cellular

compartments. To discriminate $mHTT_{ex1}$ proteins engulfed by glia from $mHTT_{ex1}$ internalized into

neuronal autophagolysosomes, we measured VA1lm ORN axonal mHApHS-associated fluorescence in

animals with glial *draper* loss-of-function (Fig. 4G-H). Glial *draper* knockout increased

pHluorin:tdTomato ratios for axonal mHApHS (Fig. 4I), suggesting that at least some portion of

$mHTT_{ex1}$ aggregates accumulate in acidic portions of the glial phagolysosomal system.

Neuronal mHTT impairs injury-responsive gene upregulation

Glia alter their transcriptional profile to elicit cellular responses to insult or injury in the brain. For

example, acute CNS injury in *Drosophila* increases transcription of many genes involved in

phagocytosis and innate immunity, including the cell surface receptors, Draper and Toll-6, and

components of their downstream signaling pathways (Fig. 5A) (Purice et al., 2017; Byrns et al., 2021;

Alphen et al., 2022). To test whether the reduced ability of glia to clear $mHTT_{ex1}$ -containing axonal

debris correlates with reduced injury responsiveness at a transcriptional level, we used qPCR and

GFP-tagged reporters to quantify changes in gene expression of key components of these phagocytic

and innate immunity pathways following $mHTT_{ex1}$ accumulation in neurons. $mHTT_{ex1}$ expression in

uninjured Or83b+ ORNs increased relative expression of *toll-6, relish*, and *drpr-I* transcripts between

1.2- and 1.5-fold (Fig. 5B) and levels of GFP-tagged Toll-6 and Jra proteins in the CNS (Fig. 5C-H),

suggesting that neuronal $mHTT_{ex1}$ aggregates activate a mild injury response in the brain. Jra-GFP

expression increased throughout the CNS and in Repo+ nuclei (Fig. 5F-H), suggesting that $mHTT_{ex1}$-

induced upregulation of this subunit of the AP-1 transcription factor can be at least partially attributed to

glia. To test whether $mHTT_{ex1}$ impairs the ability of glia to respond to acute neural injury, we monitored

gene expression changes after bilateral antennal and maxillary palp nerve ablation. Similar to previous

reports (Purice et al., 2017), expression of *toll-6, dorsal, relish, drpr-I, mmp1,* and *ets21c* genes was

significantly increased 3 hrs after injury to ORN axons expressing either mCD8-GFP or $wtHTT_{ex1}$ (Fig.

5B). However, injury-induced upregulation of each of these genes was significantly reduced in animals

expressing $mHTT_{ex1}$ in Or83b+ ORNs (Fig. 5B), suggesting that $mHTT_{ex1}$ aggregation attenuates glial

transcriptional responses to injury. We further analyzed the impact of mHTT$_{ex1}$ on downstream immune

responses in the brain by measuring induction of antimicrobial peptide (AMP) genes, well-established

transcriptional targets of activated Relish/NFκB following Toll-6 or immune-deficiency pathway

activation (Swanson et al., 2020b, 2020a; Alphen et al., 2022). Levels of five AMP genes, including

drosomycin, *attacinA*, *attacinD*, *diptericinA*, and *metchnikowin*, were significantly increased 3 hours

after ORN axotomy (Fig. 6A-E), similar to previous reports (Katzenberger et al., 2013; Swanson et al.,

2020b; Marischuk et al., 2021). Interestingly, mHTT$_{ex1}$ expression in ORNs alone was sufficient to

induce upregulation of *drosomycin* and *attacinD* (Fig. 6A & C), albeit to a lesser extent than following

acute injury. Further, injury-induced upregulation of *drosomycin* and *attacinA* was significantly reduced

by mHTT$_{ex1}$ expression in ORNs, suggesting that the activity of Relish-dependent signaling is altered by

accumulation of mHTT$_{ex1}$ aggregates. Together, these data indicate that neuronal mHTT$_{ex1}$ aggregates

trigger a mild immune response in the brain, but also inhibit the ability of glia to mount robust

transcriptional responses to neural injury.

Our findings are in agreement with previously published work demonstrating that expression of

mHTT$_{ex1}$ and Aβ in neurons activates Draper-dependent phagocytosis, likely in an effort to reduce

levels of these pathogenic proteins in the brain (Pearce et al., 2015; Ray et al., 2017). We further

investigated this by immunostaining adult fly brains expressing mHTT$_{ex1}$ in ORNs to monitor expression

levels and localization of endogenous Draper protein. In 1 week-old adult flies expressing neuronal

mHTT$_{ex1}$, Draper immunolabeling increased ~1.2-fold in the vicinity of ORN axons compared with age-

matched controls expressing wtHTT$_{ex1}$ (Fig. 7A-C). Closer analysis revealed that in some cases,

Draper immunofluorescence was directly adjacent to or surrounding mHTT$_{ex1}$-mCherry fluorescence

(Fig. 7D). To further examine these interactions, we used image segmentation and three-dimensional

reconstruction of confocal stacks to represent mHTT$_{ex1}$-mCherry+ aggregates and Draper+ glial

membranes as individual "surfaces" (Fig. 7E-F), as previously described (Donnelly et al., 2020). Close

physical association of mHTT$_{ex1}$ with Draper and other glial proteins was defined as surfaces located

≤0.2 μm from a mHTT$_{ex1}$ object. Interestingly, whereas almost no Draper signal was detected near

wtHTT$_{ex1}$ (Fig. 7E), ~14% of mHTT$_{ex1}$ aggregates were closely associated with Draper surfaces (Fig.

7F1-3). These findings are consistent with Draper+ glial membranes being recruited to neuronal

mHTT$_{ex1}$ aggregates to facilitate engulfment.

Synaptic neuropil in the *Drosophila* brain is primarily inhabited by two glial subtypes,

ensheathing glia and astrocytic glia (Doherty et al., 2009). In adult flies, ensheathing glia

compartmentalize synaptic regions and respond to CNS injury by upregulating Draper and clearing

debris via phagocytosis (Doherty et al., 2009). Consistent with our previous findings in all repo+ glia

(Pearce et al., 2015), mz0709+ ensheathing glia were vulnerable to prion-like conversion of glial

wtHTT$_{ex1}$ proteins by mHTT$_{ex1}$ aggregates generated in DA1 ORN axons (Fig. 8A, C-D). Seeded

wtHTT$_{ex1}$ aggregates were defined as wtHTT$_{ex1}$ signal that colocalized with mHTT$_{ex1}$ objects identified

by image segmentation of confocal stacks (Donnelly et al., 2020). Conversely, seeding of wtHTT$_{ex1}$

expressed in alrm+ astrocytic glia, which lack detectable Draper expression in the adult fly brain

(Doherty et al., 2009), was not observed (Fig. 8B-D). Interestingly, ensheathing glial-specific RNAi

knockdown of Toll-6, Relish, and NijA increased mHTT$_{ex1}$ aggregate numbers in DA1 ORN axons,

similar to the effects of Draper-I knockdown (Fig. 9A-B). Further, adult-specific, pan-glial knockdown of

Ets21c, which was found to be required for normal development, also increased numbers of mHTT$_{ex1}$

aggregates in DA1 ORN axons (Fig. 9C-D). Mean mHTT$_{ex1}$ aggregate volume was not affected by

ensheathing glial knockdown of these genes except for a ~20% increase following NijA depletion (Fig.

9E), suggesting that aggregate or vesicle size was unaffected by these genetic manipulations. Thus,

several glial genes with established roles in phagocytic and innate immune signaling regulate basal

turnover of mHTT$_{ex1}$ aggregates in ORN axons.

Neuronal mHTT aggregates are associated with defects in multiple endolysosomal compartments

Our findings thus far indicate that neuronal mHTT$_{ex1}$ aggregates elicit mild injury responses in

the brain and reduce the ability of phagocytic glia to respond transcriptionally and functionally to acute

nerve injury. We have previously postulated that prion-like spreading of mHTT$_{ex1}$ in the fly CNS could

be facilitated by escape of engulfed aggregates from the endolysosomal compartment of phagocytic

glia (Pearce et al., 2015; Donnelly et al., 2020). Therefore, we sought to determine whether neuronal

mHTT$_{ex1}$ aggregates are associated with defects in endolysosomal processing in non-injured brains.

We first measured effects of neuronal mHTT$_{ex1}$ expression on the quantity, size, and function of

lysosomes using dyes that label active cathepsins and low pH cellular compartments. Expression of

mHTT$_{ex1}$ in Or83b+ ORNs increased numbers of lysosomes labeled by the active cathepsin dye, Magic

Red (MR) (Fig. 10A-C), and the low pH sensor, LysoTracker Red (LTR) (Fig. 10D-F), compared with

control brains expressing wtHTT$_{ex1}$. High resolution analysis of confocal stacks and filtering for

segmented MR+ and LTR+ surfaces within 0.2 μm of a mHTT$_{ex1}$ object revealed close association of

MR+ and LTR+ signals with mHTT$_{ex1}$ aggregates (Fig. 10C and F-H).

We next employed *Drosophila* genetic tools to assess the impact of neuronal mHTT$_{ex1}$

specifically on glial lysosomes by driving expression of lysosomal-associated membrane protein 1

(LAMP1) tagged at its cytosolic C-terminus with GFP (LAMP1-GFP) in all glia. Neuronal mHTT$_{ex1}$

expression increased the overall number of glial LAMP1-GFP+ vesicles and the number of LAMP1-

GFP+ vesicles in close proximity to HTT$_{ex1}$ signal (Fig. 11A-C and G-H). We typically observed only

partial overlap of LAMP1-GFP signal with mHTT$_{ex1}$ aggregate surfaces identified by this method,

possibly due to incomplete labeling or rupture of lysosomal membranes as a result of aggregate size or

structural features. Interestingly, this subpopulation of LAMP1-GFP+ lysosomes closely associated with

mHTT$_{ex1}$ were enlarged compared with all lysosomes (Fig. 11I). To examine whether neuronal mHTT$_{ex1}$

aggregates affect lysosome integrity, we expressed a transgene encoding LAMP1 fused at its N-

terminus to GFP in glia. This construct integrates into the lysosomal membrane such that GFP is

exposed to the lumen, and loss of GFP signal can thus be used to monitor LAMP1+ lysosome

degradative activity (Pulipparacharuvil et al., 2005). Interestingly, the quantity and mean volume of

GFP-LAMP1+ vesicles were significantly increased in brains expressing mHTT$_{ex1}$ in Or83b+ ORNs

(Fig. 11D-G), suggesting lysosomal enlargement and dysfunction due to mHTT$_{ex1}$ aggregates. GFP-

LAMP1+ surfaces were also more associated with mHTT$_{ex1}$ aggregates compared to wtHTT$_{ex1}$ controls

(Fig. 11F and I).

To directly test whether glial endolysosomes experience increased membrane damage due to

neuronal mHTT$_{ex1}$ expression, we generated transgenic flies expressing mCherry-tagged Galectin-3 or

Galectin-8, lectins that translocate from the cytoplasm to the lumen of ruptured lysosomes and endosomes, respectively (Aits et al., 2015; Daussy and Wodrich, 2020; Jia et al., 2020). Neuronal $mHTT_{ex1}$ expression increased overall numbers of glial Galectin-3/8+ surfaces (Fig. 12A-E), Galectin-3/8+ surfaces closely associated with $mHTT_{ex1}$ aggregates (Fig. 12F), and mean volume of $mHTT_{ex1}$-associated Galectin-3/8+ vesicles (Fig. 12G). Together, these data suggest that neuronal $mHTT_{ex1}$ aggregates induce non cell-autonomous accumulation, enlargement, and membrane damage of endolysosome vesicles in glial cells.

"Seeding-competent" mHTT aggregates are defined by their ability to nucleate or "seed" the aggregation of normally-soluble wtHTT proteins, a defining characteristic of infectious prion and other prion-like proteins (Jucker and Walker, 2018; Donnelly et al., 2022). Many studies have pointed to a role for defective clearance by endolysosomal pathways in promoting the propagation of pathogenic aggregates (Freeman et al., 2013; Jiang et al., 2017; Chen et al., 2019; Jiang and Bhaskar, 2020; Polanco and Götz, 2022). To test whether altered glial lysosome function affects seeding competency of $mHTT_{ex1}$, we used RNAi to individually knockdown proteins with known roles in lysosome degradation in glia and examined the ability of neuronal mCherry-tagged $mHTT_{ex1}$ to seed aggregation of glial, GFP-tagged $wtHTT_{ex1}$. Depletion of two subunits of the vacuolar ATPase (V-ATPase), Vha68-3 (Portela et al., 2018) and rabconnectin-3A (Yan et al., 2009), and Spinster, a late-endosomal and lysosomal efflux permease (Rong et al., 2011), increased numbers of glial $wtHTT_{ex1}$ aggregates detected as GFP+ surfaces that colocalized with $mHTT_{ex1}$ aggregates (Fig. 13A-C) (Donnelly et al., 2020). Knockdown of Vha16-1, a V-ATPase subunit that regulates endolysosome membrane fusion (Finbow et al., 1994; Dunlop et al., 1995), did not affect $wtHTT_{ex1}$ seeding; however, it did cause accumulation of $mHTT_{ex1}$ aggregates in DA1 ORN axons (Fig. 13A and B). Together, these findings suggest that disruption of normal glial lysosome acidification and/or degradative capacity promotes formation of seeding-competent $mHTT_{ex1}$ aggregates.

The GTPase Rab10 mediates prion-like transmission of mHTT aggregates

Lysosome dysfunction could occur secondary to upstream defects in endo/phagosome

maturation. Our prior work supports a model in which a portion of mHTT$_{ex1}$ aggregates engulfed by glia

evade degradation during phagosome maturation and/or phagolysosome formation (Pearce et al.,

2015; Donnelly et al., 2020). To test this model, we used forward genetic screening to interrogate roles

for glial Rab GTPases in prion-like conversion of cytoplasmic wtHTT$_{ex1}$ proteins by engulfed neuronal

mHTT$_{ex1}$ aggregates. The *Drosophila* genome encodes 31 Rab and Rab-like proteins, all of which have

mammalian orthologs, and most of these GTPases are implicated in vesicle and target membrane

fusion in cells (Zhang et al., 2007). To determine whether any *Drosophila* Rabs mediate escape of

phagocytosed mHTT$_{ex1}$ aggregates and seeding of wtHTT$_{ex1}$ in the glial cytoplasm, we individually

knocked down each Rab in repo+ glia using RNAi. Glial-restricted silencing of 23 of the 31 Rabs

produced viable adults, and these flies were used to monitor effects of Rab knockdown on mHTT$_{ex1}$-

induced aggregation of wtHTT$_{ex1}$ in glia. Only two Rab RNAi lines, *Rab10^{RNAi}* and *Rab23$^{RNAi\#2}$*,

significantly altered numbers of induced wtHTT$_{ex1}$ aggregates (Fig. 14A-C). Of note, 3 additional Rab23

RNAi lines had no significant effects on numbers of wtHTT$_{ex1}$ aggregates, suggesting that *Rab23$^{RNAi\#2}$*

may cause off-target effects. Strikingly, Rab10 depletion reduced numbers of seeded wtHTT$_{ex1}$

aggregates, phenocopying effects of Draper knockdown (Fig. 14A and C) and suggesting that Draper

and Rab10 function in the same pathway. To test whether this effect of Rab10 loss-of-function was

mediated via a reduction in Draper expression, we measured endogenous Draper immunofluorescence

in *rab10* null flies. Draper protein levels were ~18% lower in *rab10$^{-/-}$* animals compared to wild-type

controls (Fig. 14D). This reduction is unlikely to fully account for decreased seeding of wtHTT$_{ex1}$

following Rab10 knockdown, as numbers of wtHTT$_{ex1}$ aggregates are similar between wild-type and

draper heterozygotes (Fig. 14G) (Pearce et al., 2015). To further explore this, we attempted to restore

Draper function via overexpression of Draper-I, which rescues loss of Draper function in aged flies

(Purice et al., 2016). However, transgenic expression of Draper-I in glia failed to rescue the effects of

Rab10 knockdown on wtHTT$_{ex1}$ aggregate seeding (Fig. 14E). These findings suggest that Rab10 acts

downstream of Draper rather than by regulating Draper activity. We further tested for interactions

between *drpr* and *rab10* using loss-of-function alleles to examine effects on mHTT$_{ex1}$ aggregate

transmission from presynaptic DA1 ORNs to postsynaptic projection neurons (PNs), a process

previously reported to require transport though Draper+ glia (Donnelly et al., 2020). Because reduced

survival of *rab10* null flies (Kohrs et al., 2021) was exacerbated by transgenic HTT_{ex1} expression, we

tested for genetic interaction between *drpr* and *rab10* in heterozygous and trans-heterozygous animals.

While we detected no change in $mHTT_{ex1}$ aggregate numbers (Fig. 14F), significantly fewer $wtHTT_{ex1}$

aggregates formed in PNs of $rab10^{+/-}$ $drpr^{+/-}$ transheterozygotes than in individual heterozygotes (Fig.

14G). This same effect was not observed in $rab14^{+/-}$ $drpr^{+/-}$ transheterozygous animals (Fig. 14G),

indicating that the genetic interaction is specific to *drpr* and *rab10*. Thus, Draper and Rab10 appear to

function in the same phagocytic pathway that regulates prion-like transmission of phagocytosed

$mHTT_{ex1}$ aggregates in the fly brain.

Neuronal mHTT aggregates alter numbers of early and late glial phagosomes

Our forward genetic screen indicates that at least one glial Rab GTPase, Rab10, regulates the

seeding capacity of neuronal $mHTT_{ex1}$ aggregates. Interestingly, Rab10 has been reported to regulate

phagosome maturation in mammalian cells (Cardoso et al., 2010; Seto et al., 2011; Lee et al., 2020;

Wang et al., 2023), and Rab10 expression and activity are altered in neurodegenerative diseases,

including AD and PD (Eguchi et al., 2018; Tavana et al., 2018; Yan et al., 2018). Because little is known

about Rab10's role in glia, we sought to characterize this GTPase alongside two additional Rabs with

well-established roles in endocytosis, Rab5 and Rab7, markers of early and late endo/phagosomes,

respectively (Hutagalung and Novick, 2011). Interestingly, we found that *rab10*, *rab5*, and *rab7* were

upregulated between ~1.4 and 2.4-fold following acute injury to ORN axons, and $mHTT_{ex1}$ expression in

Or83b+ ORNs alone caused upregulation of *rab10* and *rab7* genes by ~1.2-fold (Fig. 15A).

Interestingly, injury-induced upregulation of *rab5* and *rab7* was inhibited by ~50% and ~100%,

respectively, in flies expressing $mHTT_{ex1}$ compared to controls (Fig. 15A). Altogether, these results

identify *rab5*, *rab7*, and *rab10* as novel injury-response genes in the fly CNS and suggest that neuronal

$mHTT_{ex1}$ aggregates impair injury-induced responses of *rab5* and *rab7*.

We next examined effects of neuronal mHTT$_{ex1}$ expression on the localization of Rab10, Rab5,

and Rab7 proteins endogenously-tagged with YFP-myc at their N-termini, herein referred to as

YRab10, YRab5, and YRab7 (Dunst et al., 2015). YRab+ vesicles were identified in confocal stacks as

segmented YFP+ surfaces with a mean diameter of 0.3-8 μm (Fig. 15B-G), consistent with vesicle sizes

reported in other fly tissues (Prince et al., 2019). Expression of mHTT$_{ex1}$ in Or83b+ ORN axons caused

an overall increase in numbers of YRab10+ and YRab7+ vesicles, but a decrease in the number of

YRab5+ vesicles compared with wtHTT$_{ex1}$ controls (Fig. 15H). Further, each of these YRab+ vesicles

subpopulations were closely associated with mHTT$_{ex1}$ aggregates more frequently than with wtHTT$_{ex1}$

(Fig. 15I), suggesting that mHTT$_{ex1}$ protein interacts with each of these intracellular vesicle

subpopulations in the brain.

To assess effects of neuronal mHTT$_{ex1}$ aggregates specifically on glial Rab+ compartments, we

expressed YFP-tagged Rab5, 7, and 10 transgenes in all glia. Similar to our findings with endogenous

YRabs, expression of mHTT$_{ex1}$ in Or83b+ ORN axons was associated with increased numbers of glial

YFP-Rab10+ and -Rab7+ vesicles (Fig. 16A-B, E-F, and G); however, we observed a significant

decrease in glial YFP-Rab5+ vesicle abundance (Fig. 16C-D and G). YFP-Rab10+, -5+, and -7+

vesicles increased their association with axonal mHTT$_{ex1}$ aggregates compared with wtHTT$_{ex1}$ (Fig.

16H), in many cases with closely associated YFP-Rab+ signal partially surrounding a mHTT$_{ex1}$

aggregate (Fig. 17). Of note, only a small fraction of YFP-Rab+ vesicles were identified as associated

with mHTT$_{ex1}$ aggregates, possibly due to transient interactions with vesicle compartments as

aggregates transit the glial phagolysosomal system, heterogeneous labeling of phagosomes by these

markers in intact brain tissue, or because our selection filter excluded YFP-Rab+ surfaces located >0.2

μm away from an aggregate. The mean volume of YFP-Rab7+ vesicles that interacted with mHTT$_{ex1}$

aggregates was significantly increased compared with wtHTT$_{ex1}$ controls (Fig. 16I), suggesting that

mHTT$_{ex1}$ leads to enlargement of Rab7+ late phagolysosomes. Together, these data suggest that

accumulation of phagocytosed mHTT$_{ex1}$ aggregates in Rab7+ or Rab10+ late phagosomes and

decreased association with early Rab5+ phagosomes could be a key mechanism underlying protein

aggregate-induced toxicity and spreading in HD.

DISCUSSION

Toxic amyloid aggregates have been a primary target in neurodegenerative disease drug development for decades, with some recent promise using immunotherapy to reduce aggregate loads in the brain (Karran and De Strooper, 2022). Microglia and astrocytes have also emerged as attractive therapeutic targets in efforts to boost neuroprotective glial functions or reduce neuroinflammation. However, approaches that target glial cells must effectively strike a balance between amplifying beneficial and reducing harmful effects of these cells in the brain. Here, we tested for interactions between phagocytic glia and pathogenic protein aggregates in a *Drosophila* model of HD. We report that aggregates formed by mHTT$_{ex1}$ protein fragments impair glial transcriptional and functional responses to CNS injury, induce upregulation of stress response and innate immunity genes, and alter numbers of endolysosomal vesicles detected in uninjured brains. A targeted forward genetic screen revealed that Rab10, a GTPase previously reported to regulate phagosome maturation, mediates prion-like conversion of cytoplasmic wtHTT$_{ex1}$ proteins by phagocytosed mHTT$_{ex1}$ aggregates. Together, these findings suggest that neuronal mHTT$_{ex1}$ aggregates compromise intracellular membrane integrity as they transit endolysosomal systems, generating toxic, seeding-competent aggregates that propagate disease phenotypes.

Glia respond to neural injury by altering their transcriptional, morphological, and metabolic profiles to promote neuronal survival and clear debris from the brain; however, failure of glia to return to a resting state elicits harmful neuroinflammatory consequences (Liddelow et al., 2020). We have previously reported that activated phagocytic glia can have both beneficial (i.e., elimination of toxic aggregates) and harmful (i.e., as vectors for aggregate spread) effects in the brain (Pearce et al., 2015; Donnelly et al., 2020). We report here that key glial injury-responsive pathways, i.e., Draper-mediated phagocytosis and Toll-6-mediated innate immune signaling, are induced in the presence mHTT$_{ex1}$ aggregation in the adult fly brain. These findings are in line with studies from other labs demonstrating that *Drosophila* Toll-6 and mammalian Toll-Like Receptor signaling pathways are upregulated in response to dying neurons during development (McLaughlin et al., 2019) and in patient and mammalian models of neurodegenerative disease (Casula et al., 2011; Miron et al., 2018; Kouli et al., 2020).

Interestingly, increased microglial NF-κB signaling mediates tau spread and toxicity in mice, further linking innate immunity to prion-like mechanisms of disease progression (Wang et al., 2022). Thus, activation of glial immune pathways may contribute to feed-forward mechanisms involving aggregate formation, pathology propagation, and neuroinflammatory signaling.

Genome-wide association studies have revealed numerous genes associated with increased risk of AD and other neurodegenerative diseases, and many of these risk variants are enriched in pathways that control key glial cell functions. For example, rare risk-associated variants of the microglial *TREM2* gene alter amyloid aggregate accumulation and seeding in cells and animal models (Leyns et al., 2019; Parhizkar et al., 2019; Jain et al., 2023). A number of additional genes involved in endolysosomal processing are associated with increased risk of AD, PD, FTD, and/or ALS, such as the phagocytic receptor *CD33*, endosomal genes *BIN1* and *RIN3*, and *GRN*, which encodes the lysosomal progranulin protein (Podleśny-Drabiniok et al., 2020; Welikovitch et al., 2023). Although Draper/MEGF10 variants are not known risk factors in neurodegenerative disease, MEGF10 is highly expressed in phagocytic astrocytes (Chung et al., 2013), mediates Aβ aggregate engulfment (Singh et al., 2010; Fujita et al., 2020), and acts as a receptor for C1q, a mediator of early synapse loss in AD mouse models (Hong et al., 2016; Iram et al., 2016). Our finding that upregulation of *draper* and other phagocytic genes is inhibited by mHTT$_{ex1}$ expression suggests that glial responsiveness and phagocytic capacity is attenuated in the presence of protein aggregate pathology in neurons. Genetic or environmental risk factors that impact glial health could accelerate these defects and exacerbate aggregate-induced neurotoxicity in the brain.

Lysosomes are essential for cell survival, not only to clear damaged or toxic materials from cells, but also as intracellular centers for macromolecule recycling, energy metabolism, and cell-cell communication. Lysosomal abnormalities, including vesicle enlargement, deacidification, and membrane leakiness, have been reported in patient brains and animal and cell models of multiple neurodegenerative diseases, suggesting that these defects play a central role in disease progression (Bonam et al., 2019; Polanco and Götz, 2022; Udayar et al., 2022). We report here that Rab7+, Rab10+, and Lamp1+ phagolysosomes accumulate in glia as a result of mHTT$_{ex1}$ expression in

neurons and that lysosome dysfunction and membrane permeabilization increase in the presence of mHTT$_{ex1}$ aggregates. We also observed that neuronal aggregates were associated with decreased nascent phagosome formation and reduced numbers of early (Rab5+) phagosomes in glia. These findings are consistent with a "traffic jam" model (Small et al., 2017) in which mHTT$_{ex1}$ aggregates accumulate over time in glial endolysosomal compartments, preventing proper flow of materials into (i.e., engulfment) and out of (i.e., degradation) the pathway (Fig. 18). We postulate that persistence of aggregates in faulty endolysosomes promotes formation and/or release of degradation-resistant aggregates with enhanced toxicity and seeding capacity. This hypothesis is in agreement with recent studies in tauopathy models showing that Aβ aggregation occurs secondary to lysosome deacidification and membrane permeabilization (Lee et al., 2022), and hypophagocytic glia contribute to tau aggregate propagation (Hopp et al., 2018; Brelstaff et al., 2021). Release of seeding-competent aggregates from dysfunctional lysosomes could occur via active exocytosis, perhaps in an effort to alleviate cell toxicity, or passively due to vesicle rupture (Flavin et al., 2017; Falcon et al., 2018; Yuste-Checa et al., 2021).

Among the growing list of genes linked to neurodegenerative disease pathogenesis are Rab GTPases and genes that modify Rab functions (Kiral et al., 2018). Rab proteins are essential for vesicle sorting and trafficking in all cells and use GTP hydrolysis to differentially associate with and organize intracellular membranes (Homma et al., 2021). In a forward genetic screen aimed at identifying Rabs that regulate phagocytic processing of neuronal mHTT$_{ex1}$ aggregates, we uncovered glial Rab10 as a modifier of prion-like spreading of mHTT$_{ex1}$ from neurons to glia. Interestingly, several Rab proteins have been previously reported to alter secretion or cell-to-cell propagation of tau or α-synuclein aggregates (Rodriguez et al., 2017; Bae et al., 2018; Ugbode et al., 2019; Rodrigues et al., 2022), suggesting that Rab-dependent endomembrane fusion could be exploited by prion-like aggregates. Our data suggest that Rab10 acts downstream of Draper to enable engulfed mHTT$_{ex1}$ aggregates to evade lysosomal degradation, perhaps escaping to the cytoplasm during Rab10-dependent vesicle fusion. While relatively understudied compared with other Rabs, Rab10 has been reported to localize to and regulate maturing endo/phagosomes and lysosomes in human macrophages and microglia (Cardoso et al., 2010; Lee et al., 2020). Interestingly, Rab10 has already been implicated in multiple

neurodegenerative diseases: (1) *RAB10* gene expression is altered in human AD brains (Ridge et al.,

2017), (2) rare *RAB10* polymorphisms are linked to protection against AD (Ridge et al., 2017; Tavana

et al., 2018), (3) Rab10 depletion is associated with reduced Aβ levels (Udayar et al., 2013; Ridge et

al., 2017), and (4) Rab10 is a substrate of the PD risk factor gene leucine-rich repeat kinase 2 (LRRK2)

(Steger et al., 2016; Seol et al., 2019). Mutant LRRK2-mediated Rab phosphorylation leads to

endocytic defects (Liu et al., 2020; Rivero-Ríos et al., 2020; Streubel-Gallasch et al., 2021), and

LRRK2-modified Rab10 localizes to and promotes secretion from stressed lysosomes (Eguchi et al.,

2018; Kluss et al., 2022). Interestingly, phosphorylated Rab10 (pRab10) levels are elevated in the CNS

of AD and PD patients, and pRab10 has been reported to colocalize with pathological tau (Yan et al.,

2018; Tezuka et al., 2022) suggesting that post-translational modification of Rab10 modifies the

disease state. Whether Rab10's phosphorylation status affects its ability to regulate prion-like activity of

mHTT$_{ex1}$ or other amyloid aggregates warrants further investigation.

In conclusion, our findings demonstrate that axonal mHTT$_{ex1}$ aggregates activate glial

phagocytosis but also impair normal glial responses to acute neural injury. Aggregate engulfment-

induced defects in endolysosomal processing, such as Rab-mediated vesicle membrane fusion, could

facilitate the formation and spread of prion-like aggregate seeds in the brain. These findings point to

central roles for phagocytic glia in regulating pathological aggregate burden in HD and highlight the

importance of exploring therapeutic interventions that target non-neuronal cells.

REFERENCES

Aits, S., Kricker, J., Liu, B., Ellegaard, A.-M., Hämälistö, S., Tvingsholm, S., et al. (2015). Sensitive detection of lysosomal membrane permeabilization by lysosomal galectin puncta assay. *Autophagy* 11, 1408–1424. doi: 10.1080/15548627.2015.1063871

Alphen, B. van, Stewart, S., Iwanaszko, M., Xu, F., Li, K., Rozenfeld, S., et al. (2022). Glial immune-related pathways mediate effects of closed head traumatic brain injury on behavior and lethality in Drosophila. *PLOS Biology* 20, e3001456. doi: 10.1371/journal.pbio.3001456

Aman, Y., Schmauck-Medina, T., Hansen, M., Morimoto, R. I., Simon, A. K., Bjedov, I., et al. (2021). Autophagy in healthy aging and disease. *Nat Aging* 1, 634–650. doi: 10.1038/s43587-021-00098-4

Bae, E.-J., Kim, D.-K., Kim, C., Mante, M., Adame, A., Rockenstein, E., et al. (2018). LRRK2 kinase regulates α-synuclein propagation via RAB35 phosphorylation. *Nat Commun* 9, 3465. doi: 10.1038/s41467-018-05958-z

Bonam, S. R., Wang, F., and Muller, S. (2019). Lysosomes as a therapeutic target. *Nat Rev Drug Discov* 18, 923–948. doi: 10.1038/s41573-019-0036-1

Brand, A. H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes | Development | The Company of Biologists. Available at: https://journals.biologists.com/dev/article/118/2/401/37970/Targeted-gene-expression-as-a-means-of-altering (Accessed September 19, 2022).

Brelstaff, J. H., Mason, M., Katsinelos, T., McEwan, W. A., Ghetti, B., Tolkovsky, A. M., et al. (2021). Microglia become hypofunctional and release metalloproteases and tau seeds when phagocytosing live neurons with P301S tau aggregates. *Science Advances* 7, eabg4980. doi: 10.1126/sciadv.abg4980

Burbidge, K., Rademacher, D. J., Mattick, J., Zack, S., Grillini, A., Bousset, L., et al. (2022). LGALS3 (galectin 3) mediates an unconventional secretion of SNCA/α-synuclein in response to lysosomal membrane damage by the autophagic-lysosomal pathway in human midbrain dopamine neurons. *Autophagy* 18, 1020–1048. doi: 10.1080/15548627.2021.1967615

Byrns, C. N., Saikumar, J., and Bonini, N. M. (2021). Glial AP1 is activated with aging and accelerated

by traumatic brain injury. *Nat Aging* 1, 585–597. doi: 10.1038/s43587-021-00072-0

Cardoso, C. M. P., Jordao, L., and Vieira, O. V. (2010). Rab10 regulates phagosome maturation and its

overexpression rescues Mycobacterium-containing phagosomes maturation. *Traffic* 11, 221–

235. doi: 10.1111/j.1600-0854.2009.01013.x

Casula, M., Iyer, A. M., Spliet, W. G. M., Anink, J. J., Steentjes, K., Sta, M., et al. (2011). Toll-like

receptor signaling in amyotrophic lateral sclerosis spinal cord tissue. *Neuroscience* 179, 233–

243. doi: 10.1016/j.neuroscience.2011.02.001

Chan, C.-C., Scoggin, S., Wang, D., Cherry, S., Dembo, T., Greenberg, B., et al. (2011). Systematic

Discovery of Rab GTPases with Synaptic Functions in Drosophila. *Current Biology* 21, 1704–

1715. doi: 10.1016/j.cub.2011.08.058

Chen, J. J., Nathaniel, D. L., Raghavan, P., Nelson, M., Tian, R., Tse, E., et al. (2019). Compromised

function of the ESCRT pathway promotes endolysosomal escape of tau seeds and propagation

of tau aggregation. *J. Biol. Chem.*, jbc.RA119.009432. doi: 10.1074/jbc.RA119.009432

Chen, S., Berthelier, V., Yang, W., and Wetzel, R. (2001). Polyglutamine aggregation behavior in vitro

supports a recruitment mechanism of cytotoxicity11Edited by F. Cohen. *Journal of Molecular*

Biology 311, 173–182. doi: 10.1006/jmbi.2001.4850

Choi, I., Zhang, Y., Seegobin, S. P., Pruvost, M., Wang, Q., Purtell, K., et al. (2020). Microglia clear

neuron-released α-synuclein via selective autophagy and prevent neurodegeneration. *Nat*

Commun 11, 1386. doi: 10.1038/s41467-020-15119-w

Chung, W.-S., Clarke, L. E., Wang, G. X., Stafford, B. K., Sher, A., Chakraborty, C., et al. (2013).

Astrocytes mediate synapse elimination through MEGF10 and MERTK pathways. *Nature* 504,

394–400. doi: 10.1038/nature12776

Couto, A., Alenius, M., and Dickson, B. J. (2005). Molecular, Anatomical, and Functional Organization

of the Drosophila Olfactory System. *Current Biology* 15, 1535–1547. doi:

10.1016/j.cub.2005.07.034

Daussy, C. F., and Wodrich, H. (2020). "Repair Me if You Can": Membrane Damage, Response, and

Control from the Viral Perspective. *Cells* 9, 2042. doi: 10.3390/cells9092042

Dejanovic, B., Wu, T., Tsai, M.-C., Graykowski, D., Gandham, V. D., Rose, C. M., et al. (2022).

Complement C1q-dependent excitatory and inhibitory synapse elimination by astrocytes and

microglia in Alzheimer's disease mouse models. *Nat Aging* 2, 837–850. doi: 10.1038/s43587-

022-00281-1

Doherty, J., Logan, M. A., Taşdemir, O. E., and Freeman, M. R. (2009). Ensheathing glia function as

phagocytes in the adult Drosophila brain. *J Neurosci* 29, 4768–4781. doi:

10.1523/JNEUROSCI.5951-08.2009

Donnelly, K. M., Coleman, C. M., Fuller, M. L., Reed, V. L., Smerina, D., Tomlinson, D. S., et al. (2022).

Hunting for the cause: Evidence for prion-like mechanisms in Huntington's disease. *Frontiers in*

Neuroscience 16. Available at: https://www.frontiersin.org/articles/10.3389/fnins.2022.946822

(Accessed January 31, 2024).

Donnelly, K. M., DeLorenzo, O. R., Zaya, A. D., Pisano, G. E., Thu, W. M., Luo, L., et al. (2020).

Phagocytic glia are obligatory intermediates in transmission of mutant huntingtin aggregates

across neuronal synapses. *eLife* 9, e58499. doi: 10.7554/eLife.58499

Dunlop, J., Jones, P. C., and Finbow, M. E. (1995). Membrane insertion and assembly of ductin: a

polytopic channel with dual orientations. *EMBO J* 14, 3609–3616.

Dunst, S., Kazimiers, T., von Zadow, F., Jambor, H., Sagner, A., Brankatschk, B., et al. (2015).

Endogenously Tagged Rab Proteins: A Resource to Study Membrane Trafficking in Drosophila.

Dev Cell 33, 351–365. doi: 10.1016/j.devcel.2015.03.022

Duong, L., Radley, H., Lee, B., Dye, D., Pixley, F., Grounds, M., et al. (2021). Macrophage function in

the elderly and impact on injury repair and cancer. *Immunity & Ageing* 18, 4. doi:

10.1186/s12979-021-00215-2

Eapen, V. V., Swarup, S., Hoyer, M. J., Paulo, J. A., and Harper, J. W. (2021). Quantitative proteomics

reveals the selectivity of ubiquitin-binding autophagy receptors in the turnover of damaged

lysosomes by lysophagy. *eLife* 10, e72328. doi: 10.7554/eLife.72328

Eguchi, T., Kuwahara, T., Sakurai, M., Komori, T., Fujimoto, T., Ito, G., et al. (2018). LRRK2 and its

substrate Rab GTPases are sequentially targeted onto stressed lysosomes and maintain their

homeostasis. *Proc Natl Acad Sci U S A* 115, E9115–E9124. doi: 10.1073/pnas.1812196115

Evans, I. R., Rodrigues, F. S. L. M., Armitage, E. L., and Wood, W. (2015). Draper/CED-1 Mediates an

Ancient Damage Response to Control Inflammatory Blood Cell Migration In Vivo. *Curr Biol* 25,

1606–1612. doi: 10.1016/j.cub.2015.04.037

Falcon, B., Noad, J., McMahon, H., Randow, F., and Goedert, M. (2018). Galectin-8–mediated

selective autophagy protects against seeded tau aggregation. *J Biol Chem* 293, 2438–2451.

doi: 10.1074/jbc.M117.809293

Finbow, M. E., Goodwin, S. F., Meagher, L., Lane, N. J., Keen, J., Findlay, J. B. C., et al. (1994).

Evidence that the 16 kDa proteolipid (subunit c) of the vacuolar H+-ATPase and ductin from gap

junctions are the same polypeptide in Drosophila and Manduca: molecular cloning of the

Vha16k gene from Drosophila. *Journal of Cell Science* 107, 1817–1824. doi:

10.1242/jcs.107.7.1817

Flavin, W. P., Bousset, L., Green, Z. C., Chu, Y., Skarpathiotis, S., Chaney, M. J., et al. (2017).

Endocytic vesicle rupture is a conserved mechanism of cellular invasion by amyloid proteins.

Acta Neuropathol 134, 629–653. doi: 10.1007/s00401-017-1722-x

Freeman, D., Cedillos, R., Choyke, S., Lukic, Z., McGuire, K., Marvin, S., et al. (2013). Alpha-Synuclein

Induces Lysosomal Rupture and Cathepsin Dependent Reactive Oxygen Species Following

Endocytosis. *PLOS ONE* 8, e62143. doi: 10.1371/journal.pone.0062143

Fujita, Y., Maeda, T., Sato, C., Sato, M., Hatakeyama, H., Ota, Y., et al. (2020). Engulfment of Toxic

Amyloid β-protein in Neurons and Astrocytes Mediated by MEGF10. *Neuroscience* 443, 1–7.

doi: 10.1016/j.neuroscience.2020.07.016

Ginsberg, S. D., Mufson, E. J., Alldred, M. J., Counts, S. E., Wuu, J., Nixon, R. A., et al. (2011).

Upregulation of select rab GTPases in cholinergic basal forebrain neurons in mild cognitive

impairment and Alzheimer's disease. *J Chem Neuroanat* 42, 102–110. doi:

10.1016/j.jchemneu.2011.05.012

Hall, A. (1990). The Cellular Functions of Small GTP-Binding Proteins. doi: 10.1126/science.2116664

Han, C., Song, Y., Xiao, H., Wang, D., Franc, N. C., Jan, L. Y., et al. (2014). Epidermal cells are the

primary phagocytes in the fragmentation and clearance of degenerating dendrites in Drosophila.

Neuron 81, 544–560. doi: 10.1016/j.neuron.2013.11.021

Heckmann, B. L., Teubner, B. J. W., Tummers, B., Boada-Romero, E., Harris, L., Yang, M., et al.

(2019). LC3-associated endocytosis facilitates β-amyloid clearance and mitigates

neurodegeneration in murine Alzheimer's Disease. *Cell* 178, 536. doi:

10.1016/j.cell.2019.05.056

Herzog, C., Pons Garcia, L., Keatinge, M., Greenald, D., Moritz, C., Peri, F., et al. (2019). Rapid

clearance of cellular debris by microglia limits secondary neuronal cell death after brain injury in

vivo. *Development* 146, dev174698. doi: 10.1242/dev.174698

Homma, Y., Hiragi, S., and Fukuda, M. (2021). Rab family of small GTPases: an updated view on their

regulation and functions. *FEBS J* 288, 36–55. doi: 10.1111/febs.15453

Hong, S., Beja-Glasser, V. F., Nfonoyim, B. M., Frouin, A., Li, S., Ramakrishnan, S., et al. (2016).

Complement and microglia mediate early synapse loss in Alzheimer mouse models. *Science*

352, 712–716. doi: 10.1126/science.aad8373

Hopp, S. C., Lin, Y., Oakley, D., Roe, A. D., DeVos, S. L., Hanlon, D., et al. (2018). The role of

microglia in processing and spreading of bioactive tau seeds in Alzheimer's disease. *J*

Neuroinflammation 15, 269. doi: 10.1186/s12974-018-1309-z

Hutagalung, A. H., and Novick, P. J. (2011). Role of Rab GTPases in Membrane Traffic and Cell

Physiology. *Physiol Rev* 91, 119–149. doi: 10.1152/physrev.00059.2009

Iram, T., Ramirez-Ortiz, Z., Byrne, M. H., Coleman, U. A., Kingery, N. D., Means, T. K., et al. (2016).

Megf10 Is a Receptor for C1Q That Mediates Clearance of Apoptotic Cells by Astrocytes. *J*

Neurosci 36, 5185–5192. doi: 10.1523/JNEUROSCI.3850-15.2016

Jain, N., Lewis, C. A., Ulrich, J. D., and Holtzman, D. M. (2023). Chronic TREM2 activation exacerbates

Aβ-associated tau seeding and spreading. *J Exp Med* 220, e20220654. doi:

10.1084/jem.20220654

Jeong, G. R., Jang, E.-H., Bae, J. R., Jun, S., Kang, H. C., Park, C.-H., et al. (2018). Dysregulated phosphorylation of Rab GTPases by LRRK2 induces neurodegeneration. *Molecular Neurodegeneration* 13, 8. doi: 10.1186/s13024-018-0240-1

Jia, J., Claude-Taupin, A., Gu, Y., Choi, S. W., Peters, R., Bissa, B., et al. (2020). Galectin-3 coordinates a cellular system for lysosomal repair and removal. *Dev Cell* 52, 69-87.e8. doi: 10.1016/j.devcel.2019.10.025

Jiang, P., Gan, M., Yen, S.-H., McLean, P. J., and Dickson, D. W. (2017). Impaired endo-lysosomal membrane integrity accelerates the seeding progression of α-synuclein aggregates. *Sci Rep* 7, 7690. doi: 10.1038/s41598-017-08149-w

Jiang, S., and Bhaskar, K. (2020). Degradation and Transmission of Tau by Autophagic-Endolysosomal Networks and Potential Therapeutic Targets for Tauopathy. *Frontiers in Molecular Neuroscience* 13. Available at: https://www.frontiersin.org/articles/10.3389/fnmol.2020.586731 (Accessed October 31, 2023).

Jucker, M., and Walker, L. C. (2018). Propagation and spread of pathogenic protein assemblies in neurodegenerative diseases. *Nat. Neurosci.* 21, 1341–1349. doi: 10.1038/s41593-018-0238-6

Karran, E., and De Strooper, B. (2022). The amyloid hypothesis in Alzheimer disease: new insights from new therapeutics. *Nat Rev Drug Discov* 21, 306–318. doi: 10.1038/s41573-022-00391-w

Katzenberger, R. J., Loewen, C. A., Wassarman, D. R., Petersen, A. J., Ganetzky, B., and Wassarman, D. A. (2013). A Drosophila model of closed head traumatic brain injury. *Proceedings of the National Academy of Sciences* 110, E4152–E4159. doi: 10.1073/pnas.1316895110

Kiral, F. R., Kohrs, F. E., Jin, E. J., and Hiesinger, P. R. (2018). Rab GTPases and Membrane Trafficking in Neurodegeneration. *Curr Biol* 28, R471–R486. doi: 10.1016/j.cub.2018.02.010

Kluss, J. H., Beilina, A., Williamson, C. D., Lewis, P. A., Cookson, M. R., and Bonet-Ponce, L. (2022). Lysosomal positioning regulates Rab10 phosphorylation at LRRK2+ lysosomes. *Proc Natl Acad Sci U S A* 119, e2205492119. doi: 10.1073/pnas.2205492119

Knowles, T. P. J., Vendruscolo, M., and Dobson, C. M. (2014). The amyloid state and its association with protein misfolding diseases. *Nat Rev Mol Cell Biol* 15, 384–396. doi: 10.1038/nrm3810

Kohrs, F. E., Daumann, I.-M., Pavlovic, B., Jin, E. J., Kiral, F. R., Lin, S.-C., et al. (2021). Systematic

functional analysis of rab GTPases reveals limits of neuronal robustness to environmental

challenges in flies. *eLife* 10, e59594. doi: 10.7554/eLife.59594

Kouli, A., Camacho, M., Allinson, K., and Williams-Gray, C. H. (2020). Neuroinflammation and protein

pathology in Parkinson's disease dementia. *Acta Neuropathologica Communications* 8, 211.

doi: 10.1186/s40478-020-01083-5

Krasemann, S., Madore, C., Cialic, R., Baufeld, C., Calcagno, N., El Fatimy, R., et al. (2017). The

TREM2-APOE Pathway Drives the Transcriptional Phenotype of Dysfunctional Microglia in

Neurodegenerative Diseases. *Immunity* 47, 566-581.e9. doi: 10.1016/j.immuni.2017.08.008

Langemeyer, L., Fröhlich, F., and Ungermann, C. (2018). Rab GTPase Function in Endosome and

Lysosome Biogenesis. *Trends in Cell Biology* 28, 957–970. doi: 10.1016/j.tcb.2018.06.007

Larsson, M. C., Domingos, A. I., Jones, W. D., Chiappe, M. E., Amrein, H., and Vosshall, L. B. (2004).

Or83b Encodes a Broadly Expressed Odorant Receptor Essential for Drosophila Olfaction.

Neuron 43, 703–714. doi: 10.1016/j.neuron.2004.08.019

Lee, H., Flynn, R., Sharma, I., Haberman, E., Carling, P. J., Nicholls, F. J., et al. (2020). LRRK2 Is

Recruited to Phagosomes and Co-recruits RAB8 and RAB10 in Human Pluripotent Stem Cell-

Derived Macrophages. *Stem Cell Reports* 14, 940–955. doi: 10.1016/j.stemcr.2020.04.001

Lee, J.-H., Kim, J.-Y., Noh, S., Lee, H., Lee, S. Y., Mun, J. Y., et al. (2021). Astrocytes phagocytose

adult hippocampal synapses for circuit homeostasis. *Nature* 590, 612–617. doi:

10.1038/s41586-020-03060-3

Lee, J.-H., Yang, D.-S., Goulbourne, C. N., Im, E., Stavrides, P., Pensalfini, A., et al. (2022). Faulty

autolysosome acidification in Alzheimer's disease mouse models induces autophagic build-up of

Aβ in neurons, yielding senile plaques. *Nat Neurosci* 25, 688–701. doi: 10.1038/s41593-022-

01084-8

Leyns, C. E. G., Gratuze, M., Narasimhan, S., Jain, N., Koscal, L. J., Jiang, H., et al. (2019). TREM2

function impedes tau seeding in neuritic plaques. *Nat Neurosci* 22, 1217–1222. doi:

10.1038/s41593-019-0433-0

Liddelow, S. A., Guttenplan, K. A., Clarke, L. E., Bennett, F. C., Bohlen, C. J., Schirmer, L., et al.

(2017). Neurotoxic reactive astrocytes are induced by activated microglia. *Nature* 541, 481–487.

doi: 10.1038/nature21029

Liddelow, S. A., Marsh, S. E., and Stevens, B. (2020). Microglia and Astrocytes in Disease: Dynamic

Duo or Partners in Crime? *Trends Immunol* 41, 820–835. doi: 10.1016/j.it.2020.07.006

Liu, C.-C., Hu, J., Zhao, N., Wang, J., Wang, N., Cirrito, J. R., et al. (2017). Astrocytic LRP1 Mediates

Brain Aβ Clearance and Impacts Amyloid Deposition. *J Neurosci* 37, 4023–4031. doi:

10.1523/JNEUROSCI.3442-16.2017

Liu, Z., Xu, E., Zhao, H. T., Cole, T., and West, A. B. (2020). LRRK2 and Rab10 coordinate

macropinocytosis to mediate immunological responses in phagocytes. *EMBO J* 39, e104862.

doi: 10.15252/embj.2020104862

MacDonald, J. M., Beach, M. G., Porpiglia, E., Sheehan, A. E., Watts, R. J., and Freeman, M. R.

(2006). The Drosophila Cell Corpse Engulfment Receptor Draper Mediates Glial Clearance of

Severed Axons. *Neuron* 50, 869–881. doi: 10.1016/j.neuron.2006.04.028

Magaki, S. D., Williams, C. K., and Vinters, H. V. (2018). Glial function (and dysfunction) in the normal

& ischemic brain. *Neuropharmacology* 134, 218–225. doi: 10.1016/j.neuropharm.2017.11.009

Marischuk, K., Crocker, K. L., Ahern-Djamali, S., and Boekhoff-Falk, G. (2021). Innate immunity

pathways activate cell proliferation after penetrating traumatic brain injury in adult Drosophila.

2021.09.01.458615. doi: 10.1101/2021.09.01.458615

McGuire, S. E., Le, P. T., and Davis, R. L. (2001). The Role of Drosophila Mushroom Body Signaling in

Olfactory Memory. *Science* 293, 1330–1333. doi: 10.1126/science.1062622

McLaughlin, C. N., Perry-Richardson, J. J., Coutinho-Budd, J. C., and Broihier, H. T. (2019). Dying

neurons utilize innate immune signaling to prime glia for phagocytosis during development. *Dev*

Cell 48, 506-522.e6. doi: 10.1016/j.devcel.2018.12.019

Miesenböck, G., De Angelis, D. A., and Rothman, J. E. (1998). Visualizing secretion and synaptic

transmission with pH-sensitive green fluorescent proteins. *Nature* 394, 192–195. doi:

10.1038/28190

Miron, J., Picard, C., Frappier, J., Dea, D., Théroux, L., and Poirier, J. (2018). TLR4 Gene Expression

and Pro-Inflammatory Cytokines in Alzheimer's Disease and in Response to Hippocampal

Deafferentation in Rodents. *Journal of Alzheimer's Disease* 63, 1547–1556. doi: 10.3233/JAD-

171160

Monaco, A., and Fraldi, A. (2020). Protein Aggregation and Dysfunction of Autophagy-Lysosomal

Pathway: A Vicious Cycle in Lysosomal Storage Diseases. *Front Mol Neurosci* 13, 37. doi:

10.3389/fnmol.2020.00037

Neniskyte, U., Neher, J. J., and Brown, G. C. (2011). Neuronal Death Induced by Nanomolar Amyloid β

Is Mediated by Primary Phagocytosis of Neurons by Microglia*. *Journal of Biological Chemistry*

286, 39904–39913. doi: 10.1074/jbc.M111.267583

Ng, E. L., and Tang, B. L. (2008). Rab GTPases and their roles in brain neurons and glia. *Brain*

Research Reviews 58, 236–246. doi: 10.1016/j.brainresrev.2008.04.006

Paolicelli, R. C., Bolasco, G., Pagani, F., Maggi, L., Scianni, M., Panzanelli, P., et al. (2011). Synaptic

pruning by microglia is necessary for normal brain development. *Science* 333, 1456–1458. doi:

10.1126/science.1202529

Parhizkar, S., Arzberger, T., Brendel, M., Kleinberger, G., Deussing, M., Focke, C., et al. (2019). Loss

of TREM2 function increases amyloid seeding but reduces plaque-associated ApoE. *Nat*

Neurosci 22, 191–204. doi: 10.1038/s41593-018-0296-9

Pearce, M. M. P., Spartz, E. J., Hong, W., Luo, L., and Kopito, R. R. (2015). Prion-like transmission of

neuronal huntingtin aggregates to phagocytic glia in the Drosophila brain. *Nat Commun* 6, 6768.

doi: 10.1038/ncomms7768

Podleśny-Drabiniok, A., Marcora, E., and Goate, A. M. (2020). Microglial Phagocytosis: A Disease-

Associated Process Emerging from Alzheimer's Disease Genetics. *Trends Neurosci* 43, 965–

979. doi: 10.1016/j.tins.2020.10.002

Poe, A. R., Wang, B., Sapar, M. L., Ji, H., Li, K., Onabajo, T., et al. (2019). Robust CRISPR/Cas9-

Mediated Tissue-Specific Mutagenesis Reveals Gene Redundancy and Perdurance in

Drosophila. *Genetics* 211, 459–472. doi: 10.1534/genetics.118.301736

Polanco, J. C., and Götz, J. (2022). Exosomal and vesicle-free tau seeds—propagation and

convergence in endolysosomal permeabilization. *The FEBS Journal* 289, 6891–6907. doi:

10.1111/febs.16055

Portela, M., Yang, L., Paul, S., Li, X., Veraksa, A., Parsons, L. M., et al. (2018). Lgl reduces endosomal

vesicle acidification and Notch signaling by promoting the interaction between Vap33 and the V-

ATPase complex. *Sci Signal* 11, eaar1976. doi: 10.1126/scisignal.aar1976

Potter, C. J., and Luo, L. (2011). Using the Q system in Drosophila. *Nat Protoc* 6, 1105–1120. doi:

10.1038/nprot.2011.347

Preisinger, E., Jordan, B. M., Kazantsev, A., and Housman, D. (1999). Evidence for a recruitment and

sequestration mechanism in Huntington's disease. *Philos Trans R Soc Lond B Biol Sci* 354,

1029–1034. doi: 10.1098/rstb.1999.0455

Prince, E., Kroeger, B., Gligorov, D., Wilson, C., Eaton, S., Karch, F., et al. (2019). Rab-mediated

trafficking in the secondary cells of Drosophila male accessory glands and its role in fecundity.

Traffic 20, 137–151. doi: 10.1111/tra.12622

Pulipparacharuvil, S., Akbar, M. A., Ray, S., Sevrioukov, E. A., Haberman, A. S., Rohrer, J., et al.

(2005). Drosophila Vps16A is required for trafficking to lysosomes and biogenesis of pigment

granules. *Journal of Cell Science* 118, 3663–3673. doi: 10.1242/jcs.02502

Purice, M. D., Ray, A., Münzel, E. J., Pope, B. J., Park, D. J., Speese, S. D., et al. (2017). A novel

Drosophila injury model reveals severed axons are cleared through a Draper/MMP-1 signaling

cascade. *eLife* 6, e23611. doi: 10.7554/eLife.23611

Purice, M. D., Speese, S. D., and Logan, M. A. (2016). Delayed glial clearance of degenerating axons

in aged Drosophila is due to reduced PI3K/Draper activity. *Nat Commun* 7. doi:

10.1038/ncomms12871

Raiders, S., Han, T., Scott-Hewitt, N., Kucenas, S., Lew, D., Logan, M. A., et al. (2021). Engulfed by

Glia: Glial Pruning in Development, Function, and Injury across Species. *J. Neurosci.* 41, 823–

833. doi: 10.1523/JNEUROSCI.1660-20.2020

Ray, A., Speese, S. D., and Logan, M. A. (2017). Glial Draper Rescues Aβ Toxicity in a Drosophila

Model of Alzheimer's Disease. *J Neurosci* 37, 11881–11893. doi: 10.1523/JNEUROSCI.0862-

17.2017

Ridge, P. G., Karch, C. M., Hsu, S., Arano, I., Teerlink, C. C., Ebbert, M. T. W., et al. (2017). Linkage,

whole genome sequence, and biological data implicate variants in RAB10 in Alzheimer's

disease resilience. *Genome Med* 9, 100. doi: 10.1186/s13073-017-0486-1

Rivero-Ríos, P., Romo-Lozano, M., Fernández, B., Fdez, E., and Hilfiker, S. (2020). Distinct Roles for

RAB10 and RAB29 in Pathogenic LRRK2-Mediated Endolysosomal Trafficking Alterations.

Cells 9, 1719. doi: 10.3390/cells9071719

Rodrigues, P. V., de Godoy, J. V. P., Bosque, B. P., Amorim Neto, D. P., Tostes, K., Palameta, S., et

al. (2022). Transcellular propagation of fibrillar α-synuclein from enteroendocrine to neuronal

cells requires cell-to-cell contact and is Rab35-dependent. *Sci Rep* 12, 4168. doi:

10.1038/s41598-022-08076-5

Rodriguez, L., Mohamed, N.-V., Desjardins, A., Lippé, R., Fon, E. A., and Leclerc, N. (2017). Rab7A

regulates tau secretion. *J Neurochem* 141, 592–605. doi: 10.1111/jnc.13994

Rong, Y., McPhee, C. K., Deng, S., Huang, L., Chen, L., Liu, M., et al. (2011). Spinster is required for

autophagic lysosome reformation and mTOR reactivation following starvation. *Proc Natl Acad*

Sci U S A 108, 7826–7831. doi: 10.1073/pnas.1013800108

Santra, M., Dill, K. A., and de Graff, A. M. R. (2019). Proteostasis collapse is a driver of cell aging and

death. *Proceedings of the National Academy of Sciences* 116, 22173–22178. doi:

10.1073/pnas.1906592116

Scherzinger, E., Sittler, A., Schweiger, K., Heiser, V., Lurz, R., Hasenbank, R., et al. (1999). Self-

assembly of polyglutamine-containing huntingtin fragments into amyloid-like fibrils: Implications

for Huntington's disease pathology. *Proc Natl Acad Sci U S A* 96, 4604–4609.

Seol, W., Nam, D., and Son, I. (2019). Rab GTPases as Physiological Substrates of LRRK2 Kinase.

Exp Neurobiol 28, 134–145. doi: 10.5607/en.2019.28.2.134

Seto, S., Tsujimura, K., and Koide, Y. (2011). Rab GTPases regulating phagosome maturation are

differentially recruited to mycobacterial phagosomes. *Traffic* 12, 407–420. doi: 10.1111/j.1600-

0854.2011.01165.x

Singh, T. D., Park, S.-Y., Bae, J., Yun, Y., Bae, Y.-C., Park, R.-W., et al. (2010). MEGF10 functions as

a receptor for the uptake of amyloid-β. *FEBS Letters* 584, 3936–3942. doi:

10.1016/j.febslet.2010.08.050

Small, S. A., Simoes-Spassov, S., Mayeux, R., and Petsko, G. A. (2017). Endosomal traffic jams

represent a pathogenic hub and therapeutic target in Alzheimer's disease. *Trends Neurosci* 40,

592–602. doi: 10.1016/j.tins.2017.08.003

Steger, M., Tonelli, F., Ito, G., Davies, P., Trost, M., Vetter, M., et al. (2016). Phosphoproteomics

reveals that Parkinson's disease kinase LRRK2 regulates a subset of Rab GTPases. *Elife* 5,

e12813. doi: 10.7554/eLife.12813

Stein, K. C., Morales-Polanco, F., van der Lienden, J., Rainbolt, T. K., and Frydman, J. (2022). Ageing

exacerbates ribosome pausing to disrupt cotranslational proteostasis. *Nature* 601, 637–642. doi:

10.1038/s41586-021-04295-4

Streubel-Gallasch, L., Giusti, V., Sandre, M., Tessari, I., Plotegher, N., Giusto, E., et al. (2021).

Parkinson's Disease–Associated LRRK2 Interferes with Astrocyte-Mediated Alpha-Synuclein

Clearance. *Mol Neurobiol* 58, 3119–3140. doi: 10.1007/s12035-021-02327-8

Swanson, L. C., Rimkus, S. A., Ganetzky, B., and Wassarman, D. A. (2020a). Loss of the Antimicrobial

Peptide Metchnikowin Protects Against Traumatic Brain Injury Outcomes in Drosophila

melanogaster. *G3 (Bethesda)* 10, 3109–3119. doi: 10.1534/g3.120.401377

Swanson, L. C., Trujillo, E. A., Thiede, G. H., Katzenberger, R. J., Shishkova, E., Coon, J. J., et al.

(2020b). Survival Following Traumatic Brain Injury in Drosophila Is Increased by Heterozygosity

for a Mutation of the NF-κB Innate Immune Response Transcription Factor Relish. *Genetics*

216, 1117–1136. doi: 10.1534/genetics.120.303776

Tavana, J. P., Rosene, M., Jensen, N. O., Ridge, P. G., Kauwe, J. S., and Karch, C. M. (2018). RAB10: an Alzheimer's disease resilience locus and potential drug target. *Clin Interv Aging* 14, 73–79. doi: 10.2147/CIA.S159148

Tezuka, T., Taniguchi, D., Sano, M., Shimada, T., Oji, Y., Tsunemi, T., et al. (2022). Pathophysiological evaluation of the LRRK2 G2385R risk variant for Parkinson's disease. *npj Parkinsons Dis.* 8, 1–7. doi: 10.1038/s41531-022-00367-y

Udayar, V., Buggia-Prévot, V., Guerreiro, R. L., Siegel, G., Rambabu, N., Soohoo, A. L., et al. (2013). A Paired RNAi and RabGAP Overexpression Screen Identifies Rab11 as a Regulator of β-Amyloid Production. *Cell Rep* 5, 1536–1551. doi: 10.1016/j.celrep.2013.12.005

Udayar, V., Chen, Y., Sidransky, E., and Jagasia, R. (2022). Lysosomal dysfunction in neurodegeneration: emerging concepts and methods. *Trends in Neurosciences* 45, 184–199. doi: 10.1016/j.tins.2021.12.004

Ugbode, C., Fort-Aznar, L., and Sweeney, S. T. (2019). Leaky endosomes push tau over the seed limit. *Journal of Biological Chemistry* 294, 18967–18968. doi: 10.1074/jbc.H119.011687

Vieira, O. V., Botelho, R. J., and Grinstein, S. (2002). Phagosome maturation: aging gracefully. *Biochem J* 366, 689–704. doi: 10.1042/BJ20020691

Wakida, N. M., Cruz, G. M. S., Ro, C. C., Moncada, E. G., Khatibzadeh, N., Flanagan, L. A., et al. (2018). Phagocytic response of astrocytes to damaged neighboring cells. *PLOS ONE* 13, e0196153. doi: 10.1371/journal.pone.0196153

Wang, C., Fan, L., Khawaja, R. R., Liu, B., Zhan, L., Kodama, L., et al. (2022). Microglial NF-κB drives tau spreading and toxicity in a mouse model of tauopathy. *Nat Commun* 13, 1969. doi: 10.1038/s41467-022-29552-6

Wang, Y., Arnold, M. L., Smart, A. J., Wang, G., Androwski, R. J., Morera, A., et al. (2023). Large vesicle extrusions from C. elegans neurons are consumed and stimulated by glial-like phagocytosis activity of the neighboring cell. *eLife* 12, e82227. doi: 10.7554/eLife.82227

Wanker, E. E. (2000). Protein Aggregation and Pathogenesis of Huntingtons Disease: Mechanisms and Correlations. 381, 937–942. doi: 10.1515/BC.2000.114

Welikovitch, L. A., Dujardin, S., Dunn, A. R., Fernandes, A. R., Khasnavis, A., Chibnik, L. B., et al.

(2023). Rate of tau propagation is a heritable disease trait in genetically diverse mouse strains.

iScience 26, 105983. doi: 10.1016/j.isci.2023.105983

Wodrich, A. P. K., Scott, A. W., Shukla, A. K., Harris, B. T., and Giniger, E. (2022). The Unfolded

Protein Responses in Health, Aging, and Neurodegeneration: Recent Advances and Future

Considerations. *Front Mol Neurosci* 15, 831116. doi: 10.3389/fnmol.2022.831116

Yan, T., Wang, L., Gao, J., Siedlak, S. L., Huntley, M. L., Termsarasab, P., et al. (2018). Rab10

Phosphorylation is a Prominent Pathological Feature in Alzheimer's Disease. *J Alzheimers Dis*

63, 157–165. doi: 10.3233/JAD-180023

Yan, Y., Denef, N., and Schüpbach, T. (2009). The vacuolar proton pump (V-ATPase) is required for

Notch signaling and endosomal trafficking in Drosophila. *Dev Cell* 17, 387–402. doi:

10.1016/j.devcel.2009.07.001

Yuste-Checa, P., Trinkaus, V. A., Riera-Tur, I., Imamoglu, R., Schaller, T. F., Wang, H., et al. (2021).

The extracellular chaperone Clusterin enhances Tau aggregate seeding in a cellular model. *Nat*

Commun 12, 4863. doi: 10.1038/s41467-021-25060-1

Zhang, J., Schulze, K. L., Hiesinger, P. R., Suyama, K., Wang, S., Fish, M., et al. (2007). Thirty-One

Flavors of Drosophila Rab Proteins. *Genetics* 176, 1307–1322. doi:

10.1534/genetics.106.066761

Zheng, B., and Tuszynski, M. H. (2023). Regulation of axonal regeneration after mammalian spinal cord

injury. *Nat Rev Mol Cell Biol* 24, 396–413. doi: 10.1038/s41580-022-00562-y

 FIGURE LEGENDS

Figure 1. mHTT$_{ex1}$ expression in ORNs impairs clearance of injured axons. (**A-B**) Maximum

intensity projections of mCD8-GFP-labeled DA1 ORN axons expressing (**A**) HTT$_{ex1}$Q25- (wtHTT$_{ex1}$) or

(**B**) HTT$_{ex1}$Q91-mCherry (mHTT$_{ex1}$) in 7 day-old uninjured flies (*left*) or 14 day-old flies subjected to

bilateral antennal nerve axotomy 7 days earlier (*right*). Scale bars = 5 μm. (**C-D**) Quantification of (**C**)

mCD8-GFP+ and (**D**) HTT$_{ex1}$-mCherry+ DA1 ORN axons remaining in 7, 14, and 28 day-old uninjured

flies or flies at 1, 3, and 5 days post-injury. (**E**) Quantification of mCD8-GFP+ DA1 ORN axons in 7, 14,

and 28 day-old flies expressing HTT$_{ex1}$Q25- or HTT$_{ex1}$Q91-mCherry in DA1 ORNs. All quantified data

were normalized to uninjured 1 day-old adults and graphed as mean ± s.e.m.; *$p<0.05$, **$p<0.01$,

***$p<0.001$, ****$p<0.0001$ by unpaired two-tailed *t*-test.

Figure 2. mHTT$_{ex1}$ expression in glia is associated with reduced ORN axon clearance post-injury.

(**A-B**) Maximum intensity projections of antennal lobes from 5-6 day-old flies expressing HTT$_{ex1}$Q25-

(*top*) or HTT$_{ex1}$Q91-V5 (*bottom*) in glia and immunostained with anti-V5. Scale bars = 10 μm. (**B-C**)

Maximum intensity projections of mCD8-GFP-labeled DA1 ORN axons in 1 day-old flies expressing (**B**)

HTT$_{ex1}$Q25- or (**C**) HTT$_{ex1}$Q91-V5 in repo+ glia. Scale bars = 5 μm. (**D**) Quantification of mCD8-GFP+

DA1 ORN axons in flies expressing HTT$_{ex1}$Q25- or HTT$_{ex1}$Q91-V5 in repo+ glia, either uninjured or at 1

and 3 days post-injury. (**E**) Quantification of mCD8-GFP+ DA1 ORN axons in 1 day-old flies expressing

HTT$_{ex1}$Q25- or HTT$_{ex1}$Q91-mCherry in repo+ glia. All data were normalized to uninjured 1 day-old adult

flies and graphed as mean ± s.e.m.; ***$p<0.001$ by unpaired two-tailed *t*-test.

Figure 3. mHTT$_{ex1}$ expression inhibits engulfment of injured ORN axons. (**A-B**) Maximum intensity

projections of VA1lm ORN axons co-expressing the ratiometric phagocytic indicator, MApHS, and (**A**)

HTT$_{ex1}$Q25- or (**B**) HTT$_{ex1}$Q91-V5 from 7 day-old uninjured flies (*left*) and flies 25 hours post-injury

(*right*). Scale bars = 10μm. (**C**) pHluorin:tdTomato fluorescence intensity ratios calculated in VA1lm

glomeruli from 7-day-old uninjured flies and flies at 14 or 25 hours post-injury. Data were normalized to

the uninjured condition for each genotype. (**D**) pHluorin:tdTomato fluorescence intensity ratios in 7 day-

old flies co-expressing MApHS with HTT$_{ex1}$Q25- or HTT$_{ex1}$Q91-V5 in VA1lm ORNs, normalized to wtHTT$_{ex1}$ controls. Data are shown as mean ± s.e.m.; **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by unpaired two-tailed t-test. (**E-F**) Maximum intensity projections of the central brain from 6-7 day-old flies expressing (**E**) repo-Cas9 and (**F**) repo-Cas9 plus gRNAs targeting *draper* ("Draper KO"). Brains were immunostained with anti-Draper. Scale bars = 10 μm. (**G**) Quantification of Draper immunofluorescence in brains from flies shown in (**E-F**), normalized to control.

Figure 4. Neuronal mHTT$_{ex1}$ accumulates in acidic cellular compartments. (A-B, D-E) Maximum intensity projections of (**A-B**) Or83b+ ORN axons or (**D-E**) OK107+ MBN soma expressing (**A,D**) HTT$_{ex1}$Q25- or (**B,E**) HTT$_{ex1}$Q91-associated pH sensor (HApHS) from 13-14 day-old flies. Scale bars = 10 μm. (**C,F**) pHluorin:tdTomato fluorescence intensity ratios of data shown in (**A-B** and **D-E**), normalized to wtHTT$_{ex1}$ controls. (**G-H**) Maximum intensity projections of VA1lm ORN axons co-expressing mHApHS and (**G**) repo-Cas9 or (**H**) repo-Cas9 and gRNAs targeting *draper*. Scale bars = 5 μm. (**I**) pHluorin:tdTomato fluorescence intensity ratios calculated in VA1lm glomeruli from 9-10 day-old flies as shown in (**G-H**). Data were normalized to flies not expressing gRNAs. All quantified data are shown as mean ± s.e.m.; **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by unpaired two-tailed t-test.

Figure 5. mHTT$_{ex1}$ expression in ORNs upregulates phagocytic and innate immunity genes and impairs injury-induced transcriptional responses. (**A**) Diagrams of Toll-6 (purple) and Draper (blue) signaling pathways. (**B**) qPCR analysis of the indicated genes in 8-11 day-old flies expressing GFP, HTT$_{ex1}$Q25-, or HTT$_{ex1}$Q91-GFP in Or83b+ ORNs. RNA was isolated from heads of uninjured flies or flies 3 hours after bilateral antennal and maxillary palp nerve injury. Data are shown as mean ± s.e.m and normalized to the housekeeping gene *rpl32*. *$p<0.05$, **$p<0.01$, ***$p<0.001$ by one-way ANOVA; asterisks and hashtags indicate statistical significance comparing -/+ injury or genotypes, respectively. (**C-D and F-G**) Maximum intensity projections of Or83b+ ORN axons from 14-15 day-old flies expressing (**C,F**) HTT$_{ex1}$Q25- or (**D,G**) HTT$_{ex1}$Q91-V5 in (**C-D**) Toll-6^{MIMICGFP} (Toll-6-GFP) or (**F-G**) Jra-GFP genetic backgrounds. Brains were immunostained with GFP, V5, and N-Cadherin (**C-D**) or GFP,

1017 Repo, and N-Cadherin (**F-G**) antibodies. In panel (**C**), diffuse wtHTT$_{ex1}$ signal was adjusted post-

1018 acquisition for increased visibility. Scale bars = 10 µm. (**E,H**) Quantification of (**E**) Toll-6-GFP or (**H**) Jra-

1019 GFP expression from flies show in (**C-D** and **F-G**). Data are graphed as mean ± s.e.m.; *p<0.05,

1020 ****p<0.0001 by unpaired two-tailed t-test.

1021

Figure 6. mHTT$_{ex1}$ expression in ORNs increases expression and impairs injury-induced

upregulation of some AMP genes. (**A-E**) qPCR analysis of the indicated AMP genes in 8-11 day-old

flies expressing HTT$_{ex1}$Q25- or HTT$_{ex1}$Q91-GFP in Or83b+ ORNs. RNA was isolated from heads of

uninjured flies or flies 3 hours after bilateral antennal and maxillary palp nerve injury. Data are shown

as mean ± s.e.m and normalized to the housekeeping gene *rpl32*. **p<0.01, ***p<0.001 by unpaired

two-tailed t-test; asterisks and hashtags indicate statistical significance comparing -/+ injury or

genotypes, respectively.

1029

Figure 7. A subset of mHTT$_{ex1}$ aggregates are closely associated with Draper+ glial membranes.

(**A-B**) Maximum intensity projections of Or83b+ ORN axons from 7-8 day-old flies expressing (**A**)

HTT$_{ex1}$Q25- or (**B**) HTT$_{ex1}$Q91-mCherry and immunostained for Draper. Scale bars = 10 µm. (**C**)

Quantification of Draper immunofluorescence from flies show in (**A-B**). Data are normalized to control

and graphed as mean ± s.e.m.; *p<0.05 by unpaired two-tailed *t*-test; n=12. (**D**) Single 0.35 µm

confocal slice showing magnified HTT$_{ex1}$Q91-mCherry aggregate and associated Draper signal. Scale

bar = 0.5 µm. (**E-F**) High-magnification confocal stacks of Draper signal within ≤0.2 µm of either (**E**)

HTT$_{ex1}$Q25- or (**F1-3**) HTT$_{ex1}$Q91-mCherry surfaces. Raw data are shown to the left of segmented

surfaces generated from each fluorescence signal. In (**E**), diffuse wtHTT$_{ex1}$ signal was adjusted post-

acquisition for increased visibility. Scale bars = 1 µm.

1040

Figure 8. Seeded aggregation of wtHTT$_{ex1}$ protein expressed in ensheathing glia by neuronal

mHTT$_{ex1}$ aggregates. (**A-B**) Maximum intensity projections of DA1 glomeruli from 4-5 day-old flies

expressing HTT$_{ex1}$Q91-mCherry in DA1 ORNs and HTT$_{ex1}$Q25-GFP in (**A**) mz0709+ ensheathing glia or

(**B**) Alrm+ astrocytic glia. Scale bars = 5 µm. (**C-D**) Quantification of (**C**) HTT$_{ex1}$Q91-mCherry ("mHTT")

and (**D**) seeded HTT$_{ex1}$Q25-GFP ("mHTT+wtHTT") aggregates from flies shown in (**A-B**). Data are

shown as mean ± s.e.m.; ***$p<0.001$ by unpaired two-tailed t-test.

1047

Figure 9. Glial phagocytic and innate immunity genes regulate numbers of mHTT$_{ex1}$ aggregates

in ORN axons. (A) Maximum intensity projections of DA1 glomeruli from 7 day-old flies expressing

HTT$_{ex1}$Q91-mCherry in DA1 ORNs and siRNAs targeting the indicated genes in ensheathing glia. Scale

bars = 5 µm. (**B**) Quantification of HTT$_{ex1}$Q91-mCherry aggregates detected in DA1 glomeruli from flies

shown in (**A**). (**C**) Maximum intensity projections of DA1 glomeruli from 7 day-old flies expressing

HTT$_{ex1}$Q91-mCherry in DA1 ORNs and siRNAs targeting mCherry or Ets21c in repo+ glia in the

presence of tubP-Gal80^{ts}. Adult flies were raised at the permissive (18°C, *top*) or restrictive (29°C,

bottom) temperatures to regulate siRNA expression in adults. Scale bars = 5 µm. (**D**) Quantification of

HTT$_{ex1}$Q91-mCherry aggregates detected in DA1 glomeruli from flies shown in (**C**). (**E**) Mean volumes

for HTT$_{ex1}$Q91-mCherry aggregates detected in DA1 glomeruli from flies shown in (**A and C**). All

graphed data are shown as mean ± s.e.m.; *$p<0.05$, **$p<0.01$, ***$p<0.001$ by one-way ANOVA or

unpaired two-tail t-test compared to no RNAi or mCherry^{RNAi} controls.

1060

Figure 10. Neuronal mHTT$_{ex1}$ expression increases numbers of acidified and active lysosomes in

adult brains. (A-B) Maximum intensity projections of antennal lobes from 9-10 day-old adult flies

expressing (**A**) HTT$_{ex1}$Q25- or (**B**) HTT$_{ex1}$Q91-GFP in Or83b+ ORNs and stained with Magic Red (MR)

to label active cathepsins. Scale bars = 10 µm. (**C**) Quantification of MR+ surfaces (*left*) and HTT$_{ex1}$-

associated MR+ surfaces (*right*) from flies shown in (**A** and **B**); HTT$_{ex1}$-associated vesicles were defined

by filtering for MR+ surfaces ≤0.2 µm from HTT$_{ex1}$ fluorescent signal in confocal stacks. **(D-E)** Maximum

intensity projections of antennal lobes from 15 day-old flies expressing (**D**) HTT$_{ex1}$Q25- or (**E**)

HTT$_{ex1}$Q91-GFP in Or83b+ ORNs and stained with Lysotracker Red (LTR) to label low pH

compartments. Scale bars = 10 µm. Diffuse wtHTT$_{ex1}$ signal was adjusted post-acquisition for increased

visibility in panels (**A** and **D**). (**F**) Quantification of LTR+ surfaces (*left*) and HTT$_{ex1}$-associated LTR+

surfaces (*right*) from flies shown in (**D** and **E**); HTT$_{ex1}$-associated vesicles were defined by filtering for

LTR+ surfaces ≤0.2 µm from HTT$_{ex1}$ fluorescent signal in confocal stacks. Data are shown as mean ±

s.e.m.; ****$p<0.0001$ by unpaired two-tailed *t*-test. (**G-H**) High-magnification images of boxes indicated

in (**B and E**) showing co-localization of MR (**G**) or LTR (**H**) with mHTT$_{ex1}$ fluorescent signals. Scale bars

= 1 µm.

1076

Figure 11. LAMP1+ vesicle accumulation in fly brains expressing mHTT$_{ex1}$ in ORNs. (A-B)

Confocal stacks showing antennal lobes from 19-22 day-old flies expressing (**A**) HTT$_{ex1}$Q25- or (**B**)

HTT$_{ex1}$Q91-mCherry in Or83b+ ORNs and LAMP1 tagged at its cytoplasmic C-terminus with GFP

(LAMP1-GFP) in glia. Brains were immunostained with anti-GFP to amplify LAMP1-GFP signal.

LAMP1-GFP+ or HTT$_{ex1}$+ segmented surfaces are shown to the right of each set of raw fluorescence

images. Insets show magnified regions of interest from each image. Scale bars = 10 µm. (**C**) High-

magnification confocal stack showing a LAMP1-GFP+ surface within 0.2 µm of two HTT$_{ex1}$Q91-

mCherry+ aggregates. Scale bar = 1 µm. (**D-E**) Confocal stacks showing antennal lobes from 21-22

day-old flies expressing (**D**) HTT$_{ex1}$Q25- or (**E**) HTT$_{ex1}$Q91-mCherry in Or83b+ ORNs and LAMP1

tagged at its luminal N-terminus with GFP (GFP-LAMP1) in glia. GFP-LAMP1+ or HTT$_{ex1}$+ segmented

surfaces are shown to the right of each set of raw fluorescence images. Insets show a single GFP-

LAMP1+ surface of interest from each image. Scale bars = 10 µm. Diffuse wtHTT$_{ex1}$ signal was

adjusted post-acquisition for increased visibility in panels (**A** and **D**). (**F**) High-magnification confocal

stack showing a GFP-LAMP1+ surface within 0.2 µm of a HTT$_{ex1}$Q91-mCherry+ aggregate. Scale bar =

1µm. (**G-I**) Quantification of total LAMP1-GFP+ or GFP-LAMP1+ surfaces (**G**), LAMP1-GFP+ or GFP-

LAMP1+ surfaces ≤0.2 µm from HTT$_{ex1}$ surfaces (**H**), and mean LAMP1-GFP+ or GFP-LAMP1+

surface volume (**I**) in brains expressing HTT$_{ex1}$Q25- or HTT$_{ex1}$Q91-mCherry. The dark red bars in (**I**)

represent LAMP1+ surfaces that co-localized with mHTT$_{ex1}$. All graphed data are shown as mean ±

s.e.m.; *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by unpaired two-tailed *t*-test.

1096

Figure 12. Increased association of Galectins-3 and -8 expressed in glia with neuronal mHTT$_{ex1}$

aggregates. (A-D) Confocal stacks showing antennal lobes from 16-18 day-old flies expressing

HTT$_{ex1}$Q25- (**A and C**) or HTT$_{ex1}$Q91-GFP (**B and D**) in Or83b+ ORNs together with Galectin-3 (**A-B**) or

Galectin-8 (**C-D**) tagged with mCherry in glia. Segmented Galectin+ or HTT$_{ex1}$+ surfaces are shown to

the right of each set of raw fluorescence images. Insets show Galectin+ surfaces of interest from each

image. Scale bars = 10 µm. (**E-G**) Quantification of total Galectin+ surfaces (**E**), Galectin+ surfaces

≤0.2 µm from HTT$_{ex1}$ surfaces (**F**), and mean Galectin+ surface volume (**G**) in brains expressing

HTT$_{ex1}$Q25- or HTT$_{ex1}$Q91-mCherry. The dark red bars in (**G**) represent Galectin+ surfaces that co-

localized with mHTT$_{ex1}$. All graphed data are shown as mean ± s.e.m.; *$p<0.05$, **$p<0.01$, ***$p<0.001$,

****$p<0.0001$ by unpaired two-tailed t-test.

1107

Figure 13. Knockdown of genes regulating lysosome acidification alters seeded aggregation of

glial wtHTT$_{ex1}$ protein by neuronal mHTT$_{ex1}$ aggregates. (**A**) Confocal stacks showing DA1 glomeruli

from 8-9 day-old flies expressing HTT$_{ex1}$Q91-mCherry in DA1 ORNs and HTT$_{ex1}$Q25-GFP plus siRNAs

targeting the indicated genes in repo+ glia. Negative controls expressed siRNAs targeting mCherry.

Scale bars = 5 µm. (**B-C**) Quantification of (**B**) HTT$_{ex1}$Q91-mCherry or (**C**) seeded HTT$_{ex1}$Q25-GFP

aggregates from brains shown in (**A**). Data are shown as mean ± s.e.m.; *$p<0.05$, ***$p<0.005$ by

unpaired two-tailed t-test.

1115

Figure 14. Rab10 is required for seeded aggregation of glial wtHTT$_{ex1}$ by neuronal mHTT$_{ex1}$

aggregates. (**A**) Confocal stacks of DA1 glomeruli from 7 day-old flies expressing HTT$_{ex1}$Q91-mCherry

in DA1 ORNs and HTT$_{ex1}$Q25-GFP plus siRNAs targeting firefly luciferase (FFLuc), Draper, or Rab10 in

repo+ glia. Surfaces representing mHTT$_{ex1}$ aggregates (*red*) and seeded wtHTT$_{ex1}$ aggregates are

superimposed on the raw data. Scale bars = 5µm. (**B-C**) mHTT$_{ex1}$ (**B**) or mHTT$_{ex1}$+wtHTT$_{ex1}$ (**C**)

aggregates quantified from flies expressing HTT$_{ex1}$Q91-mCherry in DA1 ORNs and HTT$_{ex1}$Q25-GFP

plus siRNAs targeting Draper-I (Drpr) or 23 different Rab GTPases in repo+ glia. Negative controls

expressed no siRNAs or siRNAs targeting FFLuc or mCherry (*black bars*). (**D**) Normalized Draper

immunofluorescence in the central brain of 4-5 day-old wild-type ($rab10^{+/+}$) and $rab10$ null ($rab10^{-/-}$)

flies. (**E**) Quantification of seeded wtHTT$_{ex1}$ aggregates in flies expressing HTT$_{ex1}$Q91-mCherry in DA1

ORNs and HTT$_{ex1}$Q25-GFP plus siRNAs targeting Rab10, without or with Draper-I cDNAs in repo+glia.

(**F-G**) Quantification of mHTT$_{ex1}$ (**F**) or mHTT$_{ex1}$+wtHTT$_{ex1}$ (**G**) aggregates from 10 day-old flies

expressing HTT$_{ex1}$Q91-mCherry in DA1 ORNs and HTT$_{ex1}$Q25-GFP in PNs, either heterozygous or

trans-heterozygous for *draper* (*drpr/+*), *rab10* (*rab10/+*), or *rab14* (*rab14/+*) mutant alleles. All graphed

data are shown as mean ± s.e.m.; *$p<0.05$, **$p<0.01$, ***$p<0.005$ by one-way ANOVA or unpaired two-

tailed *t*-test comparing to controls.

Figure 15. Association of neuronal mHTT$_{ex1}$ aggregates with Rab GTPases that label early,

maturing, and late phagosomes. (A) qPCR analysis of *rab10*, *rab5*, and *rab7* expression in 8-11 day-

old flies expressing GFP, HTT$_{ex1}$Q25-, or HTT$_{ex1}$Q91-GFP in Or83b+ ORNs. RNA was isolated from

heads of uninjured flies or flies 3 hours after bilateral antennal and maxillary palp nerve injury. Data are

shown as mean ± s.e.m and normalized to the housekeeping gene *rpl32*. *$p<0.05$, **$p<0.01$, ***$p<0.001$

by one-way ANOVA, asterisks and hashtags indicate statistical significance comparing -/+ injury or

genotypes, respectively. (**B-G**) Confocal stacks of the antennal lobe from 7 day-old flies expressing (**B,**

D, & F) HTT$_{ex1}$Q25- or (**C, E, & G**) HTT$_{ex1}$Q91-V5 in Or83b+ ORNs and endogenously-tagged (**B-C**)

YRab10, (**D-E**) YRab5, or (**F-G**) YRab7 in all cells. Brains were immunostained using YFP (*green*), V5

(*magenta*), and N-Cadherin (*blue*) antibodies. Segmented YRab+ or HTT+ surfaces are shown to the

right of each set of raw fluorescent images. Insets show magnified YFP+ surfaces of interest from each

image. Diffuse wtHTT$_{ex1}$ signal was adjusted post-acquisition for increased visibility in panels (**B, D,** and

F). Scale bars = 10 µm. (**H-I**) Quantification of YRab+ surfaces (**H**) and YRab+ surfaces within 0.2µm of

HTT$_{ex1}$+ surfaces (**I**). Data are shown as mean ± s.e.m.; *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by

unpaired two-tailed *t*-test.

Figure 16. Association of neuronal mHTT aggregates with glial Rab GTPases that label early,

maturing, and late phagosomes. (A-F) Confocal stacks of the antennal lobe from 16-19 day-old flies

expressing (**A, C, E**) HTT$_{ex1}$Q25- or (**B, D, F**) HTT$_{ex1}$Q91-V5 in Or83b+ ORNs together with (**A-B**) YFP-

Rab10, (**C-D**) YFP-Rab5, or (**E-F**) YFP-Rab7 in repo+ glia. Brains were immunostained with anti-GFP

to amplify YFP-Rab signals. Segmented Rab+ or HTT$_{ex1}$+ surfaces are shown to the right of each set of

raw fluorescence images. Insets show magnified YFP-Rab+ surfaces of interest from each image.

Diffuse wtHTT$_{ex1}$ signal was adjusted post-acquisition for increased visibility in panels (**A, C,** and **E**).

Scale bars = 10 µm. (**G-I**) Quantification of total YFP-Rab+ surfaces (**G**), YFP-Rab+ surfaces ≤0.2 µm

from HTT$_{ex1}$ surfaces (**H**), and mean YFP-Rab+ surface volume (**I**) in brains expressing HTT$_{ex1}$Q25- or

HTT$_{ex1}$Q91-mCherry. The dark red bars in (**I**) represent YFP-Rab+ surfaces that co-localized with

mHTT$_{ex1}$. All graphed data are shown as mean ± s.e.m.; *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$

by unpaired two-tailed t-test.

1161

1162 **Figure 17. Glial YFP-Rab+ surfaces are closely associated with neuronal mHTT$_{ex1}$ aggregates**.

1163 High magnification confocal stacks showing examples of individual YFP+ surfaces within 0.2 µm of

1164 HTT$_{ex1}$Q91 aggregates from 21-22 day-old adult brains expressing HTT$_{ex1}$Q91-mCherry in Or83b+

1165 ORNs and YFP-Rab10 (**A1-3**), YFP-Rab5 (**B1-2**), or YFP-Rab7 (**C1-2**) in repo+ glia. Brains were

1166 immunostained with anti-GFP to amplify YFP-Rab signals. Scale bars = 1 µm.

1167

Figure 18. "Traffic jam" model illustrating effects of neuronal mHTT$_{ex1}$ aggregates on phagocytic

glia. mHTT$_{ex1}$ aggregates (*magenta circles*) generated in axons cause phagolysosomal defects in

neighboring glial cells. Neuronal mHTT$_{ex1}$ aggregates activate glial Draper and Toll-6 signaling

pathways, but impair normal phagocytic responses to injury, including reduced nascent phagosome

formation and decreased numbers of Rab5+ early phagosomes (*red arrow*). A buildup of engulfed

mHTT$_{ex1}$ aggregates in glia leads to accumulation of maturing (Rab10+ or Rab7+) phagosomes and

lysosomes (LAMP1+) (*green arrows*), possibly further slowing Draper-dependent engulfment and early

phagosome formation. Defective phagocytic clearance could enhance leak or ejection of some mHTT$_{ex1}$

aggregates from phagolysosomes to increase their toxicity and capacity to seed soluble HTT proteins

(*magenta + green circles*).

TABLE LEGENDS

Table 1. *Drosophila* **genotype and source information.**

1180

1181   **Table 2. Genotypes of flies used in all figures.**

1182

1183   **Table 3. Primer sequences used for qPCR analyses.**

1184

1185   **Table 4. Statistical information for all quantitative results.**

MApHS: pHluorin | CD4 TM | tdTomato

A wtHTT — Control | Injured   **B** mHTT — Control | Injured

pHluorin

tdTomato

Merge

C
wtHTT
mHTT
Draper KO

pHluorin:tdTomato Fluorescence

**
****
****
****
****
***

Time post-injury (hr)

D
pHluorin:tdTomato Fluorescence

***

wtHTT   mHTT

E Control   **F** Draper KO

Draper

G
Normalized Fluorescence Intensity

****

Control   Draper KO

HApHS: pHluorin HTTex1 tdTomato

ORN Axons   MBN Soma

A wtHTT   B mHTT   D wtHTT   E mHTT

pHluorin

tdTomato

Merge

C

wtHTT
mHTT

pHluorin:tdTomato Fluorescence

***

F

pHluorin:tdTomato Fluorescence

****

G Control   pHluorin   tdTomato   Merge

H Draper KO

I

pHluorin:tdTomato Fluorescence

**

mHApHS   +   +
Draper   +   −

A

FFLuc^RNAi

mHTT | wtHTT | Merge

Draper-I^RNAi

Rab10^RNAi

B

mHTT Aggregates

***   *

C

mHTT+wtHTT Aggregates

***   ***   ***

RNAi: - FFLuc mCh Drpr

2-1 2-2 3 4 5 6-1 6-2 7 8-1 8-2 9-1 9-2 9Db 10 14 18-1 18-2 19 21 23-1 23-2 23-3 23-4 27 32-1 32-2 35-1 35-2 39 40 X2 X4 X6-1 X6-2

Rab#

D

Norm. Draper levels

1.0

0.5

**

rab10^{+/+}   rab10^{-/-}

E

mHTT+wtHTT Aggregates

***

Draper-I: - - +
Rab10^RNAi: - + +

F

mHTT Aggregates

G

mHTT+wtHTT Aggregates

*

drpr/+: - + - + - +
rab10/+: - - + + - -
rab14/+: - - - - + +

Table 1. *Drosophila* genotype and source information.

Name	ID (if available)	Reference	Notes
Or67d-QF		Liang et al., 2013	
repo-Gal4	RRID:BDSC_7415	Sepp et al., 2001	
mz0709-Gal4		Ito et al., 1995	a gift from Marc Freeman (Vollum Institute)
alrm-Gal4	RRID:BDSC_67032		
orco-Gal4	RRID:BDSC_23292		
Or47b-Gal4	RRID:BDSC_9983		
QUAS-HTT$_{ex1}$Q25-mCherry^{attP24}		Donnelly et al., 2020	
QUAS-HTT$_{ex1}$Q91-mCherry^{attP24}		Donnelly et al., 2020	
QUAS-HTT$_{ex1}$Q91-mCherry^{attP3}		Donnelly et al., 2020	
QUAS-HTT$_{ex1}$Q91-mCherry$^{suHw(attP8)}$		This study	
UAS-HTT$_{ex1}$Q25-GFP		Donnelly et al., 2020	
UAS-HTT$_{ex1}$Q91-GFP		Donnelly et al., 2020	
UAS-GFP		Donnelly et al., 2020	
QUAS-HTT$_{ex1}$Q25-GFP		This study	
QUAS-HTT$_{ex1}$Q91-GFP		This study	
UAS-mCherry-LGals3		This study	
UAS-mCherry-LGals8		This study	
QUAS-HTT$_{ex1}$Q25-V5		This study	
QUAS-HTT$_{ex1}$Q91-V5		This study	
UAS-pHluorin-HTT$_{ex1}$Q25-tdTomato		This study	
UAS-pHluorin-HTT$_{ex1}$Q91-tdTomato		This study	
UAS-mCD8-GFP	RRID:BDSC_5137		

UAS-Draper^RNAi		Logan et al., 2012	a gift from Marc Freeman (Vollum Institute)
tubP-Gal80^ts	RRID:BDSC_7017		
UAS-Drs^RNAi	RRID:BDSC_55391		
UAS-PGRP-SA^RNAi	RRID:BDSC_60037		
UAS-MMP1^RNAi	RRID:BDSC_31489		
UAS-ets21c^RNAi	RRID:BDSC_39069		
UAS-mCherry^RNAi	RRID:BDSC_35785		
UAS-rel^RNAi	RRID:BDSC_33661, RRID BDSC_28943		
UAS-dl^RNAi	RRID:BDSC_38905, RRID:BDSC_32934, RRID:BDSC_34938, RRID:BDSC_27650, RRID:BDSC_36650		
UAS-NinjurinA^RNAi	RRID:BDSC_50632 RRID:BDSC_51358		
UAS-Toll-6^RNAi	RRID:BDSC_56048, RRID:BDSC_64968		
Orco-QF2	RRID:BDSC_91997		
TI{TI}Rab10^YFP	RRID:BDSC_62548		
TI{TI}Rab5^YFP	RRID:BDSC_62543		
TI{TI}Rab7^YFP	RRID:BDSC_62545		
UAS-MApHS		Han et al., 2014	a gift from Chun Han (Cornell University)
Repo-Cas9[6A]		Koreman et al., 2021	a gift from Chun Han (Cornell University)
gRNA-drpr(BR)		Sapar et al., 2018	a gift from Chun Han (Cornell University)
Jra-GFP.FLAG	RRID:BDSC_50755		
OK107-Gal4			
UAS-GFP-LAMP1		Pulipparacharuvil et al., 2005	A gift from Helmut Kramer (UT Southwestern)

UAS-LAMP1-GFP	RRID:BDSC_42714		*nSyb-Gal4* was removed from stock
UAS-Rbcn-3a^{RNAi}	RRID:BDSC_34612		
UAS-Spinster^{RNAi}	RRID:BDSC_27702		
UAS-Vha100-2^{RNAi}	RRID:BDSC_64859		
UAS-Vha16-1^{RNAi}	RRID:BDSC_40923		
UAS-Vha68-3^{RNAi}	RRID:BDSC_42954		
UAS-YFP-Rab5	RRID:BDSC_9775		
UAS-YFP-Rab7	RRID:BDSC_23641		
UAS-YFP-Rab10	RRID:BDSC_24097		
UAS-YFP-Rab10 (T23N)	RRID:BDSC_9788		
UAS-YFP-Rab10 (Q68L)	RRID:BDSC_23259		
UAS-Rab7^{RNAi}	RRID:BDSC_27051		
UAS-Rab10^{RNAi}	RRID:BDSC_26289		
UAS-Rab2^{RNAi}	RRID:BDSC_28701 RRID:BDSC_34922		
UAS-Rab6^{RNAi}	RRID:BDSC_27490 RRID:BDSC_35744		
UAS-Rab11^{RNAi}	RRID:BDSC_27730		
UAS-Rab3^{RNAi}	RRID:BDSC_34655		
UAS-Rab14^{RNAi}	RRID:BDSC_28708		
UAS-Rab4^{RNAi}	RRID:BDSC_33757		
UAS-Rab5^{RNAi}	RRID:BDSC_30518		
UAS-Rab8^{RNAi}	RRID:BDSC_27519 RRID:BDSC_34373		
UAS-Rab9^{RNAi}	RRID:BDSC_42942 RRID:BDSC_34374		
UAS-luciferase^{RNAi}	RRID:BDSC_31603		
UAS-Rab9Db^{RNAi}	RRID:BDSC_38269		
UAS-Rab32^{RNAi}	RRID:BDSC_38956 RRID:BDSC_28002		

UAS-Rab18[RNAi]	RRID:BDSC_34734 RRID:BDSC_27665		
UAS-RabX4[RNAi]	RRID:BDSC_44070		
UAS-Rab27[RNAi]	RRID:BDSC_50537		
UAS-RabX2[RNAi]	RRID:BDSC_53928		
UAS-Rab39[RNAi]	RRID:BDSC_53995		
UAS-Rab23[RNAi]	RRID:BDSC_55352 RRID:BDSC_63689 RRID:BDSC_36091 RRID:BDSC_28025		
UAS-Rab35[RNAi]	RRID:BDSC_67952 RRID:BDSC_80457		
UAS-Rab40[RNAi]	RRID:BDSC_80472		
UAS-RabX6[RNAi]	RRID:BDSC_26281 RRID:BDSC_53252		
UAS-Rab19[RNAi]	RRID:BDSC_34607		
UAS-Rab21[RNAi]	RRID:BDSC_29403		
UAS-Rab27[RNAi]	RRID:BDSC_35774		
UAS-Draper-I		Logan et al., 2012	a gift from Marc Freeman (Vollum Institute)
Rab10(CRISPR-KO)		Kohrs et al., 2021	a gift from P. Robin Heisinger (Freie Universität Berlin)

Table 2. Genotypes of flies used in all figures.

Figure	Genotype
Figure 1A & C-E	$Or67d\text{-}QF/Y;QUAS\text{-}HTT_{ex1}Q25\text{-}mCherry/QUAS\text{-}mCD8\text{-}GFP$
Figure 1B-E	$Or67d\text{-}QF/Y;QUAS\text{-}HTT_{ex1}Q91\text{-}mCherry/QUAS\text{-}mCD8\text{-}GFP$
Figure 2A,D-E	$Or67d\text{-}QF/Y;UAS\text{-}HTT_{ex1}Q25\text{-}V5/QUAS\text{-}mCD8\text{-}GFP;repo\text{-}gal4/+$
Figure 2B,D-E	$Or67d\text{-}QF/Y;UAS\text{-}HTT_{ex1}Q91\text{-}V5/QUAS\text{-}mCD8\text{-}GFP;repo\text{-}gal4/+$
Figure 2C	$+/Y;UAS\text{-}HTT_{ex1}Q25\text{-}V5/+;repo\text{-}gal4/+$ & $+/Y;UAS\text{-}HTT_{ex1}Q91\text{-}V5/+;repo\text{-}gal4/+$
Figure 3A & C-D	$w1118/Y;UAS\text{-}HTT_{ex1}Q25\text{-}V5/Or47b\text{-}Gal4;UAS\text{-}MapHS/+$
Figure 3B-D	$w1118/Y;UAS\text{-}HTT_{ex1}Q91\text{-}V5/Or47b\text{-}Gal4;UAS\text{-}MapHS/+$;
Figure 3C	$w1118/Y;repo\text{-}Cas9/Or47b\text{-}Gal4;UAS\text{-}MapHS/gRNA\text{-}draper$
Figure 4A & C	$w1118/Y;+/+;UAS\text{-}pHluorin\text{-}HTT_{ex1}Q25\text{-}tdTomato/Orco\text{-}Gal4$
Figure 4B-C	$w1118/Y;+/+;UAS\text{-}pHluorin\text{-}HTT_{ex1}Q91\text{-}tdTomato/Orco\text{-}Gal4$
Figure 4D & F	$w1118/Y;+/+;UAS\text{-}pHluorin\text{-}HTT_{ex1}Q25\text{-}tdTomato/+;OK107\text{-}Gal4/+$
Figure 4E-F	$w1118/Y;+/+;UAS\text{-} pHluorin\text{-}HTT_{ex1}Q91\text{-}tdTomato/+;OK107\text{-}Gal4/+$
Figure 4G & I	$W1118/Y;Repo\text{-}Cas9/+;+/+$
Figure 4H-I	$W1118/Y;Repo\text{-}Cas9/+;gRNA\text{-}draper/+$
Figure 4J & L	$W1118/Y;Or47b\text{-}Gal4/repo\text{-}Cas9;+/UAS\text{-} UAS\text{-}pHluorin\text{-}HTT_{ex1}Q91\text{-}tdTomato$
Figure 4J-L	$W1118/Y;Or47b\text{-}Gal4/repo\text{-}Cas9;gRNA\text{-}draper/UAS\text{-} UAS\text{-}pHluorin\text{-}HTT_{ex1}Q91\text{-}tdTomato$
Figure 5A	$w1118; UAS\text{-}GFP/+;orco\text{-}Gal4/+$ & $w1118; UAS\text{-}HTT_{ex1}Q25\text{-}GFP/+;orco\text{-}Gal4/+$ & $w1118; UAS\text{-}HTT_{ex1}Q91\text{-}GFP/+;orco\text{-}Gal4/+$
Figure 5C & E	$w1118/Y; UAS\text{-}HTT_{ex1}Q25\text{-}V5/+;orco\text{-}Gal4/Toll6^{MIMICGFP}$
Figure 5D-E	$w1118/Y; UAS\text{-}HTT_{ex1}Q91\text{-}V5/+;orco\text{-}Gal4/Toll6^{MIMICGFP}$
Figure 5F & H	$w1118/Y; UAS\text{-}HTT_{ex1}Q25\text{-}V5/+;orco\text{-}Gal4/Jra.GFP$
Figure 5G-H	$w1118/Y; UAS\text{-}HTT_{ex1}Q91\text{-}V5/+;orco\text{-}Gal4/Jra.GFP$
Figure 6A-H	$w1118; UAS\text{-}HTT_{ex1}Q25\text{-}GFP/+;orco\text{-}Gal4/+$ & $w1118; UAS\text{-}HTT_{ex1}Q91\text{-}GFP/+;orco\text{-}Gal4/+$

Figure 7A, A, C, E	*w1118; QUAS-HTT$_{ex1}$Q25-mCherry/+;orco-QF2/+*
Figure 7B-D, F-H	*w1118; QUAS-HTT$_{ex1}$Q91-mCherry/+;orco-QF2/+*
Figure 8A & C-D	*or67d-QF,QUAS-HTT$_{ex1}$Q91-mCherry/+;QUAS-HTT$_{ex1}$Q25-GFP/+;mz0709-Gal4/+*
Figure 8B-D	*or67d-QF,QUAS-HTT$_{ex1}$Q91-mCherry/+;QUAS-HTT$_{ex1}$Q25-GFP/+;alrm-Gal4/+*
Figure 9A-B, E	(control) *or67d-QF,QUAS-HTT$_{ex1}$Q91-mCherry/+;+/+;mz0709-Gal4 /+ & or67d-QF,QUAS-HTT$_{ex1}$Q91-mCherry/+;+/+;mz0709-Gal4/UAS-RNAi-X & or67d-QF,QUAS-HTT$_{ex1}$Q91-mCherry/+; UAS-RNAi-X /+;mz0709-Gal4/+*
Figure 9C-E	(control) *or67d-QF,QUAS-HTT$_{ex1}$Q91-mCherry/+;+/+;repo-Gal4,Gal80^{ts}/UAS-RNAi-mCherry & or67d-QF,QUAS-HTT$_{ex1}$Q91-mCherry/+;+/+;repo-Gal4,Gal80^{ts}/UAS-RNAi-Ets21c*
Figure 10A & C-D & F	*w1118/Y;UAS-HTT$_{ex1}$Q25-GFP/+;orco-Gal4/+*
Figure 10B-C & E-H	*w1118/Y;UAS-HTT$_{ex1}$Q91-GFP/+;orco-Gal4/+*
Figure 11A & G-I	*w1118/Y;QUAS-HTTex1Q25-mCherry/UAS-LAMP1-GFP;orco-QF2/repo-Gal4*
Figure 11B-C & G-I	*w1118/Y;QUAS-HTTex1Q91-mCherry/UAS-LAMP1-GFP;orco-QF2/repo-Gal4*
Figure 11D & G-I	*w1118/Y;QUAS-HTTex1Q25-mCherry/UAS-GFP-LAMP1;orco-QF2/repo-Gal4*
Figure 11E-I	*w1118/Y;QUAS-HTTex1Q91-mCherry/UAS-GFP-LAMP1;orco-QF2/repo-Gal4*
Figure 12A-B & E-G	Control: *QUAS-HTT$_{ex1}$Q25-GFP/+/+;repo-Gal4,orco-QF2/UAS-mCherry-LGals3 & QUAS-HTT$_{ex1}$Q91-GFP/+/+;repo-Gal4,orco-QF2/UAS-mCherry-LGals3*
Figure 12C-G	Control: *QUAS-HTT$_{ex1}$Q25-GFP/+/+;repo-Gal4,orco-QF2/UAS-mCherry-LGals8 & QUAS-HTT$_{ex1}$Q91-GFP/+/+;repo-Gal4,orco-QF2/UAS-mCherry-LGals8*
Fig 13A-C	Control*: UAS-HTT$_{ex1}$Q25-GFP/Or67d-QF,QUAS-HTT$_{ex1}$Q91-mCherry;+/+;repo-Gal4/UAS-RNAi.mCherry & UAS-HTT$_{ex1}$Q25-GFP/Or67d-QF,QUAS-mHTT-mCherry;+/+;repo-Gal4/UAS-RNAi-X or UAS-HTT$_{ex1}$Q25-GFP//Or67d-QF,QUAS-mHTT-mCherry /+;UAS-RNAi-X/+ ; repo-Gal4/+*
Figure 14A-C	*Or67d-QF,QUAS-HTT$_{ex1}$Q91-mCherry /+;QUAS- HTT$_{ex1}$Q25-GFP/UAS-RNAi-X;repo-Gal4/+ & Or67d-QF,QUAS-HTT$_{ex1}$Q91-mCherry /+;QUAS-HTT$_{ex1}$Q25-GFP/+;repo-Gal4/UAS-RNAi-X*
Figure 14D-E	*Or67d-QF,UAS-HTTex1Q25-GFP/+ ; GH146-Gal4,QUAS-HTTex1Q91-mCherry/+ & Or67d-QF,UAS-HTTex1Q25-GFP/+ ; GH146-Gal4,QUAS-HTTex1Q91-mCherry/+ ; drprD5/+ & Or67d-QF,UAS-HTTex1Q25-*

	GFP/rab10{CRISPR-KO} ; GH146-Gal4,QUAS-HTTex1Q91-mCherry/+ & Or67d-QF,UAS-HTTex1Q25-GFP/rab10{CRISPR-KO} ; GH146-Gal4,QUAS-HTTex1Q91-mCherry/+ ; drprD5/+ & Or67d-QF,UAS-HTTex1Q25-GFP/rab10CRISPR ; GH146-Gal4,QUAS-HTTex1Q91-mCherry/rab14{CRISPR-KO} & Or67d-QF,UAS-HTTex1Q25-GFP/rab10CRISPR ; GH146-Gal4,QUAS-HTTex1Q91-mCherry/rab14{CRISPR-KO} ; drprD5/+
Figure 14F	*UAS-HTT$_{ex1}$Q25-GFP,QUAS-HTT$_{ex1}$Q91-mCherry/Or67d-QF;+/+;repo-Gal4/+ & UAS-HTT$_{ex1}$Q25-GFP,QUAS-HTT$_{ex1}$Q91-mCherry/Or67d-QF;+/+;repo-Gal4/UAS-RNAi-Rab10 & UAS-HTT$_{ex1}$Q25-GFP,QUAS-HTT$_{ex1}$Q91-mCherry/Or67d-QF;UAS-draper-I/+;repo-Gal4/UAS-RNAi-Rab10*
Figure 14G	*w1118/ w1118;+/+;+/+ (w1118) & Rab10(CRISPR-KO)/w1118;+/+;+/+ & Rab10(CRISPR-KO)/ Rab10(CRISPR-KO);+/+;+/+*
Figure 15A	controls: *w1118; UAS-GFP/+;orco-Gal4/+ & w1118; UAS-HTT$_{ex1}$Q25-GFP/+;orco-Gal4/+ & w1118; UAS-HTT$_{ex1}$Q91-GFP/+;orco-Gal4/+*
Figure 15B-C & H-I	*TI{TI}Rab10[EYFP]/ Y;UAS-HTT$_{ex1}$Q25-V5/+;orco-Gal4/+ & TI{TI}Rab10[EYFP]/ Y;UAS-HTT$_{ex1}$Q91-V5/+;orco-Gal4/+*
Figure 15D-E & H-I	*w1118/Y;UAS-HTT$_{ex1}$Q25-V5/TI{TI}Rab5[EYFP];orco-Gal4/+ & w1118/Y;UAS-HTT$_{ex1}$Q91-V5/TI{TI}Rab5[EYFP];orco-Gal4/+*
Figure 15F-I	*w1118/Y;UAS-HTT$_{ex1}$Q25-V5/+;orco-Gal4/TI{TI}Rab7[EYFP] & w1118/Y;UAS-HTT$_{ex1}$Q91-V5/+;orco-Gal4/ TI{TI}Rab7[EYFP]*
Figure 16A-B & G-I	*w1118/Y;QUAS-HTTex125-mCherry/UAS-YFP-Rab10;orco-QF2,repo-Gal4/+ & w1118/Y;QUAS-HTTex1Q91-mCherry/UAS-YFP-Rab10;orco-QF2,repo-Gal4/+*
Figure 16C-D & 16G-I	*w1118/Y;QUAS-HTTex125-mCherry/UAS-YFP-Rab5;orco-QF2,repo-Gal4/+ & w1118/Y;QUAS-HTTex1Q91-mCherry/UAS-YFP-Rab5;orco-QF2,repo-Gal4/+*
Figure 16E-I	*w1118/Y;QUAS-HTTex125-mCherry/UAS-YFP-Rab7;orco-QF2,repo-Gal4/+ & w1118/Y;QUAS-HTTex1Q91-mCherry/UAS-YFP-Rab7;orco-QF2,repo-Gal4/+*
Figure 17A1-3	*w1118/Y;QUAS-HTTex1Q91-mCherry/UAS-YFP-Rab10;orco-QF2,repo-Gal4/+*
Figure 17B1-2	*w1118/Y;QUAS-HTTex1Q91-mCherry/UAS-YFP-Rab5;orco-QF2,repo-Gal4/+*
Figure 17C1-2	*w1118/Y;QUAS-HTTex1Q91-mCherry/UAS-YFP-Rab7;orco-QF2,repo-Gal4/+*

Table 3. Primer sequences used for qPCR analyses.

Gene Target	Forward (F) and Reverse (R) Primer Sequences (5'-3')
rpl32	F: CCGCTTCAAGGGACAGTATCTG
	R: ATCTCGCCGCAGTAAACGC
toll-6	F: AATATTGTGGAGTGCTCGGG
	R: GCGTTTAAGGCCACTGAAAG
dorsal	F: ATGCGAGCGGTGTTCAGTAA
	R: ACGATGCGAAAAGCCAGTCT
relish	F: ACAGGACCGCATATCG
	R: GTGGGGTATTTCCGGC
draper-I	F: TGTGATCATGGTTACGGAGGAC
	R: CAGCCGGGTGGGCAA
mmp1	F: GAAGGCTCGGACAACGAGT
	R: GTCGTTGGACTGGTGATCG
ets21c	F: CAACGACGACGAACCAAAT
	R: GTTCGCGTTGGACGAATC
rab10	F: ACATCCGCCAAGTCGAACAT
	R: CTGGTTCCGGCGATCGATAA
rab5	F: AGTCCGCTGTGGGCAAGTC
	R: CTCCTGGTACTCGTGGAACTGTC
rab7	F: AATTTTGCACGCAACCGCTG
	R: GAGTAGCCAATTCGATGGTGC
drosomycin	F: GCTGTCCTGATGCTGGTGGT
	R: CGGAAAGGGACCCTTGTATCTTC
attacin-a	F: CTCGTTTGGATCTGACCAAGG
	R: CCATGACCAGCATTGTTGTAG
attacin-d	F: CGTTGAGGTTGAGATTGCCACT
	R: CGGTCCCTCAGTTCGGCATGAC
diptericinA	F: CCACCGCAGTACCCACTCAAT
	R: CGATGACTGCAAAGCCAAAACCA
metchnikowin	F: CAGTGCTGGCAGAGCCTCAT
	R: CGTCGGTTAGGATTGAAGGGCGA