Pocillopora meandrina daytime broadcast spawning in Moorea, French Polynesia

Pierrick Harnay [1] and Hollie M Putnam [2*]

[1] University of Haifa, The Leon H. Charney School of Marine Sciences, Department of
Marine Biology, Haifa Israel
[2] University of Rhode Island, College of the Environment and Life Sciences, Department
of Biological Sciences, 120 Flagg Rd. Kingston, RI 02881 USA
*corresponding author: hputnam@uri.edu

Abstract

Pocillopora are abundant branching corals that create 3-dimensional reef structure. In
French Polynesia, when corals were decimated by crown-of-thorns, bleaching, and a
cyclone between 2007-2010, _Pocillopora_ recruitment primarily drove recovery by 2016,
yet spawning times are undocumented. To address this, we surveyed _Pocillopora_ in
Moorea between September 2022-January 2023. We documented split-spawning
around the full moons, with one colony spawning November 10th (06:32, full
moon+3days) and reef-wide spawning December 9th (06:16-06:32; full moon+2days)
and December 10th (06:21-06:35; full moon+3days). Molecular analysis (n=16)
identified 100% as _P. meandrina_, with 13 as Haplotype 1a and 3 as Haplotype 8a,
suggesting _P. verrucosa, P. grandis,_ and Haplotype 10 spawn on the new moon or in
later months. These observations provide initial documentation of _Pocillopora_ spawning
in Moorea (to our knowledge), expanding on indigenous knowledge and increasing
capacity to study _Pocillopora_ reproduction, connectivity, and adaptation.

Introduction

Natural disturbances can trigger a new successional state (Knowlton 2015) and trajectory for the community, with enduring ecological legacy effects (Cuddington 2011; Kopecky et al. 2023). However, natural disturbances such as storms and predator outbreaks have now combined with Anthropogenic disturbances, such as increasing heatwaves (Oliver et al. 2021), to degrade marine ecosystems, in particular coral reefs. Reef-building, or scleractinian corals are the ecosystem engineers that create 3D structure and provide vital resources to millions of people through critical ecosystem services, such as fisheries, food security, economic enhancement via tourism and biopharmaceuticals, indigenous cultural resources, and coastal protection from storms (Costanza et al. 2014).

Coral reef construction is based on the productive symbiotic relationship between the cnidarian host and symbiotic dinoflagellates (Symbiodiniaceae), which is susceptible to dysbiosis in response to relatively small environmental perturbations (e.g., 1°C over mean summer maxima (Lesser 2011)). Of particular concern in the Anthropocene is coral bleaching, or the coral-dinoflagellate symbiotic disruption driven by marine heatwaves that often results in substantial mortality of corals (Hughes et al. 2018) and cascading negative consequences for reef function (Courtney et al. 2018; Hughes et al. 2019). In French Polynesia, coral bleaching in 2002 contributed to reduced coral cover (Pérez-Rosales et al. 2021), with >50% of corals surveyed showing bleaching in Moorea, French Polynesia (Penin et al. 2007). In Moorea in 2019, surveys of coral bleaching of the dominant genera, *Acropora* and *Pocillopora*, revealed 89% and 46% bleaching/partial mortality, respectively for these species (Speare et al. 2022). Therefore, without coral replenishment through reproduction, reef maintenance is threatened under current and future climate conditions.

On multiple Polynesian islands, the genus *Pocillopora* has become the dominant space holder following a crown-of-thorns seastar outbreak in 2007 and a major cyclone, Oli, in 2010 (Pérez-Rosales et al. 2021). For example in Moorea, French Polynesia, when coral cover was reduced by nearly ~90% due to crown-of-thorns outbreaks, bleaching, and successive cyclones between 2006 and 2010 (Holbrook et al. 2018), *Pocillopora spp.* were responsible for rapid reef recovery (Adjeroud et al. 2007; Holbrook et al. 2018). *Pocillopora spp.* are early successional species (Grigg and Maragos 1974) due to high fecundity (Sier and Olive 1994; Tsounis and Edmunds 2016) and recruitment (Adjeroud et al. 2007). Further, with their complex branching morphologies, *Pocillopora spp.* provide extensive reef habitat (Stella et al. 2010). In light of this, it is essential that we quantify and describe the process of *Pocillopora spp.* spawning in French Polynesia, which has yet to be reported in the scientific literature.

Existing literature on *Pocillopora spp.* indicates they primarily reproduce sexually via
spawning, excluding *P. damicornis*, and *P. acuta*, which can (a)sexually brood (Yeoh
and Dai 2010), but can also spawn (Ward 1992; Schmidt-Roach et al. 2014). While not
all past studies conducted genetic species identification, there are reports of spawning
near the full moon for *P. meandrina* in Australia (Schmidt-Roach et al. 2012), Hawai☐i
(Apprill et al. 2009; Schmidt-Roach et al. 2014) and for *P. grandis* in Japan (Kinzie
1993; Hirose et al. 2000, 2001). In contrast, observations of *P. verrucosa* spawning
have been made during the new moon in the Red Sea in Saudi Arabia, Sudan
(Bouwmeester et al. 2011, 2021), and Eilat (Shlesinger and Loya 1985), as well as in
South Africa (Kruger and Schleyer 1998) and in Japan (Kinzie 1993; Hirose et al. 2000,
2001; Baird et al. 2022). In light of this, we monitored *Pocillopora spp.* colonies for signs
of spawning in the lagoon of the north shore of Mo'orea French Polynesia, with
observations for each month from September 2022 to January 2023 on 3-5 days around
the full (and new moons where possible) starting ~45 min after sunrise (~06:00 in
Moorea) for ~ 2 hours (Table 1).
96
97 **Methods**
98 In order to describe spawning times to the species level, we completed visual and
99 photographic documentation of *Pocillopora* spawning, embryo development, and
100 species level genetic identification.
101
Spawning observations and sample collection
The spawning observation site was located on the north shore lagoon reef in Moorea (-
17.475757, -149.808062), which was selected for high Pocilloporidae cover (Edmunds
2022) and capacity for safe low light transit from the UC Richard B. Gump South Pacific
Research Station. All observations were made while snorkeling at ~1-3m, with a
minimum of two observers in the water. Observers were positioned to simultaneously
visualize multiple morphologies (e.g., *P. meandrina*, *P. verrucosa*, *P. grandis*). Coral
spawning observations were documented as follows: (1) spawning was filmed and
photographs of each colony were taken (Canon G7X MarkII); (2) colonies were tagged
and geolocated via GPS (Garmin GPSMAP 64); (3) biopsies were collected from each
tagged colony in sterile Whirlpaks; and (4) where possible, gametes were collected from
the water next to the colonies with 50 mL falcon tubes. Samples were returned to the
Gump South Pacific Research Station and biopsies from the tagged colonies were fixed
in DNA/RNA Shield (Zymo Research Cat # R1100-250). To determine if successful
fertilization occurred, collected gametes were placed in a MyTemp Mini Digital incubator
(Benchmark H2200-HC) set at ambient seawater temperature (~28°C). Development
through the swimming stage was monitored with a compound microscope (LEICA
DM500).
120

<u>*Pocillopora* species genetic identification</u>

Samples (n=16, Table 2) were transported to the University of Rhode Island, USA, under CITES FR2398700017-E. DNA was extracted with Quick-DNA™ Miniprep Plus Kit (Zymo Research Cat # D4069) according to the manufacturer's instructions for samples stored in DNA/RNA Shield, including the addition of 20µl of Proteinase K (20mg mL^{-1}). Samples were eluted in 50µl of kit Elution Buffer warmed to 65°C and DNA was assessed with a 1.5% agarose gel in 1x TAE buffer run for 45 mins at 100 V and stained with Biotium GelGreen (Fisher Cat NC9728313). To identify *Pocillopora spp.* we amplified the mitochondrial open reading frame (mtORF) region using the primers of Flot et al. (2008) FatP6.1 (5′-TTTGGGSATTCGTTTAGCAG-3′) and RORF (5′-SCCAATATGTTAAACASCATGTCA-3′). PCR mixes contained 12.5 µL of EmeraldAmp GT PCR Master Mix (TaKaRa Bio USA Inc. Cat # RR310B), 0.3 µL of each of the forward and reverse primers (10 µM), 1 µL of template DNA, and 10.9 µL of deionized water to 25 µL final volume per sample. Samples were amplified with the polymerase chain reaction (PCR) profile of a single denaturation step of 60s at 94 °C, followed by 30 cycles of 30s at 94 °C for denaturation, 30s at 53 °C for annealing, and extension of 75s at 72 °C; thermocycling was followed by final incubation at 72 °C for 5 min. The products were then assessed with a 1.5% agarose gel in TAE for 45 mins at 100 V to identify bands of ~1000bp. PCR products were cleaned with Qiagen MinElute PCR Purification Kit (Qiagen Cat. 28004) according to the manufacturer's instructions. Sanger sequencing was completed in both forward and reverse at the URI Genomics and Sequencing center using Applied Biosystems BigDye Terminator v3.1 chemistry. Sequences were analyzed using Geneious Alignment in GENEIOUS PRIME 2020.2.4. Pairwise alignment of the 16 sample sequences with the reference sequences (Table 3; https://osf.io/5geqd/) was completed using Clustal Omega (Sievers et al. 2011) and Tree building was completed via a Neighbor-Joining tree (Saitou and Nei 1987) with Jukes-Cantor genetic distance model (Jukes and Cantor 1969) and no outgroup.

Where mtORF sequencing identified samples as Haplotype 1a (containing both *P. grandis* and *P. meandrina*), DNAs were also amplified with the histone 3 region using the primers of Johnston et al. (2018) (PocHistoneF: 5′-ATTCAGTCTCACTCACTCACTCAC-3′ and PocHistoneR: 5′-TATCTTCGAACAGACCCACCAAAT-3′). *P. acuta* (Sample H2, Figure 2B) and *P. grandis* (POC-198, Figure 2B) were used as positive amplification and restriction enzyme controls, respectively. PCR mixes were made following methods above (but with PocHistone primers). Thermal cycling included a denaturation step of 60 s at 94 °C, followed by 30 cycles of 30 s at 94 °C, 30 s at 53 °C, and 60 s at 72 °C; and an elongation step at 72 °C for 5 min. PCR products were then used for Restriction Fragment Length Polymorphism with *Xho*I (NEB R0146Sk) according to Johnston et al. (2018) and viewed on a 2% agarose gel in 1X TAE after running for ~110 min at 70V

161 with a GeneRuler 100 bp DNA Ladder (Thermo Fisher Cat SM0241). A lack of cutting
162 indicates the species is *P. meandrina*, while two bands at 287, 382 bp indicates the
163 species identification is *P. grandis* (Fig. 2B). Raw sequence data, alignments, and fasta
164 files are archived here: https://osf.io/5geqd/ DOI 10.17605/OSF.IO/5GEQD
165
Results and Discussion
Species specific spawning
Pocillopora meandrina spawning was clearly documented from our observations
(Table 1) and genetic identification of 16 spawning colonies (Table 2). Of these 16,
mtORF sequencing (Table 1, Figure 2A) indicated 13 samples as Haplotype 1a
(Haplotype 1, also identified as PSH 9 from Gelin et al. 2017, which contains *P.*
meandrina and *P. grandis*) followed by histone 3 region identification as *P. meandrina*.
The three remaining samples were identified as *P. meandrina* lineage Haplotype 8a.
The presence of Pocilloporid species diversity in Moorea, including *P. meandrina*, *P.*
verrucosa, P. grandis, and Haplotype 10, ((Burgess et al. 2021); proposed as *P.*
tuahiniensis, Johnston and Burgess in revision) indicate additional spawning research
should be focused on the new moon in Moorea and over additional months, or times of
day. Multiple studies indicate recruitment of Pocilloporidae peaks between December -
April each year (Gleason 1996; Adjeroud et al. 2007; Edmunds et al. 2010). This broad
window and the documentation that *Pocillopora verrucosa* (morphologically identified)
spawns in the new moon in all other locations reported to date (Shlesinger and Loya
1985; Kinzie 1993; Kruger and Schleyer 1998; Hirose et al. 2000, 2001; Bouwmeester
et al. 2011, 2021; Baird et al. 2022) indicates additional work remains in the area of
species-specific *Pocillopora spp.* gametogenesis, spawning, development, and
recruitment in French Polynesia.
186
Synchronized reef-wide spawning of Pocillopora meandrina *occurs in Moorea*
In Moorea, *P. meandrina* gametes were released early in the morning at ~06:15-
06:35 local time, which was ~ 70-80 minutes after sunrise local time. This information
supports a synchronization of *Pocillopora spp.* spawning of ~45-180 minutes after
sunrise between different locations around the world, such as Australia (~45 minutes
after sunrise (Schmidt-Roach et al. 2012)), Okinawa, Japan (~95 to 120 minutes after
sunrise (Kinzie 1993; Hirose et al. 2000, 2001; Baird et al. 2022)), Hawai□i (~120
minutes after sunrise (Schmidt-Roach et al. 2014)), and the Red Sea (~180 minutes
after sunrise (Bouwmeester et al. 2011, 2021)). Split-spawning and species-specific
windows are both possible in Moorea. While only one colony of *P. meandrina* was
observed spawning during the month of November (November 10 Full +2 at 6:32 am),
P. meandrina, *P. verrucosa,* and *P. grandis* spawning has also been documented at
similar latitude on the Great Barrier Reef in November (Schmidt-Roach et al. 2012).
200

In December, most of the colonies observed had synchronized spawning. Indeed, on
December 9 (Full+2), from 6:16 am to 6:32am, a large number of colonies were
observed spawning both eggs and sperm turning the water of the lagoon cloudy. In
addition to the observations made in the lagoon, waves of eggs and sperm appeared to
be arriving from the forereef, indicating simultaneous spawning between the two
environments (lagoon and forereef), but this was not documented by our team of
observers. Personal communications from other scientists in the region confirmed that
Pocillopora spp. were spawning on forereef sites in Tetiaroa Dec 9th ~06:30 (Camille
Gâche, Pers Comm) and Rangiroa Dec 9th ~06:20 (Camille Leonard, Pers Comm). On
December 10 (Full+3), from 6:21 to 6:35, we observed a second day of reef-wide
spawning, but with a lower abundance of colonies. Our results corroborate those from
Japan and Hawai☐i that *Pocillopora spp.* can spawn across multiple days (Kinzie 1993;
Fiene-Severns 1998).
214

Larvae contain zooxanthellae and development is rapid
Once spawning commenced on 10 December 2022, the water quickly became
cloudy with eggs and sperm (Fig. 1A, B). Both appeared to be negatively buoyant and
rapidly dispersed by the waves and current. Eggs (~90μm) and developing embryos
(~120μm) that were observed at ~10:00am via compound microscopy contained
zooxanthellae (Fig. 1C, D, E). By 14:00 (~7.5 hours post fertilization, hpf) larvae were
actively swimming, matching the timing of development in the morphologically identified
P. verrucosa and *P. eydouxi* in Japan, where ciliated swimming larvae were
documented 8 hpf (Hirose et al. 2000). The rapid development of Pocilloporidae larvae
with vertical transmission that can provide nutritional supplementation (Kopp et al. 2016)
helps to ensure high recruitment success (Gleason 1996; Adjeroud et al. 2007;
Edmunds et al. 2010). This combination of traits supports the capacity of *Pocillopora*
spp. to promote rapid recovery following disturbances on the reefs of Moorea, if enough
adult colonies are available under future conditions (Speare et al. 2022).
229

Documentation is not always discovery
Although we are the first to report scientific documentation of *Pocillopora*
meandrina spawning in French Polynesia, it is most likely that coral spawning of many
species has been known by local Polynesians for decades to centuries. Merging of
traditional ecological knowledge, citizen science, and scientific monitoring and
documentation is poised to greatly expand our understanding of reproduction in French
Polynesia, which will be critical to reef recovery and maintenance under climate change.
237
238
239
240

Acknowledgements

As guests, we recognize and give thanks for the land and water resources of Polynesia, in particular Moorea, and to the traditional owners of the land, both past and present.

We are thankful for observer support from Ariana Huffmyer, Danielle Becker, Scott Burgess, Alyssa Turner, Erica Atkins, Danielle Barnes, Holly Moeller, and Raine Detmer and for the staff of the UC Richard B. Gump South Pacific Research Station for facilitating our research. We appreciate discussions with, and feedback from, Peter Edmunds, Scott Burgess, and Ariana Huffmyer. This research was funded by the U.S. National Science Foundation Grants EF-1921465 to HMP and supported by NSF-OCE 2224354 to the Moorea Coral Reef LTER, as well as a generous gift from the Gordon and Betty Moore Foundation.

Data Availability: DOI 10.17605/OSF.IO/5GEQD

Author Contributions: Funding HMP; Conceptualization PH, HMP; Field work PH, HMP; Molecular Analysis HMP; Writing PH, HMP

Conflict of interest statement: The authors have no conflict of interest.

 Figures:

282
283
Figure 1. *Pocillopora meandrina* spawning of **A, B)** eggs and sperm on the lagoon.
Samples of the gametes were collected from the seawater revealing the **C)** interaction
of symbiotic eggs and sperm followed by **D)** later stage cell division of embryos at ~4
hours post spawning. Swimming embryos were documented at 14:00, ~7.5 hours post
spawning. More images and videos are archived here: https://osf.io/5geqd/ DOI
10.17605/OSF.IO/5GEQD

Figure 2. A) Neighbor-Joining tree (Saitou and Nei 1987) with Jukes-Cantor genetic

distance model (Jukes and Cantor 1969) and no outgroup in GENEIOUS PRIME

2020.2.4. **B)** PocHistone PCR using the primers of Johnston et al. (2018) shown in the

upper row and restriction digest of the PocHistone PCR using *Xho*I indicating *P.*

meandrina in samples with a single band of 669 bp and a positive control of *P. grandis*

in sample 198 with double bands of 287, 382 bp. Raw sequence data, alignments, and

reference fasta files are archived here: https://osf.io/5geqd/ DOI

10.17605/OSF.IO/5GEQD

314 **Table 1.** *Pocillopora spp* spawning observation timeline from September 2022 to
315 January 2023. Sunrise and sunset times were reported based on this link.

Date	Sunrise Time	Lunar Phase	Spawn	Start Time	Stop Time	Notes
26 Sept 2022	5:45:00 AM	New+1	No	05:30	7:00	NA
27 Sept 2022	5:44:00 AM	New+2	No	05:40	8:00	NA
28 Sept 2022	5:43:00 AM	New+3	No	05:45	08:15	NA
07 Oct 2022	5:36:00 AM	Full-2	No	05:45	08:00	NA
08 Oct 2022	5:35:00 AM	Full-1	No	06:00	08:05	NA
06 Nov 2022	5:18:00 AM	Full-2	No	6:00	7:45	NA
07 Nov 2022	5:17:00 AM	Full-1	No	6:10	8:00	NA
08 Nov 2022	5:17:00 AM	Full	NA	NA	NA	Bad weather with lightning precluded in water observations
09 Nov 2022	5:15:00 AM	Full+1	No	5:50	8:00	NA
10 Nov 2022	5:15:00 AM	Full+2	Yes	05:45	08:25	Spawn at 06:32, 1 colony
11 Nov 2022	5:14:00 AM	Full+3	No	6:00	7:40	NA
07 Dec 2022	5:16:00 AM	Full	No	06:00	07:28	NA
08 Dec 2022	5:16:00 AM	Full+1	No	06:20	07:52	NA
09 Dec 2022	5:17:00 AM	Full+2	yes	06:05	08:00	Spawn at 06:16 to 06:32
10 Dec 2022	5:17:00 AM	Full+3	yes	06:00	07:45	Spawn at 06:21 to 06:35

11 Dec 2022	5:17:00 AM	Full+3	NA	NA	NA	Bad weather with lightning precluded in water observations
06 Jan 2023	5:29:00 AM	Full	NA	NA	NA	Bad weather with lightning precluded in water observations
07 Jan 2023	5:30:00 AM	Full+1	No	06:09	07:37	Bad weather and strong current reduced in water time
08 Jan 2023	5:30:00 AM	Full+2	No	06:10	07:18	Bad weather and strong current reduced in water time
09 Jan 2023	5:31:00 AM	Full+3	No	06:10	06:37	Bad weather, waves and current precluded water work past ~06:40

317 **Table 2.** *Pocillopora* spawning samples collected from corals observed spawning in
318 November and December 2022.

Sample ID	mtORF Sanger ID Haplotype	mtORF Citation	PocHistone RFLP ID	Species ID	Collection Date
POC-578 (Tag 192)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	10 November 2022
POC-131 (Tag 131)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-132 (Tag 132)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-133 (Tag 133)	Haplotype 8a	PSH 6 (Gélin et al. 2017)	*NA*	*P. meandrina*	09 December 2022
POC-134 (Tag 134)	Haplotype 8a	PSH 6 (Gélin et al. 2017)	*NA*	*P. meandrina*	09 December 2022

POC-135 (Tag 135)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-176 (Tag 176)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-185 (Tag 185)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-186 (Tag 186)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-189 (Tag 189)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-191 (Tag 191)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-194 (Tag 194)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-195 (Tag 195)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-196 (Tag 196)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-199 (Tag 199)	Haplotype 1a	PSH 9 (Gélin et al. 2017)	*P. meandrina*	*P. meandrina*	09 December 2022
POC-200 (Tag 200)	Haplotype 8a	PSH 6 (Gélin et al. 2017)	*NA*	*P. meandrina*	09 December 2022

319
320

Table 3. Reference mtORF sequences and their accession numbers used for sequence alignment and tree building. Raw sequence data, fasta files, and alignment are archived at https://osf.io/5geqd/ DOI 10.17605/OSF.IO/5GEQD

Haplotype ID	Accession Number	Species	Citation
Haplotype 1a	FR729326	*Pocillopora meandrina or P. grandis*	(Flot et al. 2010)
Haplotype 1c	MW619885	*Pocillopora meandrina*	(Burgess et al. 2021)
Haplotype 1d	MW619883	*Pocillopora meandrina*	(Burgess et al. 2021)
Haplotype 1e	MW619884	*Pocillopora meandrina*	(Burgess et al. 2021)
Haplotype 2	HQ378759	*Pocillopora verrucosa*	(Pinzón and LaJeunesse 2011)
Haplotype 3a	HQ378760	*Pocillopora verrucosa*	(Pinzón and LaJeunesse 2011)
Haplotype 3b	HQ378761	*Pocillopora verrucosa*	(Pinzón and LaJeunesse 2011)
Haplotype 3c	JX994075	*Pocillopora verrucosa*	(Pinzón et al. 2013)
Haplotype 3d	JX994085	*Pocillopora verrucosa*	(Pinzón et al. 2013)
Haplotype 3e	JX994083	*Pocillopora verrucosa*	(Pinzón et al. 2013)
Haplotype 3f	JX994079	*Pocillopora verrucosa*	(Pinzón et al. 2013)
Haplotype 8a	JX994074	*Pocillopora meandrina*	(Pinzón et al. 2013)
Haplotype 10	*KF381330*	*Pocillopora sp.* Proposed as *P. tuahiniensis*	*(Johnston et al. 2022)* (Johnston and Burgess in revision)
Haplotype 11	KF381328	*Pocillopora* sp.	(Johnston et al. 2022)

References Cited

Adjeroud M, Penin L, Carroll A (2007) Spatio-temporal heterogeneity in coral
recruitment around Moorea, French Polynesia: Implications for population
maintenance. J Exp Mar Bio Ecol 341:204–218

Apprill A, Marlow HQ, Martindale MQ, Rappé MS (2009) The onset of microbial
associations in the coral *Pocillopora meandrina*. ISME J 3:685–699

Baird AH, Edwards AJ, Guest JR, Harii S, Hatta M, Lachs L, Mera H, Sinniger F,
Abrego D, Ben-Zvi O, Bronstein O, Cabaitan PC, Cumbo VR, Eyal G, Eyal-Shaham
L, Feldman B, Figueiredo J, Flot J-F, Grinblat M, Heyward A, Hidaka M, Hirose M,
Iguchi A, Isomura N, Kinzie RA, Kitanobo S, Kuba A, Levy O, Loya Y, Mezaki T,
Mohamed AR, Morita M, Nojima S, Nozawa Y, Prasetia R, Puill-Stephan E,
Ramirez-Portilla C, Rapuano H, Rosenberg Y, Sakai Y, Sakai K, Shlesinger T,
Terraneo TI, Yakovleva I, Yamamoto HH, Yamazato K (2022) A coral spawning
calendar for Sesoko Station, Okinawa, Japan. Galaxea, Journal of Coral Reef
Studies 24:41–49

Bouwmeester J, Berumen ML, Baird AH (2011) Daytime broadcast spawning of
Pocillopora verrucosa on coral reefs of the central Red Sea. Galaxea, Journal of
Coral Reef Studies 13:23–24

Bouwmeester J, Coker DJ, Sinclair-Taylor TH, Berumen ML (2021) Broadcast spawning
of *Pocillopora verrucosa* across the eastern and western coast of the central Red
Sea. Ecosphere 12:

Burgess SC, Johnston EC, Wyatt ASJ, Leichter JJ, Edmunds PJ (2021) Response
diversity in corals: hidden differences in bleaching mortality among cryptic
Pocillopora species. Ecology 102:e03324

Costanza R, de Groot R, Sutton P, van der Ploeg S, Anderson SJ, Kubiszewski I,
Farber S, Turner RK (2014) Changes in the global value of ecosystem services.
Glob Environ Change 26:152–158

Courtney TA, De Carlo EH, Page HN, Bahr KD, Barro A, Howins N, Tabata R, Terlouw
G, Rodgers KS, Andersson AJ (2018) Recovery of reef☐scale calcification following
a bleaching event in Kāne☐ohe Bay, Hawai☐i. Limnol Oceanogr Lett 3:1–9

Cuddington K (2011) Legacy effects: The persistent impact of ecological interactions.
Biol Theory 6:203–210

Edmunds PJ (2022) MCR LTER: Coral Reef: Long-term population and community
dynamics: Corals, ongoing since 2005. ver 39. Environmental Data Initiative.
https://doi.org/10.6073/pasta/913b1532594fcaf02de7671ed940b4c5

Edmunds PJ, Leichter JJ, Adjeroud M (2010) Landscape-scale variation in coral
recruitment in Moorea, French Polynesia. Mar Ecol Prog Ser 414:75–89

Fiene-Severns P (1998) A note on synchronous spawning in the reef coral *Pocillopora*
meandrina at Molokini Islet, Hawai'i. 22–24

Flot J-F, Couloux A, Tillier S (2010) Haplowebs as a graphical tool for delimiting
species: a revival of Doyle's "field for recombination" approach and its application to
the coral genus *Pocillopora* in Clipperton. BMC Evol Biol 10:372

Flot J-F, Magalon H, Cruaud C, Couloux A, Tillier S (2008) Patterns of genetic structure
among Hawaiian corals of the genus *Pocillopora* yield clusters of individuals that
are compatible with morphology. C R Biol 331:239–247

Gélin P, Postaire B, Fauvelot C, Magalon H (2017) Reevaluating species number,
distribution and endemism of the coral genus *Pocillopora* Lamarck, 1816 using
species delimitation methods and microsatellites. Mol Phylogenet Evol 109:430–
446

Gleason MG (1996) Coral recruitment in Moorea, French Polynesia:the importance of
patch type and temporal variation. J Exp Mar Bio Ecol 207:79–101

Grigg RW, Maragos JE (1974) Recolonization of hermatypic corals on submerged lava
flows in Hawai i. Ecology 55:387–395

Hirose M, Kinzie RA 3rd, Hidaka M (2000) Early development of zooxanthella-
containing eggs of the corals *Pocillopora verrucosa* and *P. eydouxi* with special
reference to the distribution of zooxanthellae. Biol Bull 199:68–75

Hirose M, Kinzie R, Hidaka M (2001) Timing and process of entry of zooxanthellae into
oocytes of hermatypic corals. Coral Reefs 20:273–280

Holbrook SJ, Adam TC, Edmunds PJ, Schmitt RJ, Carpenter RC, Brooks AJ, Lenihan
HS, Briggs CJ (2018) Recruitment drives spatial variation in recovery rates of
resilient coral reefs. Sci Rep 8:7338

Hughes TP, Kerry JT, Baird AH, Connolly SR, Chase TJ, Dietzel A, Hill T, Hoey AS,
Hoogenboom MO, Jacobson M, Kerswell A, Madin JS, Mieog A, Paley AS,
Pratchett MS, Torda G, Woods RM (2019) Global warming impairs stock–
recruitment dynamics of corals. Nature 568:387–390

Hughes TP, Kerry JT, Simpson T (2018) Large-scale bleaching of corals on the Great
Barrier Reef. Ecology 99:501

Johnston EC, Forsman ZH, Toonen RJ (2018) A simple molecular technique for
distinguishing species reveals frequent misidentification of Hawaiian corals in the
genus *Pocillopora*. PeerJ 6:e4355

Johnston EC, Wyatt ASJ, Leichter JJ, Burgess SC (2022) Niche differences in co-
occurring cryptic coral species (*Pocillopora spp.*). Coral Reefs 41:767–778

Jukes TH, Cantor CR (1969) Evolution of Protein Molecules. In: Munro H.N. (eds)
Mammalian Protein Metabolism. Elsevier,

Kinzie RA (1993) Spawning in the reef corals *Pocillopora verrucosa* and *P. eydouxi* at
Sesoko island, Okinawa. Galaxea 11:93–105

Knowlton N (2015) Thresholds and Multiple Stable States in Coral Reef Community
Dynamics1. Integr Comp Biol 32:674–682

Kopecky KL, Stier AC, Schmitt RJ, Holbrook SJ, Moeller HV (2023) Material legacies
can degrade resilience: Structure-retaining disturbances promote regime shifts on
coral reefs. Ecology 104:e4006

Kopp C, Domart-Coulon I, Barthelemy D, Meibom A (2016) Nutritional input from
dinoflagellate symbionts in reef-building corals is minimal during planula larval life
stage. Sci Adv 2:e1500681

Kruger A, Schleyer MH (1998) Sexual reproduction in the coral *Pocillopora verrucosa*
(Cnidaria: Scleractinia) in KwaZulu-Natal, South Africa. Mar Biol 132:703–710

Lesser MP (2011) Coral Bleaching: Causes and Mechanisms. In: Dubinsky Z., Stambler
N. (eds) Coral Reefs: An Ecosystem in Transition. Springer Netherlands,
Dordrecht, pp 405–419

Oliver ECJ, Benthuysen JA, Darmaraki S, Donat MG, Hobday AJ, Holbrook NJ,
Schlegel RW, Sen Gupta A (2021) Marine Heatwaves. Ann Rev Mar Sci 13:313–
342

Penin L, Adjeroud M, Schrimm M, Lenihan HS (2007) High spatial variability in coral
bleaching around Moorea (French Polynesia): patterns across locations and water
depths. C R Biol 330:171–181

Pérez-Rosales G, Brandl SJ, Chancerelle Y, Siu G, Martinez E, Parravicini V, Hédouin
L (2021) Documenting decadal disturbance dynamics reveals archipelago-specific
recovery and compositional change on Polynesian reefs. Mar Pollut Bull
170:112659

Pinzón JH, LaJeunesse TC (2011) Species delimitation of common reef corals in the
genus *Pocillopora* using nucleotide sequence phylogenies, population genetics and
symbiosis ecology. Mol Ecol 20:311–325

Pinzón JH, Sampayo E, Cox E, Chauka LJ, Chen CA, Voolstra CR, LaJeunesse TC
(2013) Blind to morphology: genetics identifies several widespread ecologically
common species and few endemics among Indo-Pacific cauliflower corals
(*Pocillopora*, Scleractinia). J Biogeogr 40:1595–1608

Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing
phylogenetic trees. Mol Biol Evol 4:406–425

Schmidt-Roach S, Johnston E, Fontana S, Jury CP, Forsman Z (2014) Daytime
spawning of *Pocillopora* species in Kaneohe Bay, Hawaiʻi. Galaxea, Journal of
Coral Reef Studies 16:11–12

Schmidt-Roach S, Miller KJ, Woolsey E, Gerlach G, Baird AH (2012) Broadcast
spawning by *Pocillopora* species on the Great Barrier Reef. PLoS One 7:e50847

Shlesinger Y, Loya Y (1985) Coral community reproductive patterns: Red Sea versus
the Great Barrier Reef. Science 228:1333–1335

Sier CJS, Olive PJW (1994) Reproduction and reproductive variability in the coral
Pocillopora verrucosa from the Republic of Maldives. Mar Biol 118:713–722

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Söding J, Thompson JD, Higgins DG (2011) Fast, scalable generation
of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst
Biol 7:539

Speare KE, Adam TC, Winslow EM, Lenihan HS, Burkepile DE (2022) Size-dependent
mortality of corals during marine heatwave erodes recovery capacity of a coral reef.
Glob Chang Biol 28:1342–1358

Stella JS, Jones GP, Pratchett MS (2010) Variation in the structure of epifaunal
invertebrate assemblages among coral hosts. Coral Reefs 29:957–973

Tsounis G, Edmunds PJ (2016) The potential for self-seeding by the coral *Pocillopora*
spp. in Moorea, French Polynesia. PeerJ 4:e2544

Ward S (1992) Evidence for broadcast spawning as well as brooding in the scleractinian
coral *Pocillopora damicornis.* Mar Biol 112:641–646

Yeoh S-R, Dai C-F (2010) The production of sexual and asexual larvae within single
broods of the scleractinian coral, *Pocillopora damicornis*. Mar Biol 157:351–359