RNA structure modulates Cas13 activity and enables mismatch detection

Ofer Kimchi[1],*, Benjamin B. Larsen[2],*, Owen R. S. Dunkley[2], Aartjan J.W. te Velthuis[2], and Cameron Myhrvold[2,3,4,5] †

[1]*Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, New Jersey, 08544, USA*
[2]*Department of Molecular Biology, Princeton University, Princeton, New Jersey, 08544, USA*
[3]*Department of Chemical and Biological Engineering, Princeton University, Princeton, New Jersey, 08544, USA*
[4]*Omenn-Darling Bioengineering Institute, Princeton University, Princeton, New Jersey, 08544, USA*
[5]*Department of Chemistry, Princeton University, Princeton, New Jersey, 08544, USA*

*These authors contributed equally
†Corresponding author: cmyhrvol@princeton.edu

Key words: Cas13, CRISPR, RNA detection, RNA structure, strand displacement

Abstract

The RNA-targeting CRISPR nuclease Cas13 has emerged as a powerful tool for applications ranging from nucleic acid detection to transcriptome engineering and RNA imaging[1–6]. Cas13 is activated by the hybridization of a CRISPR RNA (crRNA) to a complementary single-stranded RNA (ssRNA) protospacer in a target RNA[1,7]. Though Cas13 is not activated by double-stranded RNA (dsRNA) *in vitro*, it paradoxically demonstrates robust RNA targeting in environments where the vast majority of RNAs are highly structured[2,8]. Understanding Cas13's mechanism of binding and activation will be key to improving its ability to detect and perturb RNA; however, the mechanism by which Cas13 binds structured RNAs remains unknown[9]. Here, we systematically probe the mechanism of LwaCas13a activation in response to RNA structure perturbations using a massively multiplexed screen. We find that there are two distinct sequence-independent modes by which secondary structure affects Cas13 activity: structure in the protospacer region competes with the crRNA and can be disrupted via a strand-displacement mechanism, while structure in the region 3' to the protospacer has an allosteric inhibitory effect. We leverage the kinetic nature of the strand displacement process to improve Cas13-based RNA detection, enhancing mismatch discrimination by up to 50-fold and enabling sequence-agnostic mutation identification at low (<1%) allele frequencies. Our work sets a new standard for CRISPR-based nucleic acid detection and will enable intelligent and secondary-structure-guided target selection while also expanding the range of RNAs available for targeting with Cas13.

Main text

The RNA-targeting CRISPR effector protein Cas13 holds tremendous promise for numerous applications, such as RNA targeting, detection, editing, and imaging[1–5,7,10]. Cas13 is activated by the hybridization of a CRISPR RNA (crRNA) spacer sequence to a complementary region in a target RNA (the protospacer)[7,11]. Once activated, Cas13 cleaves both the target RNA (*cis* cleavage) and other RNAs in solution (*trans* cleavage)[2,3,7,9]. However, the biophysical process by which the Cas13-crRNA complex binds to and is activated by a target RNA is poorly understood.

Understanding the mechanism of Cas13 activation would solve several key challenges. First, even with perfect complementarity between crRNA and protospacer, Cas13 activity levels can vary by several orders of magnitude[12–14]. While the specifics of the crRNA and target sequences—including spacer length, nucleotide sequence, and crRNA/protospacer mismatches—are known to affect Cas13 activity, much of this variation remains unaccounted for[12–14]. RNA secondary structure is an intriguing candidate to explain this variation, since unlike the CRISPR effector proteins Cas9 and Cas12, Cas13 is believed to be incapable of unwinding structured targets[11,15]. Models including only sequence effects while excluding structural effects have had success with a binary classification of active/inactive crRNAs, but have not been able to solve the regression problem of quantitatively predicting Cas13 activity from the crRNA sequence[12,13,16]. Second, although Cas13 activation requires crRNA/protospacer complementarity, Cas13 is frequently activated to a similar or even greater degree in the presence of single mismatches[17,18]. We hypothesized that these two poorly understood characteristics of Cas13 could be addressed by studying the activation of Cas13 in more detail using structured RNAs as a model system.

RNA molecules form intramolecular base pairs (secondary structures) that compete with intermolecular RNA-RNA interactions. We sought to explore how this competition affects the crRNA-target interactions underlying Cas13 activation. RNA secondary structure has long been suspected to influence Cas13 activity due to its competition with the crRNA for target base pairing, but this has been challenging to study in isolation, as primary sequence and secondary structure are inextricably linked[19]. Abudayyeh & Gootenberg *et al.* showed a small but significant negative correlation between secondary structure and Cas13a activity for crRNAs targeting different regions of four genes in HEK293FT cells[2]. By tiling the long non-coding RNA Xist in HEK293T cells, Bandaru et al. showed that crRNAs are typically more active when targeting single-stranded regions than double-stranded regions[20]. By examining different regions of natural RNAs, prior studies have been unable to distinguish between effects on Cas13 caused by changes to target structure, and those caused by changes to target sequence. Thus, the degree to which the RNA structure affects crRNA-target binding and Cas13 activity remains unclear.

RNA structure reduces LwaCas13a activity

To isolate the effect of RNA structure on Cas13 activity, we designed an ssRNA protospacer

sequence to which we could add variable amounts of secondary structure by either
intramolecular extension of an RNA hairpin, or by adding external complementary RNA or DNA
oligonucleotides of different lengths, termed "occluders" (Fig. 1A). We designed the protospacer
to reflect viral sequence diversity and to have minimal secondary structure in the absence of
these modifications (see Methods). We tested the ability of these structured protospacers to
activate LwaCas13a using cleavage of a quenched fluorescent RNA to report activity. Increased
secondary structure decreased Cas13 activity across all three assay conditions (Fig. 1A). We
next quantified Cas13 activity by fitting the fluorescence curves to effectively first-order reaction
equations, defining activity as the rate of reporter cleavage (hr^{-1}), a proxy for the concentration
of active Cas13 in the system (see Methods). We observed a high degree of correlation
between the three types of target occlusion (Fig. 1B). Cas13 activity varied by an order of
magnitude for the same sequence with different amounts of target occlusion (Fig. 1C).

Activity reduction is quantitatively explained by a kinetic strand displacement model

An equilibrium model based on the free energy of each target RNA (see Methods) failed to
quantitatively account for the degree of structure-mediated Cas13 activity reduction (Fig. 1C
light green dotted line). Labeling the free energy of the target-occluder complex by ΔG_u, the
disagreement between the large difference in thermodynamic drives (exp (-βΔG_u) ranges over
30 orders of magnitude) and the smaller difference in activities (ranging over 2 orders of
magnitude) cannot be explained by an equilibrium RNA-RNA hybridization framework. Given
known free energies of RNA-RNA binding[21], the system temperature would have to be ~7500K
to match an equilibrium model to the measured Cas13 activity reduction.

We next sought to find a suitable alternative framework to explain the measured activity levels;
a strand displacement model presents one such framework[22]. In this model, after initial binding
to the target, the crRNA and occluding strand compete through a random-walk-like process until
either the occluding strand is fully displaced or the crRNA-Cas13 complex dissociates from the
target RNA[23,24] (Fig. 1D). We hypothesized that strand displacement must occur for Cas13 to
bind structured RNA.

In our model, Cas13 first binds non-specifically to a region of RNA[15] and then performs a local
search for a sequence complementary to its bound crRNA. If Cas13 is not activated within a
given time t$_{dwell}$ (i.e. it does not fully bind to a protospacer sequence complementary to its
crRNA) it dissociates from the RNA and repeats the search process. This sequence of events is
analogous to the process by which enzymes such as the *E. coli lac* repressor search for their
binding site on DNA[25,26]. However, the strand displacement model predicts that as secondary
structure length increases, the probability of Cas13 completing the strand displacement reaction
within the time t$_{dwell}$ decreases, leading to a proportional decrease in Cas13 activity. This model
fits our measured results with a value of t$_{dwell}$ equivalent to 100 steps of a strand displacement
reaction (~2 x 10^{-5} s), in good agreement with direct measurements of typical dwell times for
DNA-binding proteins[25] (Fig. 1C dark green dotted line; see Methods).

A massively multiplexed assay reveals structure effects are sequence independent

To probe the limits of the strand displacement model, we used a massively multiplexed assay to explore a broad range of target structure conditions for multiple sequences. Our assay uses DNA oligos to create secondary structure at defined positions in the target, having previously validated their effect as a proxy for RNA structure (Fig. 1B, 2A). We designed a single 1kb-long RNA molecule with minimal internal secondary structure[27] (Extended Data Fig. 1A-C); a NUPACK prediction estimates its minimum free energy due to intramolecular contacts to be ~-6 kcal/mol, on par with random 35-nt-long RNA sequences[28]. Our target RNA is divided into two control blocks (one at each end of the molecule) and eight experimental blocks, allowing for efficient multiplexing[12]. Each block contains a 28-nt-long protospacer flanked by two 34-nt buffer regions. For occlusion, we used DNA oligos of lengths 10, 14, 21, and 28 nucleotides (nt) in 3-nt-spaced tilings, for a total of 4,608 simultaneous conditions. We tested these conditions in parallel using a microfluidic chip-based assay[6]. Summaries of the resulting dataset are shown in Fig. 2B and Extended Data Figs. 2 and 3.

Our results demonstrate that the reduction in Cas13 activity as a result of target structure is relatively sequence-independent, with the 8 experimental target blocks showing similar activity profiles in spite of large variation in absolute activity across these blocks (Fig. 2C, Extended Data Fig. 2). While 10- and 14-nt-long occluders had negligible effects on Cas13 activity, 21- and 28-mers had a strong effect. Consistent with earlier results, occluders binding to more of the protospacer typically led to a greater activity reduction. In agreement with our dwell time model and in contrast to other strand displacement systems[24,29,30], the presence or absence of toeholds (unoccluded RNA) had little effect on Cas13 activity (Extended Data Figs. 4E-F).

The data also revealed an unexpected asymmetry among the effect of occluders on Cas13, in which occluders binding to the 5' end of the protospacer had a larger effect on Cas13 activity than occluders binding the same number of nucleotides at the 3' end (Fig. 2D, Extended Data Fig. 4A-D). We accounted for the asymmetry in our model by adding a second parameter to the model described above, creating a differential in t_{dwell} depending on whether or not the 5' end of the protospacer is occluded. We found that our revised dwell time model was able to quantitatively capture the effects of secondary structure on Cas13 activity (Fig. 2D, E).

Structure occluding the region 3' of the protospacer inhibits Cas13

Surprisingly, when occluders are placed directly 3' to the protospacer, Cas13 activity is potently inhibited. This second regime of inhibition exists across all tested crRNAs, and inhibition is strong for both 21mer and 28mer occluders (Fig. 2C). The non-monotonicity of this second activity trough cannot be explained using a strand displacement model, implying this drop in activity is not due to a reduction in crRNA-target binding.

We sought to probe whether this effect on Cas13 activity is a result of competitive or allosteric inhibition. For the protospacer occluder, but not for the 3' occluder, we observe full rescue of Cas13 activity when the crRNA and occluder are added at the same time. Increasing the

153 concentrations of 3' occluder does not increase their inhibitory effect (Fig. 2F). These results
154 indicate that the activity reduction conferred by occluding the region 3' to the protospacer is
155 likely the result of an allosteric effect.

Strand displacement enhances mismatch detection

We hypothesized that the insights from our strand displacement model could help us
dramatically improve the specificity of Cas13-based RNA detection assays. Past work has
shown that secondary structure can make nucleic acid hybridization more sensitive to
mismatches, both in CRISPR-based approaches and in other assays[31–34]; we hypothesized that
given the kinetic nature of our assays, we could take advantage of the kinetic nature of strand
displacement to similar ends without the necessity of a binding toehold required in other
approaches. With no internal structure, even a mismatched crRNA is expected to bind strongly
to the target; however, an occluding strand provides an extra kinetic barrier that is less likely to
be overcome by a mismatched crRNA than by one that is perfectly complementary, thus
improving specificity given the short dwell time of inactive Cas13 on the RNA (Fig. 3A). A
differential-equation-based strand displacement model inspired by Ref.[35] supported this
hypothesis, revealing that even when both perfectly-complementary and mismatched invader
strands bind strongly to the target in equilibrium, in the presence of an occluding oligo, the
mismatched invader takes much longer to do so than the perfectly-matched invader (Fig. 3B).

We tested Cas13's ability to differentiate between a perfectly complementary target and one
containing a single A>U mutation at position 5 of the protospacer with and without secondary
structure occlusion. We tested both occluding the target (as we did previously) and occluding
the crRNA, reasoning that strand displacement would in either case result in improved
mismatch discrimination. The presence of a crRNA occluder resulted in a ~50x enhancement of
specificity compared to the no-occluder condition, measured as the maximum ratio of
WT/mismatch fluorescence (Fig. 3C). This effect was robust to large variations in concentration,
and was maximized at higher ~1-100 nM target input concentrations (Fig. 3D, Extended Data
Fig. 7B).

After testing target-occluding and crRNA-occluding oligos, and combinations of both (Fig. 3E,
Extended Data Fig. 5), we decided to focus on crRNA occluders. These provide the added
benefit of improving mismatch detection regardless of the identity of the mismatched target, and
they do not require any sample manipulation prior to detection.

We proceeded to explore the generality of Cas13 specificity enhancement by occluders. Using 3
different targets, 4 positions on each crRNA, and 2 mutations for each position, we tested how
well a mismatch could be detected by Cas13 with and without a crRNA-occluder. We measured
Cas13 activity on the perfectly matched and mismatched targets, finding that although only
74/96 mismatches (77%) led to any activity reduction in the absence of a crRNA-occluder, all 96
(100%) led to a reduction with the occluder (Fig. 4A-B, Extended Data Fig. 6). Error was thus
reduced from 23% to <1%. We found that without occlusion, the ability of Cas13 to distinguish
between a perfectly-matched target and a mismatched target is not guaranteed for any crRNA,
for any of the mismatch positions we tested, nor for any specific type of mutation (with the

exception of G>U, which has the fewest data points). However, using occluded crRNAs, Cas13
can distinguish perfectly-matched from mismatched targets in a position- and mutation-
independent manner.

We performed a dilution series of perfectly matched target RNA and mixed it with background
RNA containing a single mismatch. Without occlusion, the perfectly matched target was
detected at allele frequencies of 11% (a 1:8 ratio) but not 6% (1:16); with occlusion, it was
detected at frequencies as low as 0.4% (1:256) for all tested targets, an order-of-magnitude
sensitivity enhancement (Fig. 4C, Extended Data Fig. 7A).

To explore nucleotide-specific effects, we mutated the crRNA and target sequences to all
possible nucleotides at position 5 of the spacer. When testing all pairwise crRNA/target
combinations, we observed extensive cross-reactivity between crRNAs and targets in the
absence of occlusion. With occlusion, we observed specific detection of each target only by its
perfect-match crRNA (Fig. 4D, Extended Data Fig. 8). Occluded crRNAs can thus discriminate
all four possible alleles at a given position, demonstrating the exquisite specificity of the
approach.

To test the efficacy of our occlusion strategy in a real-world detection scenario, we designed
crRNAs targeting a single locus in the SARS-CoV-2 spike gene containing lineage-specific
mutations in adjacent codons. In this way, it is theoretically possible to distinguish Delta and
Omicron strains with only a single crRNA designed to identify each strain. Targeting
polymorphic sequences with conventional Cas13 detection schemes is challenging, as crRNAs
targeting a given mutation are likely to be affected by nearby mutations, making it nearly
impossible to distinguish variants from one another. We found that with crRNA occlusion, but
not in its absence, targeting this single locus leads to specific detection of the target strain in
both synthetic RNA (Fig. 4E, Extended Data Fig. 9) and amplified viral seedstocks (Fig. 4F)
using a simple fluorescence readout.

Discussion

In this study, we quantified the reduction in Cas13 activity due to target RNA occlusion, showed
that our results are quantitatively consistent with a strand-displacement-based model of Cas13
activation, and used this model to improve Cas13's mismatch specificity by an order of
magnitude.

Importantly, our method is simple to implement experimentally as it merely requires annealing a
DNA oligo to the crRNA prior to mixing with the target RNA. Unlike in other studies leveraging
RNA secondary structure to improve hybridization specificity, no toehold is required, enabling
trivial construction of the DNA oligo: all crRNA occluders used occluded the entire crRNA
spacer. The main downside of using crRNA-occluder duplexes to improve mismatch detection is
a reduction in Cas13 activity. To counteract this, steps can be taken to increase overall activity
such as increasing concentrations of Cas13, reporter, or target RNA.

While our experimental study focused on one ortholog of Cas13, LwaCas13a, our results are of
interest to many Cas13-based methods. Future work will explore to what extent the mechanism

of activation described here is generic or specific to this ortholog. Furthermore, significant
questions remain regarding the mechanism by which secondary structure 3' to the protospacer
leads to an allosteric reduction in Cas13 activity. For instance, extended complementarity
between the target and the structural portion of the crRNA (known as the the tag-antitag effect)
inhibits Cas13 activity by preventing the proper formation of the active site[36,37]. Future structural
and biochemical characterization will reveal whether the inhibitory region we have discovered
affects Cas13 by a similar or a different mechanism.

Due to its negligible cost, ease of implementation, orthogonality with existing approaches, and
marked improvement in detection specificity, we anticipate the adoption of our crRNA occlusion
approach into a wide range of Cas13-based techniques. Furthermore, our proposed strand
displacement model addresses a long-standing paradox, namely how a purportedly ssRNA-
specific enzyme is able to robustly target RNAs in cellular environments where RNA structure is
ubiquitous.

Methods

General reagents
Oligonucleotides were ordered from Integrated DNA Technologies (IDT). Unless otherwise
noted, chemical reagents were ordered from Sigma. Oligonucleotide sequences are listed in
Supplementary File 1.
250
crRNA design
Most crRNA spacers were designed to be perfectly complementary to their 28 nucleotide (nt)
protospacer region. For SARS-CoV-2 targeting sequences, a single synthetic mismatch was
inserted at position 5 to improve baseline specificity. Spacers were appended to the 3' end of
the consensus LwaCas13a direct repeat sequence
(AGACUACCCCAAAAACGAAGGGGACUAAAAC) and ordered from IDT as Alt-R guide RNA.
257
Target design
For our tiling experiment, we designed an RNA molecule of length 961 nucleotides with minimal
internal secondary structure. After an initial G nucleotide, the molecule is comprised of 10 target
blocks, each defined by a 34-nt buffer region, a 28-nt protospacer, and a second 34-nt buffer
region. We sought to have as many as possible of the 28-nt protospacers resemble natural
sequences.
264
To this end, we started with a set of 18,508 28-nt-long protospacer sequences compiled from
the ADAPT dataset, which has a sequence composition representative of viral diversity[12]. 3,391
sequences with poly-A, poly-C, or poly-U stretches ≥5 nts or poly-G stretches ≥4 nts were
removed. Of the remaining sequences, we removed 6,459 which had low average measured
activity, defined as ⟨out_log_k⟩ ≤ -2 (on a logarithmic scale from -4 to 0, where 0 is high
activity) using the activity definitions and measurements from Ref[12]. We used LandscapeFold[38]
with parameter $m = 2$ (m represents the minimum allowed stem length), disallowing

pseudoknots, to predict the structure landscapes of the remaining sequences. LandscapeFold
predicted that 1,287 of these remaining sequences had extremely low intramolecular structure,
defined as all nucleotides having a ≥40% probability of being unpaired in equilibrium.
275
We then aimed to find a set of these sequences that were all dissimilar from one another. First,
given a sequence s, we found all those sequences with a Hamming distance ≤15 from s. (A pair
of sequences with a Hamming distance of h share all but h nucleotides). Of these sequences,
we chose the one with the least secondary structure to keep and removed the others, with total
amount of secondary structure quantified as $\Sigma_n p_n$ where the sum is over nucleotides and p_n is
the probability of the nucleotide being paired in equilibrium. We repeated this step for each
sequence s we had not already removed. Next, we used a Smith-Waterman alignment[39] to
check for sequence similarity in non-identical nucleotide positions, repeating the same
procedure as above but, instead of Hamming distance, using the criteria of an alignment score
≥9 to define sequence similarity, where the alignment score parameters were (+1, -2, -2) for
(match, mismatch, gap). This procedure resulted in a set of 20 sequences all distant from one
another in sequence space.
288
Finally, although we had ensured each of these sequences had low secondary structure, we
wanted to minimize binding between these sequences. For each pair of sequences, we used
LandscapeFold with parameter $m = 3$ to predict the structure of the two strands, allowing for
both intra- and inter-molecular interactions. We defined two sequences to be incompatible if the
resulting prediction had any nucleotide on either sequence with a ≤40% probability of being
unpaired in equilibrium. We exhaustively enumerated the possible ordered sets of mutually
compatible sequences, finding 60 ordered sets of 5 mutually compatible sequences, and no set
of 6 mutually compatible sequences. Of these 60 sets, we chose the one with the least
structure. Under the assumption that entropic loop closure costs will create a barrier to non-
neighbor sequence pairing (i.e. that each sequence is less likely to pair to a sequence that isn't
its neighbor), we defined structure here as the sum, over the 4 pairs of neighboring sequences,
of the maximum probability of a nucleotide being paired in that sequence pair. Thus, we arrived
at a set of 5 distinct sequences from ADAPT with minimal intra- and inter-molecular structure.
These 5 sequences became the protospacer sequences corresponding to crRNAs 2, 4, 6, 8, 9.
303
The other 5 protospacer sequences as well as the buffer regions were compiled out of 64 16-nt-
long DNA sequences with minimal internal structure from Shortreed et al.[27]. Seven of these
sequences with poly-A or poly-T stretches ≥5 nts were removed. Concatenating these
sequences resulted in a long sequence with minimal structure, which we used to construct the
rest of the 961 nt-long RNA target. We used NUPACK 3[40] to predict the structure of the resulting
target, finding various predicted stems. We then made individual point mutations by hand in the
buffer regions and non-ADAPT-derived protospacers to minimize the probabilities of the
resulting stems (ensuring NUPACK predicted no base pair forming with probability ≥60% in
equilibrium), as well as to remove sequence similarity between targets (ensuring there are no
more than 5 identical consecutive nucleotides between the protospacer regions, no more than 6
identical consecutive nucleotides between two regions spanning a protospacer and a buffer,
and no more than 8 identical consecutive nucleotides in buffer regions).

316
Finally, we created a "shuffled" version of the target, placing the target blocks (numbered 1 – 10
from 5' to 3' in the original target) in the following order: 1, 4, 2, 7, 5, 3, 9, 6, 8, 10. We ensured
NUPACK 3 did not predict any base pair forming with probability ≥60% in the resulting
sequence.

For our initial experiments (Fig. 1) we filtered the ADAPT dataset sequences to those with high
activity (<out_log_k> -2) and perfect complementarity between target and crRNA in the ADAPT
dataset. We then measured LandscapeFold's prediction of the secondary structure of each
candidate protospacer sequence. For each nucleotide, we calculated the total probability that
the nucleotide is unpaired in equilibrium. The protospacer chosen had each nucleotide with at
least a 92% probability of being unpaired in equilibrium.

RNA preparation (including structured targets)
RNA targets were ordered from Integrated DNA Technologies as DNA containing a T7 promoter
sequence. Targets were then transcribed to RNA using the T7 HiScribe High Yield RNA
Synthesis Kit in 55 µL reactions (New England Biolabs) with a 16h incubation step at 37 ºC and
purified with 1.8X volume AMPure XP beads (Beckman Coulter) with the addition of 1.6X
isopropanol, then eluted into 20 µL of nuclease free (NF) water. All RNAs were then quantified
using a NanoDrop One (Thermo Fisher Scientific) or Biotek Take3Trio (Agilent) then stored in
nuclease free (NF) water at -80 ºC for later use.

Occluded targets and crRNAs were prepared by mixing DNA/RNA oligo occluders with target
RNA or crRNA in 60mM KCl (Invitrogen) in NF water at a ratio of 2:1 (BioMark assays) or 10:1
(plate reader assays) and put through an annealing cycle consisting of a high-temperature
melting step at 85 ºC for three minutes followed by gradual cooling to 10 ºC at 0.1 ºC/sec
followed by cooling to 4 ºC. For massively multiplexed assays, occluders were first pooled by
length and start position within the target block (see Extended Data Fig. 1D) such that each
resulting oligo pool contained all 8 n-mers binding to a given position within each of the
experimental target blocks. Targets and crRNAs were then used for detection assays
immediately as described below.

Targets were input into detection reactions at various concentrations. For experiments in Fig. 1,
targets were input at 7.5 x 10^8 copies/µL (cp/µL). For experiments in Fig. 2a-e, targets were
input at 8 x 10^8 cp/µL. For Fig. 2f, targets were input at 5 x 10^9 cp/µL. For experiments in Fig. 3
and 4a-b, input concentrations of 7.5 x 10^9 cp/µL were used unless otherwise noted in figure
caption. For Fig. 4c, targets were spiked in at the indicated allele frequency into a background of
5 x 10^{10} cp/µL (for occluded conditions) or 5 x 10^8 cp/µL (for non-occluded conditions). For Fig.
4d, occluded conditions used an input concentration of 5 x 10^{10} cp/µL while non-occluded
conditions used a concentration of 5 x 10^8 cp/µL.

Viral seedstock amplification
Extracted viral genomic RNA samples were acquired from BEI Resources (hCoV-19/USA/MD-
HP05285/2021 (B.1.617.2) Delta, hCoV-19/USA/GA-EHC-2811C/2021 (B.1.1.529) Omicron).

Amplification reactions using 2uL of viral RNA as the input (50uL total reaction volume) were
performed using the Qiagen One-Step RT-PCR kit according to the manufacturer's
specifications. The forward primer contained a T7 promoter sequence; 4uL of the RT-PCR
products were used as direct input for T7 transcription (see "RNA preparation" section above).
364
Cas13 detection assays
Standard bulk detection assays were performed by mixing target RNA at a ratio of 10% v/v with
90% Cas13 detection mix. The detection mix consisted of 1X RNA Detection Buffer (20 mM
HEPES pH 8.0, 54 mM KCl, 3.5% PEG-8000 in NF water), supplemented with 45nM purified
LwaCas13a (Genscript, stored in 100 mM Tris HCl pH 7.5 and 1 mM DTT), 1 U/µL murine
RNAse Inhibitor (New England Biolabs), 62.5 nM fluorescent reporter
(/5FAM/rUrUrUrUrUrU/IABkFQ/; IDT), 22.5 nM processed crRNA (IDT), and 14 mM MgOAc. In
experiments using crRNA occlusion, crRNAs were pre-annealed to DNA occluders as described
above, and used at a final concentration of 22.5 nM. 15 µL reactions were loaded in technical
triplicate onto a Greiner 384 well clear-bottom microplate (item no. 788096) and measured on
an Agilent BioTek Cytation 5 microplate reader for 3 hours with excitation at 485 nm and
detection at 528 nm every five minutes.
377
For tiling assays, Standard Biotools genotyping IFC (192.24 format) was used in a BioMark HD
for multiplexed detection. Assay mix (10% of final reaction volume) contained 1X Assay
Detection Mix (Standard Biotools) supplemented with 100 nM crRNA, 100 nM LwaCas13a
(Genscript, stored in 100 mM Tris HCl pH 7.5 and 1 mM DTT). Sample mix (90% of final
reaction volume) contained 1X Sample Buffer (44 mM Tris-HCl pH 7.5, 5.6 mM NaCl, 10 mM
MgCl, 1.1 mM DTT, 1.1% w/v PEG-8000), supplemented with murine RNAse Inhibitor (1 U/µL,
NEB), fluorescent reporter (500 nM, IDT), 1x ROX reference dye (used for normalization of
random fluctuations in fluorescence between chambers) (Standard Biotools), 1X GE Buffer
(Standard Biotools), 20mM KCl, and occluded RNA target (9×10^8 cp/µL).
387
Sample volumes of 3.5µL and assay volumes of 3.5µL, in addition to appropriate volumes of
Control Line Fluid, Actuation Fluid, and Pressure Fluid (Standard Biotools), were loaded onto
192.24 genotyping IFC chip (Standard Biotools). Chips were then placed into the Fluidigm
Controller and loaded and mixed using the Load Mix 192.24 GE script (Standard Biotools).
After mixing, reactions were run on BioMark HD at 37°C for 8 hours with measurements taken in
the fluorescein amidite (FAM) and the carboxyrhodamine (ROX) channels every five minutes.
Normalized and background-subtracted fluorescence for a given time point was calculated as
(FAM - FAM_background) / (ROX - ROX_background) where FAM_background and
ROX_background are the FAM and ROX background measurements.
397
Activity fits
Fluorescence curves were converted to activity scores by fitting the curves to effectively first-
order reactions. With a certain amount of active Cas13, the concentration of uncleaved reporter
is expected to decrease exponentially according to the reaction
$$E^{\star} + U \rightarrow E^{\star} + P$$

where $E^\star$ is the concentration of active Cas13, U the concentration of uncleaved reporter, and P
the concentration of cleaved reporter RNA. Labeling the (second-order) rate constant of this
reaction as r, the concentration of P changes over time according to

$$P(t) = P_{tot} - (P_{tot} - P(0))e^{-rEt}.$$

Assuming that $E^\star$ is constant over time, we define an activity score $v = rE^\star$, which is an
effective first-order rate constant (with units of inverse time). Assuming that r is constant across
our assays, the activity score v is thus a proxy for the amount of active Cas13. Given measured
P(0), we find best-fit values of P_tot and v to fit the kinetic curves. To account for curves very far
from saturation (e.g. NTC data) we set a minimum value of P_tot based on data from saturating
and near-saturating curves. For tiling data, we fit the first 50 timepoints (~4 hours) to discount
occasional apparent noise appearing at very late times.

Some assays including crRNA occluders displayed fluorescence curves that did not fit well to
this effective first-order reaction (Extended Data Figs. 5, 6, 7, 8) indicating a need to relax the
assumption that A is constant over time. For data shown in Extended Data Figs. 7A and 8, we
neglected the first several timepoints measured (15 and 10 timepoints, respectively,
corresponding to 75 and 50 min) since we found that doing so increased the goodness of fit. For
other assays using crRNA occluders—and those assays being directly compared to them (i.e.
those data shown in Extended Data Figs. 5, 6, 7B)---we fit the data to a series of two effective
first-order reactions:

$$U \to I \to P$$

424 Labeling the first-order rate constant of each reaction k_1 and k_2, this model yields

$$P(t) = P_{tot} - (P_{tot} - P(0))\frac{k_1 e^{-k_2 t} - k_2 e^{-k_1 t}}{k_1 - k_2}.$$

426 We define activity in this case as $v = (1 / k_1 + 1 / k_2)^{-1}$, verifying that if the equation is first order
427 (i.e. $k_1 \gg k_2$), our previous definition of activity is recovered. We indeed find negligible change in
428 the measured activities for no-occluder control (NOC) fluorescence curves between these two
429 fits.

Tiling experiment activity correction
Each experimental condition in the tiling experiment was performed with four replicates: two
technical replicates for each of the two "shuffles" of the 961-nt-long target sequence. While we
found excellent agreement between technical replicates (Extended Data Fig. 3B), there was
some variation between the results from each of the two target "shuffles" (Extended Data Fig.
3H). This variation was apparent in and correlated between the positive controls of crRNAs 1
and 10, which were always unoccluded (Extended Data Fig. 3C, H). We hypothesized that this
variability results from small variations in target concentration in our different samples.

To correct for such variations, we sought to quantify how much each RNA sample differed from
the mean. The RNA samples were divided into 192 sample conditions, each corresponding to a
single oligo pool and one target shuffling (see "RNA preparation" section, Extended Data Fig.
1C). Each of these 192 conditions was mixed with 24 assay conditions, corresponding to 8
experimental crRNAs, 2 positive control crRNAs, one non-targeting crRNA, one no-crRNA
control (NPC) and two replicates of each.

For the two replicates of crRNAs 1 and 10 (i.e. for each of the 4 positive control assay
conditions out of the 24 total assays), we considered the activity fit from the mean fluorescence
curve, averaging over the 192 sample conditions (Extended Data Fig. 3B, black dashed lines).
Then, for each sample condition, and for each positive control, we calculated the ratio of the
control's activity to its mean activity across all samples, obtaining an estimate of the degree to
which that sample's concentration deviated from the mean. We defined a correction factor as
the average of these ratios. We then divided all activities measured for that sample by this
correction. This activity correction not only decreased the spread of activities measured by the
positive controls (Extended Data Fig. 3D-G) but also decreased the variance between
measurements made on the two target "shuffles" (Extended Data Fig. 3H).

Mismatch discrimination
In Fig. 3D we show the results of one simple metric by which to measure mismatch
discrimination: the maximum of the ratio F_{PM} / F_{MM}, where F_{PM} is the average fluorescence
measurement across the perfectly matched conditions, and F_{MM} across the mismatched
conditions. To account for the arbitrary offset of fluorescence, the minimum fluorescence
measured across the NTC experiments was subtracted from both F_{PM} and F_{MM} before taking the
ratio. Since F_{PM} and F_{MM} are each measured as the average across three replicates, each
measurement of F_{PM} and F_{MM} has an inherent error, σ_{PM} and σ_{MM} (respectively), which we
quantify as the standard deviation across the three replicates at each timepoint. The error of the
ratio is then propagated as

468
$$\frac{\sqrt{(F_{PM}\sigma_{MM})^2 + (F_{MM}\sigma_{PM})^2}}{(F_{MM})^2} .$$

Elsewhere (Fig. 3E, Figs. 4A, B) we measure mismatch discrimination by a metric that relies on
activity fits: $\log_2(v_{PM} / v_{MM})$. Thus, a mismatch discrimination of 1 indicates that the measured
activity of the perfectly matched conditions is twice that of the mismatched conditions, and a
discrimination of 3 indicates the perfectly matched conditions had eight-fold higher activity than
the mismatched conditions. We used a similar measure for discrimination at low allele
frequencies (Fig. 4C), defining activity discrimination as $\log_2(v_f / v_0)$ where v_f is the activity
measured at allele frequency f, and v_0 the activity measured in the background alone.

Equilibrium model
In an equilibrium model of crRNA-target hybridization, the target has a free energy ΔG_u that
depends on occluder conditions, and the crRNA-target complex has free energy independent of
occluder. A kinetic model describing crRNA-target hybridization would be
482
$$T + E \leftrightarrow E^\star$$
483 with the ratio of forward to reverse rate constants determined by the free energy difference
484 between the target-unbound and target-bound states. Assuming that the target is in excess, we
485 define an unknown parameter α_1 such that at steady state,
486
$$E^\star = \frac{E_{tot}}{1 + e^{-\beta(\Delta G_u - \alpha_1)}}$$

487   where $\beta = 1 / k_B T$, with k_B denoting Boltzmann's constant and T temperature (measured in
488   Kelvin), and $E_{tot} = E + E^*$ is the total concentration of Cas13 enzyme in the system (measured in
489   Molar).
490
491   Our measured activity, v, is proportional to E^* as defined previously: $v = rE^*$. Considering v on a
492   logarithmic scale (as it is plotted in Fig. 1C) yields

493
$$log\,(v) \, - \, \alpha_2 \, = \, - log\left(1 + \, e^{-\beta(\Delta G_u - \alpha_1)}\right)$$

494   with $\alpha_2 = log\,(rE_{tot})$.
495
496   The two parameters α_1 and α_2 therefore only shift the curve relating $log\,(v)$ to ΔG_u, but cannot
497   alter the shape of the curve itself.
498
499   That the relationship between $log\,(v)$ and ΔG_u is approximately linear can be seen by
500   recognizing that if

501
$$e^{-\beta(\Delta G_u - \alpha_1)} \gg 1,$$

502   as is typical in our system, we have

503
$$log\,(v) \, \approx \, \beta\Delta G_u - (\alpha_1 - \alpha_2),$$

504   indicating that there is in essence only one free parameter in the system that shifts the line but
505   cannot change its slope. The slope is set by the temperature of the system.
506
ODE model
We compared the binding rate of an invading strand to a target with and without an occluder in a
model based on Srinivas & Oulridge *et al.* (2013) and Irmisch *et al.* (2020)[29,35]. The model
consists of a set of ordinary differential equations (ODEs) representing the flux into and out of
states, where each state is defined by the set of base pairs formed. Transitions between states
occur at a rate $ke^{-\Delta G}$ where ΔG is the free energy barrier to the transition (in units of $k_B T$ where
k_B is Boltzmann's constant and T is temperature in units of Kelvin) and k is an overall rate
constant. An initial state consists of a target strand (bound to an occluding strand in the case
where an occluding strand is considered), with an invading strand unbound. Initial binding of the
invader strand to the toehold has a free energy barrier of ΔG_a. The reverse step has a barrier of
$h\Delta G_R$ where h is the toehold length and ΔG_a is the (absolute value of the) typical free energy of
an RNA-RNA base pair.
519
520   In the no-occluder case, subsequent forward steps (in which an additional base pair between
521   target and invader forms) have a free energy barrier of 0, while reverse steps have a free
522   energy barrier of ΔG_a.
523
In the occluder case, the first step of the strand displacement reaction has barrier $\Delta G_P + \Delta G_S -$
$(\Delta G_R - \Delta G_D)$, where we have subtracted $(\Delta G_R - \Delta G_D)$ from the models on which we base our
work to account for the fact that in our system, the invading strand is RNA while the occluding
strand is DNA; ΔG_D is the (absolute value of the) typical free energy of an RNA-DNA base pair.
Subsequent forward steps in the strand displacement reaction have a barrier of $\Delta G_S - (\Delta G_R -$

ΔG_D), while backward steps all have a barrier of ΔG_S. The barrier from the final state, in which
the occluder has fully dissociated, back to the penultimate state, has a barrier of ΔG_D.
531
532   Parameters were set following Irmisch *et al.* to: $\Delta G_a = 18.6$; $\Delta G_R = 2.52$; $\Delta G_s = 7.4$; $\Delta G_p = 3.5$;
533    $\Delta G_m = 9.5$ (all in units of k_BT); and an overall rate constant of $k = 6 \times 10^7$/s. We set $\Delta G_D = 1.2$ to
534   be roughly half of ΔG_R, and $\Delta G_D = 25$ to be large enough to prevent reassociation on the
535   timescales considered. In Fig. 3C we plot the results of $h = 3$, $b = 27$, with a mutation at the first
536   position after the toehold.
537

Strand displacement model

In the strand displacement model, non-specific binding of Cas13 to the RNA (independent of
RNA sequence) leads to Cas13 activation, and a corresponding decrease in Cas13 dissociation
rate, when the crRNA fully binds to the protospacer complement. We denote the typical dwell
time of Cas13 on the RNA in the absence of this activation by t_{dwell}, the main parameter in the
model. Secondary structure affects activity by modulating the probability that a strand
displacement reaction completes within this time t_{dwell}. We assume that activity is directly
proportional to this probability. To estimate this probability, for each occluder, we simulated 10^5
unbiased random walks of length equal to the number of occluded protospacer nucleotides with
a reflecting boundary at 0, measuring the number of steps taken to complete the random walk,
and therefore the probability of completing the random walk within a desired number of steps.
The number of trials chosen leads to errors in our estimate of this probability < 3% in all cases
(determined by the maximum ratio of standard deviation to the mean of our estimate across 10
replicates of 10^5 random walks).
552
To estimate a typical value of t_{dwell}, we turn to classic studies of the *E. coli lac* repressor, a DNA-
binding enzyme which searches for its binding site on the DNA by iteratively binding non-
specifically to the DNA, performing a local search, and dissociating[25,26]. The dissociation rate of
the *lac* repressor when non-specifically binding DNA (i.e. the inverse of its dwell time) has been
estimated[25] to be 5×10^4/s. This dwell time corresponds to the time it would take for ~100 steps
of a strand displacement reaction, where the rate of individual steps has been estimated[35] to be
$6 \times 10^7 \times e^{-2.5}$/s $\approx 5 \times 10^6$/s. This dwell time varies with ionic concentration (among other
factors), with dwell time decreasing anywhere from 2-10-fold upon doubling KCl concentration[25].
Given the different conditions used in the plate-reader assays and in the tiling experiments,
including different ionic conditions—with the former having ~2.5-fold higher KCl concentrations
than the latter—we fit t_{dwell} separately for the two experimental methods. We used a dwell time
of 100 steps of a strand displacement reaction for the plate-reader assays, and a dwell time of
300 steps for the tiling experiments.
566
To account for the asymmetry seen in our data (Fig. 2D, Extended Data Fig. 4A-D) we allowed
for dwell time to change depending on whether the 5' end of the protospacer was occluded or
unoccluded. For the plate reader assays, our final model has a dwell time of 100 steps for those
cases where the 5' end of the protospacer was occluded, and a dwell time of 200 steps for
those cases where it wasn't. For the tiling assays, the dwell times used are 300 and 600 steps,
respectively.

References

1. Schertzer, M. D. *et al.* Cas13d-mediated isoform-specific RNA knockdown with a unified computational and experimental toolbox. *bioRxiv* (2023) doi:10.1101/2023.09.12.557474.

2. Abudayyeh, O. O. *et al.* RNA targeting with CRISPR-Cas13. *Nature* **550**, 280–284 (2017).

3. Gootenberg, J. S. *et al.* Nucleic acid detection with CRISPR-Cas13a/C2c2. *Science* **356**, 438–442 (2017).

4. Cox, D. B. T. *et al.* RNA editing with CRISPR-Cas13. *Science* **358**, 1019–1027 (2017).

5. Wang, H. *et al.* CRISPR-mediated live imaging of genome editing and transcription. *Science* **365**, 1301–1305 (2019).

6. Welch, N. L. *et al.* Multiplexed CRISPR-based microfluidic platform for clinical testing of respiratory viruses and identification of SARS-CoV-2 variants. *Nat. Med.* **28**, 1083–1094 (2022).

7. East-Seletsky, A. *et al.* Two distinct RNase activities of CRISPR-C2c2 enable guide-RNA processing and RNA detection. *Nature* **538**, 270–273 (2016).

8. Meeske, A. J., Nakandakari-Higa, S. & Marraffini, L. A. Cas13-induced cellular dormancy prevents the rise of CRISPR-resistant bacteriophage. *Nature* **570**, 241–245 (2019).

9. Liu, L. *et al.* The Molecular Architecture for RNA-Guided RNA Cleavage by Cas13a. *Cell* **170**, 714–726.e10 (2017).

10. Yang, L.-Z. *et al.* Dynamic Imaging of RNA in Living Cells by CRISPR-Cas13 Systems. *Mol. Cell* **76**, 981–997.e7 (2019).

11. Abudayyeh, O. O. *et al.* C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector. *Science* **353**, (2016).

12. Metsky, H. C. *et al.* Designing sensitive viral diagnostics with machine learning. *Nat. Biotechnol.* **40**, (2022).

13. Wei, J. *et al.* Deep learning and CRISPR-Cas13d ortholog discovery for optimized RNA

targeting. *bioRxiv* 2021.09.14.460134 (2023) doi:10.1101/2021.09.14.460134.

14. Wessels, H.-H. *et al.* Prediction of on-target and off-target activity of CRISPR-Cas13d guide

RNAs using deep learning. *Nat. Biotechnol.* (2023) doi:10.1038/s41587-023-01830-8.

15. Smargon, A. A. *et al.* Cas13b Is a Type VI-B CRISPR-Associated RNA-Guided RNase

Differentially Regulated by Accessory Proteins Csx27 and Csx28. *Mol. Cell* **65**, 618–630.e7

(2017).

16. Mantena, S. *et al.* Model-directed generation of CRISPR-Cas13a guide RNAs designs

artificial sequences that improve nucleic acid detection. *bioRxiv* 2023.09.20.557569 (2023)

doi:10.1101/2023.09.20.557569.

17. Tambe, A., East-Seletsky, A., Knott, G. J., Doudna, J. A. & O'Connell, M. R. RNA Binding

and HEPN-Nuclease Activation Are Decoupled in CRISPR-Cas13a. *Cell Rep.* **24**, 1025–

1036 (2018).

18. Vargas, A. M. M. *et al.* New design strategies for ultra-specific CRISPR-Cas13a-based

RNA-diagnostic tools with single-nucleotide mismatch sensitivity. *bioRxiv* (2023)

doi:10.1101/2023.07.26.550755.

19. Kuo, H.-C., Prupes, J., Chou, C.-W. & Finkelstein, I. J. Massively Parallel Profiling of RNA-

targeting CRISPR-Cas13d. *bioRxiv* (2023) doi:10.1101/2023.03.27.534188.

20. Bandaru, S. *et al.* Structure-based design of gRNA for Cas13. *Sci. Rep.* **10**, 1–12 (2020).

21. Turner, D. H. & Mathews, D. H. NNDB: The nearest neighbor parameter database for

predicting stability of nucleic acid secondary structure. *Nucleic Acids Res.* **38**, 2009–2011

(2009).

22. Simmel, F. C., Yurke, B. & Singh, H. R. Principles and Applications of Nucleic Acid Strand

Displacement Reactions. *Chem. Rev.* **119**, 6326–6369 (2019).

23. Radding, C. M., Beattie, K. L., Holloman, W. K. & Wiegand, R. C. Uptake of homologous

single-stranded fragments by superhelical DNA. IV. Branch migration. *J. Mol. Biol.* **116**,

825–839 (1977).

24.   Zhang, D. Y. & Winfree, E. Control of DNA Strand Displacement Kinetics Using Toehold

Exchange. *J. Am. Chem. Soc.* **131**, 1–16 (2009).

25.   Winter, R. B., Berg, O. G. & von Hippel, P. H. Diffusion-driven mechanisms of protein

translocation on nucleic acids. 3. The Escherichia coli lac repressor--operator interaction:

kinetic measurements and conclusions. *Biochemistry* **20**, 6961–6977 (1981).

26.   Halford, S. E. & Marko, J. F. How do site-specific DNA-binding proteins find their targets?

Nucleic Acids Res. **32**, 3040–3052 (2004).

27.   Shortreed, M. R. *et al.* A thermodynamic approach to designing structure-free combinatorial

DNA word sets. *Nucleic Acids Res.* **33**, 4965–4977 (2005).

28.   Fornace, M. E. *et al.* NUPACK: Analysis and Design of Nucleic Acid Structures, Devices,

and Systems. *ChemRxiv* (2022) doi:10.26434/chemrxiv-2022-xv98l.

29.   Srinivas, N. *et al.* On the biophysics and kinetics of toehold-mediated DNA strand

displacement. *Nucleic Acids Res.* **41**, 10641–10658 (2013).

30.   Yurke, B. Using DNA to power nanostructures. *Genet. Program. Evolvable Mach.* **4**, 111–

122 (2003).

31.   Zhang, D. Y., Chen, S. X. & Yin, P. Optimizing the specificity of nucleic acid hybridization.

Nat. Chem. **4**, 208–214 (2012).

32.   Kocak, D. D. *et al.* Increasing the specificity of CRISPR systems with engineered RNA

secondary structures. *Nat. Biotechnol.* **37**, 657–666 (2019).

33.   Haley, N. E. C. *et al.* Design of hidden thermodynamic driving for non-equilibrium systems

via mismatch elimination during DNA strand displacement. *Nat. Commun.* **11**, (2020).

34.   Wu, Y. *et al.* A PAM-free CRISPR/Cas12a ultra-specific activation mode based on toehold-

mediated strand displacement and branch migration. *Nucleic Acids Res.* **50**, 11727–11737

(2022).

35.   Irmisch, P., Ouldridge, T. E. & Seidel, R. Modeling DNA-Strand Displacement Reactions in

the Presence of Base-Pair Mismatches. *J. Am. Chem. Soc.* **142**, 11451–11463 (2020).

36. Meeske, A. J. & Marraffini, L. A. RNA Guide Complementarity Prevents Self-Targeting in

Type VI CRISPR Systems. *Mol. Cell* **71**, 791–801.e3 (2018).

37. Wang, B. *et al.* Structural basis for self-cleavage prevention by tag:anti-tag pairing

complementarity in type VI Cas13 CRISPR systems. *Mol. Cell* **81**, 1100–1115.e5 (2021).

38. Kimchi, O., Cragnolini, T., Brenner, M. P. & Colwell, L. J. A Polymer Physics Framework for

the Entropy of Arbitrary Pseudoknots. *Biophys. J.* **117**, 520–532 (2019).

39. Smith, T. F. & Waterman, M. S. Identification of common molecular subsequences. *J. Mol.*

Biol. **147**, 195–197 (1981).

40. Zadeh, J. N. *et al.* NUPACK: Analysis and Design of Nucleic Acid Systems. *J. Comput.*

Chem. **32**, 170–173 (2010).

Acknowledgments

661   We thank Brian Kang, Britt Adamson, Rees Garmann, and Herman Dhaliwal, as well as the
662   members of the Myhrvold and te Velthuis labs, for useful discussions.

Author contributions

664   All authors (O.K., B.B.L., O.R.S.D., A.J.W.t.V., C.M) designed research. B.B.L. and O.R.S.D.
665   performed experiments. O.K., B.B.L., O.R.S.D., and C.M. performed data analysis. O.K. and
666   C.M. performed modeling. All authors (O.K., B.B.L., O.R.S.D., A.J.W.t.V., C.M) wrote the article.

Competing interests

668   C.M. is a co-founder and consultant to Carver Biosciences and holds equity in the company. All
669   authors (O.K., B.B.L., O.R.S.D., A.J.W.t.V., C.M) are co-inventors on a patent application
670   relating to this study.

Funding

The authors and research reported in this manuscript were supported by the National Institutes
of Health grant DP2 AI175474-01 (to A.J.W.t.V.); National Institutes of Health grant R21
AI168808-01, Centers for Disease Control and Prevention grant 75D30122C15113, and the
Princeton Catalysis Initiative (all to C.M.); the Peter B. Lewis '55 Lewis-Sigler Institute/Genomics
Fund through the Lewis-Sigler Institute of Integrative Genomics at Princeton University, and the
National Science Foundation through the Center for the Physics of Biological Function (PHY-

1734030) (O.K.). B.B.L. was supported by National Institutes of Health grants T32GM007388
and T32GM148739.

Figures

Fig. 1: RNA structure leads to a reduction in Cas13 activity due to strand displacement.
A: Different amounts of structure were introduced into the target via intramolecular structure, RNA oligos, and DNA oligos; resulting fluorescent kinetic curves. Target input concentration: 7.5 x 10^8 copies/µL. "Mut" denotes 28nt of occlusion with nine mismatches to weaken binding. Data from n=3 technical replicates are shown (dots). **B:** Scatter plot comparing the impact of the different occlusion types depicted in (A) on Cas13 activity; x axis: Cas13 activity when occluded by DNA oligos, y axis: Cas13 activity when occluded by intramolecular RNA or RNA oligos. **C:** Cas13 activity vs. occluder length for RNA occluders is compared to two single-parameter models: an equilibrium model based on crRNA-target hybridization free energies(pink) and a free-energy-independent strand displacement model (green). Effects of changing the single parameter are indicated by arrows. Gray bar is NTC. **D:** Overview of strand displacement reactions. After initial binding to part of the target (blue), the crRNA (pink) and occluder (red) undergo a random walk process until one or the other is fully displaced. Displacement of the occluder leads to Cas13 activation. Error bars in B and C show standard deviation.

Fig. 2: A massively multiplexed assay modulating secondary structure. A: Overview of the
multiplexed assay, in which a total of 4,608 simultaneous assays were performed, with oligo
occluders of lengths 10, 14, 21, 28 tiling each protospacer region in 3-nt increments. Made
using BioRender. **B:** Overview of the Cas13 activity data from the multiplexed assay. Each data
point represents the mean activity resulting from averaging four time series curves (see
Methods), normalized to the non-occluded condition; positive and negative controls are not
shown (Extended Data Fig 3). **C:** Heat map showing the degree of activity reduction (darker
greens) by each 21mer and 28mer occluder. **D:** Cumulative histogram of inhibition asymmetry,
defined as the ratio of activities when the same numbers of nucleotides are occluded at the 3'
vs. 5' ends of the protospacer (see Extended Data Fig. 4C,D). **E:** Normalized Cas13 activity for
21mer (green) and 28mer (purple) occluders with different start positions in the region around
the protospacer. Each line represents one crRNA. **F:** Bar chart showing the inhibitory effect of
28mer occluders overlapping the protospacer or the region 3' to the protospacer, at different
occluder concentrations and when annealing the occluder before or at the same time as the
crRNA. In E and F, error bars show standard deviation.

Fig. 3: Designed secondary structure enhances Cas13 mismatch detection. A: Schematic
showing strand displacement by Cas13 with a perfectly-matched target sequence versus one
containing a mismatch. **B:** ODE-based model predictions of crRNA/target hybridization kinetics
with and without occlusion and mismatches. **C:** Kinetic curves showing detection of a target
sequence with and without a single A>U mismatch at spacer position 5, in the presence and
absence of occlusion. Shaded region shows range of fluorescence measurements for each
condition across replicates. **D:** Maximum fluorescence ratios with and without occlusion at a
variety of target input concentrations (see Methods). **E:** Mismatch discrimination, defined as
$\log_2$(perfect match activity / mismatch activity), for two single mutations and one double mutation
using occluders blocking either the target, the crRNA, or both. Error bars in D and E show
standard deviations propagated through the relevant formulae (see Methods).

Fig. 4: crRNA occluders enable consistent and sensitive mismatch discrimination. A:
Violin plots showing the ability of Cas13 to distinguish between wildtype targets and targets
containing mutations at four different positions in the protospacer both with and without
occlusion; position is relative to the 5' end of the protospacer. Each data point is the
discrimination ratio of perfect-match to mismatched sequence (defined as in Fig. 3E; see
Methods). **B:** Data from A, but organized by mutation type. **C:** Heatmap showing the ability of
Cas13 to detect spiked-in target in a background of mismatched sequence at decreasing allele
frequencies, both with and without occlusion. Asterisks indicate statistically significant detection
over the no-spike in control. Significance determined by one-tailed t-test $p<0.05$. Activity
discrimination is defined analogously to mismatch discrimination (see Methods). **D:** Specificity
matrix showing Cas13 activity normalized for each target, with and without occlusion, for all
possible crRNA and target nucleotides at position 5. **E:** Detection of synthetic Delta and
Omicron SARS-CoV-2 Spike gene RNA with crRNAs specific to Delta or Omicron using total
fluorescence after 90 minutes. Fluorescence of each target was normalized independently after
subtracting minimum NTC at 90 minutes. **F:** Same as in E, but using amplified viral seedstocks.

Extended data figures

Extended Data Fig. 1: Multiplexing RNA secondary structure for Cas13-based assays. A.
Criteria used in the design of target RNAs for multiplexed detection assays. **B.** Histogram
showing the minimum and total structure free energies of randomly generated RNA sequences,
compared to the free energy of the two target RNAs used in the tiling experiment. **C.** NUPACK 3
predicted minimum free energy structures for the two experimental targets[40]. **D.** Overview of the
multiplexed tiling experiment. Made using BioRender.

Extended Data Fig. 2: Activity profiles of Cas13 targeting occluded RNAs. A-D. Cas13
activity as a function of occluder start position in each 96 nt target block. **A:** 28-mers. **B:** 21-
mers. **C:** 14-mers. **D:** 10-mers. **E-H.** Same as A-D, with activities normalized to the non-
occluded condition for each target block **I.** Activities of all negative control conditions. **J.**
Normalized activities as a function of occluder start position for a single target with all four
occlusion lengths.

Extended Data Fig. 3: Controls and data normalization for multiplexed tiling assay. A.
Fluorescence curves for the set of experimental conditions chosen for the multiplexed tiling
assay. **B.** Scatter plot showing activity correlation between technical replicates in the tiling
experiment. **C.** Raw activity curves from the control (unoccluded) targets across all conditions in
tiling assay. **D.** Swarm plot showing control target activities before and after correction. **E.**
Cumulative histogram showing activity distribution for control target 1 before and after
correction. **F.** Same as in E, but for control target 10. **G.** Dot plot showing activity correlations
between the two control targets for all conditions before and after correction. **H.** Scatter plot
showing activity correlation between the two target shufflings for all tested conditions, before
and after correction. Inset: histogram of ratio of activities of points of the two target shufflings,
before and after correction, excluding negative controls. In E, F, H (inset), σ represents standard
deviation.

Extended Data Fig. 4: Strand displacement-based t$_{dwell}$ model accounts for asymmetries in occlusion pattern. A. Activity as a function of DNA occluder length for occluders of different lengths extending inwards from 3' and 5' ends of the protospacer. Strand displacement model predictions shown as dashed lines. **B.** Data from tiling experiment plotted as in A and also including oligos occluding the middle of the protospacer (yellow). Small dots show individual data points from each crRNA; large dots show the mean; error bars show standard deviation. Model predictions shown as dashed lines. **C.** Ratio of activities resulting from 21-mer occluders targeting 3' end of protospacer to those resulting from occluders targeting 5' end, plotted as a function of occluder length. Each curve shows a different crRNA. Model prediction shown as dashed line. Cumulative histograms of these data are shown in Fig. 2D. **D.** As in C, for 28-mer occluders. **E-F.** Correlation between activities resulting from 21-mer occluders and 28-mers occluding the same nucleotides. In E, 28-mers starting at positions 1, 4, 7, etc are matched with 21-mers starting at positions 3, 6, 9, etc; in F they are matched with 21-mers starting at positions 6, 9, 12, etc.

Extended Data Fig. 5: Occlusion of target RNA and crRNA increase Cas13's sensitivity to mismatches. Note on occluder nomenclature: occluders are named according to their start position from the 5' end of the target block; all occluders here are 28nt in length. Occluder s19 binds to nucleotides 1-13 of the protospacer and extends into the 5' flanking region, s31 binds to nucleotides 1-25 and extends slightly into the 5' flanking region, s52 binds to nucleotides 18-28 and extends into the 3' flanking region, and s61 binds to nucleotide 28 of the protospacer and extends into the 3' flanking region. **A.** Bar charts corresponding to Fig. 3E, representing Cas13 activity for perfectly-matched and mismatched targets when occluded by several different target-blocking occluders, as well as crRNA occluders and combinations thereof. Upper dotted line indicates activity saturation point; curves above saturation are not fit well by exponentials and therefore bars in this region are not accurate measurements of activity. **B.** NTC (no-target control) raw curves for conditions with and without a crRNA occluder. **C.** Raw fluorescence curves and fits for the data shown in A.

Extended Data Fig. 6: Mismatch detection using different types of occlusion. Raw
fluorescence curves corresponding to Fig. 4A-B detecting various mismatches within three
different target sequences for no occlusion, target occlusion, crRNA occlusion, and
target+crRNA occlusion conditions.

Extended Data Fig. 7: Fluorescence curves from experiments probing Cas13 mismatch
detection at low allele frequencies and varying target concentration with occlusion. A.
Raw fluorescence curves corresponding to Fig. 4C detecting mismatch detection sensitivity at
decreasing allele frequency with and without occlusion. **B.** Raw fluorescence curves
corresponding to Fig. 3D detecting presence of a mismatch at different target RNA
concentrations with and without occlusion. Shaded regions show the range of fluorescence
measurements for each condition across replicates.

Extended Data Fig. 8: Fluorescence curves from Cas13 specificity matrix experiment with different types of occlusion. Raw fluorescence curves corresponding to Fig. 4D using all possible crRNA and target nucleotides at a given position. Semi-transparent dashed lines show curve fits (see Methods).

Extended Data Fig. 9: Fluorescence curves for SARS-CoV-2 variant calling experiments.
Raw fluorescence curves corresponding to Figs. 4E and F targeting both Delta and Omicron
variants with guides specific to each variant.