Characterizing cell-type spatial relationships across length scales in spatially resolved

omics data

Rafael dos Santos Peixoto[1,2,*], Brendan F. Miller[1,2,*], Maigan A. Brusko[3], Lyla Atta[1,2], Manjari

Anant[1,4], Mark A. Atkinson[3], Todd M. Brusko[3], Clive H. Wasserfall[3], Jean Fan[1,2]

[1] Center for Computational Biology, Whiting School of Engineering, Johns Hopkins University,

Baltimore, MD 21211

[2] Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD 21218,

[3] Department of Pathology, Immunology, and Laboratory Medicine, University of Florida,

Gainesville, FL 32611

[4] Department of Neuroscience, Johns Hopkins University, Baltimore, MD 21218

* These authors contributed equally

Correspondence should be addressed to:

Jean Fan (jeanfan@jhu.edu)

Abstract

Spatially resolved omics technologies provide molecular profiling of cells while preserving their

organization within tissues, allowing for the evaluation of cell-type spatial relationships. We

developed CRAWDAD to quantify cell-type spatial relationships across length scales. We

highlight the utility of such multi-scale characterization on simulated data, recapitulate expected

cell-type spatial relationships in tissues such as the mouse brain and embryo, and delineate

functionally relevant spatial-defined cell-type subsets in the human spleen.

Main

Spatially resolved omics technologies enable the identification of distinct cell types while

preserving cellular spatial organization within tissues, providing an opportunity to evaluate cell-

type organizational relationships and advance our understanding of the association between cell-

type organization and tissue function. Cell-type colocalization, defined as which cell types are

spatially near each other, is critical for our understanding of healthy tissue function[1] as well as

aberrations in disease[2]. Such cell-type colocalizations can occur at different length scales, with

some cell types colocalizing to engage in paracrine signaling and other close-range interactions

at a fine, micrometer length scale[3]; others colocalizing into distinct microenvironments and

functional tissue units at a more meso-scale[1]; while others colocalizing into anatomical structures

at a more macro-scale (Fig 1a).

To quantitatively evaluate such cell-type spatial relationships across different length scales, we

present CRAWDAD, Cell-type Relationship Analysis Workflow Done Across

Distances (available at https://github.com/JEFworks-Lab/CRAWDAD and as Supplementary

Software). Briefly, given cell positions and their cell-type annotations, CRAWDAD evaluates

the statistical enrichment of each cell-type within a spatial neighborhood around cells of a

reference cell-type compared to a series of empirical null backgrounds created by shuffling cell-

type labels within spatial grids of varying sizes in order to distinguish between cell-type spatial

relationships that may occur at different length scales (Online Methods). CRAWDAD uses a

binomial proportion testing framework to derive empirical p-values that are corrected for

multiple testing and converted to Z-scores to distinguish between significant cell-type

enrichments versus depletions. CRAWDAD is amenable to any spatially resolved omics data for

which cell positions and cell-type annotations can be obtained. We demonstrate CRAWDAD on

simulated as well as real spatial omics data for tissues assayed by seqFISH[4], Slide-Seq[5], and

CODEX[6].

First, to highlight the utility of evaluating cell-type spatial relationships at multiple length scales,

we simulated 8,000 cells representing four cell types in a 1000µm-by-1000µm tissue sample

(Online Methods). In this simulated spatial dataset, cell-types B and C are intermixed with one

another, while cell-type A forms a distinct structure around cell-types B and C, that further

isolates them from cell-type D (Fig 1b). Applying CRAWDAD to characterize the spatial

relationships between each cell-type pair at multiple length scales (Online Methods), we found

that the neighborhood around reference cell-type C is significantly enriched with cells of cell-

type B from a fine length scale of ~300µm (Fig 1c-d). Likewise, we observed that the

neighborhood around reference cell-type B is significantly enriched with cells of cell-type C

from a similarly fine length scale (Fig 1c, Supp Fig 1a). Given this mutual enrichment

relationship, we interpret cell-types B and C to be spatially colocalized from this fine length

scale. In contrast, we noted cell-types A and B to be spatially colocalized from a comparably

coarser length scale of ~600µm (Fig 1c, e, Supp Fig 1b). Notably, an evaluation of spatial

colocalization that only considers the whole tissue scale may miss these distinctions regarding

length scales (Supp Fig 1c). Further, by comparing against locally defined backgrounds,

CRAWDAD's quantified cell-type spatial relationships do not change as the global background

is broadened to a larger tissue section inclusive of other cell-types in contrast to other methods

(Supp Fig 1d-f). In this manner, CRAWDAD can identify significant spatial relationships among

cell-type pairs that occur at different length scales.

We next applied CRAWDAD to evaluate cell-type spatial relationships in spatial omics datasets

of real tissues. First, we analyzed a spatial transcriptomics dataset obtained from a 4000μm-by-

4000μm section of a highly spatially organized tissue, the mouse cerebellum, assayed by Slide-

Seq[7] (Fig 2a). As Slide-Seq uses barcoded beads to profile the gene expression within tissues in a

spatially resolved manner, we assumed that the observed spatial position and cell-type identity

associated with each bead reflects the spatial position and cell-type identity of the soma of the

cell within the immediate vicinity of that bead. We applied CRAWDAD to evaluate 10,098 cells

representing 19 annotated cell-types across length scales ranging from 100μm to 1000μm (Fig

2b, Online Methods). Among the significantly colocalized cell-type pairs identified were

Purkinje neurons and Bergmann glia, (Fig 2b-d, Supp Fig 2a), which are known to interact at

close distances within the Purkinje cell layer of the cerebellum[8]. Likewise, the neighborhood

around Purkinje neurons was significantly depleted of oligodendrocytes and vice versa (Fig 2b-d,

Supp Fig 2a). As such, we may interpret these cell-types as being spatially separated, consistent

with the known spatially distinct layer structure of the cerebellum. In contrast, other spatial

analysis approaches do not distinguish between these expected cell-type spatial relationships

(Supp Fig 2d-f, Supp Note 1).

Further, without restricting to one tissue, we assessed a single-cell resolution spatial

transcriptomics dataset of a 5000μm-by-5000μm section of a whole developing embryo assayed

by seqFISH[4] (Fig 2e). We applied CRAWDAD to evaluate cell-type spatial relationships for

19,416 cells representing 22 cell types across 14 different length scales ranging from 100μm to

1000μm (Fig 2f, Online Methods). Consistent with the original publication's observations,

CRAWDAD identified significant spatial colocalization between the endothelium and

haematoendothelial progenitors as well as between the intermediate and lateral plate mesoderm

cells at the whole tissue scale. However, given CRAWDAD's multi-scale characterization, we

were also able to further identify the endothelium and haematoendothelial progenitors as being

significantly colocalized from a finer length scale (Fig 2g-h, Supp Fig 2b, from ~250µm) than

the intermediate and lateral plate mesoderm cells (Fig 2i-j, Supp Fig 2c, from ~900µm). This

potentially reflects the developmental and functional relationship between these cell-type pairs;

where endothelium and haematoendothelial progenitors are tightly intertwined in terms of both

their developmental origins and their functional roles in the blood circulatory system[9], the

intermediate and lateral plate mesoderm cells share a developmental origin but ultimately form a

variety of spatially and functionally distinct organ structures and appendages[10]. Again, such

differences between cell-type spatial relationships are difficult to discern using on other spatial

analysis methods (Supp Fig 2g-l). In this manner, CRAWDAD can quantitatively delineate cell-

type spatial relationships across multiple length scales.

Finally, having demonstrated that CRAWDAD is able to recapitulate expected cell-type spatial

relationships in both simulated and real spatial omics data of tissues, we sought to apply

CRAWDAD to characterize cell-type spatial relationships in spatial omics data of the human

spleen. The spleen is a highly structured organ where cell types interact to filter blood and

initiate immune responses. Delineating the spatial organization of cell types in the spleen can

provide insights into how these different cellular populations may achieve such diverse

immunologic functions. Therefore, we focused on six single-cell resolution spatial proteomics

datasets of the human spleen from three individuals assayed by CODEX as part of The Human

BioMolecular Atlas Program (HuBMAP)[9,11]. We first performed graph-based clustering for one

representative 9400μm by 9100μm section of a human spleen with 154,446 cells to identify 12

cell types based on 28 protein markers (Figure 3a-c, Online Methods). Applying CRAWDAD,

we observe cell-type spatial relationships to broadly recapitulate known splenic architecture with

cell-types primarily colocalizing at a coarse length scale into two major compartments defined

by follicle B cells and red pulp B cells corresponding to the white pulp (WP) and red pulp (RP),

respectively[12] (Fig 3d). Consistent with the functional roles of these compartments, cell types

identified to colocalize with red pulp B cells include macrophages, neutrophils, and monocytes

that may remove old and dead red blood cells within the RP[13]. Likewise, cell types identified to

colocalize with follicle B cells include CD4+ memory T cells and podoplanin-expressing cells

within the WP (Fig 3d). Podoplanin-expression has been previously associated with T-cell zone

reticular cells[14] and observed to surround large arteries[11] within the WP.

To determine whether such cell-type spatial colocalization relationships are consistent across

tissue sections and individuals, we further repeated these analyses with 837,952 cells from five

additional spleen samples both within and across individuals ranging from 9400μm-by-9100μm

to 9100μm-by-12100μm in size. To ensure all datasets were annotated in a uniform manner, we

applied batch-correction[15] and used a linear discriminant analysis model to transfer cell-type

annotations to these new datasets (Online Methods, Supp Fig 3). We then applied CRAWDAD

to identify similar cell-type spatial relationships corresponding to the WP and RP compartments

both within and across individuals (Fig 3d, Supp Fig 4a). In this manner, CRAWDAD can

reproducibly identify functionally relevant cell-type spatial relationships in tissues within and

across individuals.

Although our clustering analysis identified a group of cells enriched by CD4 and CD45RO

proteins, which we interpreted as CD4+ memory T cells, we were unable to distinguish follicular

helper T cells, a specialized subset of CD4+ memory T cells that play a critical role in the

adaptive immune response, due to limitations of our 28 protein-marker panel. However, we

know that follicular helper T cells interact with B cells during the B cell maturation and

differentiation process within the B cell follicles. As such, we hypothesized that follicular helper

T cells would represent a spatially-defined subset of these CD4+ memory T cells that are

colocalized with follicular B cells[16]. To identify putative follicular helper T cells, we therefore

applied the same binomial testing framework used in CRAWDAD's pairwise spatial relationship

testing to identify CD4+ memory T cells that are statistically enriched around follicle B cells

(Fig 3e, Supp Fig 4b, Online Methods). We observed that the distribution of these putative

follicular helper T cells as a percentage of all CD4+ memory T cells (34% to 50%, Supp Fig 4c)

is consistent with previous characterizations within the spleen from healthy donors achieved

through orthogonal fluorescence-activated cell sorting approaches[17,18]. In this manner,

CRAWDAD enables spatial enrichment testing to identify functionally distinct spatially defined

cell-type subsets.

As atlasing efforts such as HuBMAP[19] and others continue to profile the spatial organization of

cells within tissues, we anticipate that CRAWDAD will be a useful tool in facilitating the

identification and characterization of cell-type colocalization relationships in complex tissues to

advance our understanding of the relationship between cell-type organization and tissue function.

Online Methods

CRAWDAD (Cell-type Relationship Analysis Workflow Done Across Distances) Overview

CRAWDAD characterizes cell-type spatial relationship trends by comparing observed pairwise

cell-type spatial relationships to a set of null distributions in which cell-type labels have been

shuffled at different length scales.

Creating null distributions at different length scales

To generate empirical null distributions against which observed cell-type spatial relationships

can be compared to evaluate for statistical significance, CRAWDAD employs a grid-based cell-

type label shuffling strategy. Given a tissue containing cells represented as x-y spatial

coordinates with cell-type annotations, we partition the tissue into square grids of area r^2. Then

for all cells that reside within the same grid, cell-type labels are shuffled to create a shuffled null

distribution dataset for the given spatial length scale r. These shuffled null distributions are

created for multiple rs (ranging from 50 to 500 by default) in order to achieve a set of empirical

null distribution at different length scales. We further create multiple permutations (10 by

default) at each length scale by using a grid-offsetting approach in order to mitigate the influence

of specific grid divisions. Specifically, when using permutations, the portion of the tissue that

does not fit perfectly into the grids is divided by the number of permutations to be used as offset.

Computing cell-type spatial relationship trends

To evaluate the statistical significance of an observed pairwise cell-type spatial relationship,

CRAWDAD uses a binomial testing framework. For a given reference cell-type, CRAWDAD

defines neighboring cells as those within a Euclidean distance of a user-defined neighbor

distance d (default: 100mm) of any reference cell-type cells. CRAWDAD then calculates the

proportion of neighboring cells that are members of a given query cell-type:

$$p_1 = \frac{y_1}{n}$$

where n is the total number of total neighboring cells for a given reference cell-type, and y_1 is

the total number of cells of a given query cell-type that are neighboring cells for a given

reference cell-type.

This proportion is also computed for the previously created null distributions:

$$p_2 = \frac{y_2}{n}$$

where y_2 is the total number of cells of the query cell-type that are neighboring cells in the

shuffled null distribution datasets.

A two-sided binomial proportion test is then performed to test the equality of p_1 and p_2

against the alternative that they are not equal:

216                    $$H_0: p_1 = p_2 \text{ versus } H_A: p_1 \neq p_2$$

218   This is defined by a test statistic Z, which is:

220                    $$Z = \frac{p_1 - p_2}{\sqrt{\frac{2p(1-p)}{n}}}$$

222   where:

224                    $$p = \frac{y_1 + y_2}{2n}$$

Defining the significance threshold

We used a significance threshold of 5% (p-value < 0.05 or |Z-score| > 1.96) corrected for

multiple testing (Bonferroni method by default) based on the number of unique reference-

neighbor cell-type pairs evaluated. In addition, when there were multiple permutations, we

considered a Z-score significant if the mean value obtained across permutations is beyond the

significance threshold.

Summarizing and visualizing spatial relationship trends

To visually summarize results for all cell-type pairs, CRAWDAD creates a dot plot to show the

length scale at which the spatial relationship trend for each cell-type pair first becomes

statistically significant, encoded as the size of dots, as well as the associated multiple-testing

corrected Z-score, encoded as the color hue of dots. Notably, such summaries do not fully

capture the dynamics of pairwise cell-type spatial relationships as a function of length scale.

Specifically, if a pairwise cell-type spatial relationship becomes significant at a scale of n, but

not significant again at a scale of m (m > n), only the n scale will be represented in the plot. If a

relationship is not significant, there will be no dot associated with it.

Therefore, to visualize results for a specific pairwise cell-type spatial relationship, CRAWDAD

further creates trend plots that show the computed multiple-testing corrected Z-score value as a

function of the evaluated length scales. When there are various permutations, the mean multiple-

testing corrected Z-score across all permutations and +/- an error bar of one standard deviation is

visualized.

Subset assignment of cells

CRAWDAD can further assign cells of a given cell-type to subsets based on the enrichment or

depletion of cell-types within their neighbors using the same binomial testing framework. Given

a tissue containing cells represented as x-y spatial coordinates with cell-type annotations, we

again define a neighbor distance d. Given a reference and a query cell type, for every cell of the

reference cell-type we perform a one-sided binomial exact test to assess if the proportion of

query cell-type among the cell's neighbors are significantly greater than the proportion of the

query cell-type in the population. Cells of the given reference cell-type with a p-value below a

certain user-defined significance threshold can then be further subtyped as cells of the reference

cell-type that are enriched with cells of the query cell type.

Analysis of simulated data

To simulate a spatially resolved omics dataset, we created 8000 cells randomly positioned from 0

to 2000 microns in the x and y coordinates. We chose four points P = [(250, 250), (250, 750),

(750, 250), (750, 750)] to be the centers of the circular neighborhoods defining the cell

populations, where d(P) represents the distance from a cell to its closest point in P. We labeled

the cells with d(P) <= 100 microns as cell types B or C (50% chance each). Then, we labeled the

cells with 100 < d(P) < 300 microns as cell type A. Finally, all the other cells were labeled cell

type D. For CRAWDAD, a set of shuffled null distribution were created at length scales of 100

to 1000 separated by 50 microns for 10 permutations. To identify significant trends with multiple

testing correction, we used a Z-score threshold of 2.96.

Analysis of the developing mouse embryo

A pre-processed subset of the seqFISH data of an 8-12 somite stage embryo[4] (Embryo 1) was

obtained from the original publication. This dataset contained 19416 cells with x-y coordinates

and 22 cell-type annotations. Coordinates were originally in micrometers and converted to

microns by multiplying by 1000.

For CRAWDAD, a set of shuffled null distributions were created at length scales of 100, 200,

300, 400, 500, 600, 700, 800, 900, 1000, 1200, 1500, 3000, 6000 microns for 10 permutations. A

neighbor distance of 100 microns was used to evaluate every pairwise combination of cell-types

at each length scale. To identify significant trends with multiple testing correction, we used a Z-

score threshold of 3.88.

Analysis of the mouse cerebellum

A pre-processed subset of a Slide-seq dataset of the mouse cerebellum was obtained from the

original publication[7]. This dataset contained 10,098 beads with x-y coordinates and 19 cell-type

annotations previously predicted by RCTD[7]. Poorly represented cell types defined as those being

annotated in less than 20 beads (Choroid, Candelabrum, Ependymal, Globular, Macrophages)

were not considered in the CRAWDAD cell type colocalization, resulting in 14 remaining cell

types.

For CRAWDAD, a set of shuffled null distributions were created at length scales of 100, 200,

300, 400, 500, 600, 700, 800, 900, 1000, 1200, 1500, 3000, 6000 microns for 10 permutations. A

neighbor distance of 100 microns was used to evaluate every pairwise combination of cell-types

at each length scale. To identify significant trends with multiple testing correction, we used a Z-

score threshold of 3.81.

Analysis of the human spleen

Pre-processed and compensated Akoya CODEX datasets of 6 human spleens tissue sections from

3 different donors were downloaded from the HuBMAP Data Portal

(https://portal.hubmapconsortium.org/) corresponding to dataset IDs: HBM389.PKHL.936,

HBM772.XXCD.697, HBM556.KSFB.592, HBM825.PBVN.284, HBM568.NGPL.345, and

HBM342.FSLD.938. These datasets contained protein expression for 28 markers, x-y

coordinates, and cell segmentation area measurements for 154,446, 150,311, 152,896, 130,584,

177,777, and 226,384 segmented cells, respectively. Protein expression was normalized by cell

area and $\log_{10}$ transformed with a pseudocount of 1. Using dataset HBM389.PKHL.936, cells

were clustered via Louvain graph-based clustering[20] based on a nearest neighbor-graph

with k=50. Protein expression values were summed together for cells assigned to each cluster, and

then the values were scaled across protein expression measurements within each cluster. The

scaled expression values were used to assign cell-type annotations, which initially resulted in 13

cell types. Outer and inner Follicular B cells annotations were combined into a single Follicular

B cells annotation. Linear discriminant analysis (LDA) was performed to transfer cell-type labels

between the datasets. Prior to label transfer, paired donor datasets HBM556.KSFB.592 and

HBM568.NGPL.345, and HBM825.PBVN.284 and HBM342.FSLD.938 were harmonized[21] due

to observed batch effects. LDA cell-type label transfers were done between the following

reference and query datasets: HBM389.PKHL.936 -> HBM772.XXCD.697,

HBM389.PKHL.936 -> HBM556.KSFB.592, HBM389.PKHL.936 -> HBM825.PBVN.284,

HBM556.KSFB.592 -> HBM568.NGPL.345, HBM825.PBVN.284 -> HBM342.FSLD.938.

For CRAWDAD, a set of shuffled null distributions were created at resolutions of 100, 200, 300,

400, 500, 600, 700, 800, 900, 1000, 1200, 1500, 3000, 6000 microns for 10 permutations. A

neighbor distance of 100 microns was used to evaluate every pairwise combination of cell-types

at each scale. The corrected Z-score threshold used was 3.58.

CD4+ memory T cells were further subsetted based on their proximity to Follicle B cells using a

one-sided exact binomial test, a neighbor distance defined as 100 microns and a p-value

threshold of 0.05. With the subsets, we calculated the proportion of CD4+ memory T cells near

Follicle B cells with respect to all CD4+ memory T cells and compared across datasets.

Comparison to other methods

Ripley's Cross-K Analysis

We used the spatstat[22] (version 3.0-6) package to compute different Ripley's Cross-K values for each pairwise combination of cell types. We subtracted the theoretical K for a Poisson homogeneous processes from the isotropic edge corrected Ripley's Cross-K in visualizing results.

Squidpy's Neighborhood Enrichment

Using Squidpy[23] (version 1.2.3), we defined the spatial neighbors using a radius of 100. We calculated Squidpy's neighborhood enrichment using default parameters. The results were plotted as a heatmap of Z-scores.

Squidpy's Cooccurrence Probability

Using Squidpy[23] (version 1.2.3) and its co_occurrence function with default parameters' values, we calculated the co-occurrence probability of clusters for each cell type.

Supplementary Note 1: Conceptual differences between CRAWDAD, Squidpy's Co-occurrence ratio, and Ripley's K

Squidpy's Co-occurrence ratio

The co-occurrence ratio implemented in Squidpy[23] is a ratio of the frequency of observing a given cell type within a specific distance interval conditioned on a reference cell type, over the frequency of observing the given cell type within a specific distance interval regardless of the reference cell type. It differs from CRAWDAD in that cell type neighbors are no longer defined by a radius distance of spatial units away from a reference cell. Instead, it assesses the frequency of each cell type within a series of specific distance intervals, which can be thought of as "rings" of a given thickness away from a reference cell. This theoretically provides information about other cell type structures or patterns that are organized at certain distances away from a given reference cell type. For expected cell type relationships, such as cell types that are known to specifically cluster together, the co-occurrence ratio is expectedly high at small intervals away from the reference cell type. However, it is difficult to define a significant co-occurrence ratio to compare spatial relationships between different reference cell types and identify novel significant cell type spatial relationships.

Ripley's K

Ripley's K is a descriptive statistic of how clustered a given point process is, or how clustered multiple point processes are with respect to each other[24]. Ripley's K calculates an observed K across multiple scales and compares this to a theoretical K, which assumes complete spatial randomness and is based on a homogenous Poisson process. However, the underlying

assumption of complete spatial randomness for this theoretical distribution is likely too relaxed

and not valid for tissues in which the cell types are inherently organized at some level. This

likely leads to the observation of all cell types appearing to be highly clustered together when

measured by Ripley's K, leading to false positives. This contrasts with CRAWDAD, which uses

a permutation-based multi-scale shuffling approach to account for the inherent underlying

structure and compartmentalization of cell types in tissues.

Acknowledgements

This material is based upon work supported by the HuBMAP Integration, Visualization &

Engagement (HIVE) Initiative under Award Number 1OT2OD026673 (JF, BFM, RdSP) and the

National Institute of General Medical Sciences of the National Institutes of Health under Award

Number R35-GM142889 (JF, MA, LA) and U54AI-142766 (MA, TB, AP, CW).

Parts of this work was carried out at the Advanced Research Computing at Hopkins (ARCH)

core facility (rockfish.jhu.edu), which is supported by the National Science Foundation (NSF)

grant number OAC1920103.

Conflicts of interest

None declared.

References

1. Mitra, P. P. The Circuit Architecture of Whole Brains at the Mesoscopic Scale. *Neuron*

83, 1273–1283 (2014).

2. Elhanani, O., Ben-Uri, R. & Keren, L. Spatial profiling technologies illuminate the tumor

microenvironment. *Cancer Cell* **41**, 404–420 (2023).

3.   Manz, B. N. & Groves, J. T. Spatial organization and signal transduction at intercellular

junctions. *Nat Rev Mol Cell Biol* **11**, 342–352 (2010).

4.   Lohoff, T. *et al.* Integration of spatial and single-cell transcriptomic data elucidates mouse

organogenesis. *Nat Biotechnol* **40**, 74–85 (2022).

5.   Rodriques, S. G. *et al.* Slide-seq: A scalable technology for measuring genome-wide

expression at high spatial resolution. *Science (1979)* **363**, 1463–1467 (2019).

6.   Goltsev, Y. *et al.* Deep Profiling of Mouse Splenic Architecture with CODEX

Multiplexed Imaging. *Cell* **174**, 968-981.e15 (2018).

7.   Cable, D. M. *et al.* Robust decomposition of cell type mixtures in spatial transcriptomics.

Nat Biotechnol **40**, 517–526 (2022).

8.   Bellamy, T. C. Interactions between Purkinje neurones and Bergmann glia. *The*

Cerebellum **5**, 116–126 (2006).

9.   Wu, Y. & Hirschi, K. K. Regulation of Hemogenic Endothelial Cell Development and

Function. *Annu Rev Physiol* **83**, 17–37 (2021).

10. Gilbert SF. *Developmental Biology*. (Sinauer Associates, 2000).

11. Currlin, S. *et al.* Immune, endothelial and neuronal network map in human lymph node

and spleen. *bioRxiv* 2021.10.20.465151 (2022) doi:10.1101/2021.10.20.465151.

12. Lewis, S. M., Williams, A. & Eisenbarth, S. C. Structure-function of the immune system

in the spleen. *Sci Immunol* **4**, (2019).

13. Nolte, M. A., 't Hoen, E. N. M., Van Stijn, A., Kraal, G. & Mebius, R. E. Isolation of the

intact white pulp. Quantitative and qualitative analysis of the cellular composition of the

splenic compartments. doi:10.1002/1521-4141.

14. Cheng, H.-W. *et al.* Origin and differentiation trajectories of fibroblastic reticular cells in

the splenic white pulp. *Nat Commun* **10**, 1739 (2019).

15. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nat Methods* **16**, 1289–1296 (2019).

16. Garside, P. *et al.* Visualization of Specific B and T Lymphocyte Interactions in the Lymph

Node. *Science (1979)* **281**, 96–99 (1998).

17. Vecchione, A. *et al.* Reduced Follicular Regulatory T Cells in Spleen and Pancreatic

Lymph Nodes of Patients With Type 1 Diabetes. *Diabetes* **70**, 2892–2902 (2021).

18. Zhao, J. *et al.* Hyperactive Follicular Helper T Cells Contribute to Dysregulated Humoral

Immunity in Patients With Liver Cirrhosis. *Front Immunol* **10**, (2019).

19. Jain, S. *et al.* Advances and prospects for the Human BioMolecular Atlas Program

(HuBMAP). *Nat Cell Biol* (2023) doi:10.1038/s41556-023-01194-w.

20. Blondel, V. D., Guillaume, J.-L., Lambiotte, R. & Lefebvre, E. Fast unfolding of

communities in large networks. (2008) doi:10.1088/1742-5468/2008/10/P10008.

21. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nat Methods* **16**, 1289–1296 (2019).

22. Baddeley, A., Rubak, E. & Turner, R. *Spatial Point Patterns: Methodology and*

Applications with R. (Chapman and Hall/CRC Press, 2015).

23. Palla, G. *et al.* Squidpy: a scalable framework for spatial omics analysis. *Nat Methods* **19**,

171–178 (2022).

Figures

Figure 1. Motivating Cell-type Relationship Analysis Workflow Done Across Distances

(CRAWDAD) using simulated data. **a.** Illustration of the cell-type spatial relationships found at

different length scales. **b.** Simulated spatial omics tissue data, visualized at different scales. Each

point is a cell, colored by cell type. **c.** Summary visualization of CRAWDAD's multi-scale cell-

type spatial relationship analysis of the simulated data. The size of the dot represents the scale in

which a neighbor cell-type first reaches a significant spatial relationship with respect to a

reference cell-type. The color of the dot is the z-score at such scale (enriched or depleted). The

yellow stars represent the mutual enrichment between this cell-type pair, and therefore, their

colocalization. **d.** The multi-scale spatial relationship trend plot for reference cell-type C and

neighbor cell-type B. The horizontal dotted lines represent the z-score significance threshold

corrected for multiple testing (Z-score = ±2.96). If a neighbor cell type has a trend that crosses

the positive z-score threshold at a certain scale, then this neighbor type is deemed to be

significantly enriched at that scale given the reference cell type. If the trend crosses the negative

z-score threshold, then the neighbor cell type is deemed to be significantly depleted at that scale

given the reference cell type. The red bars denote the standard error for the z-score estimated

using permutations. **e.** The multi-scale spatial relationship trend plot for reference cell-type A

and neighbor cell-type B.

Figure 2. CRAWDAD characterizes cell-type spatial relationships in the mouse cerebellum and

embryo assayed by spatial transcriptomics. **a.** Cell types present in the cerebellum assayed by

Slide-seq. **b.** Summary visualization of the multi-scale cell-type spatial relationship analysis of

the cerebellum data. **c.** The multi-scale cell-type spatial relationship trend plot for Purkinje

neurons as the reference cell-type and Bergman glia (top) and Oligodendrocytes (bottom) as

neighbor cell-types. **d.** Zoom-in of Purkinje neurons, Bergman glia, and Oligodendrocytes in the

cerebellum. **e.** Cell types present in the embryo assayed by seqFISH. **f.** Summary of the multi-

scale cell-type spatial relationship analysis of the embryo data. **g.** Zoom-in of Endothelium and

the Hematoendothelial Progenitor cells. **h.** The multi-scale cell-type spatial relationship trend

plot for Endothelium cells as the reference cell type and Hematoendothelial Progenitor cells as

the neighbor cell type. **i.** Zoom-in of Intermediate Mesoderm and the Lateral Plate Mesoderm

cells. **j.** The multi-scale cell-type spatial trend plot for Intermediate Mesoderm cells as the

reference cell type and Lateral Plate Mesoderm cells as the neighbor cell type.

Figure 3. CRAWDAD characterize cell-type spatial relationships in the human spleen assayed

by CODEX. **a.** Heatmap of marker protein expression for annotated cell types. **b.** UMAP

reduced-dimension visualization of annotated cell-types. **c.** Spatial visualization of annotated

cell-types. **d.** Summary visualization of the multi-scale cell-type spatial relationship analysis for

tissue sections PKHL and XXCD from patient HBM966.VNKN.965. The small yellow square

represents the white pulp, while the larger yellow square represents the red pulp. **e.** Subset of

CD4+ Memory T cells near Follicle B cells. The number of CD4+ Memory T cells (n) and the

proportion of subsets (left) and spatial visualization of subsets (right).

Supplementary Figures

Supplementary Figure 1. CRAWDAD and Squidpy's Neighborhood Enrichment on Simulated

Data. **a.** CRAWDAD's multi-scale spatial relationship trend plot for reference cell-type B and

neighbor cell-type C. **b.** CRAWDAD's multi-scale spatial relationship trend plot for reference

cell-type B and neighbor cell-type A. **c.** Squidpy's neighborhood enrichment analysis on the

simulated data. **d**. New simulated data that included cell-type E contouring the previous original

dataset. **e.** Squidpy's neighborhood enrichment analysis on the new simulated data shows

changing cell-type relationships. **f.** Summary visualization of CRAWDAD's multi-scale cell-

type spatial relationship analysis of the new simulated data. The yellow box highlights that the

cell-type relationships found in the analysis of the original simulated data were maintained.

Supplementary Figure 2. Comparison of CRAWDAD, Squidpy's co-occurrence, and Ripley's

K Cross analyses. **d-f.** Cell-type neighborhood enrichment of Purkinje neurons versus all cell

types in the cerebellum Slide-seq dataset for CRAWDAD's multi-scale spatial relationship trend

plots (**d**), Squidpy's co-occurrence (**e**) and Ripley's K Cross analysis (**f**). For CRAWDAD, the

horizontal doted lines represent the significance threshold for enrichment (positive Z-score) and

separation (negative Z-score). For Ripley`s K cross, the trends above the horizontal dotted line

indicate colocalization while the ones bellow represent separation. **g-i.** Cell-type neighborhood

enrichment of Endothelium cells as reference on the Embryo dataset for CRAWDAD's multi-

scale spatial relationship trend plots (**g**), Squidpy's co-occurrence (**h**) and Ripley's K Cross

analysis (**i**). **j-l.** Cell-type neighborhood enrichment of Intermediate Mesoderm cells as reference

on the Embryo dataset for CRAWDAD's multi-scale spatial relationship trend plots (**j**),

Squidpy's co-occurrence (**k**) and Ripley's K Cross analysis (**l**).

Supplementary Figure 3. CODEX Human Spleen Samples. a. Reduced-dimension visualization

of the samples. b. Spatial visualization of cell types in the different samples. c. Heatmaps

visualizing transcriptional correlations between label-transferred cell-types in spleen samples and

the originally annotated cell-types in reference spleen sample PKHL.

Supplementary Figure 4. Analysis of other spleen samples. **a.** Summary visualization of the

multi-scale cell-type spatial relationship analysis for datasets FSLD and PBVN from patient

545   HBM245.ZWNT.288 and for datasets KSFB and NGPL from patient HBM298.KGNJ.374. **b.**

546   On the left, the spatial visualization of the subsets of CD4+ Memory T cells near and not near

547   Follicle B cells and Follicle B cells. On the right, the number of CD4+ Memory T cells (n) and

548   the proportion of those that are near Follicle B cells. **c.** Proportion of CD4+ Memory T cells near

549   Follicle B cells over all CD4+ Memory T cells in each sample.