Title

A murine model of *Trypanosoma brucei*-induced myocarditis and cardiac dysfunction

Authors

Nathan P. Crilly[1,2], Marcelle Dina Zita[3], Alexander K. Beaver[1,2], Polina Sysa-Shah[1,4], Aashik

Bhalodia[3], Kathy Gabrielson[1], Luigi Adamo[*3,] Monica R. Mugnier*[1,2]

*Co-Senior authors

Affiliations

[1] Department of Molecular and Comparative Pathobiology, School of Medicine, Johns Hopkins

University, Baltimore, Maryland, USA

[2] Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School

of Public Health, Baltimore, Maryland, USA

[3] Division of Cardiology, Department of Medicine, School of Medicine, Johns Hopkins

University, Baltimore, Maryland, USA

[4] Molecular Imaging Service Center and Cancer Functional Imaging Core, School of Medicine,

Johns Hopkins University, Baltimore, Maryland, USA

Corresponding authors

Monica R. Mugnier. mmugnie1@jhu.edu. 410-614-4894

Luigi Adamo. ladamo2@jhmi.edu

Conflict-of-interest statement

The authors have declared that no conflicts of interest exist

Abstract

Trypanosoma brucei is a protozoan parasite that causes human and animal African

trypanosomiases (HAT and AAT). Cardiac symptoms are commonly reported in HAT patients,

and intracardiac parasites with accompanying myocarditis have been observed in both natural

hosts and animal models of *T. brucei* infection. Despite the importance of *T. brucei* as a cause of

cardiac dysfunction and the dramatic socioeconomic impact of African trypanosomiases in sub-

Saharan Africa, there are currently no reproducible murine models of *T. brucei*-associated

cardiomyopathy. We present the first clinically relevant, reproducible murine model of cardiac

dysfunction in chronic *T. brucei* infection. Similar to humans, mice showed histological

evidence of myocarditis and elevation of serum NT-proBNP with electrocardiographic

abnormalities. Serum NT-proBNP levels were elevated prior to the development of severe

ventricular dysfunction. On flow cytometry, myocarditis was associated with an increase of most

myocardial immune cell populations, including multiple T cell and macrophage subsets,

corroborating the notion that *T. brucei*-associated cardiac damage is an immune-mediated event.

This novel mouse model represents a powerful and practical tool to investigate the pathogenesis

of *T. brucei*-mediated heart damage and supports the development of therapeutic options for *T.*

brucei-associated cardiac disease.

Introduction

The threat of *Trypanosoma brucei* is a fact of life throughout sub-Saharan Africa. A protozoan parasite spread by the tsetse fly vector, *T. brucei* is the causative agent of two neglected tropical diseases, human and animal African trypanosomiasis (HAT and AAT) (1). In both humans and animals, the disease is characterized by waxing and waning fever followed by chronic wasting and a coma which gives the disease its historical name, "sleeping sickness" (2, 3). Two subspecies are responsible for human infection: *T. b. gambiense*, which is endemic in West and Central Africa and *T. b. rhodesiense,* which is restricted to East and Southern Africa (4). The disease character and progression are highly variable, although *T. b. rhodesiense* infection is generally aggressive and rapidly progressive, while *T. b. gambiense* infection tends to follow a more chronic, slowly progressive course (2). Although HAT is uncommon, with fewer than 1000 cases reported in 2019 through 2020, the ubiquity of animal reservoirs and the insect vector throughout sub-Saharan Africa means that re-emergence is a constant threat (5–8). In addition to the direct impacts on human health, AAT is a severe burden to animal agriculture throughout sub-Saharan Africa and a serious contributor to food insecurity in the region, with an estimated cost of $4.75 billion per year and an incalculable impact on individual livelihoods (9).

T. brucei is often thought of as a parasite of the hemolymphatics and central nervous system. However, both historical and recent studies have led to a renewed appreciation for *T. brucei* as a parasite of extravascular spaces (10, 11). *T. brucei* parasites invade and colonize multiple extravascular tissues, including the heart, lungs, adipose tissue, skin, eyes, gonads, pancreas, and brain (12, 13). Within these spaces, tissue-resident parasites are associated with widespread

inflammation and damage, resulting in loss of organ function that contributes to mortality (10,

11, 14).

T. brucei colonizes the heart, and cardiac disease is a likely contributor to HAT mortality (15,

16). HAT patients exhibit elevated cardiac NT-proBNP, a biomarker which is a sensitive

predictor of heart failure, and 19% report at least one symptom consistent with heart failure (16).

Although electrocardiography (ECG) is not routinely performed on HAT patients,

electrocardiographic abnormalities have been reported in up to 71% of HAT patients, with most

notable changes including prolonged QT interval and low voltage(16, 17). One study found that

patients in Cameroon with dilated cardiomyopathy were significantly more likely to have anti-

trypanosomal antibodies than healthy controls, suggesting that even clinically inapparent *T.*

brucei infection might contribute to cardiac dysfunction(18). In agreement with these clinical

findings, myocarditis has been reported in autopsies of patients who have died from HAT,

suggesting that cardiac inflammation is a contributor to *T. brucei*-associated cardiac disease (19,

20). Cardiac symptoms and resulting pathology have been reported in both *T. b. gambiense* and

T. b. rhodesiense infection (16). There is some evidence that *T. b. rhodesiense* causes more

severe cardiac pathology, but the small number of recorded cases make evaluation difficult (19,

21).

Unfortunately, while there is clear evidence that cardiac disease is a consequence of African

trypanosomiasis., *T. brucei*-associated heart disease remains understudied. In particular, there

has been a lack of well-established animal models to allow detailed investigation into the

pathology, pathogenesis, and functional consequences of *T. brucei*-associated heart disease.

Several historical studies have described the histopathology of myocarditis during *T. brucei*

infection, especially in large animal models such as cattle, dogs, and nonhuman primates (22–

25). While large animal models, especially primates, accurately recapitulate human disease, the

expense, difficulty, and ethical issues of working with such model species has prevented their

widespread adoption by *T. brucei* researchers. More recently, researchers have published a rat

model of acute cardiac disease during early *T. brucei* infection. In this acute model, infected rats

exhibited myocarditis and ventricular arrhythmias at 11 days postinfection (dpi), similar to the

pathological and EKG findings reported in human patients (16, 20, 26, 27). Although the rat

model demonstrates that *T. brucei*-related cardiac disease can be recapitulated in a rodent, there

are important limitations. Notably, studies in the rat have been limited to the first two weeks of

infection, while clinical data suggest that cardiac symptoms and myocarditis are most prominent

during chronic HAT (19, 20). In addition, studies in the rat model have been limited to analyzing

histopathology and ECG findings, although other modalities such as echocardiography and

measurement of cardiac biomarkers are important for assessing the severity and progression of

heart disease. Finally, there are relatively few tools for genetic and immunological manipulations

in the rat, potentially limiting its usefulness for investigating disease pathogenesis (28) .

A mouse model of *T. brucei*-associated cardiac disease would have multiple advantages.

Compared to other animal models, mice are relatively inexpensive, safe, and simple to handle

and house, making them practical for most investigators. In addition, there are numerous well-

characterized strains and knockout lines of mice, allowing detailed investigation into disease

mechanisms (28). Mice are also well-characterized as a model species for a variety of parasitic

diseases, including those caused by African (*T. brucei*) and New World (*Trypanosoma cruzi*)

trypanosomes (27, 28). Historical studies performed with infection protocols that cannot be

replicated in modern times showed evidence of myocarditis in *T. brucei*-infected mice via

histopathology, further supporting the utility of the mouse as a model of *T. brucei*-associated

cardiac disease, although the consequences of this inflammation were not evaluated (29, 30).

Overall, the establishment of a robust, reproducible mouse model would allow for a detailed

investigation into the mechanisms and consequences of *T. brucei*-associated cardiac disease and

would be instrumental to the development of novel therapeutic options.

In this study, we present the first reproducible, clinically relevant, murine model of African

Trypanosomiasis optimized to characterize the functional changes and immunology of *T. brucei*-

associated cardiac disease.

Results

***T. brucei* infection in mice results in infection dynamics and clinical progression which are**

comparable to HAT

Trypanosomiasis can be induced in mice by injecting viable *T. brucei* parasites into the tail vein

of the mouse, resulting in a disease which is commonly used as a model for HAT (10, 31). To

demonstrate that the mouse model recapitulates major features of natural infection, we infected

adult male and female C57Bl/6J mice intravenously with 5-25 parasites of the *T. b. brucei*

subspecies, a non-human infective subspecies of *T. brucei* which is most closely related to *T. b.*

rhodesiense (4).

Following infection, parasitemia was measured every two days by hemocytometer. Parasitemia

first became detectable in the bloodstream at 4-6 dpi, and the first peak of parasitemia occurred

at 6-7 dpi (Fig 1A). At 8-10 dpi, the first trough of parasitemia occurred, when parasitemia

reached undetectable levels. For the remainder of infection, parasitemia fluctuated and was

highly variable between individual mice (Fig 1A).

T. brucei infection causes severe, multisystemic inflammation in natural hosts, with associated

splenomegaly (2, 32). To confirm that *T. brucei* infection caused systemic inflammation in our

model, mice were euthanized and the spleens were weighed at 28 dpi. Infected mice exhibited

significant splenomegaly, with a mean splenic weight of 466 g, compared to 80.1 g in age-

matched control mice (Fig 1B). Mice were euthanized when they reached 10^9 parasites per mL of

blood, lost >20% of body weight, or were assessed to be clinically terminal. Median survival

time was 39 dpi, with the first mouse reaching terminal endpoint at 31 dpi and the last surviving

until 49 dpi. Male mice exhibited a shorter survival time than female mice, although this

difference was not statistically significant (Supp. Fig 2.1A).

The dynamics of *T. brucei* infection in our model were in general agreement with other

published studies which have used similar mouse models and demonstrate that the mouse

recapitulates natural infection (10, 31, 33).

***T. brucei* infection induces elevation in NT-ProBNP and ECG abnormalities prior to a**

reduction in left ventricular ejection fraction

At 28 days postinfection (dpi)—a commonly used timepoint to represent chronic *T. brucei*

infection—we evaluated cardiac biomarkers and function (31).

We measured cardiac NT-proBNP as a biomarker of cardiac dysfunction. Cardiac NT-proBNP is

a sensitive marker of heart failure (34). Released from cardiomyocytes under conditions of

excess stretch due to pressure or volume overload, NT-proBNP is elevated in HAT patients (16).

Plasma NT-proBNP concentrations were significantly increased in infected mice at 28 dpi (Fig.

2A). NT-proBNP levels were slightly more elevated in male than in female mice, although this

difference was not statistically significant (Supp. Fig. 2A).

We used echocardiography under sedation at 28 dpi to evaluate heart function. Infected mice had

significantly increased ejection fraction and fractional shortening at 28 dpi (Fig 2B). Ejection

fraction was slightly higher in infected males than in females, although this difference was not

statistically significant (Supp. Fig. 2B). To evaluate the cardiac conduction system, we used

ECG. Infected mice exhibited several ECG changes. As compared to uninfected controls,

infected animals had a lower heart rate. A review of the EKG tracings showed that this was due

both to a reduction in the intrinsic heart rate and to a significant burden of arrhythmias, including

grouped beating and dropped beats, with significantly elevated QT intervals, including when QT

interval was corrected for heart rate (QTc) (Fig. 2C). Overall, ECG findings were highly

suggested of AV node dysfunction and intermittent high-degree atrioventricular block. ECG

abnormalities were more exaggerated in infected males than females, although the differences

were not statistically significant (Supp. Fig. 2C). All measured ECG parameters are shown in

Table 1.

To further explore our observations, we repeated our functional assessments at 33 dpi, the latest

timepoint at which mice can be experimentally manipulated before they begin to succumb to

disease. Unsurprisingly, plasma NT-proBNP was again significantly elevated, with the mean

NT-proBNP concentration (173.5 pg/mL) being similar to that observed at 28 dpi (165.4 pg/mL)

(Fig. 3A). Due to the fragility of the mice at this timepoint, echocardiography was performed

awake. Even so, 2 male mice and 1 female mouse had to be excluded due to reaching terminal

endpoint before or during the echocardiographic evaluation. Interestingly, at 33 dpi mice

exhibited a decreased ejection fraction, opposite to what was seen at 28 dpi (Fig. 3B). This could

indicate cardiac decompensation as mice approach terminal status. It should be noted that, due to

the fragility of mice at 33 dpi, echocardiography was performed awake, a technique that typically

results in higher measured ejection fraction. Therefore, our findings indicate a significant drop in

cardiac contractility between 28 dpi and 33 dpi.

***T. brucei* colonizes the myocardial interstitial space and causes myocarditis**

Within the mammalian host, parasites live extracellularly and colonize multiple extravascular

spaces, including the heart, ultimately causing fatal disease (13, 14). To confirm the presence of

intracardiac parasites in our model and to demonstrate that these parasites are extravascular, we

infected adult C57Bl/6J mice with genetically modified *T. brucei* parasites that constitutively

express the tdTomato fluorescent protein (35)(35). This parasite line has previously been shown

to exhibit similar infection dynamics as wild-type *T. brucei* parasites (13, 35). Mice were

sacrificed at 14 dpi, a timepoint when extravascular parasite populations have been established

and can be demonstrated on microscopy (10, 13). After sacrifice, we perfused mice to remove

intravascular parasites, collected the heart, and performed immunofluorescence for CD31 to

visualize the vasculature. *T. brucei* parasites were identified within the interstitial spaces of the

heart. Intracardiac parasites localized separately from CD31-lined spaces, indicating that

intracardiac parasites are extravascular (Fig 4, Supp. Fig 3).

To evaluate cardiac pathology due to *T. brucei* infection, we performed histopathology on the

hearts of 6 *T. brucei*-infected mice at 28 dpi. Of the 6 hearts evaluated, 5 met the Dallas Criteria

for lymphocytic myocarditis, while the sixth exhibited borderline myocarditis (36). To further

characterize histopathological changes, we used an established quantitative grading system to

evaluate the severity of myocarditis (Table 2) (37). At 28 dpi, all mice exhibited myocarditis,

with an average histological grade of 2, indicating moderate multifocal myocarditis (Fig 5A).

Inflammation affected all layers of the heart, including endocardium, myocardium, and

epicardium. On histopathology, observed cells included large numbers of lymphocytes, plasma

cells, and macrophages. Necrosis was not a major feature. The severity of inflammation was

most severe in the atria and at the atrioventricular (AV) junction (Fig 5B). The pattern and

severity of inflammation observed were consistent with previous histopathology findings in

humans and natural hosts (20, 22–25, 29, 38, 39). To evaluate cardiac fibrosis, longitudinal

sections of the heart were stained with Masson's Trichrome, and the percentage of collagen in

the myocardium was quantified using the Automated Fibrosis Analysis Tool (AFAT) (40). At 28

dpi, infected mice had a higher percentage of collagen in the heart compared to uninfected mice,

although this difference was not statistically significant (Fig 5C).

Immune cell populations in the heart are consistent with immune-mediated myocarditis

We had identified myocarditis as a consequence of *T. brucei* infection, and previous research has

indicated that the host immune response may contribute to *T. brucei*-associated cardiac

pathology (29). To better understand the intracardiac immune response to *T. brucei*, we

performed flow cytometry at 28 dpi to characterize the immune cell population in the heart.

Numbers of CD45+ cells/mg of tissue were markedly elevated in the hearts of all infected mice

at 28 dpi, supporting our histopathological finding of myocarditis (Fig 6A).

Examining individual cell populations, most immune cell lineages were more abundant in *T.*

brucei infected hearts than in age-matched controls (Fig 6B-C). In agreement with the functional

data, most immune cell populations were slightly more elevated in males than in females,

although the difference was not statistically significant for any cell lineage (Supp. Fig 2E-F).

Numbers of both CD4+ T cells and macrophages were significantly increased in infected hearts,

which is unsurprising, since these cell types are necessary for an effective immune response to

African trypanosomes (Fig 6B-C) (41). Of the different macrophage subsets, the CCR2^{hi}MHC-

II^{hi} population experienced the largest increase in numbers. This subset represents pro-

inflammatory, monocyte-derived macrophages, indicating that *T. brucei* infection causes an

influx of pro-inflammatory macrophages into the heart (42, 43). There was also a significant

elevation of both CD8+ T cells and natural killer (NK) cells, even though cell-mediated

immunity is relatively ineffective at responding to *T. brucei,* which is an extracellular parasite

(42, 43).

<u>Discussion</u>

T. brucei-associated cardiac disease is an underappreciated and understudied aspect of African

trypanosomiasis. Here we report a reproducible murine model of *T. brucei* infection optimized to

investigate acute and chronic *T. brucei*-associated cardiac disease. We demonstrate that this

model recapitulates key aspects of *T. brucei*-associated cardiac disease in humans and natural

hosts, including the presence of intracardiac parasites, myocarditis, EKG changes and elevated

NT-proBNP. Using this model, we learned that *T. brucei* infection results in parasite colonization

of the myocardial extracellular space and elevation of NT-proBNP (Figs 2-3). NT-proBNP

elevation was initially present in the context of a preserved ejection fraction, but the infection

eventually resulted in severe cardiac dysfunction as mice approached terminal endpoint (Figs 2-

3). At a histological level, we found that myocardial colonization by the parasite triggered a

severe myocarditis characterized by cardiomyocyte damage (Fig 5) and a multilineage

inflammatory infiltrate (Fig 6). These findings advance our understanding of the host-pathogen

interactions in the heart, suggesting that cardiac dysfunction associated with *T. brucei* infection

is an immune-mediated event, potentially providing valuable information to inform the

development of novel targeted therapies for animals and humans affected by African

trypanosomiases.

Animal models have been used to study *T. brucei* infection since the early 1900s, and the first

attempts to use a murine model to investigate *T. brucei*-associated cardiac disease were

published in the late 1970s (15, 29, 30) . In these historical murine studies, investigators

demonstrated cardiac lesions in infected mice and showed that immunosuppression was

associated with a reduction in the size and frequency of inflammatory cardiac lesions (29).

However, despite their seminal nature, these studies were limited in the scope of their

investigation, primarily focusing on histopathology and lacking more detailed functional or

immunological assessments. Moreover, the studies were performed using infection protocols and

parasite lines which were not standardized, thus making their methods challenging to replicate.

Our present work expands significantly on those historical studies as it presents a reproducible

method to generate cardiac damage and dysfunction in a murine model, demonstrably

recapitulating multiple aspects of natural disease.

We have found that mice infected with *T. brucei* initially exhibit elevation in NT-proBNP and

electrocardiographic abnormalities in the context of a preserved left ventricular ejection fraction.

At 28 dpi, when we initially assessed BNP, the mice had clear histological evidence of

myocarditis. NT-proBNP is elevated in response to an increase in intracardiac filling pressures

(34). In the context of a normal ejection fraction and cardiac pathology, BNP elevation is highly

suggestive of Heart Failure with Preserved Ejection fraction (HFpEF), a clinical condition

characterized by reduced cardiac performance and symptoms of heart failure despite a seemingly

normal cardiac contractility (44). However, BNP elevation in the context of a normal ejection

fraction has also been described in patients with acute myocarditis, likely as a result of stiffening

of cardiac walls triggered by the intracardiac immune response (45, 46). The potential overlap

between myocarditis with preserved ejection fraction and elevated BNP and HFpEF is an area

that has not been properly investigated so far in the clinical setting. However, taken together, our

findings suggest that colonization of the cardiac interstitium by *T. brucei* might result in a

stiffening of cardiac walls and elevation of cardiac pressures compatible with HFpEF, and that

the myocarditic immune response triggered by the parasites likely plays a key role in this

process. We hypothesize that as mice approach terminal endpoint, the cardiac pathology

becomes severe enough to cause a decrease in ejection fraction. However, more definite

conclusions require more detailed investigation into the functional impact of *T. brucei* infection

at multiple timepoints.

At 28 dpi, electrocardiographic abnormalities included marked bradycardia, increased QT

interval, grouped beating and dropped beats. These findings are in agreement with reports from

HAT patients, in which increased QT interval is the most common findings (16, 17). AV blocks

have also been reported in a subset of HAT patients, while autopsies have found inflammation of

the sinoatrial node, suggesting a disruption of the cardiac conduction system (47). Interestingly,

we did not observe premature ventricular contractions (PVCs), which have been observed in a rat

model of African trypanosomiasis (26). However, the relatively short recording period in our

study (1 minute per animal) may have hampered measurements of PVCs compared to the rat

model, in which 30-minute ECG recordings were analyzed (26).

We used flow cytometry to investigate the intracardiac immune milieu during *T. brucei*

infection. Previous studies in humans and large animal models have found that the major

immune effectors during *T. brucei* infection are B cells, CD4+ T cells, and macrophages, which

are all necessary for a protective anti-trypanosomal immune response (41). On the other hand,

CD8+ T cells and natural killer (NK) cells are thought to be mediators of pathological

inflammation in *T. brucei* infection, and their activation is associated with worse clinical

outcomes (41, 48, 49). In agreement with our histopathological finding of myocarditis, we found

that most intracardiac immune cell populations were elevated, especially CD4+ and CD8+ T

cells, macrophages, and NK cells. The most highly elevated macrophage subset was the

CCR2^{hi}MHC-II^{hi} population, which represents pro-inflammatory, monocyte-derived

macrophages (42, 43). Such pro-inflammatory macrophages are necessary for an effective

immune response to *T. brucei* (41). However, increased activation of anti-inflammatory,

alternatively-activated macrophages during chronic *T. brucei* infection is important for long-term

survival in experimentally infected animals, due to their ability to attenuate pathological

inflammation (47, 48). The large numbers of intracardiac CCR2^{hi}MHC-II^{hi} in our model suggests

that an imbalance between pro-inflammatory and anti-inflammatory macrophages could be a

contributor to myocarditis in our model. The elevation of CD8+ T cells and NK cells in the heart

are further suggestive of immunopathology as a contributor to cardiac damage. Increases in these

cell populations are suggestive of a response to intracellular antigens, although *T. brucei* is

entirely extracellular throughout its life cycle (48). Since CD8+ T cells are not a major feature of

the immune response to extracellular pathogens, we hypothesize that their elevation in our model

indicates a response to cardiac self-antigens, indicating an auto-immune component of the

observed myocarditis. This hypothesis is corroborated by other evidence suggesting that *T.*

brucei-associated organ damage has an auto-immune component. A recent study found that

meningeal B cells produce high-affinity autoantibodies during *T. brucei* infection, suggesting

that autoimmune responses may be a component of *T. brucei*-associated pathology in multiple

tissues(50). Clinically, cardiomyopathy patients in Cameroon have been found to have high

levels of both anti-trypanosomal and anti-heart antibodies, further indicating a connection

between *T. brucei* infection and autoimmune heart disease (18).

Furthermore, we hypothesize that immunopathology in the heart during *T. brucei* infection

follows a similar process to the pathogenesis of Chagas cardiomyopathy, caused by the New

World trypanosome *Trypanosoma cruzi.* The leading theory to explain the pathogenesis of

Chagas disease posits that parasite colonization of the heart induces inflammation and release of

self-antigens, resulting in a breakdown of self-tolerance and immune destruction of

cardiomyocytes (51). With chronicity, loss of cardiac muscle and replacement with scar tissue

causes a collapse in cardiac function, ultimately ending in the characteristic Chagas

cardiomyopathy (51). Our findings suggest that similar mechanisms might be at play during *T.*

brucei infection.

There was no increase in the B cell population in the heart. This is somewhat surprising, since

antibodies are the major mediators of immune clearance of *T. brucei*, and B cells are crucial for

systemic anti-trypanosomal immunity (41). However, our findings are consistent with other

tissue-specific flow cytometry experiments during *T. brucei* infection, which have identified no

changes in number of B cells in the lungs or adipose bodies of *T. brucei*-infected mice (14, 33).

In contrast, other models of myocarditis and heart failure show that intracardiac B cell

populations are increased and play a major pathologic role during viral myocarditis and

myocardial infarction (52–57). It is possible that B cell activation plays a role in *T. brucei*-

associated cardiac disease, even if there is no expansion of local populations.

We found limited evidence of a sex-specific difference in *T. brucei*-associated cardiac disease.

Male mice exhibited higher NT-proBNP at 33 dpi, higher numbers of intracardiac immune cells,

and more severe echocardiographic and ECG changes than females, although these differences

were not statistically significant. Our findings are in agreement with evidence that HAT is more

severe in men than women, as well as recent findings which indicate that T. brucei causes more

severe weight loss and adipose wasting in male rodents (58, 59). Our findings are also consistent

with data indicating that myocarditis is more common in men than women (60). Further work

will be needed to expand on these observations and dissect the mechanistic basis of sex

differences in HAT.

Much about *T. brucei*-associated cardiac disease remains unknown, and there are serious gaps in

our current knowledge, which this animal model could be used to investigate. Our findings raise

concerns about long-term, post-treatment consequences of *T. brucei* infection. Most HAT

patients are identified and treated, permanently clearing infection. However, the severity of the

myocarditis reported by ourselves and others suggests that heart dysfunction might be only

loosely connected with parasite presence. (20, 22–25, 29, 38, 39). If so, anti-parasitic treatment

would not fully alleviate the cardiac symptoms of HAT, since self-directed inflammation could

continue even with the removal of the inciting stimulus. Moreover, this suggests that

immunomodulatory therapies could be considered for treatment of both humans and cattle to

suppress inflammation that could contribute to persistent heart disease. There is a necessity to

investigate the long-term and post-treatment consequences of *T. brucei*-associated myocarditis to

determine whether HAT patients are at increased lifetime risk of heart failure. In addition, the

effect of *T. brucei* subspecies on cardiac disease remains to be investigated. Our mouse model

most closely recapitulates HAT due to *T. b. rhodesiense* infection. However, currently the

majority of human cases are caused by *T. b. gambiense*, which often causes a disease of a more

chronic character than *T. b. rhodesiense* (2). Although there is clinical evidence that all T. brucei

subspecies cause cardiac disease, the differences in cardiac manifestations between subspecies

are uncertain (16, 21, 61).

In summary, we describe the first reproducible, optimized, clinically relevant murine model of *T.*

brucei-associated cardiac disease, a feature of African trypanosomiasis that is often overlooked

in current textbooks and clinical resources. In the context of this model's characterization, we

produced evidence suggesting that *T. brucei*-associated cardiac damage is an immune-mediated

event. Future studies with this model will be instrumental in identifying potential treatments for

cardiac manifestations of African trypanosomiasis, a disease which continues to take an

extensive human and economic toll in large regions of the world.

Methods

Sex as a biological variable

For all experiments, data from both male and female mice were collected and analyzed.

Mouse and parasite strains and infection

Male and female C57Bl/6J mice (WT, strain# 000664 Jackson Laboratory) between 7-10 weeks

old were each infected by intravenous tail vein injection with ~5 pleiomorphic EATRO 1125

AnTat1.1E 90-13 *T. brucei brucei* parasites. To quantify parasitemia, blood was collected from

the tail vein every two 2 days starting at 4 days post-infection (dpi), and parasites were counted

via hemocytometer (Supplemental Figure 1). Blood (125 uL) was collected by a submandibular

bleed at terminal endpoint (28 or 33 dpi). Unless otherwise described, mice were euthanized at

terminal endpoint by intraperitoneal injection of ketamine/xylazine and perfused with 50 mL of

PBS-Glucose (0.055M D-glucose) with heparin to remove intravascular parasites.

For immunofluorescence experiments, mice were infected intravenously with ~5 AnTat1.1E

chimeric triple reporter *T. brucei* parasites which express tdTomato (35). Parasitemia was

measured as previously described. At 14 dpi, mice were sacrificed as previously described.

For flow cytometry experiments, mice were infected and parasitemia was measured as

previously described. Mice were sacrificed at 28 dpi using CO_2 euthanasia.

Echocardiography

At 28 dpi, echocardiography was performed on sedated mice by a blinded investigator using a

Vevo 2100 Ultrasound System (VisualSonics Inc.) as described previously (62).

At 33 dpi, echocardiography was performed on awake mice using a 40-MHz transducer

(VisualSonics Inc.) as described previously (63).

Electrocardiography

At 28 dpi, ECG was performed on unanesthetized mice as previously described (64). The entire

procedure was performed in approximately 1 minute per mouse using a PowerLab data

acquisition system (model ML866, ADInstruments, Colorado Springs, CO) and Animal Bio

Amp (model ML136, ADInstruments). ECG data were analyzed using LabChart software

(ADInstruments). The lead II tracings were used for analysis.

Heart digestion

Heart digestion was performed using a technique previously (65). Briefly, after sacrifice, mice

were perfused with 3 mL Hank's Balanced Salt Solution (HBSS). 60 mg of heart tissue was

placed in a 15 mL tube with 3 mL of HBSS. Digestion was performed for half an hour with 300

units of DNase I (Sigma-Aldrich), 625 units of collagenase II (Sigma-Aldrich), and 50 units of

hyaluronidase II (Sigma-Aldrich). After digestion, the tubes were centrifuged at 250xg for 5

minutes at 4 °C, the supernatant was removed, and the pellet was resuspended in 5 mL ACK

lysis buffer (Invitrogen) and incubated for 5 minutes at room temperature to remove red blood

cells. Suspended cells were passed through a 50 μm filter and then prepared for flow cytometry.

Flow cytometry

Flow cytometry was performed using a technique previously described (65). Cell suspensions

were labeled with fluorescently conjugated antibodies (Supplemental Table 1). FACS was

performed on an Aurora Spectral Flow Cytometer (Cytek) and analyses were performed using

FlowJo version 10.8.1 (Becton Dickinson).

NT-proBNP ELISA

At 28 and 33 dpi, plasma was collected from mice via cheek-bleeding. NT-proBNP plasma

levels were measured using the LS-F34395 Mouse NT-proBNP Sandwich ELISA Kit (LSBio)

according to the manufacturer's instructions.

Histopathology

For histopathology experiments, animals were infected and sacrificed at 28 dpi as previously

described. Hearts were placed in 10% neutral buffered formalin and fixed at room temperature

for 48 hours. The heart was cut in half longitudinally and sectioned by the Johns Hopkins

Oncology Tissue and Imaging Services Core. To evaluate inflammation, hearts were stained with

hematoxylin and eosin and imaged with 4x, 10x, and 40x objectives using a Nikon Eclipse E400

light microscope (Nikon). Cardiac lesions were graded by a blinded veterinary pathologist using

a previously established both the Dallas Criteria and a semiquantitative scoring system for

myocarditis in rodents (Table 2) (36, 37). Briefly, the severity and extent of lesions was graded

using a five-point system from 0 to 4. A zero score indicates no infiltration. A '1' score indicates

very limited focal distribution of myocardial lesions. A '2' score indicates multiple lesions. A '3'

score indicates multiple regions with confluence and extensive necrosis. A '4' score indicates

lesions affecting most of the observed tissue. Images were captured using an Excelis 4K camera

(Unitron) and processed using ImageJ v1.53 image analysis software (NIH). All methods

followed published guidelines for experimental pathology (66). To evaluate cardiac fibrosis,

additional longitudinal sections of hearts were stained with Masson's Trichrome and scanned at

200x magnification using an Axios Z.1 scanner (Zeiss). The percentage of collagen in each heart

section was quantified using the AFAT (40).

Immunofluorescence

For immunofluorescence experiments, 8-week-old female C57Bl/6J mice were infected

intravenously with 5 triple marker *T. brucei* parasites (35). At 14 dpi, mice were euthanized and

perfused as previously described. Heart was collected and fixed in 4% paraformaldehyde in PBS

for 12 hours at 4 C. Post-fixation, tissues were sectioned longitudinally, frozen embedded in

O.C.T. Compound (Tissue-Tek), and cut using cryostat microtome into 10 μm sections by the

Johns Hopkins Oncology Tissue and Imaging Services Core.

The following antibodies were applied to sections: rat anti-mouse CD-31 (SCBT) followed by

goat anti-rat Alexa Fluor 488 (CST). Coverslips were mounted using ProLong Gold (Life

technologies). Tissues were imaged with 4x, 10x and 20x objectives using a Nikon Eclipse 90i

fluorescence microscope (Nikon) and X-Cite 120 fluorescent lamp (Excelitas) with an ORCA-

ER digital CCD camera (Hammamatsu) and ImageJ v1.53 image analysis software (NIH). Image

collection and analysis followed published guidelines for rigor and reproducibility in

immunofluorescence (67).

Statistics

Unless otherwise described, figures were generated and statistical analyses were performed using

the Tidyverse package in R version 4.2.3 (68).

Study approval

Animal experiments were performed with the approval of the IACUC of Johns Hopkins

University under protocol MO22H163

Data availability

Values for all data points in graphs are reported in the Supporting Data Values file.

<u>Author contributions</u>

NPC, MRM and LA conceptualized the study. NPC, MDC, AKB, PS, and KG conducted

experiments and acquired the data. NPC, MDC, AL, PS, AB, and KG analyzed the data. KG,

LA, and MRM provided resources. NC wrote the original draft of the manuscript. NPC, AKB,

LA, and MRM reviewed and edited the manuscript. MRM acquired funding. MRM and LA

supervised the study.

<u>Acknowledgements</u>

We would like to kindly thank Brice Rotureau for providing us with the chimeric triple reporter

cell line.

The Aurora Flow Cytometer used in this study was funded by NIH Grant S10OD026859. We

acknowledge the support of the JHU Ross Flow Cytometry Core.

Echocardiograms were performed by Nadan Wang of the JHU Small Animal Cardiovascular

Phenotyping and Animal Model Core.

Histology support was provided by the JHU Oncology Tissue Services Core, funded by JHU

Cancer Center Core Grant P30 CA006973.

This project was supported by the Johns Hopkins Specialized Center for Research Excellence in

Sex Differences (U54AG062333) and The Foundation for Gender-Specific Medicine. NPC was

supported by T32 OD011089.

References

1. Holmes P. Tsetse-transmitted trypanosomes - Their biology, disease impact and control
[preprint]. *J Invertebr Pathol*. 2013;112(SUPPL.1). https://doi.org/10.1016/j.jip.2012.07.014.
2. Kennedy PGE, Rodgers J. Clinical and neuropathogenetic aspects of human African
trypanosomiasis [preprint]. *Front Immunol*. 2019;10(JAN).
https://doi.org/10.3389/fimmu.2019.00039.
3. Spickler AR. African Animal Trypanosomiasis [Internet]. *IO State University Center for Food*
Security and Public Health. 2018.
4. Balmer O, et al. Phylogeography and taxonomy of Trypanosoma brucei. *PLoS Negl Trop Dis*.
2011;5(2):e961.
5. Franco JR, et al. The elimination of human African trypanosomiasis: Achievements in relation
to WHO road map targets for 2020. *PLoS Negl Trop Dis*. 2022;16(1).
https://doi.org/10.1371/JOURNAL.PNTD.0010047.
6. Funk S, et al. Identifying Transmission Cycles at the Human-Animal Interface: The Role of
Animal Reservoirs in Maintaining Gambiense Human African Trypanosomiasis. *PLoS Comput*
Biol. 2013;9(1). https://doi.org/10.1371/journal.pcbi.1002855.
7. Njiokou F, et al. Wild fauna as a probable animal reservoir for Trypanosoma brucei gambiense
in Cameroon. *Infection, Genetics and Evolution*. 2006;6(2):147–153.
8. Njiokou F, et al. Domestic animals as potential reservoir hosts of Trypanosoma brucei
gambiense in sleeping sickness foci in Cameroon. *Parasite*. 2010;17(1):61–6.
9. Abro Z, et al. The potential economic benefits of controlling trypanosomiasis using waterbuck
repellent blend in sub-Saharan Africa. *PLoS One*. 2021;16(7 July).
https://doi.org/10.1371/journal.pone.0254558.
10. Trindade S, et al. Trypanosoma brucei Parasites Occupy and Functionally Adapt to the
Adipose Tissue in Mice. *Cell Host Microbe*. 2016;19(6):837–848.
11. Poltera AA. Pathology of human African trypanosomiasis with reference to experimental
African trypanosomiasis and infections of the central nervous system. *Br Med Bull*.
1985;41(2):169–174.
12. Crilly NP, Mugnier MR. Thinking outside the blood: Perspectives on tissue-resident
Trypanosoma brucei [preprint]. *PLoS Pathog*. 2021;17(9).
https://doi.org/10.1371/journal.ppat.1009866.
13. De Niz M, et al. Organotypic endothelial adhesion molecules are key for Trypanosoma
brucei tropism and virulence. *Cell Rep*. 2021;36(12).
https://doi.org/10.1016/j.celrep.2021.109741.
14. Machado H, et al. Trypanosoma brucei triggers a broad immune response in the adipose
tissue. *PLoS Pathog*. 2021;17(9). https://doi.org/10.1371/journal.ppat.1009933.
15. Wolbach SB, Binger CA. A Contribution to the Parasitology of Trypanosomiasis. *J Med Res*.
1912;27(1):83-108.13.
16. Blum JA, et al. Sleeping hearts: The role of the heart in sleeping sickness (human African
trypanosomiasis). *Tropical Medicine and International Health*. 2007;12(12):1422–1432.
17. Blum JA, et al. Cardiac alterations in human African trypanosomiasis (T.b. gambionse) with
respect to the disease stage and antiparasitic treatment. *PLoS Negl Trop Dis*. 2009;3(2).
https://doi.org/10.1371/journal.pntd.0000383.

18. Tsala Mbala P, et al. [Functional and immunologic involvement in human African
trypanosomiasis caused by Trypanosoma gambiense]. *Bull Soc Pathol Exot Filiales*. 1988;81(3
BIS):490–501.
19. Hawking F, Greenfield JG. Two autopsies on rhodesiense sleeping sickness; Visceral lesions
and significance of changes in cerebrospinal fluid. *Trans R Soc Trop Med Hyg*. 1941;35(3):155–
156.
20. Poltera AA, Cox JN, Owor R. Pancarditis affecting the conducting system and all valves in
human African trypanosomiasis. *Heart*. 1976;38(8):827–837.
21. Koten JW, de Raadt P. Myocarditis in Trypanosoma rhodesiense infections. *Trans R Soc
Trop Med Hyg*. 1969;63(4):485–489.
22. Poltera AA, et al. Trypanosomal cardiac valvulitis in vervet monkeys. *Tropical Medicine and
Parasitology*. 1985;36(2):77–80.
23. Ikede BO, Losos GJ. Pathology of the disease in sheep produced experimentally by
Trypanosoma brucei. *Vet Pathol*. 1972;9(4):278–289.
24. Ikede BO, Losos GJ. Pathological changes in cattle infected with Trypanosoma brucei. *Vet
Pathol*. 1972;9(4):272–277.
25. Morrison WI, et al. The pathogenesis of experimentally induced Trypanosoma brucei
infection in the dog I. Tissue and organ damage. *American Journal of Pathology*.
1981;102(2):168–181.
26. McCarroll CS, et al. A Pre-clinical Animal Model of Trypanosoma brucei Infection
Demonstrating Cardiac Dysfunction. *PLoS Negl Trop Dis*. 2015;9(5).
https://doi.org/10.1371/journal.pntd.0003811.
27. Elliott EB, et al. Trypanosoma brucei cathepsin-L increases arrhythmogenic sarcoplasmic
reticulum-mediated calcium release in rat cardiomyocytes. *Cardiovasc Res*. 2013;100(2):325–
335.
28. Wildner G. Are rats more human than mice? *Immunobiology*. 2019;224(1):172–176.
29. Poltera AA, Hochmann A, Lambert PH. A model for cardiopathy induced by Trypanosoma
brucei brucei in mice. A histologic and immunopathologic study. *American Journal of
Pathology*. 1980;99(2):325–352.
30. Galvao-Castro B, Hochmann A, Lambert PH. The role of the host immune response in the
development of tissue lesions associated with African trypanosomiasis in mice. *Clin Exp
Immunol*. 1978;33(1):12–24.
31. Beaver AK, et al. Extravascular spaces are the primary reservoir of antigenic diversity in
Trypanosoma brucei infection. *bioRxiv*. 2023;2022.06.27.497797.
32. Spickler AR. African Animal Trypanosomiasis [Internet]. 2018;2020.
http://www.cfsph.iastate.edu/DiseaseInfo/factsheets.php.
33. Mabille D, et al. Impact of pulmonary African trypanosomes on the immunology and
function of the lung. *Nat Commun*. 2022;13(1). https://doi.org/10.1038/s41467-022-34757-w.
34. Yancy CW, et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA
Guideline for the Management of Heart Failure: A Report of the American College of
Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the
Heart Failure Society of America. *J Am Coll Cardiol*. 2017;70(6):776–803.
35. Calvo-Alvarez E, et al. A new chimeric triple reporter fusion protein as a tool for in vitro and
in vivo multimodal imaging to monitor the development of African trypanosomes and
Leishmania parasites. *Infection, Genetics and Evolution*. 2018;63:391–403.
36. Thomas Aretz H. Myocarditis: The Dallas criteria. *Hum Pathol*. 1987;18(6):619–624.

37. Aly M, et al. Complex genetic control of host susceptibility to coxsackievirus B3-induced
myocarditis. *Genes Immun*. 2007;8(3):193–204.
38. Dupont B, Charmot G, Lapresle C. [Trypanosomiasis presenting with trypanids and
complicated by myopericarditis (author's transl)]. *Nouv Presse Med*. 1979;8(19):1579–81.
39. Büngener W, Mehlitz D. [Experimental Trypanosoma infections in Cameroon dwarf goats:
histopathological observations (author's transl)]. *Tropenmed Parasitol*. 1976;27(4):405–10.
40. Gratz D, et al. Computational tools for automated histological image analysis and
quantification in cardiac tissue. *MethodsX*. 2020;7:22–34.
41. Onyilagha C, Uzonna JE. Host Immune Responses and Immune Evasion Strategies in
African Trypanosomiasis [preprint]. *Front Immunol*. 2019;10.
https://doi.org/10.3389/fimmu.2019.02738.
42. Dick SA, et al. Self-renewing resident cardiac macrophages limit adverse remodeling
following myocardial infarction. *Nat Immunol*. 2019;20(1):29–39.
43. Cohen CD, et al. Myocardial Immune Cells: The Basis of Cardiac Immunology. *The Journal
of Immunology*. 2023;210(9):1198–1207.
44. Ranek M, et al. Pathophysiology of heart failure and an overview of therapies. In: Buja L,
Butany J, eds. *Cardiovascular Pathology*. London: Academic Press; .
45. Sara B, et al. Are high NT-proBNP levels more related to inflammation than to left
ventricular systolic dysfunction in acute myocarditis? *Eur Heart J Acute Cardiovasc Care*.
2021;10(Supplement_1). https://doi.org/10.1093/ehjacc/zuab020.189.
46. Moady G, Perlmutter S, Atar S. The Prognostic Value of Natriuretic Peptides in Stable
Patients with Suspected Acute Myocarditis: A Retrospective Study. *J Clin Med*.
2022;11(9):2472.
47. Bertrand E, et al. La symptomatologie cardio-vasculaire dans la trypanosomiase humaine
Africaine a Trypanosoma gambiense.
48. Liu G, et al. Distinct contributions of CD4+ and CD8+ T cells to pathogenesis of
trypanosoma brucei infection in the context of gamma interferon and interleukin-10. *Infect
Immun*. 2015;83(7):2785–2795.
49. Frenkel D, et al. Trypanosoma brucei Co-opts NK Cells to Kill Splenic B2 B Cells. *PLoS
Pathog*. 2016;12(7). https://doi.org/10.1371/journal.ppat.1005733.
50. Quintana JF, et al. The murine meninges acquire lymphoid tissue properties and harbour
autoreactive B cells during chronic Trypanosoma brucei infection. *bioRxiv*.
2023;2023.04.30.538710.
51. Bonney KM, Engman DM. Autoimmune pathogenesis of chagas heart disease: Looking
back, looking ahead [preprint]. *American Journal of Pathology*. 2015;185(6):1537–1547.
52. Zouggari Y, et al. B lymphocytes trigger monocyte mobilization and impair heart function
after acute myocardial infarction. *Nat Med*. 2013;19(10):1273–1280.
53. Tschöpe C, et al. Myocarditis and inflammatory cardiomyopathy: current evidence and future
directions [preprint]. *Nat Rev Cardiol*. 2021;18(3):169–193.
54. Tschöpe C, et al. Targeting CD20+ B-lymphocytes in inflammatory dilated cardiomyopathy
with rituximab improves clinical course: a case series. *Eur Heart J Case Rep*. 2019;3(3).
https://doi.org/10.1093/ehjcr/ytz131.
55. Mo F, et al. Are activated B cells involved in the process of myocardial fibrosis after acute
myocardial infarction? An in vivo experiment. *BMC Cardiovasc Disord*. 2021;21(1).
https://doi.org/10.1186/s12872-020-01775-9.

56. Bermea K, et al. The Role of B Cells in Cardiomyopathy and Heart Failure [preprint]. *Curr*
Cardiol Rep. 2022;24(8):935–946.
57. Adamo L, et al. Modulation of subsets of cardiac B lymphocytes improves cardiac function
after acute injury. *JCI Insight*. 2018;3(11). https://doi.org/10.1172/jci.insight.120137.
58. Sinton MC, et al. IL-17 signalling is critical for controlling subcutaneous adipose tissue
dynamics and parasite burden during chronic murine Trypanosoma brucei infection. *Nat*
Commun. 2023;14(1). https://doi.org/10.1038/s41467-023-42918-8.
59. Pépin J, Mpia B, Iloasebe M. Trypanosoma brucei gambiense African trypanosomiasis:
Differences between men and women in severity of disease and response to treatment. *Trans R*
Soc Trop Med Hyg. 2002;96(4):421–426.
60. Fairweather DL, et al. Sex and gender differences in myocarditis and dilated
cardiomyopathy: An update [preprint]. *Front Cardiovasc Med*. 2023;10.
https://doi.org/10.3389/fcvm.2023.1129348.
61. Losos GJ, Ikede BO. Review of pathology of diseases in domestic and laboratory animals
caused by Trypanosoma congolense, T. vivax, T. brucei, T. rhodesiense and T. gambiense. *Vet*
Pathol. 1972;Sup. 9. https://doi.org/10.1177/030098587200901s01.
62. Fitch ML, et al. Cadmium exposure induces a sex-dependent decline in left ventricular
cardiac function. *Life Sci*. 2023;324. https://doi.org/10.1016/j.lfs.2023.121712.
63. Sørensen LL, et al. Echocardiographic characterization of a murine model of hypertrophic
obstructive cardiomyopathy induced by cardiac-specific overexpression of epidermal growth
factor receptor 2. *Comp Med*. 2016;66(4):268–277.
64. Sysa-Shah P, et al. Electrocardiographic characterization of cardiac hypertrophy in mice that
overexpress the ErbB2 receptor tyrosine kinase. *Comp Med*. 2015;65(4):295–307.
65. Bermea KC, Rousseau ST, Adamo L. Flow Cytometry-Based Quantification and Analysis of
Myocardial B-cells. *Journal of Visualized Experiments*. 2022;2022(186).
https://doi.org/10.3791/64344.
66. Scudamore CL, et al. Recommendations for minimum information for publication of
experimental pathology data: MINPEPA guidelines [preprint]. *Journal of Pathology*.
2016;238(2):359–367.
67. Lee JY, Kitaoka M. A beginner's guide to rigor and reproducibility in fluorescence imaging
experiments [preprint]. *Mol Biol Cell*. 2018;29(13):1519–1525.
68. Wickham H, et al. Welcome to the Tidyverse. *J Open Source Softw*. 2019;4(43):1686.

718   **<u>Figures</u>**

720   **<u>Figure 1</u>**. *T. brucei* infection causes waxing and waning waves of parasitemia and massive

721   splenomegaly, with terminal endpoint reached within 50 dpi

722   A. Quantification of median and individual parasitemia in a representative experiment.

723   Parasitemia first becomes detectable at 4-6 dpi.

724   B. Weight of spleens of infected mice at 28 dpi(n=6) compared to uninfected, age-matched

725   controls. Infected mice exhibit significant splenomegaly (Student's t-test, p=3.7E-06).

726   C. Kaplan-Meier survival curve of mice infected with T. brucei vs. uninfected control

727   animals. Infected mice have a median survival time=39 dpi (p=4.32E-05).

<u>Figure 2</u>. *T. brucei* infection causes elevated NT-proBNP, electrocardiographic abnormalities

and heart failure with preserved ejection fraction

    A. Measurement of plasma NT-proBNP at 28 dpi. Plasma NT-proBNP is significantly

        elevated at 28 dpi (n=12) compared to age-matched uninfected controls (Student's t-test,

        p=0.00037).

    B. Echocardiographic measurements at 28 dpi. At 28 dpi, infected mice (n=12) exhibit

        significantly increased ejection fraction compared to uninfected age-matched controls, as

        measured by sedated echocardiography (Student's t-test, p=0.0029).

739   C. Electrocardiographic measurements at 28 dpi. At 28 dpi, infected mice (n=10) exhibit

740   significantly decreased heart rate (HR, Student's t-test, p= 0.00011), increased QRS

741   interval, (Student's t-test, p=0.00281), increased QT interval (Student's t-test,

742   p=0.00493), and increased QTc (Student's t-test, p=0.01294) compared to uninfected

743   age-matched controls.

<u>Figure 3.</u> T. brucei infection causes elevated NT-proBNP and decreased heart function at

terminal endpoint (33 dpi).

A. Measurement of plasma NT-proBNP at 33 dpi. Plasma NT-proBNP is significantly

elevated at 33 dpi in infected mice (n=6) compared to uninfected age-matched controls

(Student's t-test, p=2.8e-05).

B. Echocardiographic measurements at 33 dpi. At 33 dpi, infected mice (n=6) exhibit

significantly decreased ejection fraction compared to uninfected age-matched controls, as

measured by awake echocardiography (Student's t-test, p=0.011)

Hoechst (nuclei)   **Fluor 488 (CD31)**

TdTomato (parasites)   **Merged**

Figure 4. *T. brucei* parasites occupy extravascular spaces in the heart

Representative immunofluorescence microphotographs of the cardiac ventricle at 14 dpi at 20x

magnification. *T. brucei* parasites (tdTomato—red) localize separately from CD31-lined vascular

structures (Alexa Fluor 488—green), confirming the extravascular localization of parasites.

Nuclei are stained with Hoechst (blue).

Figure 5. *T. brucei* infection causes myocarditis at 28 dpi

A. Representative brightfield microphotographs of the heart of infected mice at 28 dpi,

compared to uninfected age-matched controls at 10x magnification. Inflammation is

present in all layers and regions of the heart and is composed largely of mononuclear

cells, primarily lymphocytes, plasma cells, and histiocytes. Inflammation is most severe

at the atrioventricular junction.

B. Histological grading of myocarditis (n=6) at 28 dpi, compared to equal numbers of age-

matched controls. The average grade at 28 dpi is '2', indicating multifocal moderate

myocarditis without extensive associated necrosis. Histological grade is significantly

higher in infected than uninfected mice (Student's t test, p=1.5e-8).

C. Measurement of the percentage of collagen in longitudinal sections of heart (n=6) at 28

dpi, compared to equal numbers of age-matched controls. There is no significant

difference between infected and control mice (Student's t test, p=0.31).

Figure 6. The intracardiac immune cell population during *T. brucei* infection is characteristic of immune-mediated myocarditis

A. Flow cytometric quantification of CD45+ cells per mg. At 28 dpi, CD45+ cells are markedly increased in the hearts of infected mice (n=12) compared to uninfected age-matched controls, supportive of myocarditis (Student's t-test, p= 4.6e-7).

B. Flow cytometric quantification of myeloid and lymphoid cell populations. Most immune cell populations are significantly increased in the hearts of infected mice at 28 dpi. Statistical comparisons were made using One-way Anova. *p<0.05, **p<0.01, ***p<0.001.

C. Visualization of magnitude of changes in immune cell populations using the log2-fold change in cells/mg compared to uninfected age-matched control mice calculated based on the average cells/mg in the hearts of uninfected control mice.

813　**<u>Tables</u>**

815　**Table 1:** Comparison of the mean values of all ECG parameters between infected animals and

816　age-matched controls at 28 dpi (NS, P>0.05. *, P ≤0.05. **, P ≤0.01. ***,P ≤0.001)

	Control	Infected	P value	Significance
RR Interval (ms)	104.98	166.77	0.00111	**
Heart Rate (BPM)	616.18	394.40	0.00011	***
QRS Interval (ms)	7.82	10.34	0.00281	**
QT Interval (ms)	39.43	58.80	0.00493	***
QTc (ms)	39.17	48.66	0.01294	*
JT Interval (ms)	31.77	48.25	0.00758	**
Tpeak Tend Interval (ms)	28.97	42.24	0.00985	**
Q Amplitude (mV)	0.01	0.02	0.42396	NS
R Amplitude (mV)	0.89	0.78	0.19872	NS
S Amplitude (mV)	-0.52	-0.30	0.00438	**
ST Height (mV)	0.10	0.07	0.32718	NS
T Amplitude (mV)	0.19	0.11	0.00670	**

817
818　NS, P>0.05. *, P ≤0.05. **, P ≤0.01. ***,P ≤0.001
819
820
821
822　**Table 2**: Histological grading criteria
823

Grade	Description
0	No inflammatory infiltrates
1	Limited focal distribution of myocardial lesions
2	Multiple lesions with inflammatory infiltrates and some confluence
3	Multiple lesions with confluence and extensive necrosis
4	Lesions affecting most of the observed tissue

834   **<u>Supplemental Materials</u>**
835
836   Supplemental data is available in the associated Supplemental Data document
837
838   Supporting data values are available in the associated Supporting Data Values document