Aperiodic component of EEG power spectrum and cognitive performance in aging: the role of education

Sonia Montemurro[1]*, Daniel Borek[2]*, Daniele Marinazzo[2], Sara Zago[1], Fabio Masina[1], Nicola Filippini[1], Giorgio Arcara[1]

[1]*IRCCS San Camillo Hospital, Venice (Italy)*
[2]*Department of Data-Analysis, Faculty of Psychology and Educational Sciences, Ghent University, Belgium*

*These authors share the first authorship

Abstract

Aging is associated with changes in the oscillatory (periodic) brain activity in the alpha band (8-12 Hz), as measured with resting-state EEG (rsEEG), and it is characterized by significantly lower alpha frequency and power. Aging can influence the aperiodic component of the power spectrum: at a higher age, the slope flattens, and it may show a significant relationship with cognitive performance. However, it is unclear whether education, a cognitive reserve proxy known for its modulatory role on cognition, may influences such relationship. N=179 healthy participants from the LEMON dataset (Babayan et al., 2019) were classified into three groups based on age and education: young adults (N=123) 20-35 years old with high education, older adults 60-77 years old with high (N=24) and low (N=32) education. Eyes-closed rsEEG power spectrum was decomposed at the occipital level with specparam method. Individual Alpha Peak Frequency, exponent, and offset were analyzed in their relationship with cognitive abilities.

Results show lower IAPF, exponent, and offset in older as compared to younger adults. Furthermore, visual attention and working memory were differently predicted by the aperiodic rsEEG component depending on educational levels: in older adults with higher education, increasing exponent predicted slower processing speed and less working memory capacity, with an opposite trend in those with lower education.

Although further investigations are necessary, this study shows a potential modulatory role of education and other cognitive reserve proxies in the relationship between aperiodic rsEEG and cognition in aging.

Keywords: education, processing speed, working memory

1. INTRODUCTION

As we age, many changes occur in individuals' behavior and cognition, worsening in memory (Nyberg et al., 2012), attention span (Gazzaley et al., 2005), executive functions (Buckner, 2004), and processing speed (Salthouse, 2010), reflecting just a few of the consequences of the natural aging process. These behavioral changes are accompanied by (and correlated with) changes in the brain's structural anatomy (Habes et al., 2016; Taki et al., 2011), metabolism (Camandola & Mattson, 2017) and functionality (Rossini et al., 2007) which produce a significant effect on its neurophysiological activity (Rossini et al., 2007). Electroencephalogram (EEG) studies on aging have shown changes in neural oscillatory activity, especially in the alpha band (8-12 Hz) (Babiloni et al., 2006; Michels et al., 2013; Scally et al., 2018). Researchers have reported that older adults display slower alpha oscillatory activity and lower alpha power than their younger counterparts (Klimesch, 1999; Rossini et al., 2007; Sghirripa et al., 2020). Moreover, individual alpha peak frequency (IAPF), i.e., the frequency where EEG activity exhibits the maximum power in the alpha range, tends to decrease from adulthood to midlife (Michels et al., 2013; Scally et al., 2018). In most studies, EEG activity in specific frequency bands has been traditionally measured as the average of the power in the frequency bands of interest as calculated from the power spectrum (Donoghue et al., 2020). However, this approach (and subsequent interpretation of the results) has been challenged by a renewed interest in the non-oscillatory, aperiodic component of the EEG signal.

The aperiodic component does not necessarily entail specific frequency bands; it exhibits a 1/f-like distribution in the log-log space of a Power Spectrum Density (PSD), meaning its power exponentially decreases as frequency increases.

Aperiodic activity can be parametrized by values of the exponent, which describe the negativity of the power spectrum slope and the offset, the broadband shift of power across frequencies (Donoghue et al., 2020). Importantly, change in the spectrum's aperiodic component may occur without changes in the oscillatory components and may affect the power values calculated for each frequency. For

example, a change in aperiodic slope may influence the power values in the alpha band as calculated from the power spectrum, when no change in alpha's oscillatory component occurred. This may lead to spurious results and wrong interpretations and highlight the importance of taking into account the aperiodic component when interpreting power spectrum data (see Donoghue et al., 2020 for further explanations and visualizations of the effect of the aperiodic component on the oscillatory component; see also Schaworonkow & Voytek, 2021).

For a long time, researchers ignored the aperiodic component of brain activity, considering it to be merely background noise; however, recently, this component has gained attention not only for its methodological impact but also because of its potential role as a marker of neurophysiological mechanisms underlying cognition. For instance, the aperiodic offset has been associated with neuronal population spiking (Manning et al., 2009; Miller et al., 2013) and blood-oxygen-level-dependent (BOLD) signal of functional magnetic resonance (fMRI) (Winawer et al., 2013). Meanwhile, the exponent has been linked to changes in cortical excitation/inhibition balance (Gao et al., 2017), which means that increased power in higher frequency (associated with increased excitation at the cerebral level) produced a flattening of the PSD; thus a diminished exponent; the exponent has also been correlated to local neural tissue properties (Bédard & Destexhe, 2009; Merkin et al., 2021). In light of the investigations mentioned above, narrowly associating 1/f characteristics with specific cerebral function, alternative explanations should be acknowledged. A possible interpretation of the functional meaning of the aperiodic component posits that this variable indicates an underlying self-organized critical system (Bak et al., 1987). Within this theoretical framework, the critical system is considered as the optimal state of brain's functionality (O'Byrne & Jerbi, 2022) where large-scale reorganization occurs quickly in response to stimuli, reflecting the brain's adaptation to changes in the endogenous and exogenous environment (Bullmore & Sporns, 2009). Aperiodic activity may arise from spatial integration of asynchronous spiking of neural populations (Miller et al., 2009). Therefore, a reduction of the spectral exponent suggests functional neural decoupling.

In the context of aging, a first study from Voytek and collaborators (2015) showed that the aperiodic slope of EEG and electrocorticography (ECoG) spectra flattened in a group of older people compared with a younger one, with decreased power between 8 to 14 Hz and increased power between 14 and 25 Hz. The changes in EEG spectral slope were also correlated with age differences in working memory performance. Interestingly, the aperiodic exponent appeared to mediate this relationship, suggesting that the slope effect alone could account for behavioral differences between older and young adults.

Voytek and colleagues explained these results with the "Neural Noise Hypothesis". Initially proposed by Voytek and collaborators (2015), this theory suggests that as people age, there is an increase in spontaneous desynchronized neural activity, resulting in a decreased fidelity of neural communication and a flatter power spectrum. Recent studies have replicated Voytek findings and added new insights to the relationship between aging and aperiodic spectral components, highlighting their impact on cognitive performance. Flatter slopes in older adults were linked to poorer performance during spatial attention tasks (Tran et al., 2020; Waschke et al., 2017) and short-term memory tasks (Thuwal et al., 2021). Recent work from Pathania and collaborators (2021) has associated the flattening of the aperiodic slope in frontal regions with worse performance on tasks involving processing speed and executive functions. Moreover, Tran and collaborators (2020) measured changes of the aperiodic spectra at the baseline period in younger and older adults. They examined to what extent 1/f like exponent was related to alpha trial-by-trial consistency, during a spatial discrimination task. The authors considered these signal properties as measures of "neural noise". Interestingly, they found that older adults with the highest baseline noise levels also had the worse alpha trial-by-trial consistency, suggesting that age-related increases in baseline noise might diminish sensory processing and cognitive performance.

Understanding the impact of neural noise could unveil new perspectives in capturing the effects of aging on neurophysiological functioning. Critically, studies on the aperiodic component of the

spectrum and its functional role have neglected the possible modulatory role of "cognitive reserve" in modulating the effects on cognition in aging. Cognitive reserve is a concept proposed by Stern (2002; 2023), and it describes the capacity of the brain to actively cope with age-related or pathological changes by employing pre-existing resources. Such reserve accumulates since early life through exposure to cognitively stimulating life (Lojo-Seoane et al., 2018; Mondini et al., 2022; Stern et al., 1994). One of the most widely recognized variables associated with the level of cognitive reserve is education, typically operationalized as a numerical variable, indicating years of successful education or as an educational level operationalized as an ordered factor (e.g., high school, University etc.). Education has been largely used to examine the aging population's potential compensatory processes (e.g., Lövdén et al., 2020; Montemurro et al., 2021). While education modulates cognitive performance (i.e., the higher the education, the better cognitive performance), its impact on the intricate relationship between neural activity in the aging brain and cognitive capacity remains uncertain. Aperiodic activity, conceived as a possible pattern of neural communication impairment, is associated with a decline in cognitive performance in older adults. Critically, although education can be expected to favor the preservation/maintenance of neurobehavioral functionality in aging (Cabeza et al., 2018), how this is reflected at the level of aperiodic brain activity is not clear yet. With reference to the most recently updated framework targeting aging compensatory mechanisms, a specific set of variables were combined to investigate: i) possible life course-related neural change (i.e., aperiodic values), ii) expected age-related cognitive changes (test scores), iii) and education as a variable expected to moderate the association between i) and ii) (see Stern et al., 2023 for further information). Based on such a framework, older adults with higher education were expected to preserve a more youth-like profile than older adults with lower education.

2. METHODS

2.1. Participants and Materials

All participants included in this study were taken from the "Leipzig Study for Mind-Body-Emotion Interactions" (LEMON; Babayan et al., 2019). The final sample consisted of N=179 individuals. Socio-demographic information like age and education was shared in bins by Babayan and colleagues (2019), not continuous. Participants were grouped based on age (older vs. younger adults) and education (high vs. lower).Participants included in the young group (N=123) aged 20-35 years and had high education levels (12 years of *lyceum/gymnasium*), whereas the group of older adults (N=56), age range 60-77 years, was divided in two groups: one with high education (12 years of *lyceum/*gymnasium, N=24) and the other with low education (10 years of *technical high school/Realschule*, N=32). Participants with no availability of EEG data and those who were indicated as with alcohol abuse or dyslexia problems were not included in the final data sample. A small subgroup of young adults with low education was not included in the sample according to the study purpose (N=7); one participant resulted as outlier on both visual inspection of residuals and Cook's threshold and it was removed. Data collection was performed upon the Declaration of Helsinki and approved by the local ethics committee (Babayan et al., 2019) ; all participants provided written informed consent before data acquisition for the study.

2.2. Cognitive Assessment

Processing Speed and Memory capacity were investigated in relationship with periodic and aperiodic EEG components.

- Processing Speed included alertness and visual attention and it was assessed using two tasks: the Test of Attentional Performance (TAP_A, Zimmermann & Fimm, 2012) and the Trail Making Test (TMT, Reitan, 1992). The former estimated alertness: i.e., participants were asked to respond, as fast as possible, to the appearance of a visual stimulus on the screen. The TMT estimated visual attention w,

i.e., participants were asked to connect as fast as possible a series of visual stimuli, alternatively with a definition order: numerical and alphabetical orders.

- Memory included working and delayed memory tasks; it was assessed with a working memory task (WM_TAP, Zimmermann & Fimm, 2012) and the California Verbal Learning Task (CVLT, Niemann et al., 2008). For the working memory task, participants had to simultaneously provide a response only when a given stimulus was equal to the second last one perceived in the series, while keeping track of a series of different stimuli. In the delayed memory task, participants were asked to retain and correctly recall a series of 16 words belonging to their vocabulary.

2.3 Neural correlates

2.3.1. Gray Matter Volume

A 3 Tesla scanner (MAGNETOM Verio, Siemens Healthcare GmbH, Erlangen, Germany) with a 32-channel head coil was used to conduct Magnetic Resonance Imaging (MRI); (Babayan et al., 2019). The pre-processing pipeline included a series of steps: a) re-orientating images to the standard (MNI) template, b) bias field correction, c) registration to the MNI template using both linear (FLIRT) and nonlinear (FNIRT) registration tools, and d) brain extraction. Brain tissues were segmented using FMRIB's Automated Segmentation Tool (FAST) that allowed extracting measures of total Gray Matter, White Matter and Cerebrospinal Fluid. Brain tissues were visually inspected by a trained neuroscientist (NF) to ensure an accurate segmentation.

2.3.2. EEG preprocessing and source reconstruction

A 16-min rs-EEG was recorded with a BrainAmp MR plus amplifier in an electrically shielded and sound-attenuated EEG booth using 62-channel (61 scalp electrodes plus 1 electrode recording the VEOG below the right eye) active ActiCAP electrodes (Brain Products GmbH, Gilching, Germany; international standard 10–20 localization system, and referenced to FCz). EEG was recorded with a

band-pass filter between 0.015 Hz and 1 kHz and digitized with a sampling rate of 2500 Hz. Raw EEG data were down-sampled from 2500 Hz to 250 Hz and band-pass filtered within 1-45 Hz. Outlier channels were rejected after visual inspection for frequent jumps/shifts in voltage and poor signal quality. Data intervals containing extreme peak-to-peak deflections or large bursts of high-frequency activity were identified by visual inspection and removed. Independent component analysis (ICA) was performed using the Infomax algorithm (runica function from MATLAB). On pre-processed files, source reconstruction was run by using a standard head model. A 3-shell boundary element model was constructed via Brainstorm (Tadel et al., 2011). The default current density maps were normalized through Standardized LOw Resolution brain Electromagnetic TomogrAphy approach (sLORETA, Pascual-Marqui, 2002). Welch's method was used to calculate the power spectrum at the level of the reconstructed sources. Window-overlap was 50%. Due to the small number of EEG channels, we grouped cortical vertices into major regions (ROIs), aggregated accordingly to Desikan-Killiany atlas by following a similar approach (Cesnaite et al., 2023).

2.3.3. *Periodic and aperiodic components of the power spectral density*

The specparam algorithm (version 1.0.0 (Donoghue et al., 2020) was used to parametrise power spectra of ROIs. In specparam algorithm, the power spectrum PSD is modeled as a combination of aperiodic component AP and a sum of N oscillatory peak modeled with a Gaussian:

$$P = AP + \sum_{n=0}^{N} G_n.$$

The component $AP(f)$ for frequency f is expressed by the formula:

$$AP(f) = b - log(k + f^{\chi}),$$

where b is the broadband offset, χ is the exponent and k is the knee parameter, controlling the "bend", When $k = 0$ this the component AP will be a line fitted in the log-log space (this is later referred as a fixed mode). in this case, the slope of the line a in log-log space is directly related to exponent χ, $\chi =$

$-a$ (Donoghue et al., 2020). The output of algorithm for estimated peaks are the mean of the Gaussian G_n for the center frequency of the peak, aperiodic-adjusted power (the distance between the peak of the Gaussian and the aperiodic fit at this frequency) and bandwidth as 2 *SD* of the fitted Gaussian.

In the current analysis, power spectra were parameterized across the frequency range 3 to 48 Hz (the maximal frequency range to avoid the line noise frequency) using the "fixed" mode (Figure 1). Additional algorithm settings were set as: peak width limits: [2.5 8]; max number of peaks: 6; minimum peak height: 0.5; peak threshold: 2. All the parameters describing identified peaks, offset, exponent and the parameters describing how well the model was fit were extracted.

Figure 1. Example plot with fitted parameters (associated with participant 'sub-032302' right cuneus). The x-axis shows the frequency band in Hz units. The y-axis shows power in arbitrary units (a.u.) associated with resting-state activity in the eyes-closed condition at the right cuneus ROI (referred to the Desikan-Killiany atlas) for this participant.

The parameters were extracted for every PSDs in the Eyes Closed condition, then ROIs that belong to the occipital regions were selected (that is where dominant activity in alpha is expected to exhibit age-related patterns; (Grandy et al., 2013; Knyazeva et al., 2018; Mizukami & Katada, 2018); thus the parameters from all ROIs belonging to this region were averaged. The choice of parameters gave a median goodness-of-fit measure of $r^2 = 0.981$, IQR= [0.971,0.978] across all regions within the occipital lobe. Model fits were not statistically different between the two groups: YA median r2 = 0.983, IQR= [0.974,0.988]; OA median r2 = 0.976, IQR= [0.962, 0.984]. Thus, although other processing parameters could have been chosen, we achieved suitable spectral parameterization across participants and regions. Compared to previous studies using the specparam algorithm, where frequency range varies, many used 40 Hz as the upper frequency range (Cesnaite et al., 2023; Merkin et al., 2021; Tröndle et al., 2023). For sake of clarity, using 48 Hz as upper frequency is also a widely adopted option (Iemi et al., 2019; van Nifterick et al., 2023), and stays within the recommendations of the algorithm's authors.. All participants showed a discernible alpha peak in the PSD (see an example in Figure 1). Individual alpha peak frequency values per subject were created from periodic components fitted by algorithm in the alpha range (8-12 Hz), but instead of averaging several of them per subject per ROI only the one that had the highest value of PW was taken per ROI and then averaged across ROIs within the occipital region. Figure 2 offers a qualitative overview of the differences in EEG power spectra among our groups. The figure visually confirms differences in IAPF (Individual Alpha Peak Frequency) values between the young and old populations.

Figure 2. Age- and education-related differences in resting EEG power spectra in the occipital region, in the eyes closed condition. The plot shows a median for each group (Young – High Edu = younger adults with high education, Old – High Edu = older adults with high education; Old – Low Edu = older adults with low education) and a 50% percentile interval, ranging from the 25 to 75 percentiles.

2.4. Statistical Analyses

Analyses were performed with the R software (R Core Team, 2022). Standardized scores on cognitive tests (z-scaled) were first correlated with gray matter and EEG parameters: IAPF, exponent, and offset, Figure S2 in Supplementary Material. Shapiro-Wilk tests on each dependent variable, inspection of residuals, and Kolmogorov-Smirnov analysis were performed to build up the regression models. Results indicated General Linear regression Models as suitable; they included processing speed and accuracy scores on cognitive tests as dependent variables, the variable group as a factor: older adults

with high education vs. older adults with lower education vs. young adults (all high education). Other continuous predictors of interest were the periodic and the aperiodic EEG components: IAPF, exponent, and offset values. Sex and normalized gray matter volume were accounted for in all regression models. A simplified syntax of the R linear models is reported below:

$$test\ score \sim EEG\ parameter\ x\ group + sex + Gray\ Matter\ volume$$

Sex and Gray matter volume were added as covariates as they were two relevant variables that could also be associated with cognitive performance.

3. RESULTS

3.1. Descriptive Analyses

Young adults performed significantly better on all cognitive tasks than older adults (Table S1 and Figure S1). IAPF, exponent, and offset values were shown in young adults compared to older adults (Figure 3 and S1). Older adults with different education did not differ on most cognitive tasks except for the working memory one, where older adults with high education performed better than older adults with lower education and more similarly to the younger adults (Table 1).

	Young Adults		Older Adults with high education		Older Adults with low education		Group comparison Young – High Edu vs Old – Low Edu		Group comparison Young – High Edu vs Old – Low Edu		Group comparison Old – Low Edu vs Old – Low Edu	
	Range	M(SD)	Range	M (SD)	Range	M (SD)	χ^2	p	χ^2	p	χ^2	p
PS_Allertness	(-1.47 - 1.50)	-0.43 (0.56)	(-0.99 - 3.35)	0.69 (1.07)	(-0.66 - 5.29)	0.77 (1.07)	23.15	<0.001	30.44	<0.001	<0.01	0.92

	Young Adults		Older Adults with high education		Older Adults with low education		Group comparison Young – High Edu vs Old – Low Edu		Group comparison Young – High Edu vs Old – Low Edu		Group comparison Old – Low Edu vs Old – Low Edu	
PS_Visual Attention	(-1.47 - 1.50)	-0.43 (0.56)	(-0.71 - 3.62)	0.94 (1.07)	(-0.84 - 3.59)	0.97 (1.07)	42.84	<0.001	42.11	<0.001	0.08	0.77
MEM_WM	(-2.83 - 0.80)	0.24 (0.71)	(-3.27 - 0.80)	-0.04 (1.28)	(-3.72 - 0.80)	-0.92 (1.28)	0.64	0.42	26.30	<0.001	8.13	<0.01
MEM_Del	(-2.46 - 1.27)	0.30 (0.84)	(-3.51 - 0.79)	-0.85 (0.97)	(-2.46 - 1.08)	-0.54 (0.97)	24.16	<0.001	23.20	<0.001	0.98	0.32
GM_Volume	(0.42 - 0.50)	0.46 (0.01)	(0.36 - 0.44)	0.41 (0.01)	(0.38 - 0.45)	0.42 (0.01)	56.63	<0.001	65.30	<0.001	2.07	0.14
IAPF	(8.95 - 11.98)	10.38 (0.68)	(8.81 - 11.85)	10.04 (0.67)	(8.69 - 11.44)	10.00 (0.67)	3.96	0.04	5.46	0.01	0.17	0.67
Exponent	(0.59 - 2.71)	2.00 (0.36)	(1.11 - 2.07)	1.67 (0.38)	(0.61 - 2.48)	1.61 (0.38)	22.19	<0.001	20.77	<0.001	0.21	0.64
Offset	(-17.21 - 14.55)	-15.72 (0.54)	(-16.89 - 15.28)	-16.23 (0.51)	(-17.52 - 15.22)	-16.24 (0.51)	18.33	<0.001	18.61	<0.001	0.02	0.88

Table 1. Descriptive information about the sample and group comparisons based on participants' age and education. Table depicts: Processing Speed alertness (PS_Allertness), Processing Speed Visual Attention (PS_Attention), Working Memory accuracy (MEM_WM) and Delayed Memory accuracy (MEM_Del); gray matter volume normalized (GM_Volume), Individual Alpha Peak Frequency (IAPF), Exponent, and Offset.

Figure 3. Age- and Education-related differences of the EEG components. The Individual Alpha Peak Frequency, the exponent, and the offset of each group (Young – High Edu = younger adults with high education, Old – High Edu = older adults with high education; Old – Low Edu = older adults with low education) are shown on the y-axis.

3.2. EEG spectral parameters and cognitive performance

In the whole sample (179) a higher exponent and offset predicted significantly a better performance on the visual attention task (exp: B = -0.44, p < 0.01; offset: B = -0.32, p < 0.01). The exponent and the offset values predicted working memory capacity (exp: B = 0.37, p = 0.04; offset: B = 0.31, p = 0.01); see Section C of Supplementary Materials. Significant results emerged when considering the three groups for both the exponent and the offset, in the visual attention and working memory tasks. More specifically, compared to the group of young adults, where exponent and offset did not predict any variation in cognitive performance, older adults showed significant associations, different across educational levels. In the visual attention task, low educated older adults had a better (faster) performance at the higher aperiodic values (exp: B = -

0.67, p = 0.04; offset: B = -0.56, p=0.03) while highly educated ones had a worse performance (slower) at the higher values of the exponent (exp: B = 1.41, p=0.02), a result confirmed post-hoc through slope comparisons, which confirmed a general age-related effect and also a significant difference between older adults with different education, at the third quartile of the exponent values (t=2.45; p = 0.03), see Figure 4.

Figure 4. Relationship between exponent and response times in the visual attention task (processing speed). On the x-axis, the exponent value parameterized at the occipital level and in the Alpha band (8-12 Hz) is reported. On the y-axis, the z-scores associated with the Processing Speed response times: visual attention (i.e., "Trail Making Test-B") is reported. The higher the exponent, the slower older adults with higher education, the faster the older adults with lower education. A non-significant relationship was shown between exponent and response times in the group of young adults.

Results show that, also in the working memory task, older adults with high education had worse performance with increasing exponent values, as compared with those with low education (B = -1.71, p=0.03). Post-hoc slope comparison showed no main education-related difference among older adults. Young adults and highly educated older adults did not differ between each other at

different quartiles of exponent values (25%: t=1.16, p = 0.47; 50%: t=1.94, p = 0.12; 75%: t=2.12, p = 0.08); Figure S6 in Supplementary Materials and Tables S2-5 for more details.

4. DISCUSSION

The present study aimed to investigate the relationship between aperiodic activity and cognitive performance by accounting for the level of education in older individuals, and comparing them with a control group of healthy younger adults highly educated (with an expected high neural and cognitive efficiency carried out by young age and high education).

Older adults were cognitively less efficient than young adults across all cognitive tasks, which aligns with the well-established literature about cognitive decline in healthy aging (Harada et al., 2013). Upon stratifying older adults based on their education, the results indicated that those with higher education exhibited comparable performance to older adults with lower education, except for working memory performance. Indeed, older adults with higher education showed better working memory performance that made this group more similar to the young group, suggesting a potential role of education in mitigating age-related cognitive decline.

Consistent with previous evidence (Dustman et al., 1993; Hill et al., 2022; Voytek et al., 2015), in this study the periodic and aperiodic components of EEG could differentiate between young and older adults: older adults indeed exhibited lower values across components, compared to younger adults. More specifically, in relation to the periodic component of the EEG signal, IAPF showed the pattern of an age-related slowing, as confirmed in recent studies (Knyazeva et al., 2018; Mizukami & Katada, 2018). Several interpretations have been advocated to link periodic EEG components with aging. In addition to slowing with age, structural alterations in the brain have also been associated with the decline in power and peak frequency of alpha oscillations, particularly in older individuals (Babiloni et al., 2006; Kumral et al., 2022). Additionally, the stability of IAPF over the life course reflects the preserved functionality of the central nervous

system (Grandy et al., 2013). Regarding cognition, IAPF has previously demonstrated a positive relationship with interference resolution in working memory performance, primarily observed in the temporal lobes (Cesnaite et al., 2023). This may indicate that IAPF plays a functional role in the ability to disregard and suppress interfering information. However, our study did not confirm the relationship between IAPF and working memory, suggesting that other factors, such as the specific nature of the task may mediate it.

In relation to the aperiodic EEG components, we found that both exponent and offset values significantly decreased with age. This result corroborates previous evidence suggesting aperiodic EEG component can serve as neurophysiological marker of aging. Furthermore, recent studies have revealed that aperiodic activity is influenced by various factors, including drugs (Colombo et al., 2019; Waschke et al., 2021) and level of arousal (Lendner et al., 2020). However, the potential mediation of education, specifically its influence on the relationship between aperiodic components and cognitive performance, has remained unexplored in the literature.

In our study, education might help in interpreting the relationship between aperiodic components and performance on some tasks, that is visual attention (processing speed) tasks and working memory tasks but not on a delayed memory task. Such a relationship varied depending on education level, with a reversed pattern between exponent and cognitive performance in older adults across higher vs lower education. Older adults with lower education displayed a positive relationship between exponent and cognitive performance, while those with higher education (counterintuitively) exhibited the opposite trend. In this sense, the neurobiological bases of aperiodic activity are still uncertain. However, evidence suggests that a flatter exponent (when the exponent approximates zero) may reflect an increase of asynchronous background neuronal firing, commonly called neural noise (Hill et al., 2022).

Moreover, in non-linear systems like the brain, the notion of stochastic resonance proposes that information at the threshold level can be better processed within an optimal noise range than

without noise (Miniussi et al., 2013). If different exponent values represent varying levels of neural noise, it is possible that noise also has different effects on performance according to the system. In older adults with lower education levels, higher exponents - corresponding to lower noise values - may contribute to better performance. On the other hand, older adults with higher education would exhibit the opposite pattern. In this latter group, higher exponents (lower noise values) would reduce performance efficiency. These two scenarios may depend on the fact that, according to the stochastic resonance framework, there is no ideal level of noise and its effect on performance may not follow a linear pattern and can vary based on the specific system.

Although this result may seem counterintuitive, a similar reversed pattern has also been observed in a previous study that examined the relationship between mathematical achievement and glutamate concentrations. Glutamate has the effect of flattening the power spectrum, leading to exponent values closer to zero (Colombo et al., 2019). A study by Zacharopoulos and colleagues (2021) has demonstrated that the concentration of glutamate and the exponent levels could result in reversed cognitive performance outcomes depending on the participants' age. Specifically, in their study, the concentration of glutamate (in the intraparietal sulcus) was negatively associated with mathematical achievement in younger participants, but it was positively associated with mathematical achievement in older participants. In our study, where the observed effects on cognitive performance cannot be influenced by age within the group of older adults, they could be associated with educational levels, which is a distinguishing factor between the two aging groups.

Although tantalizing, this proposed relationship between exponent values, noise levels, and cognitive performance in older adults with different education levels should be interpreted cautiously. Indeed, education may mediate the relationship between the aperiodic component and cognitive performance differently. For example, it may impact cognitive strategies, task engagement, and compensatory mechanisms, leading individuals with higher education to have

a high cognitive performance independently of the specific neural mechanisms reflected by the aperiodic EEG components. We did observe a relationship between the exponent and processing speed, in line with Ouyang and colleagues (2020). This study also partially replicates what it was found in some previous studies where a relationship between exponent and working memory performance was identified (Donoghue et al., 2020; Voytek et al., 2015).

Our results cannot be interpreted as exhaustive, rather they should emphasize the importance of considering the aperiodic component of EEG signal as a marker of neurophysiological mechanisms related to cognitive performance, which could be mediated by various aspects. Our study, in particular, focused on education as one of these aspects. Future longitudinal studies may explore the ontogenetic trajectory of the exponent further to investigate its role in cognitive performance during aging. In the present study, the availability of age and education variables in a categorical form might have limited the assessment of neurobehavioral relationships and the potential use of finer analysis modeling.

Future studies may explore the intricate connection between EEG parameters and cognition by encompassing a broader range of variables that could modulate such a relationship. Another relevant aspect is related to the correlational nature of this study: although it could provide some insights into relationships among the variables, it cannot establish causation, and results should be interpreted as potential incipit for further investigations about the modulatory mechanisms of cognitive reserve in the context of EEG parametrization.

This study may open to future research about the modulatory role of education and other cognitive reserve proxies in influencing the crucial relationship between aperiodic EEG and cognitive capacity in aging.

Bibliography

Babayan, A., Erbey, M., Kumral, D., Reinelt, J. D., Reiter, A. M. F., Röbbig, J., Schaare, H. L., Uhlig,

    M., Anwander, A., Bazin, P.-L., Horstmann, A., Lampe, L., Nikulin, V. V., Okon-Singer, H.,

    Preusser, S., Pampel, A., Rohr, C. S., Sacher, J., Thöne-Otto, A., … Villringer, A. (2019). A

    mind-brain-body dataset of MRI, EEG, cognition, emotion, and peripheral physiology in

    young and old adults. *Scientific Data*, *6*, 180308. https://doi.org/10/gfvkp5

Babiloni, C., Binetti, G., Cassarino, A., Dal Forno, G., Del Percio, C., Ferreri, F., Ferri, R., Frisoni,

    G., Galderisi, S., Hirata, K., Lanuzza, B., Miniussi, C., Mucci, A., Nobili, F., Rodriguez, G.,

    Luca Romani, G., & Rossini, P. M. (2006). Sources of cortical rhythms in adults during

    physiological aging: A multicentric EEG study. *Human Brain Mapping*, *27*(2), 162–172.

    https://doi.org/10.1002/hbm.20175

Bak, P., Tang, C., & Wiesenfeld, K. (1987). Self-organized criticality: An explanation of the 1/f noise.

    Physical Review Letters, *59*(4), 381–384. https://doi.org/10.1103/PhysRevLett.59.381

Bédard, C., & Destexhe, A. (2009). Macroscopic Models of Local Field Potentials and the Apparent

    1/f Noise in Brain Activity. *Biophysical Journal*, *96*(7), 2589–2603.

    https://doi.org/10.1016/j.bpj.2008.12.3951

Buckner, R. L. (2004). Memory and Executive Function in Aging and AD. *Neuron*, *44*(1), 195–208.

    https://doi.org/10.1016/j.neuron.2004.09.006

Bullmore, E., & Sporns, O. (2009). Complex brain networks: Graph theoretical analysis of structural

    and functional systems. *Nature Reviews Neuroscience*, *10*(3), Article 3.

    https://doi.org/10.1038/nrn2575

Cabeza, R., Albert, M., Belleville, S., Craik, F. I. M., Duarte, A., Grady, C. L., Lindenberger, U.,

    Nyberg, L., Park, D. C., Reuter-Lorenz, P. A., Rugg, M. D., Steffener, J., & Rajah, M. N.

    (2018). Maintenance, reserve and compensation: The cognitive neuroscience of healthy

    ageing. *Nature Reviews Neuroscience*, *19*(11), Article 11. https://doi.org/10.1038/s41583-

    018-0068-2

Camandola, S., & Mattson, M. P. (2017). Brain metabolism in health, aging, and neurodegeneration. *The EMBO Journal*, *36*(11), 1474–1492. https://doi.org/10.15252/embj.201695810

Cesnaite, E., Steinfath, P., Jamshidi Idaji, M., Stephani, T., Kumral, D., Haufe, S., Sander, C., Hensch, T., Hegerl, U., Riedel-Heller, S., Röhr, S., Schroeter, M. L., Witte, A., Villringer, A., & Nikulin, V. V. (2023). Alterations in rhythmic and non-rhythmic resting-state EEG activity and their link to cognition in older age. *NeuroImage*, *268*, 119810. https://doi.org/10.1016/j.neuroimage.2022.119810

Colombo, M. A., Napolitani, M., Boly, M., Gosseries, O., Casarotto, S., Rosanova, M., Brichant, J.-F., Boveroux, P., Rex, S., Laureys, S., Massimini, M., Chieregato, A., & Sarasso, S. (2019). The spectral exponent of the resting EEG indexes the presence of consciousness during unresponsiveness induced by propofol, xenon, and ketamine. *NeuroImage*, *189*, 631–644. https://doi.org/10.1016/j.neuroimage.2019.01.024

Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A. H., Wallis, J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing neural power spectra into periodic and aperiodic components. *Nature Neuroscience*, *23*(12), 1655–1665. https://doi.org/10.1038/s41593-020-00744-x

Dustman, R. E., Shearer, D. E., & Emmerson, R. Y. (1993). EEG and event-related potentials in normal aging. *Progress in Neurobiology*, *41*(3), 369–401. https://doi.org/10.1016/0301-0082(93)90005-d

Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance from field potentials. *NeuroImage*, *158*, 70–78. https://doi.org/10.1016/j.neuroimage.2017.06.078

Gazzaley, A., Cooney, J. W., Rissman, J., & D'Esposito, M. (2005). Top-down suppression deficit underlies working memory impairment in normal aging. *Nature Neuroscience*, *8*(10), 1298–1300. https://doi.org/10.1038/nn1543

Grandy, T. H., Werkle-Bergner, M., Chicherio, C., Schmiedek, F., Lövdén, M., & Lindenberger, U. (2013). Peak individual alpha frequency qualifies as a stable neurophysiological trait marker in healthy younger and older adults. *Psychophysiology*, *50*(6), 570–582. https://doi.org/10.1111/psyp.12043

Habes, M., Erus, G., Toledo, J. B., Zhang, T., Bryan, N., Launer, L. J., Rosseel, Y., Janowitz, D., Doshi, J., Van der Auwera, S., von Sarnowski, B., Hegenscheid, K., Hosten, N., Homuth, G., Völzke, H., Schminke, U., Hoffmann, W., Grabe, H. J., & Davatzikos, C. (2016). White matter hyperintensities and imaging patterns of brain ageing in the general population. *Brain: A Journal of Neurology*, *139*(Pt 4), 1164–1179. https://doi.org/10.1093/brain/aww008

Harada, C. N., Natelson Love, M. C., & Triebel, K. L. (2013). Normal cognitive aging. *Clinics in Geriatric Medicine*, *29*(4), 737–752. https://doi.org/10.1016/j.cger.2013.07.002

Hill, A. T., Clark, G. M., Bigelow, F. J., Lum, J. A. G., & Enticott, P. G. (2022). Periodic and aperiodic neural activity displays age-dependent changes across early-to-middle childhood. *Developmental Cognitive Neuroscience*, *54*, 101076. https://doi.org/10.1016/j.dcn.2022.101076

Iemi, L., Busch, N. A., Laudini, A., Haegens, S., Samaha, J., Villringer, A., & Nikulin, V. V. (2019). Multiple mechanisms link prestimulus neural oscillations to sensory responses. *ELife*, *8*, e43620. https://doi.org/10.7554/eLife.43620

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: A review and analysis. *Brain Research. Brain Research Reviews*, *29*(2–3), 169–195. https://doi.org/10.1016/s0165-0173(98)00056-3

Knyazeva, M. G., Barzegaran, E., Vildavski, V. Y., & Demonet, J.-F. (2018). Aging of human alpha rhythm. *Neurobiology of Aging*, *69*, 261–273. https://doi.org/10.1016/j.neurobiolaging.2018.05.018

Kumral, D., Cesnaite, E., Beyer, F., Hofmann, S. M., Hensch, T., Sander, C., Hegerl, U., Haufe, S., Villringer, A., Witte, A. V., & Nikulin, V. V. (2022). Relationship between regional white matter hyperintensities and alpha oscillations in older adults. *Neurobiology of Aging*, *112*, 1–11. https://doi.org/10.1016/j.neurobiolaging.2021.10.006

Lendner, J. D., Helfrich, R. F., Mander, B. A., Romundstad, L., Lin, J. J., Walker, M. P., Larsson, P. G., & Knight, R. T. (2020). An electrophysiological marker of arousal level in humans. *ELife*, *9*, e55092. https://doi.org/10.7554/eLife.55092

Lojo-Seoane, C., Facal, D., Guàrdia-Olmos, J., Pereiro, A. X., & Juncos-Rabadán, O. (2018). Effects of Cognitive Reserve on Cognitive Performance in a Follow-Up Study in Older Adults With Subjective Cognitive Complaints. The Role of Working Memory. *Frontiers in Aging Neuroscience*, *10*, 189. https://doi.org/10.3389/fnagi.2018.00189

Lövdén, M., Fratiglioni, L., Glymour, M. M., Lindenberger, U., & Tucker-Drob, E. M. (2020). Education and Cognitive Functioning Across the Life Span. *Psychological Science in the Public Interest: A Journal of the American Psychological Society*, *21*(1), 6–41. https://doi.org/10.1177/1529100620920576

Manning, J. R., Jacobs, J., Fried, I., & Kahana, M. J. (2009). Broadband shifts in local field potential power spectra are correlated with single-neuron spiking in humans. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *29*(43), 13613–13620. https://doi.org/10.1523/JNEUROSCI.2041-09.2009

Merkin, A., Sghirripa, S., Graetz, L., Smith, A. E., Hordacre, B., Harris, R., Pitcher, J., Semmler, J., Rogasch, N. C., & Goldsworthy, M. (2021). *Age differences in aperiodic neural activity measured with resting EEG* (p. 2021.08.31.458328). bioRxiv. https://doi.org/10.1101/2021.08.31.458328

Michels, L., Muthuraman, M., Lüchinger, R., Martin, E., Anwar, A. R., Raethjen, J., Brandeis, D., & Siniatchkin, M. (2013). Developmental changes of functional and directed resting-state connectivities associated with neuronal oscillations in EEG. *NeuroImage*, *81*, 231–242. https://doi.org/10.1016/j.neuroimage.2013.04.030

Miller, J. F., Neufang, M., Solway, A., Brandt, A., Trippel, M., Mader, I., Hefft, S., Merkow, M., Polyn, S. M., Jacobs, J., Kahana, M. J., & Schulze-Bonhage, A. (2013). Neural activity in human hippocampal formation reveals the spatial context of retrieved memories. *Science (New York, N.Y.)*, *342*(6162), 1111–1114. https://doi.org/10.1126/science.1244056

Miller, K. J., Sorensen, L. B., Ojemann, J. G., & den Nijs, M. (2009). Power-law scaling in the brain surface electric potential. *PLoS Computational Biology*, *5*(12), e1000609. https://doi.org/10.1371/journal.pcbi.1000609

Miniussi, C., Harris, J. A., & Ruzzoli, M. (2013). Modelling non-invasive brain stimulation in cognitive neuroscience. *Neuroscience and Biobehavioral Reviews*, *37*(8), 1702–1712. https://doi.org/10.1016/j.neubiorev.2013.06.014

Mizukami, K., & Katada, A. (2018). EEG frequency characteristics in healthy advanced elderly. *Journal of Psychophysiology*, *32*(3), 131–139. https://doi.org/10.1027/0269-8803/a000190

Mondini, S., Pucci, V., Montemurro, S., & Rumiati, R. I. (2022). Protective factors for subjective cognitive decline individuals: Trajectories and changes in a longitudinal study with Italian elderly. *European Journal of Neurology*, *29*(3), 691–697. https://doi.org/10.1111/ene.15183

Montemurro, S., Mondini, S., & Arcara, G. (2021). Heterogeneity of effects of cognitive reserve on performance in probable Alzheimer's disease and in subjective cognitive decline. *Neuropsychology*, *35*(8), 876–888. https://doi.org/10.1037/neu0000770

Niemann, H., Sturm, W., Thöne-Otto, A. I. T., & Willmes, K. (2008). *CVLT California Verbal Learning Test. German adaptation. Manual*. https://pure.mpg.de/pubman/faces/ViewItemOverviewPage.jsp?itemId=item_721063

Nyberg, L., Lövdén, M., Riklund, K., Lindenberger, U., & Bäckman, L. (2012). Memory aging and brain maintenance. *Trends in Cognitive Sciences*, *16*(5), 292–305. https://doi.org/10.1016/j.tics.2012.04.005

O'Byrne, J., & Jerbi, K. (2022). How critical is brain criticality? *Trends in Neurosciences*, *45*(11), 820–837. https://doi.org/10.1016/j.tins.2022.08.007

Ouyang, G., Hildebrandt, A., Schmitz, F., & Herrmann, C. S. (2020). Decomposing alpha and 1/f brain activities reveals their differential associations with cognitive processing speed. *NeuroImage*, *205*, 116304. https://doi.org/10.1016/j.neuroimage.2019.116304

Pascual-Marqui, R. D. (2002). Standardized low-resolution brain electromagnetic tomography (sLORETA): Technical details. *Methods and Findings in Experimental and Clinical Pharmacology*, *24 Suppl D*, 5–12.

Pathania, A., Schreiber, M., Miller, M. W., Euler, M. J., & Lohse, K. R. (2021). Exploring the reliability and sensitivity of the EEG power spectrum as a biomarker. *International Journal of Psychophysiology*, *160*, 18–27. https://doi.org/10.1016/j.ijpsycho.2020.12.002

Podell, J. E., Sambataro, F., Murty, V. P., Emery, M. R., Tong, Y., Das, S., Goldberg, T. E., Weinberger, D. R., & Mattay, V. S. (2012). Neurophysiological correlates of age-related changes in working memory updating. *NeuroImage*, *62*(3), 2151–2160. https://doi.org/10.1016/j.neuroimage.2012.05.066

R Core Team. (2022). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.

Reitan, R. M. (1992). *Trail Making Test: Manual for administration and scoring*.

Rossini, P. M., Rossi, S., Babiloni, C., & Polich, J. (2007a). Clinical neurophysiology of aging brain: From normal aging to neurodegeneration. *Progress in Neurobiology*, *83*(6), 375–400. https://doi.org/10.1016/j.pneurobio.2007.07.010

Rossini, P. M., Rossi, S., Babiloni, C., & Polich, J. (2007b). Clinical neurophysiology of aging brain: From normal aging to neurodegeneration. *Progress in Neurobiology*, *83*(6), 375–400. https://doi.org/10.1016/j.pneurobio.2007.07.010

Salthouse, T. A. (2010). Selective review of cognitive aging. *Journal of the International Neuropsychological Society: JINS*, *16*(5), 754–760. https://doi.org/10.1017/S1355617710000706

Scally, B., Burke, M. R., Bunce, D., & Delvenne, J.-F. (2018). Resting-state EEG power and connectivity are associated with alpha peak frequency slowing in healthy aging. *Neurobiology of Aging*, *71*, 149–155. https://doi.org/10.1016/j.neurobiolaging.2018.07.004

Schaworonkow, N., & Voytek, B. (2021). Longitudinal changes in aperiodic and periodic activity in electrophysiological recordings in the first seven months of life. *Developmental Cognitive Neuroscience*, *47*, 100895. https://doi.org/10.1016/j.dcn.2020.100895

Sghirripa, S., Graetz, L., Merkin, A., Rogasch, N. C., Ridding, M. C., Semmler, J. G., & Goldsworthy, M. R. (2020). *The role of alpha power in the suppression of anticipated distractors during verbal working memory* (p. 2020.07.16.207738). bioRxiv. https://doi.org/10.1101/2020.07.16.207738

Stern, Y. (2002). What is cognitive reserve? Theory and research application of the reserve concept. *Journal of the International Neuropsychological Society: JINS*, *8*(3), 448–460.

Stern, Y., Albert, M., Barnes, C. A., Cabeza, R., Pascual-Leone, A., & Rapp, P. R. (2023). A framework for concepts of reserve and resilience in aging. *Neurobiology of Aging*, *124*, 100–103. https://doi.org/10.1016/j.neurobiolaging.2022.10.015

Stern, Y., Gurland, B., Tatemichi, T. K., Tang, M. X., Wilder, D., & Mayeux, R. (1994). Influence of education and occupation on the incidence of Alzheimer's disease. *JAMA*, *271*(13), 1004–1010.

Tadel, F., Baillet, S., Mosher, J. C., Pantazis, D., & Leahy, R. M. (2011). Brainstorm: A user-friendly application for MEG/EEG analysis. *Computational Intelligence and Neuroscience*, *2011*, 879716. https://doi.org/10.1155/2011/879716

Taki, Y., Thyreau, B., Kinomura, S., Sato, K., Goto, R., Kawashima, R., & Fukuda, H. (2011). Correlations among brain gray matter volumes, age, gender, and hemisphere in healthy individuals. *PloS One*, *6*(7), e22734. https://doi.org/10.1371/journal.pone.0022734

Thuwal, K., Banerjee, A., & Roy, D. (2021). Aperiodic and Periodic Components of Ongoing Oscillatory Brain Dynamics Link Distinct Functional Aspects of Cognition across Adult Lifespan. *ENeuro*, *8*(5), ENEURO.0224-21.2021. https://doi.org/10.1523/ENEURO.0224-21.2021

Tran, T. T., Rolle, C. E., Gazzaley, A., & Voytek, B. (2020). Linked Sources of Neural Noise Contribute to Age-related Cognitive Decline. *Journal of Cognitive Neuroscience*, *32*(9), 1813–1822. https://doi.org/10.1162/jocn_a_01584

Tröndle, M., Popov, T., Pedroni, A., Pfeiffer, C., Barańczuk-Turska, Z., & Langer, N. (2023). Decomposing age effects in EEG alpha power. *Cortex*, *161*, 116–144. https://doi.org/10.1016/j.cortex.2023.02.002

van Nifterick, A. M., Mulder, D., Duineveld, D. J., Diachenko, M., Scheltens, P., Stam, C. J., van Kesteren, R. E., Linkenkaer-Hansen, K., Hillebrand, A., & Gouw, A. A. (2023). Resting-state oscillations reveal disturbed excitation–inhibition ratio in Alzheimer's disease patients. *Scientific Reports*, *13*(1), Article 1. https://doi.org/10.1038/s41598-023-33973-8

Voytek, B., Kramer, M. A., Case, J., Lepage, K. Q., Tempesta, Z. R., Knight, R. T., & Gazzaley, A. (2015). Age-Related Changes in 1/f Neural Electrophysiological Noise. *The Journal of*

Neuroscience: The Official Journal of the Society for Neuroscience, *35*(38), 13257–13265. https://doi.org/10.1523/JNEUROSCI.2332-14.2015

Waschke, L., Donoghue, T., Fiedler, L., Smith, S., Garrett, D. D., Voytek, B., & Obleser, J. (2021). Modality-specific tracking of attention and sensory statistics in the human electrophysiological spectral exponent. *ELife*, *10*, e70068. https://doi.org/10.7554/eLife.70068

Waschke, L., Wöstmann, M., & Obleser, J. (2017). States and traits of neural irregularity in the age-varying human brain. *Scientific Reports*, *7*(1), 17381. https://doi.org/10.1038/s41598-017-17766-4

Winawer, J., Kay, K. N., Foster, B. L., Rauschecker, A. M., Parvizi, J., & Wandell, B. A. (2013). Asynchronous broadband signals are the principal source of the BOLD response in human visual cortex. *Current Biology: CB*, *23*(13), 1145–1153. https://doi.org/10.1016/j.cub.2013.05.001

Zacharopoulos, G., Sella, F., Cohen Kadosh, K., Hartwright, C., Emir, U., & Cohen Kadosh, R. (2021). Predicting learning and achievement using GABA and glutamate concentrations in human development. *PLoS Biology*, *19*(7), e3001325. https://doi.org/10.1371/journal.pbio.3001325

Zimmermann, P., & Fimm, V. (2012). *Estbatterie zur Aufmerksamkeitsprüfung (TAP)* (Psytest). Psytest. https://www.psytest.net/index.php?page=TAP-2-2&hl=en_US