Multiple origins of dorsal ecdysial sutures in trilobites and their relatives

Kun-sheng Du [a,b,1], Jin Guo [b,c,1], Sarah R. Losso [d ✉], Stephen Pates [e, f,✉], Ming Li [g] and Ai-lin Chen [a,h,✉]

[a] Research Center of Paleobiology, Yuxi Normal University, Yuxi 653100, China.

[b] Key Laboratory for Palaeobiology and MEC International Joint Laboratory for Palaeoenvironment,

Institute of Palaeontology, Yunnan University, Kunming 650091, China.

[c] Management Committee of the Chengjiang Fossil Site World Heritage, Chengjiang, China.

[d] Museum of Comparative Zoology and Department of Organismic and Evolutionary Biology, Harvard

University, Cambridge, Massachusetts 02138, USA

[e] Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Penryn, Cornwall, TR10

9FE, UK

[f] Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, UK

[g] Institute of Geology, Chinese Academy of Geological Sciences, Beijing 100037, China.

[h] State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Paleontology,

CAS, Nanjing, 210008, China.

✉e-mail: ailinchen@yxnu.edu.cn (A.L.C.), sarahlosso@g.harvard.edu (S.R.L.) or s.pates@exeter.ac.uk

(S.P.).

[1] Joint first authors.

Keywords: Cambrian; arthropod; Chengjiang Biota; ecdysis; dorsal suture

Abstract

Euarthropods are an extremely diverse phylum in the modern, and have been since their origination in the

early Palaeozoic. They grow through moulting the exoskeleton (ecdysis) facilitated by breaking along lines

of weakness (sutures). Artiopodans, a group that includes trilobites and their non-biomineralizing relatives,

dominated arthropod diversity in benthic communities during the Palaeozoic. Most trilobites – a

hyperdiverse group of tens of thousands of species - moult by breaking the exoskeleton along cephalic

sutures, a strategy that has contributed to their high diversity during the Palaeozoic. However, the recent

description of similar sutures in early diverging non-trilobite artiopodans mean that it is unclear whether

these sutures evolved deep within Artiopoda, or convergently appeared multiple times within the group.

Here we describe new well-preserved material of *Acanthomeridion*, a putative early diverging artiopodan,

including hitherto unknown details of its ventral anatomy and appendages revealed through CT scanning,

highlighting additional possible homologous features between the ventral plates of this taxon and trilobite

free cheeks. We used three coding strategies treating ventral plates as homologous to trilobite free cheeks,

to trilobite cephalic doublure, or independently derived. If ventral plates are considered homologous to free

cheeks, *Acanthomeridion* is recovered sister to trilobites however dorsal ecdysial sutures are still recovered

at many places within Artiopoda. If ventral plates are considered homologous to doublure or non-

homologous, then *Acanthomeridion* is not recovered as sister to trilobites, and thus the ventral plates

represent a distinct feature to trilobite doublure/free cheeks.

1. Introduction

Euarthropods, more specifically members of the clade Artiopoda, dominated diversity in Palaeozoic animal communities (Daley et al., 2018). This group, which includes trilobites and their non-biomineralizing relatives, are united by a morphology consisting of a headshield covering a pair of uniramous antennae and at least three pairs of biramous limbs, followed by a dorsoventrally flattened body with a series of biramous appendages (Stein and Selden, 2012; Giribet and Edgecombe, 2019). The origins of key morphological features facilitating the diversity of trilobites can be found in their non-biomineralizing artiopodan relatives (e.g. Du et al., 2019; Schmidt et al., 2022). The cephalon, as an informative and functionally specialized body region, affords pivotal data for resolving the phylogenetic relationships and evolutionary history of arthropods, artiopodans, (Budd and Telford, 2009; Yang et al., 2013; Strausfeld et al., 2022) and trilobites in particular (e.g. Stubblefield 1936; Lieberman 2002; Paterson et al. 2019).

The presence, shape, and character of sutures in the trilobite cephalon have been central to classification schemes of the group for nearly 200 years (e.g. Emmrich 1839; Salter 1864; Beecher 1897a, b; Stubblefield 1936; Rasetti 1945; Whittington et al. 1997), while the similarity in the cranidial outlines of Cambrian suture-bearing groups supported a single origin or strong evolutionary constraint on its origins (Foote 1991; Hughes 2007). These sutures provided lines of weakness along which the cephalon split during moulting, creating an anterior ecdysial gape through which the animal exited the old exoskeleton (e.g. Henningsmoen 1975; Whittington 1990; Daley and Drage, 2016). Sutures may have served additional purposes (Stubblefield 1936) and were under additional functional demands, including feeding (Fortey & Owens 1999) and burrowing behaviours (Esteve et al. 2021). Different trilobite clades display distinct suture patterns, with two sets, the circumocular and marginal sutures, relevant for the creation of this ecdysial gape (Henningsmoen 1975). For the earliest diverging trilobites, olenellines (Palmer & Repina 1993; Lieberman 2002) which a have been recovered as a paraphyletic grade rather than a monophyletic group (e.g. Paterson et al. 2019) circumocular sutures facilitated shedding of the cornea during ecdysis (e.g. Ramskold & Edgecombe 1991) while a marginal suture (**Fig. 1e**) facilitated anterior egression through separation of the lateral cephalic doublure from the dorsal cephalon (e.g. Stubblefield 1936; Henningsmoen 1975). In later diverging trilobites the circumocular and marginal sutures combined into a

single system (**Fig. 1d**). Here the cornea is fused to the free cheek (e.g. Stubblefield 1936; Ramsköld &

Edgecombe 1991) and these facial sutures – the fused circumocular and marginal sutures - facilitated both

ecdysis of the visual surface and anterior egression of the exoskeleton following withdrawal from the free

cheeks (e.g. Henningsmoen 1975).

Fig. 1 Hypotheses of homology in sutures between *Acanthomeridion* and trilobites. (a) Cephalic

morphology of *Acanthomeridion serratum*. (b) Cephalic morphology of a proetid trilobite (modified from

Daley and Drage, 2016). (c) Cephalic morphology of a redlichid trilobite (modified from Whittington

1989). (d) Ventral plates and librigenae as homologous with dorsal suture. (e) Ventral plates and doublure

as homologous with marginal suture. (f) Non-homology between ventral plates of *Acanthomeridion*,

librigenae and doublure.

The origin of these fused dorsal ecdysial sutures – facial sutures – has traditionally been considered to fall

within Trilobita (e.g. Fortey and Whittington, 1989; Edgecombe and Ramsköld, 1999; Lieberman, 2002).

However, the presence of dorsal ecdysial sutures in earlier diverging artiopodans – the Protosutura (Du et

al. 2019) and eye slits in Petalopleura (Chen et al., 2019) raised questions of the homology and origins

within Artiopoda (e.g. Hou et al. 2017; Du et al. 2019). An alternative hypothesis for the origins of the

dorsal cephalic sutures emerged: that these had a deep root within Artiopoda and were subsequently lost in

some groups including multiple times within Trilobita (Hou et al., 2017a; Du et al., 2019), rather than the

traditional view that dorsal cephalic sutures in trilobites were derived within the clade, and thus these eye

slits and facial sutures were acquired independently. Support for the deep root hypothesis comes from the

variability of facial sutures in trilobites, and the recognition that convergence and loss of features is

common when they have a function or allow adapation to a particular niche (e.g. Moore & Wilmer 1997).

Further evidence for the variability of suture morphology comes from ontogenetic studies, such as the

fusion of dalmanitinid facial sutures during ontogeny (Drage et al., 2018), and the loss of this feature in

other trilobite such as 'Cedaria' woosteri (Hughes et al. 1997). This indicates that there is scope for a

facial suture to have been lost repeatedly in artiopodan groups earlier diverging than Trilobita, should this

feature prove to have a deeper origin within Artiopoda (Hou et al. 2017; Du et al. 2019). A deep root for

facial sutures within Artiopoda would have repercussions for the importance of the facial sutures in

determining trilobite relationships (e.g. Jell 2003) and the position of a grade of olenelline trilobites as the

earliest diverging members of the group (e.g. Paterson et al. 2019). To date, a third possibility – that the

ventral plates of *Acanthomeridion* are homologous to the doublure of olenellids, and thus *Acanthomeridion*

and olenelline trilobites share presence of an unfused marginal suture – has not received broad attention in

the literature, but should also be considered in light of the similarities in position of the ventral plates and

olenelline doublure, and the marginal rather than dorsal position of the suture in *Acanthomeridion.*.

Resolving the complex history of dorsal ecdysial sutures evolution within Artiopoda, requires the nature of

cephalic sutures in artiopodans such as *Acanthomeridion,* protosuturans, and petalopleurans to be resolved,

and their phylogenetic position confidently determined (Hou et al., 2017a; Du et al., 2019). However, the

morphology of many earlier diverging artiopodans is not completely understood (Ortega Hernández et al.,

2013; Lerosey-Aubril et al., 2017), contributing a significant amounts of missing data to character matrices

used in phylogenetic analyses. Up to now, *Acanthomeridion* was arguably the least well known of these

early diverging artiopodans, with its ventral anatomy and non-biomineralized structures very poorly known

(Hou and Bergström, 1997; Hou et al., 2017a).

Here we describe new specimens of *Acanthomeridion serratum*, using micro-CT to reveal details of the

ventral anatomy, eyes, and appendages for the first time, and reconstruct the changing shape of the body

during ontogeny. We use these new data to show that *A. anacanthus* is a junior synonym of *A. serratum.*

We then assess the phylogenetic position of the species considering two hypotheses relating to the dorsal

ecdysial sutures – as homologous to the dorsal ecdysial sutures (facial sutures combined of a fused

marginal and circumocular sutures) of trilobites (**Fig. 1d**), that they are homologous to marginal sutures of

olenellines (**Fig. 1e**), and that these structures are unique to *Acanthomeridion* (**Fig. 1f**), to evaluate the

different hypothesis of origination within Artiopoda.

2. Materials and methods

New specimens of *Acanthomeridion* were collected from Jiucun town, Chengjiang (**Fig. 2**) and

accessioned at the Management Committee of the Chengjiang Fossil Site World Heritage (CJHMD 00052–

00062), and Research Center of Paleobiology, Yuxi Normal University (YRCP 0016–0020). New

specimens of *Wutingaspis tingi* (YRCP 0021) and *Xandarella spectaculum* (YRCP 0022) were collected

from Haikou town, Kunming (**Fig. 2**) and housed at Research Center of Paleobiology, Yuxi Normal

University. The holotype of *Zhiwenia coronata* (YKLP 12370) is deposited at the Key Laboratory for

Palaeobiology, Yunnan University which was excavated from Xiaoshiba area, Kunming (**Fig. 2**). The

specimens were photographed by a LEICA DFC 550 digital camera mounted to a Stereoscope LEICA

M205 C, and scanned with a Zeiss Xradia 520 Versa X-ray Microscope and Nikon XTH 225 ST micro-

focus X-ray tomography machine. All micro-CT images were processed with the software Drishti (Zhai et

al., 2019). Figures were processed with CorelDRAW X8, Inkscape 1.0 and Illustrator.

Given that specimens with thorax longitudinally curved and specimens with flattened thorax are

considered as two species of *Acanthomeridion*, despite having the same exoskeletal morphology, we

measured their axial length and attempted to determined their true maximum width in all specimens,

including those described in this study and previously documented measurable specimens. The data was

137 initially analyzed using a bivariate regression fitting function in Microsoft Office Excel. Subsequently, the

138 software Past (Hammer et al. 2001) was employed for testing and validation, obtaining the same function

139 and conducting a rational analysis of the function.

140 The character matrix used for the phylogenetic analyses was adapted from Schmidt et al., (2022), itself

141 adapted from Du et al. (2019) and Chen et al. (2020). *Olenellus getzi* was added, to facilitate comparison

142 with olenelline trilobites, while *Anacheirurus adserai*, *Cryptolithus tesselatus* and *Olenoides serratus* were

143 added to provide more trilobite terminals. *Olenellus getzi* was chosen as it represents the olenelline with

144 best known soft tissues (antennae only; Dunbar 1925). Trunk appendages are known in *A. adserai, C.*

145 *tesselatus* and *O. serratus* (Walcott, 1911; Whittington, 1975) , with descriptions by Bicknell et al.,

146 2021; Losso and Ortega-Hernández 2022 for coding the matrix. Our updated dataset includes 70 taxa

147 (*Acanthomeridion anacanthus* was removed, four trilobites were added). Three matrices were used to test

148 the alternative hypotheses of sutures, one with 100 characters considering the ventral plates unique, and

149 not homologous to trilobite librigenae ('ventral plates under head' as a new character in the dataset) and

150 two with 99 characters. The first of these considered the ventral plates homologous to trilobite librigenae,

151 and the second to the doublure of olenellines (see Morphobank project). We added the character state

152 'ringed distribution lamellae' to Ch. 15 in all matrices (Chen et al., 2019; Du et al., 2019), and altered the

153 characters relating to the suture pattern (Ch. 31, 32) to include character states suitable for *Olenellus getzi*.

154 Given the difficulties of identifying circumocular sutures in non-biomineralizing artiopodans, Ch. 31

155 indicated the presence or absence of ecdysial sutures in the cephalon, while Ch. 32 indicated the type of

156 suture. This could either be marginal and/or circumocular, but unfused (state 0) or marginal and

157 circumocular fused into a facial suture (state 1). This facilitated differentiation between the state in

158 *Olenellus getzi* and the other trilobites in the dataset, and thus allowed the two hypotheses relating to

159 homology between the ventral plates and the doublure or free cheeks of trilobites to be tested. Eight

160 additional characters relating to the trilobite glabellar region were added, taken from Paterson et al. (2019),

161 in order to provide data to distinguish internal trilobite relationships. These were Ch. 5, 6, 7, 10, 11, 12, 14

162 and 15 in Paterson et al. (2019), which are Ch. 37 – 44 in all three matrices used for this study. A further

163 character (calcified thorax separated into prothorax and opisthothorax) was also added (Ch. 99 or 100 in

164 matrices used for this study). Matrices used in phylogenetic analyses are provided through morphobank

Phylogenetic analyses were performed with MrBayes (Bayesian Inference) (Ronquist et al., 2012). Both

the 'maximum information' and 'minimum assumptions' strategies of ref (Bapst et al., 2018) were utilised

on an earlier version of the matrix, in order to confirm that model choice was not a major driver in

differences for the results (Du et al. 2023). For the final version, the 'maximum information' strategy was

used, as model choice did not play a large influence on the topologies recovered. Bayesian analyses were

run for 20 million generations using four chains, every 1000^{th} sample stored and 25% burn-in (Lewis,

2001; Ronquist et al., 2012). Convergence was diagnosed using Tracer (Rambaut et al., 2018).

For each coding strategy, two sets of Bayesian analyses were run. The first was unconstrained, whereas for

the second all trilobites except for *Olenellus getzi* were constrained as a monophyletic group to recover the

relationships of trilobites (*Olenellus* as the earliest diverging member) comparable to the comprehensive

trilobite phylogenetic analysis of Paterson et al. (2019). Resulting posterior tree samples were compared by

comparisons in the consensus trees, and in multidimensional treespace (using R code and method from

Pates et al. 2022). The number of trees where *Acanthomeridion serratum* was sister to trilobites was

quantified for each coding strategy and for constrained and unconstrained analyses, and the area of

treespace occupied by the posterior samples – and trees where *A. serratum* was sister to triobites within

those samples – were visualized.

Legend
Road — Railway — Lake — River — Cities — Vallige — Fossil site — Cambrian Terreneuvian — Cambrian Series 2

Fig. 2 Sites yielding *Acanthomeridion* from the Cambrian Stage 3 Chengjiang Biota (red circles). All

the specimens of *A. serratum* used here are collected from Jiucun town, Chengjiang County.

3. Results

3.1 Systematic palaeontology

ARTIOPODA Hou and Bergström, 1997

Acanthomeridion Hou et al., 1989

Type and only species. *Acanthomeridion serratum* Hou et al., 1989

Diagnosis. Non-biomineralized artiopodan with elongate dorsal exoskeleton that gives body an

elliptical outline. Subtriangular head shield with deep lateral notches that accommodate stalked elliptical

eyes, rounded genal angles of the head shield, an axe-like hypostome, and paired teardrop-shaped plates on

either side of hypostome. Head bears large eyes and long multi-segmented antennae consisting of over 40

podomeres anterior to three pairs of small cephalic limbs. Trunk composed of 11 tergites bearing expanded

tergopleurae with well-developed distal spines, and a terminal spine. The ninth tergite bears a pleural spine

more elongate than others, eleventh tergite reduced and expanded into leaf-like outline. Each tergite bears a

pair of biramous limbs with long and dense spines on the endopods, and slender stick-like exopods with

long and dense bristles. Modified from Hou et al., 1989, 2017a.

Acanthomeridion serratum Hou et al., 1989 **Syn. nov.**

Figs. 3–5; 7g-i; S1–6

1989 *Acanthomeridion serratum* Hou et al., pl. Ⅲ, figs. 1-5; pl. Ⅳ, figs. 1-5; p. 46, text-figs. 3, 4.

1996 *Acanthomeridion serratum* Chen et al., p. 157, figs 199-203.

1997 *Acanthomeridion serratum* Hou and Bergström, p. 38.

1999 *Acanthomeridion serratum* Luo et al., p. 51, pl. 6 fig. 5.

1999 *Acanthomeridion serratum* Hou et al., p. 130, fig. 188.

2004 *Acanthomeridion serratum* Chen, p. 280, fig. 442.

2004 *Acanthomeridion serratum* Hou et al., p. 176, fig. 16.63; p. 177, fig. 16.64.

2017a *Acanthomeridion serratum*, *A*. *anacanthus*, and *Acanthomeridion* sp in Hou et al., p. 734, figs.

1-4.

2017b *Acanthomeridion serratum* Hou et al., p. 204, fig. 20.34; p. 205, fig. 20.35.

Type material. *Holotype*, CN 108305. *Paratype*, CN 108306–108310.

New material. Sixteen new specimens were collected from Jiucun **(Fig. 2)**, Chengjiang County and housed

in the Management Committee of the Chengjiang Fossil Site World Heritage (CJHMD 00052–00062), and

Research Center of Paleobiology, Yuxi Normal University (YRCP 0016–0020).

Occurrence. Yu'anshan Member, Qiongzhusi Formation, *Wutingaspis–Eoredlichia* biozone, Cambrian

Stage 3. Jiucun and Maotianshan, Chengjiang County, Yunnan, China; Mafang and Ercaicun, Haikou town,

Kunming, Yunnan, China **(Fig. 2)**.

Diagnosis. As for genus, by monotypy.

Description. The dorsal exoskeleton of *Acanthomeridion serratum* displays weak trilobation and elliptical

outline. Specimens measure from 20 to 75 mm along the sagittal axis (**Figs. 3a, c; 4a; 5; S1**–**6**). The head

shield is subtriangular in dorsal view, with rounded anterior margin and small spines on posterior margin

(**Figs. 3a, c; 4a; 5; S6a, c**). A pair of lateral posterior notches accommodate stalked elliptical eyes. The

lateral margin of the head shield is a suture that attaches paired ventral plates. Ventral plates have a

teardrop outline, occupy the entire length of the cephalon and terminate in a posterior spine (maximum

length 3.4 mm; **Fig. S2b**) that projects into the thorax as far as the anteriormost tergite (**Fig. S1a**). The

medial margin is curved towards the axially line and sits adjacent to a conterminant axe-shaped hypostome

(maximum dimensions 9.6 mm x 11.6 mm; **Fig. 3a, b**). Four pairs of appendages are present in the head

(**Figs. 3f**–**j; S5a**). A pair of antennae with at least 43 articles (**Figs. 4a, b; S2a, f; S4b**) sits anterior to three

pairs of small cephalic limbs (**Fig. 3f**–**j**). The cephalic appendages are poorly preserved and only the gross

morphology can be discerned as no individual podomeres are visible. The protopodite of the cephalic

limbs are subtriangular in outline. Endopodites and exopodites are not clearly preserved. Appendage three

may include the endopodite, but no podomeres are visible.

Fig. 3 *Acanthomeridion serratum* from the Cambrian Stage 3 Chengjiang Biota. (a, b) CJHMD

00052a/b respectively, individual with hypostome, ventral plates, and 11 tergites. (c, d) CJHMD 00053a,

showing ventral plate, dorsal sutures, and 11 tergites. (e–g) YRCP 0016a, showing the ventral plates, and

three post-antennal limbs. (h–k) Showing the post-antennal appendages under head, gnathobases of trunk

limbs, stick-like exopodites with bristles (black arrows), and endopodites with long spines (white arrows).

(f) Micro-CT image of YRCP 0016a; (h, i, k) Micro-CT images of CJHMD 00053a. Abbreviations: ant,

antenna; ca*n*, post-antennal appendage *n* beneath head; dbl, doublure; ds, dorsal suture; en, endopodites;

ex, exopodites; ey, eye; hs, head shield; hy, hypostome; L, left; lam, lamellae; no, notch; p*n*, podomere *n*;

R, right; T*n*, tergite *n*; ts, terminal spine; vp, ventral plate.

Fig. 4 *Acanthomeridion serratum* from the Cambrian Stage 3 Chengjiang Biota. (a–e) CJHMD 00055,

showing the antenna, ventral plate, endopodites with long spines (arrows in d), 11 tergites, and paddle-like

structure (red arrow). (a) Overview of whole specimen. (b) Detail of long left antenna. (c) Close-up of

ventral plate. (d) Details of long spines (arrows) of right endopodites. (e) Close-up of paddle-like structure

(red arrow). Abbreviations same as Figs. 2, 3.

The trunk is composed of 11 tergites which extend laterally into spinose tergopleurae. Tergites are all

subequal in length. Tergites 8-11 curve towards the posterior, and reduce in width progressively, so that

T11 is approximately 25% the width of T8 and 33% the width of T9 (**Figs. 3a, c; 4a; S1a, c, d**). Each

tergite has small spines on its lateral and posterior margins (**Figs. 6; S1a, c–f; S2i, k; S3a, j; S5b, e, f;**

S6a–f). The shape of tergites and the convexity of the trunk changes through ontogeny (**Fig. 5**). In the

smaller specimens, the trunk appears slender from a dorsal view as the body displays a high convexity

(**Fig. 5t**). Pleurae curve ventrally, and only slightly towards the posterior (**Fig. 5a–k**). In larger specimens,

the trunk is less convex than for the smaller specimens (**Fig. 5s**), ventral curvature of pleurae is decreased,

and posterior curvature increased (**Fig. 5o–r**). A linear relationship between trunk axial length and width is

obtained (**Fig. 6**). Where trunk appendages are preserved, each tergite is associated with a pair of biramous

trunk appendages (**Figs. 3a, c, h, k; 4; S1a, b**). The protopodite is sub triangular with a broad attachment

to the body wall and short robust gnathobasal spines along the medial margin (**Figs. 3h–j; S3a–f, i**). The

ventral margin of the protopodite has endites longer than the gnathobasal spins. The dorsal edge of the

protopodite is poorly preserved and does not clearly show the attachment of the exopodite. Exopodites are

slender, stick-like and long (8.3 mm, ca. 3x protopodite length), bearing two rows of lamellae (**Figs. 3h, k;**

S1a, b). Exopods are shorter than the width of corresponding tergites (**Figs. 3h, S1a, b**). The lamellae

appear short but may not be completely preserved. There is no evidence that the exopodite divided into

lobes or has articulations. Endopodites are composed of seven trapezoidal podomeres. Podomeres 1–6 bear

spiniferous endites arranged in rows, six on pd1, 2, four on pd3, 4, and three on pd5, 6. Podomere seven

displays a terminal claw of three elongate spines (**Figs. 3h, k; 4d**). From the preserved appendages, no

significant antero-posterior differentiation or dramatic size reduction can be discerned (**Figs. 3, S1**) One

specimen shows evidence for paired midgut diverticulae (**Fig. S4a, f**). The body terminates in a long,

slender spine, dorsal to a paired paddle-like structure (**Figs. 4a, e; S5b, g**).

Remarks. The eyes were previously identified as genal projections (see Fig. 1, Fig. 2A, and Fig. 3A in Hou

et al., 2017a) because of the limited available specimens and the posterior location of the eyes.

Acanthomeridion anacanthus Hou et al. 2017a is here proposed to be a junior subjective synonym of

Acanthomeridion serratum Hou et al.1989, an interpretation supported by:

1) The paddle-like structure under the terminal spine (**Figs. 4a, e; S5b, g**) was previously interpreted as the

twelfth tergite. The specimen YKLP 11115 (see Fig.4 in Hou et al., 2017a) which is a similar size to YKLP

11116 (holotype of *Acanthomeridion anacanthus*) and YKLP 11113 (see Fig.2A, C and Fig.3A, C in Hou

et al., 2017a) show 11 tergites.

2) Specimens CJHMD 00052-55 and YRCP 0016 show the evidently pleural spines of T8 and T9 **(Figs.**

3a, c; 4a; S1a, c, d; S2a, f; S4b, g).

3) The orientation of pleurae makes some specimens appear narrower than others. Specifically, those

where pleurae curve ventrally look more slender than those where pleurae are curved posteriorly.

Specimen YRCP 0017 preserves right ventrally curving pleura and left flat pleura, which lead to its right

pleura narrower than the left **(Figs. 5j, o; S2i)**. Moreover, most small specimens (1cm–3cm) are preserved

with ventrally curving pleurae **(Figs. 5a–e; S3a–c; S4a; S5a-c; S6g, i)** and these appear narrower than the

few small specimens with flat pleurae **(Figs. 5f; S4b**. See also Fig.2A–D, and Fig.4 in Hou et al., 2017a).

Larger specimens are typically preserved with flattened, posteriorly curving pleurae.

4) Twenty specimens show a good ontogenetic series of *Acanthomeridion* including some that have

previously have been assigned to '*Acanthomeridion anacanthus*' **(Figs. 5, 6)**. Linear measurements do not

support distinct species of different sizes, nor with different proportions of thorax axial length to width.

The length and width data fit well with a linear equation; most of the data fall within the 95% forecast

(**Fig. 6b**). Data fall close to the fitted line, as indicated by the R^2 value of 0.95. Lastly, specimens

previously assigned to '*A. anacanthus*' are recovered as smaller *A. serratum* specimens with similar

thoracic shape.

The new specimens and CT data presented herein allow the description of cephalic and trunk limbs, and a

conterminant hypostome adjacent to ventral plates in *Acanthomeridion* for the first time. The presence of a

terminal spine on these ventral plates is identified . The ventral plate is separated from the dorsal part of

the cephalon by a suture that runs close to the stalked eyes. The possible affinities of these ventral plates,

and the phylogenetic implications of these, are discussed below (Subsections 3.2, 3.3) The presence of four

appendages in the head (or three in the head and one underneath the articulation) is comparable to

Zhiwenia (Du et al., 2019), as well as other artiopodans including trilobites (Edgecombe and Ramsköld,

1999). However, the morphology of the three non-antenniform cephalic limbs is difficult to determine with

only the protopodite clearly visible. The lack of observed endopodites and exopodites on these three

appendages may be a preservational artefact. Possibly the absence of clear endopodites and exopodites

represents specialization of the cephalic appendages with the structures being significantly reduced. Other

Cambrian artiopodans have been shown to have appendage specialization (Losso and Ortega-Hernández,

2022) which is frequently found in the cephalon (Chen et al., 2019; Schmidt et al., 2022), but none show

reduction of both the exopodite or endopodite. In *Sinoburius lunaris*, the first two post-antennal

appendages have reduced endopodites with elongated exopodites (Chen et al., 2019), and in

Pygmaclypeatus daziensis the exopodite is significantly reduced in the cephalic appendages (Schmidt et

al., 2022).

Recently, the head of *Retifacies abnormalis* was interpreted to develop three pairs of uniramous post-

antennal appendages and following by one biramous appendage pair (Zhang et al., 2022). It would be

unclear the function of an appendage with both distal elements being significantly reduced until more clear

appendages are found.

The exopodites of *Acanthomeridion* also differ substantially from those known in other artiopodans.

Specifically, the stick-like nature of the exopodite, the rows of lamellae combined with the apparent lack

of segmentation, differ from the paddle shaped exopodites thought to characterise the remainder of the

group (e.g. Schmidt et al., 2022, and fig. 4 in Ortega Hernández et al., 2013).

Fig. 5 Ontogenetic series (a–r) of *Acanthomeridion* and their ventrally curling pleurae (s, t). (a–r)

Showing the individuals from smallest to largest with same scale bar. (s) Lateral view of (o), note the right

curling pleurae and left flat pleurae. (t) Lateral view of (f), showing the left curling pleurae.

Fig. 6 Scatterplot of axial length and width of thorax of *Acanthomeridion serratum.* (a) scatterplot of

raw data. (b) data with 95% forecast shown in pink. Specimens previously assigned to *A. anacanthus*

indicated by blue lettering.

3.2 Possible affinities of the ventral plates in *Acanthomeridion.*

3.2.1. ventral plates as homologues of trilobite free cheeks

The ventral plates of *Acanthomeridion* have previously been suggested to be homologues to the free

cheeks of trilobites, with the suture between these features and the remainder of the cephalon interpreted as

a facial suture **(Fig. 1d)** (e.g. Hou et al. 2017). Morphological similarities are apparent between the ventral

plates and free cheeks in many trilobites **(Fig. 7a, b)** (Whittington et al., 1997) as well as the path of suture

separating them from the rest of the cephalon. Our new data provide additional support for interpreting the

ventral plates as homologues of the librigenae. The ventral plates of *Acanthomeridion* are here shown to be

transversely broad, tapering posteriorly to a spine, in a position directly comparable to the genal spine of

many trilobites **(Figs. 1d; 7a, b)**. Furthermore, the ventral plate is separated from the rest of the cephalon

by a suture that passes near the compound eye. This path is similar to the facial suture (fused circumocular

and marginal sutures) of many trilobites especially those bearing opisthoparian facial sutures, which

separate the free cheeks from the cranidium passing through the visual surface **(Fig. 7a, b)** (Whittington et

al., 1997). The homology of eye notches (e.g., *Luohuilinella* and *Zhiwenia* **(Fig. 7e, f)**, and eye slits (e.g.

Phytophilaspis (Ivantsov, 1999) of petalopleurans, dorsal ecdysial sutures in trilobites and

Acanthomeridion has been suggested previously (Du et al. 2019). These eye slits join the dorsal eye with

the anterolateral margin of the head shield, possibly forming a single suture that facilitated release of the

visual surface and created an anterior gape for the animal to leave the old exoskeleton during ecdysis.

However, the evolution from eye notches to eye slits and finally to the loss of the eye slit can been seen,

calling into question the affinity between these structures and sutures (Chen et al. 2019). This hypothesis

was tested explicitly in the first matrix, where the ventral plates of *Acanthomeridion* were considered

homologous to the free cheeks of trilobites. In this analysis, the dorsal slits and sutures of other artiopodans

were also treated as homologous ecdysial sutures in the cephalon (Fig. 8a).

3.2.2. ventral plates as homologues to doublure in olenelloid trilobites

In the above scenario, the ventral plates of *Acanthomeridion* would be homologous to the free cheeks both

dorsally and ventrally in trilobites, including the doublure. However it is possible that the ventral plates are

only homologous to the doublure, and thus that the suture represents the marginal suture, rather than a

fused facial suture. The difficulty lies in the marginal position of the eyes, which could support the

interpretation that these are fused sutures, but might also represent a position that facilitated ecdysis

through a single, unfused, suture. Considered homologous only to the doublure, the suture in

Acanthomeridion separating the ventral plates from the cephalon would be homologous to the unfused,

marginal suture in olenelloids rather than the fused facial suture of most other trilobites **(Fig. 1e)**   (e.g.

Stubblefield 1936).

Few morphological features can be drawn to support this hypothesis beyond the ventral position of the

plates. In particular, the lateral position of the eyes in *Acanthomeridion*, and the path of the suture close to

these eyes, suggests that the suture functioned to release both the ventral plate and the eye. Although some

artiopodans are known with expansive doublure such as *Squamacula buckorum* (Paterson et al., 2012) this

structure usually follows the outline of the margin and lacks spines (Whittington et al., 1997).

This hypothesis was tested in the second matrix, where the ecdysial suture of *Acanthomeridion* was

considered unfused (**Fig. 8b**). In this scenario, the presence of eye slits and other dorsal sutures in other

artiopodans would not represent homologous structures, as the proposed evolution is from a ventral-lateral

structure to a dorsal one. Thus, the eye slits of the petalopleurans who bear them were not considered as

ecdysial sutures for this analysis.

3.2.3 ventral plates and eye slits of artiopodans as unique structures without homologues in trilobites

Eye slits of petalopleuran artiopodans as homologues to facial sutures of trilobites has drawn support from

comparisons such as *Loganopeltoides* (Rasetti 1948), where a suture charts a path from the dorsal eye to

the anterolateral margin of the head shield. However, in *Loganopeltoides* this suture represents the vestiges

of the free cheeks (Edgecombe & Ramksold 1999), a derived state within Trilobita. Given that the features

in support of homology between the ventral plates of *Acanthomeridion* and trilobite free cheeks, and

implicit within this a deep root of the facial suture within Artiopoda rather than Trilobita, may not

represent true homologues (e.g. spines are often evolved convergently, and the dorsal eye slits of

artiopodans have previously been considered unique characters – see e.g. Edgecombe & Ramskold 1999;

Chen et al., 2019) it is important to consider the hypothesis that these features are unique to

Acanthomeridion. Thus for this third analysis, the ventral plates of *Acanthomeridion* were treated as their

own character in the matrix (**Fig. 8c**).

3.3 Phylogenetic analyses

All six Bayesian analyses produced the same broad relationships within Artiopoda, with the exception of

the position of *Acanthomeridion* (**Fig. 8**). Petalopleurans were recovered sister to Nektaspida,

Australimicola sister to Conciliterga, and Vicissicaudata was not completely resolved. The positions of

Bailongia, Emeraldella, Kwanyinaspis, Squamacula and *Zhiwenia* were not resolved, and the internal

relationships of Artiopodans were also not well resolved. *Acanthomeridion* was recovered as sister to

Trilobita when the ventral plates were considered homologous to the free cheeks of trilobites, but

otherwise its position was not well resolved. Within Trilobita, when the analysis was unconstrained,

Olenellus and *Eoredlichia* formed a group sister to the other trilobites, or these two taxa were recovered in

a polytomy with a clade containing the other trilobites (**SFigs. 8—10**). When the group *Anacherirurus,*

Cryptolithus, Eoredlichia, Olenoides and *Triarthrus* was constrained (with the exclusion of *Olenellus*),

then *Olenellus* was recovered as sister to all other trilobites (**SFigs. 8—10**). This result is important for

considering the hypothesis that the ventral plates of *Acanthomeridion* were homologous to the doublure of

trilobites.

When visualised in treespace, constrained and unconstrained analyses produce very similar results (**Fig. 9**).

Trees in the posterior sample of the analysis where the ventral plates of *Acanthomeridion* were treated as

homologous to the free cheeks of trilobites fall in the area of treespace supporting *Acanthomeridion* as

sister to trilobites. This result is corroborated by the raw counts in the full sample (Table 1), and by the

support for this node as seen in the consensus tree (**Fig. 9**). Posterior samples of analyses treating ventral

plates as homologous to the doublure and those treating them as non-homologous with any trilobite feature

overlapped in treespace, over a much broader region. These display a very similar confidence ellipse to the

sample of trees where *Acanthomeridion* is not resolved as sister to trilobites, but instead in a different

position in the tree (**Fig. 9**). Again, these treespace results are corroborated by the raw counts in the full

sample (Table 1) and by the poorly resolved position of *Acanthomeridion* in the consensus trees (**Fig. 9**).

	Unconstrained	Constrained
Homologous to free cheeks	7832 / 10504 (75%)	8083/10504
Homologous to doublure	1496 / 12004 (12%)	1836/10504
Not homologous	761 / 12004 (6%)	926/10504

Table 1: number of trees in posterior sample supporting *Acanthomeridion* as sister to trilobites / number of trees in the posterior sample total. Results tabulated for each coding strategy, with unconstrained and constrained analyses (constrained analyses where a monophyletic group of all trilobites except for *Olenellus getzi* was forced, in order to recover trilobite relationships compatible with Paterson et al. 2019).

Fig. 7. Four representative artiopodans from early Cambrian with their dorsal sutures, free cheeks

and ventral plates. (a, b) The iconic trilobite *Wutingaspis tingi* from Chengjiang Biota, note its free

cheeks and dorsal or facial sutures. (c, d) The symbolic xandarellid artiopodan *Xandarella spectaculum*

from Chengjiang Biota bearing the distinctive dorsal sutures. (e, f) Holotype of the protosuturan artiopodan

Zhiwenia coronata from Xiaoshiba Biota developing dorsal sutures. (g) The left dorsal suture of

Acanthomeridion serratum from Chengjiang Biota, showing the morphological and positional similarities

to that of *W. tingi* (a, b), *X. spectaculum* (c, d), and *Z. coronata* (e, f). (h, i) Right ventral plates of *A.*

serratum from Chengjiang Biota bearing a terminal spine, which is similar to free cheek of trilobite like *W.*

tingi (a, b).

4. Discussion and conclusion

All consensus trees, of parsimony and Bayesian results from both coding strategies, support multiple

origins of cephalic sutures in Artiopoda (**Fig. 8, SFigs. 8—10**). When the sutures of *Acanthomeridion* and

trilobites are considered as homologous (e.g., Hou et al., 2017), as well as to those of other artiopodans

(e.g., Du et al., 2019), terminals with ecdysial sutures in the cephalon are found in multiple groups in

Artiopoda (**Fig. 8a**), although *Acanthomeridion* is recovered sister to trilobites. If instead the sutures in

Acanthomeridion are considered marginal, and homologous to those of olenelline trilobites,

Acanthomeridion and trilobites do not form a clade (**Fig 8b, Table 1**), and trees occupy a different region

of the treespace (**Fig. 9**). Indeed, the posterior samples of analyses treating the ventral plates as not

homologous to any other artiopodan feature (**Fig 8c**) are extremely similar to those treating them as

homologous to the doublure of trilobites, as illustrated by their broad overlap in treespace (**Fig. 9**) and

similar support for nodes in the consensus trees (**SFigs. 8—10**). Thus, two of the three coding strategies

receive some support from the phylogenetic results. Either *Acanthomeridion* shares the presence of facial

sutures with trilobites, and these form a clade, but the sutures in petalopleurans and *Zhiwenia* are distinct

(multiple origins of dorsal ecdysial sutures) or the ventral lateral plates of *Acanthomeridion* are a distinct

feature to the free cheeks of trilobites (and petalopeluran and *Zhiwenia* features are also distinct; multiple

origins of sutures). The position of *Acanthomeridion* as not sister to trilobites in the analyses treating the

ventral plates as homologous to doublure mean that this hypothesis did not gain support, given the

assumptions and character matrices of this study.

Support for *Acanthomeridion* as sister to trilobites comes from morphological similarity between the

spinose termination of the ventral plates of *Acanthomeridion* (similar to trilobite genal spines) and the

suture that follows an oblique line and passes near the eye both for *Acanthomeridion* (similar to trilobite

opisthoparian facial sutures; Whittington et al., 1997). Taken together with trilobate exoskeleton of large

Acanthomeridion specimens, the non-biomineralised *Acanthomeridion* bears many morphological

similarities to biomineralised trilobites that further support a sister relationship. However, this sister

relationship rests on the treatment of the ventral plates as free cheeks. If this interpretation is not favored,

then these similarities can be considered convergent, and the position of *Acanthomeridion* within

Artiopoda is not well resolved by these analyses. For this interpretation of the ventral plates, the path of the

suture between the ventral plates and cephalon passing close to the eye may represent a way to minimise

damage to the eye during ecdysis, which is comparable to what is observed in trilobites with facial sutures.

Fig. 8. Simplified results of phylogenetic analyses. Comparison of results from matrices with different coding strategies, with coding of key characters for terminals within the analysis indicated by colored circles. (a) Ventral plates considered homologous to free cheeks. (b) Ventral plates considered homologous to trilobite doublure. (c) Ventral plates not considered homologous to any artiopodan character.

Other features observed in *Acanthomeridion* for the first time – uniramous deutocerebral appendages, an

axe-shaped conterminant hypostome, two librigenal-like plates and three additional head appendages – are

consistent with a phylogenetic position for the species as an early diverging artiopodan, or as sister to

trilobites. Multipodomerous uniramous deutocerebral antennae are known in nearly all other members of

the clade (Zhang et al., 2007; Du et al., 2019; Zhai et al., 2019), and thus this feature is not very

informative. Similarly, a conterminant hypostome was already known in numerous other artiopodans

(Zhang et al., 2004; Stein and Selden, 2012). As this type of hypostomal attachment is considered ancestral

within Trilobita (Fortey, 1990), its presence in the possible sister to trilobites (if dorsal sutures are

considered homologous) is consistent with these expectations from trilobite anatomy. If *Acanthomeridion*

is instead an early diverging artiopodan, the presence of natant hypostomes in both trilobitomorphs and

vicissicaudates (Fortey, 1990; Chen et al., 2019) demonstrates that a natant hypostome has most likely

evolved repeatedly within Artiopoda.

In summary, new fossils and CT-scan data reveals the ventral anatomy and appendages of

Acanthomeridion serratum for the first time, and additional specimens demonstrate that *A. 'anacanthus'*

represents a junior synonym of *A. serratum,* with some features thought to distinguish the two instead

ontogenetic in origin. The presence of a posteriorly orientated spine on the cephalic ventral plates of

Acanthomeridion might be homologous to the genal spine of trilobite librigenae, providing additional

support for a.sister group relationship between these taxa. Phylogenetic analyses interrogating this

hypothesis recover a sister relationship between *Acanthomeridion* and trilobites, but still require multiple

origins of cephalic sutures within the group. If the ventral plates are instead treated as homologous to the

doublure of trilobites, or as not homologous to any trilobite feature, then no close relationship is recovered,

and the morphological similarities between *Acanthomeridion* and trilobites such as the path of the suture

separating the ventral plates from the cephalon, and details of the ventral plates, should instead be treated

as convergent. Thus, regardless of how the ventral plates of *Acanthomeridion* are interpreted, multiple

origins of cephalic sutures in artiopodans are recovered by phylogenetic analyses.

Figure 9. Treespace visualisation of Bayesian Inference phylogenetic analyses. Multidimensional

treespace plotted by bipartition relating to coding strategies (color of point). (a) Analyses were

unconstrained. (b) Trilobites were constrained so that *Olenellus getzi* was resolved as the earliest diverging

member of the group. Area included within a 95% confidence interval (CI) indicated by points inside the

line of different colors. Area where *Acanthomeridion* is recovered sister to trilobites (95% CI) also shown

by area within black line. Where *Acanthomeridion* is not recovered sister to trilobites (95% CI), shown by

area within grey line. Note the overlap in the areas of the black and purple lines.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

The comments and suggestions of three anonymous reviewers greatly improved the manuscript. We thank Jian Han and Jie Sun for scanning the fossils and Javier Ortega-Hernández for helpful discussion. This work was supported by the National Natural Science Foundation of China (grant numbers 42262004, 42202003, 41662003), State Key Laboratory of Palaeobiology and Stratigraphy (Nanjing Institute of Geology and Palaeontology, CAS) (No. 193104). Stephen Pates acknowledges support from a Herchel Smith Postdoctoral Fellowship (University of Cambridge) and NERC Fellowship NE/X017745/1.

CRediT authorship contribution statement

Kunsheng Du: Conceptualization, Data Curation, Investigation, Methodology, Visualization, Writing – original draft, Writing – review and editing. **Jin Guo:** Resources, Data curation, Validation. **Sarah R. Losso:** Conceptualization, Data Curation, Formal analysis, Writing – original draft, Writing – review and editing. **Stephen Pates:** Conceptualization, Data Curation, Investigation, Formal analysis, Methodology, Visualization, Writing – original draft, Writing – review and editing, Supervision. **Ming Li:** Methodology, Data curation. **Ailin Chen:** Resources, Funding acquisition, Project administration, Visualization, Supervision.

References

Aguinaldo, A.M.A., Turbeville, J.M., Linford, L.S., Rivera, M.C., Garey, J.R., Raff, R.A., Lake, J.A.,

1997. Evidence for a clade of nematodes, arthropods and other moulting animals. Nature. 387, 489-

493.

Bapst, D.W., Schreiber, H.A., Carlson, S.J., 2018. Combined analysis of extant Rhynchonellida

(Brachiopoda) using morphological and molecular data. Syst. Biol. 67, 32-48.

Beecher C.E. 1897 Outline of a natural classification of the trilobites, part I. American Journal of Science

3, 89-106.

Beecher C.E. 1897 Outline of a natural classification of the trilobites, part II. American Journal of Science

3, 181-207.

Bicknell, R.D.C., Holmes, J.D., Edgecombe, G.D., Losso, S.R., Ortega-Hernández, J., Wroe, S., Paterson,

J.R., 2021. Biomechanical analyses of Cambrian euarthropod limbs reveal their effectiveness in

mastication and durophagy. Proc. Roy. Soc. B. Biol. Sci. 288, 20202075.

Brusca, R.C., Moore, W., Shuster, S.M., 2016. Invertebrates, Third 3rd Edition. Sinauer Associates Inc.,

Sunderland, Massachusetts U.S.A, 711-961 pp.

Budd, G.E., Telford, M.J., 2009. The origin and evolution of arthropods. Nature. 457, 812-817.

Chen, J.Y., 2004. The dawn of animal world. Jiangsu Science and Technology Publishing House, Nanjing,

365 pp [in Chinese].

Chen, J.Y., Zhou, G.Q., Zhu, M.Y., Yeh, K.Y., 1996. The Chengjiang biota: a unique window of the

Cambrian explosion, Taichung, Taiwan, 222 pp [in Chinese].

Chen, X.H., Ortega-Hernández, J., Wolfe, J.M., Zhai, D.Y., Hou, X.G., Chen, A.L., Mai, H.J., Liu, Y.,

2019. The appendicular morphology of *Sinoburius lunaris* and the evolution of the artiopodan clade

Xandarellida (Euarthropoda, early Cambrian) from South China. BMC Evol. Biol. 19, 165.

Daley, A.C., Antcliffe, J.B., Drage, H.B., Pates, S., 2018. Early fossil record of Euarthropoda and the

Cambrian Explosion. Proc. Natl. Acad. Sci. USA. 115, 5323-5331.

Daley, A.C., Drage, H.B., 2016. The fossil record of ecdysis, and trends in the moulting behaviour of

trilobites. Arthropod Struct. Dev. 45, 71-96.

Drage, H.B., 2019. Quantifying intra-and interspecific variability in trilobite moulting behaviour across the

Palaeozoic. Palaeontol. Electron. 22, 1-39. https://doi.org/10.26879/940

Drage, H.B., 2022. Trilobite moulting behaviour variability had little association with morphometry.

bioRxiv. https://doi.org/10.1101/2022.12.12.520015

Drage, H.B., Laibl, L., Budil, P., 2018. Postembryonic development of Dalmanitina, and the evolution of

facial suture fusion in Phacopina. Paleobiology. 44, 638-659.

Du, K.S., Bruton, D.L., Yang, J., Zhang, X.G., 2023. An early Cambrian *Sidneyia* (Arthropoda) resolves

the century-long debate of its head organization. Sci. China Earth Sci. 66, 521-527.

Du, K.S., Ortega Hernández, J., Yang, J., Zhang, X.G., 2019. A soft-bodied euarthropod from the early

Cambrian Xiaoshiba Lagerstätte of China supports a new clade of basal artiopodans with dorsal

ecdysial sutures. Cladistics. 35, 269-281.

Du, K.S., Guo, J., Losso, S.R., Pates, S., Li, M., Chen, A. 2023. Multiple origins of cephalic sutures in

trilobites and their relatives. bioRxiv. 2023-10. https://doi.org/10.1101/2023.10.05.560990 [first

version of this article]

Dunbar, C. O. 1925. Antennae in *Olenellus getzi* n. sp. Am. J. of Science. 5, 303—308.

Edgecombe, G.D., 2020. Arthropod origins: integrating paleontological and molecular evidence. Annu.

Rev. Ecol. Evol. Syst. 51, 1-25.

Edgecombe, G.D., Ramsköld, L., 1999. Relationships of Cambrian Arachnata and the systematic position

of Trilobita. J. Paleontol. 73, 263-287.

Emmrich, H.F. 1839. De trilobitis. 56 Berlin : Berolini. Dissertatio petrefactologica etc.

Esteve, J., Marcé Nogué, J., Pérez Peris, F., Rayfield, E., 2021. Cephalic biomechanics underpins the

evolutionary success of trilobites. Palaeontology. 64, 519-530.

Foote, M. 1991. Morphologic patterns of diversification: examples from trilobites. Palaeontology. 24, 461-

85.

Fortey, R.A., 1990. Ontogeny, hypostome attachment and trilobite classification. Palaeontology. 33, 529-

576.

Fortey, R.A., Owens, R.M. 1999. Feeding habits in trilobites. Palaeontology. 42, 429-465.

Fortey, R.A., Whittington, H.B. 1989. Trilobita as a natural group. Historical Biology. 2, 125-138.

Giribet, G., Edgecombe, G.D., 2019. The Phylogeny and Evolutionary History of Arthropods. Curr. Biol.

29, R592-R602.

Hammer, Ø, Harper, D.A.T., Ryan, P.D. 2001. Past: palaeontological statistics software package for

education and data analysis. Palaeontologia electronica 4, 1—9.

Henningsmoen, G. 1975. Moulting in trilobites. Fossils and Strata. 4, 179-200.

Hou, X.G., Aldridge, R.J., Bergström, J., Siveter, D.J., Siveter, D.J., Feng, X.H., 2004. The Cambrian

fossils of Chengjiang, China: the flowering of early animal life, 233. Wiley Online Library, Oxford,

233 pp.

Hou, X.G., Bergström, J., 1997. Arthropods of the Lower Cambrian Chengjiang fauna, southwest China.

Foss. Strat. 45, 1-116.

Hou, X.G., Bergström, J., Feng, X.H., Wang, H.F., Chen, A.L., 1999. The Chengjiang Fauna:

exceptionally well-preserved animals from 530 million years ago. Yunnan Science and Technology

Press, Kunming, 170 pp [in Chinese with English summary].

Hou, X.G., Chen, J.Y., Lu, H.Z., 1989. Early Cambrian new arthropods from Chengjiang, Yunnan. Acta

Palaeontol. Sin. 28, 42-57.

Hou, X.G., Siveter, D.J., Siveter, D.J., Aldridge, R.J., Cong, P.Y., Gabbott, S.E., Ma, X.Y., Purnell, M.A.,

Williams, M., 2017b. The Cambrian fossils of Chengjiang, China: The flowering of early animal life.

John Wiley & Sons, Oxford.

Hou, X.G., Williams, M., Gabbott, S., Siveter, D.J., Siveter, D.J., Cong, P.Y., Ma, X.Y., Sansom, R.,

2017a. A new species of the artiopodan arthropod *Acanthomeridion* from the lower Cambrian

Chengjiang Lagerstätte, China, and the phylogenetic significance of the genus. J. Syst. Palaeontol. 15,

733-740.

Hou, X.G., Williams, M., Sansom, R., Siveter, D.J., Siveter, D.J., Gabbott, S., Harvey, T.H.P., Cong, P.Y.,

Liu, Y., 2019. A new xandarellid euarthropod from the Cambrian Chengjiang biota, Yunnan

Province, China. Geol. Mag. 156, 1375-1384.

Hughes, N.C., Gunderson, G.O, Wheedon, M.J. 1997. Circumocular suture and visual surface of '*Cedaria'*

woosteri (Trilobita, Late Cambrian) from the Eau Claire Formation, Wisconsin. Journal of

Paleontology. 71, 103-107.

Hughes, N.C. 2007. The evolution of trilobite body patterning. Annual Review of Earth and Planetary

Sciences. 35, 401-34.

Ivantsov, A.Y., 1999. Trilobite-like arthropod from the Lower Cambrian of the Siberian Platform. Acta

Palaeontol. Pol. 44, 455-466.

Jell, P.A. 2003. Phylogeny of early Cambrian trilobites. Special Papers in Palaeontology. 70, 45-57.

Jiao, D.G., Du, K.S., Zhang, X.G., Yang, J., Eggink, D., 2022. A new small soft-bodied non-trilobite
artiopod from the Cambrian Stage 4 Guanshan Biota. Geol. Mag. 159, 730-734.

Lerosey-Aubril, R., Zhu, X.J., Ortega-Hernández, J., 2017. The Vicissicaudata revisited-insights from a
new aglaspidid arthropod with caudal appendages from the Furongian of China. Sci. Rep. 7, 1-18.

Lewis, P.O., 2001. A likelihood approach to estimating phylogeny from discrete morphological character
data. Syst. Biol. 50, 913-925.

Lieberman, B.S., 2002. Phylogenetic analysis of some basal early Cambrian trilobites, the biogeographic
origins of the Eutrilobita, and the timing of the Cambrian radiation. J. Paleontol. 76, 692-708.

Liu, Y., Ortega-Hernández, J., Zhai, D.Y., Hou, X.G., 2020. A reduced labrum in a Cambrian great-
appendage euarthropod. Curr. Biol. 30, 3057-3061.

Losso, S.R., Ortega-Hernández, J., 2022. Claspers in the mid-Cambrian *Olenoides serratus* indicate
horseshoe crab – like mating in trilobites. Geology. 50, 897-901.

Luo, H.L., Hu, S.X., Chen, L.Z., Zhang, S.S., Tao, Y.H., 1999. Early Cambrian Chengjiang Fauna From
Kunming Region China. Yunnan Science and Technology Press, Kunming, 181 pp [in Chinese with
English summary].

Moore, J., Willmer, P. 1997. Convergent evolution in invertebrates. Biol. Rev. 72, 1—60.

Moysiuk, J., Caron, J., 2022. A three-eyed radiodont with fossilized neuroanatomy informs the origin of
the arthropod head and segmentation. Curr. Biol. 32, 3302-3316.

Ortega Hernández, J., Legg, D.A., Braddy, S.J., 2013. The phylogeny of aglaspidid arthropods and the
internal relationships within Artiopoda. Cladistics. 29, 15-45.

Palmer, A.R., Repina, L.N. 1993. Through a glass darkly: taxonomy, phylogeny and biostratigraphy of the
Olenellina. The University of Kansas Paleontological Contributions. 3, 1-35.

Paterson, J.R., García-Bellido, D.C., Edgecombe, G.D., 2012, New artiopodan arthropods from the early
Cambrian Emu Bay Shale *Konservat-Lagerstätte* of South Australia. Journal of Paleontology. 86,
340-357.

Pates, S., Wolfe, J.M., Lerosey-Aubril, R., Daley A.C., Ortega-Hernández, J. 2022. New opabiniid
diversifies the weirdest wonders of the euarthropod stem group. *Proceedings of the Royal Society B*.
289:20212093.

Rambaut, A., Drummond, A.J., Xie, D., Baele, G., Suchard, M.A., 2018. Posterior summarization in

Bayesian phylogenetics using Tracer 1.7. Syst. Biol. 67, 901-904.

Ramsköld, L. Edgecombe, G.D. 1991. Trilobite monophyly revisited. Historical Biology. 4, 267-283.

Ramsköld, L. Edgecombe, G.D. 1999. Relationships of Cambrian Arachnata and the systematic position of

Trilobita. Journal of Paleontology. 73, 263-287.

Rasetti, F. 1945. Evolution of the facial sutures in the trilobites *Loganopeltoides* and *Loganopeltis*

(Quebec). American Journal of Science. 243, 44-50.

Rasetti, F. 1948. Cephalic sutures in *Loganopeltoides* and the origin of 'hypoparian' trilobites. Journal of

Paleontology. 22, 25-29.

Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L.,

Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and

model choice across a large model space. Syst. Biol. 61, 539-542.

Salter, J.W. 1864. A monograph of British trilobites. Monographs of the Palaeontographical Society. 16, 1-

83.

Schmidt, M., Hou, X.G., Zhai, D.Y., Mai, H.J., Belojevi, J., Chen, X.H., Melzer, R.R., Ortega-Hernández,

J., Liu, Y., 2022. Before trilobite legs: *Pygmaclypeatus daziensis* reconsidered and the ancestral

appendicular organization of Cambrian artiopods. Philos. T. R. Soc. B. 377.

https://doi.org/10.1098/rstb.2021.0030

Stein, M., Selden, P.A., 2012. A restudy of the Burgess Shale (Cambrian) arthropod *Emeraldella brocki*

and reassessment of its affinities. J. Syst. Palaeontol. 10, 361-383.

Strausfeld, N.J., Hou, X., Sayre, M.E., Hirth, F., 2022. The lower Cambrian lobopodian Cardiodictyon

resolves the origin of euarthropod brains. Science. 378, 905-909.

Stubblefield, C.J. 1936. Cephalic sutures and their bearing on current classifications of trilobites. Bio. Rev.

11, 407—440.

Walcott, C.D., 1911, Middle Cambrian Merostomata. Cambrian Geology and Paleontology II. Smithsonian

Miscellaneous Collections. 57, 17-40.

Whittington, H.B., 1975, Trilobites with appendages from the Middle Cambrian, Burgess Shale, British

Columbia. Fossils and Strata. 4. 97-136.

Whittington, H.B. 1989. Olenelloid trilobites: type species, functional morphology and higher

classification. Philosophical Transactions of the Royal Society of London B. 324, 111-147.

Whittington, H.B. 1990. Articulation and exuviation in Cambrian trilobites. Philosophical Transactions of

the Royal Society of London B. 329, 27-46.

Whittington, H.B., Chatterton, B., Speyer, S.E., Fortey, R.A., Owens, R.M., Chang, W.T., Dean, W.T.,

Jell, P.A., Laurie, J.R., Palmer, A.R., 1997. Treatise on Invertebrate Paleontology, Part O, Arthropoda

1, Trilobita, Revised, pp. 530.

Whittington, H.B., 1989, Olenelloid trilobites: type species, functional morphology and higher

classification. Philos. T. R. Soc. B. 324, 111-147.

Wu, Y., Pates, S., Ma, J., Lin, W., Wu, Y., Zhang, X., Fu, D., 2022. Addressing the Chengjiang

conundrum: A palaeoecological view on the rarity of hurdiid radiodonts in this most diverse early

Cambrian Lagerstätte. Geosci. Front. 13, 101430. https://doi.org/10.1016/j.gsf.2022.101430

Yang, J., Ortega-Hernández, J., Butterfield, N.J., Zhang, X.G., 2013. Specialized appendages in

fuxianhuiids and the head organization of early euarthropods. Nature. 494, 468-471.

Yang, J., Ortega-Hernández, J., Drage, H.B., Du, K.S., Zhang, X.G., 2019. Ecdysis in a stem-group

euarthropod from the early Cambrian of China. Sci. Rep. 9. https://doi.org/10.1038/s41598-019-

41911-w

Yang, J., Ortega-Hernández, J., Legg, D.A., Lan, T., Hou, J.B., Zhang, X.G., 2018. Early Cambrian

fuxianhuiids from China reveal origin of the gnathobasic protopodite in euarthropods. Nat. Commun.

9. https://doi.org/10.1038/s41467-017-02754-z

Zhai, D.Y., Edgecombe, G.D., Bond, A.D., Mai, H.J., Liu, Y., 2019. Fine-scale appendage structure of the

Cambrian trilobitomorph *Naraoia spinosa* and its ontogenetic and ecological implications. P. Roy.

Soc. B-Biol. Sci. 286, 20192371. https://doi.org/10.1098/rspb.2019.2371

Zhang, C.X., Liu, Y., Ortega-Hernández, J., Wolfe, J.M., Jin, C.F., Mai, H.J., Hou, X.G., Guo, J., Zhai,

D.Y., 2023. Three-dimensional morphology of the biramous appendages in *Isoxys* from the early

Cambrian of South China, and its implications for early euarthropod evolution. P. Roy. Soc. B-Biol.

Sci. 290, 20230335. https://doi.org/10.1098/rspb.2023.0335

Zhang, M.Y., Liu, Y., Hou, X.G., Ortega-Hernández, J., Mai, H.J., Schmidt, M., Melzer, R.R., Guo, J.,

2022. Ventral Morphology of the Non-Trilobite Artiopod *Retifacies abnormalis* Hou, Chen & Lu,

1989, from the Early Cambrian Chengjiang Biota, China. Biology. 11, 1235.

https://doi.org/10.3390/biology11081235

Zhang, X.L., Fu, D.J., Dai, T., 2012. A new xandarellid arthropod from the Chengjiang Lagerstätte, Lower

Cambrian of Southwest China. Geobios. 45, 335-338.

Zhang, X.L., Han, J., Zhang, Z.F., Liu, H.Q., Shu, D.G., 2004. Redescription of the Chengjiang arthropod

Squamacula clypeata Hou and Bergström, from the Lower Cambrian, south-west China.

Palaeontology. 47, 605-617.

Zhang, X.L., Shu, D.G., Erwin, D.H., 2007. Cambrian naraoiids (Arthropoda): morphology, ontogeny,

systematics, and evolutionary relationships. J. Paleontol. 81, 1-52.