Multiple origins of dorsal ecdysial sutures in trilobites and their relatives

Kun-sheng Du [a,b,1], Jin Guo [b,c,1], Sarah R. Losso [d ✉], Stephen Pates [e, f,✉], Ming Li [g] and Ai-lin Chen

[a,h,✉]

[a] Research Center of Paleobiology, Yuxi Normal University, Yuxi 653100, China.

[b] Key Laboratory for Palaeobiology and MEC International Joint Laboratory for Palaeoenvironment,

Institute of Palaeontology, Yunnan University, Kunming 650091, China.

[c] Management Committee of the Chengjiang Fossil Site World Heritage, Chengjiang, China.

[d] Museum of Comparative Zoology and Department of Organismic and Evolutionary Biology,

Harvard University, Cambridge, Massachusetts 02138, USA

[e] Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Penryn, Cornwall,

TR10 9FE, UK

[f] Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, UK

[g] Institute of Geology, Chinese Academy of Geological Sciences, Beijing 100037, China.

[h] State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and

Paleontology, CAS, Nanjing, 210008, China.

✉e-mail: ailinchen@yxnu.edu.cn (A.L.C.), sarahlosso@g.harvard.edu (S.R.L.) or

s.pates@exeter.ac.uk (S.P.).

[1] Joint first authors.

Keywords: Cambrian; arthropod; Chengjiang Biota; ecdysis; dorsal suture

Abstract

Euarthropods are an extremely diverse phylum in the modern, and have been since their origination in

the early Palaeozoic. They grow through moulting the exoskeleton (ecdysis) facilitated by breaking

along lines of weakness (sutures). Artiopodans, a group that includes trilobites and their non-

biomineralizing relatives, dominated arthropod diversity in benthic communities during the

Palaeozoic. Most trilobites – a hyperdiverse group of tens of thousands of species - moult by breaking

the exoskeleton along cephalic sutures, a strategy that has contributed to their high diversity during

the Palaeozoic. However, the recent description of similar sutures in early diverging non-trilobite

artiopodans mean that it is unclear whether these sutures evolved deep within Artiopoda, or

convergently appeared multiple times within the group. Here we describe new well-preserved material

of *Acanthomeridion*, a putative early diverging artiopodan, including hitherto unknown details of its

ventral anatomy and appendages revealed through CT scanning, highlighting additional possible

homologous features between the ventral plates of this taxon and trilobite free cheeks. We used three

coding strategies treating ventral plates as homologous to trilobite free cheeks, to trilobite cephalic

doublure, or independently derived. If ventral plates are considered homologous to free cheeks,

Acanthomeridion is recovered sister to trilobites however dorsal ecdysial sutures are still recovered at

many places within Artiopoda. If ventral plates are considered homologous to doublure or non-

homologous, then *Acanthomeridion* is not recovered as sister to trilobites, and thus the ventral plates

represent a distinct feature to trilobite doublure/free cheeks.

1. Introduction

Euarthropods, more specifically members of the clade Artiopoda, dominated diversity in Palaeozoic animal communities (Daley et al., 2018). This group, which includes trilobites and their non-biomineralizing relatives, are united by a morphology consisting of a headshield covering a pair of uniramous antennae and at least three pairs of biramous limbs, followed by a dorsoventrally flattened body with a series of biramous appendages (Stein and Selden, 2012; Giribet and Edgecombe, 2019). The origins of key morphological features facilitating the diversity of trilobites can be found in their non-biomineralizing artiopodan relatives (e.g. Du et al., 2019; Schmidt et al., 2022). The cephalon, as an informative and functionally specialized body region, affords pivotal data for resolving the phylogenetic relationships and evolutionary history of arthropods, artiopodans, and trilobites in particular (e.g. Stubblefield 1936; Lieberman 2002; Budd and Telford, 2009; Yang et al., 2013; Paterson et al. 2019; Strausfeld et al., 2022).

The presence, shape, and character of sutures in the trilobite cephalon have been central to classification schemes of the group for nearly 200 years (e.g. Emmrich 1839; Salter 1864; Beecher 1897a, b; Stubblefield 1936; Rasetti 1945; Whittington et al. 1997), while the similarity in the cranidial outlines of Cambrian suture-bearing groups supported a single origin or strong evolutionary constraint on its origins (Foote 1991; Hughes 2007). These sutures provided lines of weakness along which the cephalon split during moulting, creating an anterior ecdysial gape through which the animal exited the old exoskeleton (e.g. Henningsmoen 1975; Whittington 1990; Daley and Drage, 2016). Sutures may have served additional purposes (Stubblefield 1936) and were under additional functional demands, including feeding (Fortey & Owens 1999) and burrowing behaviours (Esteve et al. 2021). Different trilobite clades display distinct suture patterns, with two sets, the circumocular and marginal sutures, relevant for the creation of this ecdysial gape (Henningsmoen 1975). For the earliest diverging trilobites, olenellines (Palmer & Repina 1993; Lieberman 2002) which a have been recovered as a paraphyletic grade rather than a monophyletic group (e.g. Paterson et al. 2019) circumocular sutures facilitated shedding of the cornea during ecdysis (e.g. Ramskold & Edgecombe 1991) while a marginal suture (**Fig. 1e**) facilitated anterior egression through separation of the lateral

cephalic doublure from the dorsal cephalon (e.g. Stubblefield 1936; Henningsmoen 1975). In later

diverging trilobites the circumocular and marginal sutures combined into a single system (**Fig. 1d**).

Here the cornea is fused to the free cheek (e.g. Stubblefield 1936; Ramsköld & Edgecombe 1991) and

these facial sutures – the fused circumocular and marginal sutures - facilitated both ecdysis of the

visual surface and anterior egression of the exoskeleton following withdrawal from the free cheeks

(e.g. Henningsmoen 1975).

Ventral structures
Eye
Homologous with librigenae
Homologous with doublure
Non-homologous

Circumocular suture
Dorsal suture
Proposed facial suture
Marginal suture
Proposed marginal suture

Fig. 1 Hypotheses of homology in sutures between *Acanthomeridion* and trilobites. (a) Cephalic morphology of *Acanthomeridion serratum*. (b) Cephalic morphology of a proetid trilobite (modified from Daley and Drage, 2016). (c) Cephalic morphology of a redlichid trilobite (modified from Whittington 1989). (d) Ventral plates and librigenae as homologous with dorsal suture. (e) Ventral plates and doublure as homologous with marginal suture. (f) Non-homology between ventral plates of *Acanthomeridion*, librigenae and doublure.

The origin of these fused dorsal ecdysial sutures – facial sutures – has traditionally been considered to

fall within Trilobita (e.g. Fortey and Whittington, 1989; Edgecombe and Ramsköld, 1999; Lieberman,

2002). However, the presence of dorsal ecdysial sutures in earlier diverging artiopodans – the

Protosutura (Du et al. 2019) and eye slits in Petalopleura (Chen et al., 2019) raised questions of the

homology and origins within Artiopoda (e.g. Hou et al. 2017; Du et al. 2019). An alternative

hypothesis for the origins of the dorsal cephalic sutures emerged: that these had a deep root within

Artiopoda and were subsequently lost in some groups including multiple times within Trilobita (Hou

et al., 2017a; Du et al., 2019), rather than the traditional view that dorsal cephalic sutures in trilobites

were derived within the clade, and thus these eye slits and facial sutures were acquired independently.

Support for the deep root hypothesis comes from the variability of facial sutures in trilobites, and the

recognition that convergence and loss of features is common when they have a function or allow

adapation to a particular niche (e.g. Moore & Wilmer 1997). Further evidence for the variability of

suture morphology comes from ontogenetic studies, such as the fusion of dalmanitinid facial sutures

during ontogeny (Drage et al., 2018), and the loss of this feature in other trilobite such as '*Cedaria*'

woosteri (Hughes et al. 1997). This indicates that there is scope for a facial suture to have been lost

repeatedly in artiopodan groups earlier diverging than Trilobita, should this feature prove to have a

deeper origin within Artiopoda (Hou et al. 2017; Du et al. 2019). A deep root for facial sutures within

Artiopoda would have repercussions for the importance of the facial sutures in determining trilobite

relationships (e.g. Jell 2003) and the position of a grade of olenelline trilobites as the earliest

diverging members of the group (e.g. Paterson et al. 2019). To date, a third possibility – that the

ventral plates of *Acanthomeridion* are homologous to the doublure of olenellids, and thus

Acanthomeridion and olenelline trilobites share presence of an unfused marginal suture – has not

received broad attention in the literature, but should also be considered in light of the similarities in

position of the ventral plates and olenelline doublure, and the marginal rather than dorsal position of

the suture in *Acanthomeridion.*

Resolving the complex history of dorsal ecdysial sutures evolution within Artiopoda, requires the

nature of cephalic sutures in artiopodans such as *Acanthomeridion,* protosuturans, and petalopleurans

to be resolved, and their phylogenetic position confidently determined (Hou et al., 2017a; Du et al.,

2019). However, the morphology of many earlier diverging artiopodans is not completely understood

(Ortega Hernández et al., 2013; Lerosey-Aubril et al., 2017), contributing a significant amounts of

missing data to character matrices used in phylogenetic analyses. Up to now, *Acanthomeridion* was

arguably the least well known of these early diverging artiopodans, with its ventral anatomy and non-

biomineralized structures very poorly known (Hou and Bergström, 1997; Hou et al., 2017a).

Here we describe new specimens of *Acanthomeridion serratum*, using micro-CT to reveal details of

the ventral anatomy, eyes, and appendages for the first time, and reconstruct the changing shape of the

body during ontogeny. We use these new data to show that *A. anacanthus* is a junior synonym of *A.*

serratum. We then assess the phylogenetic position of the species considering two hypotheses relating

to the dorsal ecdysial sutures – as homologous to the dorsal ecdysial sutures (facial sutures combined

of a fused marginal and circumocular sutures) of trilobites (**Fig. 1d**), that they are homologous to

marginal sutures of olenellines (**Fig. 1e**), and that these structures are unique to *Acanthomeridion*

(**Fig. 1f**), to evaluate the different hypothesis of origination within Artiopoda.

2. Materials and methods

New specimens of *Acanthomeridion* were collected from Jiucun town, Chengjiang (**Fig. 2**) and

accessioned at the Management Committee of the Chengjiang Fossil Site World Heritage (CJHMD

00052–00062), and Research Center of Paleobiology, Yuxi Normal University (YRCP 0016–0020).

New specimens of *Wutingaspis tingi* (YRCP 0021) and *Xandarella spectaculum* (YRCP 0022) were

collected from Haikou town, Kunming (**Fig. 2**) and housed at Research Center of Paleobiology, Yuxi

Normal University. The holotype of *Zhiwenia coronata* (YKLP 12370) is deposited at the Key

Laboratory for Palaeobiology, Yunnan University which was excavated from Xiaoshiba area,

Kunming (**Fig. 2**). The specimens were photographed by a LEICA DFC 550 digital camera mounted

to a Stereoscope LEICA M205 C, and scanned with a Zeiss Xradia 520 Versa X-ray Microscope and

Nikon XTH 225 ST micro-focus X-ray tomography machine. All micro-CT images were processed

with the software Drishti (Zhai et al., 2019). Figures were processed with CorelDRAW X8, Inkscape

1.0 and Illustrator.

Given that specimens with thorax longitudinally curved and specimens with flattened thorax are

considered as two species of *Acanthomeridion*, despite having the same exoskeletal morphology, we

measured their axial length and attempted to determine their true maximum width in all specimens,

including those described in this study and previously documented measurable specimens. The data

were initially analyzed using a bivariate regression fitting function in Microsoft Office Excel.

Subsequently, the software Past (Hammer et al. 2001) was employed for testing and validation,

obtaining the same function and conducting a rational analysis of the function.

The character matrix used for the phylogenetic analyses was adapted from Schmidt et al., (2022),

itself adapted from Du et al. (2019) and Chen et al. (2020). *Olenellus getzi* was added, to facilitate

comparison with olenelline trilobites, while *Anacheirurus adserai*, *Cryptolithus tesselatus* and

Olenoides serratus were added to provide more trilobite terminals. *Olenellus getzi* was chosen as it

represents the olenelline with best known soft tissues (antennae only; Dunbar 1925). Trunk

appendages are known in *A. adserai, C. tesselatus* and *O. serratus* (Walcott, 1911; Whittington, 1975;

Perez-Peris et al., 2021), with further descriptions by Bicknell et al., 2021; Losso and Ortega-

Hernández 2022 for coding the matrix. Our updated dataset includes 70 taxa (*Acanthomeridion*

anacanthus was removed, four trilobites were added). Three matrices were used to test the alternative

hypotheses of sutures, one with 100 characters considering the ventral plates unique, and not

homologous to trilobite librigenae ('ventral plates under head' as a new character in the dataset) and

two with 99 characters. The first of these considered the ventral plates homologous to trilobite

librigenae, and the second to the doublure of olenellines (see Morphobank project). We added the

character state 'ringed distribution lamellae' to Ch. 15 in all matrices (Chen et al., 2019; Du et al.,

2019), and altered the characters relating to the suture pattern (Ch. 31, 32) to include character states

suitable for *Olenellus getzi*. Given the difficulties of identifying circumocular sutures in non-

biomineralizing artiopodans, Ch. 31 indicated the presence or absence of ecdysial sutures in the

cephalon, while Ch. 32 indicated the type of suture. This could either be marginal and/or

circumocular, but unfused (state 0) or marginal and circumocular fused into a facial suture (state 1).

This facilitated differentiation between the state in *Olenellus getzi* and the other trilobites in the

dataset, and thus allowed the two hypotheses relating to homology between the ventral plates and the

doublure or free cheeks of trilobites to be tested. Eight additional characters relating to the trilobite

glabellar region were added, taken from Paterson et al. (2019), in order to provide data to distinguish

internal trilobite relationships. These were Ch. 5, 6, 7, 10, 11, 12, 14 and 15 in Paterson et al. (2019),

which are Ch. 37 – 44 in all three matrices used for this study. A further character (calcified thorax

separated into prothorax and opisthothorax) was also added (Ch. 99 or 100 in matrices used for this

study). Matrices used in phylogenetic analyses are provided through Morphobank (Project #P4290.

Go to morphobank.org, then Log In. Use credentials: Email address: P4290, Password:

Acanthomeridion2023).

Phylogenetic analyses were performed with MrBayes (Bayesian Inference) (Ronquist et al., 2012).

Both the 'maximum information' and 'minimum assumptions' strategies of ref (Bapst et al., 2018)

were utilised on an earlier version of the matrix, in order to confirm that model choice was not a

major driver in differences for the results (Du et al. 2023). For the final version, the 'maximum

information' strategy was used, as model choice did not play a large influence on the topologies

recovered. Bayesian analyses were run for 20 million generations using four chains, every 1000th

sample stored and 25% burn-in (Lewis, 2001; Ronquist et al., 2012). Convergence was diagnosed

using Tracer (Rambaut et al., 2018).

For each coding strategy, two sets of Bayesian analyses were run. The first was unconstrained,

whereas for the second all trilobites except for *Olenellus getzi* were constrained as a monophyletic

group to recover the relationships of trilobites (*Olenellus* as the earliest diverging member)

comparable to the comprehensive trilobite phylogenetic analysis of Paterson et al. (2019). Resulting

posterior tree samples were compared by comparisons in the consensus trees, and in multidimensional

treespace (using R code and method from Pates et al. 2022). The number of trees where

Acanthomeridion serratum was sister to trilobites was quantified for each coding strategy and for

constrained and unconstrained analyses, and the area of treespace occupied by the posterior samples –

and trees where *A. serratum* was sister to triobites within those samples – were visualized.

Fig. 2 Sites yielding *Acanthomeridion* from the Cambrian Stage 3 Chengjiang Biota (red circles).

196 All the specimens of *A. serratum* used here are collected from Jiucun town, Chengjiang County.

3. Results

3.1 Systematic palaeontology

ARTIOPODA Hou and Bergström, 1997

Acanthomeridion Hou et al., 1989

Type and only species. *Acanthomeridion serratum* Hou et al., 1989

Diagnosis. Non-biomineralized artiopodan with elongate dorsal exoskeleton that gives body an elliptical outline. Subtriangular head shield with deep lateral notches that accommodate stalked elliptical eyes, rounded genal angles of the head shield, an axe-like hypostome, and paired teardrop-shaped plates on either side of hypostome. Head bears large eyes and long multi-segmented antennae consisting of over 40 articles anterior to three pairs of small cephalic limbs. Trunk composed of 11 tergites bearing expanded tergopleurae with well-developed distal spines, and a terminal spine. The ninth tergite bears a pleural spine more elongate than others, eleventh tergite reduced and expanded into leaf-like outline. Each tergite bears a pair of biramous limbs with long and dense spines on the endopodites, and slender stick-like exopodites with long and dense bristles. Modified from Hou et al., 1989, 2017a.

Acanthomeridion serratum Hou et al., 1989 **Syn. nov.**

Figs. 3–7; 7g-i; S1–6

1989   *Acanthomeridion serratum* Hou et al., pl. Ⅲ, figs. 1-5; pl. Ⅳ, figs. 1-5; p. 46, text-figs. 3, 4.

1996   *Acanthomeridion serratum* Chen et al., p. 157, figs 199-203.

1997   *Acanthomeridion serratum* Hou and Bergström, p. 38.

1999   *Acanthomeridion serratum* Luo et al., p. 51, pl. 6 fig. 5.

1999   *Acanthomeridion serratum* Hou et al., p. 130, fig. 188.

2004   *Acanthomeridion serratum* Chen, p. 280, fig. 442.

2004   *Acanthomeridion serratum* Hou et al., p. 176, fig. 16.63; p. 177, fig. 16.64.

2017a   *Acanthomeridion serratum*, *A. anacanthus,* and *Acanthomeridion* sp in Hou et al., p. 734,

figs. 1-4.

2017b   *Acanthomeridion serratum* Hou et al., p. 204, fig. 20.34; p. 205, fig. 20.35.

Type material. *Holotype*, CN 108305. *Paratype*, CN 108306–108310.

New material. Sixteen new specimens were collected from Jiucun **(Fig. 2)**, Chengjiang County and

housed in the Management Committee of the Chengjiang Fossil Site World Heritage (CJHMD 00052–

00062), and Research Center of Paleobiology, Yuxi Normal University (YRCP 0016–0020).

Occurrence. Yu'anshan Member, Qiongzhusi Formation, *Wutingaspis–Eoredlichia* biozone,

Cambrian Stage 3. Jiucun and Maotianshan, Chengjiang County, Yunnan, China; Mafang and

Ercaicun, Haikou town, Kunming, Yunnan, China **(Fig. 2)**.

Diagnosis. As for genus, by monotypy.

Description. The dorsal exoskeleton of *Acanthomeridion serratum* displays weak trilobation and

elliptical outline. Specimens measure from 20 to 75 mm along the sagittal axis (**Figs. 3a, c; 4, 5a; 6;**

S1–**6**). The head shield is subtriangular in dorsal view, with rounded anterior margin and small spines

on posterior margin (**Figs. 3a, c; 6; 8a; S6a, c**). A pair of lateral posterior notches accommodate

stalked elliptical eyes. The head shield becomes wider relative to its length during ontogeny (**Fig. 6**).

The lateral margin of the head shield is a suture that attaches paired ventral plates. Ventral plates have

a teardrop outline, occupy the entire length of the cephalon and terminate in a posterior spine

(maximum length 3.4 mm; **Fig. S2b**) that projects into the thorax as far as the anteriormost tergite

(**Fig. 4a**). The medial margin is curved towards the axially line and sits adjacent to a conterminant

axe-shaped hypostome (maximum dimensions 9.6 mm x 11.6 mm; **Fig. 3a, b**). Four pairs of

appendages are present in the head (**Figs. 3f**–**j; S5a**). A pair of antennae with at least 43 articles (**Figs.**

5a, b; S2a, f; S4b) sits anterior to three pairs of small cephalic limbs (**Fig. 3f**–**j**). Articles of the

antennae decrease in length and width distally. The preserved part of the antennae are slightly shorter

than the trunk (**Fig. 5**). The cephalic appendages are poorly preserved and only the gross morphology

can be discerned as no individual podomeres are visible. The protopodite of the cephalic limbs are

subtriangular in outline. Endopodites and exopodites are not clearly preserved. Appendage three may

include the endopodite, but no podomeres are visible.

Fig. 3 *Acanthomeridion serratum* **from the Cambrian Stage 3 Chengjiang Biota.** (a, b) CJHMD

00052a/b respectively, individual with hypostome, ventral plates, and 11 thoracic tergites. (c, d)

CJHMD 00053a, showing ventral plate, dorsal sutures, and 11 thoracic tergites. (e–g) YRCP 0016a,

showing the ventral plates, and three post-antennal limbs. (h–k) Showing the post-antennal

appendages under head, gnathobases of trunk limbs, stick-like exopodites with bristles (black arrows),

and endopodites with long spines (white arrows). (f) Micro-CT image of YRCP 0016a; (h, i, k) Micro-

CT images of CJHMD 00053a. Abbreviations: ant, antenna; ca*n*, post-antennal appendage *n* beneath

head; dbl, doublure; ds, dorsal suture; en, endopodites; ex, exopodites; ey, eye; hs, head shield; hy,

hypostome; L, left; lam, lamellae; no, notch; p*n*, podomere *n*; R, right; T*n*, tergite *n*; ts, terminal spine;

vp, ventral plate.

Fig. 4 *Acanthomeridion serratum* **from the Cambrian Stage 3 Chengjiang Biota** (a) CJHMD

00054, complete individual with head, eye, and 11 tergites. (b) Micro-CT image of panel (a) showing

the stick-like exopods. Abbreviations: ex, exopod; ey, eye; es, eyestalk; hs, head shield; T*n*, tergite *n*.

Fig. 5 *Acanthomeridion serratum* from the Cambrian Stage 3 Chengjiang Biota. (a–e) CJHMD

00055, showing the antenna, ventral plate, endopodites with long spines (arrows in d), 11 tergites, and

paddle-like structure (red arrow). (a) Overview of whole specimen. (b) Detail of long left antenna. (c)

Close-up of ventral plate. (d) Details of long spines (arrows) of right endopodites. (e) Close-up of

paddle-like structure (red arrow). Abbreviations same as Figs. 3, 4.

The trunk is composed of 11 tergites which extend laterally into spinose tergopleurae. Tergites are all

subequal in length. Tergites 8-11 curve towards the posterior, and reduce in width progressively, so

that T11 is approximately 25% the width of T8 and 33% the width of T9 (**Figs. 3a, c; 4, 5a; S1**). Each

tergite has small spines on its lateral and posterior margins (**Figs. 4a; 7; S1; S2i, k; S3a, j; S5b, e, f;**

S6a–f). The shape of tergites and the convexity of the trunk changes through ontogeny (**Fig. 7**). In the

smaller specimens, the trunk appears slender from a dorsal view as the body displays a high convexity

(**Fig. 6t**). Pleurae curve ventrally, and only slightly towards the posterior (**Fig. 6a–k**). In larger

specimens, the trunk is less convex than for the smaller specimens (**Fig. 6s**), ventral curvature of

pleurae is decreased, and posterior curvature increased (**Fig. 6o–r**). A linear relationship between

trunk axial length and width is obtained (**Fig. 7**). Where trunk appendages are preserved, each tergite

is associated with a pair of biramous trunk appendages (**Figs. 3a, c, h, k; 4; 5**). The protopodite is sub

triangular with a broad attachment to the body wall and short robust gnathobasal spines along the

medial margin (**Figs. 3h–j; S3a–f, i**). The ventral margin of the protopodite has endites longer than

the gnathobasal spines. The dorsal edge of the protopodite is poorly preserved and does not clearly

show the attachment of the exopodite. Exopodites are slender, stick-like and long (8.3 mm, ca. 3x

protopodite length), bearing two rows of lamellae (**Figs. 3h, i, k; 4**). Exopodites are shorter than the

width of corresponding tergites (**Figs. 3h; 4**). The lamellae appear short but may not be completely

preserved. There is no evidence that the exopodite divided into lobes or has articulations. Endopodites

are composed of seven trapezoidal podomeres. Podomeres 1–6 bear spiniferous endites arranged in

rows, six on pd1, 2, four on pd3, 4, and three on pd5, 6. Podomere seven displays a terminal claw of

three elongate spines (**Figs. 3h, k; 5d**). From the preserved appendages, no significant antero-

posterior differentiation or dramatic size reduction can be discerned (**Figs. 3, 4**). One specimen shows

evidence for paired midgut diverticulae (**Fig. S4a, f**). The body terminates in a long, slender spine,

dorsal to a paired paddle-like structure (**Figs. 5a, e; S5b, g; S6a, e**).

Remarks. The eyes were previously identified as genal projections (see Fig. 1, Fig. 2A, and Fig. 3A in

Hou et al., 2017a) because of the limited available specimens and the posterior location of the eyes.

Acanthomeridion anacanthus Hou et al. 2017a is here proposed to be a junior subjective synonym of

Acanthomeridion serratum Hou et al.1989, an interpretation supported by:

1) The paddle-like structure under the terminal spine **(Figs. 5a, e; S5b, g)** was previously interpreted

as the twelfth tergite. The specimen YKLP 11115 (see Fig.4 in Hou et al., 2017a) which is a similar

size to YKLP 11116 (holotype of *Acanthomeridion anacanthus*) and YKLP 11113 (see Fig.2A, C and

Fig.3A, C in Hou et al., 2017a) show 11 tergites.

2) Specimens CJHMD 00052-55 and YRCP 0016 show the evidently pleural spines of T8 and T9

(Figs. 3a, c; 4a; 5a; S1; S2a, f; S4b, g).

3) The orientation of pleurae makes some specimens appear narrower than others. Specifically, those

where pleurae curve ventrally look more slender than those where pleurae are curved posteriorly.

Specimen YRCP 0017 preserves right ventrally curving pleura and left flat pleura, which lead to its

right pleura narrower than the left **(Figs. 6j, o, s; S2i)**. Moreover, most small specimens (1cm–3cm)

are preserved with ventrally curving pleurae **(Figs. 6a–i; S3a–c; S4a; S5a-c; S6g, i)** and these appear

narrower than the few small specimens with flat pleurae **(Figs. 6j; S4b**. See also Fig.2A–D, and Fig.4

in Hou et al., 2017a). Larger specimens are typically preserved with flattened, posteriorly curving

pleurae.

4) Twenty specimens show a good ontogenetic series of *Acanthomeridion* including some that have

previously have been assigned to '*Acanthomeridion anacanthus*' **(Figs. 6, 7)**. Linear measurements do

not support distinct species of different sizes, nor with different proportions of thorax axial length to

width. The length and width data fit well with a linear equation; most of the data fall within the 95%

forecast (**Fig. 7b**). Data fall close to the fitted line, as indicated by the R^2 value of 0.95. Lastly,

specimens previously assigned to '*A. anacanthus*' are recovered as smaller *A. serratum* specimens

with similar thoracic shape.

The new specimens and CT data presented herein allow the description of cephalic and trunk limbs,

and a conterminant hypostome adjacent to ventral plates in *Acanthomeridion* for the first time

(reconstruction in **Fig. 8**). The presence of a terminal spine on these ventral plates is identified. The

ventral plate is separated from the dorsal part of the cephalon by a suture that runs close to the stalked

eyes. The possible affinities of these ventral plates, and the phylogenetic implications of these, are

discussed below (Subsections 3.2, 3.3). The presence of four appendages in the head (or three in the

head and one underneath the articulation) is comparable to *Zhiwenia* (Du et al., 2019), as well as other

artiopodans including trilobites (Edgecombe and Ramsköld, 1999; El Albani et al., 2024). However,

the morphology of the three non-antenniform cephalic limbs is difficult to determine with only the

protopodite clearly visible. The lack of observed endopodites and exopodites on these three

appendages may be a preservational artefact. Possibly the absence of clear endopodites and

exopodites represents specialization of the cephalic appendages with the structures being significantly

reduced. Other Cambrian artiopodans have been shown to have appendage specialization (Losso and

Ortega-Hernández, 2022) which is frequently found in the cephalon (Chen et al., 2019; Schmidt et al.,

2022), but none show reduction of both the exopodite or endopodite. In *Sinoburius lunaris*, the first

two post-antennal appendages have reduced endopodites with elongated exopodites (Chen et al.,

2019), and in *Pygmaclypeatus daziensis* the exopodite is significantly reduced in the cephalic

appendages (Schmidt et al., 2022).

Recently, the head of *Retifacies abnormalis* was interpreted to develop three pairs of uniramous post-

antennal appendages and following by one biramous appendage pair (Zhang et al., 2022). It would be

unclear the function of an appendage with both distal elements being significantly reduced until more

clear appendages are found.

The exopodites of *Acanthomeridion* also differ substantially from those known in other artiopodans.

Specifically, the stick-like nature of the exopodite, the rows of lamellae combined with the apparent

lack of segmentation, differ from the paddle shaped exopodites thought to characterise the remainder

of the group (e.g. Schmidt et al., 2022, and fig. 4 in Ortega Hernández et al., 2013).

Fig. 6 Ontogenetic series (a–r) of *Acanthomeridion* and their ventrally curling pleurae (s, t). (a–r)
Showing the individuals from smallest to largest with same scale bar. (s) Lateral view of (o), note the
right curling pleurae and left flat pleurae. (t) Lateral view of (f), showing the left curling pleurae.

Fig. 7 Scatterplot of axial length and width of thorax of *Acanthomeridion serratum*. (a) scatterplot

of raw data. (b) data with 95% forecast shown in pink. Specimens previously assigned to *A.*

anacanthus indicated by blue lettering.

Fig. 8. Reconstruction of *Acanthomeridion serratum* in dorsal and ventral view. Only the

protopodite of the head appendages have been observed. There appear to be three post-antennal limbs

with the fourth pair below the cephalic-thoracic boundary.

3.2 Possible affinities of the ventral plates in *Acanthomeridion*.

3.2.1. ventral plates as homologues of trilobite free cheeks

The ventral plates of *Acanthomeridion* have previously been suggested to be homologues to the free

cheeks of trilobites, with the suture between these features and the remainder of the cephalon

interpreted as a facial suture **(Fig. 1d)** (e.g. Hou et al. 2017). Morphological similarities are apparent

between the ventral plates and free cheeks in many trilobites **(Fig. 9a, b)** (Whittington et al., 1997) as

well as the path of suture separating them from the rest of the cephalon. Our new data provide

additional support for interpreting the ventral plates as homologues of the librigenae. The ventral

plates of *Acanthomeridion* are here shown to be transversely broad, tapering posteriorly to a spine, in

a position directly comparable to the genal spine of many trilobites **(Figs. 1d; 9a, b)**. Furthermore, the

ventral plate is separated from the rest of the cephalon by a suture that passes near the compound eye.

This path is similar to the facial suture (fused circumocular and marginal sutures) of many trilobites

especially those bearing opisthoparian facial sutures, which separate the free cheeks from the

cranidium passing through the visual surface **(Fig. 9a, b)** (Whittington et al., 1997). The homology of

eye notches (e.g., *Luohuilinella* and *Zhiwenia* **[Fig. 9e, f]**)*,* and eye slits (e.g. *Phytophilaspis*

[Ivantsov, 1999]**[Fig. 9c, d]**) of petalopleurans, dorsal ecdysial sutures in trilobites and

Acanthomeridion has been suggested previously (Du et al. 2019). These eye slits join the dorsal eye

with the anterolateral margin of the head shield, possibly forming a single suture that facilitated

release of the visual surface and created an anterior gape for the animal to leave the old exoskeleton

during ecdysis. However, the evolution from eye notches to eye slits and finally to the loss of the eye

slit has been inferred, calling into question the affinity between these structures and sutures (Chen et

al. 2019). This hypothesis was tested explicitly in the first matrix, where the ventral plates of

Acanthomeridion were considered homologous to the free cheeks of trilobites. In this analysis, the

dorsal slits and sutures of other artiopodans were also treated as homologous ecdysial sutures in the

cephalon (**Fig. 10a**).

3.2.2. ventral plates as homologues to doublure in olenelloid trilobites

In the above scenario, the ventral plates of *Acanthomeridion* would be homologous to the free cheeks

both dorsally and ventrally in trilobites, including the doublure. However, it is possible that the

ventral plates are only homologous to the doublure, and thus that the suture represents the marginal

suture, rather than a fused facial suture. The difficulty lies in the marginal position of the eyes, which

could support the interpretation that these are fused sutures but might also represent a position that

facilitated ecdysis through a single, unfused, suture. Considered homologous only to the doublure, the

suture in *Acanthomeridion* separating the ventral plates from the cephalon would be homologous to

the unfused, marginal suture in olenelloids rather than the fused facial suture of most other trilobites

(Fig. 1e)   (e.g. Stubblefield 1936).

Few morphological features can be drawn to support this hypothesis beyond the ventral position of the

plates. In particular, the lateral position of the eyes in *Acanthomeridion*, and the path of the suture

close to these eyes, suggests that the suture functioned to release both the ventral plate and the eye.

Although some artiopodans are known with expansive doublure such as *Squamacula buckorum*

(Paterson et al., 2012) this structure usually follows the outline of the margin and lacks spines

(Whittington et al., 1997).

This hypothesis was tested in the second matrix, where the ecdysial suture of *Acanthomeridion* was

considered unfused (**Fig. 10b**). In this scenario, the presence of eye slits and other dorsal sutures in

other artiopodans would not represent homologous structures, as the proposed evolution is from a

ventral-lateral structure to a dorsal one. Thus, the eye slits of the petalopleurans who bear them were

not considered as ecdysial sutures for this analysis.

3.2.3 ventral plates and eye slits of artiopodans as unique structures without homologues in

trilobites

Eye slits of petalopleuran artiopodans as homologues to facial sutures of trilobites has drawn support

from comparisons such as *Loganopeltoides* (Rasetti 1948), where a suture charts a path from the

dorsal eye to the anterolateral margin of the head shield. However, in *Loganopeltoides* this suture

represents the vestiges of the free cheeks (Edgecombe & Ramksold 1999), a derived state within

Trilobita. Given that the features in support of homology between the ventral plates of

Acanthomeridion and trilobite free cheeks, and implicit within this a deep root of the facial suture

within Artiopoda rather than Trilobita, may not represent true homologues (e.g. spines are often

evolved convergently, and the dorsal eye slits of artiopodans have previously been considered unique

characters – see e.g. Edgecombe & Ramskold 1999; Chen et al., 2019) it is important to consider the

hypothesis that these features are unique to *Acanthomeridion*. Thus for this third analysis, the ventral

plates of *Acanthomeridion* were treated as their own character in the matrix (**Fig. 10c**).

3.3 Phylogenetic analyses

All six Bayesian analyses produced the same broad relationships within Artiopoda, with the exception

of the position of *Acanthomeridion* (**Fig. 10**). Petalopleurans were recovered sister to Nektaspida,

Australimicola sister to Conciliterga, and Vicissicaudata was not completely resolved. The positions

of *Bailongia, Emeraldella, Kwanyinaspis, Squamacula* and *Zhiwenia* were not resolved, and the

internal relationships of Artiopodans were also not well resolved. *Acanthomeridion* was recovered as

sister to Trilobita when the ventral plates were considered homologous to the free cheeks of trilobites,

but otherwise its position was not well resolved. Within Trilobita, when the analysis was

unconstrained, *Olenellus* and *Eoredlichia* formed a group sister to the other trilobites, or these two

taxa were recovered in a polytomy with a clade containing the other trilobites (**SFigs. 7—9**). When

the group *Anacherirurus, Cryptolithus, Eoredlichia, Olenoides* and *Triarthrus* was constrained (with

the exclusion of *Olenellus*), then *Olenellus* was recovered as sister to all other trilobites (**SFigs. 7—9**).

This result is important for considering the hypothesis that the ventral plates of *Acanthomeridion* were

homologous to the doublure of trilobites.

When visualised in treespace, constrained and unconstrained analyses produce very similar results

(**Fig. 11**). Trees in the posterior sample of the analysis where the ventral plates of *Acanthomeridion*

were treated as homologous to the free cheeks of trilobites fall in the area of treespace supporting

Acanthomeridion as sister to trilobites. This result is corroborated by the raw counts in the full sample

(Table 1), and by the support for this node as seen in the consensus tree (**Fig. 11**). Posterior samples

of analyses treating ventral plates as homologous to the doublure and those treating them as non-

homologous with any trilobite feature overlapped in treespace, over a much broader region. These

display a very similar confidence ellipse to the sample of trees where *Acanthomeridion* is not resolved

as sister to trilobites, but instead in a different position in the tree (**Fig. 11**). Again, these treespace

results are corroborated by the raw counts in the full sample (Table 1) and by the poorly resolved

position of *Acanthomeridion* in the consensus trees (**Fig. 11**).

	Unconstrained	Constrained
Homologous to free cheeks	7832 / 10504 (75%)	8083/10504
Homologous to doublure	1496 / 12004 (12%)	1836/10504
Not homologous	761 / 12004 (6%)	926/10504

Table 1: number of trees in posterior sample supporting *Acanthomeridion* as sister to trilobites / number of trees in the posterior sample total. Results tabulated for each coding strategy, with unconstrained and constrained analyses (constrained analyses where a monophyletic group of all trilobites except for *Olenellus getzi* was forced, in order to recover trilobite relationships compatible with Paterson et al. 2019).

Fig. 9. Four representative artiopodans from early Cambrian with their dorsal sutures, free cheeks and ventral plates. (a, b) Trilobite *Wutingaspis tingi* from Chengjiang Biota, note its free cheeks and dorsal or facial sutures. (c, d) Xandarellid artiopodan *Xandarella spectaculum* from Chengjiang Biota bearing the distinctive dorsal sutures. (e, f) Holotype of the protosuturan artiopodan *Zhiwenia coronata* from Xiaoshiba Biota developing dorsal sutures. (g) The left dorsal suture of *Acanthomeridion serratum* from Chengjiang Biota, showing the morphological and positional similarities to that of *W. tingi* (a, b), *X. spectaculum* (c, d), and *Z. coronata* (e, f). (h, i) Right ventral plates of *A. serratum* from Chengjiang Biota bearing a terminal spine, which is similar to free cheek of trilobite like *W. tingi* (a, b).

4. Discussion and conclusion

All consensus trees support multiple origins of cephalic sutures in Artiopoda (**Fig. 10, SFigs. 7—9**). When the sutures of *Acanthomeridion* and trilobites are considered as homologous (e.g., Hou et al., 2017), as well as to those of other artiopodans (e.g., Du et al., 2019), terminals with ecdysial sutures in the cephalon are found in multiple groups in Artiopoda (**Fig. 10a**), although *Acanthomeridion* is recovered sister to trilobites. If instead the sutures in *Acanthomeridion* are considered marginal, and homologous to those of olenelline trilobites, *Acanthomeridion* and trilobites do not form a clade (**Fig 10b, Table 1**), and trees occupy a different region of the treespace (**Fig. 11**). Indeed, the posterior samples of analyses treating the ventral plates as not homologous to any other artiopodan feature (**Fig. 10c**) are extremely similar to those treating them as homologous to the doublure of trilobites, as illustrated by their broad overlap in treespace (**Fig. 11**) and similar support for nodes in the consensus trees (**SFigs. 7—9**). Thus, two of the three coding strategies receive some support from the phylogenetic results. Either *Acanthomeridion* shares the presence of facial sutures with trilobites, and these form a clade, but the sutures in petalopleurans and *Zhiwenia* are distinct (multiple origins of dorsal ecdysial sutures) or the ventral lateral plates of *Acanthomeridion* are a distinct feature to the free cheeks of trilobites (and petalopeluran and *Zhiwenia* features are also distinct; multiple origins of sutures). The position of *Acanthomeridion* as not sister to trilobites in the analyses treating the ventral plates as homologous to doublure mean that this hypothesis did not gain support, given the assumptions and character matrices of this study.

Support for *Acanthomeridion* as sister to trilobites comes from morphological similarity between the spinose termination of the ventral plates of *Acanthomeridion* (similar to trilobite genal spines) and the suture that follows an oblique line and passes near the eye both for *Acanthomeridion* (similar to trilobite opisthoparian facial sutures; Whittington et al., 1997). Taken together with trilobate exoskeleton of large *Acanthomeridion* specimens, the non-biomineralised *Acanthomeridion* bears many morphological similarities to biomineralised trilobites that further support a sister relationship. However, this sister relationship rests on the treatment of the ventral plates as free cheeks. If this interpretation is not favored, then these similarities can be considered convergent, and the position of

Acanthomeridion within Artiopoda is not well resolved by these analyses. For this interpretation of

the ventral plates, the path of the suture between the ventral plates and cephalon passing close to the

eye may represent a way to minimise damage to the eye during ecdysis, which is comparable to what

is observed in trilobites with facial sutures.

Fig. 10. Simplified results of phylogenetic analyses. Comparison of results from matrices with

different coding strategies, with coding of key characters for terminals within the analysis indicated

by colored circles. (a) Ventral plates considered homologous to free cheeks. (b) Ventral plates

considered homologous to trilobite doublure. (c) Ventral plates not considered homologous to any

artiopodan character.

Other features observed in *Acanthomeridion* for the first time – uniramous deutocerebral appendages,

an axe-shaped conterminant hypostome, two librigena-like plates and three additional head

appendages – are consistent with a phylogenetic position for the species as an early diverging

artiopodan, or as sister to trilobites. Multipodomerous uniramous deutocerebral antennae are known in

nearly all other members of the clade (Zhang et al., 2007; Du et al., 2019; Zhai et al., 2019), and thus

this feature is not very informative. Similarly, a conterminant hypostome was already known in

numerous other artiopodans (Zhang et al., 2004; Stein and Selden, 2012). As this type of hypostomal

attachment is considered ancestral within Trilobita (Fortey, 1990), its presence in the possible sister to

trilobites (if dorsal sutures are considered homologous) is consistent with these expectations from

trilobite anatomy. If *Acanthomeridion* is instead an early diverging artiopodan, the presence of natant

hypostomes in both trilobitomorphs and vicissicaudates (Fortey, 1990; Chen et al., 2019)

demonstrates that a natant hypostome has most likely evolved repeatedly within Artiopoda.

In summary, new fossils and CT-scan data reveal the ventral anatomy and appendages of

Acanthomeridion serratum for the first time, and additional specimens demonstrate that *A.*

'anacanthus' represents a junior synonym of *A. serratum,* with some features thought to distinguish

the two instead ontogenetic in origin. The presence of a posteriorly orientated spine on the cephalic

ventral plates of *Acanthomeridion* might be homologous to the genal spine of trilobite librigenae,

providing additional support for a.sister group relationship between these taxa. Phylogenetic analyses

interrogating this hypothesis recover a sister relationship between *Acanthomeridion* and trilobites, but

still require multiple origins of cephalic sutures within the group. If the ventral plates are instead

treated as homologous to the doublure of trilobites, or as not homologous to any trilobite feature, then

no close relationship is recovered, and the morphological similarities between *Acanthomeridion* and

trilobites such as the path of the suture separating the ventral plates from the cephalon, and details of

the ventral plates, should instead be treated as convergent. Thus, regardless of how the ventral plates

of *Acanthomeridion* are interpreted, multiple origins of cephalic sutures in artiopodans are recovered

by phylogenetic analyses.

Figure 11. Treespace visualisation of Bayesian Inference phylogenetic analyses. Multidimensional

treespace plotted by bipartition relating to coding strategies (color of point). (a) Analyses were

unconstrained. (b) Trilobites were constrained so that *Olenellus getzi* was resolved as the earliest

diverging member of the group. Area included within a 95% confidence interval (CI) indicated by

points inside the line of different colors. Area where *Acanthomeridion* is recovered sister to trilobites

(95% CI) also shown by area within black line. Where *Acanthomeridion* is not recovered sister to

trilobites (95% CI), shown by area within grey line. Note the overlap in the areas of the black and

purple lines.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

The comments and suggestions of three anonymous reviewers greatly improved the manuscript. We thank Jian Han and Jie Sun for scanning the fossils and Javier Ortega-Hernández for helpful discussion. This work was supported by the National Natural Science Foundation of China (grant numbers 42262004, 42202003, 41662003), State Key Laboratory of Palaeobiology and Stratigraphy (Nanjing Institute of Geology and Palaeontology, CAS) (No. 193104). Stephen Pates acknowledges support from a Herchel Smith Postdoctoral Fellowship (University of Cambridge) and NERC Fellowship NE/X017745/1.

CRediT authorship contribution statement

Kunsheng Du: Conceptualization, Data Curation, Investigation, Methodology, Visualization, Writing – original draft, Writing – review and editing. **Jin Guo:** Resources, Data curation, Validation. **Sarah R. Losso:** Conceptualization, Data Curation, Formal analysis, Writing – original draft, Writing – review and editing. **Stephen Pates:** Conceptualization, Data Curation, Investigation, Formal analysis, Methodology, Visualization, Writing – original draft, Writing – review and editing, Project administration, Visualization, Supervision. **Ming Li:** Methodology, Data curation. **Ailin Chen:** Resources, Funding acquisition, Project administration, Visualization, Supervision.

References

References

Aguinaldo, A.M.A., Turbeville, J.M., Linford, L.S., Rivera, M.C., Garey, J.R., Raff, R.A., Lake, J.A.,

1997. Evidence for a clade of nematodes, arthropods and other moulting animals. Nature. 387,

489-493.

Bapst, D.W., Schreiber, H.A., Carlson, S.J., 2018. Combined analysis of extant Rhynchonellida

(Brachiopoda) using morphological and molecular data. Syst. Biol. 67, 32-48.

Beecher C.E. 1897 Outline of a natural classification of the trilobites, part I. Am. J. Sci. 3, 89-106.

Beecher C.E. 1897 Outline of a natural classification of the trilobites, part II. Am. J. Sci. 3, 181-207.

Bicknell, R.D.C., Holmes, J.D., Edgecombe, G.D., Losso, S.R., Ortega-Hernández, J., Wroe, S.,

Paterson, J.R., 2021. Biomechanical analyses of Cambrian euarthropod limbs reveal their

effectiveness in mastication and durophagy. Proc. Roy. Soc. B. Biol. Sci. 288, 20202075.

Brusca, R.C., Moore, W., Shuster, S.M., 2016. Invertebrates, Third 3rd Edition. Sinauer Associates

Inc., Sunderland, Massachusetts U.S.A, 711-961 pp.

Budd, G.E., Telford, M.J., 2009. The origin and evolution of arthropods. Nature. 457, 812-817.

Chen, J.Y., 2004. The dawn of animal world. Jiangsu Science and Technology Publishing House,

Nanjing, 365 pp [in Chinese].

Chen, J.Y., Zhou, G.Q., Zhu, M.Y., Yeh, K.Y., 1996. The Chengjiang biota: a unique window of the

Cambrian explosion, Taichung, Taiwan, 222 pp [in Chinese].

Chen, X.H., Ortega-Hernández, J., Wolfe, J.M., Zhai, D.Y., Hou, X.G., Chen, A.L., Mai, H.J., Liu,

Y., 2019. The appendicular morphology of *Sinoburius lunaris* and the evolution of the artiopodan

clade Xandarellida (Euarthropoda, early Cambrian) from South China. BMC Evol. Biol. 19, 165.

Daley, A.C., Antcliffe, J.B., Drage, H.B., Pates, S., 2018. Early fossil record of Euarthropoda and the

Cambrian Explosion. Proc. Natl. Acad. Sci. USA. 115, 5323-5331.

Daley, A.C., Drage, H.B., 2016. The fossil record of ecdysis, and trends in the moulting behaviour of

trilobites. Arthropod Struct. Dev. 45, 71-96.

Drage, H.B., 2019. Quantifying intra-and interspecific variability in trilobite moulting behaviour

across the Palaeozoic. Palaeontol. Electron. 22, 1-39. https://doi.org/10.26879/940

Drage, H.B., 2022. Trilobite moulting behaviour variability had little association with morphometry.

bioRxiv. https://doi.org/10.1101/2022.12.12.520015

Drage, H.B., Laibl, L., Budil, P., 2018. Postembryonic development of Dalmanitina, and the evolution

of facial suture fusion in Phacopina. Paleobiology. 44, 638-659.

Du, K.S., Bruton, D.L., Yang, J., Zhang, X.G., 2023. An early Cambrian *Sidneyia* (Arthropoda)

resolves the century-long debate of its head organization. Sci. China Earth Sci. 66, 521-527.

Du, K.S., Ortega Hernández, J., Yang, J., Zhang, X.G., 2019. A soft-bodied euarthropod from the

early Cambrian Xiaoshiba Lagerstätte of China supports a new clade of basal artiopodans with

dorsal ecdysial sutures. Cladistics. 35, 269-281.

Du, K.S., Guo, J., Losso, S.R., Pates, S., Li, M., Chen, A. 2023. Multiple origins of cephalic sutures in

trilobites and their relatives. bioRxiv. 2023-10. https://doi.org/10.1101/2023.10.05.560990 [first

version of this article]

Dunbar, C. O. 1925. Antennae in *Olenellus getzi* n. sp. Am. J. Sci. 5, 303—308.

Edgecombe, G.D., 2020. Arthropod origins: integrating paleontological and molecular evidence.

Annu. Rev. Ecol. Evol. Syst. 51, 1-25.

Edgecombe, G.D., Ramsköld, L., 1999. Relationships of Cambrian Arachnata and the systematic

position of Trilobita. J. Paleontol. 73, 263-287.

El Albani, A., Mazurier, A., Edgecombe, G.D., Azizi, A., El Bakhouch, A., Berks, H.O., Bouougri,

E.H., Chraiki, I., Donoghue, P.C., Fontaine, C., Gaines, R.R., Ghnahalla, M., Meunier, A.,

Trentesaux, A., Paterson, J.R. 2024. Rapid volcanic ash entombment reveals the 3D anatomy of

Cambrian trilobites. Science. 384, 1429—1435.

Emmrich, H.F. 1839. De trilobitis. 56 Berlin : Berolini. Dissertatio petrefactologica etc.

Esteve, J., Marcé Nogué, J., Pérez Peris, F., Rayfield, E., 2021. Cephalic biomechanics underpins the

evolutionary success of trilobites. Palaeontology. 64, 519-530.

Foote, M. 1991. Morphologic patterns of diversification: examples from trilobites. Palaeontology. 24,

461-85.

Fortey, R.A., 1990. Ontogeny, hypostome attachment and trilobite classification. Palaeontology. 33,

529-576.

Fortey, R.A., Owens, R.M. 1999. Feeding habits in trilobites. Palaeontology. 42, 429-465.

Fortey, R.A., Whittington, H.B. 1989. Trilobita as a natural group. Hist. Biol. 2, 125-138.

Giribet, G., Edgecombe, G.D., 2019. The Phylogeny and Evolutionary History of Arthropods. Curr.

Biol. 29, R592-R602.

Hammer, Ø, Harper, D.A.T., Ryan, P.D. 2001. Past: palaeontological statistics software package for

education and data analysis. Palaeontol. Electron. 4, 1—9.

Henningsmoen, G. 1975. Moulting in trilobites. Foss. Strat. 4, 179-200.

Hou, X.G., Aldridge, R.J., Bergström, J., Siveter, D.J., Siveter, D.J., Feng, X.H., 2004. The Cambrian

fossils of Chengjiang, China: the flowering of early animal life, 233. Wiley Online Library,

Oxford, 233 pp.

Hou, X.G., Bergström, J., 1997. Arthropods of the Lower Cambrian Chengjiang fauna, southwest

China. Foss. Strat. 45, 1-116.

Hou, X.G., Bergström, J., Feng, X.H., Wang, H.F., Chen, A.L., 1999. The Chengjiang Fauna:

exceptionally well-preserved animals from 530 million years ago. Yunnan Science and

Technology Press, Kunming, 170 pp [in Chinese with English summary].

Hou, X.G., Chen, J.Y., Lu, H.Z., 1989. Early Cambrian new arthropods from Chengjiang, Yunnan.

Acta Palaeontol. Sin. 28, 42-57.

Hou, X.G., Siveter, D.J., Siveter, D.J., Aldridge, R.J., Cong, P.Y., Gabbott, S.E., Ma, X.Y., Purnell,

M.A., Williams, M., 2017b. The Cambrian fossils of Chengjiang, China: The flowering of early

animal life. John Wiley & Sons, Oxford.

Hou, X.G., Williams, M., Gabbott, S., Siveter, D.J., Siveter, D.J., Cong, P.Y., Ma, X.Y., Sansom, R.,

2017a. A new species of the artiopodan arthropod *Acanthomeridion* from the lower Cambrian

Chengjiang Lagerstätte, China, and the phylogenetic significance of the genus. J. Syst.

Palaeontol. 15, 733-740.

Hou, X.G., Williams, M., Sansom, R., Siveter, D.J., Siveter, D.J., Gabbott, S., Harvey, T.H.P., Cong,

P.Y., Liu, Y., 2019. A new xandarellid euarthropod from the Cambrian Chengjiang biota,

Yunnan Province, China. Geol. Mag. 156, 1375-1384.

Hughes, N.C., Gunderson, G.O, Wheedon, M.J. 1997. Circumocular suture and visual surface of

'*Cedaria' woosteri* (Trilobita, Late Cambrian) from the Eau Claire Formation, Wisconsin. J. Paleontol. 71, 103-107.

Hughes, N.C. 2007. The evolution of trilobite body patterning. Ann. Rev. Earth Planet. Sci. 35, 401-34.

Ivantsov, A.Y., 1999. Trilobite-like arthropod from the Lower Cambrian of the Siberian Platform. Acta Palaeontol. Pol. 44, 455-466.

Jell, P.A. 2003. Phylogeny of early Cambrian trilobites. Special Pap. Palaeontol. 70, 45-57.

Jiao, D.G., Du, K.S., Zhang, X.G., Yang, J., Eggink, D., 2022. A new small soft-bodied non-trilobite artiopod from the Cambrian Stage 4 Guanshan Biota. Geol. Mag. 159, 730-734.

Lerosey-Aubril, R., Zhu, X.J., Ortega-Hernández, J., 2017. The Vicissicaudata revisited-insights from a new aglaspidid arthropod with caudal appendages from the Furongian of China. Sci. Rep. 7, 1-18.

Lewis, P.O., 2001. A likelihood approach to estimating phylogeny from discrete morphological character data. Syst. Biol. 50, 913-925.

Lieberman, B.S., 2002. Phylogenetic analysis of some basal early Cambrian trilobites, the biogeographic origins of the Eutrilobita, and the timing of the Cambrian radiation. J. Paleontol. 76, 692-708.

Liu, Y., Ortega-Hernández, J., Zhai, D.Y., Hou, X.G., 2020. A reduced labrum in a Cambrian great-appendage euarthropod. Curr. Biol. 30, 3057-3061.

Losso, S.R., Ortega-Hernández, J., 2022. Claspers in the mid-Cambrian *Olenoides serratus* indicate horseshoe crab‐like mating in trilobites. Geology. 50, 897-901.

Luo, H.L., Hu, S.X., Chen, L.Z., Zhang, S.S., Tao, Y.H., 1999. Early Cambrian Chengjiang Fauna From Kunming Region China. Yunnan Science and Technology Press, Kunming, 181 pp [in Chinese with English summary].

Moore, J., Willmer, P. 1997. Convergent evolution in invertebrates. Biol. Rev. 72, 1—60.

Moysiuk, J., Caron, J., 2022. A three-eyed radiodont with fossilized neuroanatomy informs the origin of the arthropod head and segmentation. Curr. Biol. 32, 3302-3316.

Ortega Hernández, J., Legg, D.A., Braddy, S.J., 2013. The phylogeny of aglaspidid arthropods and the

internal relationships within Artiopoda. Cladistics. 29, 15-45.

Palmer, A.R., Repina, L.N. 1993. Through a glass darkly: taxonomy, phylogeny and biostratigraphy

of the Olenellina. U. Kansas Paleontol. Contrib. 3, 1-35.

Paterson, J.R., García-Bellido, D.C., Edgecombe, G.D., 2012, New artiopodan arthropods from the

early Cambrian Emu Bay Shale *Konservat-Lagerstätte* of South Australia. J. Paleontol. 86, 340-

357.

Pates, S., Wolfe, J.M., Lerosey-Aubril, R., Daley A.C., Ortega-Hernández, J. 2022. New opabiniid

diversifies the weirdest wonders of the euarthropod stem group. Proc. Roy. Soc. B. Biol. Sci.

289:20212093.

Perez-Peris, F., Laibl, L., Muriel, V., Daley, A.C. 2021. Systematics, morphology, and appendages of

an Early Ordovician pilekiine trilobite *Anacheirurus* from Fezouata Shale and the early

diversification of Cheiruridae. Acta Palaeontol. Pol. 66:857—877.

Rambaut, A., Drummond, A.J., Xie, D., Baele, G., Suchard, M.A., 2018. Posterior summarization in

Bayesian phylogenetics using Tracer 1.7. Syst. Biol. 67, 901-904.

Ramsköld, L. Edgecombe, G.D. 1991. Trilobite monophyly revisited. Hist. Biol. 4, 267-283.

Ramsköld, L. Edgecombe, G.D. 1999. Relationships of Cambrian Arachnata and the systematic

position of Trilobita. J. Paleontol. 73, 263-287.

Rasetti, F. 1945. Evolution of the facial sutures in the trilobites *Loganopeltoides* and *Loganopeltis*

(Quebec). Am. J. Sci. 243, 44-50.

Rasetti, F. 1948. Cephalic sutures in *Loganopeltoides* and the origin of 'hypoparian' trilobites. J.

Paleontol. 22, 25-29.

Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu,

L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: efficient Bayesian phylogenetic

inference and model choice across a large model space. Syst. Biol. 61, 539-542.

Salter, J.W. 1864. A monograph of British trilobites. Monog. Pal. Soc. 16, 1-83.

Schmidt, M., Hou, X.G., Zhai, D.Y., Mai, H.J., Belojevi, J., Chen, X.H., Melzer, R.R., Ortega-

Hernández, J., Liu, Y., 2022. Before trilobite legs: *Pygmaclypeatus daziensis* reconsidered and

701  the ancestral appendicular organization of Cambrian artiopods. Philos. T. R. Soc. B. 377.

702  https://doi.org/10.1098/rstb.2021.0030

703 Stein, M., Selden, P.A., 2012. A restudy of the Burgess Shale (Cambrian) arthropod *Emeraldella*

704  *brocki* and reassessment of its affinities. J. Syst. Palaeontol. 10, 361-383.

705 Strausfeld, N.J., Hou, X., Sayre, M.E., Hirth, F., 2022. The lower Cambrian lobopodian

706  Cardiodictyon resolves the origin of euarthropod brains. Science. 378, 905-909.

707 Stubblefield, C.J. 1936. Cephalic sutures and their bearing on current classifications of trilobites. Bio.

708  Rev. 11, 407—440.

709 Walcott, C.D., 1911, Middle Cambrian Merostomata. Cambrian Geology and Paleontology II. Smith.

710  Misc. Coll. 57, 17-40.

711 Whittington, H.B., 1975, Trilobites with appendages from the Middle Cambrian, Burgess Shale,

712  British Columbia. Foss. Strat. 4. 97-136.

713 Whittington, H.B. 1989. Olenelloid trilobites: type species, functional morphology and higher

714  classification. Philos. T. R. Soc. B. 324, 111-147.

715 Whittington, H.B. 1990. Articulation and exuviation in Cambrian trilobites. Philos. T. R. Soc. B. 329,

716  27-46.

717 Whittington, H.B., Chatterton, B., Speyer, S.E., Fortey, R.A., Owens, R.M., Chang, W.T., Dean,

718  W.T., Jell, P.A., Laurie, J.R., Palmer, A.R., 1997. Treatise on Invertebrate Paleontology, Part O,

719  Arthropoda 1, Trilobita, Revised, pp. 530.

720 Whittington, H.B., 1989, Olenelloid trilobites: type species, functional morphology and higher

721  classification. Philos. T. R. Soc. B. 324, 111-147.

722 Yang, J., Ortega-Hernández, J., Butterfield, N.J., Zhang, X.G., 2013. Specialized appendages in

723  fuxianhuiids and the head organization of early euarthropods. Nature. 494, 468-471.

724 Yang, J., Ortega-Hernández, J., Drage, H.B., Du, K.S., Zhang, X.G., 2019. Ecdysis in a stem-group

725  euarthropod from the early Cambrian of China. Sci. Rep. 9. https://doi.org/10.1038/s41598-019-

726  41911-w

727 Yang, J., Ortega-Hernández, J., Legg, D.A., Lan, T., Hou, J.B., Zhang, X.G., 2018. Early Cambrian

728  fuxianhuiids from China reveal origin of the gnathobasic protopodite in euarthropods. Nat.

Commun. 9. https://doi.org/10.1038/s41467-017-02754-z

Zhai, D.Y., Edgecombe, G.D., Bond, A.D., Mai, H.J., Liu, Y., 2019. Fine-scale appendage structure

of the Cambrian trilobitomorph *Naraoia spinosa* and its ontogenetic and ecological implications.

P. Roy. Soc. B. Biol. Sci. 286, 20192371. https://doi.org/10.1098/rspb.2019.2371

Zhang, C.X., Liu, Y., Ortega-Hernández, J., Wolfe, J.M., Jin, C.F., Mai, H.J., Hou, X.G., Guo, J.,

Zhai, D.Y., 2023. Three-dimensional morphology of the biramous appendages in *Isoxys* from the

early Cambrian of South China, and its implications for early euarthropod evolution. P. Roy. Soc.

B. Biol. Sci. 290, 20230335. https://doi.org/10.1098/rspb.2023.0335

Zhang, M.Y., Liu, Y., Hou, X.G., Ortega-Hernández, J., Mai, H.J., Schmidt, M., Melzer, R.R., Guo,

J., 2022. Ventral Morphology of the Non-Trilobite Artiopod *Retifacies abnormalis* Hou, Chen &

Lu, 1989, from the Early Cambrian Chengjiang Biota, China. Biology. 11, 1235.

https://doi.org/10.3390/biology11081235

Zhang, X.L., Fu, D.J., Dai, T., 2012. A new xandarellid arthropod from the Chengjiang Lagerstätte,

Lower Cambrian of Southwest China. Geobios. 45, 335-338.

Zhang, X.L., Han, J., Zhang, Z.F., Liu, H.Q., Shu, D.G., 2004. Redescription of the Chengjiang

arthropod *Squamacula clypeata* Hou and Bergström, from the Lower Cambrian, south-west

China. Palaeontology. 47, 605-617.

Zhang, X.L., Shu, D.G., Erwin, D.H., 2007. Cambrian naraoiids (Arthropoda): morphology,

ontogeny, systematics, and evolutionary relationships. J. Paleontol. 81, 1-52.