1

A comparative approach for selecting orthologous candidate genes underlying signal in genome-wide association studies across multiple species

Lauren Whitt[1,#a], Elizabeth H. Mahood[1], Greg Ziegler[1], Collin Luebbert[1], Jason D. Gillman[2], Gareth J. Norton[3], Adam H. Price[3], David E. Salt[4], Brian P. Dilkes[5], Ivan Baxter[1*]

[1]Donald Danforth Plant Science Center, St. Louis, Missouri, United States of America

[2]USDA-ARS Plant Genetics Research Unit, Columbia, Missouri, United State of America

[3] School of Biological Sciences, University of Aberdeen, Aberdeen, United Kingdom

[4]Future Food Beacon of Excellence and School of Biosciences, University of Nottingham, Nottingham, United Kingdom

[5]Department of Biochemistry, Purdue University, West Lafayette, Indiana, United States of America

[#a]Plant Science and Technology, University of Missouri, Columbia, Missouri, United States of America

* Corresponding author

Email: ibaxter@danforthcenter.org (IB)

Abstract

Advances in quantitative genetics have enabled researchers to identify genomic regions associated

with changes in phenotype. However, genomic regions can contain hundreds to thousands of genes,

and progressing from genomic regions to candidate genes is still challenging. In genome-wide

association studies (GWAS) measuring elemental accumulation (ionomic) traits, a mere 5% of loci are

associated with a known ionomic gene - indicating that many causal genes are still unknown. To

select candidates for the remaining 95% of loci, we developed a method to identify conserved genes

underlying GWAS loci in multiple species. For 19 ionomic traits, we identified 14,336 candidates

across Arabidopsis, soybean, rice, maize, and sorghum. We calculated the likelihood of candidates

with random permutations of the data and determined that most of the top 10% of candidates were

orthologous genes linked to GWAS loci across all five species. The candidate list also includes

orthologous genes with previously established ionomic functions in Arabidopsis and rice. Our

methods highlight the conserved nature of ionomic genetic regulators and enable the identification

of previously unknown ionomic genes.

Author summary

Identifying the genes contributing to changes in a given trait is challenging. Many genes can be near

the region of interest, but proximity does not always translate to causality. We use other methods to

narrow our focus to the genes most likely involved in our trait of interest before confirming their

involvement through confirmation experiments. However, these other methods are often time,

labor, and resource-consuming. We developed an approach to narrow these gene lists before these

laborious methods are required. Through testing with genetic markers for elemental (i.e., calcium,

iron, zinc) uptake, we found that comparing markers across multiple species for nearby

evolutionarily conserved genes is a successful approach. We've produced a list of candidate genes

likely to be involved in elemental uptake traits, including previously known elemental uptake genes

and genes whose potential elemental uptake function has yet to be observed. Some of these genes

would not have been considered according to the significance threshold within a single GWAS, but

combining comparable datasets across species has collectively boosted their signal. Methods like our

approach are useful for reducing candidate lists to conserve resources spent in functional

characterization experiments and encouraging the discovery of new functional roles of these genes.

Introduction

High-throughput phenotyping (HTP) paired with genome-wide association studies (GWAS) has

identified many genetic loci associated with variation in the accumulation of elements in many plant

species [1–5]. However, identifying the causal gene associated with a group of linked variants, often

called a GWAS peak, is challenging for any trait or organism. Typically, many genes are linked to a

GWAS peak, especially in species with large linkage disequilibrium ranges, and each is a candidate

for the allele affecting variation in the phenotype. Complex loci, multiple alleles, rare alleles, and

incomplete variant information (due to technology or sparse marker data) can also result in the most

statistically significant variant being encoded at a position displaced from the causal gene [6]. As a

result, additional and complementary information about allele consequences and gene function is

necessary to identify the molecular cause of detected quantitative genetic variation. Bridging these

trait-to-gene gaps is necessary to deliver molecular mechanistic insights into natural variation.

Current gene selection methods range from quick functional annotation checks to more extensive

experimental confirmations. Gene ontologies (GO) and annotations provide knowledge of our

current understanding of known (or predicted) genes but are prone to human biases [7,8] and

oversights from automatic computational algorithms [9]. Excluding a supremely significant GWAS

peak where the single nucleotide polymorphism (SNP) resides within a coding region, known genes

are more likely to be selected for further studies than unknown genes [7,10]. Additionally, these

methods are most effective in species with the most annotations, like *Arabidopsis thaliana*

(Arabidopsis), with well-annotated phenotypes. GO terms are largely uninformative for novel or

under-described traits, like some elemental accumulation phenotypes. GO terms require ongoing

curation and submission from researchers to maintain relevance and utility.

More in-depth approaches look for candidates with additional signals that indicate their candidacy,

regardless of a gene's existing annotations or ontologies. For example, linkage mapping in

experimental populations has been used to differentiate causal candidates in trait-associated GWAS

peaks [11,12]. However, fine mapping to find a causal gene requires expensive and laborious

methods (for example, high-resolution sequencing and cloning) [13,14] and will not work for alleles

with moderate effect sizes. Candidate gene testing by working with knockout and overexpression

mutants is an alternative approach to differentiate candidates from causal loci. Despite advances in

gene editing, these methods are much easier in species like Arabidopsis and *Zea mays* (maize) with

substantial mutant resources. Model systems like Arabidopsis also benefit from short generation

times and small growth requirements [15]. Developments in whole genome and next-generation

sequencing have improved *de novo* genome assembly of organisms with complex genomes and

identification of genetic markers (i.e., SNPs, indels) within populations [16]. Now, most agriculturally

important crop species have genome assembly and annotations. Still, Arabidopsis has the highest

quality plant genome, annotations, and resources that inform annotations in other organisms.

Indeed, for several agricultural species (e.g., soybean), gene annotations are inferred from homology

searches against Arabidopsis. Several approaches have been used to identify causal genes from

genes of unknown function. Guided methods that can work without gene annotations include

machine learning techniques like PILGRM [17], GenAMap [18], network analysis, gene expression

experiments, or a combination of techniques [19]. These methods rely on extensive computation

and resources like gene expression data. As a result, they are most valuable in model systems with

large collections of preexisting data.

The ionome describes the composition of elements in plant tissue acquired from its environment

[20]. The challenge of obtaining, excluding, and homeostatically maintaining the concentration of

elements in cells is a key function shared by organisms across all domains of life. Elements obtained

from the environment are critical cofactors and substrates for multiple biochemical reactions, and

solving these challenges was a necessary step in the evolution of cellular life. As a result, many of the

mechanisms that achieve elemental homeostasis are ancient, and experimental approaches for

mechanistic validation, such as cross-kingdom molecular complementation (e.g., yeast

complementation, bacterial complementation), are highly effective. Moreover, as elements are the

same regardless of which species they are measured in, we can directly compare elemental

accumulation traits across species. As a result, the field has repeatedly demonstrated that

homologous genes affect the levels of the same elements in multiple plants [21–30]. The

independent evolution of alleles with similar phenotypic consequences in multiple species is the

hallmark of conserved critical regulators. The direct comparability of ionomics traits and the deep

conservation of mechanisms affecting its variation raises the possibility that we could identify

homologous genes affecting variation in ionomic traits linked to GWAS peaks across multiple

species.

We created a computational method to take the genes linked to GWAS data sets from multiple

species and detect orthologous genes when present among the linked candidates. With this, we built

a permutation-based assessment of the effectiveness of this tool that calculates the expected

background distribution of orthologs for each collection of GWAS results. Together, this

demonstrated that GWAS detects alleles at orthologous genes more often than expected by chance,

including genes of previously unknown function in regulating the ionome. We further validated

these results using lists of genes of known ionomic function [24] and demonstrated that these are

also detected by our combination of GWAS and orthology at a higher rate than expected by chance.

One key advantage of our approach is that it is powered by phylogenetic analysis and, therefore,

detects high-quality candidate genes for traits without requiring prior demonstration of function.

This approach utilizes existing datasets and encourages exploration into novel functional

annotations.

Results

Known Ionomic Genes as GWAS candidates

Elemental accumulation, or ionomic, datasets from association mapping populations of five species

(Table 1): Arabidopsis [31], *Glycine max* (soybean; see methods), *Oryza sativa* (rice) *[32–35]*,

Sorghum bicolor (sorghum) [1], and maize [3] were reanalyzed to identify alleles at conserved

orthologous genes. These GWAS were selected because they are publicly available and contain 19

independent traits measured with the same type of instrument (ICP-MS). In all cases, element

accumulation was measured in the seed, and in all cases except Arabidopsis, data sets were

generated from plants grown in multiple environments. SNPs were filtered by p-values specific to

each dataset (see methods) and, when applicable, the SNP's significance across multiple

environments (Table 1). Loci from the maize NAM dataset from Ziegler et al. 2017 were filtered

more stringently than the other GWAS datasets due to the large number of SNPs. General linkage

ranges for each dataset were used to determine loci and linked genes.

Table 1. Ionomic loci from GWAS across five species.					
	Arabidopsis	soybean	rice	sorghum	maize
Environments	single	multiple	multiple	multiple	multiple
boron (B)	0	401	0	63	10
sodium (Na)	118	331	40	53	0
magnesium (Mg)	65	355	10	163	54
phosphorus (P)	58	372	14	87	39
sulfur (S)	57	362	13	78	54

potassium (K)	49	380	12	93	53
calcium (Ca)	54	360	16	102	43
iron (Fe)	55	354	20	210	67
manganese (Mn)	59	380	15	175	72
cobalt (Co)	149	373	0	106	105
nickel (Ni)	0	374	0	73	56
copper (Cu)	53	353	26	82	72
zinc (Zn)	51	384	24	103	58
arsenic (As)	68	360	20	0	28
selenium (Se)	126	359	0	114	7
rubidium (Rb)	72	352	0	53	59
strontium (Sr)	60	387	0	0	37
molybdenum (Mo)	109	362	70	115	73
cadmium (Cd)	114	381	152	98	11
Total loci	1404	7308	432	1768	905

To help identify candidate genes underlying elemental uptake loci in plants, we had previously

compiled a list of known ionomic genes (KIG) experimentally shown to alter elemental uptake in

plants [24]. As these genes were sourced from the primary literature, we called this our "primary"

KIG list. Since the original publication, we have added 19 new primary genes to the list (S1 Table), for

a total of 201 primary genes. Most primary genes are in Arabidopsis (59%), with smaller

contributions from rice (33%) and a small number in maize, *Medicago truncatula*, and *Triticum*

aestivum (wheat). The overlap between the identified GWAS loci and the KIG list was used to

determine if these GWAS datasets identified alleles at genes already known to affect changes in the

ionome (Fig 1). Loci and gene overlaps were considered on a trait-by-trait basis, i.e., Fe uptake GWAS

loci were only compared to KIG genes listed for Fe. Permutations of our GWAS loci were created by

selecting an equal number of random base pair locations across the genome and assigning a "locus

range" randomly selected from the actual loci. The primary KIG list covered a low percentage of

GWAS loci (<2%), but overlapped with ionomic loci more often than the random locations (80th

percentile in Arabidopsis, 95th in rice, >99th in maize).

Fig 1. Primary KIG list coverage of ionomic GWAS loci. Solid, vertical lines represent the percentage

of GWAS loci explained by a gene from the primary KIG list. Kernel Density Estimate (KDE) plots

display the distribution of 1000 random permutations with the 5th/95th percentiles shown as

dashed lines. Permutations were made with random genome selections of the same size and species

composition as the actual GWAS datasets (S3 Table).

Can orthology improve the KIG list?

While KIGs overlapped with ionomic loci more often than random genome locations, a large portion

of GWAS loci are left unexplained by the KIG list. We hypothesized that inferring orthologs across

species would increase the overlap between GWAS loci and expand the list of KIGs and orthologs.

Orthologs are genes across species inherited from a common ancestor. Although now in separate

species, orthologs tend to retain similar functionality [36], making them good candidate genes when

their function is known in one species. Indeed, for many KIGs, orthologs have been tested in multiple

species and found to encode conserved functions affecting element accumulation for S, Fe, Cd, Na,

K, Rb, Mo, Ni, Cu, P, Zn, As, B, Mn, and aluminum (Al) (Whitt et al., 2020). For example, alleles at

orthologous genes encoding the *MOT1* transporter affect variation in the level of Mo in multiple

species [26,33,37,38] and the same has been observed at *HKT1* orthologs for Na [27,39,40]. Inferring

the orthologs of KIGs across all of our study species should allow us to discover alleles in genes

affecting the ionome, including genes that have not yet been tested in every species in our list.

There are various ortholog inference algorithms that vary by their use of pairwise similarity scores,

phylogenetic analysis, and multiple sequence alignment - all of which produce variations in the

ortholog groups. The original KIG list [24] was made with InParanoid [41] because the ortholog

tables were readily available on Phytozome [42]. For this analysis, we switched to OrthoFinder v2.0

[43] because it performed better than other ortholog group inference methods in benchmarking

tests [43,44], and it can form ortholog groups between 3 or more species in one comparison. With

the ortholog tables from OrthoFinder, we produce "inferred" KIG lists by identifying orthologs of the

primary KIGs in the original species and any species with a sequenced genome, including the five we

are studying here. Expanding the list of taxa in the inferred KIG list created soybean and sorghum

gene lists so we could repeat the KIG/GWAS overlap comparison in all five of our ionomic GWAS

species.

The inferred KIG lists improved loci coverage across all five species and resulted in significant overlap

between the observed GWAS loci and the KIG ortholog lists (>95th percentile) for all species (Fig 2).

As expected, this is true whether we randomized the loci (Figs 1 and 2) or randomized the genes in

the list (Supplemental Fig 5). While the likelihood that we observed an enrichment of true positives

was significant (Fig 2), per-gene assessment of false discovery rates (FDR) was relatively high (Table

2), partly due to the large number of genes linked to GWAS peaks. Thus, the overlaps between KIG

and GWAS loci represent high-value candidates for alleles affecting ionomic variation at conserved

genes (Table 2; S6 Table).

Fig 2. Inferred KIG list coverage of ionomic GWAS loci. Solid, vertical lines represent the percentage

206   of GWAS loci explained by a gene from OrthoFinder inferred KIG lists. Kernel Density Estimate (KDE)

207   plots display the distribution of 1000 random permutations with the 5th/95th percentiles shown as

208   dashed lines. Permutations were made with random genome selections of the same size and species

209   composition as the actual GWAS datasets (Table 2).

Table 2. Summary of inferred KIG list overlaps with GWAS loci.									
				KIG/GWAS overlaps		KIG/random overlaps		False discovery rate (FDR)	
Species	GWAS loci	random locations	Inferred genes	genes	loci	genes	loci	genes	loci
Arabidopsis	1404	1404	163	45	40	32.86	30.55	0.73	0.76
soybean	7308	7308	378	264	223	170.03	166.23	0.64	0.75
rice	432	432	181	22	18	11.57	10.63	0.53	0.59
sorghum	1768	1768	201	23	21	13.36	12.13	0.58	0.58
maize	905	905	222	13	12	4.89	4.59	0.38	0.38

211
Among the genes that overlap between the inferred KIG orthologs and GWAS loci are conserved

genes known to regulate elemental accumulation across these plant species (Table 3; S4 Table).

Identification of alleles linked to orthologous candidate genes via GWAS for the same element

provides evidence that these genes not only encode conserved determinants of elemental

homeostasis but also indicate that phenotypically relevant, standing variation exists in these genes

that could be used to enhance plant mineral nutrition (e.g., P, Fe, Zn, Mn, Mo), improved mineral

nutrition for humans (e.g., Fe and Zn), or increased food safety (e.g., Cd).

Table 3. KIG ortholog groups overlapping GWAS loci for their cited traits. KIG citations are listed in S1 Table.			
KIG family	Cited traits	GWAS trait	Orthologs
AtZIF1, AtZIFL2 *OsZIFL12* *ZmYS3*	Cs Zn Fe	Fe	*AtZIFL2* *Glyma.06G313600, Glyma.06G313700* *Sobic.008G015700*
AtPHT1;1, AtPHT1;4, *AtPHT1;5* *OsPT2*	P As Se	P	*AtPHT1;4* *Glyma.10G186400, Glyma.10G186500* *LOC_Os08g45000*
AtPEN3	Cd	Cd	*AT3G16340* *Glyma.07G015800*

			LOC_Os01g52560, LOC_Os06g36090
AtNRAMP3	Fe Mn Zn	Mn	AtNRAMP3 Glyma.04G044000 Sobic.001G170000 Zm00001d030846
AtMTP11	Mn	Mn	AtMTP11 Glyma.18G286100 Sobic.003G349200 Zm00001d042939
AtMOT2	Mo	Mo	AtMOT2 Glyma.08G208700 LOC_Os01g45830 Sobic.003G237000 Zm00001d044068
AtMOT1 MtMOT1.2, MtMOT1.3 OsMOT1;1	Mo	Mo	AtMOT1 Glyma.06G074100, Glyma.17G203000, Glyma.17G203200, Glyma.17G203500 Zm00001d033053
AtBTS	Fe Mn Zn	Fe	Glyma.05G237500, Glyma.08G044700 Sobic.009G222200 Zm00001d038916
AtESB1	Ca, Mn, Zn, Na, S, K, As, Se, Mo	Mn	AT1G07730, AT2G39430 Glyma.08G258300, Glyma.12G067400, Glyma.18G282600 Sobic.003G064800
AtHMA2, AtHMA3, AtHMA4 OsHMA3	Zn Cd	Cd	AtHMA4, AtHMA2, AtHMA3 Glyma.09G055600 OsHMA3 Sobic.002G083000, Sobic.002G083100 Zm00001d005189, Zm00001d005190

Among the orthologs of KIGs linked to GWAS peaks for the same element are several transporters

with known roles in element uptake, sequestration, and remobilization. Notably, alleles were

discovered in orthologs of the *ZIFL1* and *ZIFL2* genes that affect the accumulation of Fe in seeds

(Table 3), a human nutrient that is limited in many diets [45–47]—making these alleles high-priority

targets in breeding materials for improved nutrient accumulation in food. KIGs *MOT1* and *MOT2*

encode molybdate transport proteins [26,48–51] and overlap Mo loci in multiple species. Alleles

affecting the accumulation of the toxic heavy metal Cd were discovered in all five species at the

heavy metal ATPases *AtHMA2*, *AtHMA3*, *AtHMA4*, and *OsHMA3*, which belong to a subset of *HMAs*

specific to Zn/Cd transport [11,52–54]. In addition to direct Cd transport, alleles at *PEN3/PDR8*,

implicated in Cd transport via chelation with the glutathione-derived phytochelatin [55], affected Cd

in three species. The alleles in these transporters represent additional breeding targets for improved

human food safety. The phosphate transporter (*PHT1*) family has been shown to affect phosphate

and As accumulation in Arabidopsis [56–58], Se uptake in rice [59] and was linked to P loci in

Arabidopsis, soybean, and rice. Thus, these alleles represent possible targets to enhance nutrient

uptake for the crop-limiting nutrient P. *MTP11* encodes a Mn transporter involved in Mn tolerance

and detoxification with the Golgi apparatus [60], and orthologs to *AtMTP11* are linked to Mn loci in

four species. Alleles in the orthologs of yet another transporter family, including *AtNRAMP3*, which is

involved in Mn homeostasis via the mesophyll vacuoles [61], were associated with Mn accumulation

in Arabidopsis, soybean, sorghum, and maize.

Alleles at regulatory genes from conserved ortholog groups were also identified among multiple

species. These included alleles at orthologs of the transcription factor BRUTUS (*BTS*), which affects

the expression of genes affecting Fe accumulation in Arabidopsis, including the transporter *IRT1* and

the regulatory transcription factor *FIT1* [29,62,63]. Alleles linked to the *BTS* orthologs affected Fe

accumulation in soybean, sorghum, and maize, demonstrating that this approach can identify

regulators of transporters and the proteins themselves. Thus, these alleles and genes represent high-

value breeding targets for improving seed Fe content in these three crops. The enhanced suberin 1

(*ESB1*) gene involved in Casparian strip biosynthesis [64] was previously known to affect Ca, Mn, Zn,

Na, S, K, As, Se, and Mo levels in Arabidopsis [65], but GWAS detected alleles at this gene affecting

Mn in Arabidopsis, soybean, and sorghum, suggesting that disruption of the Casparian strip has

similar impacts on the ionome across these species with dramatically different root systems.

The utility of combining GWAS across species using orthology and conservation for the known genes

was clear. The number of GWAS loci overlapping a KIG ortholog (Fig 2, Table 2) was greater than the

overlap with a primary KIG gene (Fig 1, S3 Table). However, this overlap only captured less than 5%

of the ionomic GWAS loci. Prior research bias has resulted in gene discovery focused on a handful of

elements due to researcher interests, leaving a majority of elements under-studied [24]. The overlap

of less than 5% may indicate that the KIG list is vastly incomplete [8], and more plant ionome genes

are waiting to be discovered. The permutation results in Figs 1 and 2 also demonstrate the naive

assumption of using GWAS results as independent experiments. Jointly considering the p-values

returned for orthologous genes across all five GWAS would increase false positives and result in

spurious conclusions. To overcome this and identify high-quality candidates for the remainder of the

GWAS results, we sought a formal procedure for assessing the likelihood of GWAS results at

orthologous loci across these five species regardless of previously demonstrated functions.

Orthology as a key factor for candidate selection

We hypothesized that conservation of orthologs within GWAS peaks across multiple species could be

a criterion to identify genes of previously unknown function. In the expanded KIG ortholog/GWAS

overlap comparison (Fig 2, Tables 2 and 3, S4 Table), we frequently detected phenotypically-

impactful alleles in multiple species GWAS. GWAS is "forward" genetics and will also detect alleles at

genes of previously unknown functions. Therefore, ionomic GWAS datasets will detect alleles at

unknown, conserved regulators of elemental accumulation. If alleles at previously unknown

conserved regulators of the ionome are present, comparisons between GWAS loci from multiple

species will occur at orthologous genes more than expected by chance.

To determine if this was possible and identify conserved ionomic genes, we developed a method we

call FiReMAGE: Filtering Results of Multi-species, Analogous, GWAS Experiments. This approach

identifies the orthologous genes linked to GWAS associations in multiple species and compares their

occurrence to a background distribution to assess the false discovery rate. Additionally, it increases

confidence in a candidate gene linked to a GWAS locus without relying on previously known

functions. FiReMAGE works by tabulating the genes underlying trait loci in each species, determining

their orthologous relationships via OrthoFinder, and retaining ortholog groups with candidate hits

from at least three species (Fig 3). The threshold of three or more species is to reduce the false

positives due to the likelihood of increased synteny between two closely related species. Like the

permutations used with the KIG/GWAS comparisons, FiReMAGE creates 1000 permutations of our

actual GWAS data by randomly selecting an equal number of genomic locations and ranges from

each species' genome. FiReMAGE runs the random permutations through the same orthology

comparison as the actual data and uses the permutation distributions to evaluate the likelihood that

loci overlap orthologs. This background distribution is used to evaluate the significance of the

overlap and produces a false discovery rate for candidate genes identified this way.

Fig 3. Overview of the FiReMAGE pipeline. (A) FiReMAGE orthology comparison identifies genes

linked to SNPs in each GWAS that are also orthologous to each other. (B) FiReMAGE starts by

identifying genes within range of GWAS loci intervals. These genes are recorded and filtered through

the orthology comparison to identify orthologous candidates. The original GWAS loci are used to

generate 1000 random permutations, which go through the same process as the actual data.

Distributions from the permutations are used to assess genes in the real dataset and return a

prioritized candidate list. Created with BioRender.com.

With FiReMAGE, we identified candidates in ortholog groups with hits in at least three of the five

species for over 30% of soybean and maize loci and over 50% in Arabidopsis, rice, and sorghum (Fig

4). The signal from the actual datasets was above the 95th percentile of the permutations in each

species, although the false discovery rate was still relatively high. Across the 19 elemental traits for

the 3 out of 5 species comparison, 3866 unique orthologs were linked to loci in Arabidopsis, 7852 in

soybean, 2630 in rice, 2982 in sorghum, and 872 in maize (S5 Fig; S7 Table).

Fig 4. FiReMAGE identifies candidates for a larger poriton of GWAS loci than the KIG list.

Percentage of ionomic GWAS intervals with a candidate from FiReMAGE's comparative approach

(left) and the OrthoFinder KIG list (right). The actual GWAS dataset is in green; 1000 random

permutations of the actual dataset are in blue, with error bars representing 5th/95th percentiles.

The size of candidate lists and false discovery rates vary between species and traits (Figs 5 and 6, S1

and S2 Figs), but overall, requiring more species to be present in an ortholog group returns a shorter

candidate list (Fig 5) with lower false discovery rates (Fig 6). While still encompassing a large number

of genes, the 3/5 list is a significant reduction in the number of potential candidates.

Fig 5. Orthologs from ortholog groups linked to Ca, Cd, Fe, Mn, or Mo loci in 3/5, 4/5, and 5/5

species. Actual dataset counts are in green, and 1000 permutations of the actual data are in blue,

with the point representing the mean and error bars representing the 5th/95th percentiles. Columns

are by species; rows are the number of species represented in the returned genes' ortholog groups.

Distribution plots for ionomic traits found only in 3/5 or 4/5 species ortholog groups are in S1 and S2

Figs.

Fig 6. False discovery rate (FDR) of FiReMAGE's selected candidate genes from ortholog groups

with representation from 3/5, 4/5, and all five species for each ionomic trait. FDR was calculated

by taking the average number of orthologous candidates returned in the 1000 permutation datasets

and dividing it by the number of orthologous candidates returned in the actual dataset for each

combination of species, ionomic trait, and species representation in the ortholog group.

Including maize and sorghum in our FiReMAGE comparison enabled the observation of how two

closely related species impacted the results. We expected the increased synteny between maize and

sorghum to increase the number of orthologous genes found, especially in the 3/5 ortholog groups

where both maize and sorghum are used (highlighted red in Fig 7). However, these ortholog groups

from species with increased synteny do not make up the majority of the results. The largest

highlighted intersection was between maize, sorghum, and soybean, with 245 ortholog groups.

However, the FDR of these genes is comparable to 3/5 ortholog groups with less syntenic species

compositions, suggesting that this is not a major concern. Including a perfectly matched permutation

set enabled the weighting of candidates (through calculation of FDR) across all comparisons.

Fig 7. UpSet plot of FiReMAGE's selected candidates' ortholog group composition. The intersection

matrix shows the unique compositions of all ortholog groups determined by OrthoFinder. Ortholog

groups are ordered by the degree (number of species) of the intersection and then by group size.

Intersection size represents the number of ortholog groups matching the species composition below.

Set size shows the total number of instances in which a species appears in all ortholog groups. The

top panel displays the individual genes' false discovery rate (FDR) distribution in their respective

ortholog groups for all ionomic traits. Note that some genes may be represented twice in the boxplot

distribution if they appear for multiple traits. Thus, they may have multiple FDRs due to the different

SNP distributions and FDR of the ionomic traits. Because of the theory that high levels of synteny

349   between sorghum and maize may increase the number of false positives returned, ortholog groups

350   with two of the three species being sorghum and maize are highlighted red.

Notable candidates

For each ionomic trait, FiReMAGE produced a list of orthologous candidates prioritized by likelihood:

the FDRs as estimated by permutation (S7 Table). If elemental accumulation is controlled by

conserved genes, then the FiReMAGE procedure should identify more orthologous gene/loci

overlaps than expected by chance via the permutations. FDRs differed by element accumulation trait

and species combinations. Generally, high-likelihood candidates were observed for elements with a

previously demonstrated high heritability [66–70]. These elements also had ortholog groups with

GWAS hits in all five species: Cd, Ca, Fe, Mn, and Mo (Fig 5). Of 23,117 FiReMAGE candidates, only 81

belong to these 5/5 ortholog groups. The value of orthology is highlighted by looking at these genes'

positions in ranked candidate likelihoods. When ranked by likelihood, the top 56 FiReMAGE GWAS

candidates are from the 81 candidates with GWAS hits in all five study species (S7 Table). When the

top 10% of candidate genes were considered (232 genes; S8 Table), candidates 1-56 and 115-116 are

linked to GWAS hits at orthologous candidates in all five species. The GWAS hits linked to these 5/5

candidate genes (n = 64 SNPs) were also enriched for the top 10% of the most significant SNPs from

their respective GWAS at a Chi-squared estimated p-value of 1.25 e-10 (obs: 21 True, 43 False; exp: 6

True, 58 False). By design, the signal in GWAS is determining portions of our top 10% of candidates.

Considered individually, candidates linked to the 21 GWAS hits that are the most significant SNPs

would be targets for candidate verification. Using orthology, we can filter through the GWAS SNPs

with "middling" signal to identify the remaining conserved candidates linked to the other 43 GWAS

hits.

371
The identities of the genes encoded by the FIReMAGE candidates in our 5/5 species lists implicate a

wide variety of biological processes in elemental homeostasis. The 81 candidates belong to nine

orthologous groups. For Fe, among the 5/5 candidates with no previously known role in Fe

homeostasis, is an ortholog group that encodes the Mevalonate kinases across these five species,

suggesting a currently unknown role for the mevalonic acid pathway in regulating Fe homeostasis in

flowering plants. Among the top Cd candidates, we find the *HMA* ortholog group mentioned above

in the KIG list (Table 3), RAB GTPase homologs of *RABA1i* that are involved in regulating cell wall

composition [71], and an ortholog group of Domains of Unknown Function 567 (DUF567) protein

family. In Arabidopsis, DUF567 is expressed in the roots and identified among a list of 129 genes

associated with metal hyperaccumulation in *Arabidopsis halleri* [72]. Thus, FireMAGE provides

additional experimental evidence, leveraged across five species, that this DUF567 family affects ion

homeostasis and may underlie ecological adaptation via hyperaccumulation in Arabidopsis relatives.

For Mo, the 5/5 candidates are the molybdate transporters, *MOT2*, listed in the KIG list for Mo

accumulation [48], and adenine nucleotide transporters (*NTT1* and *NTT2*). Nucleotide metabolism is

tied to Mo via molybdopterin, which is synthesized from nucleotide precursors [73] to form reaction

cofactors for four plant enzymes [74]. For Ca, the orthologous candidates are annotated as encoding

a Ring/Ubox superfamily with no known role in Ca accumulation. Lastly, for Mn, the candidate

ortholog groups are annotated as histone proteins and cellulose synthase A9 (*CESA9*). *CESA9* has

been shown to influence the cellulose content and cell wall biosynthesis in the epidermal seed coat

in Arabidopsis seeds [75]. In cellulose synthesis, Mn ions are used to catalyze glycosidic bond

formation, and in Arabidopsis, *AtCESA3* coordinates Mn ions with the glycosyltransferase substrate

uridine diphosphate glucose [76].

We also compared the FiReMAGE candidates to another candidate selection method performed in

maize: Camoco [19]. Out of over 47,000 candidate observations over multiple expression networks

and elemental trait combinations, we selected the top 200 observations determined by FDR rankings

from Camoco's multiple expression-network-based prioritizations and ionomics trait analysis. This

selection contained 122 unique candidate genes to compare against FiReMAGE's evolutionary-based

prioritizations. Of these, only two affected variations for the same trait in both methods, indicating

that single-species expression networks and orthology selection methods boost different genetic

signals to tie GWAS results to candidate genes. One gene, *Zm00001d04461*, identified through both

methods as affecting Mo, encodes an F-box domain containing protein; the other,

Zm00001d012274, affecting Fe accumulation, is an enzyme encoding an Omega-hydroxypalmitate O-

feruloyl transferase involved in the production of suberinized hydrophobic barriers [77].

Discussion

We have demonstrated that combining orthology and GWAS from multiple species detects

conserved regulators of phenotypic variation. Using the Known Ionomic Genes (KIG) as a set of

curated annotations validates our hypothesis that combining GWAS and orthology will correctly

identify variation affecting elemental accumulation. This resulted in the discovery of phenotypically-

impactful alleles affecting ionomic variation in genes of previously known functions encoded in these

five species. Additionally, the FiReMAGE approach goes beyond the KIG list by identifying numerous

candidates in genes without previously determined roles in element accumulation (S7 Table). This

ability to identify and prioritize candidate genes for conserved roles in trait variation using the

natural variation of multiple species transcends one of the key limitations of the post-genomic era:

ascertainment bias due to prior study. Instead of selecting candidates with prior annotations, these

orthologs are annotated as high-value candidate genes for further molecular function

characterization through the demonstrated orthology of genes linked to natural variation in multiple

species. The overlap of GWAS results with genes of previously known function and the identification

of orthologous variants using FiReMAGE produced a comprehensive public list of alleles at candidate

conserved regulators of 19 elemental accumulation for future molecular mechanistic studies or

marker-assisted selection (S2 and S7 Tables). Together, these two approaches can help identify

candidate genes for traits that are common between species.

Orthology-based FiReMAGE is highly effective for identifying gene candidates

To detect unknown ionomic candidates that do not meet the criteria of the KIG list [24], we

implemented the multi-species comparison we called FiReMAGE. Orthology clearly provides a good

estimate of function, as demonstrated by the improved GWAS/KIG overlap (Figs 1 and 2, Table 2, S3

Table). As a result of this success, we hypothesized that orthology, even in the absence of gene

function annotations, would also identify conserved genes as candidates without any prior

demonstration of ionomic function. FiReMAGE identified more orthologous genes underlying GWAS

for the same trait than expected by chance (Figs 4 and 5), demonstrating that this approach does

work. In addition to KIGs (Table 3), FiReMAGE also returned non-KIG ortholog groups that encoded

natural variation in elemental accumulation (S7 Table). These FiReMAGE candidates are likely to

encode conserved regulators of elemental accumulation, for which our analysis is the first genetic

demonstration of this role. Finding conserved critical regulators of ion homeostasis by GWAS

validates the expectation that conserved regulators (e.g., hub genes) will repeatedly accumulate

phenotypically effective variation. It also dramatically broadens the capacity of GWAS to work as a

gene function discovery tool in the forward-genetics mode. Thus, comparing element accumulation

association studies across organisms can help us discover additional element homeostasis

mechanisms. Our approach relies on both annotated genomes and GWAS data, but it should be

generalizable to any homologous traits across taxa. FiReMAGE does not require any information

other than a list of GWAS SNPs and gene positions in genomes, thus avoiding prior knowledge bias

and enabling accelerated gene function discovery for traits and species with little to no functional

annotations or prior work.

One of the non-KIG ortholog groups with alleles in all five species detected by FiReMAGE for Cd

encodes a domain of unknown function 567 protein (DUF567) family. This DUF567 is just one of 198

elemental accumulation candidate genes identified by FiReMAGE in Arabidopsis that has no

previously known function in any biological processes (S9 Table). The remaining unknown genes are

not in 5/5 ortholog groups, but when ranked by likelihood (permutation estimated FDRs), 19 are in

the top 10% of Arabidopsis candidates for all traits, and 46 unknown genes are in the top 25%. None

of these unknown genes would be considered candidates if relying only on gene annotations. The

likelihood rankings returned by FiReMAGE and the fact that known conserved regulators also

underlie significant GWAS peaks in multiple species (Table 3) suggest these are conserved,

previously unknown regulators of the ionome. Further single-gene mechanistic experiments may

result in the annotation of these genes as elemental regulators. Problems with annotating plant

genes and Gene Ontology (GO) terms in general, and for plants in particular, have been extensively

discussed [7–9]. FiReMAGE side-steps this problem, as is illustrated by the observation that it finds

602 orthologous gene candidates for Mo and 116 for Rb (S7 Table), yet there is no GO term explicitly

for either Mo or Rb.

Using the data from multiple experiments, FiReMAGE emphasizes loci that would be passed over in a

single experiment but have a strong signal when considered across taxa. Five traits, Ca, Fe, Mn, Mo,

and Cd (Fig 5, S7 Table), have ortholog groups with candidate gene hits in all five species.

Interestingly, these ortholog groups are substantially enriched for hits within the top 10% of most

significant associations (p = 1.25e-10). FiReMAGE uses the federation of data across multiple species

to boost signal below the p-value thresholds typically employed for single-species GWAS

assessment. This is illustrated by the results underlying GWAS hits at the ortholog groups containing

MOT2 and the Cd/Zn transporting *HMAs* (Table 3). The worst performing taxa for these ortholog

groups were soybean for Mo accumulation at *MOT2* (p-value = 0.0104) and Arabidopsis for Cd

accumulation at the tandem gene duplication comprising AtHMA2, AtHMA3, and AtHMA4 (p-value =

5.6e-05) - neither meeting strict stringency cutoffs in their respective GWAS assessments (S7 Table).

FiReMAGE approach and utility for gene discovery

Our FiReMAGE approach combines a wide array of information. It has many steps (functions) and

dozens to hundreds of user-defined parameters with no obvious correct choice. These include the p-

value thresholds for each GWAS, the linkage range for SNP-to-gene relationships, and the outgroups

for constructing ortholog relationships. In addition, several variables are inherent in the data and

experimentation that also affect FiReMAGE, including differences in measurement error in each

experiment, variation in the statistical power of each GWAS panel, the total genetic variation in a

species, and the association panel selection. By using permutations instead of an idealized null

distribution, we can estimate how likely the results are to happen by random chance and provide

estimates of the likelihood that our predictions are true. The FiReMAGE approach calculates these

permutations for every trait, species, and dataset combination. As researchers get better estimates

of parameters or add additional data, the permutations must also be updated. Preliminary testing

determined that random selection parameters were replicating real data structure and composition,

and our permutations serve as consistent internal controls for benchmarking our real results.

We expected that the high synteny between the three Poaceae species, rice, sorghum, and maize,

would inflate false positives. To the extent that multiple linked genes are shared across taxa due to

conservation of gene order, we expected the ortholog groups detected as 3/5 hits in FiReMAGE in

only the Poaceae to have high FDR. However, these ortholog groups did not have increased FDR

compared to other 3/5 ortholog groups (Fig 7). The combination of the recent genome duplications

and uneven loss of duplicate genes well demonstrated in maize, Arabidopsis, and others [78–83] and

the prevalence of chromosomal rearrangements and transposition of genes in plant genomes may

have shuffled these genomes enough to permit the detection of single conserved genes within our

selected window sizes. Although we did not see a huge influence of synteny in our comparison,

other species combinations may find controlling for synteny critical [84]. In all cases, the undue

influence of synteny will be reflected in the permutation FDR calculations, highlighting the value of

empirically estimated expectations over naive null hypotheses.

The success of FiReMAGE with the particular data sets and user-set parameters we chose here

demonstrates that the approach is viable. One feature of this analysis was the reliance on user-

reported p-values for a number of the data sets. This resulted in uneven p-value thresholds and

presumably false positive and negative rates across the input experiments. Because FiReMAGE

federates all the data together, the optimal p-value thresholds for each experiment are dependent

on the performance of the comparison. One interesting feature of FiReMAGE is that it is highly

sensitive to false-negative rates, a problem made larger as more species are brought together. Using

too stringent of a p-value cutoff for the GWAS results input into FiReMAGE can fail to include a true

association of modest effect near a conserved regulator. FiReMAGE cannot then include this gene in

an overlap or loses one true overlap as a result. We chose p-value thresholds considered permissive

by community standards (see methods) to reasonably limit false negative rates. As candidate genes

suggested by FiReMAGE are tested, the genes that are confirmed and genes that were false positives

will provide an additional mechanism to compare and refine parameters.

The high FDRs from the KIG comparison demonstrate why common practices filter SNPs with

stringent thresholds (Bonferroni, FDR, etc.) before referencing functional annotations of linked

genes. However, these recommendations do not address that even with generous genome coverage,

KIG candidates cannot explain a large portion of ionomic variation (Figs 1 and 2). While some GWAS

loci are due to spurious associations and some KIGs do not drive elemental phenotypes in all

populations in all species, we believe the majority of loci lack KIG candidates because the KIG list is

incomplete [8]. Candidate selection methods reliant on prior information have inherently narrow

scopes that increase false negatives and inhibit gene discovery. The KIG list is slightly better at

discovery, as it is valuable in transferring information between species with few and abundant

annotations. However, identifying novel ionomic candidates or functional annotations requires novel

approaches, such as FiReMAGE.

Limitations of FiReMAGE

One effect of the FiReMAGE approach on candidate gene assessment is that it rejects alleles and

false positives that occur in GWAS results from a single species or in a limited number of species.

While of great potential utility, this approach can potentially contribute to false negative

assessments in at least two ways. 1) The FiReMAGE approach will necessarily miss taxonomically-

restricted solutions and limitations to element homeostasis. For example, glucosinolates in

Arabidopsis, benzoxazinoids in maize, and nodulation in soybeans, all have impacts on the ionome

and are not evolutionarily conserved processes across these taxa. As a result, alleles affecting

variation in these processes, even if they have substantial impacts in one taxon, would not be

prioritized by FiReMAGE. The KIG list or more involved experiments like gene transcript expression

would be better selection methods in these cases. 2) The requirement to appear in multiple lists and

the stringency of the criteria for inclusion in the GWAS outputs contribute to false negatives intrinsic

in FiReMAGE. Researchers particularly worried about false positive assessments in their GWAS

outputs may not have considered the effect that the high rate of false negatives will have on

downstream meta-analysis and data federation approaches, like FiReMAGE. Because detecting

conserved candidates by FiReMAGE relies on true positives in multiple species, providing too-

stringent lists of loci will result in poor performance as more species are added to the comparison.

Another possible limitation is that FiReMAGE requires strict trait matches across taxa and we only

compared KIGs to loci of traits for which they had been cited. However, many elemental uptake

traits are related and often co-transported by the same transporters. The same gene could have

different elemental phenotypes in taxa due to many factors, including the interaction with species-

specific mechanisms (see above). The rules governing elemental interactions are poorly understood

[85,86]. Rather than attempting to relate ortholog groups to similar elements they have yet to be

tested for with our (likely) biased understanding of the ionome, we chose a strict comparison

between traits across species. We have provided a list of KIGs overlapping any trait loci (S4 Table) for

exploring their potential functions in related traits.

A functional requirement for FiReMAGE is the collection of the same trait across multiple diverse

taxa. Many traits, such as flower morphology and resistance to specific pathogens, are not

comparable across species. Additionally, collecting trait data from GWAS size populations across

multiple species can be time and resource-intensive. However, advances in phenotyping

technologies are enabling the creation of multi-species datasets, and more datasets are publicly

available in GWAS databases like AraGWAS [87] for Arabidopsis, RiceVarMap2 [88] for rice, and

GWAS Atlas [89] for studies in cotton, maize, soybean, rice, and sorghum. FiReMAGE is able to

identify candidates for any trait where the data is available in three or more species. We note that

the threshold of three or more species excludes eudicot-specific candidates in this five-species

comparison, and we may include more eudicot species as their ionomic GWAS experiments become

available.

Lastly, because FiReMAGE relies on orthology, the limitations surrounding current ortholog group

inference methods apply to the candidate lists produced by FiReMAGE. While OrthoFinder v2.0 is

the best algorithm for whole genome comparisons [43], there are likely some gene trees with

evolutionary histories that make accurate ortholog inference challenging [44]. We recommend

following best practices for operating OrthoFinder (see methods), including proper species sampling

and validating gene tree phylogenies for FiReMAGE candidates of interest when appropriate.

Future mutant screens

Unbiased candidate selection methods are necessary to transition into research using non-model

species and with genes of unknown function. Our approach has identified the conserved genes

underlying genetic variation via analogous GWAS in multiple species and their likelihood as causal

genes in ionomic phenotypic variation. In future studies, testing mutant knock-out libraries in

Arabidopsis [90], sorghum [91], and maize [92] will allow us to validate FiReMAGE predictions and

improve candidate selection parameters.

Our research demonstrates that the disparate approaches of the KIG list and FiReMAGE both

present valuable ionomic candidates, each with distinct advantages and weaknesses, and highlight

the importance of strategic decisions surrounding candidate gene identification for downstream

quantitative genetic analysis. Future experiments will further explore best practices and outcomes of

comparative GWAS methods, including parameters to balance false negatives and false positives.

We will also explore implementing a ranking equation that considers other variables besides the

number of species in ortholog groups. We hope this tool facilitates gene discovery in plant genetics.

Methods

Data availability

R scripts created for this version of the KIG list can be found at

https://github.com/danforthcenter/KIG_v2. FiReMAGE scripts, GWAS SNPs used in this paper,

ortholog tables, and gene coordinates can be found at

https://github.com/danforthcenter/FiReMAGE. Gene coordinates can alternatively be downloaded

from Phytozome (https://phytozome-next.jgi.doe.gov/phytomine/begin.do).

GWAS datasets

598   **Single nucleotide polymorphism (SNP) filtering**
599

The SNPs from each GWAS dataset were selected by p-value thresholds that were separately

determined for each species: *Arabidopsis thaliana* (Arabidopsis; $p < 0.001$), *Glycine max* (soybean; p

< 0.05), *Oryza sativa* (rice; $p < 1e-5$), *Sorghum bicolor* (sorghum; $p < 1e-4$), and *Zea mays* (maize; p $<$

0.01). The p-value thresholds used are intentionally permissive, especially in soybean, to limit

spurious SNPs but mostly to reduce false negatives in our GWAS/ortholog overlaps.

SNPs from each dataset are collapsed into loci using general linkage ranges for each species (Table

4). The p-value from the most significant SNP is retained as the collapsed loci's p-value. Collapsing

SNPs reduces computational load and avoids genes overlapping neighboring/redundant SNPs. The

pipeline also accepts files already in loci intervals, like the maize NAM dataset.

Table 4. Overview of GWAS datasets.				
Species Annotation	general linkage range	input as loci	SNPs	collapsed SNPs/loci
A. thaliana_TAIR10	75000	FALSE	7167	1404
G. max_Wm82.a2.v1	200000	FALSE	37854	7308
O. sativa_v7.0	243000	FALSE	17109	432
S. bicolor_v3.1.1	100000	FALSE	2139	1768
Z. mays_RefGen_V4	100000	TRUE	905	905

610
611   **Soybean GWAS and 2015-2017 field experiments**
612
Glycine max genotypes were obtained from the USDA-ARS GRIN collection (https://npgsweb.ars-

[grin.gov/gringlobal/](https://npgsweb.ars-grin.gov/gringlobal/)). Field experiments were performed over three years (2015, 2016, and 2017)

with three field replicates at each field experiment in a random complete block design. The 2015 and

2016 field experiments were performed at the University of Missouri Bradford Research farm

located near Columbia, Missouri, (38°53'38.4"N 92°11'47.1" W) which has a Mexico silt loam (fine,

smectitic, and mesic Vertic Epiaqualf) and the 2017 field experiment was performed at the Rollins

Bottom research farm located near Columbia, MO, ((38°55'37.5" N, 92°20'44.6" W) on a Haymond

silt loam soil (course-silty, mixed, superactive, mesic Dystric Fluventic Eutrudepts). Seeds were

planted at a rate of 8 seed/ft, over 6 ft with a 2 ft gap between plots in 2015 and 2016, and 3 seed/ft

over 8 ft with a 4 ft gap between plots in 2017. Row spacing was 30" between rows in all

experiments. Field conditions were typical of soybean production in the Midwest USA, with NPK

fertilizer applied at rates appropriate per soil analyses, and pre-emergent and post-emergent

herbicides were applied as recommended to control weeds for conventional soybeans. Field plots

were planted on 06/30/2015, 05/06/2016, and 06/06/2017. Plots were hand-harvested and

threshed using a small plot thresher by 10/29/2015, 09/23/2016, and 11/17/2017. Flower color for

each field plot was recorded and compared to known colors for each genotype. Similarly, seeds were

inspected after threshing and compared to known hila and seedcoat coloration, with non-matching

samples discarded. Approximately 25 seeds were lyophilized for each field plot and ground to a fine

powder using a Perten 3310 lab mill using a "type 0" grinding disc (Perten Instruments, Sweden).

Powdered samples from each grow-out were analyzed concurrently for concentrations of 20

elements using an inductively coupled plasma mass spectrometry (ICP-MS) pipeline previously

described [93]. Batch effects inherent in the ICP-MS process were confounded with grow-out in the

initial analysis. Thus, a subset of samples from each batch of the initial ICP-MS analysis were re-

analyzed for elemental concentrations, and a batch-specific correction factor was generated and

subtracted from the first run's values. Analytical outliers identified as > 10 median absolute

deviations away from the median were removed within grow out.

For analysis within each grow-out, phenotypic values for each genotype were taken as the median

value within that grow-out. To better meet normality assumptions both for the GWAS and for

aggregation of the grow-outs using a mixed model method, phenotypes were Box-Cox transformed

within the out and across grow-out and then standardized. Lambda values for the Box-Cox were

chosen using a 95% confidence interval. To aggregate data from different grow-outs, best linear

unbiased predictors (BLUPs) for each genotype were calculated from a random intercept mixed

model: $Phenotype \sim (1|Genotype) + (1|Grow\ out) + (1|Genotype:Grow\ out)$.

Association mapping was performed for each phenotype in the three grow-outs and the combined

grow-out BLUPs. Genotype sample sizes for each grow-out are 355, 363, and 364 for 2015, 2016, and

2017, respectively. To account for population structure, kinship matrix and principal components

were included in the mixed model as fixed effects. The kinship matrix was calculated using all 365

genotypes present in at least one grow-out following the realizedAB method in the synbreed R

package [94] as implemented in the R package ionomicsUtils (v1.0

https://github.com/gziegler/ionomicsUtils). Principal components included in the model were

selected for each phenotype if they were correlated significantly with that phenotype at the alpha =

.001 level.

The multi-locus mixed model (MLMM) used performs forward inclusion and backwards elimination

until all of the heritable variation is explained or a maximum number of cofactors has been reached,

which in this study was 40. Model selection was performed using the extended Bayesian information

criterion [95] and the multiple-Bonferroni criterion [96]. A third set of SNP's was also returned based

on inclusion in the final forward inclusion step (max cofactor). Effect sizes for SNPs returned were

calculated using the mlmm.gwas package (S10 Table) [97]. A resulting SNP was said to overlap with a

neighboring SNP if the r^{-2} linkage disequilibrium was > .2 as calculated using the LD function in the

genetics R package [98] and implemented in the R package ionomicsUtils (v1.0

https://github.com/gziegler/ionomicsUtils). Average linkage disequilibrium values for all pairs of

SNPs were calculated across the genome by first separating SNP pairs into 100 kB bins based on

distance and then calculating LD within these bins [99].

OrthoFinder orthologs

OrthoFinder v2.0 [43] was downloaded from https://github.com/davidemms/OrthoFinder and used

with default parameters and best practices outlined here:

https://davidemms.github.io/orthofinder_tutorials/orthofinder-best-practices.html. Protein reads of

primary transcripts were obtained from Phytozome (v13) [42]. To make the KIG list, we used files for

all species in the list except for wheat due to its low-quality genome annotation. *Liriodendron*

tulipifera was included as the most closely related outgroup to the eudicots and monocots in

Phytozome v13. The resulting ortholog table included one-to-one, one-to-many, and many-to-many

ortholog relationships.

Loci/gene overlaps

We obtained all the Phytozome (v13) [42] gene coordinates for each species via the BioMart tool

(v0.7) located on the Phytozome homepage (https://phytozome-next.jgi.doe.gov). Using the

coordinates, we search for genes overlapping trait loci that are also in our 5-species ortholog groups.

For each trait, FiReMAGE retains genes from ortholog groups with overlaps in at least 3/5 species

and produces a list of prioritized candidate genes. The number of overlapping orthologs is recorded

for each locus to calculate false discovery rates.

Random permutations

GWAS loci permutations were 1000 selections of random locations equal to the actual datasets'

species and trait composition. Each species' chromosome start and end coordinates were

downloaded from the European Nucleotide Archive (ENA) database (GCA_000001735,

GCA_000004515, GCA_001433935, GCA_000003195, GCA_000005005). Because random base pair

locations are unlikely to be in range for collapsing into "loci", we assigned range lengths randomly

chosen from the actual GWAS loci. If random locations are close together, their randomly assigned

ranges may overlap. FiReMAGE runs the random permutations through the same loci/gene overlap

pipeline as the actual data, calculates the permutation's distributions to evaluate the likelihood of a

locus overlapping any nearby ortholog, and produces a false discovery rate (FDR) to prioritize

candidate genes. FDR was calculated as the mean number of orthologs overlapping random

locations divided by the number of actual orthologs overlapping loci.

The primary KIG list permutations (S3 and S4 Figs, S6 Table) were formed by selecting random genes

to replicate the same species and element compositions of the actual primary KIG list and repeating

1000 times. The inferred KIG list permutations were formed by putting the primary genes

permutations through the same ortholog-finding process as the actual KIG list, with one deviation. In

the actual inferred lists, a primary KIG's paralogs (if any) and orthologs are sourced from its ortholog

group. In the permutations, we only take the random gene's orthologs from its ortholog group. After

finding all the orthologs for the random genes, we add a random selection of genes from

Arabidopsis, rice, and maize that equal the number of paralogs in their actual inferred KIG lists.

Specifically, it is the number of paralogs only in the list as inferred genes - not primary genes that are

also paralogs to each other. This deviation was made to avoid random gene selections in large,

single-species gene families, as this would no longer randomly replicate the structure of the KIG list.

Funding
{.707}

This study was supported by the National Science Foundation (www.nsf.gov) grant IOS-

2309932 (IB, BPD), the United States Department of Energy Office of Science (BER;

https://science.osti.gov/ber) grant DE-SC0023305 (BPD), the Biotechnology and the

Biological Sciences Research Council (https://www.ukri.org/councils/bbsrc/) grants

BB/J003336/1 (AHP, GJN, DES) and BB/L000113/1 (DES), and Donald Danforth Plant

Science Center (https://www.danforthcenter.org) funds (LW). The funders had no role in

study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

Acknowledgements

717 We thank Ryan Hartsock for contributions to the early development of our orthology comparison

718 methods.

References

1. Shakoor N, Ziegler G, Dilkes BP, Brenton Z, Boyles R, Connolly EL, et al. Integration of

Experiments across Diverse Environments Identifies the Genetic Determinants of Variation in

Sorghum bicolor Seed Element Composition. Plant Physiol. 2016;170: 1989–1998.

2. Ziegler G, Nelson R, Granada S, Krishnan HB, Gillman JD, Baxter I. Genomewide association

study of ionomic traits on diverse soybean populations from germplasm collections. Plant

Direct. 2018;2: e00033.

3. Ziegler G, Kear PJ, Wu D, Ziyomo C, Lipka AE, Gore M, et al. Elemental Accumulation in Kernels

of the Maize Nested Association Mapping Panel Reveals Signals of Gene by Environment

Interactions. bioRxiv. 2017. p. 164962. doi:10.1101/164962

4. Zhang M, Pinson SRM, Tarpley L, Huang X-Y, Lahner B, Yakubova E, et al. Mapping and

validation of quantitative trait loci associated with concentrations of 16 elements in unmilled

rice grain. Theor Appl Genet. 2014;127: 137–165.

5. Atwell S, Huang YS, Vilhjálmsson BJ, Willems G, Horton M, Li Y, et al. Genome-wide association

study of 107 phenotypes in Arabidopsis thaliana inbred lines. Nature. 2010;465: 627–631.

6. Cubillos FA, Coustham V, Loudet O. Lessons from eQTL mapping studies: non-coding regions

735 and their role behind natural phenotypic variation in plants. Curr Opin Plant Biol. 2012;15: 192–

736 198.

737 7. Stoeger T, Gerlach M, Morimoto RI, Nunes Amaral LA. Large-scale investigation of the reasons

738 why potentially important genes are ignored. PLoS Biol. 2018;16: e2006643.

739 8. Baxter I. We aren't good at picking candidate genes, and it's slowing us down. Curr Opin Plant

740 Biol. 2020;54: 57–60.

741 9. Wimalanathan K, Friedberg I, Andorf CM, Lawrence-Dill CJ. Maize GO Annotation—Methods,

742 Evaluation, and Review (maize-GAMER). Plant Direct. 2018;2: e00052.

743 10. Stoeger T, Nunes Amaral LA. COVID-19 research risks ignoring important host genes due to pre-

744 established research patterns. Elife. 2020;9. doi:10.7554/eLife.61981

745 11. Chao D-Y, Silva A, Baxter I, Huang YS, Nordborg M, Danku J, et al. Genome-wide association

746 studies identify heavy metal ATPase3 as the primary determinant of natural variation in leaf

747 cadmium in Arabidopsis thaliana. PLoS Genet. 2012;8: e1002923.

748 12. Chao D-Y, Chen Y, Chen J, Shi S, Chen Z, Wang C, et al. Genome-wide association mapping

749 identifies a new arsenate reductase enzyme critical for limiting arsenic accumulation in plants.

750 PLoS Biol. 2014;12: e1002009.

751 13. Gupta PK, Kulwal PL, Jaiswal V. Association mapping in plants in the post-GWAS genomics era.

752 Adv Genet. 2019;104: 75–154.

753 14. Spain SL, Barrett JC. Strategies for fine-mapping complex traits. Hum Mol Genet. 2015;24:

754 R111–9.

755 15. Koornneef M, Meinke D. The development of Arabidopsis as a model plant. Plant J. 2010;61:

756 909–921.

757   16.   Park ST, Kim J. Trends in Next-Generation Sequencing and a New Era for Whole Genome

758   Sequencing. Int Neurourol J. 2016;20: S76–83.

759   17.   Greene CS, Troyanskaya OG. PILGRM: an interactive data-driven discovery platform for expert

760   biologists. Nucleic Acids Res. 2011;39: W368–74.

761   18.   Xing EP, Curtis RE, Schoenherr G, Lee S, Yin J, Puniyani K, et al. GWAS in a box: statistical and

762   visual analytics of structured associations via GenAMap. PLoS One. 2014;9: e97524.

763   19.   Schaefer RJ, Michno J-M, Jeffers J, Hoekenga O, Dilkes B, Baxter I, et al. Integrating

764   Coexpression Networks with GWAS to Prioritize Causal Genes in Maize. Plant Cell. 2018;30:

765   2922–2942.

766   20.   Salt DE, Baxter I, Lahner B. Ionomics and the study of the plant ionome. Annu Rev Plant Biol.

767   2008;59: 709–733.

768   21.   Chen A, Hu J, Sun S, Xu G. Conservation and divergence of both phosphate- and mycorrhiza-

769   regulated physiological responses and expression patterns of phosphate transporters in

770   solanaceous species. New Phytol. 2007;173: 817–831.

771   22.   Chen A, Gu M, Sun S, Zhu L, Hong S, Xu G. Identification of two conserved cis-acting elements,

772   MYCS and P1BS, involved in the regulation of mycorrhiza-activated phosphate transporters in

773   eudicot species. New Phytol. 2011;189: 1157–1169.

774   23.   Ram H, Sardar S, Gandass N. Vacuolar Iron Transporter (Like) proteins: Regulators of cellular

775   iron accumulation in plants. Physiol Plant. 2021;171: 823–832.

776   24.   Whitt L, Ricachenevsky FK, Ziegler GZ, Clemens S, Walker E, Maathuis FJM, et al. A curated list

777   of genes that affect the plant ionome. Plant Direct. 2020;4: e00272.

25.   Véry A-A, Nieves-Cordones M, Daly M, Khan I, Fizames C, Sentenac H. Molecular biology of K+

transport across the plant cell membrane: what do we learn from comparison between plant

species? J Plant Physiol. 2014;171: 748–769.

26. Baxter I, Muthukumar B, Park HC, Buchner P, Lahner B, Danku J, et al. Variation in molybdenum

content across broadly distributed populations of Arabidopsis thaliana is controlled by a

mitochondrial molybdenum transporter (MOT1). PLoS Genet. 2008;4: e1000004.

27. Baxter I, Brazelton JN, Yu D, Huang YS, Lahner B, Yakubova E, et al. A coastal cline in sodium

accumulation in Arabidopsis thaliana is driven by natural variation of the sodium transporter

AtHKT1;1. PLoS Genet. 2010;6: e1001193.

28. Rubio F, Nieves-Cordones M, Alemán F, Martínez V. Relative contribution of AtHAK5 and

AtAKT1 to K+ uptake in the high-affinity range of concentrations. Physiol Plant. 2008;134: 598–

608.

29. Eide D, Broderius M, Fett J, Guerinot ML. A novel iron-regulated metal transporter from plants

identified by functional expression in yeast. Proc Natl Acad Sci U S A. 1996;93: 5624–5628.

30. Huang X-Y, Deng F, Yamaji N, Pinson SRM, Fujii-Kashino M, Danku J, et al. A heavy metal P-type

ATPase OsHMA4 prevents copper accumulation in rice grain. Nat Commun. 2016;7: 12138.

31. Campos ACA, van Dijk WFA, Ramakrishna P, Giles T, Korte P, Douglas A, et al. Full species-wide

leaf and seed ionomic diversity of Arabidopsis thaliana. bioRxiv. 2020. p. 2020.11.09.373282.

doi:10.1101/2020.11.09.373282

32. Norton GJ, Travis AJ, Douglas A, Fairley S, Alves EDP, Ruang-Areerate P, et al. Genome Wide

Association Mapping of Grain and Straw Biomass Traits in the Rice Bengal and Assam Aus Panel

(BAAP) Grown Under Alternate Wetting and Drying and Permanently Flooded Irrigation. Front

Plant Sci. 2018;9: 1223.

33. Norton GJ, Douglas A, Lahner B, Yakubova E, Guerinot ML, Pinson SR, et al. Genome wide

association mapping of grain arsenic, copper, molybdenum and zinc in rice (Oryza sativa L.)

grown at four international field sites. PLoS One. 2014;9: e89685.

34. Norton GJ, Travis A, Ruang-Areerate P, Nicol GW, Adeosun AA, Hossain M, et al. Genetic loci

regulating cadmium content in rice grains. Euphytica. 2021;217: 35.

35. Talukdar P, Travis AJ, Hossain M, Islam MR, Norton GJ, Price AH. Identification of genomic loci

regulating grain iron content in aus rice under two irrigation management systems. Food

Energy Secur. 2022;11: e329.

36. Altenhoff AM, Studer RA, Robinson-Rechavi M, Dessimoz C. Resolving the ortholog conjecture:

orthologs tend to be weakly, but significantly, more similar in function than paralogs. PLoS

Comput Biol. 2012;8: e1002514.

37. Asaro A, Dilkes B, Baxter I. Multivariate analysis reveals environmental and genetic

determinants of element covariation in the maize grain ionome. bioRxiv. 2017. p. 241380.

doi:10.1101/241380

38. Ziegler G, Kear P, Wu D, Ziyomo C, Lipka A, Gore M. Elemental Accumulation in Kernels of the

Maize Nested Association Mapping Panel Reveals Signals of Gene by Environment Interactions.

BioRxiv. 2017. Available: https://www.biorxiv.org/content/10.1101/164962v1.abstract

39. Campbell MT, Bandillo N, Al Shiblawi FRA, Sharma S, Liu K, Du Q, et al. Allelic variants of

OsHKT1;1 underlie the divergence between indica and japonica subspecies of rice (Oryza

sativa) for root sodium content. PLoS Genet. 2017;13: e1006823.

40. Rus A, Baxter I, Muthukumar B, Gustin J, Lahner B, Yakubova E, et al. Natural variants of AtHKT1

enhance Na+ accumulation in two wild populations of Arabidopsis. PLoS Genet. 2006;2: e210.

41. Remm M, Storm CE, Sonnhammer EL. Automatic clustering of orthologs and in-paralogs from

pairwise species comparisons. J Mol Biol. 2001;314: 1041–1052.

42. Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, et al. Phytozome: a comparative

platform for green plant genomics. Nucleic Acids Res. 2012;40: D1178–86.

43. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics.

Genome Biol. 2019;20: 238.

44. Emms DM, Kelly S. Benchmarking Orthogroup Inference Accuracy: Revisiting Orthobench.

Genome Biol Evol. 2020;12: 2258–2266.

45. Haydon MJ, Kawachi M, Wirtz M, Hillmer S, Hell R, Krämer U. Vacuolar nicotianamine has

critical and distinct roles under iron deficiency and for zinc sequestration in Arabidopsis. Plant

Cell. 2012;24: 724–737.

46. Remy E, Cabrito TR, Batista RA, Teixeira MC, Sá-Correia I, Duque P. The major facilitator

superfamily transporter ZIFL2 modulates cesium and potassium homeostasis in Arabidopsis.

Plant Cell Physiol. 2015;56: 148–162.

47. Che J, Yokosho K, Yamaji N, Ma JF. A Vacuolar Phytosiderophore Transporter Alters Iron and

Zinc Accumulation in Polished Rice Grains. Plant Physiol. 2019;181: 276–288.

48. Gasber A, Klaumann S, Trentmann O, Trampczynska A, Clemens S, Schneider S, et al.

Identification of an Arabidopsis solute carrier critical for intracellular transport and inter-organ

allocation of molybdate. Plant Biol . 2011;13: 710–718.

49. Gil-Díez P, Tejada-Jiménez M, León-Mediavilla J, Wen J, Mysore KS, Imperial J, et al. MtMOT1.2

is responsible for molybdate supply to Medicago truncatula nodules. Plant Cell Environ. 2018.

doi:10.1111/pce.13388

50. Tejada-Jiménez M, Gil-Díez P, León-Mediavilla J, Wen J, Mysore KS, Imperial J, et al. Medicago

truncatula Molybdate Transporter type 1 (MtMOT1.3) is a plasma membrane molybdenum

transporter required for nitrogenase activity in root nodules under molybdenum deficiency.

New Phytol. 2017;216: 1223–1235.

51. Huang X-Y, Liu H, Zhu Y-F, Pinson SRM, Lin H-X, Guerinot ML, et al. Natural variation in a

molybdate transporter controls grain molybdenum concentration in rice. New Phytol.

2019;221: 1983–1997.

52. Hussain D, Haydon MJ, Wang Y, Wong E, Sherson SM, Young J, et al. P-type ATPase heavy metal

transporters with roles in essential zinc homeostasis in Arabidopsis. Plant Cell. 2004;16: 1327–

1339.

53. Olsen LI, Hansen TH, Larue C, Østerberg JT, Hoffmann RD, Liesche J, et al. Mother-plant-

mediated pumping of zinc into the developing seed. Nat Plants. 2016;2: 16036.

54. Tanaka N, Nishida S, Kamiya T, Fujiwara T. Large-scale profiling of brown rice ionome in an

ethyl methanesulphonate-mutagenized hitomebore population and identification of high- and

low-cadmium lines. Plant Soil. 2016;407: 109–117.

55. Kim D-Y, Bovet L, Maeshima M, Martinoia E, Lee Y. The ABC transporter AtPDR8 is a cadmium

extrusion pump conferring heavy metal resistance. Plant J. 2007;50: 207–218.

56. Shin H, Shin H-S, Dewbre GR, Harrison MJ. Phosphate transport in Arabidopsis: Pht1;1 and

Pht1;4 play a major role in phosphate acquisition from both low- and high-phosphate

environments. Plant J. 2004;39: 629–642.

57. Nagarajan VK, Jain A, Poling MD, Lewis AJ, Raghothama KG, Smith AP. Arabidopsis Pht1;5

mobilizes phosphate between source and sink organs and influences the interaction between

phosphate homeostasis and ethylene signaling. Plant Physiol. 2011;156: 1149–1163.

58.   Catarecha P, Segura MD, Franco-Zorrilla JM, García-Ponce B, Lanza M, Solano R, et al. A mutant

of the Arabidopsis phosphate transporter PHT1;1 displays enhanced arsenic accumulation.

Plant Cell. 2007;19: 1123–1133.

59.   Zhang L, Hu B, Li W, Che R, Deng K, Li H, et al. OsPT2, a phosphate transporter, is involved in

the active uptake of selenite in rice. New Phytol. 2014;201: 1183–1191.

60.   Peiter E, Montanini B, Gobert A, Pedas P, Husted S, Maathuis FJM, et al. A secretory pathway-

localized cation diffusion facilitator confers plant manganese tolerance. Proc Natl Acad Sci U S

A. 2007;104: 8532–8537.

61.   Lanquar V, Ramos MS, Lelièvre F, Barbier-Brygoo H, Krieger-Liszkay A, Krämer U, et al. Export of

vacuolar manganese by AtNRAMP3 and AtNRAMP4 is required for optimal photosynthesis and

growth under manganese deficiency. Plant Physiol. 2010;152: 1986–1999.

62.   Hindt MN, Akmakjian GZ, Pivarski KL, Punshon T, Baxter I, Salt DE, et al. BRUTUS and its

paralogs, BTS LIKE1 and BTS LIKE2, encode important negative regulators of the iron deficiency

response in Arabidopsis thaliana. Metallomics. 2017;9: 876–890.

63.   Yuan Y, Wu H, Wang N, Li J, Zhao W, Du J, et al. FIT interacts with AtbHLH38 and AtbHLH39 in

regulating iron uptake gene expression for iron homeostasis in Arabidopsis. Cell Res. 2008;18:

385–397.

64.   Hosmani PS, Kamiya T, Danku J, Naseer S, Geldner N, Guerinot ML, et al. Dirigent domain-

containing protein is part of the machinery required for formation of the lignin-based Casparian

strip in the root. Proc Natl Acad Sci U S A. 2013;110: 14498–14503.

65.   Baxter I, Hosmani PS, Rus A, Lahner B, Borevitz JO, Muthukumar B, et al. Root suberin forms an

extracellular barrier that affects water relations and mineral nutrition in Arabidopsis. PLoS

Genet. 2009;5: e1000492.

66.   Asaro A, Ziegler G, Ziyomo C, Hoekenga OA, Dilkes BP, Baxter I. The Interaction of Genotype

and Environment Determines Variation in the Maize Kernel Ionome. G3 . 2016;6: 4175–4183.

67.   Baxter IR, Ziegler G, Lahner B, Mickelbart MV, Foley R, Danku J, et al. Single-kernel ionomic

profiles are highly heritable indicators of genetic and environmental influences on elemental

accumulation in maize grain (Zea mays). PLoS One. 2014;9: e87628.

68.   Campos ACAL, van Dijk WFA, Ramakrishna P, Giles T, Korte P, Douglas A, et al. 1,135 ionomes

reveal the global pattern of leaf and seed mineral nutrient and trace element diversity in

Arabidopsis thaliana. Plant J. 2021;106: 536–554.

69.   Taliercio E, Scaboo A, Baxter I, Locke AM. The Ionome of a Genetically Diverse Set of Wild

Soybean Accessions. Crop Sci. 2019. doi:10.2135/cropsci2019.02.0079

70.   Norton GJ, Deacon CM, Xiong L, Huang S, Meharg AA, Price AH. Genetic mapping of the rice

ionome in leaves and grain: identification of QTLs for 17 elements including arsenic, cadmium,

iron and selenium. Plant Soil. 2010;329: 139–153.

71.   Lunn D, Gaddipati SR, Tucker GA, Lycett GW. Null mutants of individual RABA genes impact the

proportion of different cell wall components in stem tissue of Arabidopsis thaliana. PLoS One.

2013;8: e75724.

72.   Filatov V, Dowdle J, Smirnoff N, Ford-Lloyd B, Newbury HJ, Macnair MR. Comparison of gene

expression in segregating families identifies genes and genomic regions involved in a novel

adaptation, zinc hyperaccumulation. Mol Ecol. 2006;15: 3045–3059.

73.   Hover BM, Loksztejn A, Ribeiro AA, Yokoyama K. Identification of a cyclic nucleotide as a cryptic

intermediate in molybdenum cofactor biosynthesis. J Am Chem Soc. 2013;135: 7019–7032.

74.   Schwarz G, Mendel RR. Molybdenum cofactor biosynthesis and molybdenum enzymes. Annu

Rev Plant Biol. 2006;57: 623–647.

75. Stork J, Harris D, Griffiths J, Williams B, Beisson F, Li-Beisson Y, et al. CELLULOSE SYNTHASE9

serves a nonredundant role in secondary cell wall synthesis in Arabidopsis epidermal testa cells.

Plant Physiol. 2010;153: 580–589.

76. Qiao Z, Lampugnani ER, Yan X-F, Khan GA, Saw WG, Hannah P, et al. Structure of *Arabidopsis*

CESA3 catalytic domain with its substrate UDP-glucose provides insight into the mechanism of

cellulose synthesis. Proc Natl Acad Sci U S A. 2021;118. doi:10.1073/pnas.2024015118

77. Mertz RA, Ellsworth P, Ellsworth P, Lori Tausta S, von Caemmerer S, Howard Berg R, et al. The

maize Aliphatic Suberin Feruloyl Transferase genes affect leaf water movement but are

dispensable for bundle sheath CO2 concentration. bioRxiv. 2020. p. 2020.09.30.321265.

doi:10.1101/2020.09.30.321265

78. Birchler JA, Yang H. The multiple fates of gene duplications: Deletion, hypofunctionalization,

subfunctionalization, neofunctionalization, dosage balance constraints, and neutral variation.

Plant Cell. 2022;34: 2466–2474.

79. Gout J-F, Lynch M. Maintenance and Loss of Duplicated Genes by Dosage Subfunctionalization.

Mol Biol Evol. 2015;32: 2141–2148.

80. Schnable JC, Springer NM, Freeling M. Differentiation of the maize subgenomes by genome

dominance and both ancient and ongoing gene loss. Proc Natl Acad Sci U S A. 2011;108: 4069–

4074.

81. Edger PP, Pires JC. Gene and genome duplications: the impact of dosage-sensitivity on the fate

of nuclear genes. Chromosome Res. 2009;17: 699–717.

82. Panchy N, Lehti-Shiu M, Shiu S-H. Evolution of Gene Duplication in Plants. Plant Physiol.

2016;171: 2294–2316.

83. Thomas BC, Pedersen B, Freeling M. Following tetraploidy in an Arabidopsis ancestor, genes

were removed preferentially from one homeolog leaving clusters enriched in dose-sensitive

genes. Genome Res. 2006;16: 934–946.

84. Murat F, Van de Peer Y, Salse J. Decoding plant and animal genome plasticity from differential

paleo-evolutionary patterns and processes. Genome Biol Evol. 2012;4: 917–928.

85. Baxter I. Should we treat the ionome as a combination of individual elements, or should we be

deriving novel combined traits? J Exp Bot. 2015;66: 2127–2131.

86. Baxter I. Ionomics: studying the social network of mineral nutrients. Curr Opin Plant Biol.

2009;12: 381–386.

87. Togninalli M, Seren Ü, Meng D, Fitz J, Nordborg M, Weigel D, et al. The AraGWAS Catalog: a

curated and standardized Arabidopsis thaliana GWAS catalog. Nucleic Acids Res. 2018;46:

D1150–D1156.

88. Zhao H, Yao W, Ouyang Y, Yang W, Wang G, Lian X, et al. RiceVarMap: a comprehensive

database of rice genomic variations. Nucleic Acids Res. 2015;43: D1018–22.

89. Tian D, Wang P, Tang B, Teng X, Li C, Liu X, et al. GWAS Atlas: a curated resource of genome-

wide variant-trait associations in plants and animals. Nucleic Acids Res. 2020;48: D927–D932.

90. Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, et al. Genome-wide insertional

mutagenesis of Arabidopsis thaliana. Science. 2003;301: 653–657.

91. Addo-Quaye C, Tuinstra M, Carraro N, Weil C, Dilkes BP. Whole-Genome Sequence Accuracy Is

Improved by Replication in a Population of Mutagenized Sorghum. G3 . 2018;8: 1079–1094.

92. McCarty DR, Settles AM, Suzuki M, Tan BC, Latshaw S, Porch T, et al. Steady-state transposon

mutagenesis in inbred maize. Plant J. 2005;44: 52–61.

93. Ziegler G, Terauchi A, Becker A, Armstrong P, Hudson K, Baxter I. Ionomic screening of field-

grown soybean identifies mutants with altered seed elemental composition. Plant Genome.

2013;6: lantgenome2012.07.0012.

94. Wimmer V, Albrecht T, Auinger H-J, Schön C-C. synbreed: a framework for the analysis of

genomic prediction data using R. Bioinformatics. 2012;28: 2086–2087.

95. Chen J, Chen Z. Extended Bayesian information criteria for model selection with large model

spaces. Biometrika. 2008;95: 759–771.

96. Segura V, Vilhjálmsson BJ, Platt A, Korte A, Seren Ü, Long Q, et al. An efficient multi-locus

mixed-model approach for genome-wide association studies in structured populations. Nat

Genet. 2012;44: 825–830.

97. mlmm.gwas: Pipeline for GWAS Using MLMM. In: Comprehensive R Archive Network (CRAN)

[Internet]. [cited 23 Mar 2023]. Available: https://cran.r-

project.org/web/packages/mlmm.gwas/index.html

98. Warnes G, Warnes MG. The genetics package. Locus. 2007;26: 1.

99. Barria A, López ME, Yoshida G, Carvalheiro R, Lhorente JP, Yáñez JM. Population Genomic

Structure and Genome-Wide Linkage Disequilibrium in Farmed Atlantic Salmon (Salmo salar L.)

Using Dense SNP Genotypes. Front Genet. 2018;9: 649.

100. Krishnakumar V, Hanlon MR, Contrino S, Ferlanti ES, Karamycheva S, Kim M, et al. Araport: the

Arabidopsis information portal. Nucleic Acids Res. 2015;43: D1003–9.

Supplemental Files

S1 Fig. Orthologs in groups with 3/5 species representation. Error bars represent the 5th/95th percentiles of the permutations.

S2 Fig. Orthologs in groups with 4/5 species representation. Error bars represent the 5th/95th percentiles of the permutations.

S3 Fig. Distribution of list sizes in the inferred KIG permutations depending on ortholog tables. The old distribution of permutations' list sizes was from using OrthoFinder to call ortholog groups only between the species in the inferred KIG. The new distribution of permutations' list sizes shows the list size when *Liriodendron tulipifera* is added to our OrthoFinder run as an outgroup to our target species. The green dot represents the number of inferred KIGs in the actual list.

S4 Fig. Inferred KIG and its permutations compared to GWAS loci. Solid, vertical lines represent the percentage of GWAS loci explained by a gene from OrthoFinder inferred KIG lists. Kernel Density Estimate (KDE) plots display the distribution of 1000 random permutations with the 5th/95th percentiles shown as dashed lines. Inferred KIG list permutations were made from the orthologs of a list of random genes equal to the species and elemental composition of the primary KIG list (see Random permutations methods section).

S5 Fig. Distribution of the number of orthologous candidates returned per loci.

S1 Table. The current primary KIG list. Genes with functional characterizations from primary literature sources that are linked to changes in the ionome.

1002   **S2 Table. Updated KIG list with OrthoFinder inferred orthologs.** *T. aestivum* (wheat) is not included

1003   in this version of the inferred KIG list.

1004

1005   **S3 Table. False discovery rates of the genes and loci overlaps in Fig 1.** False discovery rates are

1006   calculated by taking the average number of genes/loci in random permutation overlaps and dividing

1007   by the number of genes/loci in the actual GWAS datasets.

1008

1009   **S4 Table. KIGs overlapping GWAS loci.** KIGs compared to all elemental uptake trait loci regardless of

1010   KIG list citations.

1011

1012   **S5 Table. KIGs in FiReMAGE's candidate lists.**

1013

1014   **S6 Table. False discovery rate table for gene and loci overlaps in S4 Fig.** False discovery rates are

1015   calculated by taking the average number of genes/loci in random permutation overlaps and dividing

1016   by the number of genes/loci in the actual GWAS datasets.

1017

1018   **S7 Table. The complete FiReMAGE priority lists for each trait and species.**

1019

1020   **S8 Table. Top 10% of all FiReMAGE candidates.** FiReMAGE candidates were sorted by their

1021   permutation calculated false discovery rates, and taking the top 10% with the lowest FDR.

1022

1023   **S9 Table. FiReMAGE candidates in Arabidopsis with no previously known biological processes**

1024   **function.** We used ThaleMine [100] to analyze Arabidopsis candidates for each trait to identify genes

1025   with unknown function.

1026

1027   **S10 Table. Soybean SNPs associated with ionomic traits.** SNP associations generated with methods

1028   outlined in "Soybean GWAS and 2015-2017 field experiments".