FMNL2 regulates actin for ER and mitochondria distribution in oocyte meiosis

Meng-Hao Pan[1, 3], Zhen-Nan Pan[1], Ming-Hong Sun[1], Xiao-Han Li[1], Jia-Qian Ju[1], Shi-Ming Luo[2], Xiang-Hong Ou[2], Shao-Chen Sun[1*]

[1]College of Animal Science and Technology, Nanjing Agricultural University, Nanjing 210095, China.

[2]Fertility Preservation Lab, Reproductive Medicine Center, Guangdong Second Provincial General Hospital, Guangzhou, China.

[3]College of Veterinary Medicine, Northwest A&F University, Yangling, Shaanxi, China.

Running Title: FMNL2 in mouse oocytes

[*]**Correspondence to:** Shao-Chen Sun, College of Animal Science and Technology, Nanjing Agricultural University, Nanjing 210095, China. E-mail: sunsc@njau.edu.cn

Funding: This work was supported by the National Natural Science Foundation of China (32170857); the National Key Research and Development Program of China (2021YFC2700100).

There is no conflict of interest to declare.

Abbreviations: FMNL2, Formin-like 2; FMNLs, Formin-likes; GV, germinal vesicle; GVBD, germinal vesicle breakdown; ATI, anaphase-telophase I; MI, metaphase I; MII, metaphase II; ER, Endoplasmic reticulum.

Abstract

During mammalian oocyte meiosis, spindle migration and asymmetric cytokinesis

are unique steps for the successful polar body extrusion. The asymmetry defects of

oocytes will lead to the failure of fertilization and embryo implantation. In present

study we reported that an actin nucleating factor formin-like 2 (FMNL2) played

critical roles in the regulation of spindle migration and organelle distribution. Our

results showed that FMNL2 mainly localized at the oocyte cortex and periphery of

spindle. Depletion of FMNL2 led to the failure of polar body extrusion and large

polar bodies in oocytes. Live-cell imaging revealed that the spindle failed to migrate

to the oocyte cortex, which caused polar body formation defects, and this might be

due to the decreased polymerization of cytoplasmic actin by FMNL2 depletion.

Furthermore, mass spectrometry analysis indicated that FMNL2 was associated with

mitochondria and endoplasmic reticulum-related proteins, and FMNL2 depletion

disrupted the function and distribution of mitochondria and endoplasmic reticulum,

showing with decreased mitochondrial membrane potential and the occurrence of

endoplasmic reticulum stress. Microinjecting Fmnl2-EGFP mRNA into FMNL2-

depleted oocytes significantly rescued these defects. Thus, our results indicate that

FMNL2 is essential for the actin assembly, which further involves into meiotic

spindle migration and ER/mitochondria functions in mouse oocytes.

Keywords: FMNL2; oocyte; actin; Endoplasmic reticulum; Mitochondria

Introduction

Mammalian oocyte maturation is an asymmetric division process that generates a

large egg and a small polar body. This asymmetry is critical for the following

fertilization and early embryo development. After germinal vesicle breakdown

(GVBD), the meiotic spindle is organized at the center of the oocyte, and then it

migrates to the oocyte cortex at the late metaphase I (MI). The oocytes are arrested at

metaphase II (MII) after the extrusion of first polar body (*1, 2*). Actin filaments, as the

most widely distributed cytoskeleton in cells, regulate various dynamic events during

oocyte meiotic maturation (*3*), and two key events are the spindle migration and

cortical reorganization in mammalian oocytes (*1, 4, 5*). Small GTPases and actin

nucleation factors are shown to promote the assembly and function of actin. The actin

nucleation factors are the molecules that directly promote the actin assembly: Arp2/3

complex control the assembly of branched actin, and formin family member Formin2

(FMN2) and Spire1/2 control the assembly of linear actin. These proteins are all

proposed to play a role in actin-related spindle migration and cytokinesis during

mammalian oocyte maturation (*6-8*). The cortex protein Arp2/3 complex nucleates the

actin to produce a hydrodynamic force to move the spindle toward the cortex, and

regulates cytokinesis during oocyte maturation (*1, 8*). FMN2 and Spire1/2 nucleates

actin around the spindle in the cytoplasm to give the meiotic spindle an initial power

for migration (*7, 9*).

Besides Formin2, the DRFs (diaphanous-related formins) subfamily in the

formin family has been extensively studied. The DRFs family consists of mDia, Daam,

FHOD and FMNLs (*10*). The "Formin-like" proteins (FMNLs) subfamily includes

FMNL1 (FRL1), FMNL2 (FRL3), and FMNL3 (FRL2). Like other Formin family

proteins, FMNLs play important roles in cell migration, cell division, and cell polarity

(*10, 11*). While FMNL2 is widely expressed in multiple human tissues, especially in

the gastrointestinal and mammary epithelia, lymphatic tissues, placenta, and

reproductive tract (*12*). As an important actin assembly factor, FMNL2 accelerates the

elongation of actin filaments branched by Arp2/3 complex (*13*). In invasive cells,

FMNL2 is mainly localized in the leading edge of the cell, lamellipodia and filopodia

tips, to improve cell migration ability by actin-based manner (*13-15*). FMNL2 is also

involved in the maintenance of epithelial-mesenchymal transition (EMT) in human

colorectal carcinoma cell (*16*). Besides its roles on the actin assembly, emerging

evidences indicate that FMNL2 may interact with organelle dynamics. It is shown that

FMNL2 is related with the Golgi apparatus, since the absence of FMNL2/3 can cause

the Golgi fragmentation (*17*). However, till now the roles of FMNLs especially

FMNL2 on oocyte meiosis are still largely unknown.

In the present study, we disturbed the FMNL2 expression and explored the roles

of FMNL2 during mouse oocyte meiosis. Our results showed that FMNL2 was

essential for the polar body size control and successful extrusion; and these abnormal

phenotypes might be due to aberrant actin-based meiotic spindle migration.

Meanwhile, we also found that FMNL2 was essential for the functions and

distribution of mitochondria and endoplasmic reticulum. Therefore, this study

provided the evidence for the critical roles of FMNL2-mediated actin on spindle

movement and organelle dynamics in mammalian oocytes.

Materials and Methods

Antibodies and chemicals

Rabbit monoclonal anti-FMNL2 antibody, rabbit monoclonal anti-Arp2 antibody,

mouse monoclonal anti-profilin1 antibody were from Santa Cruz (Santa Cruz, CA,

USA). Rabbit monoclonal anti-Fascin antibody was purchased form Abcam

(Cambridge, UK). Rabbit polyclonal anti-INF2 antibody was purchased from

Proteintech (Proteintech, CHI, USA). Rabbit monoclonal anti-α-tubulin (11H10)

antibody, rabbit monoclonal anti-Grp78 antibody, rabbit monoclonal anti-cofilin

antibody and rabbit monoclonal anti-Chop antibody were from Cell Signaling

Technology (Beverly, MA, USA). Mouse monoclonal anti-α-tubulin-FITC antibody

was from Sigma-Aldrich Corp. (St. Louis, MO, USA). FITC-conjugated goat anti-

rabbit IgG were from Zhongshan Golden Bridge Biotechnology (Beijing). ER-Tracker

Red, Mito-Tracker Green and enhanced mitochondrial membrane potential assay Kit

were from Beyotime Biotechnology (Shanghai). All other chemicals and reagents

were from Sigma-Aldrich Corp., unless otherwise stated.

Ethics statement and oocyte culture

We followed the guidelines of Animal Research Institute Committee of Nanjing

Agricultural University to conduct the operations. The animal facility had license

authorized by the experimental animal committee of Jiangsu Province (SYXK-Su-

20170007). The Female Institute of Cancer Research (ICR) mice, aged 4–6 weeks,

were kept in a room with a regulated temperature of 22 °C and provided with a

standard diet. Fully developed germinal vesicle stage (GV) oocytes were retrieved

from the ovaries of mice, and then cultured in M16 medium with para□n oil at 37 °C

and in the presence of 5% CO_2 for in vitro maturation. At specific intervals, the

oocytes were collected for various tests and analyses. The oocytes were placed at

37°C with an atmosphere of 5% CO_2, and cultured to different time points for

immunostaining, microinjection and western blot.

Plasmid construct and in *vitro* transcription

Template RNA was generated from mouse ovaries with RNA Isolation Kit

(Thermos fisher), then we reversed transcription of these RNA to create cDNA by a

PrimeScript 1st strand cDNA synthesis kit (Takara, Japan). FMNL2-EGFP vector was

generated by Wuhan GeneCreate Biological Engineering Co, Ltd. mRNA was

synthesized from linearized plasmid using HiScribe T7 high yield RNA synthesis kit

(NEB), then capped with m7G (5') ppp (5') G (NEB) and tailed with a poly(A)

polymerase tailing kit (Epicentre) and purified with RNA clean & concentrator-25 kit

(Zymo Research).

Microinjection of FMNL2 siRNA and mRNA

The FMNL2 siRNA working solution was dissolved in RNase-free water,

achieving a concentration of 20 μM. For FMNL2 knockdown (KD), three individual

siRNA strands were precisely mixed and subjected to centrifugation to obtain the

supernatant, approximately 5-10 pL of supernatant was microinjected into the

cytoplasm of GV stage oocytes. In contrast, an equal volume of a negative control

solution was microinjected into the cytoplasm of oocytes in the control group.

FMNL2 siRNA: 5′- GCU GAA UGC UAU GAA CCU ATT-3′, 5′- GCC AUU GAU

CUU UCU UCA ATT-3′, 5′- GGA AUU AAG AAG GCG ACA ATT-3′; Negative

control siRNA: 5′-UUC UCC GAA CGU GUC ACG UTT-3′. Then, the oocytes were

arrested in the GV stage for 18h in M16 medium with 5μM milrinone. This was done

to optimize the effectiveness of the siRNA and aid in the depletion of FMNL2. For the

rescue experiment, 5–10 pL of 200ng/μL FMNL2-EGFP mRNA was injected into the

GV oocytes 18h after FMNL2 siRNA injection. Following that, the GV stage oocytes

were cultured in M16 medium with 5μM milrinone for 4h. Then, the oocytes being

cultured in fresh M16 medium for subsequent experiments.

Immunofluorescent staining and confocal microscopy

Oocytes were fixed in 4% paraformaldehyde (in PBS) for 30 min and

permeabilized with 0.5% Triton X-100 in PBS for 20min then blocked in blocking

buffer (1% BSA-supplemented PBS) at room for 1 h. For FMNL2 staining, the

oocytes at various stages (GV, GVBD, MI, and MII) were incubated with Rabbit

monoclonal anti-FMNL2 antibody (1:100) at 4 °C overnight, then oocytes were

washed by wash buffer (0.1% Tween 20 and 0.01% Triton X-100 in PBS) for 3 times

(5 min each time). Next the oocytes were labeled with secondary antibody coupled to

FITC-conjugated goat anti-rabbit IgG (1:100) at room temperature for 1 h. For α-

tubulin staining, MI stage oocytes were incubated with anti-α-tubulin-FITC antibody

(1:200). For actin staining, GV and MI stage oocytes were incubated with Phalloidin-

TRITC at room temperature for 2 h. Then the oocytes were washed as the same way.

Finally, oocytes were incubated with Hoechst 33342 at room temperature for 10-20

min. After staining, samples were mounted on glass slides and observed with a

confocal laser-scanning microscope (Zeiss LSM 800 META, Germany).

ER and Mito-tracker staining

To study ER and mitochondria distribution during mouse oocyte meiosis, MI

stage oocytes were incubated with ER-Tracker Red (1:3000) or 200 nM Mito-tracker

green (Red) in M16 medium for 30 min at 37℃ and 5% CO_2. Then the oocytes were

washed three times with M2 medium, finally the samples were examined with

confocal microscopy. During the MI stage of oocyte development, both the ER and

mitochondria evenly distributed in the cytoplasm and accumulated at the spindle

periphery in MI stage. This clustering phenomenon is considered as the normal

localization pattern of these organelles. On the other hand, when the ER and

mitochondria cluster randomly within the cytoplasmic region, this localization is

defined as organelle abnormal localization pattern.

JC-1 detection

165　The enhanced mitochondrial membrane potential assay Kit was employed to

166　analyze the mitochondrial membrane potential of oocytes. The MI stage oocytes were

167　transferred from the M16 medium to JC-1 for 30 min at 37℃ and 5% CO_2. Following

168　three washes with M2 medium, the oocytes were examined using a fluorescent

169　microscope (OLYMPUS IX71, Japan) for the presence of a fluorescent signal.

Time lapse microscopy

170　To image the dynamic changes that occurred during oocyte maturation, oocytes

171 were cultured in M16 medium, then transferred to the Leica SD AF confocal imaging

172 system equipped with 37 °C incubator and 5% CO2 supply (H301-K-FRAME). The

173 spindle in oocytes was labeled by α-tubulin-EGFP.

Immunoprecipitation

4-6 ovaries were put into RIPA Lysis Buffer contained phosphatase inhibitor

cocktail (100×) (Kangwei Biotechnology, China), and were completely cleaved on ice

block. We collected supernatant after centrifugation (13200 rps, 20 min) and then took

out 50 μl as input sample at 4 °C. The rest of the supernatant was incubated with

primary antibody (FMNL2 or INF2 antibody) overnight at 4 °C. 30 μl conjugated

beads (washed five times in PBS) were added to the supernatant/antibody mixture and

incubated at 4 °C for 4-6 h, after three times wash by immune complexes, the samples

were then released from the beads by mixing in 2× SDS loading buffer for 10min at

30 °C.

Western blot analysis

Approximate 100-150 GV or MI stage mouse oocytes were placed in Laemmli

sample buffer and heated at 85□ for 7-10 min. Proteins were separated by SDS-PAGE

at 165V for 70-80 min and then electrophoretically transferred to polyvinylidene

fluoride (PVDF) membranes (Millipore, Billerica, MA, USA) at 20 V for 1 hour.

After transfer, the membranes were then blocked with TBST (TBS containing 0.1%

Tween 20) containing 5% non-fat milk at room temperature for 90 min. After

blocking, the membranes were incubated with rabbit monoclonal anti-FMNL2

antibody (1:500), rabbit monoclonal anti-Arp2 antibody (1:500), mouse monoclonal

anti-profilin1 antibody (1:500), rabbit monoclonal anti-Fascin antibody (1:5000),

rabbit polyclonal anti-INF2 antibody (1:500), rabbit monoclonal anti-Grp78 antibody

(1:1000), rabbit monoclonal anti-cofilin antibody (1:2000), rabbit monoclonal anti-

CHOP antibody (1:1000), or rabbit monoclonal anti-tubulin antibody (1:2000) at 4 °C

overnight. After washing 5 times in TBST (5 min each), membranes were incubated

for 1h at room temperature with HRP-conjugated Pierce Goat anti-Rabbit IgG (1:5000)

or HRP-conjugated Pierce Goat anti-mouse IgG (1:5000). After washing for 5 times,

the membranes were visualized using chemiluminescence reagent (Millipore,

Billerica, MA). Every experiment repeated at least 3 times with different samples.

Fluorescence Intensity Analysis

Immunofluorescence experiments were conducted simultaneously and with

consistent parameters in both the control and treatment groups. The images were

consistently captured using identical confocal microscope settings. Subsequently, the

average intensity of fluorescence per unit area in the designated region of interest was

quantified following the fluorescence staining. The acquired fluorescence data was

analyzed employing ZEN 2011 and ImageJ software.

Statistical analysis

All statistical analyses were performed using GraphPad Prism7.00 software

(GraphPad, CA, USA), employing the t-test to assess the statistical significance

between the control and treatment groups. The results were represented as the mean ±

standard error of the mean (SEM). Statistical significance was defined as a P-value <

0.05, denoted as *, ** for P < 0.01, ***, and **** for P < 0.001 and P < 0.0001,

respectively. Every experiment was conducted with a minimum of three biological

replicates.

Results

Expression and subcellular localization of FMNL2 during mouse oocyte maturation

We first examined FMNL2 expression in mouse oocytes at different stages. The

results indicated that FMNL2 all expressed in GV, MI and MII stages during mouse

oocyte maturation (GV, 1; MI, 0.82 ± 0.07; MII, 0.61 ± 0.10, Figure 1A). Next, we

performed FMNL2-EGFP mRNA microinjection to examine the localization of

FMNL2. As shown in Figure 1B, FMNL2 accumulated at the oocyte cortex during the

GV, GVBD MI and MII stages. Besides, FMNL2 also localized at the spindle

periphery during MI stages. The FMNL2 antibody staining results also confirmed this

localization pattern. In addition, we co-stained FMNL2 antibody with F-actin, and the

results revealed that both FMNL2 and F-actin are localized in the cortex region of

oocytes (Figure 1C). The FMNL2 localization pattern indicated that FMNL2 might

interact with actin dynamics during oocyte meiosis.

FMNL2 is essential for polar body extrusion and asymmetric division in mouse oocytes

To investigate the functional roles of FMNL2 in mouse oocytes, we employed

FMNL2 siRNA microinjection to knockdown FMNL2 protein expression. A

significant decrease of FMNL2 protein level was shown in FMNL2-KD oocytes

compared to control group by western blot (1 vs. 0.48 ± 0.08, $P < 0.01$, Figure 2A).

We then examined the maturation of oocytes, and the results indicated that

knockdown of FMNL2 had an impact on the extrusion of the first polar body.

Moreover, a significant proportion of oocytes exhibited larger polar bodies upon

extrusion (Figure 2B). Based on the size of the extruded polar bodies, those with a

diameter exceeding one-third of the oocyte's diameter were categorized as large polar

bodies. Consequently, we proceeded to calculate the rates of polar body extrusion and

the generation of large polar bodies in the oocytes. The quantitative results also

confirmed this phenotype (rate of polar body extrusion: $74.26 \pm 1.44\%$, n = 439 vs.

$59.5 \pm 2.82\%$, n = 398, $P < 0.001$, Figure 2C; rate of large polar bodies: $19.05 \pm$

1.97%, n = 311 vs. $37.16 \pm 1.87\%$, n = 257, $P < 0.0001$, Figure 2D). In addition, live-

cell imaging was used to determine the dynamic changes that occurred during oocyte

maturation, and the results showed that the oocytes either failed to undergo

cytokinesis or divided from the central axis of the oocytes and formed big polar

bodies (Figure 2E). To further confirm the phenotype, we performed FMNL2 rescue

experiments by expressing exogenous Fmnl2 mRNA in FMNL2-depleted oocytes

(Figure 2F), we found that exogenous Fmnl2 mRNA expression rescued first polar

body extrusion and large polar body defects (Figure 2G). The quantitative results also

confirmed this phenotype (rate of polar body extrusion: $48.34 \pm 4.2\%$, n = 355 vs.

$62.62 \pm 3.6\%$, n = 377, $P < 0.01$, Figure 2H; rate of large polar bodies: $30.93 \pm 2\%$, n

= 193 vs. $9.58 \pm 2.4\%$, n = 203, $P < 0.01$, Figure 2I). It is known that knockdown of

FMNL3 leads to inhibition of oocyte maturation. To investigate whether FMNL2

exhibits an additive effect with FMNL3 in terms of functionality, we simultaneously

knocked down both FMNL2 and FMNL3. The results demonstrated that simultaneous

knockdown of the two FMNL proteins, compared to the control group, resulted in a

decrease in oocyte maturation rate. However, when compared to the single

knockdown of FMNL2, the double knockdown of FMNL2 and FMNL3 did not cause

more severe defects in polar body extrusion. (polar body extrusion, Control: 70.97 ±

1.23%, n=261 vs FMNL2+3-KD: 60.42 ± 2.99%, n=198, P < 0.05, Figure 2J. Large

polar body, Control: 10.85 ± 0.97%, n=172 vs FMNL2+3-KD: 32.90 ± 1.88%, n=118,

P < 0.001, Figure 2K). These results suggested that FMNL2 played critical roles for

the polar body extrusion and asymmetric division during mouse oocyte maturation.

FMNL2 regulates spindle migration during mouse oocyte maturation

To investigate the causes for polar body extrusion defects, we examined the

spindle migration by time-lapse microscopy during oocyte meiosis. As shown in

Figure 3A, in the control oocyte, the meiotic spindle formed in the center of the

oocyte after culture 8 h and moved to the oocyte cortex at 9.5h; and the polar body

was extruded at 11-12h, with a spindle formed near the cortex at MII stage. However,

in FMNL2-KD oocytes, two phenotypes were observed: 1) the meiotic spindle

remained in the center of the oocyte until 10 h, and then the oocytes initiated the

cytokinesis at 10.5 h but failed to extrude the polar body; 2) some oocytes with

arrested spindles initiated the cytokinesis but extruded a big polar body (Figure 3A).

This indicated the failure of spindle migration after FMNL2 depletion. We analyzed

the rate of cortex-localized spindle in oocytes by cultured for 9.5 h, and the result

showed that the rate of migrated spindles in control oocytes was significantly higher

than that in FMNL2-KD oocytes (59.94 ± 3.42%, n = 78 vs. 38.97 ± 6.34%, n = 64, P

< 0.05, Figure 3B). We also performed FMNL2 rescue experiments. Supplementing

with exogenous Fmnl2 rescued the spindle migration defects compared with the

FMNL2-depletion group (40.27 ± 3.19%, n = 181 vs. 57.01± 2.72%, n = 57, P < 0.01,

Figure 3C). These results suggested that FMNL2 might be involved in spindle

migration in mouse oocytes.

FMNL2 promotes cytoplasmic actin assembly during mouse oocyte maturation

As FMNL2 is a key actin assembly factor, we further investigated actin assembly

after deleting FMNL2 in mouse oocytes. Surprisingly there was no significant

difference for the signals of cortex actin was observed between control oocytes and

FMNL2-KD oocytes, which was confirmed by the fluorescence intensity analysis

(30.88 ± 1.10, n = 28 vs. 30.58 ± 1.12, n = 28, P > 0.05, Figure 4A, 4B). However, we

found a significant decrease of cytoplasmic actin signals in the FMNL2-KD oocytes,

and the statistical analysis for the cytoplasmic actin fluorescent signals also confirmed

our findings (58.25 ± 2.05, n = 26 vs. 37.92 ± 2.02, n = 24, P < 0.0001, Figure 4C,

4D). Moreover, the rescue experiments showed that exogenous FMNL2 rescued the

decrease of cytoplasmic actin filaments compared with the FMNL2-depletion group

(37.98 ± 1.98, n = 16 vs. 54.72± 2.88, n = 15, P < 0.0001, Figure 4E, 4F). We next

explored how FMNL2 regulates cytoplasmic actin assembly in oocytes. By mass

spectrometry analysis we found there were several actin-related potential candidates

which might be related with FMNL2 (Figure 4G). Co-immunoprecipitation results

showed that FMNL2 precipitated Arp2 and Formin2 but not Profilin and fascin

(Figure 4H). To further verify the correlation between FMNL2 and Arp2 and Formin2,

we then examined Arp2 and Formin2 protein expression after FMNL2 knockdown.

The results showed Arp2 protein expression increased significantly after FMNL2

knockdown (1 vs. 1.56 ± 0.07, P < 0.001, Figure 4I) but Formin2 decreased after

FMNL2 knockdown (1 vs. 0.62 ± 0.04, P < 0.001, Figure 4J). Exogenous FMNL2

rescued these alterations compared with that in the FMNL2-KD group (Arp2 protein

expression: 1 vs. 0.65 ± 0.06, P < 0.01, Figure 4I; Formin2 protein expression: 1 vs.

1.24 ± 0.05, P < 0.01, Figure 4J). These results indicated that FMNL2 may be

associated with Formin2 and Arp2 for actin assembly in mouse oocytes.

FMNL2 regulates endoplasmic reticulum distribution during mouse oocyte maturation

The mass spectrometry analysis data indicated that several ER-related potential

candidates which might be related with FMNL2 (Figure 5A), while INF2, a typical

protein which mediates actin polymerization at ER showed high confidence level. We

then examined the relationship between FMNL2 and INF2, and the co-

immunoprecipitation results showed that FMNL2 precipitated INF2 and INF2 also

precipitated FMNL2 (Figure 5B), indicating that FMNL2 interacted with INF2 in

mouse oocytes. We then examined the ER distribution in FMNL2-KD oocytes. As

shown in Figure 5C, in control oocytes the ER evenly distributed in the cytoplasm and

accumulated at the spindle periphery in MI stage; however, ER agglomerated in

cytoplasm in FMNL2-KD oocytes (Figure 5C). The statistical analysis showed that

the abnormal distribution of ER increased significantly in the FMNL2-KD group

(28.91 ± 5.62, n = 27 vs. 59.64 ± 6.95, n = 28, P < 0.05, Figure 5D). The localization

pattern of ER indicated its functions might be disturbed. In FMNL2-KD oocytes, we

found the expressions of ER-stress related proteins Grp78 and Chop were

significantly increased (Grp78: 1 vs. 1.42 ± 0.12, P < 0.05; Chop: 1 vs. 1.53 ± 0.16, P

< 0.05. Figure 5E), indicating the occurrence of ER stress. We also performed

FMNL2 rescue experiments. Supplementing with exogenous Fmnl2 rescued the ER

distribution defects caused by FMNL2 knockdown (Figure 5F), which was supported

by the statistical analysis showing that the abnormal distribution rate of ER decreased

significantly in the rescue group (52.04 ± 5.29, n = 70 vs. 34.91 ± 3.37, n = 78, P <

0.05, Figure 5G). Moreover, Grp78 protein expression decreased in the rescue group

(1 vs. 0.78 ± 0.05, P < 0.01. Figure 5H). These results indicated that the depletion of

FMNL2 affected ER distribution and caused ER stress in mouse oocytes.

FMNL2 regulates mitochondrial distribution during mouse oocyte maturation

As INF2 is also related to the mitochondrial connection of ER, we further

screened up the mass spectrometry analysis data and we found many mitochondria-

related potential candidates which might be related with FMNL2 (Figure 6A).

Therefore, we further examined the distribution of mitochondria in FMNL2-KD

oocytes. In control oocytes, the mitochondria evenly distributed in the cytoplasm and

accumulated at the spindle periphery in MI stage; however, in FMNL2-KD oocytes,

mitochondria presented clumped aggregation distribution in cytoplasm (Figure 6B).

We counted the number of clumps and found that the uniform distribution of

mitochondria decreased significantly in the FMNL2-KD group (59.66 ± 8.48, n = 31

vs. 20.83 ± 4.17, n = 32, P < 0.05, Figure 6C). A large number of FMNL2-KD oocytes

agglomerated into one to three clumps (22.73 ± 4.27, n = 31 vs. 42.50 ± 1.25, n = 32, P

< 0.05, Figure 6C). Supplementing with exogenous Fmnl2 rescued the mitochondria

distribution (Figure 6D), the statistical analysis showed that the uniform distribution

of mitochondria increased significantly in the rescue group (36.49 ± 3.97, n = 53 vs.

53.90 ± 2.09, n = 79, P < 0.05, Figure 6E). We also examined mitochondrial

membrane potential, and the results showed that FMNL2 depletion caused the

alterations of mitochondrial membrane potential (MMP) by JC-1 staining. The

fluorescence intensity of JC-1 red channel was decreased compared with the control

group (Figure 6F). We also calculated the ratio for red/green fluorescence intensity,

and the results also confirmed this (control group: 0.40 vs. FMNL2-KD: 0.21 ± 0.01,

P < 0.01) (Figure 6G). Cofilin is an important factor of actin assembly and regulates

mitochondrial function. We also examined cofilin protein expression after FMNL2

knockdown. The results showed cofilin protein expression decreased significantly

after FMNL2 knockdown (1 vs. 0.81 ± 0.03, P < 0.01, Figure 6H). These results

indicated that FMNL2 regulated mitochondria distribution and function during mouse

oocyte maturation.

Discussion

In this study, we explored the functions of FMNL2 during mouse oocyte meiosis.

Our results indicated that FMNL2 regulated actin-based spindle migration for

asymmetric cell division of oocytes, and more importantly FMNL2 was critical for

maintaining the distribution of the ER and mitochondria, which set up a link for actin-

related spindle migration and organelle dynamics in oocytes.

As a subfamily of Formin family, FMNLs play an important role in regulating

actin filaments (*18*), while FMNL2 is most widely expressed in variety of cell models

among the members of FMNLs. In this study, we showed that FMNL2 expressed in

mouse oocytes and it mainly accumulated at the oocyte cortex and spindle periphery,

which was similar with the actin distribution pattern in oocytes. This specific

localization is also similar to FMN2, a well-studied factor in the formin family for

spindle migration during oocyte meiosis (*7, 19*). In addition, another FMNLs family

member, FMNL1 is also localized at the cortex and is essential for actin

polymerization and spindle assembly during oocyte meiosis (*20*). Based on the

localization pattern of FMNL2, we speculated that the functions of FMNL2 might be

also involved in actin-related process during mouse oocyte meiosis.

To confirm our hypothesis, we depleted FMNL2 protein expression and we found

that absence of FMNL2 caused the aberrant first polar body extrusion. The oocytes

either failed to form the polar body or extruded large polar bodies. These phenotypes

caused by FMNL2 depletion are similar to the other actin-related proteins during

oocyte maturation such as Arp2/3 complex (*8, 21*) and FMN2 (*22, 23*). We next

examined the actin distribution in oocytes since it is reported that FMNL2 promotes

actin filament assembly in many models. FMNL2 is required for cell-cell adhesion

formation by regulating the actin assembly (*24*), and FMNL2 could directly drives

actin elongation (*15*). In CRC cells, cortactin bind to FMNL2 to active the actin

polymerization, and FMNL2 is important for invadopodia formation and functions

(*25*). Our results showed that the FMNL2 depletion caused significantly decrease in

cytoplasmic actin, indicating the conserved roles of FMNL2 on actin assembly in

mammalian oocyte model. Other Formin family proteins such as Daam1, FHOD1,

and Formin-homology family protein mDia1 are also reported to affect oocyte

meiosis by regulating actin polymerization (*26-28*).

We then tried to explore how FMNL2 involves into the actin assembly in oocytes.

Mass spectrometry analysis data indicated that FMNL2 associated with several actin-

related proteins, and we found that a potential association between FMNL2 and Arp2

/Formin2. This could be confirmed by the altered expression of these two molecules

after FMNL2 depletion. Therefore, we speculated FMNL2 could regulate cytoplasmic

actin assembly in oocytes through the association with Formin2 since it is reported to

be an important protein for cytoplasmic actin assembly in oocytes (*22*). Interestingly,

our results showed that unlike the reduction of cytoplasmic actin, cortex actin was not

affected by the absence of FMNL2. We speculate that FMNL2 and Arp2/3 both

contribute to the cortex actin dynamics, when FMNL2 decreases, ARP2 increases to

compensate for this, which maintains the cortex actin level. As an actin nucleator

Arp2/3 complex localizes at the cortex and is essential for actin polymerization during

oocyte meiosis (*8, 29*). These results suggested that FMNL2 might be involved in

cytokinesis and asymmetric division by regulating actin assembly during mouse

oocyte maturation.

The spindle migration is a key step in ensuring the asymmetric division for

oocytes (*30*). In mitosis, spindle position is decided by cortical actin and astral

microtubules; in contrast, spindle migration is mainly mediated by actin filaments

during oocyte meiosis (*30, 31*). Due to the effects of FMNL2 on asymmetric division

and cytoplasmic actin, we analyzed the spindle positioning at late MI, we found that

the spindle migration was disturbed after FMNL2 depletion, no matter the cytokinesis

occurred or not. Several formin proteins are shown to regulate spindle migration

during oocytes meiosis. For example, FMN2 nucleates actin surrounding the spindle,

pushing force generated by actin to trigger the spindle migration (*19, 22*), and cyclin-

dependent kinase 1 (Cdk1) induces cytoplasmic Formin-mediated F-actin

polymerization to propel the spindle into the cortex (*32*). Our previous studies also

showed that absence of the formin family member FMNL1 or FHOD1 could lead to

the decrease of cytoplasmic actin to prevent the spindle migration (*20, 27*). We

speculated that FMNL2 together with other Formin proteins, conservatively regulate

actin-mediated spindle migration during oocyte meiosis.

Another important finding is that through the mass spectrometry analysis we

found many candidate proteins which were related with ER, and our results indicated

that FMNL2 was essential for the maintenance of ER distribution in the cytoplasm.

Moreover, the loss of FMNL2 induced ER stress, showing with altered expression of

GRP78 and CHOP. Proper distribution of ER is important for the oocyte quality. ER

displays a homogeneous distribution pattern throughout the entire ooplasm during

development of oocytes and embryos from diabetic mice(*33*). During the transition of

mouse oocytes from MI to MII phase, actin regulates cortical ER aggregation(*34*). In

addition, Formin2 is shown to colocalize with the ER during oocyte meiosis and the

ER-associated Formin2 at the spindle periphery is required for MI chromosome

migration(*6*). In our results we showed that FMNL2 associated with INF2 protein in

oocytes. INF2 is an ER-associated protein, and the expression of GFP-INF2 which

containing DAD/WH2 mutations causes the ER to collapse around the nucleus (*35*).

We concluded that FMNL2 might regulate INF2 for the distribution of ER in

cytoplasm of oocytes.

Besides its roles of ER distribution, it is shown that INF2 also affects

mitochondrial length and ER-mitochondrial interaction in an actin-dependent manner

(*35, 36*). It is shown that INF2 regulates Drp1 for mitochondrial fission, and INF2-

induced actin filaments may drive initial mitochondrial constriction, which allows

Drp1-driven secondary constriction(*36, 37*). In addition, we also found many

candidate proteins which were related with mitochondria from mass spectrometry

analysis. During oocyte meiosis, mitochondria gradually accumulated around the

spindle after GVBD, and the spindle-peripheral FMN2 and its actin nucleation

activity are important for the accumulation of mitochondria in this region (*19*). Our

results found that FMNL2 depletion caused agglutination of mitochondria and altered

MMP level in the cytoplasm, indicating its roles on the mitochondria distribution and

functions. Another formin protein mDia1 is shown to be necessary to induce the

anchoring of mitochondria along the cytoskeletal in mammalian CV-1 cells and

Drosophila BG2-C2 neuronal cells(*38*). Moreover, the formin interaction protein

Spire1C binds INF2 to promote actin assembly on mitochondrial surfaces, and

Spire1C disruption could reduce mitochondrial constriction and division(*39*). In

addition, our result indicated that cofilin expression decreased in FMNL2 depletion

oocytes. Cofilin is an actin-depolymerizing factor and its localization at the

mitochondrial fission site is crucial for inducing mitochondrial fission and mitophagy

(*40*). Depleting of cofilin resulted in abnormal interconnection and elongation of

mitochondria (*41*). Together with its roles on ER, these data indicated that FMNL2

might associate with INF2 and cofilin for the actin-based organelle distribution during

oocyte meiosis.

Collectively, we provide a body of evidence showing that FMNL2 associates with

Formin2 and Arp2/3 complex for actin assembly, which further regulates spindle

migration and INF2/Cofilin-related organelle dynamics during mouse oocyte

maturation.

Data Availability

All data generated or analyzed during this study are included in this published

article

Acknowledgement

We are particularly grateful to Xiao-Yan Fan and Xing-Hua Wang from Fertility

Preservation Laboratory, Reproductive Medicine Center, Guangdong Second

Provincial General Hospital for their technical assistance of live cell imaging system.

Contributions

MHP and SCS designed the study. MHP performed the majority of the experiments.

SML, ZNP, MHS, XHL, JQJ, YZ contributed to the regents and materials. MHP,

XHO and SCS analyzed the data. MHP and SCS wrote the manuscript.

Competing interests

466 There is no conflict of interest to declare.

Ethics approval and consent to participate

468 Not applicable.

Consent for publication

470 Not applicable.

References

1. K. Yi, B. Rubinstein, R. Li, Symmetry breaking and polarity establishment during

mouse oocyte maturation. *Philosophical transactions of the Royal Society of London.*

Series B, Biological sciences **368**, 20130002 (2013).

2. B. Pan, J. Li, The art of oocyte meiotic arrest regulation. *Reproductive biology and*

endocrinology : RB&E **17**, 8 (2019).

3. Q. Y. Sun, H. Schatten, Regulation of dynamic events by microfilaments during

oocyte maturation and fertilization. *Reproduction* **131**, 193-205 (2006).

4. S. C. Sun, N. H. Kim, Molecular mechanisms of asymmetric division in oocytes.

Microscopy and microanalysis : the official journal of Microscopy Society of America,

Microbeam Analysis Society, Microscopical Society of Canada **19**, 883-897 (2013).

5. X. Duan, S. C. Sun, Actin cytoskeleton dynamics in mammalian oocyte meiosis. *Biol*

Reprod **100**, 15-24 (2019).

6.   K. Yi, B. Rubinstein, J. R. Unruh, F. Guo, B. D. Slaughter, R. Li, Sequential actin-

based pushing forces drive meiosis I chromosome migration and symmetry breaking

in oocytes. *The Journal of cell biology* **200**, 567-576 (2013).

7.   H. Li, F. Guo, B. Rubinstein, R. Li, Actin-driven chromosomal motility leads to

symmetry breaking in mammalian meiotic oocytes. *Nature cell biology* **10**, 1301-1308

(2008).

8.   S. C. Sun, Z. B. Wang, Y. N. Xu, S. E. Lee, X. S. Cui, N. H. Kim, Arp2/3 complex

regulates asymmetric division and cytokinesis in mouse oocytes. *PloS one* **6**, e18392

(2011).

9.   S. Pfender, V. Kuznetsov, S. Pleiser, E. Kerkhoff, M. Schuh, Spire-type actin

nucleators cooperate with Formin-2 to drive asymmetric oocyte division. *Current*

biology : CB **21**, 955-960 (2011).

10.   S. Kuhn, M. Geyer, Formins as effector proteins of Rho GTPases. *Small GTPases* **5**,

e29513 (2014).

11.   M. Katoh, M. Katoh, Identification and characterization of human FMNL1, FMNL2 and

FMNL3 genes in silico. *International journal of oncology* **22**, 1161-1168 (2003).

12.   M. Gardberg, K. Talvinen, K. Kaipio, K. Iljin, C. Kampf, M. Uhlen, O. Carpen,

Characterization of Diaphanous-related formin FMNL2 in human tissues. *BMC cell*

biology **11**, 55 (2010).

13.   F. Kage, M. Winterhoff, V. Dimchev, J. Mueller, T. Thalheim, A. Freise, S. Bruhmann,

J. Kollasser, J. Block, G. Dimchev, M. Geyer, H. J. Schnittler, C. Brakebusch, T. E.

Stradal, M. F. Carlier, M. Sixt, J. Kas, J. Faix, K. Rottner, FMNL formins boost

507   lamellipodial force generation. *Nature communications* **8**, 14832 (2017).

508   14.   X. L. Zhu, Y. F. Zeng, J. Guan, Y. F. Li, Y. J. Deng, X. W. Bian, Y. Q. Ding, L. Liang,

509   FMNL2 is a positive regulator of cell motility and metastasis in colorectal carcinoma.

510   *The Journal of pathology* **224**, 377-388 (2011).

511   15.   J. Block, D. Breitsprecher, S. Kuhn, M. Winterhoff, F. Kage, R. Geffers, P. Duwe, J. L.

512   Rohn, B. Baum, C. Brakebusch, M. Geyer, T. E. Stradal, J. Faix, K. Rottner, FMNL2

513   drives actin-based protrusion and migration downstream of Cdc42. *Current biology :*

514   *CB* **22**, 1005-1012 (2012).

515   16.   Y. Li, X. Zhu, Y. Zeng, J. Wang, X. Zhang, Y. Q. Ding, L. Liang, FMNL2 enhances

516   invasion of colorectal carcinoma by inducing epithelial-mesenchymal transition.

517   *Molecular cancer research : MCR* **8**, 1579-1590 (2010).

518   17.   F. Kage, A. Steffen, A. Ellinger, C. Ranftler, C. Gehre, C. Brakebusch, M. Pavelka, T.

519   Stradal, K. Rottner, Author Correction: FMNL2 and -3 regulate Golgi architecture and

520   anterograde transport downstream of Cdc42. *Scientific reports* **9**, 18008 (2019).

521   18.   D. Breitsprecher, B. L. Goode, Formins at a glance. *Journal of cell science* **126**, 1-7

522   (2013).

523   19.   X. Duan, Y. Li, K. Yi, F. Guo, H. Wang, P. H. Wu, J. Yang, D. B. Mair, E. A. Morales,

524   P. Kalab, D. Wirtz, S. X. Sun, R. Li, Dynamic organelle distribution initiates actin-

525   based spindle migration in mouse oocytes. *Nature communications* **11**, 277 (2020).

526   20.   F. Wang, L. Zhang, X. Duan, G. L. Zhang, Z. B. Wang, Q. Wang, B. Xiong, S. C. Sun,

527   RhoA-mediated FMNL1 regulates GM130 for actin assembly and phosphorylates

528   MAPK for spindle formation in mouse oocyte meiosis. *Cell cycle* **14**, 2835-2843

(2015).

21.   K. Yi, J. R. Unruh, M. Deng, B. D. Slaughter, B. Rubinstein, R. Li, Dynamic

maintenance of asymmetric meiotic spindle position through Arp2/3-complex-driven

cytoplasmic streaming in mouse oocytes. *Nature cell biology* **13**, 1252-1258 (2011).

22.   J. Dumont, K. Million, K. Sunderland, P. Rassinier, H. Lim, B. Leader, M. H. Verlhac,

Formin-2 is required for spindle migration and for the late steps of cytokinesis in

mouse oocytes. *Developmental biology* **301**, 254-265 (2007).

23.   B. Leader, H. Lim, M. J. Carabatsos, A. Harrington, J. Ecsedy, D. Pellman, R. Maas,

P. Leder, Formin-2, polyploidy, hypofertility and positioning of the meiotic spindle in

mouse oocytes. *Nature cell biology* **4**, 921-928 (2002).

24.   K. Grikscheit, T. Frank, Y. Wang, R. Grosse, Junctional actin assembly is mediated

by Formin-like 2 downstream of Rac1. *The Journal of cell biology* **209**, 367-376

(2015).

25.   X. L. Ren, Y. D. Qiao, J. Y. Li, X. M. Li, D. Zhang, X. J. Zhang, X. H. Zhu, W. J. Zhou,

J. Shi, W. Wang, W. T. Liao, Y. Q. Ding, L. Liang, Cortactin recruits FMNL2 to

promote actin polymerization and endosome motility in invadopodia formation.

Cancer letters **419**, 245-256 (2018).

26.   Y. Lu, Y. Zhang, M. H. Pan, N. H. Kim, S. C. Sun, X. S. Cui, Daam1 regulates fascin

for actin assembly in mouse oocyte meiosis. *Cell cycle* **16**, 1350-1356 (2017).

27.   M. H. Pan, F. Wang, Y. Lu, F. Tang, X. Duan, Y. Zhang, B. Xiong, S. C. Sun, FHOD1

regulates cytoplasmic actin-based spindle migration for mouse oocyte asymmetric

cell division. *Journal of cellular physiology* **233**, 2270-2278 (2018).

28.   Y. Zhang, F. Wang, Y. J. Niu, H. L. Liu, R. Rui, X. S. Cui, N. H. Kim, S. C. Sun,

Formin mDia1, a downstream molecule of FMNL1, regulates Profilin1 for actin

assembly and spindle organization during mouse oocyte meiosis. *Biochimica et*

biophysica acta **1853**, 317-327 (2015).

29.   E. D. Goley, M. D. Welch, The ARP2/3 complex: an actin nucleator comes of age.

Nature reviews. Molecular cell biology **7**, 713-726 (2006).

30.   S. Brunet, B. Maro, Cytoskeleton and cell cycle control during meiotic maturation of

the mouse oocyte: integrating time and space. *Reproduction* **130**, 801-811 (2005).

31.   S. Reinsch, P. Gonczy, Mechanisms of nuclear positioning. *Journal of cell science*

111 (Pt 16), 2283-2295 (1998).

32.   Z. Wei, J. Greaney, C. Zhou, A. H. H, Cdk1 inactivation induces post-anaphase-onset

spindle migration and membrane protrusion required for extreme asymmetry in

mouse oocytes. *Nature communications* **9**, 4029 (2018).

33.   C. H. Zhang, W. P. Qian, S. T. Qi, Z. J. Ge, L. J. Min, X. L. Zhu, X. Huang, J. P. Liu, Y.

C. Ouyang, Y. Hou, H. Schatten, Q. Y. Sun, Maternal diabetes causes abnormal

dynamic changes of endoplasmic reticulum during mouse oocyte maturation and

early embryo development. *Reproductive biology and endocrinology : RB&E* **11**, 31

(2013).

34.   G. FitzHarris, P. Marangos, J. Carroll, Changes in endoplasmic reticulum structure

during mouse oocyte maturation are controlled by the cytoskeleton and cytoplasmic

dynein. *Dev Biol* **305**, 133-144 (2007).

35.   E. S. Chhabra, V. Ramabhadran, S. A. Gerber, H. N. Higgs, INF2 is an endoplasmic

573  reticulum-associated formin protein. *Journal of cell science* **122**, 1430-1440 (2009).

574 36. F. Korobova, V. Ramabhadran, H. N. Higgs, An actin-dependent step in mitochondrial

575  fission mediated by the ER-associated formin INF2. *Science* **339**, 464-467 (2013).

576 37. W. K. Ji, R. Chakrabarti, X. Fan, L. Schoenfeld, S. Strack, H. N. Higgs, Receptor-

577  mediated Drp1 oligomerization on endoplasmic reticulum. *The Journal of cell biology*

578  **216**, 4123-4139 (2017).

579 38. A. A. Minin, A. V. Kulik, F. K. Gyoeva, Y. Li, G. Goshima, V. I. Gelfand, Regulation of

580  mitochondria distribution by RhoA and formins. *Journal of cell science* **119**, 659-670

581  (2006).

582 39. U. Manor, S. Bartholomew, G. Golani, E. Christenson, M. Kozlov, H. Higgs, J.

583  Spudich, J. Lippincott-Schwartz, A mitochondria-anchored isoform of the actin-

584  nucleating spire protein regulates mitochondrial division. *eLife* **4**, (2015).

585 40. G. B. Li, H. W. Zhang, R. Q. Fu, X. Y. Hu, L. Liu, Y. N. Li, Y. X. Liu, X. Liu, J. J. Hu, Q.

586  Deng, Q. S. Luo, R. Zhang, N. Gao, Mitochondrial fission and mitophagy depend on

587  cofilin-mediated actin depolymerization activity at the mitochondrial fission site.

588  *Oncogene* **37**, 1485-1502 (2018).

589 41. S. Li, S. Xu, B. A. Roelofs, L. Boyman, W. J. Lederer, H. Sesaki, M. Karbowski,

590  Transient assembly of F-actin on the outer mitochondrial membrane contributes to

591  mitochondrial fission. *The Journal of cell biology* **208**, 109-123 (2015).

593 **Figure legends**

594 **Figure 1. Expression and subcellular localization of FMNL2 during mouse oocyte**

meiosis. (A) Western blotting results of FMNL2 protein expression at different stages.

FMNL2 expressed at the GV, MI, and MII stages. **(B)** Subcellular localization of

FMNL2-EGFP and FMNL2 antibody during mouse oocyte meiosis. FMNL2 was

enriched at the cortex (GV, GVBD, MI and MII stage) and spindle periphery (MI

stage). Green, FMNL2-EGFP; blue, DNA. Negative control: Green, EGFP; blue,

DNA. Bar =20 μm. **(C)** Co-staining of oocytes for FMNL2 and actin. FMNL2 and

actin both localization in cortex. Green, FMNL2-antibody; red, actin; blue, DNA. Bar

=20 μm.

Figure 2. Knockdown of FMNL2 affects first polar body extrusion and

asymmetric division. (A) Western blot analysis for FMNL2 expression in the

FMNL2-KD group and control group. Relative intensity of FMNL2 and tubulin was

assessed by densitometry. **(B)** DIC images of control oocytes and FMNL2-KD

oocytes after 12 h culture. FMNL2-KD caused large polar bodies (black arrows) and

some oocytes failed to extrude the polar bodies (white arrows). **(C)** Rate of polar body

extrusion after 12 h culture of the control group and FMNL2-KD group. **(D)** Rate of

large polar body extrusion after 12 h culture in the control group and FMNL2-KD

group. **(E)** Time-lapse microscopy showed that polar body extrusion failed after

FMNL2-KD. Bar = 10 μm. **(F)** Western blot analysis for FMNL2 expression in the

control group, FMNL2-KD group and rescue group. Relative intensity of FMNL2 and

tubulin was assessed by densitometry. **(G)** DIC images of FMNL2-KD oocytes and

rescue oocytes after 12 h culture. **(H)** Rate of polar body extrusion after 12 h culture

of the FMNL2-KD group and rescue group. **(I)** Rate of large polar body extrusion

after 12 h culture in the FMNL2-KD group and rescue group. (**J**) Rate of polar body

extrusion after 12 h culture of the control group and FMNL2+3-KD group. (**K**) Rate

of large polar body extrusion after 12 h culture in the control group and FMNL2+3-

KD group. The data are presented as mean ± SEM from at least three independent

experiments. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Figure 3. Knockdown of FMNL2 disrupts spindle localization during mouse

oocyte meiosis. (**A**) Time-lapse microscopy showed that spindle migration failed after

FMNL2-KD. Green, tubulin-EGFP. Bar = 10 μm. (**B**) Representative images and the

proportion of spindle migration after 9.5 h of culture in the control group and

FMNL2-KD oocyte group. White, actin; green, tubulin; magenta, DNA. Bar = 10 μm.

(**C**) Representative images and the proportion of spindle migration after 9.5 h of

culture in the FMNL2-KD group and rescue oocyte group. magenta, DNA. Bar = 10

μm. The data are presented as mean ± SEM from at least three independent

experiments. * $P < 0.05$, ** $P < 0.01$.

Figure 4. Knockdown of FMNL2 disrupts actin assembly during mouse oocyte

meiosis. (**A, B**) Representative images of actin distribution at the oocyte cortex and

the fluorescent intensities in the control group and FMNL2-KD group ($P > 0.1$).

White, actin; green, tubulin; magenta, DNA. Bar = 10 μm. (**C, D**) Representative

images of actin distribution in the oocyte cytoplasm and the fluorescent intensities in

the control group and FMNL2-KD group. White, actin; green, tubulin; magenta, DNA.

Bar = 10 μm. (**E, F**) Representative images of actin distribution in the oocyte

cytoplasm and the fluorescent intensities in the FMNL2-KD group and rescue group.

White, actin; magenta, DNA. Bar = 10 μm. **(G)** Mass spectrometry results showed

that FMNL2 was related to many actin-related proteins. **(H)** Co-IP results showed that

FMNL2 was correlated with Arp and Formin2 but not with Profiling and Fascin. **(I)**

Arp2 protein expression significantly increased in the FMNL2-KD oocytes compared

with the control oocytes. Arp2 protein expression significantly decreased in the

rescue oocytes compared with the FMNL2-KD oocytes. **(J)** Formin2 protein

expression significantly decreased in the FMNL2-KD oocytes compared with the

control oocytes. Formin2 protein expression significantly increased in the rescue

oocytes compared with the FMNL2-KD oocytes. The data are presented as mean **±**

SEM from at least three independent experiments. * P < 0.05, ** P < 0.01, **** P <

0.0001.

Figure 5. FMNL2 regulates endoplasmic reticulum distribution during mouse

oocytes maturation. (A) Co-IP results showed that FMNL2 was correlated with INF2.

(B) Mass spectrometry results showed that FMNL2 was associated with ER-related

proteins. **(C)** Representative images of ER distribution in the oocyte cytoplasm in the

control group and FMNL2-KD group. In FMNL2-KD oocytes, ER agglomerated in

cytoplasm (white arrow). Red, ER; Blue, DNA. Bar = 10 μm. **(D)** Abnormal

distribution of ER significantly increased in the FMNL2-KD oocytes compared with

the control oocytes. **(E)** Grp78 and Chop protein expression significantly increased in

the FMNL2-KD oocytes compared with the control oocytes. The band intensity

analysis also confirmed this finding. **(F)** Representative images of ER distribution in

the oocyte cytoplasm in the FMNL2-KD group and rescue group. Red, ER; Blue,

DNA. Bar = 10 μm. **(G)** Abnormal distribution of ER significantly decreased in the

rescue oocytes compared with the FMNL2-KD oocytes. **(H)** Grp78 protein expression

significantly decreased in the rescue oocytes compared with the FMNL2-KD oocytes.

The band intensity analysis also confirmed this finding. The data are presented as

mean ± SEM from at least three independent experiments. * P < 0.05, ** P < 0.01.

Figure 6. FMNL2 regulates mitochondrial distribution during mouse oocytes

maturation. **(A)** Mass spectrometry results showed that FMNL2 was related to many

mitochondria-related proteins. **(B)** Representative images of mitochondrial

distribution in the oocyte cytoplasm in the control group and FMNL2-KD group. In

FMNL2-KD oocytes, mitochondrial agglomerated in cytoplasm (white arrow). Green,

Mito. Bar = 20 μm. **(C)** Abnormal distribution of mitochondrial significantly

increased in the FMNL2-KD oocytes compared with the control oocytes. **(D)**

Representative images of mitochondrial distribution in the oocyte cytoplasm in the

FMNL2-KD group and rescue group. In FMNL2-KD oocytes, mitochondrial

agglomerated in cytoplasm (white arrow). Red, Mito; Blue, DNA. Bar = 20 μm. **(E)**

Abnormal distribution of mitochondrial significantly decreased in the rescue oocytes

compared with the FMNL2-KD oocytes. **(F)** The typical picture for JC1 green

channel and red channel after FMNL2-KD. **(G)** The JC1 signal (red/green ratio) after

FMNL2-KD compare with the control group, the JC-1 red/green fluorescence ratio

was significantly reduced in FMNL2-KD groups. Bar = 20 μm. **(H)** cofilin protein

expression significantly decreased in the FMNL2-KD oocytes compared with the

control oocytes. The band intensity analysis also confirmed this finding. The data are

presented as mean ± SEM from at least three independent experiments. * P < 0.05, **

P < 0.01.

A

Relative intensity (FMNL2/Tubulin)

FMNL2

Tubulin

GV | MI | MII

B

DIC | FMNL2-EGFP | DNA | FMNL2-Ab

GV

GVBD

MI

MII

20μm

EGFP | DNA

Negative control

20μm

C

FMNL2 | Actin | DNA

GV

MI

A

| | 8h | 9h | 9.5h | 10h | 10.5h | 11h | 12h |

Control

FMNL2-KD

FMNL2-KD

DIC Tubulin

B

Control   FMNL2-KD

Actin

Tubulin
DNA

Rate of spindle migration

Percentage %

Control   FMNL2-KD

C

FMNL2-KD   Rescue

DIC

DNA

Rate of spindle migration

Percentage %

FMNL2-KD   Rescue

A Control   FMNL2-KD

Tubulin Actin DNA

B Cortex actin fluorescense intensity after cultured of 9 h

C Control   FMNL2-KD

Tubulin Actin DNA

D Cytoplasm actin fluorescense intensity after cultured of 9 h

****

E FMNL2-KD   Rescue

Actin

Actin DNA

F Cytoplasm actin fluorescense intensity after cultured of 9 h

****

G Actin-related proteins

Clf1 Actl6a
Smarce1 Tpm3 Arpc4 Capza1
Acta2 Fscn1 Arpc2 Capg Actr3
Vcl Actr1a Tpm1 Actn1 Arpc5l
Pls3 Ctnna1 Arpc2 Arpc5 Capza2 Dctn2
Ctnnb1 FMNL2 Nptn
Clf2 Ctnnd1 Actb
Inf2 Tpm2 Mylk Actg1 Dctn1
Pfn2 Serpinf2 Pfn1 Capzb Arfip1
Syne2 Actn4 Actr2

H Input IP:FMNL2

Arp2

Profilin

fascin

Formin2

I

Control FMNL2-KD
Arp2
Tubulin

FMNL2-KD Rescue
Arp2
Tubulin

J

Control FMNL2-KD
Formin2
Tubulin

FMNL2-KD Rescue
Formin2
Actin