FMNL2 regulates actin for ER and mitochondria distribution in oocyte meiosis

Meng-Hao Pan[1, 2], Zhen-Nan Pan[1], Ming-Hong Sun[1], Xiao-Han Li[1], Jia-Qian Ju[1], Shi-Ming Luo[2], Xiang-Hong Ou[2], Shao-Chen Sun[1*]

[1]College of Animal Science and Technology, Nanjing Agricultural University, Nanjing 210095, China.

[2]Fertility Preservation Lab, Reproductive Medicine Center, Guangdong Second Provincial General Hospital, Guangzhou, China.

Running Title: FMNL2 in mouse oocytes

[*]**Correspondence to:** Shao-Chen Sun, College of Animal Science and Technology, Nanjing Agricultural University, Nanjing 210095, China. E-mail: sunsc@njau.edu.cn

Funding: This work was supported by the National Natural Science Foundation of China (32170857); the National Key Research and Development Program of China (2021YFC2700100).

There is no conflict of interest to declare.

Abbreviations: FMNL2, Formin-like 2; FMNLs, Formin-likes; GV, germinal vesicle; GVBD, germinal vesicle breakdown; ATI, anaphase-telophase I; MI, metaphase I; MII, metaphase II; ER, Endoplasmic reticulum.

Abstract

During mammalian oocyte meiosis, spindle migration and asymmetric cytokinesis

are unique steps for the successful polar body extrusion. The asymmetry defects of

oocytes will lead to the failure of fertilization and embryo implantation. In present study

we reported that an actin nucleating factor formin-like 2 (FMNL2) played critical roles

in the regulation of spindle migration and organelle distribution. Our results showed

that FMNL2 mainly localized at the oocyte cortex and periphery of spindle. Depletion

of FMNL2 led to the failure of polar body extrusion and large polar bodies in oocytes.

Live-cell imaging revealed that the spindle failed to migrate to the oocyte cortex, which

caused polar body formation defects, and this might be due to the decreased

polymerization of cytoplasmic actin by FMNL2 depletion. Furthermore, mass

spectrometry analysis indicated that FMNL2 was associated with mitochondria and

endoplasmic reticulum-related proteins, and FMNL2 depletion disrupted the function

and distribution of mitochondria and endoplasmic reticulum, showing with decreased

mitochondrial membrane potential and the occurrence of endoplasmic reticulum stress.

Microinjecting Fmnl2-EGFP mRNA into FMNL2-depleted oocytes significantly

rescued these defects. Thus, our results indicate that FMNL2 is essential for the actin

assembly, which further involves into meiotic spindle migration and ER/mitochondria

functions in mouse oocytes.

Keywords: FMNL2; oocyte; actin; Endoplasmic reticulum; Mitochondria

Introduction

Mammalian oocyte maturation is an asymmetric division process that generates a

large egg and a small polar body. This asymmetry is critical for the following

fertilization and early embryo development. After germinal vesicle breakdown (GVBD),

the meiotic spindle is organized at the center of the oocyte, and then it migrates to the

oocyte cortex at the late metaphase I (MI). The oocytes are arrested at metaphase II

(MII) after the extrusion of first polar body (*1, 2*). Actin filaments, as the most widely

distributed cytoskeleton in cells, regulate various dynamic events during oocyte meiotic

maturation (*3*), and two key events are the spindle migration and cortical reorganization

in mammalian oocytes (*1, 4, 5*). Small GTPases and actin nucleation factors are shown

to promote the assembly and function of actin. The actin nucleation factors are the

molecules that directly promote the actin assembly: Arp2/3 complex control the

assembly of branched actin, and formin family member Formin2 (FMN2) and Spire1/2

control the assembly of linear actin. These proteins are all proposed to play a role in

actin-related spindle migration and cytokinesis during mammalian oocyte maturation

(*6-8*). The cortex protein Arp2/3 complex nucleates the actin to produce a

hydrodynamic force to move the spindle toward the cortex, and regulates cytokinesis

during oocyte maturation (*1, 8*). FMN2 and Spire1/2 nucleates actin around the spindle

in the cytoplasm to give the meiotic spindle an initial power for migration (*7, 9*).

Besides Formin2, the DRFs (diaphanous-related formins) subfamily in the formin

family has been extensively studied. The DRFs family consists of mDia, Daam, FHOD

and FMNLs (*10*). The "Formin-like" proteins (FMNLs) subfamily includes FMNL1

(FRL1), FMNL2 (FRL3), and FMNL3 (FRL2). Like other Formin family proteins,

FMNLs play important roles in cell migration, cell division, and cell polarity (*10, 11*).

While FMNL2 is widely expressed in multiple human tissues, especially in the

gastrointestinal and mammary epithelia, lymphatic tissues, placenta, and reproductive

tract (*12*). As an important actin assembly factor, FMNL2 accelerates the elongation of

actin filaments branched by Arp2/3 complex (*13*). In invasive cells, FMNL2 is mainly

localized in the leading edge of the cell, lamellipodia and filopodia tips, to improve cell

migration ability by actin-based manner (*13-15*). FMNL2 is also involved in the

maintenance of epithelial-mesenchymal transition (EMT) in human colorectal

carcinoma cell (*16*). Besides its roles on the actin assembly, emerging evidences

indicate that FMNL2 may interact with organelle dynamics. It is shown that FMNL2 is

related with the Golgi apparatus, since the absence of FMNL2/3 can cause the Golgi

fragmentation (*17*). However, till now the roles of FMNLs especially FMNL2 on oocyte

meiosis are still largely unknown.

In the present study, we disturbed the FMNL2 expression and explored the roles of

FMNL2 during mouse oocyte meiosis. Our results showed that FMNL2 was essential

for the polar body size control and successful extrusion; and these abnormal phenotypes

might be due to aberrant actin-based meiotic spindle migration. Meanwhile, we also

found that FMNL2 was essential for the functions and distribution of mitochondria and

endoplasmic reticulum. Therefore, this study provided the evidence for the critical roles

of FMNL2-mediated actin on spindle movement and organelle dynamics in mammalian

oocytes.

Materials and Methods

Antibodies and chemicals

Rabbit monoclonal anti-FMNL2 antibody, rabbit monoclonal anti-Arp2 antibody, mouse monoclonal anti-profilin1 antibody were from Santa Cruz (Santa Cruz, CA, USA). Rabbit monoclonal anti-Fascin antibody was purchased form Abcam (Cambridge, UK). Rabbit polyclonal anti-INF2 antibody was purchased from Proteintech (Proteintech, CHI, USA). Rabbit monoclonal anti-α-tubulin (11H10) antibody, rabbit monoclonal anti-Grp78 antibody, rabbit monoclonal anti-cofilin antibody and rabbit monoclonal anti-Chop antibody were from Cell Signaling Technology (Beverly, MA, USA). Mouse monoclonal anti-α-tubulin-FITC antibody was from Sigma-Aldrich Corp. (St. Louis, MO, USA). FITC-conjugated goat anti-rabbit IgG were from Zhongshan Golden Bridge Biotechnology (Beijing). ER-Tracker Red and Mito-Tracker Green were from Beyotime Biotechnology (Shanghai). All other chemicals and reagents were from Sigma-Aldrich Corp., unless otherwise stated.

Ethics statement and oocyte culture

We followed the guidelines of Animal Research Institute Committee of Nanjing Agricultural University to conduct the operations. The animal facility had license authorized by the experimental animal committee of Jiangsu Province (SYXK-Su-20170007). These mice were housed in considerably ideal conditions which consisted of controlled temperature, regular diet and appropriate light. Female mice (4 - 6 weeks) were used to collect germinal vesicle oocytes. The oocytes were placed at 37°C with an atmosphere of 5% CO_2, and cultured to different time points for immunostaining,

111   microinjection and western blot.

Plasmid construct and in *vitro* transcription

Template RNA was generated from mouse ovaries with RNA Isolation Kit

(Thermos fisher), then we reversed transcription of these RNA to create cDNA by a

PrimeScript 1st strand cDNA synthesis kit (Takara, Japan). Fmnl2-EGFP vector was

generated by Wuhan GeneCreate Biological Engineering Co, Ltd. mRNA was

synthesized from linearized plasmid using HiScribe T7 high yield RNA synthesis kit

(NEB), then capped with m7G(5')ppp(5')G (NEB) and tailed with a poly(A)

polymerase tailing kit (Epicentre) and purified with RNA clean & concentrator-25 kit

(Zymo Research).

Microinjection of Fmnl2 siRNA and mRNA

Fmnl2 siRNA microinjection was used to knock down endogenous Fmnl2 in

mouse oocytes. Fmnl2 siRNA 5′- GCU GAA UGC UAU GAA CCU ATT-3′, 5′- GCC

AUU GAU CUU UCU UCA ATT-3′, 5′- GGA AUU AAG AAG GCG ACA ATT-3′,

(Genepharma, Shanghai, China) were diluted with DEPC water to give a 20μM stock

solution respectively, and the three siRNA were mixed in equal proportions before

microinjection. The control group was microinjected with negative control siRNA 5′-

UUC UCC GAA CGU GUC ACG UTT-3′. After injection, the oocytes were cultured

in M16 medium containing 5μM milrinone for 18-20 h, and then washed five times (2

min each) in fresh M2 medium. We transferred the oocytes to fresh M16 medium, and

cultured for following experiments. When microinjecting the α-tubulin-EGFP mRNA

and rescue experiments, GV oocytes need to be cultured in M16 medium with 5μM

milrinone for 2h. DNase/RNase-free water microinjected as the control.

Immunofluorescent staining and confocal microscopy

Oocytes were fixed in 4% paraformaldehyde (in PBS) for 30 min and

permeabilized with 0.5% Triton X-100 in PBS for 20min then blocked in blocking

buffer (1% BSA-supplemented PBS) at room for 1 h. For FMNL2 staining, the oocytes

after blocking were incubated with Rabbit monoclonal anti-FMNL2 antibody (1:100)

at 4 °C overnight, then oocytes were washed by wash buffer (0.1% Tween 20 and 0.01%

Triton X-100 in PBS) for 3 times (5 min each time). Next the oocytes were labeled with

secondary antibody coupled to FITC-conjugated goat anti-rabbit IgG (1:100) at room

temperature for 1 h. For α-tubulin staining, oocytes were incubated with anti-α-tubulin-

FITC antibody (1:200). For actin staining, oocytes were incubated with Phalloidin-

TRITC at room temperature for 2 h. Then the oocytes were washed as the same way.

Finally, oocytes were incubated with Hoechst 33342 at room temperature for 10-20 min.

After staining, samples were mounted on glass slides and observed with a confocal

laser-scanning microscope (Zeiss LSM 800 META, Germany).

ER and Mito-tracker staining

145 To study ER and mitochondria distribution during mouse oocyte meiosis, MI stage

146 oocytes were incubated with ER-Tracker Red (1:3000) or 200 nM Mito-tracker green

147 (Red) in M16 medium for 30 min at 37ºC and 5% CO_2. Then the oocytes were washed

148 three times with M2 medium, finally the samples were examined with confocal

149 microscopy.

Time lapse microscopy

To image the dynamic changes that occurred during oocyte maturation, oocytes were cultured in M16 medium, then transferred to the Leica SD AF confocal imaging system equipped with 37 °C incubator and 5% CO2 supply (H301-K-FRAME). The spindle in oocytes was labeled by α-tubulin-EGFP.

Immunoprecipitation

4-6 ovaries were put into RIPA Lysis Buffer contained phosphatase inhibitor cocktail (100×) (Kangwei Biotechnology, China), and were completely cleaved on ice block. We collected supernatant after centrifugation (13200 rps, 20 min) and then took out 50 μl as input sample at 4 °C. The rest of the supernatant was incubated with primary antibody (FMNL2 or INF2 antibody) overnight at 4 °C. 30 μl conjugated beads (washed five times in PBS) were added to the supernatant/antibody mixture and incubated at 4 °C for 4-6 h, after three times wash by immune complexes, the samples were then released from the beads by mixing in 2× SDS loading buffer for 10min at 30 °C.

Western blot analysis

Approximate 100-150 mouse oocytes were placed in Laemmli sample buffer and heated at 85°C for 7-10 min. Proteins were separated by SDS-PAGE at 165V for 70-80 min and then electrophoretically transferred to polyvinylidene fluoride (PVDF) membranes (Millipore, Billerica, MA, USA) at 20 V for 1 hour. After transfer, the membranes were then blocked with TBST (TBS containing 0.1% Tween 20) containing 5% non-fat milk at room temperature for 90 min. After blocking, the membranes were incubated with rabbit monoclonal anti-FMNL2 antibody (1:500), rabbit monoclonal anti-Arp2 antibody (1:500), mouse monoclonal anti-profilin1 antibody (1:500), rabbit

monoclonal anti-Fascin antibody (1:5000), rabbit polyclonal anti-INF2 antibody

(1:500), rabbit monoclonal anti-Grp78 antibody (1:1000), rabbit monoclonal anti-

cofilin antibody (1:2000), rabbit monoclonal anti-CHOP antibody (1:1000), or rabbit

monoclonal anti-tubulin antibody (1:2000) at 4 °C overnight. After washing 5 times in

TBST (5 min each), membranes were incubated for 1h at room temperature with HRP-

conjugated Pierce Goat anti-Rabbit IgG (1:5000) or HRP-conjugated Pierce Goat anti-

mouse IgG (1:5000). After washing for 5 times, the membranes were visualized using

chemiluminescence reagent (Millipore, Billerica, MA). Every experiment repeated at

least 3 times with different samples.

Intensity analysis

To analyze the fluorescence intensity of actin filaments, the control group and

treated group were mounted on the same glass slide and tested with same parameters.

Image J was used to determine the average fluorescence intensity per unit area within

the region of interest (ROI). The independent measures were taken for the cell cortex

and cytoplasm. For quantification of the western blot results, the band intensity was

measured by Image J.

Statistical analysis

At least three biological replicates were performed for each analysis. The results

were endowed as means ± SEM. All analyses were performed using GraphPad

Prism7.00 software (GraphPad, CA, USA). Results of P<0.05 were considered

statistically significant (differences P<0.05 denoted by *, P<0.01 denoted by **,

P<0.001 denoted by *** and P<0.0001 denoted by ****).

Results

Expression and subcellular localization of FMNL2 during mouse oocyte maturation

We first examined FMNL2 expression in mouse oocytes at different stages. The results indicated that FMNL2 all expressed in GV, MI and MII stages during mouse oocyte maturation (GV, 1; MI, 0.82 ± 0.07; MII, 0.61 ± 0.10, Figure 1A). Next, we performed Fmnl2-EGFP mRNA microinjection to examine the localization of FMNL2. As shown in Figure 1B, FMNL2 accumulated at the oocyte cortex during the GV, GVBD and MI stages. Besides, FMNL2 also localized at the spindle periphery during GVBD and MI stages. At anaphase-telophase I stage (ATI), FMNL2 was mainly at the midbody. The FMNL2 antibody staining results also confirmed this localization pattern. We also co-stained FMNL2 antibody with actin, and the result showed that the signals of FMNL2 and actin overlapped at the cortex in oocytes (Figure 1C). The FMNL2 localization pattern indicated that FMNL2 might interact with actin dynamics during oocyte meiosis.

FMNL2 is essential for polar body extrusion and asymmetric division in mouse oocytes

To investigate the functional roles of FMNL2 in mouse oocytes, we employed Fmnl2 siRNA microinjection to knockdown FMNL2 protein expression. A significant decrease of FMNL2 protein level was shown in FMNL2-KD oocytes compared to control group by western blot (1 vs. 0.48 ± 0.08, $P < 0.01$, Figure 2A). We then examined first polar body extrusion, and the results showed that deleting FMNL2

disturbed first polar body extrusion, while a large proportion of oocytes showed big

polar bodies among the oocytes which extruded polar bodies (Figure 2B). The

quantitative results also confirmed this phenotype (rate of polar body extrusion: 74.26

± 1.44%, n = 439 vs. 59.5 ± 2.82%, n = 398, P < 0.001, Figure 2C; rate of large polar

bodies: 19.05 ± 1.97%, n = 311 vs. 37.16 ± 1.87%, n = 257, P < 0.0001, Figure 2D). In

addition, live-cell imaging was used to determine the dynamic changes that occurred

during oocyte maturation, and the results showed that the oocytes either failed to

undergo cytokinesis or divided from the central axis of the oocytes and formed big polar

bodies (Figure 2E). To further confirm the phenotype, we performed FMNL2 rescue

experiments by expressing exogenous Fmnl2 mRNA in FMNL2-depleted oocytes

(Figure 2F), we found that exogenous Fmnl2 mRNA expression rescued first polar body

extrusion and large polar body defects (Figure 2G). The quantitative results also

confirmed this phenotype (rate of polar body extrusion: 48.34 ± 4.2%, n = 355 vs. 62.62

± 3.6%, n = 377, P < 0.01, Figure 2H; rate of large polar bodies: 30.93 ± 2%, n = 193

vs. 9.58 ± 2.4%, n = 203, P < 0.01, Figure 2I). To testify whether the functions of

FMNL2 were associated with other FMNLs, we also compared the polar body extrusion

with the knock down of both FMNL2 and FMNL3, and the results showed that double

knock down of FMNL2 and FMNL3 did not cause a severe polar body extrusion defects

compared with the single knock down of FMNL2 (polar body extrusion, Control: 70.97

± 1.23%, n=261 vs FMNL2+3-KD: 60.42 ± 2.99%, n=198, P < 0.05, Figure 2J. Large

polar body, Control: 10.85 ± 0.97%, n=172 vs FMNL2+3-KD: 32.90 ± 1.88%, n=118,

P < 0.001, Figure 2K). These results suggested that FMNL2 played critical roles for the

229 polar body extrusion and asymmetric division during mouse oocyte maturation.

FMNL2 regulates spindle migration during mouse oocyte maturation

To investigate the causes for polar body extrusion defects, we examined the

spindle migration by time-lapse microscopy during oocyte meiosis. As shown in Figure

4A, in the control oocyte, the meiotic spindle formed in the center of the oocyte after

culture 8 h and moved to the oocyte cortex at 9.5h; and the polar body was extruded at

11-12h, with a spindle formed near the cortex at MII stage. However, in FMNL2-KD

oocytes, two phenotypes were observed: 1) the meiotic spindle remained in the center

of the oocyte until 10 h, and then the oocytes initiated the cytokinesis at 10.5 h but

failed to extrude the polar body; 2) some oocytes with arrested spindles initiated the

cytokinesis but extruded a big polar body (Figure 3A). This indicated the failure of

spindle migration after FMNL2 depletion. We analyzed the rate of cortex-localized

spindle in oocytes by cultured for 9.5 h, and the result showed that the rate of migrated

spindles in control oocytes was significantly higher than that in FMNL2-KD oocytes

$(59.94 \pm 3.42\%$, n = 78 vs. $38.97 \pm 6.34\%$, n = 64, P < 0.05, Figure 3B). We also

performed FMNL2 rescue experiments. Supplementing with exogenous Fmnl2 rescued

the spindle migration defects compared with the Fmnl2-depletion group $(40.27 \pm 3.19\%$,

n = 181 vs. $57.01 \pm 2.72\%$, n = 57, P < 0.01, Figure 3C). These results suggested that

FMNL2 might be involved in spindle migration in mouse oocytes.

FMNL2 promotes cytoplasmic actin assembly during mouse oocyte maturation

247   As FMNL2 is a key actin assembly factor, we further investigated actin assembly

248 after deleting FMNL2 in mouse oocytes. Surprisingly there was no significant

difference for the signals of cortex actin was observed between control oocytes and FMNL2-KD oocytes, which was confirmed by the fluorescence intensity analysis (30.88 ± 1.10, n = 28 vs. 30.58 ± 1.12, n = 28, P > 0.05, Figure 4A). However, we found a significant decrease of cytoplasmic actin signals in the FMNL2-KD oocytes, and the statistical analysis for the cytoplasmic actin fluorescent signals also confirmed our findings (58.25 ± 2.05, n = 26 vs. 37.92 ± 2.02, n = 24, P < 0.0001, Figure 4B). Moreover, the rescue experiments showed that exogenous Fmnl2 rescued the decrease of cytoplasmic actin filaments compared with the Fmnl2-depletion group (37.98 ± 1.98, n = 16 vs. 54.72 ± 2.88, n = 15, P < 0.0001, Figure 3C). We next explored how FMNL2 regulates cytoplasmic actin assembly in oocytes. By mass spectrometry analysis we found there were several actin-related potential candidates which might be related with FMNL2 (Figure 4D). Co-immunoprecipitation results showed that FMNL2 precipitated Arp2 and Formin2 but not Profilin and fascin (Figure 4E). To further verify the correlation between FMNL2 and Arp2 and Formin2, we then examined Arp2 and Formin2 protein expression after FMNL2 knockdown. The results showed Arp2 protein expression increased significantly after FMNL2 knockdown (1 vs. 1.56 ± 0.07, P < 0.001, Figure 4F) but decreased after FMNL2 knockdown (1 vs. 0.62 ± 0.04, P < 0.001, Figure 4G). Exogenous Fmnl2 rescued these alterations compared with that in the FMNL2-KD group (Arp2 protein expression: 1 vs. 0.65 ± 0.06, P < 0.01, Figure 4F; Formin2 protein expression: 1 vs. 1.24 ± 0.05, P < 0.01, Figure 4G). These results indicated that FMNL2 associated with Formin2 and Arp2 for actin assembly in mouse oocytes.

FMNL2 regulates endoplasmic reticulum distribution during mouse oocyte maturation

The mass spectrometry analysis data indicated that several ER-related potential

candidates which might be related with FMNL2 (Figure 5A), while INF2, a typical

protein which mediates actin polymerization at ER showed high confidence level. We

then examined the relationship between FMNL2 and INF2, and the co-

immunoprecipitation results showed that FMNL2 precipitated INF2 and INF2 also

precipitated FMNL2 (Figure 5B), indicating that FMNL2 interacted with INF2 in

mouse oocytes. We then examined the ER distribution in FMNL2-KD oocytes. As

shown in Figure 5C, in control oocytes the ER evenly distributed in the cytoplasm and

accumulated at the spindle periphery in MI stage; however, ER agglomerated in

cytoplasm in FMNL2-KD oocytes (Figure 5C). The statistical analysis showed that the

abnormal distribution of ER increased significantly in the FMNL2-KD group (28.91 $\pm$

5.62, n = 27 vs. 59.64 $\pm$ 6.95, n = 28, P < 0.05, Figure 5D). The localization pattern of

ER indicated its functions might be disturbed. In FMNL2-KD oocytes, we found the

expressions of ER-stress related proteins Grp78 and Chop were significantly increased

(Grp78: 1 vs. 1.42 $\pm$ 0.12, P < 0.05; Chop: 1 vs. 1.53 $\pm$ 0.16, P < 0.05. Figure 5E),

indicating the occurrence of ER stress. We also performed FMNL2 rescue experiments.

Supplementing with exogenous Fmnl2 rescued the ER distribution defects caused by

FMNL2 knockdown (Figure 5F), which was supported by the statistical analysis

showing that the abnormal distribution rate of ER decreased significantly in the rescue

group (52.04 $\pm$ 5.29, n = 70 vs. 34.91 $\pm$ 3.37, n = 78, P < 0.05, Figure 5G). Moreover,

Grp78 protein expression decreased in the rescue group (1 vs. 0.78 ± 0.05, $P < 0.01$.

Figure 5H). These results indicated that the depletion of FMNL2 affected ER

distribution and caused ER stress in mouse oocytes.

FMNL2 regulates mitochondrial distribution during mouse oocyte maturation

As INF2 is also related to the mitochondrial connection of ER, we further screened

up the mass spectrometry analysis data and we found many mitochondria-related

potential candidates which might be related with FMNL2 (Figure 6A). Therefore, we

further examined the distribution of mitochondria in FMNL2-KD oocytes. In control

oocytes, the mitochondria evenly distributed in the cytoplasm and accumulated at the

spindle periphery in MI stage; however, in FMNL2-KD oocytes, mitochondria

presented clumped aggregation distribution in cytoplasm (Figure 6B). We counted the

number of clumps and found that the uniform distribution of mitochondria decreased

significantly in the FMNL2-KD group (59.66 ± 8.48, n = 31 vs. 20.83 ± 4.17, n = 32, P

< 0.05, Figure 6C). A large number of FMNL2-KD oocytes agglomerated into one to

three clumps (22.73 ± 4.27, n = 31 vs. 42.50 ± 1.25, n = 32, P < 0.05, Figure 6C).

Supplementing with exogenous Fmnl2 rescued the mitochondria distribution (Figure

6D), the statistical analysis showed that the uniform distribution of mitochondria

increased significantly in the rescue group (36.49 ± 3.97, n = 53 vs. 53.90 ± 2.09, n =

79, P < 0.05, Figure 6E). We also examined mitochondrial membrane potential, and the

results showed that FMNL2 depletion caused the alterations of mitochondrial

membrane potential (MMP) by JC-1 staining. The fluorescence intensity of JC-1 red

channel was decreased compared with the control group (Figure 6F). We also calculated

the ratio for red/green fluorescence intensity, and the results also confirmed this (control

group: 0.40 vs. FMNL2-KD: 0.21 ± 0.01, $P < 0.01$) (Figure 6G). Cofilin is an important

factor of actin assembly and regulates mitochondrial function. We also examined cofilin

protein expression after FMNL2 knockdown. The results showed cofilin protein

expression decreased significantly after FMNL2 knockdown (1 vs. 0.81 ± 0.03, $P <$

0.01, Figure 6H). These results indicated that FMNL2 regulated mitochondria

distribution and function during mouse oocyte maturation.

Discussion

In this study, we explored the functions of FMNL2 during mouse oocyte meiosis.

Our results indicated that FMNL2 regulated actin-based spindle migration for

asymmetric cell division of oocytes, and more importantly FMNL2 was critical for

maintaining the distribution of the ER and mitochondria, which set up a link for actin-

related spindle migration and organelle dynamics in oocytes.

As a subfamily of Formin family, FMNLs play an important role in regulating actin

filaments (*18*), while FMNL2 is most widely expressed in variety of cell models among

the members of FMNLs. In this study, we showed that FMNL2 expressed in mouse

oocytes and it mainly accumulated at the oocyte cortex and spindle periphery, which

was similar with the actin distribution pattern in oocytes. This specific localization is

also similar to FMN2, a well-studied factor in the formin family for spindle migration

during oocyte meiosis (*7, 19*). In addition, another FMNLs family member, FMNL1 is

also localized at the cortex and is essential for actin polymerization and spindle

assembly during oocyte meiosis (*20*). Based on the localization pattern of FMNL2, we

speculated that the functions of FMNL2 might be also involved in actin-related process

during mouse oocyte meiosis.

To confirm our hypothesis, we depleted FMNL2 protein expression and we found

that absence of FMNL2 caused the aberrant first polar body extrusion. The oocytes

either failed to form the polar body or extruded large polar bodies. These phenotypes

caused by FMNL2 depletion are similar to the other actin-related proteins during oocyte

maturation such as Arp2/3 complex (*8, 21*) and FMN2 (*22, 23*). We next examined the

actin distribution in oocytes since it is reported that FMNL2 promotes actin filament

assembly in many models. FMNL2 is required for cell-cell adhesion formation by

regulating the actin assembly (*24*), and FMNL2 could directly drives actin elongation

(*15*). In CRC cells, cortactin bind to FMNL2 to active the actin polymerization, and

FMNL2 is important for invadopodia formation and functions (*25*). Our results showed

that the FMNL2 depletion caused significantly decrease in cytoplasmic actin, indicating

the conserved roles of FMNL2 on actin assembly in mammalian oocyte model. Other

Formin family proteins such as Daam1, FHOD1, and Formin-homology family protein

mDia1 are also reported to affect oocyte meiosis by regulating actin polymerization

(*26-28*).

We then tried to explore how FMNL2 involves into the actin assembly in oocytes.

Mass spectrometry analysis data indicated that FMNL2 associated with several actin-

related proteins, and we found that FMNL2 was associated with Arp2 and Formin2.

This could be confirmed by the altered expression of these two molecules after FMNL2

depletion. Therefore, we speculated FMNL2 could regulate cytoplasmic actin assembly

in oocytes through the association with Formin2 since it is reported to be an important

protein for cytoplasmic actin assembly in oocytes (*22*). Interestingly, our results showed

that unlike the reduction of cytoplasmic actin, cortex actin was not affected by the

absence of FMNL2. We speculated that Arp2/3 complex has a compensation effect on

the depleting of FMNL2 during oocyte meiosis, ensuring the cortex actin assembly in

oocytes since Arp2 protein expression significantly increased after FMNL2 depletion.

As an actin nucleator Arp2/3 complex localizes at the cortex and is essential for actin

polymerization during oocyte meiosis (*8, 29*). These results suggested that FMNL2

might be involved in cytokinesis and asymmetric division by regulating actin assembly

during mouse oocyte maturation.

The spindle migration is a key step in ensuring the asymmetric division for oocytes

(*30*). In mitosis, spindle position is decided by cortical actin and astral microtubules; in

contrast, spindle migration is mainly mediated by actin filaments during oocyte meiosis

(*30, 31*). Due to the effects of FMNL2 on asymmetric division and cytoplasmic actin,

we analyzed the spindle positioning at late MI, we found that the spindle migration was

disturbed after FMNL2 depletion, no matter the cytokinesis occurred or not. Several

formin proteins are shown to regulate spindle migration during oocytes meiosis. For

example, FMN2 nucleates actin surrounding the spindle, pushing force generated by

actin to trigger the spindle migration (*19, 22*), and cyclin-dependent kinase 1 (Cdk1)

induces cytoplasmic Formin-mediated F-actin polymerization to propel the spindle into

the cortex (*32*). Our previous studies also showed that absence of the formin family

member FMNL1 or FHOD1 could lead to the decrease of cytoplasmic actin to prevent

the spindle migration (*20, 27*). We speculated that FMNL2 together with other Formin

proteins, conservatively regulate actin-mediated spindle migration during oocyte

meiosis.

Another important finding is that through the mass spectrometry analysis we found

many candidate proteins which were related with ER, and our results indicated that

FMNL2 was essential for the maintenance of ER distribution in the cytoplasm.

Moreover, the loss of FMNL2 induced ER stress, showing with altered expression of

GRP78 and CHOP. Proper distribution of ER is important for the oocyte quality. ER

displays a homogeneous distribution pattern throughout the entire ooplasm during

development of oocytes and embryos from diabetic mice(*33*). During the transition of

mouse oocytes from MI to MII phase, actin regulates cortical ER aggregation(*34*). In

addition, Formin2 is shown to colocalize with the ER during oocyte meiosis and the

ER-associated Formin2 at the spindle periphery is required for MI chromosome

migration(*6*). In our results we showed that FMNL2 associated with INF2 protein in

oocytes. INF2 is an ER-associated protein, and the expression of GFP-INF2 which

containing DAD/WH2 mutations causes the ER to collapse around the nucleus (*35*).

We concluded that FMNL2 might regulate INF2 for the distribution of ER in cytoplasm

of oocytes.

Besides its roles of ER distribution, it is shown that INF2 also affects

mitochondrial length and ER-mitochondrial interaction in an actin-dependent manner

(*35, 36*). It is shown that INF2 regulates Drp1 for mitochondrial fission, and INF2-

induced actin filaments may drive initial mitochondrial constriction, which allows

Drp1-driven secondary constriction(*36, 37*). In addition, we also found many candidate

proteins which were related with mitochondria from mass spectrometry analysis.

During oocyte meiosis, mitochondria gradually accumulated around the spindle after

GVBD, and the spindle-peripheral FMN2 and its actin nucleation activity are important

for the accumulation of mitochondria in this region (*19*). Our results found that FMNL2

depletion caused agglutination of mitochondria and altered MMP level in the cytoplasm,

indicating its roles on the mitochondria distribution and functions. Another formin

protein mDia1 is shown to be necessary to induce the anchoring of mitochondria along

the cytoskeletal in mammalian CV-1 cells and Drosophila BG2-C2 neuronal cells(*38*).

Moreover, the formin interaction protein Spire1C binds INF2 to promote actin assembly

on mitochondrial surfaces, and Spire1C disruption could reduce mitochondrial

constriction and division(*39*). In addition, our result indicated that cofilin expression

decreased in FMNL2 depletion oocytes. Cofilin is an actin-depolymerizing factor and

its localization at the mitochondrial fission site is crucial for inducing mitochondrial

fission and mitophagy (*40*). Depleting of cofilin resulted in abnormal interconnection

and elongation of mitochondria (*41*). Together with its roles on ER, these data indicated

that FMNL2 might associate with INF2 and cofilin for the actin-based organelle

distribution during oocyte meiosis.

Collectively, we provide a body of evidence showing that FMNL2 associates with

Formin2 and Arp2/3 complex for actin assembly, which further regulates spindle

migration and INF2/Cofilin-related organelle dynamics during mouse oocyte

maturation.

Data Availability

All data generated or analyzed during this study are included in this published article

Acknowledgement

We are particularly grateful to Xiao-Yan Fan and Xing-Hua Wang from Fertility Preservation Laboratory, Reproductive Medicine Center, Guangdong Second Provincial General Hospital for their technical assistance of live cell imaging system.

Contributions

MHP and SCS designed the study. MHP performed the majority of the experiments. SML, ZNP, MHS, XHL, JQJ, YZ contributed to the regents and materials. MHP, XHO and SCS analyzed the data. MHP and SCS wrote the manuscript.

Competing interests

There is no conflict of interest to declare.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

References

1. K. Yi, B. Rubinstein, R. Li, Symmetry breaking and polarity establishment during mouse oocyte maturation. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **368**, 20130002 (2013).

2. B. Pan, J. Li, The art of oocyte meiotic arrest regulation. *Reproductive biology and*
endocrinology : RB&E **17**, 8 (2019).

3. Q. Y. Sun, H. Schatten, Regulation of dynamic events by microfilaments during oocyte
maturation and fertilization. *Reproduction* **131**, 193-205 (2006).

4. S. C. Sun, N. H. Kim, Molecular mechanisms of asymmetric division in oocytes.
Microscopy and microanalysis : the official journal of Microscopy Society of America,
Microbeam Analysis Society, Microscopical Society of Canada **19**, 883-897 (2013).

5. X. Duan, S. C. Sun, Actin cytoskeleton dynamics in mammalian oocyte meiosis. *Biol*
Reprod **100**, 15-24 (2019).

6. K. Yi, B. Rubinstein, J. R. Unruh, F. Guo, B. D. Slaughter, R. Li, Sequential actin-based
pushing forces drive meiosis I chromosome migration and symmetry breaking in
oocytes. *The Journal of cell biology* **200**, 567-576 (2013).

7. H. Li, F. Guo, B. Rubinstein, R. Li, Actin-driven chromosomal motility leads to symmetry
breaking in mammalian meiotic oocytes. *Nature cell biology* **10**, 1301-1308 (2008).

8. S. C. Sun, Z. B. Wang, Y. N. Xu, S. E. Lee, X. S. Cui, N. H. Kim, Arp2/3 complex regulates
asymmetric division and cytokinesis in mouse oocytes. *PloS one* **6**, e18392 (2011).

9. S. Pfender, V. Kuznetsov, S. Pleiser, E. Kerkhoff, M. Schuh, Spire-type actin nucleators
cooperate with Formin-2 to drive asymmetric oocyte division. *Current biology : CB* **21**,
955-960 (2011).

10. S. Kuhn, M. Geyer, Formins as effector proteins of Rho GTPases. *Small GTPases* **5**,
e29513 (2014).

11. M. Katoh, M. Katoh, Identification and characterization of human FMNL1, FMNL2 and
FMNL3 genes in silico. *International journal of oncology* **22**, 1161-1168 (2003).

12. M. Gardberg, K. Talvinen, K. Kaipio, K. Iljin, C. Kampf, M. Uhlen, O. Carpen,
Characterization of Diaphanous-related formin FMNL2 in human tissues. *BMC cell*
biology **11**, 55 (2010).

13. F. Kage, M. Winterhoff, V. Dimchev, J. Mueller, T. Thalheim, A. Freise, S. Bruhmann, J.
Kollasser, J. Block, G. Dimchev, M. Geyer, H. J. Schnittler, C. Brakebusch, T. E. Stradal, M.
F. Carlier, M. Sixt, J. Kas, J. Faix, K. Rottner, FMNL formins boost lamellipodial force
generation. *Nature communications* **8**, 14832 (2017).

14. X. L. Zhu, Y. F. Zeng, J. Guan, Y. F. Li, Y. J. Deng, X. W. Bian, Y. Q. Ding, L. Liang, FMNL2 is
a positive regulator of cell motility and metastasis in colorectal carcinoma. *The Journal*
of pathology **224**, 377-388 (2011).

15. J. Block, D. Breitsprecher, S. Kuhn, M. Winterhoff, F. Kage, R. Geffers, P. Duwe, J. L. Rohn,
B. Baum, C. Brakebusch, M. Geyer, T. E. Stradal, J. Faix, K. Rottner, FMNL2 drives actin-
based protrusion and migration downstream of Cdc42. *Current biology : CB* **22**, 1005-
1012 (2012).

16. Y. Li, X. Zhu, Y. Zeng, J. Wang, X. Zhang, Y. Q. Ding, L. Liang, FMNL2 enhances invasion
of colorectal carcinoma by inducing epithelial-mesenchymal transition. *Molecular cancer*
research : MCR **8**, 1579-1590 (2010).

17. F. Kage, A. Steffen, A. Ellinger, C. Ranftler, C. Gehre, C. Brakebusch, M. Pavelka, T. Stradal,
K. Rottner, Author Correction: FMNL2 and -3 regulate Golgi architecture and
anterograde transport downstream of Cdc42. *Scientific reports* **9**, 18008 (2019).

18. D. Breitsprecher, B. L. Goode, Formins at a glance. *Journal of cell science* **126**, 1-7

488   (2013).

489   19.   X. Duan, Y. Li, K. Yi, F. Guo, H. Wang, P. H. Wu, J. Yang, D. B. Mair, E. A. Morales, P. Kalab,
490   D. Wirtz, S. X. Sun, R. Li, Dynamic organelle distribution initiates actin-based spindle
491   migration in mouse oocytes. *Nature communications* **11**, 277 (2020).

492   20.   F. Wang, L. Zhang, X. Duan, G. L. Zhang, Z. B. Wang, Q. Wang, B. Xiong, S. C. Sun,
493   RhoA-mediated FMNL1 regulates GM130 for actin assembly and phosphorylates MAPK
494   for spindle formation in mouse oocyte meiosis. *Cell cycle* **14**, 2835-2843 (2015).

495   21.   K. Yi, J. R. Unruh, M. Deng, B. D. Slaughter, B. Rubinstein, R. Li, Dynamic maintenance of
496   asymmetric meiotic spindle position through Arp2/3-complex-driven cytoplasmic
497   streaming in mouse oocytes. *Nature cell biology* **13**, 1252-1258 (2011).

498   22.   J. Dumont, K. Million, K. Sunderland, P. Rassinier, H. Lim, B. Leader, M. H. Verlhac,
499   Formin-2 is required for spindle migration and for the late steps of cytokinesis in mouse
500   oocytes. *Developmental biology* **301**, 254-265 (2007).

501   23.   B. Leader, H. Lim, M. J. Carabatsos, A. Harrington, J. Ecsedy, D. Pellman, R. Maas, P.
502   Leder, Formin-2, polyploidy, hypofertility and positioning of the meiotic spindle in
503   mouse oocytes. *Nature cell biology* **4**, 921-928 (2002).

504   24.   K. Grikscheit, T. Frank, Y. Wang, R. Grosse, Junctional actin assembly is mediated by
505   Formin-like 2 downstream of Rac1. *The Journal of cell biology* **209**, 367-376 (2015).

506   25.   X. L. Ren, Y. D. Qiao, J. Y. Li, X. M. Li, D. Zhang, X. J. Zhang, X. H. Zhu, W. J. Zhou, J. Shi,
507   W. Wang, W. T. Liao, Y. Q. Ding, L. Liang, Cortactin recruits FMNL2 to promote actin
508   polymerization and endosome motility in invadopodia formation. *Cancer letters* **419**,
509   245-256 (2018).

510   26.   Y. Lu, Y. Zhang, M. H. Pan, N. H. Kim, S. C. Sun, X. S. Cui, Daam1 regulates fascin for
511   actin assembly in mouse oocyte meiosis. *Cell cycle* **16**, 1350-1356 (2017).

512   27.   M. H. Pan, F. Wang, Y. Lu, F. Tang, X. Duan, Y. Zhang, B. Xiong, S. C. Sun, FHOD1
513   regulates cytoplasmic actin-based spindle migration for mouse oocyte asymmetric cell
514   division. *Journal of cellular physiology* **233**, 2270-2278 (2018).

515   28.   Y. Zhang, F. Wang, Y. J. Niu, H. L. Liu, R. Rui, X. S. Cui, N. H. Kim, S. C. Sun, Formin
516   mDia1, a downstream molecule of FMNL1, regulates Profilin1 for actin assembly and
517   spindle organization during mouse oocyte meiosis. *Biochimica et biophysica acta* **1853**,
518   317-327 (2015).

519   29.   E. D. Goley, M. D. Welch, The ARP2/3 complex: an actin nucleator comes of age. *Nature*
520   *reviews. Molecular cell biology* **7**, 713-726 (2006).

521   30.   S. Brunet, B. Maro, Cytoskeleton and cell cycle control during meiotic maturation of the
522   mouse oocyte: integrating time and space. *Reproduction* **130**, 801-811 (2005).

523   31.   S. Reinsch, P. Gonczy, Mechanisms of nuclear positioning. *Journal of cell science* **111**
524   **(Pt 16)**, 2283-2295 (1998).

525   32.   Z. Wei, J. Greaney, C. Zhou, A. H. H, Cdk1 inactivation induces post-anaphase-onset
526   spindle migration and membrane protrusion required for extreme asymmetry in mouse
527   oocytes. *Nature communications* **9**, 4029 (2018).

528   33.   C. H. Zhang, W. P. Qian, S. T. Qi, Z. J. Ge, L. J. Min, X. L. Zhu, X. Huang, J. P. Liu, Y. C.
529   Ouyang, Y. Hou, H. Schatten, Q. Y. Sun, Maternal diabetes causes abnormal dynamic
530   changes of endoplasmic reticulum during mouse oocyte maturation and early embryo
531   development. *Reproductive biology and endocrinology : RB&E* **11**, 31 (2013).

34.   G. FitzHarris, P. Marangos, J. Carroll, Changes in endoplasmic reticulum structure during

mouse oocyte maturation are controlled by the cytoskeleton and cytoplasmic dynein.

Dev Biol **305**, 133-144 (2007).

35.   E. S. Chhabra, V. Ramabhadran, S. A. Gerber, H. N. Higgs, INF2 is an endoplasmic

reticulum-associated formin protein. *Journal of cell science* **122**, 1430-1440 (2009).

36.   F. Korobova, V. Ramabhadran, H. N. Higgs, An actin-dependent step in mitochondrial

fission mediated by the ER-associated formin INF2. *Science* **339**, 464-467 (2013).

37.   W. K. Ji, R. Chakrabarti, X. Fan, L. Schoenfeld, S. Strack, H. N. Higgs, Receptor-mediated

Drp1 oligomerization on endoplasmic reticulum. *The Journal of cell biology* **216**, 4123-

4139 (2017).

38.   A. A. Minin, A. V. Kulik, F. K. Gyoeva, Y. Li, G. Goshima, V. I. Gelfand, Regulation of

mitochondria distribution by RhoA and formins. *Journal of cell science* **119**, 659-670

(2006).

39.   U. Manor, S. Bartholomew, G. Golani, E. Christenson, M. Kozlov, H. Higgs, J. Spudich, J.

Lippincott-Schwartz, A mitochondria-anchored isoform of the actin-nucleating spire

protein regulates mitochondrial division. *eLife* **4**,   (2015).

40.   G. B. Li, H. W. Zhang, R. Q. Fu, X. Y. Hu, L. Liu, Y. N. Li, Y. X. Liu, X. Liu, J. J. Hu, Q. Deng,

Q. S. Luo, R. Zhang, N. Gao, Mitochondrial fission and mitophagy depend on cofilin-

mediated actin depolymerization activity at the mitochondrial fission site. *Oncogene* **37**,

1485-1502 (2018).

41.   S. Li, S. Xu, B. A. Roelofs, L. Boyman, W. J. Lederer, H. Sesaki, M. Karbowski, Transient

assembly of F-actin on the outer mitochondrial membrane contributes to mitochondrial

fission. *The Journal of cell biology* **208**, 109-123 (2015).

Figure legends

Figure 1. Expression and subcellular localization of FMNL2 during mouse oocyte meiosis. (A) Western blotting results of FMNL2 protein expression at different stages. FMNL2 expressed at the GV, MI, and MII stages. **(B)** Subcellular localization of FMNL2-EGFP during mouse oocyte meiosis. FMNL2 was enriched at the cortex (GV, GVBD and MI stage) and spindle periphery (MI stage). Green, FMNL2-EGFP; blue, DNA. Negative control: Green, EGFP; blue, DNA. Bar =20 μm. **(C)** Co-staining of oocytes for FMNL2 and actin. FMNL2 and actin overlapped in cortex and spindle surrounding. Green, FMNL2-antibody; red, actin; blue, DNA. Bar =20 μm.

Figure 2. Knockdown of FMNL2 affects first polar body extrusion and asymmetric

division. **(A)** Western blot analysis for FMNL2 expression in the FMNL2-KD group and control group. Relative intensity of FMNL2 and tubulin was assessed by densitometry. **, significant difference ($P < 0.01$). **(B)** DIC images of control oocytes and FMNL2-KD oocytes after 12 h culture. FMNL2-KD caused large polar bodies (black arrows) and some oocytes failed to extrude the polar bodies (white arrows). **(C)** Rate of polar body extrusion after 12 h culture of the control group and FMNL2-KD group. ***, significant difference ($P < 0.001$). **(D)** Rate of large polar body extrusion after 12 h culture in the control group and FMNL2-KD group. ****, significant difference ($P < 0.0001$). **(E)** Time-lapse microscopy showed that polar body extrusion failed after FMNL2-KD. Bar $= 10$ μm. **(F)** Western blot analysis for FMNL2 expression in the control group, FMNL2-KD group and rescue group. Relative intensity of FMNL2 and tubulin was assessed by densitometry. **(G)** DIC images of FMNL2-KD oocytes and rescue oocytes after 12 h culture. **(H)** Rate of polar body extrusion after 12 h culture of the FMNL2-KD group and rescue group. *, significant difference ($P < 0.05$). **(I)** Rate of large polar body extrusion after 12 h culture in the FMNL2-KD group and rescue group. ***, significant difference ($P < 0.001$). **(J)** Rate of polar body extrusion after 12 h culture of the control group and FMNL2+3-KD group. *, significant difference ($P < 0.05$). **(K)** Rate of large polar body extrusion after 12 h culture in the control group and FMNL2+3-KD group. ***, significant difference ($P < 0.001$).

Figure 3. Knockdown of FMNL2 disrupts spindle localization during mouse oocyte meiosis. (A) Time-lapse microscopy showed that spindle migration failed after FMNL2-KD. Green, tubulin-EGFP. Bar $= 10$ μm. **(B)** Representative images and the

proportion of spindle migration after 9.5 h of culture in the control group and FMNL2-

KD oocyte group. White, actin; green, tubulin; magenta, DNA. Bar = 10 μm. *,

significant difference (P < 0.05). **(C)** Representative images and the proportion of

spindle migration after 9.5 h of culture in the FMNL2-KD group and rescue oocyte

group. magenta, DNA. Bar = 10 μm. *, significant difference (P < 0.01).

Figure 4. Knockdown of FMNL2 disrupts actin assembly during mouse oocyte

meiosis. (A) Representative images of actin distribution at the oocyte cortex and the

fluorescent intensities in the control group and FMNL2-KD group (P > 0.1). White,

actin; green, tubulin; magenta, DNA. Bar = 10 μm. **(B)** Representative images of actin

distribution in the oocyte cytoplasm and the fluorescent intensities in the control group

and FMNL2-KD group. White, actin; green, tubulin; magenta, DNA. Bar = 10 μm.

****, significant difference (P < 0.0001). **(C)** Representative images of actin

distribution in the oocyte cytoplasm and the fluorescent intensities in the FMNL2-KD

group and rescue group. White, actin; magenta, DNA. Bar = 10 μm. ****, significant

difference (P < 0.0001) **(D)** Mass spectrometry results showed that FMNL2 was related

to many actin-related proteins. **(E)** Co-IP results showed that FMNL2 was correlated

with Arp and Formin2 but not with Profiling and Fascin. **(F)** Arp2 protein expression

significantly increased in the FMNL2-KD oocytes compared with the control oocytes.

Arp2 protein expression significantly decreased in the rescue oocytes compared with

the FMNL2-KD oocytes. **, significant difference (P < 0.01). **(G)** Formin2 protein

expression significantly decreased in the FMNL2-KD oocytes compared with the

control oocytes. Formin2 protein expression significantly increased in the rescue

oocytes compared with the FMNL2-KD oocytes. **, significant difference (P < 0.01).

Figure 5. FMNL2 regulates endoplasmic reticulum distribution during mouse

oocytes maturation. (A) Co-IP results showed that FMNL2 was correlated with INF2.

(B) Mass spectrometry results showed that FMNL2 was associated with ER-related

proteins. **(C)** Representative images of ER distribution in the oocyte cytoplasm in the

control group and FMNL2-KD group. In FMNL2-KD oocytes, ER agglomerated in

cytoplasm (white arrow). Red, ER; Blue, DNA. Bar = 10 μm. **(D)** Abnormal distribution

of ER significantly increased in the FMNL2-KD oocytes compared with the control

oocytes. *, significant difference (P < 0.05). **(E)** Grp78 and Chop protein expression

significantly increased in the FMNL2-KD oocytes compared with the control oocytes.

The band intensity analysis also confirmed this finding. *, significant difference (P <

0.05). **(F)** Representative images of ER distribution in the oocyte cytoplasm in the

FMNL2-KD group and rescue group. Red, ER; Blue, DNA. Bar = 10 μm. **(G)** Abnormal

distribution of ER significantly decreased in the rescue oocytes compared with the

FMNL2-KD oocytes. *, significant difference (P < 0.05). **(H)** Grp78 protein expression

significantly decreased in the rescue oocytes compared with the FMNL2-KD oocytes.

The band intensity analysis also confirmed this finding. **, significant difference (P <

0.01).

Figure 6. FMNL2 regulates mitochondrial distribution during mouse oocytes

maturation. (A) Mass spectrometry results showed that FMNL2 was related to many

mitochondria-related proteins. **(B)** Representative images of mitochondrial distribution

in the oocyte cytoplasm in the control group and FMNL2-KD group. In FMNL2-KD

oocytes, mitochondrial agglomerated in cytoplasm (white arrow). Green, Mito. Bar =

20 μm. **(C)** Abnormal distribution of mitochondrial significantly increased in the

FMNL2-KD oocytes compared with the control oocytes. *, significant difference ($P <$

0.05). **(D)** Representative images of mitochondrial distribution in the oocyte cytoplasm

in the FMNL2-KD group and rescue group. In FMNL2-KD oocytes, mitochondrial

agglomerated in cytoplasm (white arrow). Red, Mito; Blue, DNA. Bar = 20 μm. **(E)**

Abnormal distribution of mitochondrial significantly decreased in the rescue oocytes

compared with the FMNL2-KD oocytes. **, significant difference ($P < 0.01$). **(F)** The

typical picture for JC1 green channel and red channel after FMNL2-KD. **(G)** The JC1

signal (red/green ratio) after FMNL2-KD compare with the control group, the JC-1

red/green fluorescence ratio was significantly reduced in FMNL2-KD groups. Bar = 20

μm. **, $P < 0.01$. **(H)** cofilin protein expression significantly decreased in the FMNL2-

KD oocytes compared with the control oocytes. The band intensity analysis also

confirmed this finding. *, significant difference ($P < 0.05$).

A FMNL2 protein level after RNAi

B Control | FMNL2-KD

C Polar body extrusion rate

D Large polar body

E DIC 6h 7h 8h 9h 10h 11h 12h
Control
FMNL2-KD

F Control FMNL2-KD Rescue
FMNL2
Tubulin

G FMNL2-KD | Rescue

H Polar body extrusion

I Large polar body

J Polar body extrusion

K Large polar body

A

DIC 8h 9h 9.5h 10h 10.5h 11h 12h

Control

Tubulin

FMNL2-KD

B

Control   FMNL2-KD

Actin

Tubulin   DNA

Rate of spindle migration

Percentage %

Control   FMNL2-KD

*

C

FMNL2-KD   Rescue

DIC

DNA

Rate of spindle migration

Percentage %

FMNL2-KD   Rescue

**

A

Control | FMNL2-KD

Actin | Tubulin

DNA

Actin

Cortex actin fluorescense intensity after cultured of 9 h

Relative fluorescence intensity

Control | FMNL2-KD

B

Control | FMNL2-KD

Actin

Tubulin | DNA

Actin

Cytoplasm actin fluorescense intensity after cultured of 9 h

Relative fluorescence intensity

****

Control | FMNL2-KD

C

FMNL2-KD | Rescue

Actin

DNA

Actin

Cytoplasm actin fluorescense intensity after cultured of 9 h

Relative fluorescence intensity

****

FMNL2-KD | Rescue

D

Actin-related proteins

Clf1 | Actl6a
Smarce1 | Tpm3 | Arpc4 | Capza1
Acta2 | Fscn1 | Tpm1 | Capg | Actr3
Vcl | Pls3 | Actr1a | Arpc2 | Arpc5 | Actn1 | Arpc5l
Ctnnb1 | Ctnna1 | Capza2 | Dctn2
Clf2 | Ctnnd1 | **FMNL2** | Actb | Nptn
Inf2 | Tpm2 | Mylk | Actg1 | Dctn1
Pfn2 | Serpinf2 | Pfn1 | Capzb
Syne2 | Actn4 | Actr2 | Arfip1

E

Input | IP:FMNL2

Arp2

Profilin

fascin

Formin2

F

Relative Intensity(Arp2/Tubulin)

**

Control | FMNL2-KD

Arp2

Tubulin

Relative Intensity(Arp2/Tubulin)

**

FMNL2-KD | Rescue

Arp2

Tubulin

G

Relative Intensity(Formin2/Tubulin)

*

Control | FMNL2-KD

Formin2

Tubulin

Relative Intensity(Formin2/Tubulin)

**

FMNL2-KD | Rescue

Formin2

Actin

A Mitochondria-related proteins

Ndufs1 · Hibch · Aldh5a1 · Myg1
Me2 · Acat1 · Fh · Cyp11a1 · Ak2 · Aldh6a1
Slc25a11 · Acaa2 · Hadha · Vcp · Aldh2 · Echs1 · Aifm1
Prdx5 · Acadvl · Idh2 · Dld · Fdxr · Mdh2 · Glud1 · Pcx · Suclg2
Idh3a · Acadl · Uqcrc2 · **FMNL2** · Hspa9 · Atp5f1 · Aldh1b1
Got2 · Pitrm1 · Abat · Atp5o · · Hmgcs2 · Lonp1 · Acadm
Sucla2 · Atp5c1 · Atp5f1a · Slc25a12 · Trap1 · Cs · Hadhb · Tufm
Oxct1 · Etfdh · Lrpprc · Atp5f1b · Aco2 · Etfa · Slc25a3 · Hadh
Sdhb · Eci1 · Uqcrc1 · Pccb · Iars2 · Prdx3 · Sdha · Acads
Acsf2 · Hspe1 · Pck2 · Ndufv1
Inf2 · Cyc1

B Mito / DIC — Control, FMNL2-KD

H Relative Intensity(Cofilin/Tubulin) — Control, FMNL2-KD; Cofilin; Tubulin

D DIC / Mito / Merge — FMNL2-KD, Rescue; Mito / DNA

C Percentage % — Control, FMNL2-KD; normal, 1≤clumps≤3, 4≤clumps≤6, clumps>6

E Percentage % — FMNL2-KD, Rescue; normal, 1≤clumps≤3, 4≤clumps≤6, clumps>6

F Red channel / Green channel / JC1 Merge — Control, FMNL2-KD

G JC1 signal (Red/Green Ratio) — Control, FMNL2-KD