1   Neutrophil-derived IL-17 limits protective host responses and promotes tuberculosis
2   pathogenesis
3
4

5   Priya Sharma[1], Raman Deep Sharma[1], Mrinmoy Das[1], Binayak Sarkar[2], Lakshyaveer Singh[1],
6   Neharika Jain[1], Shivam Chaturvedi[1], Lalita Mehra[3], Aditya Rathee[1], Shilpa Sharma[1], Shihui
7   Foo[4], Andrea Lee[5], N Pavan kumar[6], Prasenjit Das[3], Vijay Viswanathan[7], Hardy Kornfeld[8],
8   Shanshan W. Howland[4], Subash Babu[9,10], Vinay Nandicoori[2,11], Amit Singhal[5,12] and Dhiraj
9   Kumar[1*]
10

13   [1]Cellular Immunology Group, International Centre for Genetic Engineering and
14   Biotechnology, Aruna Asaf Ali Marg, New Delhi, India-110067
15   [2] National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi, India-110067
16   [3]Department of Pathology, All India Institute of Medical Sciences, New Delhi, India-110012
17   [4]Singapore Immunology Network (SIgN), Agency for Science, Technology and Research
18   (A*STAR), Singapore, Republic of Singapore.
19   [5]Infectious Diseases Labs (ID labs), Agency for Science, Technology and Research
20   (A*STAR), Singapore, Republic of Singapore
21   [6]Department of Immunology, ICMR-National Institute for Research in Tuberculosis,
22   Chennai, India – 600031
23   [7]Prof. M. Viswanathan Diabetes Research Center, Chennai, India
24   [8]Department of Medicine, University of Massachusetts Medical School, Worcester, MA,
25   USA
26   [9]NIH-International Center of Excellence in Research, Chennai, India
27   [10]Laboratory of Parasitic Diseases, NIAID-NIH, Bethesda, Maryland, USA
28   [11]Centre for Cellular and Molecular Biology, Hyderabad, India
29   [12]Lee Kong Chian School of Medicine, Nanyang Technological University (NTU),
30   Singapore, Republic of Singapore.
31

*Correspondence: dhiraj@icgeb.res.in

Running title: IL-17 and PMNs in Tuberculosis Pathogenesis

Abstract

The protective correlates of *Mycobacterium tuberculosis* (*Mtb*) infection-elicited host

immune responses are incompletely understood. Here we report pro-pathogenic crosstalk

between IL-17, COX2, and Neutrophils. We show that most intracellular bacilli in the wild-

type (WT) mice's lungs are neutrophil-resident. In BCG-vaccinated WT mice, *Mtb* continues

to persist inside neutrophils, which is sensitive to COX2 inhibition. Using a genetically

susceptible *IFNγ* $^{-/-}$ mice model, we report that excessive neutrophil infiltration and

corresponding bacteremia are dependent on neutrophil-derived IL-17. COX2 inhibition and

IL-17 neutralization reverse neutrophil infiltration, associated pathology and death in *IFNγ* $^{-/-}$

mice. The IL-17-Neutrophils interplay also operates in WT mice, and RORγt inhibition

enhances BCG's protective efficacy. In a prospective clinical cohort of pulmonary TB, high

neutrophil count and IL-17 precipitate adverse treatment outcomes. We propose targeting the

pro-pathogenic IL-17- Neutrophils axis to enhance TB prevention and therapy in normal and

vulnerable subjects.

Introduction

Tuberculosis (TB) is the major cause of mortality and morbidity due to an infectious disease. A lack of an effective vaccine, prognostic diversity among the patients, and the emergence of drug resistance greatly hamper the efforts to control TB. BCG, the only available vaccine, has limited efficacy in adults, although it works well in preventing childhood TB and TB meningitis. A comprehensive understanding of how the host immune system responds to vaccination, infection and treatment is fundamental to the success of TB elimination programs (Chandra et al., 2022). Not all individuals exposed to *Mtb* follow a common trajectory on parameters like developing latent or active disease, disease severity and responsiveness to anti-TB chemotherapy (Barry et al., 2009; Cadena et al., 2017; Dartois and Rubin, 2022; Heyckendorf et al., 2021; Pai et al., 2016; Thompson et al., 2017). For example, the average treatment success rate for TB is ~80%, which is further lower in the case of drug-resistant TB (DR-TB, 50%) (Chaves Torres et al., 2019; Rockwood et al., 2016). Immunological features that contribute to treatment success and failure in non-DR-TB patients, presents a unique opportunity to understand the host immunity associated with protection against TB. Unfortunately, the immune parameters that define vaccine efficacy or treatment outcomes are poorly understood. There is a consensus that host-driven mechanisms play a critical role in the differential BCG response between young children and adults (Brazier and McShane, 2020). For example, the BCG-induced memory T-cell responses are believed to fade over time (Soares et al., 2013). For intracellular pathogens like *Mtb*, understandably, the cell-mediated immunity arm acquires greater importance in terms of vaccine efficacy (Brazier and McShane, 2020). However, since the bacteria mostly reside inside the innate immune cells, it is pertinent to ask- whether innate immune-based mechanisms also contribute to the differential protective responses. To further elaborate, the presence of *Mtb* has been shown in unconventional cells like mesenchymal stem cells

73 (MSCs), bone cells, adipocytes etc. (Das et al., 2013; Fatima et al., 2020; Jain et al., 2020).

74 The bacteria residing in the MSCs can escape immune cytokines like IFNγ and TNFα, which

75 could effectively blunt the T-cell-mediated protective immune responses (Jain et al., 2020).

76 TB granulomas also comprise other innate immune cells, like granulocytes and dendritic cells

77 (Cadena et al., 2017). Whether different cellular constituents confound the protective host

78 responses remains poorly understood. Neutrophils are the most extensively studied

79 granulocytes in the context of TB (Dallenga and Schaible, 2016; Kimmey et al., 2015;

80 Lovewell et al., 2021; Mishra et al., 2017). They represent the most common *Mtb*-containing

81 innate cells in the airways of active TB patients (Eum et al., 2010). Recent studies point to a

82 more pathological role of neutrophils in TB (Berry et al., 2010; Han et al., 2018; Lowe et al.,

83 2013; Nwongbouwoh Muefong et al., 2022) although some studies also report their protective

84 roles in specific contexts (Lowe et al., 2018; Martineau et al., 2007; Silva et al., 1989).

85 Using the mice model of tuberculosis, we explored neutrophils as a distinctive and preferred

86 niche for *Mtb* in this study. Our results show that neutrophil-resident *Mtb* contributes to lower

87 the efficacy of BCG vaccines and precipitates incidences of severe pathology and treatment

88 failure. We propose targeting this population can significantly enhance the preventive

89 responses of BCG, enable better control in genetically vulnerable subjects, and provide an

90 unexpected opportunity for host-directed therapy against TB.

Results

Neutrophils serve as the major niche for *Mycobacterium tuberculosis* in mice lungs.

To assess diverse cellular niches in the lungs of *Mtb*-infected C57BL/6 (WT) mice, we investigated intracellular burden in Ly6G$^+$CD11b$^+$ neutrophils (herein after also referred as polymorphonuclear neutrophils or PMNs), CD64$^+$MerTK$^+$ macrophages and Sca1$^+$CD90.1$^+$CD73$^+$ mesenchymal stem cells (MSCs). Mice were infected with 100-200 H37Rv colony forming units (CFUs) via aerosol challenge, and at 4, 8 and, 12 weeks post-infection (p.i.), the total bacterial burden in the lung and spleen was estimated by CFU counting (Fig. S1A-C). Half of the lungs from each animal was processed to flow-sort PMNs, Macrophages and MSCs following the gating strategy shown in Fig. S1D. Among the three cell populations, MSCs were the most abundant cell type across each of the time points post *Mtb* infection (2-6x10^4/million sorted cells), followed by macrophages (4-9x10^3/million sorted cells) and PMNs (1-4x10^3 /million sorted cells, Fig. S1E). However, PMNs contained, on an average, 4-5 times more total bacillary load (from the total sorted pool of cells) as compared to either macrophages or MSCs (Fig. S1F). To account for the differences in the total number of the three cell types, we calculated CFU/10^4 cells. This also showed the highest bacillary load in the PMNs (Fig. S1G). Thus, a disproportionately large number of *Mtb* in the lungs of infected animals reside inside the PMNs, which outnumber those present in macrophages or MSCs.

The residual *Mtb* population in BCG-vaccinated animals are largely neutrophil-resident.

To understand the relevance of neutrophil-resident *Mtb* and its contribution towards TB pathogenesis, we repeated the above experiments in BCG-vaccinated animals. Mice were administered either PBS or BCG (Pasteur strain) intradermally, followed by aerosol challenge with H37Rv at eight weeks post-BCG vaccination (Fig. 1A). As expected, animals that

received BCG showed nearly ten-fold lower bacterial burden in the lungs and spleen

compared to the unvaccinated PBS controls (Fig. 1B-C). Consequently, BCG-vaccinated

animals also showed less pathology as compared to the unvaccinated PBS controls (Fig.

S1H). Among the sorted cells from the lungs, except for a marginal decline in PMNs number

at eight weeks post-infection in the BCG-vaccinated group, the relative distribution of PMNs,

macrophages and MSCs in both control and BCG-vaccinated animals remained unperturbed

over the course of infection (Fig. 1D). The bacterial burden, however, in each of the three cell

types dropped significantly (by 80-90%) in the BCG-vaccinated animals (Fig. 1E). Curiously,

lung PMNs in BCG-vaccinated mice still contained 10-20-fold more bacilli compared to the

other two cell types for each time points p.i. investigated (Fig. 1E). These results indicate that

BCG-induced protective host immunity is unable to control the PMNs -resident *Mtb*

efficiently. Thus, it appears that PMNs serve as a protective niche for *Mtb* against

immunological assaults such as BCG vaccination in mice.

RNAseq analysis of neutrophils sorted from the lungs of infected animals.

We next analyzed the global gene expression profile of flow-sorted PMNs from the lungs of

unvaccinated or BCG-vaccinated mice, either uninfected or *Mtb*-challenged (Fig. 1F). Only

27 differentially regulated genes were found between unvaccinated, uninfected (UI) and

BCG-vaccinated, uninfected animals, and therefore, the BCG-vaccinated, uninfected mice

group was excluded from the heatmap in figure 1F (Table S1). Compared to the UI control

group, 944 genes were differentially expressed in the *Mtb*-infection group alone, whereas

there were 620 DEGs in the BCG-vaccinated *Mtb*-infected group, 373 genes being common

to both the lists (Fig. S1I). Compared to the *Mtb*-infected group, the BCG-vaccinated *Mtb*-

infected group had 209 DEGs (Fig. S1I). The union of these DEGs (n=1332) formed seven

distinct clusters (Fig. 1F, Table S2). Table S2 shows Gene Ontology Biological Process (GO-

BP) enriched among genes in each cluster. Clusters 2 and 6 comprised of downregulation and

upregulation, respectively, upon *Mtb* infection alone, the expression of which were not

altered by BCG-vaccination (Fig. 1F). For the remaining five clusters, a few selected GO-BP

classes and corresponding genes are shown in figure 1F. Each of these clusters showed a

distinctive pattern in the BCG-vaccinated group compared to the *Mtb*-infected group.

Compared to the unvaccinated *Mtb*-infected group the expression of genes for clusters 1 and

3 was reverted in the BCG-vaccinated *Mtb*-infected group to the level observed in uninfected

animals, suggesting BCG vaccination-mediated reversal in the neutrophil transcriptional

program. The cluster 1 genes (upregulated in BCG-vaccinated *Mtb*-infected group) belonged

to functional classes like xenophagy, T cell proliferation, Eosinophil activation etc., whereas

genes in cluster 3 (downregulated in BCG-vaccinated-*Mtb*-infected group) were involved in

lactate metabolism, mitophagy, microtubule polymerization etc. Similarly, genes in cluster 4

(downregulated in BCG-vaccinated-*Mtb*-infected group) belonged to functional classes like

blood vessel remodelling, inflammatory response to wounding, TGFβ signalling and Th17

cell chemotaxis (Fig. 1F). Th17 cell chemotaxis has been associated with exacerbated TB

pathology and therefore a decline in this pathway in BCG-vaccinated group concurred with

earlier studies(Jung et al., 2022) (Basile et al., 2011; Jurado et al., 2012; Mills, 2023). Genes

in cluster 5 were upregulated only in the BCG-vaccinated-*Mtb*-infected group and belonged

to steroid metabolism, TLR2 signaling pathway, epithelium cell activation and leukotriene

production (Fig. 1F). To note, increased production of leukotriene is also associated with TB

protection (Pavan Kumar et al., 2019; Peres et al., 2007; Tobin et al., 2010). Cluster 7 genes,

expression of which was marginally lower in the BCG-vaccinated *Mtb*-infected group

compared to the *Mtb*-infected group alone, belonged to IL1β production, response to type I

IFN, macrophage activation and nitric oxide production (Fig. 1F). Taken together, RNAseq

analysis provided insights into the association of the functional states of PMNs with either

pathology (*Mtb*-infected) or protection (BCG-vaccinated, *Mtb*-infected) from TB.

Celecoxib potentiates the effect of BCG by targeting the neutrophil-resident bacterial population.

The RNAseq data indicated an association of enhanced leukotriene pathway with the protective functional state of neutrophils in the BCG-vaccinated *Mtb*-infected group (cluster 5, Fig. 1F). Neutrophil recruitment to the site of inflammation is driven by the chemotactic signaling via receptor CXCR2 (Kolaczkowska and Kubes, 2013) and the eicosanoids downstream to COX2 (Funk, 2001; Lemos et al., 2009). The COX2 enzyme competes with 5-lipoxygenase (5-LO) enzyme for the substrate arachidonic acid to promote prostaglandin and thromboxane biosynthesis (Funk, 2001). On the other hand, 5-LO promotes leukotriene synthesis pathways. We found increased prostaglandin E2 (PGE2) level in the lung homogenates of *Mtb*-infected mice compared to the uninfected mice (Fig. S2A). Since BCG vaccination was associated with the activation of leukotriene pathway (Fig. 1F), we hypothesized that skewing the balance away from PGE2 biosynthesis may impact the survival of neutrophil-resident *Mtb* pool. Therefore, we treated unvaccinated and BCG-vaccinated *Mtb*-infected mice with celecoxib, a specific COX2 inhibitor, and evaluated total lung and spleen CFU at 8- and 12-week p.i. and PMNs-resident *Mtb* pool in the lungs at 12-week p.i. (Fig. 2A). Celecoxib (50mg/kg/day) significantly reduced the bacterial burden in the lungs of unvaccinated and BCG-vaccinated *Mtb*-infected mice, with a greater impact in BCG-vaccinated mice (Fig. 2B). While the responses were similar in both lungs and spleen, they were more pronounced in the lungs (Fig. 2B). This corresponded to celecoxib-mediated decline in PMNs-resident pool of *Mtb* (by ~45%) in the lung of BCG-vaccinated *Mtb*-infected mice (Fig. 2C). To note, celecoxib treatment had no significant effect on the bacterial burden in macrophages and MSCs (Fig. 2C). Furthermore, histological analysis revealed significant recovery in the lung pathology in BCG-vaccinated animals that received celecoxib (Fig. 2D). Finally, neither BCG or celecoxib treatment had any significant impact

on the number of PMNs, macrophages or MSCs in the lungs at 12 weeks p.i. (Fig. S2B).

These results suggest a role of COX2 in determining the intracellular *Mtb* pool in the

neutrophils.

To assess the effect of celecoxib on the transcriptional landscape of neutrophils, we

performed RNAseq on sorted lung PMNs from celecoxib-treated unvaccinated or BCG-

vaccinated *Mtb*-infected mice. Here we compared the transcriptome of PMNs from

uninfected mice with those from *Mtb*-infected, celecoxib-treated *Mtb*-infected and celecoxib-

treated, BCG-vaccinated and *Mtb*-infected mice (Fig. 2E, Table S3). A total of 2190 genes

were differentially regulated in this comparison group, which could be clustered in four major

clusters (Fig. 2E, Table S3). Cluster 2 was of particular interest since it was majorly

upregulated only in the group that was BCG-vaccinated, *Mtb*-infected and received celecoxib

(Fig. 2E). Functionally, these genes belonged to biological processes like wound healing,

muscle tissue development, phagocytosis, mononuclear cell proliferation and differentiation

(Fig. 2F, Table S3). To understand the direct targets of celecoxib in the PMNs, we compared

genes from celecoxib-treated and *Mtb*-infected mice with those from celecoxib-treated, BCG-

vaccinated and *Mtb*-infected mice. Of around 23 common genes, a majority of them belonged

to defence mechanisms like response to IFNγ, TNF, LPS and production of cytokines in

inflammatory response (Fig S2C, Table S4). It, therefore, appears that celecoxib activates

anti-bacterial effector function in the neutrophils of *Mtb*-infected mice. These results

underscore two important points- i) PMN's effector functions appear to be perturbed during

Mtb infection, and ii) inhibiting COX2 triggers anti-bacterial pathways in the PMNs.

In the *IFNγ*$^{-/-}$ animals, neutrophils account for high bacterial burden and disease

pathology.

The results so far demonstrate a key role of PMNs-resident *Mtb* in TB pathology. While

COX2 inhibition helped clear this pool of *Mtb*, the mechanism of PMNs-driven pathology

was unclear. To understand the mechanism, we used *IFNγ*$^{-/-}$ animals, which are known to

develop severe neutrophilia and pathology upon *Mtb* infection (Cooper et al., 1993; Flynn et

al., 1993; Pearl et al., 2001). We first confirmed the severe disease phenotype in *Mtb*-infected

IFNγ$^{-/-}$ mice (Fig. 3A), which included severe gross pathology in the lungs, a very high

bacterial burden in the lungs and spleen and high mortality, compared to WT mice (Fig. S3A-

C). Histopathology analysis of lung sections revealed massive infiltration of the immune cells

(Fig. 3B), especially PMNs, as shown by immunofluorescence staining in the *IFNγ*$^{-/-}$ mice

compared to the WT mice (Fig. 3C). *Mtb*-infected *IFNγ*$^{-/-}$ mice showed a nearly 40-fold

higher number of PMNs (Fig. 3D) and ~2.5-fold higher macrophage numbers than the WT

animals (Fig. S3D). Unlike PMNsand macrophages, a decline in the number of MSCs was

observed in *IFNγ*$^{-/-}$ mice than the WT animals (Fig. S3D). In terms of resident *Mtb* in the

sorted cells, bacterial burden increased several folds in each of the three cell types (Fig. 3E

and S3E). However, due to the sheer number, there were more bacteria in the PMN

compartment than any other cell types in the *IFNγ*$^{-/-}$ mice, especially at 10 weeks p.i. Thus,

massive infiltration of PMNs and a very high intracellular *Mtb* burden in PMNs signifies

severe pathology in *IFNγ*$^{-/-}$ mice.

Cytokine dysregulation in *IFNγ*$^{-/-}$ animals suggests a role of IL17 in disease pathology.

To understand what contributed to the increased immune cells infiltration in the *Mtb*-infected

IFNγ$^{-/-}$ animals, we analyzed cytokine levels in the lung homogenates of the infected mice.

Several cytokines/chemokines were significantly dysregulated in the *Mtb*-infected *IFNγ*$^{-/-}$

mice compared to the *Mtb*-infected WT mice (Fig. 3F-G, S3G). Among those CXCL1,

CXCL2, IL-1β, IL-1α, IL-6, IL-17A (broadly known as IL-17), M-CSF, CCL3 and TNFα

were significantly upregulated (Fig. 3F) whereas IFNγ, IL-12, IL-9, IL-10, IL-15, IL-18, IL-

27 and IL-28 were downregulated (Fig. 3G). Among the upregulated cytokines in *Mtb*-

infected *IFNγ⁻ᐟ⁻* animals, CXCL1 and CXCL2 are the known chemoattractant for the PMNs

recruitment (Kolaczkowska and Kubes, 2013), whereas IL-1α, IL-1β and IL-6 are key

players in differentiation and expansion of the Th17 cells (Ketelut-Carneiro et al., 2019; Lee

et al., 2010; Mills, 2023). Interestingly, IL-17 was also significantly upregulated in the *Mtb*-

infected *IFNγ⁻ᐟ⁻* animals (Fig. 3F). Thus, more than one cytokine increased in the lungs of

IFNγ⁻ᐟ⁻ animals upon *Mtb* infection converged to IL-17 regulation, indicating a key role for

IL-17 in disease pathology, which also concurs with earlier studies (Cruz et al., 2010).

Neutrophils are the key source of IL-17 in TB granulomas

IL-17 is a classical pro-inflammatory cytokine predominantly secreted by the Th_{17} cells,

which are important for TB pathology (Mills, 2023). Also, IL-17 has been shown to be

responsible for the recruitment and activation of neutrophils (Allen et al., 2015). To

investigate the source of IL-17, we first analyzed the IL-17-producing cells in the lungs of

Mtb-infected WT mice (Fig. 4A). To our surprise, >80% of PMNs were also positive for IL-

17 in the lungs of *Mtb*-infected WT mice (Fig. 4A). In the $CD45^+CD11b^-Ly6G^-$ population,

which mostly represented adaptive immune cells like T cells, B cells, NK cells etc., ~48%

were also positive for IL-17 (Fig. 4A). Immuno-fluorescence analysis of lung sections from

Mtb-infected WT mice revealed similar observation when stained for CD4 and IL-17. While

we could witness CD4 and IL-17 co-expressing cells, several $IL-17^+$ cells lacked CD4

expression (Fig. 4B). Staining for Ly6G and IL-17 revealed intense IL-17-Ly6G co-staining

in the lung tissue sections of *Mtb*-infected WT mice, suggesting PMNs as the source of lung

IL-17 (Fig. 4C). A similar IL-17-Ly6G co-staining was also observed in the *Mtb*-infected

IFNγ⁻ᐟ⁻ mice (Fig. 4D). Interestingly, even in human TB granulomas, we observed significant

co-staining of CD66b (the human neutrophil marker) and IL-17 (Fig. 4E). Thus, it appears

that there is an IL-17-PMNs axis operational during *Mtb* infection, which may contribute

towards TB pathology.

IL-17 neutralization controls bacterial burden and disease pathology in *IFNγ⁻/⁻* animals

The impact of IL-17 on TB pathogenesis is not unequivocally established. For example, IL-17 is believed to exacerbate TB pathology during the chronic phase of infection, partly through excessive neutrophil recruitment (Torrado and Cooper, 2010). On the other hand, a more protective role of IL-17 in the WT mice was reported when infected with a hypervirulent strain of *Mtb* (Domingo-Gonzalez et al., 2017). Since the results in the above sections suggested a pro-pathogenic role of IL-17, we decided to neutralize IL-17 in *Mtb*-infected *IFNγ⁻/⁻* mice. The *Mtb*-infected *IFNγ⁻/⁻* mice were administered anti-IL-17 monoclonal Ab (mAb) intra-peritoneally every 3-4 days, starting a day before the infection (Fig. 5A). Similar to the WT mice, the lung homogenates of *IFNγ⁻/⁻* mice also showed a copious amount of PGE2 upon *Mtb* infection, prompting us to include a celecoxib arm in this experiment (Fig. S4A, Fig. 5). A significant decline in IL-17 levels in the lung homogenates was observed in IL-17 mAb-administered mice (Fig. S4B). Neutralizing IL-17 led to a significant decline in the number of infiltrating PMNs in the lungs of *Mtb*-infected *IFNγ⁻/⁻* mice (Fig. 5B). Unlike PMNs, there was no effect of IL-17 neutralization on the number of macrophages or MSCs (Fig. S4C). *Mtb*-infected *IFNγ⁻/⁻* mice that received celecoxib alone also showed a significant decline in PMN recruitment (Fig. 5B). There was no additive effect on the recruitment of PMNs when celecoxib treatment was given in addition to IL-17 mAb (Fig. 5B). While IL-17 neutralization alone reduced PMN recruitment, it could not bring down the *Mtb* burden in the lungs of *IFNγ⁻/⁻* mice (Fig. 5C). Combined with celecoxib, IL-17 neutralization significantly brought down the *Mtb* burden in the lungs of *IFNγ⁻/⁻* mice (Fig. 5C). Unlike in the lungs, IL-17 neutralization alone was sufficient to reduce bacterial dissemination to the spleen (Fig. 5D), suggesting PMN's active role in *Mtb* dissemination. Moreover, IL-17 neutralization or celecoxib treatment alone did not impact bacterial burden in the sorted lung PMNs (Fig. 5E). However, the combination of IL-17 neutralization and

celecoxib treatment significantly brought down bacterial load in the PMNs (Fig. 5E). On the

contrary, celecoxib alone significantly reduced the bacterial burden in the sorted

macrophages and the MSCs (Fig. S4D). Combined with IL-17 neutralization, macrophage-

resident but not MSC-resident *Mtb* numbers further declined compared to celecoxib treatment

alone (Fig. S4D). Accordingly, IL-17 neutralization alone or in combination with celecoxib

treatment led to a significant rescue in the survival of $IFN\gamma^{-/-}$ animals (Fig. 5F) and recovery

in lung pathology (Fig. 5G-H). Taken together, while celecoxib alone is insufficient to

control bacterial burden in $IFN\gamma^{-/-}$ animals, combined with IL-17 neutralization, it effectively

controls *Mtb* growth, tissue pathology and dissemination.

Pathological role of PMN-derived IL-17 in the lungs of *Mtb*-infected $IFN\gamma^{-/-}$ animals.

To check the effect of IL-17 neutralization or celecoxib treatment on local IL-17-producing

cells, we first assessed the number of IL-17-positive T helper cells (Th$_{17}$) in the single-cell

preparations from the lungs. While there was a trend of a decline in the number of Th$_{17}$ cells

upon either IL-17 neutralization alone or in combination to COX2 inhibition in the lungs of

Mtb-infected $IFN\gamma^{-/-}$ mice, those changes were not found significant (Fig. 5I). In the lung

sections stained for IL-17 along with Ly6G, IL-17 neutralization, either alone or in

combination with COX2 inhibition, led to a consistent decline in IL-17$^+$ Ly6G$^+$ cells (Fig.

5J). The results from IL-17 ELISA in the lung homogenates, Th$_{17}$ cell count in the lungs and

IL-17-Ly6G co-staining together suggest that while IL-17 contributes to high PMN

infiltration, PMNs itself are a significant source of IL-17, thereby establishing a vicious loop.

Since PMNs are also known to express IL-17 receptors (Tan et al., 2013; Taylor et al., 2014),

these results raise the possibility that IL-17 and PMNs could establish a self-amplification

loop, leading to severe TB pathology.

Two distinct populations of PMNs with contrasting behaviours determine the fate of

infection.

Both pro- and anti-bacterial roles are attributed to PMNs largely due to some heterogeneity in

the PMN population (Leliefeld et al., 2018; Tsuda et al., 2004). Results in earlier sections

showed a non-linear association between PMN numbers, associated bacterial burden and

overall bacteremia in the lung. For example, BCG or BCG and celecoxib did not reduce PMN

numbers, but the PMN-resident bacterial pool and total tissue bacterial load showed a

significant decline (Fig. 2). On the other hand, in *IFNγ*[-/-] animals, IL-17 neutralization and

celecoxib treatment led to different outcomes. For example, IL-17 neutralization or celecoxib

treatment alone reduced PMN infiltration but had no impact on CFU in PMNs and whole

lung, while celecoxib treatment, in addition to IL-17 neutralization, led to both - decline in

PMN infiltration and bacterial burden (Fig. 4). We re-analyzed our FACS data to explore

whether there are different subtypes of PMNs which differently associated with the bacterial

burden. We observed at least two distinct PMN sub-populations, grossly separated as low and

high FSC on the FSC-SSC plot (Fig. S5A). Interestingly, animals that received anti-IL-17

mAb and celecoxib together had a higher ratio of FSC^{high} to FSC^{low} PMNs than other groups

(Fig. S5B-D). Since this group showed the least PMN-resident *Mtb* and lower burden in the

lungs and spleen, FSC^{high} PMNs appear to be more restrictive towards *Mtb*, whereas FSC^{low}

PMNs appear more permissive or pathological. These sub-types need further characterization

to understand their true nature.

Inhibiting RORγt breaks the PMN-IL-17 axis in WT animals and facilitates celecoxib-

mediated control of TB.

IL-17 production by Th17 cells and PMNs is driven by the transcription factor RORγt (Berry

et al., 2010; Tan et al., 2013). While in *Mtb*-infected *IFNγ*[-/-] animals, IL-17 levels were very

high, even the WT animals showed an increase in IL-17 levels upon *Mtb* infection, albeit to a

lower magnitude than the *IFNγ*[-/-] mice (Fig. S5E, Fig. 3F). Since the WT mice, like *IFNγ*[-/-]

mice, also showed substantial IL-17-Ly6G co-staining (Fig. 4C), we reasoned that a similar

IL-17-PMN axis could also be operating in the WT animals and contribute to sustained PMN-

resident pool of *Mtb* in the lungs. To test this hypothesis, we decided to inhibit RORγt in

Mtb-infected WT mice (Fig. 6A). Treatment with RORγt inhibitor SR2211 (inverse agonist)

led to a significant decline in IL-17 levels in the lung homogenates (Fig. 6B). Interestingly,

unlike the *Mtb*-infected *IFNγ* $^{-/-}$ mice, celecoxib alone also was able to reduce the level of IL-

17 in the lung homogenates of *Mtb*-infected WT mice, and so was the combination of both

SR2211 and celecoxib (Fig. 6B). To our surprise, SR2211 treatment also led to a loss of IL-

17-Ly6G co-staining, suggesting the role of PMN-derived IL-17 in disease pathology in the

WT mice (Fig. 6C). Treatment with SR2211 did not impact the PMN count in the lungs but

showed reduced PMN-resident *Mtb* population (Fig. 6D-E). Celecoxib, either alone or in

combination with SR2211, also reduced the intra-PMN *Mtb* load without impacting PMN

numbers (Fig. 6D-E). At the whole tissue level, SR2211 alone did not reduce *Mtb* burden in

the lungs. However, it was highly efficient in controlling dissemination to the spleen (Fig.

6F). Similarly, celecoxib treatment, either alone or in combination with SR2211, significantly

reduced bacterial CFUs in lung and spleen; however, the effects were more pronounced in the

spleen (Fig. 6F). The histopathology analysis of the lungs reflected the protective effects of

SR2211 and celecoxib (Fig. S6A-B). Treatment with SR2211 and celecoxib also impacted

the numbers of lung macrophages and MSCs and the resident bacterial population in

macrophages (Fig. S6C-F). Our data show that an IL-17-PMN axis gets established during

Mtb infection, which, if uncontrolled, could lead to severe pathology and disease. At the same

time, the results also provide a mechanistic basis for exploring RORγt inhibition as an adjunct

to preventive and therapeutic approaches against TB.

RORγt inhibition enhances the protective efficacy of BCG in WT mice

Since BCG vaccination failed to control the PMN-resident *Mtb* population, and our

subsequent results showed a strong influence of IL-17 on PMN recruitment and function, we

wondered whether blocking IL-17 could also enhance BCG efficacy, specifically by targeting

the PMN-resident *Mtb* population. So next, we treated BCG-vaccinated, *Mtb*-infected WT

mice with SR2211 (Fig. 6G). Treating BCG-vaccinated, *Mtb*-infected WT mice with SR2211

led to a ~3-fold reduction in the PMN-resident pool of *Mtb* (Fig. 6H). This also led to a more

than 7-fold decline in the total bacterial burden seen in the lungs of these mice (Fig. 6I).

Thus, together, these results indicate that the pro-pathogenic role of IL-17 also negatively

impacts the efficacy of BCG vaccination.

The IL-17-Neutrophil axis in human TB subjects

The results so far establish the role of the IL-17-PMN axis in TB pathology in mice. To

verify the existence of a similar axis in human TB, we took help from a prospective cohort

study of pulmonary tuberculosis (PTB). This study comprised 201 subjects of PTB, who were

treated for 6-8 months and followed up for an additional one year to track any recurrence

(Kumar et al., 2021; Kumar et al., 2020). Among those, 133 subjects were cured of PTB after

the full anti-TB treatment (ATT) regime (Fig. 7A). The cohort also had 68 subjects that failed

the standard ATT regime and showed manifestation of TB recurrence and death (Fig. 7A).

We compared this cohort's IL-17 and neutrophil levels at the time of recruitment (under

treatment-naïve conditions). The failed treatment group had significantly higher serum IL-17

levels at the time of recruitment (Fig. 7B). Moreover, the treatment failure group also had a

significantly higher absolute blood-neutrophil count and an increased neutrophil-to-

lymphocyte ratio (NLR) at the time of recruitment (Fig. 7C-D). Thus, the IL-17-PMN axis

appears to influence TB pathogenesis in both mice and humans, thereby providing an

opportunity to develop a protective strategy by targeting this axis.

Discussions

The immune response elicited by *Mtb* infection triggers the recruitment of diverse innate and adaptive immune cells to the site of infection, leading to the formation of granulomas (Ramakrishnan, 2012). A variety of infiltrating innate cells are expected to phagocytose and kill the pathogen. However, *Mtb* coopts these cells and uses them as a niche to evade adaptive immune response during the chronic phase (Chandra et al., 2022; Lee et al., 2020; Lovewell et al., 2021). For example, different monocyte and macrophage populations in the lungs are reported as major immune cells harbouring the intracellular *Mtb* in the lungs of infected mice, which may also impact *Mtb* growth and survival differentially (Lee et al., 2020)· (Huang et al., 2018). Similarly, MSCs are shown to offer an immune-privileged niche to the *Mtb*, protecting them from inflammatory cytokines (Jain et al., 2020). In *Mtb*-infected mice lungs, neutrophils were reported to have the highest number of bacilli early during the infection(Huang et al., 2018), while another study showed Ly6G$^+$ granulocytes as a permissive niche to *Mtb*(Lovewell et al., 2021; Mishra et al., 2017). In TB patients, neutrophils are the most predominant infected phagocytes in the sputum samples, BAL fluid and lung cavities (Eum et al., 2010). Extending the results from human patient studies and mice experiments described above, counting CFUs from the cells sorted from infected mice's lungs, we also observed that PMNs contained the largest pool of intracellular bacilli.

The role of neutrophils in impeding the protective response of BCG vaccination was unexpected. While certain innate immune responses of BCG-vaccinated animals are attributed to the trained immunity (Kaufmann et al., 2018), it is also reported that BCG vaccination results in neutrophil reprogramming, leading to an enhanced anti-bacterial state (Moorlag et al., 2020). We did observe depletion in the neutrophil-resident *Mtb* population, suggesting an anti-bacterial activation state. However, they continued to represent the largest, if not the only, pool of intracellular bacilli in those animals. Could the anti-bacterial state of

neutrophils in BCG-vaccinated animals be further enhanced? It is reported that BCG-dependent COX-2/PGE2 pathway activation blunts macrophage, DC and PMN responses (Ballinger et al., 2006; Serezani et al., 2007; Sheppe and Edelmann, 2021). This has led to explorations on a combination of BCG with selective COX2 inhibitors to ameliorate the efficacy of BCG (EP2763666B1) (Ibrahim et al., 2021). Curiously, selective COX-2 inhibition by celecoxib had no impact on neutrophil numbers in the lungs; it significantly reduced the *Mtb* load in the neutrophils in BCG-vaccinated animals. Thus, our results show that the COX2 pathway contributes to lower the efficacy of BCG-mediated protection by preventing the anti-bacterial functions of neutrophils. The COX-2-mediated negative regulation of neutrophil anti-bacterial response is also shown in the case of *P. aeruginosa* infection (Sadikot et al., 2007).

Neutrophil infiltration to the site of infection depends on several factors. Chemotactic stimuli like CXCL1, CXCL2, CCL2 and CCL7 can trigger neutrophil chemotaxis under resting or activated states (Souto et al., 2011). Each of the four chemokines was upregulated to a varying degree upon *Mtb* infection. In addition, IL-17 is also implicated in neutrophil recruitment across different settings (Flannigan et al., 2017; Gelderblom et al., 2012). In TB, IL-17 from CD4[+] T cells is believed to trigger neutrophil infiltration (Nandi and Behar, 2011). Interestingly, IFNγ can block IL-17 production and, therefore, serves as the mechanism for high neutrophilia in *IFNγ* [-/-] mice upon *Mtb* infection (Nandi and Behar, 2011). We also observed very high lung neutrophil infiltration and lung IL-17 levels in *Mtb*-infected *IFNγ* [-/-] mice. However, instead of Th$_{17}$ cells, our results show that neutrophils are a major source of IL-17 in *IFNγ* [-/-] mice. Since neutrophils also express IL-17 receptors, it appears to establish an autocrine self-amplification loop contributing to severe disease pathology. Surprisingly, COX2 inhibition, which helps control *Mtb* in the WT mice, fails to do the same in *IFNγ* [-/-] mice unless the high IL-17 levels were also neutralized. The crosstalk

between prostaglandins and IL-17 is reported as they help Th$_{17}$ cell differentiation

(Napolitani et al., 2009). However, our results show that IL-17, in addition to being produced

by Th$_{17}$ cells, was also produced by neutrophils in *IFNγ$^{-/-}$* mice and therefore, COX2

inhibition could have an independent mechanism through which it synergized with IL-17

neutralization to control *Mtb*. Neutrophils are known to express COX2 under inflammatory

conditions and secrete PGE2 (Maloney et al., 1998). Also, PGE2 can regulate neutrophil

migration, specifically when induced by IL-23/IL-17 (Lemos et al., 2009). However, we

suspect PGE2 also somehow curtail the antibacterial responses of neutrophils since we did

observe conditions where without a drop in neutrophil count, the intra-PMN bacillary load

reduced significantly upon COX2 inhibition.

Inhibiting RORγt with SR2211 provided further insights into the crosstalk between IL-17 and

COX2. RORγt is a classic nuclear receptor implicated with the Th$_{17}$ cell differentiation

(Ivanov et al., 2006). Neutrophils express RORγt and IL-17 and can be activated through

autocrine IL-17A-IL-17RC interactions during fungal infections (Taylor et al., 2014).

However, the IL-17-mediated autocrine activation of neutrophils is believed to elicit strong

anti-microbial functions in the neutrophils (Taylor et al., 2014). While our data does indicate

a potential autocrine IL-17 signaling in the neutrophils of both *Mtb*-infected WT or *IFNγ$^{-/-}$*

mice, it does not appear to induce an antibacterial state but rather facilitates bacterial survival

and pathology. Inhibiting RORγt almost entirely wiped-out IL-17 levels in the lung

homogenates, and, to a large extent, it appeared to impact via inhibiting IL-17 production by

the neutrophils. Moreover, similar to the results obtained with IL-17 neutralization and COX2

inhibition in *IFNγ$^{-/-}$* mice, inhibiting both RORγt and COX2 worked even in WT mice,

leading to significant control of bacterial load in the sorted cells as well as the whole lung and

spleen. Thus overall, IL-17-PMN crosstalk appears to restrain the protective host responses in

normal as well as highly susceptible models of TB. Finally, results with RORγt inhibition in

BCG-vaccinated animals suggest potential ways to enhance the efficacy of TB vaccines.

Whether a prophylactic regimen involving RORγt inhibition to high-risk individuals could

protect them from active disease is worth exploring. The results in this study also

inadvertently provided evidence for an anecdote, which is not usually proven experimentally

in TB disease. In all conditions where the neutrophil population or resident PMN were

targeted, either by celecoxib, IL-17 neutralization or SR2211 treatment, irrespective of the

impact on the bacillary load in the lungs, dissemination to the spleen was significantly

curtailed. This is in agreement with the reports that neutrophils play a direct role in the

dissemination of *Mtb* to other extrapulmonary sites (Hult et al., 2021).

Finally, the results on the human cohort re-emphasize the primacy of neutrophils in

determining the disease outcome in humans. We found that high neutrophils and IL-17 can

predict treatment failure. This underscores a similar neutrophil-IL-17 axis operational in

humans underlying disease severity, as reported in the mouse model in this study.

Interestingly, a neutrophil-driven transcriptional signature was earlier reported to be

associated with active TB (Berry et al., 2010). It is noteworthy that high neutrophilia alone

has been shown to predict death in TB patients (Lowe et al., 2013).

Finally, the results from mice lacking IFNγ have immediate clinical application. A relatively

small fraction, but a sizeable number of TB cases, belong to a category known as Mendelian

Susceptibility to Tubercular Diseases or MSMD, which typically is caused by a mutation in

the IFNγ/IL12p70 pathway, which also acts via regulating IFNγ (Filipe-Santos et al., 2006;

Kerner et al., 2020; Levin et al., 1995). These patients develop serious diseases even upon

exposure to BCG and respond poorly to anti-TB treatment (Casanova et al., 1995;

Mahdaviani et al., 2022). We believe the MSMD phenotype could be a manifestation of the

IL-17-PMN axis observed in *IFNγ $^{-/-}$* animals. Thus, IL-17 neutralization or RORγt inhibition

could facilitate better treatment outcomes in these patients.

To conclude, through this study, we experimentally establish PMNs as a niche for *Mtb* under

normal and severe disease conditions, which is largely regulated by PMN-derived IL-17 as

well as the COX2 pathway. Targeted inhibition of IL-17 and COX2 can be exploited to

enhance the efficacy of BCG or other vaccine candidates and as adjunct therapeutic strategies

for the normal population and among those with genetic vulnerability towards severe TB.

Materials and Methods:

Ethical clearance

ITB biopsy samples and control samples were obtained from the Department of Pathology, AIIMS, New Delhi. These biopsy samples were taken from the patients for diagnostic purposes after taking their written informed consent. The use of the archived leftover biopsy samples was approved by the institute's EC of AIIMS (Ref no. IEC-304/02-06-2017), and ICGEB (Ref no. ICGEB/IEC/2017/06-verII and ICGEB/IEC/2016/03) and the same were accessed according to the institutional guidelines. Animal experiments were approved by Institutional Animal Ethics Committee, ICGEB (ICGEB/IAEC/280718/CI-14), (ICGEB/IAEC/07032020/CI-18), and (ICGEB/IAEC/18092021/CI-16). The study on human PTB subjects was approved by the Ethics Committees of the Prof. M. Viswanathan Diabetes Research Center (ECR/51/INST/TN/ 2013/MVDRC/01) and NIRT (NIRT-INo:2014004). Informed written consent was obtained from all participants. All the methods were performed in accordance with the relevant institutional ethical committee guidelines.

Reagents and antibodies

Tween-20 (P1379), Vectashield (F6182), Triton-X-100 (9036-19-5), DNase 1 (D5025), Collagenase D (11088858001), Propidium Iodide (P4170), HBSS (H6648), Trizol (T9424), Hematoxylin (51275), Eosin (318906), Formalin (HT501128), Celecoxib (SML3031-50MG), DAPI (28718-90-3), Xylene (CAS-1330-20-7), Absolute ethanol (CAS- 64-17-5), Sodium Citrate (CAS--6132-04-3) were obtained from Sigma-Aldrich. SR 2211, Selective RORγt inverse agonist (4869) was procured from TOCRIS. 7H9 (271310), 7H11 (212203), ADC (211887), OADC (211886) FACS Flow Sheath Fluid (342003), Accudrop Fluorescent Beads (345249), CS&T Research Beads Kit (655050), all FACS antibodies used for acquisition and sorting (details are provided in the separate table) were procured from Becton Dickinson

Private Limited. Fixable Viability Dye eFluor™ 780 used for live/dead staining was obtained from eBioscience™. InVivoMAb anti-mouse IL-17A (BE0173), InVivoPure pH 7.0 Dilution Buffer (IP0070), InVivoMAb mouse IgG1 isotype control, unknown specificity (BE0083), InVivoPure pH 6.5 Dilution Buffer (IP0065) were purchased from BioXcell. Mouse IL-17A ELISA Kit (ab199081), mouse PGE2 ELISA Kit (ab287802and Cytokine & Chemokine 36-Plex Mouse ProcartaPlexTM Panel 1A (EPX360-26092-901) for Luminex assay were obtained from Abcam and Thermofisher respectively. Penta antibiotic mix (FD260-5VL) was procured from Himedia.

Mycobacterial cultures

Bacterial strains: Lab-adapted strain of *Mycobacterium tuberculosis*, H37Rv, seed stock was received from Colorado State University. BCG (Pasteur strain) was obtained from the National Institute of Immunology, New Delhi. For *Mtb* ex-vivo and in-vivo infection experiments, H37Rv and BCG mycobacterial cultures were grown in 7H9 media (BD Difco) supplemented with 10% Albumin–Dextrose–Catalase (ADC, BD, Difco) and incubated in an orbiter shaker at 100 rpm, 37 °C till the growth reach to the log phase. The log phase cultures were pelleted down, and the pellet was resuspended in either saline (for aerosol infection in mice) or culture media (for ex-vivo infection in cells). Single-cell suspensions of H37Rv and BCG were prepared by passing bacterial cultures seven times through 23-gauge needles, five times through 26-gauge needles, and three times through 30-gauge needles.

Animals

WT C57BL/6 mice were procured from animal house, ICGEB, New Delhi and *IFNγ$^{-/-}$* C57BL/6 mice were procured from animal house, NII, New Delhi. Both male and female, six to eight weeks old mice were used for all the in vivo experiments. *Mtb*-infected mice were housed in our biosafety level-3 laboratory and kept pathogen-free by routine serological and

histopathological examinations. Animal protocols employed in this study were approved by the Institutional Animal Ethics Committee.

Aerosol challenge in C57BL/6 mice with H37Rv strain

All mice experiments were carried out in the Tuberculosis Aerosol Challenge Facility (TACF, ICGEB, New Delhi, India). C57BL/6 mice were housed in cages within a biosafety level 3 laminar flow enclosure. According to the standardized protocol, mice were infected with 100-200 CFU (unless otherwise stated) with H37Rv in a Wisconsin-Madison chamber. To confirm the infection establishment and to enumerate the exact CFU delivered, two animals were selected randomly and humanely euthanized 24 hours post-aerosol challenge. The lung tissues were harvested and homogenized in 1X PBS, followed by plating on Petri plates containing Middlebrook 7H11 agar (Difco) supplemented with 10% OADC (Becton, Dickinson) 0.5% glycerol. The colonies were counted after 21 days to enumerate the *Mtb* load.

Drug treatment and BCG vaccination of mice

Cox-2 inhibitor Celecoxib (50mg/kg) was administered to the mice through oral gavaging once per day, according to the timeline of the experiment. RORγt inhibitor SR2211 (20mg/kg) was administered to the mice through intra-peritoneal route every alternate day according to the timeline of the experiment. Mice were vaccinated with BCG intradermally at a dose of 1 million bacilli per mouse.

CFU plating post *Mtb*-Infection

At various time-points post *Mtb*-infection, lung, and spleen tissues were harvested from the infected mice. The left half lung of the mice was homogenized and serial diluted in 1X PBS. Different dilutions were plated in duplicates on 7H11 agar plates supplemented with 10% OADC to get *Mtb* colonies within the countable range. Similarly, spleens were also harvested,

homogenized, and serially diluted in 1X PBS, followed by plating on 7H11 agar plates. The

colonies were counted after 21 days to enumerate the *Mtb* load.

FACS sample preparation

The right half lung of the infected mice was used in the preparation of single cell suspension

followed by FACS sorting and subsequent plating of sorted cells (i.e., Neutrophils,

Macrophages, and Mesenchymal stem cells. For sorting, the tissue was washed with 1X PBS

and chopped into small pieces. 20U/ml DNase 1 and 1mg/ml collagenase D were added to the

chopped tissue and incubated for 30-45 minutes at 37ºC. After incubation, the chopped lung

tissues were passed through the nylon mesh of a 70-micron cell strainer to get a single-cell

suspension. The cells were pelleted at 1200 rpm, and the cell pellet was treated with 1X ACK

RBC lysis buffer to get rid of RBCs and washed with 1X PBS. The cells were incubated with

Fc-Block antibody (CD16/32 FCR-γ antibody) on ice for 15 minutes to inhibit the binding of

surface antibodies to Fc receptors on the immune cell surface. The cells were washed with PBS

and subsequentially stained with the antibody cocktail of CD45.2, CD11b, Ly6G, MHCII,

CD64, MerTK, Sca-1, CD90.1, and CD73 and placed on ice for 45 minutes. Dead cells were

stained with fixable viability dye eFluor^{TM} 780 (FVD eFluor 780) or Propidium Iodide (PI)at

five µg/ml concentration just before acquisition.

FACS cell sorting

After staining and washing, the lung cells were again passed through a 40-micron strainer

before acquisition into the FACS machine to ensure formation of single-cell suspension. The

whole sample was acquired to get the maximum number of live sorted cells (Neutrophils,

Macrophages, and Mesenchymal stem cells). The sorted cells were then pelleted (at 13000

RPM), lysed, and plated on 7H11 plates. The colonies were counted after 21 days to enumerate

the *Mtb* burden within each sorted cell type.

RNA-Seq analysis

Reads were quality filtered using Trimmomatic (v.0.39, LEADING:10 TRAILING:10

SLIDING WINDOW:4:15 MINLEN:75) (Bolger et al., 2014) and were assessed for quality

using FastQC (v.10.1) (Andrews, 2010) and visualised with MultiQC (v.1.9) (Ewels et al.,

2016). Salmon (v.1.8, –numGibbsSamples 30) (Patro et al., 2017) was used to quantify quality

filtered reads against mouse genome (GRCm39, primary assembly) with Gencode annotation

(v.M30) (Frankish et al., 2019). Profiles were imported using tximeta (v.1.10) (Love et al.,

2020) and SummarizedExperiment (v.1.22) (Morgan et al., 2018) which were then summarised

to gene-level expression. Following biotypes were filtered out: pseudogenes, TEC, snRNA,

lncRNA, miRNA, snoRNA, misc_RNA, rRNA, ribozyme, scaRNA, sRNA, scRNA,

Mt_tRNA, Mt_rRNA. Dataset was further filtered to remove features not expressed across the

whole dataset. The dataset was normalised to scaled TPM values. Downstream analyses are

described in the statistical analyses section.

<u>Multiplex Microbead-based Immunoassay (Luminex)</u>

Mouse lung homogenates were collected for Luminex analysis using the Cytokine &

Chemokine 36-Plex Mouse ProcartaPlexTM Panel 1A (ThermoFisher, # EPX360-26092-

901). Samples were incubated with fluorescent-coded magnetic beads pre-coated with

respective antibodies in a black 96-well clear-bottom plate overnight at 4 °C. After

incubation, plates were washed 5 times with wash buffer (PBS with 1% BSA (Capricorn

Scientific) and 0.05% Tween-20 (Promega)). Sample- antibody-bead complexes were

incubated with Biotinylated detection antibodies for 1h and washed 5 times. Subsequently,

Streptavidin-PE was added and incubated for another 30 mins. Plates were washed 5 times,

before sample-antibody-bead complexes were re-suspended in sheath fluid for acquisition on

the FLEXMAP® 3D (Luminex) using xPONENT® 4.0 (Luminex) software. Data analysis

was done using Bio-Plex ManagerTM 6.1.1 (Bio-Rad). Standard curves were generated with

a 5-PL (5-parameter logistic) algorithm, reporting values for both mean florescence intensity

(MFI) and concentration data.

Tissue Immunofluorescence Assay (IFA)

Five-micron thick sections of paraffin-embedded tissue sections were taken in poly-L-Lysine

coated slide. Deparaffinization was performed by heating the slides at 50 °C for 20 seconds

till the wax melts, followed by the subsequent steps, 100% xylene for 10 minutes, xylene and

absolute ethanol for 10 minutes, 100% ethanol for 10 minutes, 70% percent ethanol for 5

minutes, 50% ethanol, distilled water for 5 minutes and a final wash in 1x PBS for 5 minutes.

Antigen retrieval was performed in antigen retrieval buffer (10mM Sodium Citrate, 0.05%

tween-20, pH: 6) by heating the slides at 100 °C for 15 minutes. After antigen retrieval,

permeabilization was performed with 0.04% Triton-X 100 in 1X PBS (0.04% PBST) for 20

minutes followed by proper washing with 1X PBS. Blocking was done by 3% BSA for 1

hour. Sequential addition of primary antibody was performed at 1:100 dilution at 4 °C

overnight. Primary antibody was washed with 1X PBS followed by counter stain with DAPI

nuclear stain at 1 ug/ml concentration. Mounting was done with a drop of vectashield.

Histopathological studies

Formalin fixed tissue samples (<5mm thick) were washed under running water for 4-12

hours. Tissue dehydration was performed in ascending grades of alcohol (30%, 50%, 70%,

90%, 95%) followed by two washes in absolute alcohol, one hour each and two washes in

xylene for one hour each. Paraffinization in melted paraffin wax (57-60 °C) was performed

three times sequentially for one hour each. FFPE (formalin fixed paraffin embedded) tissue

blocks were prepared by placing tissue and melted paraffin in steel moulds. Five micrometres

thick sections of tissue were cut with the help of microtome. Tissue sections were allowed to

float on water bath (45-50 °C) and placed on clean glass slide. Deparaffinization was

performed by three sequential washes in xylene for 5 minutes each followed by hydration in

descending grades of alcohol (absolute alcohol, 90% and 70%) followed by one wash in

water, 5 minutes each. For H&E staining, slides were first stained with haematoxylin (it

stains the nucleus blue) for 5-10 minutes followed by one wash with water. The slides were

subsequently stained with Eosin (it stains the cytoplasm pink) for 5-10 minutes followed by

one wash with absolute alcohol and one wash with xylene. Slides were mounted with

coverslips followed by sealing with DPX. Images were acquired on Zeiss microscope.

In vivo IL-17 Neutralization experiment

IFNγ knock out C57BL/6 mice were challenged with 100-200 CFU with H37Rv via the aerosol

route (as described previously). The anti-mouse IL-17 monoclonal Ab was administered via

intra-peritoneal route at a dose of 100 milligrams per mouse every 3–4 days starting one day

before *M. tuberculosis* infection for the duration of the study, i.e., six weeks. Control groups

received matched IgG isotypes. Six weeks post *Mtb* infection, mice were treated with 50mg/kg

celecoxib through oral route for the duration of the experiments. Mice in all experimental

groups were sacrificed at 6 weeks post infection, according to the experimental timeline. The

left half lungs and whole spleen of the mice were homogenized and serially diluted in 1X PBS.

Different dilutions were plated in duplicates on 7H11 agar plates and the colonies were counted

after 21 days to enumerate the *Mtb* load. The middle lobe of the right lung was preserved in

formalin for Histopathology and IFA studies. The rest part of the right half lung of the infected

mice was used in the preparation of single cell suspension. The single cell suspension was

divided into two parts and stained with different antibody cocktails. One part was used for

FACS sorting and subsequent plating of sorted cells (i.e., Neutrophils, Macrophages, and

Mesenchymal stem cells) and the other half was used for FACS acquisition to enumerate Th17

cells inside the lungs. Sera was also collected to quantify IL17 levels.

IL-17/PGE2 ELISA

IL-17 levels and PGE2 levels in the lung homogenates of uninfected mice, *Mtb*-infected mice,

Mtb-infected treated mice from various experimental groups were measured using ELISA kits

procured from Abcam using the manufacturer's protocol.

Study Population (PTB subjects)

Participants were enrolled from the Effect of Diabetes on Tuberculosis Severity (EDOTS)

study, a prospective cohort study conducted in Chennai, India. A total of 68 had unfavourable

treatment outcomes, comprising treatment failure, relapse, or death. Inclusion criterion was

new smear and culture-positive adults (age 18-75 years). Exclusion criteria were previous

treatment for TB, drug resistant TB, HIV positivity or taking immunosuppressive drugs, and

>7 days of anti-TB treatment prior to enrolment. As part of our study protocol HIV screening

was done for all the study participants. The diagnosis of PTB was established by positive

sputum culture on solid media with compatible chest X-ray. Anti-TB treatment (ATT) was

provided by government TB clinics according the Revised National Tuberculosis Control

Programme standards in effect at that time. Participants were followed up monthly through the

six months course of treatment and every three months thereafter till one year after treatment

completion Cure was defined as negative sputum cultures at months 5 and 6 of treatment.

Adverse treatment outcomes included treatment failure defined as positive sputum culture at

months 5 or 6, all-cause mortality over the duration of treatment, or recurrent TB demonstrated

by positive sputum smear and culture with compatible symptoms or chest X-ray within twelve

months after initial cure. There was a total of 18 TB treatment failures, 16 deaths and 34 TB

recurrences. Case-control matching was based on age, gender, body mass index (BMI), and

diabetic status. Peripheral blood was collected in heparinized tubes. Following centrifugation,

plasma was collected and stored at -80 $^{\circ}$C till further analysis. Baseline sample collection was

performed at enrolment. The demographic and epidemiological data for this cohort have been

previously reported (Kumar et al., 2021; Kumar et al., 2020).

Multiplex assays

689 Circulating plasma levels of cytokines and chemokines were measured using the Luminex

690 MAGPIX Multiplex Assay system (Bio-Rad, Hercules, CA). Luminex Human Magnetic Assay

691 Kit 45 Plex (R&D Systems) was used to measure the cytokines and chemokine levels

Statistical analysis

Transcriptomics analyses were done in R statistical software environment (v.4.1) (Team,

2021). Statistical analysis was done using swish methodology from fishpond (v.1.8) (Zhu et

al., 2019) and significantly differentially expressed genes (DEGs) were defined based on

qvalue $\leq$ 0.1 and absolute $\log_2$ fold change $\geq$ 1. Heatmap of the DEGs was generated with

ComplexHeatmap (v.2.8) using mean z-scaled values and clustered based on Euclidean

distance and Ward.D2 methodology. Enrichment analysis was done using clusterProfiler (v.

4.0.5) (Wu et al., 2021) and simplify function was used to reduce the GO BP terms at a

similarity cutoff of 0.5 as applicable. Significance of enrichment defined by nominal p-value

$\leq$ 0.05 unless stated otherwise. Other figures from RNASeq analyses were generated using

ggplot2 (v.3.3.5) (Wickham, 2016).

Data availability

The raw RNAseq data used for this study is available at the GEO database with the accession

number: **GSE238265.**

Funding: The work on this project was supported by Department of Biotechnology (BT/IC-

06/003/91-Flagship program to DK), MDR-TB project from Department of Biotechnology

(BT/PR23092/NER/95/589/2017, DK) and SR-National Bioscience Award from Department

of Biotechnology (BT/HRD/NBA-NWB/39/2020-21 (4), DK). The research in DK lab is also

supported by DBT-Wellcome Trust India Alliance Senior Fellowship (IA/S/17/1/503071) and

extramural research grants from SERB (EMR/2016/005296 to DK). Priya Sharma is a

recipient of a fellowship from the Indian Council of Medical Research (3/1/3/JRF-

2017/HRD-LS/55193/79). Raman Deep Sharma is a recipient of a fellowship from

Department of Biotechnology (DBT/JRF/BET-18/I/2018/AL/160).

717   Acknowledgements

We thank the Tuberculosis Aerosol Challenge Facility (TACF), supported by the Department

of Biotechnology, Govt of India, for supporting the BSL3 facility. We thank the staff of the

Department of Clinical Research and the Department of Bacteriology, NIRT for valuable

assistance in bacterial cultures and radiology and the staff of Prof. M. Viswanathan Diabetes

Research Center, RNTCP and Chennai corporation for valuable assistance in recruiting the

patients for this study. Data in this manuscript were collected as part of the Regional

Prospective Observational Research for Tuberculosis (RePORT) India Consortium.

725
726 **References:**
727

Allen, J.E., T.E. Sutherland, and D. Ruckerl. 2015. IL-17 and neutrophils: unexpected players
in the type 2 immune response. *Curr Opin Immunol* 34:99-106.
Andrews, S. 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data. In.
Ballinger, M.N., D.M. Aronoff, T.R. McMillan, K.R. Cooke, K. Olkiewicz, G.B. Toews, M.
Peters-Golden, and B.B. Moore. 2006. Critical role of prostaglandin E2
overproduction in impaired pulmonary host response following bone marrow
transplantation. *J Immunol* 177:5499-5508.
Barry, C.E., 3rd, H.I. Boshoff, V. Dartois, T. Dick, S. Ehrt, J. Flynn, D. Schnappinger, R.J.
Wilkinson, and D. Young. 2009. The spectrum of latent tuberculosis: rethinking the
biology and intervention strategies. *Nat Rev Microbiol* 7:845-855.
Basile, J.I., L.J. Geffner, M.M. Romero, L. Balboa, Y.G.C. Sabio, V. Ritacco, A. Garcia, M.
Cuffre, E. Abbate, B. Lopez, L. Barrera, M. Ambroggi, M. Aleman, M.C. Sasiain, and
S.S. de la Barrera. 2011. Outbreaks of mycobacterium tuberculosis MDR strains
induce high IL-17 T-cell response in patients with MDR tuberculosis that is closely
associated with high antigen load. *J Infect Dis* 204:1054-1064.
Berry, M.P., C.M. Graham, F.W. McNab, Z. Xu, S.A. Bloch, T. Oni, K.A. Wilkinson, R.
Banchereau, J. Skinner, R.J. Wilkinson, C. Quinn, D. Blankenship, R. Dhawan, J.J. Cush,
A. Mejias, O. Ramilo, O.M. Kon, V. Pascual, J. Banchereau, D. Chaussabel, and A.
O'Garra. 2010. An interferon-inducible neutrophil-driven blood transcriptional
signature in human tuberculosis. *Nature* 466:973-977.
Bolger, A.M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30:2114-2120.
Brazier, B., and H. McShane. 2020. Towards new TB vaccines. *Semin Immunopathol* 42:315-
331.
Cadena, A.M., S.M. Fortune, and J.L. Flynn. 2017. Heterogeneity in tuberculosis. *Nat Rev*
Immunol 17:691-702.
Casanova, J.L., E. Jouanguy, S. Lamhamedi, S. Blanche, and A. Fischer. 1995. Immunological
conditions of children with BCG disseminated infection. *Lancet* 346:581.
Chandra, P., S.J. Grigsby, and J.A. Philips. 2022. Immune evasion and provocation by
Mycobacterium tuberculosis. *Nat Rev Microbiol* 20:750-766.
Chaves Torres, N.M., J.J. Quijano Rodriguez, P.S. Porras Andrade, M.B. Arriaga, and E.M.
Netto. 2019. Factors predictive of the success of tuberculosis treatment: A
systematic review with meta-analysis. *PLoS One* 14:e0226507.
Cooper, A.M., D.K. Dalton, T.A. Stewart, J.P. Griffin, D.G. Russell, and I.M. Orme. 1993.
Disseminated tuberculosis in interferon gamma gene-disrupted mice. *The Journal of*
experimental medicine 178:2243-2247.
Cruz, A., A.G. Fraga, J.J. Fountain, J. Rangel-Moreno, E. Torrado, M. Saraiva, D.R. Pereira, T.D.
Randall, J. Pedrosa, A.M. Cooper, and A.G. Castro. 2010. Pathological role of
interleukin 17 in mice subjected to repeated BCG vaccination after infection with
Mycobacterium tuberculosis. *The Journal of experimental medicine* 207:1609-1616.
Dallenga, T., and U.E. Schaible. 2016. Neutrophils in tuberculosis--first line of defence or
booster of disease and targets for host-directed therapy? *Pathog Dis* 74:
Dartois, V.A., and E.J. Rubin. 2022. Anti-tuberculosis treatment strategies and drug
development: challenges and priorities. *Nat Rev Microbiol* 20:685-701.

Das, B., S.S. Kashino, I. Pulu, D. Kalita, V. Swami, H. Yeger, D.W. Felsher, and A. Campos-
Neto. 2013. CD271(+) bone marrow mesenchymal stem cells may provide a niche for
dormant Mycobacterium tuberculosis. *Sci Transl Med* 5:170ra113.
Domingo-Gonzalez, R., S. Das, K.L. Griffiths, M. Ahmed, M. Bambouskova, R. Gopal, S. Gondi,
M. Munoz-Torrico, M.A. Salazar-Lezama, A. Cruz-Lagunas, L. Jimenez-Alvarez, G.
Ramirez-Martinez, R. Espinosa-Soto, T. Sultana, J. Lyons-Weiler, T.A. Reinhart, J.
Arcos, M. de la Luz Garcia-Hernandez, M.A. Mastrangelo, N. Al-Hammadi, R.
Townsend, J.M. Balada-Llasat, J.B. Torrelles, G. Kaplan, W. Horne, J.K. Kolls, M.N.
Artyomov, J. Rangel-Moreno, J. Zuniga, and S.A. Khader. 2017. Interleukin-17 limits
hypoxia-inducible factor 1alpha and development of hypoxic granulomas during
tuberculosis. *JCI Insight* 2:
Eum, S.Y., J.H. Kong, M.S. Hong, Y.J. Lee, J.H. Kim, S.H. Hwang, S.N. Cho, L.E. Via, and C.E.
Barry, 3rd. 2010. Neutrophils are the predominant infected phagocytic cells in the
airways of patients with active pulmonary TB. *Chest* 137:122-128.
Ewels, P., M. Magnusson, S. Lundin, and M. Kaller. 2016. MultiQC: summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics* 32:3047-
3048.
Fatima, S., S.S. Kamble, V.P. Dwivedi, D. Bhattacharya, S. Kumar, A. Ranganathan, L. Van
Kaer, S. Mohanty, and G. Das. 2020. Mycobacterium tuberculosis programs
mesenchymal stem cells to establish dormancy and persistence. *J Clin Invest*
130:655-661.
Filipe-Santos, O., J. Bustamante, A. Chapgier, G. Vogt, L. de Beaucoudrey, J. Feinberg, E.
Jouanguy, S. Boisson-Dupuis, C. Fieschi, C. Picard, and J.L. Casanova. 2006. Inborn
errors of IL-12/23- and IFN-gamma-mediated immunity: molecular, cellular, and
clinical features. *Semin Immunol* 18:347-361.
Flannigan, K.L., V.L. Ngo, D. Geem, A. Harusato, S.A. Hirota, C.A. Parkos, N.W. Lukacs, A.
Nusrat, V. Gaboriau-Routhiau, N. Cerf-Bensussan, A.T. Gewirtz, and T.L. Denning.
2017. IL-17A-mediated neutrophil recruitment limits expansion of segmented
filamentous bacteria. *Mucosal Immunol* 10:673-684.
Flynn, J.L., J. Chan, K.J. Triebold, D.K. Dalton, T.A. Stewart, and B.R. Bloom. 1993. An
essential role for interferon gamma in resistance to Mycobacterium tuberculosis
infection. *The Journal of experimental medicine* 178:2249-2254.
Frankish, A., M. Diekhans, A.M. Ferreira, R. Johnson, I. Jungreis, J. Loveland, J.M. Mudge, C.
Sisu, J. Wright, J. Armstrong, I. Barnes, A. Berry, A. Bignell, S. Carbonell Sala, J. Chrast,
F. Cunningham, T. Di Domenico, S. Donaldson, I.T. Fiddes, C. Garcia Giron, J.M.
Gonzalez, T. Grego, M. Hardy, T. Hourlier, T. Hunt, O.G. Izuogu, J. Lagarde, F.J.
Martin, L. Martinez, S. Mohanan, P. Muir, F.C.P. Navarro, A. Parker, B. Pei, F. Pozo,
M. Ruffier, B.M. Schmitt, E. Stapleton, M.M. Suner, I. Sycheva, B. Uszczynska-
Ratajczak, J. Xu, A. Yates, D. Zerbino, Y. Zhang, B. Aken, J.S. Choudhary, M. Gerstein,
R. Guigo, T.J.P. Hubbard, M. Kellis, B. Paten, A. Reymond, M.L. Tress, and P. Flicek.
2019. GENCODE reference annotation for the human and mouse genomes. *Nucleic*
Acids Res 47:D766-D773.
Funk, C.D. 2001. Prostaglandins and leukotrienes: advances in eicosanoid biology. *Science*
294:1871-1875.
Gelderblom, M., A. Weymar, C. Bernreuther, J. Velden, P. Arunachalam, K. Steinbach, E.
Orthey, T.V. Arumugam, F. Leypoldt, O. Simova, V. Thom, M.A. Friese, I. Prinz, C.
Holscher, M. Glatzel, T. Korn, C. Gerloff, E. Tolosa, and T. Magnus. 2012.

819   Neutralization of the IL-17 axis diminishes neutrophil invasion and protects from
820   ischemic stroke. *Blood* 120:3793-3802.

821   Han, Y., S.J. Kim, S.H. Lee, Y.S. Sim, Y.J. Ryu, J.H. Chang, S.S. Shim, Y. Kim, and J.H. Lee. 2018.
822   High blood neutrophil-lymphocyte ratio associated with poor outcomes in miliary
823   tuberculosis. *J Thorac Dis* 10:339-346.

824   Heyckendorf, J., S. Marwitz, M. Reimann, K. Avsar, A.R. DiNardo, G. Gunther, M. Hoelscher,
825   E. Ibraim, B. Kalsdorf, S.H.E. Kaufmann, I. Kontsevaya, F. van Leth, A.M. Mandalakas,
826   F.P. Maurer, M. Muller, D. Nitschkowski, I.D. Olaru, C. Popa, A. Rachow, T. Rolling, J.
827   Rybniker, H.J.F. Salzer, P. Sanchez-Carballo, M. Schuhmann, D. Schaub, V. Spinu, I.
828   Suarez, E. Terhalle, M. Unnewehr, J. Weiner, 3rd, T. Goldmann, and C. Lange. 2021.
829   Prediction of anti-tuberculosis treatment duration based on a 22-gene
830   transcriptomic model. *Eur Respir J* 58:

831   Huang, L., E.V. Nazarova, S. Tan, Y. Liu, and D.G. Russell. 2018. Growth of Mycobacterium
832   tuberculosis in vivo segregates with host macrophage metabolism and ontogeny. *The*
833   *Journal of experimental medicine* 215:1135-1152.

834   Hult, C., J.T. Mattila, H.P. Gideon, J.J. Linderman, and D.E. Kirschner. 2021. Neutrophil
835   Dynamics Affect Mycobacterium tuberculosis Granuloma Outcomes and
836   Dissemination. *Front Immunol* 12:712457.

837   Ibrahim, O.M., P.H. Basse, W. Jiang, K. Guru, G. Chatta, and P. Kalinski. 2021. NFkappaB-
838   Activated COX2/PGE(2)/EP4 Axis Controls the Magnitude and Selectivity of BCG-
839   Induced Inflammation in Human Bladder Cancer Tissues. *Cancers (Basel)* 13:

840   Ivanov, II, B.S. McKenzie, L. Zhou, C.E. Tadokoro, A. Lepelley, J.J. Lafaille, D.J. Cua, and D.R.
841   Littman. 2006. The orphan nuclear receptor RORgammat directs the differentiation
842   program of proinflammatory IL-17+ T helper cells. *Cell* 126:1121-1133.

843   Jain, N., H. Kalam, L. Singh, V. Sharma, S. Kedia, P. Das, V. Ahuja, and D. Kumar. 2020.
844   Mesenchymal stem cells offer a drug-tolerant and immune-privileged niche to
845   Mycobacterium tuberculosis. *Nat Commun* 11:3062.

846   Jung, B.G., B. Samten, K. Dean, R.J. Wallace, Jr., B.A. Brown-Elliott, T. Tucker, S. Idell, J.V.
847   Philley, and R. Vankayalapati. 2022. Early IL-17A production helps establish
848   Mycobacterium intracellulare infection in mice. *PLoS Pathog* 18:e1010454.

849   Jurado, J.O., V. Pasquinelli, I.B. Alvarez, D. Pena, A.I. Rovetta, N.L. Tateosian, H.E. Romeo,
850   R.M. Musella, D. Palmero, H.E. Chuluyan, and V.E. Garcia. 2012. IL-17 and IFN-
851   gamma expression in lymphocytes from patients with active tuberculosis correlates
852   with the severity of the disease. *J Leukoc Biol* 91:991-1002.

853   Kaufmann, E., J. Sanz, J.L. Dunn, N. Khan, L.E. Mendonca, A. Pacis, F. Tzelepis, E. Pernet, A.
854   Dumaine, J.C. Grenier, F. Mailhot-Leonard, E. Ahmed, J. Belle, R. Besla, B. Mazer, I.L.
855   King, A. Nijnik, C.S. Robbins, L.B. Barreiro, and M. Divangahi. 2018. BCG Educates
856   Hematopoietic Stem Cells to Generate Protective Innate Immunity against
857   Tuberculosis. *Cell* 172:176-190 e119.

858   Kerner, G., J. Rosain, A. Guerin, A. Al-Khabaz, C. Oleaga-Quintas, F. Rapaport, M.J. Massaad,
859   J.Y. Ding, T. Khan, F.A. Ali, M. Rahman, C. Deswarte, R. Martinez-Barricarte, R.S.
860   Geha, V. Jeanne-Julien, D. Garcia, C.Y. Chi, R. Yang, M. Roynard, B. Fleckenstein, F.
861   Rozenberg, S. Boisson-Dupuis, C.L. Ku, Y. Seeleuthner, V. Beziat, N. Marr, L. Abel, W.
862   Al-Herz, J.L. Casanova, and J. Bustamante. 2020. Inherited human IFN-gamma
863   deficiency underlies mycobacterial disease. *J Clin Invest* 130:3158-3171.

Ketelut-Carneiro, N., C.O.S. Souza, L. Benevides, L.G. Gardinassi, M.C. Silva, L.A. Tavares, D.S.
Zamboni, and J.S. Silva. 2019. Caspase-11-dependent IL-1alpha release boosts Th17
immunity against Paracoccidioides brasiliensis. *PLoS Pathog* 15:e1007990.

Kimmey, J.M., J.P. Huynh, L.A. Weiss, S. Park, A. Kambal, J. Debnath, H.W. Virgin, and C.L.
Stallings. 2015. Unique role for ATG5 in neutrophil-mediated immunopathology
during M. tuberculosis infection. *Nature* 528:565-569.

Kolaczkowska, E., and P. Kubes. 2013. Neutrophil recruitment and function in health and
inflammation. *Nat Rev Immunol* 13:159-175.

Kumar, N.P., K. Moideen, A. Nancy, V. Viswanathan, K. Thiruvengadam, D. Nair, V.V.
Banurekha, S. Sivakumar, S. Hissar, H. Kornfeld, and S. Babu. 2021. Plasma
Chemokines Are Baseline Predictors of Unfavorable Treatment Outcomes in
Pulmonary Tuberculosis. *Clin Infect Dis* 73:e3419-e3427.

Kumar, N.P., K. Moideen, A. Nancy, V. Viswanathan, K. Thiruvengadam, S. Sivakumar, S.
Hissar, D. Nair, V.V. Banurekha, H. Kornfeld, and S. Babu. 2020. Association of Plasma
Matrix Metalloproteinase and Tissue Inhibitors of Matrix Metalloproteinase Levels
With Adverse Treatment Outcomes Among Patients With Pulmonary Tuberculosis.
JAMA Netw Open 3:e2027754.

Lee, J., S. Boyce, J. Powers, C. Baer, C.M. Sassetti, and S.M. Behar. 2020. CD11cHi monocyte-
derived macrophages are a major cellular compartment infected by Mycobacterium
tuberculosis. *PLoS Pathog* 16:e1008621.

Lee, W.W., S.W. Kang, J. Choi, S.H. Lee, K. Shah, E.E. Eynon, R.A. Flavell, and I. Kang. 2010.
Regulating human Th17 cells via differential expression of IL-1 receptor. *Blood*
115:530-540.

Leliefeld, P.H.C., J. Pillay, N. Vrisekoop, M. Heeres, T. Tak, M. Kox, S.H.M. Rooijakkers, T.W.
Kuijpers, P. Pickkers, L.P.H. Leenen, and L. Koenderman. 2018. Differential
antibacterial control by neutrophil subsets. *Blood Adv* 2:1344-1355.

Lemos, H.P., R. Grespan, S.M. Vieira, T.M. Cunha, W.A. Verri, Jr., K.S. Fernandes, F.O. Souto,
I.B. McInnes, S.H. Ferreira, F.Y. Liew, and F.Q. Cunha. 2009. Prostaglandin mediates
IL-23/IL-17-induced neutrophil migration in inflammation by inhibiting IL-12 and
IFNgamma production. *Proc Natl Acad Sci U S A* 106:5954-5959.

Levin, M., M.J. Newport, S. D'Souza, P. Kalabalikis, I.N. Brown, H.M. Lenicker, P.V. Agius, E.G.
Davies, A. Thrasher, N. Klein, and et al. 1995. Familial disseminated atypical
mycobacterial infection in childhood: a human mycobacterial susceptibility gene?
Lancet 345:79-83.

Love, M.I., C. Soneson, P.F. Hickey, L.K. Johnson, N.T. Pierce, L. Shepherd, M. Morgan, and R.
Patro. 2020. Tximeta: Reference sequence checksums for provenance identification
in RNA-seq. *PLoS Comput Biol* 16:e1007664.

Lovewell, R.R., C.E. Baer, B.B. Mishra, C.M. Smith, and C.M. Sassetti. 2021. Granulocytes act
as a niche for Mycobacterium tuberculosis growth. *Mucosal Immunol* 14:229-241.

Lowe, D.M., A.K. Bandara, G.E. Packe, R.D. Barker, R.J. Wilkinson, C.J. Griffiths, and A.R.
Martineau. 2013. Neutrophilia independently predicts death in tuberculosis. *Eur*
Respir J 42:1752-1757.

Lowe, D.M., J. Demaret, N. Bangani, J.K. Nakiwala, R. Goliath, K.A. Wilkinson, R.J. Wilkinson,
and A.R. Martineau. 2018. Differential Effect of Viable Versus Necrotic Neutrophils
on Mycobacterium tuberculosis Growth and Cytokine Induction in Whole Blood.
Front Immunol 9:903.

Mahdaviani, S.A., M. Fallahi, M. Jamee, M. Marjani, P. Tabarsi, A. Moniri, P. Farnia, Z.
Daneshmandi, N. Parvaneh, J.L. Casanova, J. Bustamante, D. Mansouri, and A.A.
Velayati. 2022. Effective anti-mycobacterial treatment for BCG disease in patients
with Mendelian Susceptibility to Mycobacterial Disease (MSMD): a case series. *Ann*
Clin Microbiol Antimicrob 21:8.

Maloney, C.G., W.A. Kutchera, K.H. Albertine, T.M. McIntyre, S.M. Prescott, and G.A.
Zimmerman. 1998. Inflammatory agonists induce cyclooxygenase type 2 expression
by human neutrophils. *J Immunol* 160:1402-1410.

Martineau, A.R., S.M. Newton, K.A. Wilkinson, B. Kampmann, B.M. Hall, N. Nawroly, G.E.
Packe, R.N. Davidson, C.J. Griffiths, and R.J. Wilkinson. 2007. Neutrophil-mediated
innate immune resistance to mycobacteria. *J Clin Invest* 117:1988-1994.

Mills, K.H.G. 2023. IL-17 and IL-17-producing cells in protection versus pathology. *Nat Rev*
Immunol 23:38-54.

Mishra, B.B., R.R. Lovewell, A.J. Olive, G. Zhang, W. Wang, E. Eugenin, C.M. Smith, J.Y.
Phuah, J.E. Long, M.L. Dubuke, S.G. Palace, J.D. Goguen, R.E. Baker, S. Nambi, R.
Mishra, M.G. Booty, C.E. Baer, S.A. Shaffer, V. Dartois, B.A. McCormick, X. Chen, and
C.M. Sassetti. 2017. Nitric oxide prevents a pathogen-permissive granulocytic
inflammation during tuberculosis. *Nat Microbiol* 2:17072.

Moorlag, S., Y.A. Rodriguez-Rosales, J. Gillard, S. Fanucchi, K. Theunissen, B. Novakovic, C.M.
de Bont, Y. Negishi, E.T. Fok, L. Kalafati, P. Verginis, V.P. Mourits, V. Koeken, L.C.J. de
Bree, G.J.M. Pruijn, C. Fenwick, R. van Crevel, L.A.B. Joosten, I. Joosten, H. Koenen,
M.M. Mhlanga, D.A. Diavatopoulos, T. Chavakis, and M.G. Netea. 2020. BCG
Vaccination Induces Long-Term Functional Reprogramming of Human Neutrophils.
Cell Rep 33:108387.

Morgan, M., V. Obenchain, J. Hester, and H. Pagès. 2018. SummarizedExperiment:
SummarizedExperiment container. In.

Nandi, B., and S.M. Behar. 2011. Regulation of neutrophils by interferon-gamma limits lung
inflammation during tuberculosis infection. *The Journal of experimental medicine*
208:2251-2262.

Napolitani, G., E.V. Acosta-Rodriguez, A. Lanzavecchia, and F. Sallusto. 2009. Prostaglandin
E2 enhances Th17 responses via modulation of IL-17 and IFN-gamma production by
memory CD4+ T cells. *Eur J Immunol* 39:1301-1312.

Nwongbouwoh Muefong, C., O. Owolabi, S. Donkor, S. Charalambous, A. Bakuli, A. Rachow,
C. Geldmacher, and J.S. Sutherland. 2022. Neutrophils Contribute to Severity of
Tuberculosis Pathology and Recovery From Lung Damage Pre- and Posttreatment.
Clin Infect Dis 74:1757-1766.

Pai, M., M.A. Behr, D. Dowdy, K. Dheda, M. Divangahi, C.C. Boehme, A. Ginsberg, S.
Swaminathan, M. Spigelman, H. Getahun, D. Menzies, and M. Raviglione. 2016.
Tuberculosis. *Nat Rev Dis Primers* 2:16076.

Patro, R., G. Duggal, M.I. Love, R.A. Irizarry, and C. Kingsford. 2017. Salmon provides fast and
bias-aware quantification of transcript expression. *Nat Methods* 14:417-419.

Pavan Kumar, N., K. Moideen, A. Nancy, V. Viswanathan, B.S. Shruthi, S. Shanmugam, S.
Hissar, H. Kornfeld, and S. Babu. 2019. Plasma Eicosanoid Levels in Tuberculosis and
Tuberculosis-Diabetes Co-morbidity Are Associated With Lung Pathology and
Bacterial Burden. *Front Cell Infect Microbiol* 9:335.

Pearl, J.E., B. Saunders, S. Ehlers, I.M. Orme, and A.M. Cooper. 2001. Inflammation and
lymphocyte activation during mycobacterial infection in the interferon-gamma-
deficient mouse. *Cell Immunol* 211:43-50.

Peres, C.M., L. de Paula, A.I. Medeiros, C.A. Sorgi, E.G. Soares, D. Carlos, M. Peters-Golden,
C.L. Silva, and L.H. Faccioli. 2007. Inhibition of leukotriene biosynthesis abrogates the
host control of Mycobacterium tuberculosis. *Microbes Infect* 9:483-489.

Ramakrishnan, L. 2012. Revisiting the role of the granuloma in tuberculosis. *Nat Rev*
Immunol 12:352-366.

Rockwood, N., E. du Bruyn, T. Morris, and R.J. Wilkinson. 2016. Assessment of treatment
response in tuberculosis. *Expert Rev Respir Med* 10:643-654.

Sadikot, R.T., H. Zeng, A.C. Azim, M. Joo, S.K. Dey, R.M. Breyer, R.S. Peebles, T.S. Blackwell,
and J.W. Christman. 2007. Bacterial clearance of Pseudomonas aeruginosa is
enhanced by the inhibition of COX-2. *Eur J Immunol* 37:1001-1009.

Serezani, C.H., J. Chung, M.N. Ballinger, B.B. Moore, D.M. Aronoff, and M. Peters-Golden.
2007. Prostaglandin E2 suppresses bacterial killing in alveolar macrophages by
inhibiting NADPH oxidase. *Am J Respir Cell Mol Biol* 37:562-570.

Sheppe, A.E.F., and M.J. Edelmann. 2021. Roles of Eicosanoids in Regulating Inflammation
and Neutrophil Migration as an Innate Host Response to Bacterial Infections. *Infect*
Immun 89:e0009521.

Silva, M.T., M.N. Silva, and R. Appelberg. 1989. Neutrophil-macrophage cooperation in the
host defence against mycobacterial infections. *Microb Pathog* 6:369-380.

Soares, A.P., C.K. Kwong Chung, T. Choice, E.J. Hughes, G. Jacobs, E.J. van Rensburg, G.
Khomba, M. de Kock, L. Lerumo, L. Makhethe, M.H. Maneli, B. Pienaar, E. Smit, N.G.
Tena-Coki, L. van Wyk, W.H. Boom, G. Kaplan, T.J. Scriba, and W.A. Hanekom. 2013.
Longitudinal changes in CD4(+) T-cell memory responses induced by BCG vaccination
of newborns. *J Infect Dis* 207:1084-1094.

Souto, F.O., J.C. Alves-Filho, W.M. Turato, M. Auxiliadora-Martins, A. Basile-Filho, and F.Q.
Cunha. 2011. Essential role of CCR2 in neutrophil tissue infiltration and multiple
organ dysfunction in sepsis. *Am J Respir Crit Care Med* 183:234-242.

Tan, Z., R. Jiang, X. Wang, Y. Wang, L. Lu, Q. Liu, S.G. Zheng, B. Sun, and B. Ryffel. 2013.
RORgammat+IL-17+ neutrophils play a critical role in hepatic ischemia-reperfusion
injury. *J Mol Cell Biol* 5:143-146.

Taylor, P.R., S. Roy, S.M. Leal, Jr., Y. Sun, S.J. Howell, B.A. Cobb, X. Li, and E. Pearlman. 2014.
Activation of neutrophils by autocrine IL-17A-IL-17RC interactions during fungal
infection is regulated by IL-6, IL-23, RORgammat and dectin-2. *Nat Immunol* 15:143-
151.

Team, R.C. 2021. R: A Language and Environment for Statistical Computing. In Vienna,
Austria.

Thompson, E.G., Y. Du, S.T. Malherbe, S. Shankar, J. Braun, J. Valvo, K. Ronacher, G. Tromp,
D.L. Tabb, D. Alland, S. Shenai, L.E. Via, J. Warwick, A. Aderem, T.J. Scriba, J. Winter,
G. Walzl, D.E. Zak, and T.B.B.C. Catalysis. 2017. Host blood RNA signatures predict
the outcome of tuberculosis treatment. *Tuberculosis (Edinb)* 107:48-58.

Tobin, D.M., J.C. Vary, Jr., J.P. Ray, G.S. Walsh, S.J. Dunstan, N.D. Bang, D.A. Hagge, S.
Khadge, M.C. King, T.R. Hawn, C.B. Moens, and L. Ramakrishnan. 2010. The lta4h
locus modulates susceptibility to mycobacterial infection in zebrafish and humans.
Cell 140:717-730.

Torrado, E., and A.M. Cooper. 2010. IL-17 and Th17 cells in tuberculosis. *Cytokine Growth*
Factor Rev 21:455-462.
Tsuda, Y., H. Takahashi, M. Kobayashi, T. Hanafusa, D.N. Herndon, and F. Suzuki. 2004. Three
different neutrophil subsets exhibited in mice with different susceptibilities to
infection by methicillin-resistant Staphylococcus aureus. *Immunity* 21:215-226.
Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. In Springer-Verlag New York,
ISBN: 978-3-319-24277-4.
Wu, T., E. Hu, S. Xu, M. Chen, P. Guo, Z. Dai, T. Feng, L. Zhou, W. Tang, L. Zhan, F. xiaochong,
S. Liu, X. Bo, and G. Yu. 2021. clusterProfiler 4.0: A universal enrichment tool for
interpreting omics data. *The Innovation* 2:100141-100141.
Zhu, A., A. Srivastava, J.G. Ibrahim, R. Patro, and M.I. Love. 2019. Nonparametric expression
analysis using inferential replicate counts. *Nucleic Acids Res* 47:e105.
1013

Figures and figure Legends:

1017 **Figure 1: PMNs harbours substantial amount of *Mtb* and provide an immune-privileged**
1018 **niche to *Mtb*.**

(A) Study design for BCG vaccination experiment. C57BL/6 mice were vaccinated intradermally with 10^6 BCG bacilli, followed by aerosol infection with H37Rv (100-200 CFU) 8 weeks post BCG vaccination. At 4,8,12 weeks post-infection, mice were euthanized for various studies. Time course of the lung (B) and spleen (C) bacterial burden in unvaccinated and BCG vaccinated groups (unpaired student's t-test was applied to compare both experimental groups at different time points. (D) Quantification of the sorted cells normalized to per 1 million of total acquired cells from the lungs of vaccinated and unvaccinated mice. (E) Determination of *Mtb* burden in individual sorted cells, plotted as CFU/10^4 sorted cells at 4,8,12 weeks post-infection, from the lungs of vaccinated and unvaccinated mice. Experimental groups were evaluated using a 2-way ANOVA statistical test. n= 5 mice/group in all treatment groups. (F) Heatmap of DEGs representing the z-scaled normalised average expression in each group. The DEGs are clustered using Euclidean distance and ward D2 methodology. Each cluster reflects significantly enriched GO BP terms unique to the cluster.

Figure 2: PMN-resident *Mtb* in BCG-vaccinated mice is susceptible to celecoxib treatment.

(A) Study plan. C57BL/6 mice were vaccinated intradermally with 10^6 BCG bacilli. After 8 weeks, vaccinated and unvaccinated mice were subjected to aerosol infection with H37Rv (100 CFU). 4 weeks post *Mtb* infection, mice received celecoxib (50mg/kg, orally) treatment for 4 and 8 weeks in with and without BCG vaccination group keeping the untreated arm as control. At 4,8,12 weeks post-infection, mice were euthanized, and lung and spleen were harvested. (B) Enumeration of lung and spleen bacterial burden in all four treatment groups (Control, BCG-vaccinated, celecoxib-treated and BCG-vaccinated celecoxib-treated) at 8- and 12-weeks post-infection. (C) *Mtb* burden in individual sorted cells (PMNs, Macrophages and MSCs), plotted as CFU/10^4 sorted cells at 12 weeks post-infection time point, from the lungs of all 4 treatment groups of mice. A 2-way ANOVA statistical test was applied to compare all 4 treatment groups at different time points. Sample size (n) = 5 mice/group in all

treatment groups. (D) H&E-stained lung sections of Control, BCG-vaccinated, celecoxib-
treated and BCG-vaccinated celecoxib-treated group at 10X magnification. (E) The average
z-scaled expression of the DEGs identified between each pairwise comparison is represented
in the heatmap. The data for UI and *Mtb* groups are the same as used in Fig. 1F, being part of
the common RNAseq experiment. (F) Significantly enrichment GO BP (p-adjusted ≤ 0.1)
highlights immune response against pathogens from the DEGs set of cluster 3.
1054

Figure 3: Loss of IFNγ leads to uncontrolled neutrophilia, huge *Mtb* burden, severe disease pathology and death.

(A) Experimental design and timeline. C57BL/6 mice (WT and *IFNγ* [-/-]) were infected with H37Rv, 100 CFU, via the aerosol route. Mice from both groups were euthanized at 4,7- and 10 weeks post-infection time-point. Lung and spleen tissues were harvested and processed for

diverse assays, as shown in the study design. (B) H&E-stained lung sections of WT and *IFNγ* $^{-/-}$ mice at 10X magnification, showing severe histopathology in the latter group of mice. (C) Lung IFA images from the same group of mice showing the extensive infiltration of PMNs (Ly6G$^+$ cells stained in red colour) in *Mtb*-infected *IFNγ* $^{-/-}$ mice compared to WT mice. (D) Number of PMNs in the lungs of *Mtb* infected WT and *IFNγ* $^{-/-}$, normalized to per 1 million of total acquired cells, at 4,7- and 10 weeks post infection timepoint. (E) *Mtb* burden in the PMNs, sorted from the lung of *Mtb* infected WT and *IFNγ* $^{-/-}$ (plotted as CFU/10^4 sorted cells) at 4,7- and 10 weeks post-infection. Experimental groups were evaluated using a 2-way ANOVA statistical test. Sample size (n)= 5-6 mice/group in all treatment groups. Luminex data: Quantification of several cytokines and chemokines from the lung supernatant of uninfected and *Mtb*-infected WT and *IFNγ* $^{-/-}$ mice. (F) Cytokines and chemokines that are upregulated in *Mtb*-infected *IFNγ* $^{-/-}$ mice compared to WT mice (G) Cytokines and chemokines that are downregulated in *Mtb*-infected *IFNγ* $^{-/-}$ mice compared to WT mice. Data was evaluated using a 2-way ANOVA statistical test. Sample size (n)= 5-6 mice/group in all treatment groups.

Figure 4: PMNs are one of the major sources of IL-17 in the lungs of *Mtb*-infected mice.
(A) Lung cells from WT *Mtb*-infected mice were stained with different antibodies and were assayed through flow cytometry to elucidate the cellular source for elevated IL-17. FACS gating strategy showing PMNs (Ly6G$^+$ CD11b$^+$ cells)) as the dominant producer of IL-17 in the lungs of *Mtb*-infected mice compared to Ly6G$^-$ CD11b$^-$ cells. (B) IFA images of the lungs of *Mtb*-infected WT mice at 6 weeks post-infection time-point showing the co-localization of CD4$^+$ T cells (shown in red colour) and IL-17 (shown in green colour). IFA images of the lungs of *Mtb*-infected WT (C) and *IFNγ* $^{-/-}$ (D) mice at 6 weeks post-infection time-point showing the co-localization of PMNs (Ly6G$^+$ cells shown in red colour) and IL-17 (shown in green colour). (E) Co-localization of PMNs (CD66b$^+$ cells shown in red colour) and IL-17 (shown in green colour) in the gut granuloma sections of intestinal TB (ITB) patients.

1091

Figure 5: Celecoxib treatment with IL-17 neutralization reverses IL-17-dependent neutrophilia and controls TB disease in *IFNγ⁻/⁻* mice.

(A) Study plan. *IFNγ⁻/⁻*C57BL/6 mice infected with H37Rv at a dose of 100 CFU via the aerosol route. One arm of the *Mtb*-infected mice received an αIL-17 neutralizing antibody (100μg/mouse), and the other arm received an isotype control antibody (100μg/mouse) intraperitoneally twice in a week, starting a day before *Mtb* infection. After 3 weeks of infection, both the groups of mice were further divided into two groups, one that received celecoxib (50mg/kg, orally) for 3 weeks and the other that received vehicle control (DMSO here). Mice from all four groups of mice were euthanized at 6 weeks post *Mtb* infection, and their lungs and spleen were harvested, and serum was stored for ELISA. (B) PMN's count,

normalized to 1 million total acquired cells, in the lungs of infected mice in all four treatment groups at 6 weeks post-infection. Bacterial burden in lungs (C) and spleen (D) of *Mtb* infected mice from described treatment groups at 6 weeks post-infection time-point sacrifice. (E) CFU (normalized to per 10^4 sorted cells) inside PMNs sorted out from the lungs of infected mice at 6 weeks post-infection sacrifice. (n= 3-4 mice/group). The statistical significance of the treatment groups was assessed by applying a one-way ANOVA test. (F) Survival curve of the group of *Mtb* infected *IFNγ$^{-/-}$* mice that received αIL-17 neutralizing antibody, celecoxib alone or together, along with the untreated control group and infected WT mice group. Both mice strains were infected with 100 CFU of H37Rv through an aerosol route. One arm of the *Mtb*-infected mice received an αIL-17 neutralizing antibody, and the other arm received an isotype control antibody twice a week, starting a day before the *Mtb* infection. After 3 weeks of infection, both the groups of mice were further divided into two groups, one that received celecoxib for 3 weeks and the other that received vehicle control (DMSO). Mice were incubated till their natural death. The time of death of each mouse was noted and plotted for survival analysis (n= 7 mice/group). (G) H&E-stained lung sections showing pathology in the αIL-17Ab+Celecoxib group. (H) Granuloma scoring for lung pathology from all 4 treatment groups. (I) %Th$_{17}$ cells (gated on CD4 T cells) in the lungs of mice from the same treatment groups described above. (J) Lung IFA images from the above-stated treated group of mice show the colocalization (yellow) of PMNs (Ly6G$^+$ cells shown in red colour) and IL17 (shown in green colour).

Figure 6: RORγt inhibition brings down *Mtb* burden and augments BCG vaccine efficacy in WT mice.

(A) Study plan. C57BL/6 mice were infected with H37Rv at a 100-200 CFU dose via the aerosol route. At 3 weeks post-infection, treatment with celecoxib and SR 2211 (inverse agonist for RORγt) was started for further 3 weeks. Celecoxib (50mg/kg) was administered

orally once a day, and SR 2211 (20mg/kg) was administered intraperitoneally thrice in a
week. At 6 weeks post-infection, mice from the treatment and untreated control groups were
sacrificed, lung and spleen were harvested for CFU, histopathology and FACS analysis, and
lung homogenates were stored for ELISA. (B) Levels of IL-17 in the lung homogenates of all
four-treatment groups of mice. (C) Lung IFA images from the above-stated treated group of
mice show the colocalization (yellow) of PMNs (Ly6G$^+$ cells shown in red colour) and IL17
(shown in green colour). (D) Lung PMN's count normalized to 1 million of total acquired
cells, (E) Lung PMN's *Mtb* burden normalized to per 10^4 sorted cells, (F) mycobacterial
burden in lung and spleen, in the above-stated treatment groups of mice. The statistical
significance of the treatment groups was assessed by applying a one-way ANOVA test, n=5
mice/group. (G) Experimental plan. C57BL/6 mice were vaccinated intradermally with 10^6
BCG bacilli for 8 weeks, followed by aerosol infection with H37Rv at a dose of 100-200
CFU. 4 weeks post-infection, treatment with SR2211 was started for additional 3 weeks,
intraperitoneally, thrice a week. At 7 weeks post-infection, mice from the described four
treatment groups were sacrificed, and lung and spleen were harvested for CFU and FACS
analysis. (H) CFU (normalized to per 10^4 sorted cells) inside PMNs sorted from the lungs of
infected mice at 7 weeks post-infection time-point. (I) *Mtb* burden in the lungs of infected
mice at 7 weeks post-infection (n= 5-6 mice/group). The statistical significance of the
treatment groups was assessed by applying a one-way ANOVA test.
1148

1149
Figure 7: High IL-17 precipitates adverse treatment outcomes in pulmonary TB
patients.
(A) Pulmonary TB (PTB) patients-cohort study design: It includes 133 subjects that were
cured of TB after the full standard ATT regime and 68 subjects that have failed the standard
ATT regime (independent of drug resistance) or have experienced recurrence of symptoms.
At the time of recruitment in the cohort study, the baseline blood levels of IL17, neutrophils,
total lymphocytes in both the treatment groups were measured. (B) blood IL17 levels, (C)
blood neutrophils level and (D) blood neutrophil/total lymphocyte ratio in treatment failure
and treatment success cases of pulmonary TB patients. Unpaired t-test was applied to
calculate significance.
1160