Title: The common *TMEM173 HAQ, AQ* alleles rescue CD4 T cell death, restore T-regs, and

prevent *SAVI (N153S)* inflammatory disease in mice

Authors/Affiliations: Alexandra Aybar-Torres[1], Lennon A Saldarriaga[1], Ann T. Pham[1], Amir

M. Emtiazjoo[1], Ashish K Sharma[2], Andrew J. Bryant[1] and Lei Jin[1*]

[1]Division of Pulmonary, Critical Care and Sleep Medicine, Department of Medicine,

[2]Division of Vascular Surgery & Endovascular Therapy, Department of Surgery, University of

Florida, Gainesville, FL 32610, U.S.A

*Corresponding Author:

Lei Jin, Ph.D.

1600 SW Archer Road, P.O.Box 100225, Gainesville, FL, 32610

Fax : 352-273-9154

Phone: 352-294-8495

Email: lei.jin@medicine.ufl.edu

Abstract

STING activation induces lymphocyte cell death that is independent of type I IFNs. The *in vivo* significance and mechanism of STING-mediated cell death is unclear. Using STING knock-in mice, we found that lymphocytes from the *HAQ*, *AQ*, and *Q293* mice are resistant to STING-mediated cell death *ex vivo*, establishing a critical role of STING residue 293 in cell death. CD4 T cellpenia is evident in STING-associated vasculopathy with onset in infancy (SAVI), an autosomal dominant, fatal inflammatory disease caused by gain-of-function human STING mutations. The *HAQ/SAVI(N153S)* and *AQ/SAVI(N153S)* mice have similar spleen CD4 T cells as the *WT* mice reversing the CD4 T cellpenia by the gain-of-function N153S STING. The *HAQ/SAVI(N153S), AQ/SAVI(N153S)* mice have more (~10-fold, ~20-fold, respectively) T-regs than *WT/SAVI(N153S)* mice. Remarkably, while have comparable TBK1, IRF3, NFκB activation as the *WT/SAVI*, the *AQ/SAVI* mice have no tissue inflammation, regular body weight, and normal lifespan. The type I IFNs-independent function of STING in health and diseases has been increasingly recognized. We propose that STING activation promotes tissue inflammation by depleting T-regs cells *in vivo*. Billions of modern humans have the dominant *HAQ, AQ* alleles. STING research and STING-targeting immunotherapy should consider *TMEM173* heterogeneity in humans.

Keywords: STING, *HAQ AQ TMEM173* alleles, tissue inflammation, SAVI, T-regs

One-sentence summary: The Common *TMEM173* alleles *HAQ, AQ* prevent SAVI disease.

Introduction

STING drives cytosolic DNA-induced type I IFNs production [1]. Recent research revealed that STING promotes inflammation in a variety of inflammatory diseases including nonalcoholic fatty liver disease, nonalcoholic steatohepatitis, kidney injury, neurodegenerative diseases, cardiovascular diseases, obesity, diabetes, and aging [2-9]. The type I IFNs-independent function of STING has also emerged [10,11]. For example, initially described as a type I interferonopathy [12], the symptoms of STING-associated vasculopathy with onset in infancy (SAVI) are largely independent of type I IFNs [13-16]. In a STING N153S mouse model of SAVI, crossing N153S mice to cGAS, IRF3/IRF7, and IFNAR1 knockout mice, N153S mice still developed spontaneous lung diseases [13]. In another example of STING-dependent, type I IFNs-independent inflammatory disease, DNase II-deficient mice developed autoinflammatory arthritis that can be rescued by STING deficiency [17]. But $Dnase\ II^{-/-}$ mice with the STING-S365A mutation still exhibited severe polyarthritis [18,19]. The STING-S365A mutant cannot recruit IRF3 and produce IFNβ [10,11,18,19]. How STING drives inflammation *in vivo,* independent of type I IFNs, remains unknown. The activation of NFκB or NLRP3 inflammasome by STING has been proposed [9,20].

Characterized as an innate immune sensor, STING expression is, paradoxically, high in CD4 T cells [12,21]. Furthermore, STING activation kills mouse and human CD4 T cells *ex vivo* [22-24]. SAVI patients and mouse model had CD4 T cellpenia [12,24]. STING was first discovered as MPYS for its cell growth inhibition and cell death function in mouse B lymphoma cells [25]. STING-mediated cell death is cell type dependent. For example, while STING activation kills human endothelial cells, primary and cancerous T cells, it does not kill mouse MEFs, BMDCs, or BMDMs [26-28]. Second, STING-mediated cell death is type I IFNs-independent [24,26,28]. Multiple cell death pathways, i.e., apoptosis, necroptosis, pyroptosis, ferroptosis, and PANoptosis, are triggered [28-30]. Last, the *in vivo* biological significance of STING-mediated CD4 T cell death is not clear [24,28]. WT and STING$^{-/-}$ mice have similar thymic and pLN CD4$^-$CD8$^-$, CD4$^+$CD8$^+$, and single positive CD4 and CD8 T cell populations [22,23]. The expression of CD4 and CD8 TCR are comparable [22,23]. The naive, memory, or regulatory T cell populations are similar in WT and STING$^{-/-}$ mice that expressed comparable levels of CD69 and CD25 following TCR activation [22,23]. Together, the absence of STING in mice does not alter T cell development or TCR signaling [22]. In humans, SAVI patients with constitutively activated STING have low CD4 T cell numbers [12], and type I

IFNs are dispensable for STING-mediated human CD4 T cell death [24]. Notably, SAVI patients

(N154S or V155M) had similar counts of CD8 T and B cells [12].

Also, unlike other innate immune genes, the human *TMEM173* gene is highly heterogeneous

[31,32]. ~50% of people in the U.S. carry at least one copy of non-*WT (R232) TMEM173* allele [31].

Among them, the R71H-G230A-R293Q (*HAQ*) is the 2nd most common *TMEM173* allele carried

by ~23% of people in the U.S. [31]. However, in East Asians, *WT/HAQ* (34.3%), not *WT/WT*

(22.0%), is the most common *TMEM173* genotype [32]. Critically, the *HAQ* allele was positively

selected in modern humans outside Africa [33]. Anatomically modern humans outside Africa are

descendants of a single Out-of-Africa Migration 50,000~70,000 years ago. ~1.4% of Africans have

the *HAQ* allele, while ~63.9% of East Asians are *HAQ* carriers [33]. Haplotype analysis revealed that

HAQ was derived from G230A-R293Q (*AQ*) allele [33]. Importantly, the *AQ* allele was negatively

selected outside Africa [33]. ~40.1% of Africans are *AQ* carriers, while ~0.4% of East Asians have

the *AQ* allele [33]. Importantly, *TMEM173* alleles often have a dominant negative effect because the

protein product STING exists as a homodimer [25,34]. For example, SAVI is an autosomal dominant

inflammatory disease [35]. *WT/HAQ* individuals had reduced Pneumovax23®-induced antibody

responses compared to *WT/WT* individuals (NCT02471014) [36]. Notably, *AQ* responds to CDNs

and produce type I IFNs *in vivo* and *in vitro* [33,37,38], but the *AQ* allele was negatively selected in

non-Africans [33]. In contrast, the *HAQ* allele, defective in CDNs-type I IFNs responses [31,32,36,39,40],

was positively selected in non-Africans[33] indicating that the CDNs-type I IFNs independent

function of STING was essential for the survival of early modern humans outside of Africa.

In this study, we discovered, surprisingly, that the *HAQ*, *AQ* cells are resistant to STING-

mediated lymphocyte death. We generated *HAQ/SAVI(N153S)* and *AQ/SAVI(N153S)* mice and

found that the *HAQ, AQ* alleles prevent CD4 T cellpenia, increase/restore T-regs and alleviate/stop

tissue inflammation in SAVI mice, thus providing evidence for the *in vivo* significance of type I

IFNs-independent, STING-mediated CD4 T cell death.

Results

STING activation kills mouse spleen CD4, CD8 T and CD19 B cells *ex vivo*.

We first used the synthesized non-CDNs STING agonist diABZI [41] to induce lymphocyte death because diABZI induces cell death [27,42] and diABZI is in clinical trials (NCT05514717). Splenocytes from C57BL/6N mice were treated with diABZI in culture, and cell death was determined by Annexin V and Propidium Iodide stain. Splenocyte cell death could be detected as early as 5hrs post diABZI treatment (Figure S1A), indicating that diABZI likely directly kills splenocytes rather than via the secondary cytokines, *e.g.* TNF, type I IFNs, et.al. Dosage responses showed that ~25ng/ml diABZI could kill 70% of splenocytes (Figure S1B). Similarly, STING agonists 2'3'-cGAMP, DMXAA, and synthetic CDNs RpRpssCDA, killed mouse spleen CD4, CD8 T cells, and CD19 B cells (Figure 1A). Thus, STING activation induces mouse lymphocyte death *ex vivo*.

TBK1 activation is required for STING-mediated mouse spleen cell death *ex vivo*.

STING activation can lead to apoptosis, pyroptosis, necroptosis, or ferroptosis [24-30,42,43]. We then treated mouse splenocytes with apoptosis, pyroptosis, necroptosis inhibitors, STING inhibitors H151, C176, palmitoylation inhibitor 2-bromopalmitate (2-BP), followed by diABZI stimulation. Inhibitors for NLRP3 (MCC950), RIPK1 (Necrostatin-1), RIPK3 (GSK872), Caspase-1 (VX-795), Caspase-3 (Z-DEVD-FMK), Caspase 1,3,8,9 (Q-VD-Oph), ferroptosis (liproxstatin-1) did not affect diABZI-induced splenocyte cell death *ex vivo* (Figure S1C, 1B). The STING inhibitors H151, C175, and 2-BP also could not prevent diABZI-induced cell death (Figure S1C). Instead, the TBK1 inhibitor Bx-795 treatment abolished diABZI-induced splenocyte death (Figure S1A).

BX-795 is a multi-kinase inhibitor including 3-phosphoinositide-dependent protein kinase 1 (PDK1) and TBK1 (IC_{50s} = 6 and 11 nM, respectively). However, the treatment of PDK1 inhibitor GSK2334470 (IC50 = 10nM) did not prevent diABZI-induced splenocyte death (Figure 1B). In contrast, GSK8612, a highly potent and selective inhibitor for TBK1, prevented diABZI-induced splenocyte death (Figure 1B). Thus, TBK1 activation is likely critical for STING-mediated lymphocyte cell death in mice.

***HAQ, AQ, Q293 STING* knock-in mouse splenocytes are resistant to STING-mediated cell death *ex vivo*.**

HAQ and *AQ* are common human *TMEM173* alleles [31-33]. Previously, we reported that *HAQ* knock-

in mice are defective in CDNs-induced immune responses, while CDNs responses in *AQ* knock-

in mice are similar to WT mice [33]. We treated splenocytes from *HAQ* and *AQ* mice with diABZI

ex vivo and found, surprisingly, that both *HAQ* and *AQ* splenocytes were resistant to diABZI-

induced cell death (Figure 1C, 1D). In comparison, IFNAR1$^{-/-}$ splenocytes were killed by diABZI,

confirming that STING-mediated lymphocytes death are type I IFNs-independent (Figure 1C, 1D)

[24,26,28].

HAQ and *AQ* share the common A230 and Q293 residues changes. We thus generated a *Q293*

TMEM173 knock-in mouse. Notably, the Q293 splenocytes were resistant to STING agonists 2'3'-

cGAMP, RpRpssCDA, and diABZI-induced cell death (Figure 1E, 1F). Thus, the residue 293 of

STING is critical for its cell death function.

WT/HAQ, WT/AQ mouse splenocytes are partially resistant to STING-mediated cell death

***ex vivo*.**

WT/HAQ (34.3%), not *WT/WT* (22.0%), is the most common human *STING* genotype in East

Asians, while *WT/AQ* (28.2%) is the 2nd most common *STING* genotype in Africans [32]. We

generated *WT/HAQ*, *WT/AQ* mice and treated the splenocytes with mouse STING agonist

DMXAA. *WT/HAQ* and *WT/AQ* splenocytes were protected from 25µg/ml DMXAA-induced cell

death (Figure 1G). 100µg/ml DMXAA could kill *WT/HAQ* and *WT/AQ* splenocytes albeit less than

WT/WT cells (Figure 1G). Thus, the *HAQ* and *AQ* alleles are not recessive and likely impact

STING activation even in heterozygosity.

STING activation kills primary human CD4 T cells, but not CD8 T or CD19 B cells

STING agonists-based clinical trials in humans have been disappointing (NCT02675439,

NCT03010176, NCT05514717) [44,45]. We also showed that the human *TMEM173* gene might

undergo natural selection during the out-of-Africa migration [33] sensitive to evolutionary pressure.

Thus, we investigated STING-mediated death in primary human lymphocytes.

Human explant lung cells from the *WT(R232)/WT(R232)* donors were treated with STING

agonists 2'3-c GAMP, RpRpssCDA, diABZI for 24hrs in culture. Lymphocyte cell death was

determined by Propidium Iodide staining. Different from mouse lymphocytes, diABZI and

RpRpssCDA killed human CD4 T but not CD8 T or CD19 B cells (Figure 2A, 2B). Human CD8

T and CD19 B cells are resistant to 500ng/ml diABZI-induced cell death (Figure S2A).

WT/HAQ human CD4 T cells are resistant to low-dose of diABZI-induced cell death.

WT/HAQ mouse splenocytes are resistant to low dose diABZI-induced cell death. We obtained

lung explants from *WT/WT* and *WT/HAQ* individuals (Figure S2B) and treated them with diABZI

in culture. 25ng/ml diABZI killed *WT/WT*, but not *WT/HAQ*, lung CD4 T cells (Figure 2C).

diABZI induces cell death in STING-KO human THP-1 cells reconstituted with *WT* human

STING (R232) but not *HAQ, AQ or Q293* human *STING* allele.

To further determine cell death influenced by human *TMEM173* alleles *HAQ, AQ* and Q293, we

used the *STING-KO* THP-1 cell line because STING agonist induces type I IFNs and cell death in

STING-KO THP-1 cells expressing WT human STING [30] (Figure S2C, S2D). We, thus, generated

stable THP-1 *STING-KO* lines expressing *HAQ, AQ, WT* or *Q293 TMEM173* allele. Cell death

was determined by Annexin-V staining. diABZI killed THP-1 *STING-KO* lines expressing *WT* but

not *HAQ, AQ,* or *Q293 TMEM173* allele (Figure 2D, 2E). No cell death was induced in the *Q293*

THP-1 cells stimulated by 20 to 200 ng/ml of diABZI (Figure 2F). diABZI also did not induce

STING activatin in *Q293* THP-1 cells (Figure 2G). Notably, 50ng/ml diABZI induced p-IRF3

activation and type I IFNs in *AQ* THP-1 cells, but not *HAQ* THP-1 cells (Figure 2H, 2I, 2J)

indicating that the STING-cell death and STING-IRF3-Type I IFNs pathways can be uncoupled.

HAQ and AQ alleles rescue the lymphopenia and suppress myeloid cell expansion in

SAVI(N153S) mice.

The *in vivo* significance of the STING/MPYS-cell death is unclear. SAVI is an autosomal

dominant, inflammatory disease caused by gain-of-function STING mutation [12]. CD4 T cellpenia

was found in SAVI patients and SAVI mouse models [12,13]. We thus generated *HAQ/SAVI(N153S)*

and *AQ/SAVI(N153S)* mice aiming to restore CD4 T cells in SAVI mice to establish the *in vivo*

significance of STING/MPYS-cell death.

First, *HAQ/SAVI(N153S)* and *AQ/SAVI(N153S)* mice had reduced splenomegaly compared to

WT/SAVI(N153S) mice though their spleens were still larger than the littermates *WT/HAQ* and

WT/AQ (Figure 3A, 3B). Next, *HAQ/SAVI(N153S)* and *AQ/SAVI(N153S)* mice had similar spleen

B cells and CD4 T cells numbers as the *WT/HAQ*, *WT/AQ* mice (Figure 2B, 2C). Their CD8⁺ T

cells were lower than their WT littermates but much higher than the *WT/SAVI(N153S)* mice (Figure

3D). Third, spleen myeloid cell numbers, i.e., neutrophils, Ly6C^{hi} monocytes and F4/80

macrophages, were all reduced by half compared to *WT/SAVI(N153S)* mice (Figure 3E~3H).

Notably, the *HAQ/SAVI(N153S)* and *AQ/SAVI(N153S)* mice also had restored bone marrow

monocyte (Figure S3). Thus, *HAQ* and *AQ* alleles prevent lymphopenia and suppress myeloid cells

expansion in *SAVI(N153S)* mice.

The *HAQ* allele alleviates and the *AQ* allele prevents *SAVI(N153S)* disease in mice.

SAVI(N153S) disease is characterized by early onset, failure to thrive (low body weight), persistent

lung inflammation, decreased lung function, and young death in humans and mouse models

[12,16,46,47]. The *HAQ/SAVI(N153S)* mice weighed more and had an improved lifespan than the

WT/SAVI(N153S) mice (Figure 4A, 4B). The lifespan, airway resistance and tissue inflammation

were also improved in *HAQ/SAVI(N153S)* mice compared to the *WT/SAVI(N153S)* mice (Figure

4C, 4D, 4J, 4K, S4). However, the pulmonary artery pressure was still elevated in

HAQ/SAVI(N153S) mice (Figure 4E). Remarkably, the *AQ/SAVI(N153S)* mice had similar body

weight and lifespan as the *WT/AQ* mice (Figure 4F, 4G). The airway resistance, pulmonary artery

pressure, and tissue inflammation in *AQ/SAVI(N153S)* were similar to the *WT/AQ* mice (Figure

4H, 4I, 4J, 4K, S4). Thus, the *AQ* allele prevents inflammatory SAVI disease in mice.

diABZI induces similar STING, TBK1, IRF3, NFκB activation in the *AQ/SAVI(N153S)* and

***WT/SAVI(N153S)* BMDM.**

SAVI was characterized as type I interferonopathy [16]. However, several studies showed that type

I IFN signaling and IRF3 activation were dispensable for SAVI disease [13,14,47,48]. *AQ* allele prevents

SAVI disease (Figure 4). However, diABZI-treated *AQ/SAVI(N153S)* and *WT/SAVI(N153S)*

BMDM had similar TBK1-IRF3 activation (Figure 5A). Furthermore, diABZI activation led to

STING protein degradation in *WT/SAVI* and *AQ/SAVI* BMDM (Figure 5B). In addition, diABZI

treatment caused IκBα degradation in *WT/SAVI* and *AQ/SAVI* BMDM (Figure 5C). Last, using

cleavable crosslinker dithiobis(succinimidyl propionate (DSP), we showed that STING in

WT/SAVI, AQ/SAVI BMDM forms similar dimer *in situ*. Thus, the *AQ/SAVI* BMDM had similar

STING degradation, TBK1, IRF3, NFκB activation, and dimerization as the *WT/SAVI* BMDM.

216 ***The HAQ* allele increased, and the *AQ* allele restored T-regs in *SAVI(N153S)* mice.**

217 IFNγ was proposed to drive SAVI disease [14,34,48]. We confirmed that *WT/SAVI* CD4 T cells were

218 enriched with IFNγ⁺ cells (Figure 6A). However, *WT/SAVI* mice have CD4 T cellpenia. Thus, the

219 total numbers of spleen IFNγ⁺ CD4 T cells were comparable in *WT/SAVI* and *AQ/SAVI* mice

220 (Figure 6A). In contrast, the *HAQ/SAVI mice* had decreased IFNγ⁺ CD4 T cells (Figure 6A).

221 CD4 T cellpenia depletes CD4 T-regs. Indeed, *WT/SAVI* mice had ~20-fold reduction of spleen

222 FoxP3⁺ T-regs compared to *AQ/SAVI* or *WT/WT* mice (Figure 6B). The *HAQ/SAVI* mice also had

223 ~10-fold more T-regs than the *WT/SAVI* mice (Figure 6B).

Discussion

This study, using the *HAQ, AQ, SAVI(N153S) TMEM173* knock-in mice, reveals the *in vivo* significance of STING-mediated CD4 T cell death. *HAQ, AQ* alleles prevent CD4 T cellpenia, and increase/restore CD4 T-regs in SAVI mice. The results are consistent with previous finding that the impaired CD4 T cell proliferation by the *SAVI(V155M)* mutant could be rescued by the addition of the *HAQ* allele *in vitro* [23]. Besides SAVI, STING has been increasingly implicated in inflammatory diseases such as nonalcoholic fatty liver disease, nonalcoholic steatohepatitis, cardiomyopathy, obesity, diabetes, neurodegenerative diseases, aging, and kidney injury, many of which are independent of type I IFNs [2,3,9]. It is tempting to suggest that STING activation in CD4 T cells leads to CD4 T-regs depletion that break tissue tolerance and exacerbates tissue inflammation.

Human immunodeficiency virus (HIV) primarily infects CD4 T cells and might activate the STING pathway in CD4 T cells [49-54]. The loss of CD4 T cells is the hallmark of untreated HIV infection [55,56], and the measurement of CD4 T cell count is a central part of HIV care. We found that *HAQ* and *AQ* CD4 T cells are resistant to STING-mediated cell death. Mogensen and colleagues reported that *HAQ/HAQ* was enriched in HIV-infected long-term nonprogressors [39]. These *HAQ/HAQ* individuals had reduced inhibition of CD4 T cell proliferation and a reduced immune response to DNA and HIV [39]. It is likely that HIV infection activates STING-cell death pathway in CD4 T cells. Furthermore, in *HAQ/SAVI* and *AQ/SAVI* mice, one copy of *HAQ, AQ* allele suppressed CD4 T cell death. *HAQ, AQ* carriers might have fewer HIV-induced CD4 T cell death and be long-term nonprogressors in HIV infection-induced Acquired immunodeficiency syndrome (AIDS).

Activating the STING pathway is a promising strategy for cancer immunotherapy [57-63]. Multiple STING agonists are in clinical trials [44,45]. Recently, the safety issue emerged in some STING agonist trials [44,45]. For example, in the STING agonist, ADU-S100, clinical trial, Grade 3/4 treatment-related adverse events were reported in 12.2% of 41 pretreated patients (NCT02675439)[44,45]. The National Institutes of Health defines grade 3 as "incapacitating; unable to perform usual activities; requires absenteeism or bed rest." In a clinical trial using STING antibody-drug-conjugate (ADC) that conjugates diABZI to anti-HER2 Ab, a Grade 5 (fatal) serious adverse event was recorded and deemed related to the STING-ADC (NCT05514717). SAVI disease, driven by overreacting STING, is often fatal. *AQ*, to a less degree, *HAQ*, suppress

mortality in SAVI mice. Future STING clinical trials should be based on human *TMEM173*
genotype to achieve safe and effective responses.

Mechanistically, apoptosis, pyroptosis, ferroptosis, necroptosis, and PANoptosis have all been
reported in STING-mediated cell death [24-30,42,43]. Different cell types and STING agonists used
likely contributed to the inconsistency and complexity. Here, we focused on lymphopenia in the
SAVI mice that avoids ligand-dependent, non-physiological dosage in STING-mediated cell
death. *HAQ* and *AQ* alleles could prevent CD4 T cellpenia in the SAVI mice strongly indicating
that residue A230 or Q293 prevent STING-mediated CD4 T cell death *in vivo*. Splenocyte from
Q293 mice were resistant to STING agonists-induced cell death *ex vivo*. Thus, it is likely that the
Q293 residue is critical for STING-mediate lymphopenia. Notably, Q293 is outside the C-terminal
tail (CTT) (residues 341–379 of human STING) critical for TBK1 recruitment and IRF3
phosphorylation [64] or miniCTT domain (aa343–354) [23] , or the UPR motif (aa322–343) [46]
important for T cells death *in vitro*. Further studies are needed to understand how the aa293 of
STING mediates cell death *in vivo*. Noteworthy, *AQ/SAVI* cells had similar TBK1-IRF3, NFκB
activation and STING degradation as the *WT/SAVI* cells. Yet, *AQ/SAVI* mice did not have CD4 T
cellpenia as *WT/SAVI* mice suggesting that the canonical STING-TBK1-IRF3/NFκB pathway,
likely STING oligomerization, is not sufficient for the induction of cell death at the physiological
condition.

We used the *WT/N153S* knock-in SAVI mouse model that spontaneously develop lung
inflammation, T cell cytopenia, and early mortality, mimicking pathological findings in human
SAVI patients[15]. Using the *WT/N153S* SAVI mouse model and human Jurkat T cell line, it was
proposed that STING activation causes chronic ER stress and unfolded protein response, leading
to T cell death by apoptosis [46]. Furthermore, the study showed that crossing *WT/N153S* mice to the
OT-I mice reduced ER stress and restored CD8$^+$, but not CD4$^+$, T cells [46]. The restoration of CD8$^+$T
cells reduces inflammation and lung disease [46]. However, human *WT/N154S* SAVI patients have
normal CD8$^+$ T cells numbers [12], and primary human CD8$^+$ T cells are largely resistant to STING-
agonists-induced cell death *ex vivo* (Figure 2A) [24]. Thus, it is puzzling how restoring CD8$^+$ T cells
can rescue SAVI phenotypes since the SAVI patients already have normal CD8$^+$ T cells numbers.

Finally, it is unexpected that both *HAQ* and *AQ* alleles are resistant to cell death. Our previous
studies showed that the *HAQ* and *AQ* alleles have opposite functions [33]. AQ-STING, not HAQ-
STING, responds to CDNs [31-33,36-40,65]. *AQ* mice are lean while *HAQ* mice are fat [33]. Most

importantly, *HAQ* was positively selected, while *AQ* was negatively selected, in modern humans

outside Africans [33]. Thus, the death pathway of STING is also distinct from the STING function

that was naturally selected. Together, we propose that there are at least three unique pathways of

STING/MPYS: STING-Type I IFNs, STING-cell death, and STING-fatty acid metabolism.

Materials and Methods

Study Design. The study was designed to reveal (i) the *in vivo* significance of the type I IFNs-independent, STING-dependent cell death function; (ii) the interplay between common *STING* alleles *HAQ, AQ* and the rare, gain-of-function SAVI STING mutation; (iii). The driver for the inflammatory SAVI disease. Mouse splenocytes, primary human lung cells, human THP-1 cells and *HAQ, AQ, SAVI* knock-in mice were used to establish the *in vivo* significance and human relevance. All the repeats were biological replications that involve the same experimental procedures on different mice. Where possible, treatments were assigned blindly to the experimenter by another individual in the lab. When comparing samples from different groups, samples from each group were analyzed in concert, thereby preventing any biases that might arise from analyzing individual treatments on different days. All experiments were repeated at least twice.

Mice. *WT/SAVI(N153S)* mice were purchased from The Jackson Laboratory. *HAQ*, *AQ* mice were previously generated in the lab [32,33]. The *Q293* mice were generated by Cyagen Biosciences. Briefly, the linearized targeting vector was transfected into JM8A3.N1 C57BL/6N embryonic stem cells. A positive embryonic stem clone was subjected to the generation of chimera mice by injection using C57BL/6J blastocysts as the host. Successful germline transmission was confirmed by PCR sequencing. The heterozygous mice were bred to Actin-flpase mice [The Jackson Laboratory, B6.Cg-Tg (ACTFLPe)9205Dym/J] to remove the neo gene and make the *Q293* knock-in mouse. Age- and gender-matched mice (2 – 6 month old, both male and female) were used for indicated experiments. *WT/SAVI* (male) x *WT/HAQ* (female), *WT/SAVI* (male) x *WT/AQ* (female) breeders were set up to generate *HAQ/SAVI, AQ/SAVI* mice. Mice were housed at 22°C under a 12-h light-dark cycle with ad libitum access to water and a chow diet (3.1 kcal/g, Teklad 2018, Envigo, Sommerset, NJ) and bred under pathogen-free conditions in the Animal Research Facility at the University of Florida. Littermates of the same sex were randomly assigned to experimental groups. All mouse experiments were performed by the regulations and approval of the Institutional Animal Care and Use Committee at the University of Florida, IACUC202200000058.

Reagents. Recombinant human IFNβ (R&D, cat no. 8499-IF-010/CF), diABZI (Invivogen, cat no. 2138299-34-8), 2'3'-cGAMP (Invivogen, cat no. tlrl-nacga23-02), DMXAA (Invivogen, cat no. tlrl-dmx), H151 (Invivogen, cat no. inh-h151), RpRpSSCDA (Biolog, cat no. C118), THP1-Dual™ KO-STING Cells (Invivogen, cat no. thpd-kostg). All other chemical inhibitors are from Selleckchem.

Histology. Lungs and livers were fixed in 10% formalin, paraffin-embedded, and cut into 4-μm sections. Lung, liver sections were then stained for hematoxylin-eosin. All staining procedures were performed by the histology core at the University of Florida. Briefly, tissue sectins were immersed Harris Hematoxylin for 10 seconds, then washed with tap water. Cleard sections were re-immersed in EOSIN stain for ~30 seconds. The sections were washed with tap water until clear, then dehydrate in ascending alcohol solutions (50%,70%,80%,95% x 2, 100% x 2). Afterwards, the sections werer cleared with xylene (3 - 4 x). The sections were mounted on glass slide with permount organic mounting medium for visulization.

Lung Function. Pulmonary function was evaluated using an isolated, buffer-perfused mouse lung apparatus (Hugo Sachs Elektronik, March-Huggstetten, Germany), as previously described [66]. Briefly, mice were anesthetized with ketamine and xylazine and a tracheostomy was performed, and animals were ventilated with room air at 100 breaths/min at a tidal volume of 7 μl/g body weight with a positive end-expiratory pressure of 2 cm H2O using a pressure-controlled ventilator (Hugo Sachs Elektronik, March-Huggstetten, Germany).

Isolation of lung cells. Cells were isolated from the lung as previously described [67]. The lungs were perfused with ice-cold PBS and removed. Lungs were digested in DMEM containing 200μg/ml DNase I (Roche, 10104159001), 25μg/ml Liberase TM (Roche, 05401119001) at 37°C for 2 hours. Red blood cells were then lysed and a single cell suspension was prepared by filtering through a 70-μm cell strainer.

BMDM activation
BMDMs were induced from mouse bone marrow cells cultured in RPMI 1640 (cat#11965; Invitrogen) with 10% FBS, 2 mM L-glutamine, 1 mM sodium pyruvate, 10 mM HEPES buffer,

1% nonessential amino acids, 50 mM 2-ME, 1% Pen/Strep, with 20ng/ml M-CSF (Kingfisher,

RP0407M) for 10 days [32].

Flow cytometry. Single cell suspensions were stained with fluorescent-dye-conjugated antibodies

in PBS containing 2% FBS and 1mM EDTA. Surface stains were performed at 4°C for 20 min.

For intracellular cytokine or transcription factor staining of murine and human cells, cells were

fixed and permeabilized with the Foxp3 staining buffer set (eBioscience, cat no 00-5523-00). Cells

were washed and stained with surface markers. Cells were then fixed and permeabilized

(eBioscience, cat no. 00-5523-00) for intracellular cytokine stain. Data were acquired on a BD

LSRFortessa and analyzed using the FlowJo software package (FlowJo, LLC). Cell sorting was

performed on the BD FACSAriaIII Flow Cytometer and Cell Sorter. More information on the

antibodies used can be found in supplementary table S1.

Human lung explants

Human lung explants were procured at the Lung Transplant Center, Division of Pulmonary,

Critical Care and Sleep Medicine, Department of Medicine, University of Florida. Donor and

patients consent was obtained for a research protocol (UF IRB201902955-Treatment with IFNβ

Induces Tolerogenic Lung Dendritic Cells in Human advanced lung disease). Healthy donor lungs

were surgically removed postmortem, perfused, small pieces were cut from the right middle and

lower lobes for research purpose, and stored in cold Perfadex ® at 4°C for no more than 12 hrs

before processing. Ex planted lungs from emphysema lung transplant patients were stored in cold

Perfadex ® at 4°C for no more than 12 hrs before the process. No lung explants were procured

from prisoners.

Statistical Analysis. To gain statistical power, we employ three~four mice/groups to characterize

lung immunity. The statistical justification for group size was calculated using the SAS program

to calculate the animal numbers. The analysis was carried out using a standard error of 0.5 for

immunological assays, and a power of 0.9. All data are expressed as means ± SEM. Statistical

significance was evaluated using Prism 7.0 software. Comparisons between two groups were

analyzed by performing an unpaired Student's *t* test. Comparisons between more than two groups

were analyzed by performing a one-way analysis of variance (ANOVA) with Tukey's multiple

comparisons test.

Supplemental materials

Figure S1: Bx-795 inhibits diABZI-induced mouse splenocyte death.

Figure S2: STING activation in primary human cells and THP-1 cells reconstituted with WT

human STING.

Figure S3: *HAQ* and *AQ* restore bone marrow monocytes in SAVI(N153S) mice.

Figure S4: *HAQ* and *AQ* suppress lung myeloid infiltration in SAVI(N153S) mice.

References

1   Barber, G. N. STING: infection, inflammation and cancer. *Nat Rev Immunol* **15**, 760-770
394   (2015). https://doi.org:10.1038/nri3921

395   2   Skopelja-Gardner, S., An, J. & Elkon, K. B. Role of the cGAS-STING pathway in systemic
396   and   organ-specific   diseases.   *Nat   Rev   Nephrol*   **18**,   558-572   (2022).
397   https://doi.org:10.1038/s41581-022-00589-6

398   3   Bai, J. & Liu, F. cGASSTING signaling and function in metabolism and kidney diseases.
399   *J Mol Cell Biol* **13**, 728-738 (2021). https://doi.org:10.1093/jmcb/mjab066

400   4   Li, H., Guo, X., Aquino, E. & Wu, C. Mini review: STING activation during non-alcoholic
401   fatty   liver   disease.   *Front   Nutr*   **10**,   1139339   (2023).
402   https://doi.org:10.3389/fnut.2023.1139339

403   5   Gulen,   M.   F.   *et al.*   cGAS-STING   drives   ageing-related   inflammation   and
404   neurodegeneration. *Nature* **620**, 374-380 (2023). https://doi.org:10.1038/s41586-023-
405   06373-1

406   6   Szego, E. M. *et al.* Constitutively active STING causes neuroinflammation and
407   degeneration   of   dopaminergic   neurons   in   mice.   *Elife*   **11**   (2022).
408   https://doi.org:10.7554/eLife.81943

409   7   Gao, L. *et al.* STING/ACSL4 axis-dependent ferroptosis and inflammation promote
410   hypertension-associated   chronic   kidney   disease.   *Mol   Ther*   (2023).
411   https://doi.org:10.1016/j.ymthe.2023.07.026

412   8   Yan, M. *et al.* Mitochondrial damage and activation of the cytosolic DNA sensor cGAS-
413   STING pathway lead to cardiac pyroptosis and hypertrophy in diabetic cardiomyopathy
414   mice. *Cell Death Discov* **8**, 258 (2022). https://doi.org:10.1038/s41420-022-01046-w

415   9   Gao, K. M., Marshak-Rothstein, A. & Fitzgerald, K. A. Type-1 interferon-dependent and
416   -independent mechanisms in cyclic GMP-AMP synthase-stimulator of interferon genes-
417   driven   auto-inflammation.   *Curr   Opin   Immunol*   **80**,   102280   (2023).
418   https://doi.org:10.1016/j.coi.2022.102280

419   10   Yamashiro, L. H. *et al.* Interferon-independent STING signaling promotes resistance to
420   HSV-1 in vivo. *Nat Commun* **11**, 3382 (2020). https://doi.org:10.1038/s41467-020-17156-
421   x

Wu, J., Dobbs, N., Yang, K. & Yan, N. Interferon-Independent Activities of Mammalian STING Mediate Antiviral Response and Tumor Immune Evasion. *Immunity* **53**, 115-126 e115 (2020). https://doi.org:10.1016/j.immuni.2020.06.009

Liu, Y. *et al.* Activated STING in a vascular and pulmonary syndrome. *N Engl J Med* **371**, 507-518 (2014). https://doi.org:10.1056/NEJMoa1312625

Luksch, H. *et al.* STING-associated lung disease in mice relies on T cells but not type I interferon. *J Allergy Clin Immunol* **144**, 254-266 e258 (2019). https://doi.org:10.1016/j.jaci.2019.01.044

Stinson, W. A. *et al.* The IFN-gamma receptor promotes immune dysregulation and disease in STING gain-of-function mice. *JCI Insight* **7** (2022). https://doi.org:10.1172/jci.insight.155250

Warner, J. D. *et al.* STING-associated vasculopathy develops independently of IRF3 in mice. *J Exp Med* **214**, 3279-3292 (2017). https://doi.org:10.1084/jem.20171351

Fremond, M. L. *et al.* Overview of STING-Associated Vasculopathy with Onset in Infancy (SAVI) Among 21 Patients. *J Allergy Clin Immunol Pract* **9**, 803-818 e811 (2021). https://doi.org:10.1016/j.jaip.2020.11.007

Rodero, M. P. *et al.* Type I interferon-mediated autoinflammation due to DNase II deficiency. *Nat Commun* **8**, 2176 (2017). https://doi.org:10.1038/s41467-017-01932-3

Li, T. *et al.* TBK1 recruitment to STING mediates autoinflammatory arthritis caused by defective DNA clearance. *J Exp Med* **219** (2022). https://doi.org:10.1084/jem.20211539

Yum, S., Li, M., Fang, Y. & Chen, Z. J. TBK1 recruitment to STING activates both IRF3 and NF-kappaB that mediate immune defense against tumors and viral infections. *Proc Natl Acad Sci U S A* **118** (2021). https://doi.org:10.1073/pnas.2100225118

Gaidt, M. M. *et al.* The DNA Inflammasome in Human Myeloid Cells Is Initiated by a STING-Cell Death Program Upstream of NLRP3. *Cell* **171**, 1110-1124 e1118 (2017). https://doi.org:10.1016/j.cell.2017.09.039

Jin, L. *et al.* STING/MPYS mediates host defense against Listeria monocytogenes infection by regulating Ly6C(hi) monocyte migration. *J Immunol* **190**, 2835-2843 (2013). https://doi.org:10.4049/jimmunol.1201788

22   Larkin, B. *et al.* Cutting Edge: Activation of STING in T Cells Induces Type I IFN
Responses   and   Cell   Death.   *J   Immunol*   **199**,   397-402   (2017).
https://doi.org:10.4049/jimmunol.1601999

23   Cerboni, S. *et al.* Intrinsic antiproliferative activity of the innate sensor STING in T
lymphocytes. *J Exp Med* **214**, 1769-1785 (2017). https://doi.org:10.1084/jem.20161674

24   Kuhl, N. *et al.* STING agonism turns human T cells into interferon-producing cells but
impedes   their   functionality.   *EMBO   Rep*   **24**,   e55536   (2023).
https://doi.org:10.15252/embr.202255536

25   Jin, L. *et al.* MPYS, a novel membrane tetraspanner, is associated with major
histocompatibility complex class II and mediates transduction of apoptotic signals. *Mol*
Cell Biol **28**, 5014-5026 (2008). https://doi.org:10.1128/MCB.00640-08

26   Gulen, M. F. *et al.* Signalling strength determines proapoptotic functions of STING. *Nat*
Commun **8**, 427 (2017). https://doi.org:10.1038/s41467-017-00573-w

27   Kabelitz, D. *et al.* Signal strength of STING activation determines cytokine plasticity and
cell death in human monocytes. *Sci Rep* **12**, 17827 (2022). https://doi.org:10.1038/s41598-
022-20519-7

28   Murthy, A. M. V., Robinson, N. & Kumar, S. Crosstalk between cGAS-STING signaling
and cell death. *Cell Death Differ* **27**, 2989-3003 (2020). https://doi.org:10.1038/s41418-
020-00624-8

29   Li, C., Liu, J., Hou, W., Kang, R. & Tang, D. STING1 Promotes Ferroptosis Through
MFN1/2-Dependent Mitochondrial Fusion. *Front Cell Dev Biol* **9**, 698679 (2021).
https://doi.org:10.3389/fcell.2021.698679

30   Song, C. *et al.* SHR1032, a novel STING agonist, stimulates anti-tumor immunity and
directly induces AML apoptosis. *Sci Rep* **12**, 8579 (2022). https://doi.org:10.1038/s41598-
022-12449-1

31   Jin, L. *et al.* Identification and characterization of a loss-of-function human MPYS variant.
Genes Immun **12**, 263-269 (2011). https://doi.org:10.1038/gene.2010.75

32   Patel, S. *et al.* The Common R71H-G230A-R293Q Human TMEM173 Is a Null Allele. *J*
Immunol **198**, 776-787 (2017). https://doi.org:10.4049/jimmunol.1601585

33 Mansouri, S. *et al.* MPYS Modulates Fatty Acid Metabolism and Immune Tolerance at
Homeostasis Independent of Type I IFNs. *J Immunol* **209**, 2114-2132 (2022).
https://doi.org:10.4049/jimmunol.2200158

34 Patel, S. & Jin, L. TMEM173 variants and potential importance to human biology and
disease. *Genes Immun* **20**, 82-89 (2019). https://doi.org:10.1038/s41435-018-0029-9

35 Liu, Y. *et al.* Activated STING in a vascular and pulmonary syndrome. *N Engl J Med* **371**,
507-518 (2014). https://doi.org:10.1056/NEJMoa1312625

36 Sebastian, M. *et al.* Obesity and STING1 genotype associate with 23-valent pneumococcal
vaccination efficacy. *JCI Insight* **5** (2020). https://doi.org:10.1172/jci.insight.136141

37 Yi, G. *et al.* Single nucleotide polymorphisms of human STING can affect innate immune
response to cyclic dinucleotides. *PLoS One* **8**, e77846 (2013).
https://doi.org:10.1371/journal.pone.0077846

38 Patel, S. *et al.* Response to Comment on "The Common R71H-G230A-R293Q Human
TMEM173 Is a Null Allele". *J Immunol* **198**, 4185-4188 (2017).
https://doi.org:10.4049/jimmunol.1700322

39 Nissen, S. K. *et al.* Multiple Homozygous Variants in the STING-Encoding TMEM173
Gene in HIV Long-Term Nonprogressors. *J Immunol* **200**, 3372-3382 (2018).
https://doi.org:10.4049/jimmunol.1701284

40 Ruiz-Moreno, J. S. *et al.* The common HAQ STING variant impairs cGAS-dependent
antibacterial responses and is associated with susceptibility to Legionnaires' disease in
humans. *PLoS Pathog* **14**, e1006829 (2018). https://doi.org:10.1371/journal.ppat.1006829

41 Ramanjulu, J. M. *et al.* Design of amidobenzimidazole STING receptor agonists with
systemic activity. *Nature* **564**, 439-443 (2018). https://doi.org:10.1038/s41586-018-0705-
y

42 Messaoud-Nacer, Y. *et al.* STING agonist diABZI induces PANoptosis and DNA mediated
acute respiratory distress syndrome (ARDS). *Cell Death Dis* **13**, 269 (2022).
https://doi.org:10.1038/s41419-022-04664-5

43 Tang, C. H. *et al.* Agonist-Mediated Activation of STING Induces Apoptosis in Malignant
B Cells. *Cancer Res* **76**, 2137-2152 (2016). https://doi.org:10.1158/0008-5472.CAN-15-
1885

44   Meric-Bernstam, F. *et al.* Combination of the STING Agonist MIW815 (ADU-S100) and
PD-1 Inhibitor Spartalizumab in Advanced/Metastatic Solid Tumors or Lymphomas: An
Open-Label, Multicenter, Phase Ib Study. *Clin Cancer Res* **29**, 110-121 (2023).
https://doi.org:10.1158/1078-0432.CCR-22-2235

45   Meric-Bernstam, F. *et al.* Phase I Dose-Escalation Trial of MIW815 (ADU-S100), an
Intratumoral STING Agonist, in Patients with Advanced/Metastatic Solid Tumors or
Lymphomas. *Clin Cancer Res* **28**, 677-688 (2022). https://doi.org:10.1158/1078-
0432.CCR-21-1963

46   Wu, J. *et al.* STING-mediated disruption of calcium homeostasis chronically activates ER
stress and primes T cell death. *J Exp Med* **216**, 867-883 (2019).
https://doi.org:10.1084/jem.20182192

47   Motwani, M. *et al.* Hierarchy of clinical manifestations in SAVI N153S and V154M mouse
models. *Proc Natl Acad Sci U S A* **116**, 7941-7950 (2019).
https://doi.org:10.1073/pnas.1818281116

48   Gao, K. M., Motwani, M., Tedder, T., Marshak-Rothstein, A. & Fitzgerald, K. A.
Radioresistant cells initiate lymphocyte-dependent lung inflammation and IFNgamma-
dependent mortality in STING gain-of-function mice. *Proc Natl Acad Sci U S A* **119**,
e2202327119 (2022). https://doi.org:10.1073/pnas.2202327119

49   Monroe, K. M. *et al.* IFI16 DNA sensor is required for death of lymphoid CD4 T cells
abortively infected with HIV. *Science* **343**, 428-432 (2014).
https://doi.org:10.1126/science.1243640

50   Doitsh, G. *et al.* Cell death by pyroptosis drives CD4 T-cell depletion in HIV-1 infection.
Nature **505**, 509-514 (2014). https://doi.org:10.1038/nature12940

51   Jakobsen, M. R., Olagnier, D. & Hiscott, J. Innate immune sensing of HIV-1 infection.
Curr Opin HIV AIDS **10**, 96-102 (2015). https://doi.org:10.1097/COH.0000000000000129

52   Silvin, A. & Manel, N. Innate immune sensing of HIV infection. *Curr Opin Immunol* **32**,
54-60 (2015). https://doi.org:10.1016/j.coi.2014.12.003

53   Altfeld, M. & Gale, M., Jr. Innate immunity against HIV-1 infection. *Nat Immunol* **16**, 554-
562 (2015). https://doi.org:10.1038/ni.3157

54 Krapp, C., Jonsson, K. & Jakobsen, M. R. STING dependent sensing - Does HIV actually
care? *Cytokine Growth Factor Rev* **40**, 68-76 (2018).
https://doi.org:10.1016/j.cytogfr.2018.03.002

55 Gubser, C., Pitman, M. C. & Lewin, S. R. CD4(+) T cell signatures in HIV infection. *Nat*
Immunol **20**, 948-950 (2019). https://doi.org:10.1038/s41590-019-0447-5

56 Morou, A. *et al.* Altered differentiation is central to HIV-specific CD4(+) T cell
dysfunction in progressive disease. *Nat Immunol* **20**, 1059-1070 (2019).
https://doi.org:10.1038/s41590-019-0418-x

57 Hines, J. B., Kacew, A. J. & Sweis, R. F. The Development of STING Agonists and
Emerging Results as a Cancer Immunotherapy. *Curr Oncol Rep* **25**, 189-199 (2023).
https://doi.org:10.1007/s11912-023-01361-0

58 Samson, N. & Ablasser, A. The cGAS-STING pathway and cancer. *Nat Cancer* **3**, 1452-
1463 (2022). https://doi.org:10.1038/s43018-022-00468-w

59 Liu, Y. *et al.* STING, a promising target for small molecular immune modulator: A review.
Eur J Med Chem **211**, 113113 (2021). https://doi.org:10.1016/j.ejmech.2020.113113

60 Zheng, J. *et al.* Comprehensive elaboration of the cGAS-STING signaling axis in cancer
development and immunotherapy. *Mol Cancer* **19**, 133 (2020).
https://doi.org:10.1186/s12943-020-01250-1

61 Corrales, L. *et al.* Direct Activation of STING in the Tumor Microenvironment Leads to
Potent and Systemic Tumor Regression and Immunity. *Cell Rep* **11**, 1018-1030 (2015).
https://doi.org:10.1016/j.celrep.2015.04.031

62 Fu, J. *et al.* STING agonist formulated cancer vaccines can cure established tumors
resistant to PD-1 blockade. *Sci Transl Med* **7**, 283ra252 (2015).
https://doi.org:10.1126/scitranslmed.aaa4306

63 Barber, G. N. Innate immune DNA sensing pathways: STING, AIMII and the regulation
of interferon production and inflammatory responses. *Curr Opin Immunol* **23**, 10-20
(2011). https://doi.org:10.1016/j.coi.2010.12.015

64 Liu, S. *et al.* Phosphorylation of innate immune adaptor proteins MAVS, STING, and TRIF
induces IRF3 activation. *Science* **347**, aaa2630 (2015).
https://doi.org:10.1126/science.aaa2630

65   Movert, E. *et al.* Interplay between human STING genotype and bacterial NADase activity
regulates   inter-individual   disease   variability.   *Nat   Commun*   **14**,   4008   (2023).
https://doi.org:10.1038/s41467-023-39771-0

66   Cai, J. *et al.* MicroRNA-206 antagomiRenriched extracellular vesicles attenuate lung
ischemiareperfusion injury through CXCL1 regulation in alveolar epithelial cells. *J Heart*
Lung Transplant **39**, 1476-1490 (2020). https://doi.org:10.1016/j.healun.2020.09.012

67   Mansouri, S. *et al.* Lung IFNAR1(hi) TNFR2(+) cDC2 promotes lung regulatory T cells
induction and maintains lung mucosal tolerance at steady state. *Mucosal Immunol* **13**, 595-
608 (2020). https://doi.org:10.1038/s41385-020-0254-1

Acknowledgments This work was supported by NIH grants HL152163 (to L.J.).

Author contributions L.J. conceived and designed the research. A.A.T, L.A.S, A.T.P, A.K.S, A,

J,B, L.J. performed experiments and analyzed the data. A.M.E helped with human lung explants.

L.J. wrote the manuscript. L.J. supervised the overall research.

Competing interests, The authors declare that they have no competing interests.

Data and materials availability All data needed to evaluate the conclusions in the paper are

present in the paper or the supplementary materials.

    Figure Captions

Figure 1. Splenocytes from *HAQ, AQ*, and *Q293* mice are resistant to STING-mediated cell death *ex vivo*. A. C57BL/6N splenocytes were treated directly (no transfection) with diABZI (100ng/ml), RpRpssCDA (5 μg/ml) or 2′3′-cGAMP (10μg/ml), DMXAA (25μg/ml) for 24hrs in

culture. CD4, CD8 T cells and CD19 B cells death were determined by PI staining. **B**. Splenocytes

from C57BL/6N mice were pre-treated with indicated small molecules, GSK2334470 (1.25µM),

GSK8612 (2.5µM), Bx-795 (0.5µM), QVD-OPH (25µM) for 2hrs. diABZI (100ng/ml) was added

in culture for another 24hrs. Dead cells were determined by PI staining. **C-D.** Flowcytometry of

HAQ, AQ, IFNAR1$^{-/-}$ or C57BL/6N splenocytes treated with diABZI (100ng/ml) for 24hrs. Cell

death was determined by PI staining. **E-F**. *Q293* or the *WT* littermates splenocytes were treated

with diABZI (100ng/ml), RpRpssCDA (5 µg/ml) or 2′3′-cGAMP 10µg/ml) for 24h. Cell death was

determined by PI staining. **G.** *WT/HAQ*, *WT/AQ*, or *WT/WT* littermates splenocytes were treated

with DMXAA (10, 25 or 100µg/ml) for 24h. Cell death was determined by PI staining. Data are

representative of three independent experiments. Graphs represent the mean with error bars

indication s.e.m. P values are determined by one-way ANOVA Tukey's multiple comparison test

(A, E, G) or unpaired student T-test (B, D). * $p<0.05$. n.s: not significant.

Figure S1. Bx-795 inhibits diABZI-induced mouse splenocyte death. A. Splenocytes from

C57BL/6N mice were treated with 100ng/ml diABZI in culture for the indicated time. Dead cells

were determined by PI and Annexin V staining. **B.** Splenocytes from C57BL/6N mice were treated

with indicated dose of diABZi in culture for 24hrs. Dead cells were determined by PI staining. **C.**

Splenocytes from C57BL/6N mice were pre-treated with indicated small molecules, H151

(10µg/ml), C176 (1µM), MCC950 (1.25µM), 2-BP (20µM), GSK872 (312.5nM), Liproxstatin-1

(1µM), necrostatin-1 (10µM), VX-765 (1µM), Z-DEVD-FMK (25µM), Bx-795 (0.5µM) for 2hrs.

diABZi (100ng/ml) was added in culture for another 24hrs0 1Dead cells were determined by PI

staining. Data are representative of three independent experiments. Graphs represent the mean with

error bars indication s.e.m. n=3~5 mice/group. *p* values are determined by one-way ANOVA

Tukey's multiple comparison test (A) or unpaired student T-test (C). * $p < 0.05$. n.s: not significant.

A Human Lung Lymphocytes (*R232/R232*) (24hrs)

Untreated | diABZi | RpRpssCDA | 2'3'-cGAMP

CD4 T cells: 29.0 | 73.8 | 54.0 | 31.5

CD8 T cells: 0.017 | 0.49 | 0.94 | 0.22

CD19 B cells: 25.3 | 21.0 | 25.9 | 26.6

PI / FSC

B STING agonists (24hrs)

% of Dead Splenocytes

Untreated / diABZi / RpRpssCDA / 2'3'-c GAMP

CD4 T cells, CD8 T cells, B cells

C CD4 T cells, diABZI (24hrs)

% of Dead Splenocytes

Untreated / 25ng/ml / 100ng/ml

WT/WT, WT/HAQ, n.s

D diABZI-induced cell death in STING-KO THP-1 cells reconstituted with human *STING* alleles (200ng/ml, 24hrs)

WT | KO | HAQ | AQ | Q293

PBS: 40.6 / 12.5 | 27.4 / 8.84 | 33.8 / 12.0 | 31.1 / 13.3 | 31.4 / 11.1

diABZI: 55.6 / 17.4 | 26.0 / 11.5 | 36.8 / 17.0 | 20.9 / 19.3 | 34.2 / 13.2

PI / Annexin V

E THP-1 cell death [diABZI, 24hrs]

% of Annexin V+ THP-1 cells

PBS / diABZI

WT, KO (n.s), HAQ (n.s), AQ (n.s), Q293 (n.s)

F THP-1 cell death [diABZI, 24hrs]

% of Annexin V+ THP-1 cells

WT / KO / Q293 / Activated STING

0, 5, 20, 100, 200 ng/ml

G THP-1 cell activation [diABZI, 2hrs]

WT / Q293

PBS | diABZI 50ng/ml | diABZI 150ng/ml | PBS | diABZI 50ng/ml | diABZI 150ng/ml

Activated STING / STING / STING / 37kDa

Stain-free gel, Loading control / 50kDa / 37kDa

H ISG54 induction [diABZI, 50ng/ml, 24hrs]

Fold of ISG54 induction normalized by IFNβ stimulation

WT, HAQ, AQ, Q293, KO

I THP-1 cell activation [diABZI, 2hrs]

WT / HAQ / AQ

PBS diABZI | PBS diABZI | PBS diABZI

p-IRF3 / 37kDa

Stain-free gel, Loading control / 50kDa / 37kDa

J THP-1 cell activation [diABZI, 2hrs]

Activated STING / WT / HAQ / AQ

PBS diABZI | PBS diABZI | PBS diABZI

STING / 37kDa

Stain-free gel, Loading control / 50kDa / 37kDa

Figure 2. *HAQ*, *AQ*, and *Q293* human cells are resistant to STING agonists-induced death.

A-B. Total human Lung cells from *WT/WT* individuals were treated with diABZi (100ng/ml) for 24hrs. Cell death in CD4, CD8 T cells and CD19 B cells were determined by PI staining. **C.** Total lung cells from a *WT/HAQ* (2 individuals) and a *WT/WT* (3 individuals) were treated with diABZi (25, 100ng/ml) for 24h. Cell death in CD4 T cells was determined by PI staining. **D-E.** STING-KO THP-1 cells (Invivogen,, cat no. thpd-kostg) were stably reconstituted with human *WT (R232), HAQ, AQ, Q293*. Cells were treated with diABZI (200ng/ml) in culture for 24 hrs. Dead cells were determined by Annexin V staining. **F.** STING-KO THP-1 cells stably reconstituted with human *WT (R232), Q293* were treated with indicated dose of diABZI for 24hs in culture. Dead cells were determined by Annexin V staining. G. STING-KO THP-1 cells stably reconstituted with human *WT (R232), Q293* were treated with indicated dose of diABZI for 2hs in culture. STING activation was detected by anti-STING antibody (Proteintech, #19851-1-AP). **H.** STING-KO THP-1 cells stably reconstituted with human *WT (R232), HAQ, AQ, Q293* were treated with 50ng/ml diABZI in culture for 24hrs. ISG-54 reporter luciferase activity was determined in cell supernatant and normalized to 10ng/ml IFNβ-stimulated ISG-54 luciferase activity. **I-J.** STING-KO THP-1 cells stably reconstituted with human *WT (R232), HAQ, AQ* were treated with 50ng/ml diABZI in culture for 2hrs. STING and IRF3 activation were determined by anti-STING antibody and anti-p IRF3 antibody (CST, Ser396, clone 4D4G). Data are representative of at least two independent experiments. Graphs represent the mean with error bars indication s.e.m. *p* values determined by one-way ANOVA Tukey's multiple comparison test (**B, C, F, G**) or unpaired student T-test (**E**). * *p*<0.05, n.s: not significant.

Figure S2. STING activation in primary human cells and THP-1 cells reconstituted with _WT_ human _STING_. A. Total human Lung cells from _WT/WT_ individuals were treated with indicated dose of diABZi for 24hrs. Cell death in CD4, CD8 T cells and CD19 B cells were determined by PI staining. **B.** Sequencing results of human _TMEM173_ gene in _WT/WT_ and _WT/HAQ_ individuals. **C-D**. STING-KO THP-1 cells stably reconstituted with human _WT (R232)_ were treated with indicated dose of diABZI for indicated time in culture. ISG-54 reporter luciferase activity was determined in cell supernatant and normalized to 10ng/ml IFNβ-stimulated ISG-54 luciferase activity. Data are representative of at least two independent experiments. Graphs represent the mean with error bars indication s.e.m. _p_ values determined by one-way ANOVA Tukey's multiple comparison test. * _p<0.05,_

Figure 3. *HAQ* **and** *AQ* **rescue the lymphopenia and suppress myeloid cell expansion in** *SAVI*

(N153S) **mice. A.** The size and weight of spleens from WT/HAQ, HAQ/SAVI, AQ/SAVI,

WT/AQ, WT/SAVI. **B-D.** Spleen $CD19^+$ B cells, CD4, CD8 T cells were determined in the

indicated mice by Flow. **E-H.** Spleen $Ly6G^+$ neutrophils, $Ly6C^{hi}$ monocytes and $F4/80^+$

macrophage were determined in the indicated mice by Flow. Data are representative of three

independent experiments. n=3~5 mice/group. Graphs represent the mean with error bars indication

s.e.m. *P* values are determined by one-way ANOVA Tukey's multiple comparison test. * *p*<0.05,

n.s: not significant.

Figure S3. *HAQ* **and** *AQ* **restore bone marrow monocytes in** *SAVI (N153S)* **mice. A-D.** Flow analysis of Ly6G⁺CD11B⁺ neutrophils and Ly6G⁻Ly6C⁺CD11B⁺ bone marrow monocytes from HAQ/SAVI, AQ/SAVI, WT/SAVI and their littermates. n=3~5 mice/group. Data are representative of three independent experiments. Graphs represent the mean with error bars indication s.e.m. *P* values are determined by one-way ANOVA Tukey's multiple comparison test. * $p < 0.05$, n.s: not significant.

Figure 4. ***HAQ*** **and** ***AQ*** **alleles prevent** ***SAVI(N153S)*** **disease in mice. A, B, G.** The size and

body weight of *HAQ/SAVI, AQ/SAVI, WT/SAVI* and their littermates *WT/HAQ, WT/AQ* mice. **D,**

E, H, I. Airway resistance, and pulmonary artery pressure were determined as described in

Materials and Methods. **C, F.** *HAQ/SAVI, AQ/SAVI, WT/SAVI* (10 mice/group), were monitored

for survival by Kaplan-Meier. **J, K.** Representative hematoxylin and eosin (H&E) staining of lung,

676　liver sections from indicated mice. n=3~5 mice/group. Data are representative of at least 2

677　independent experiments. Graphs represent the mean with error bars indication s.e.m. *P* values are

678　determined by one-way ANOVA Tukey's multiple comparison test. * $p<0.05$, **$p<0.01$. n.s.: not

679　significant. WBC: white blood cells; H: hepatocytes; K: Kupper cells; PV: portal vein; CV: central

680　vein.

Figure S4. ***HAQ, AQ* suppress lung myeloid cells infiltration in *SAVI(N153S)* mice**. **A-B.** Flow analysis of lung Ly6G$^+$CD11B$^+$ neutrophils and Ly6G$^-$Ly6C$^+$CD11B$^+$ monocytes from *HAQ/SAVI, AQ/SAVI, WT/SAVI and their littermates WT/HAQ, WT/AQ* mice. n=3~5 mice/group. Data are representative of three independent experiments. Graphs represent the mean with error bars indication s.e.m. *P* values are determined by one-way ANOVA Tukey's multiple comparison test. * $p < 0.05$.

Figure 5. *AQ/SAVI(N153S)* **cells had similar TBK1-IRF3, NFκB activation and STING**

degradation as the *WT/SAVI(N153S)* **cells. A-C.** BMDM from *WT/WT, WT/SAVI, HAQ/SAVI*

and *AQ/SAVI* were treated with 100ng/ml diABZi in culture for 2 hrs. Cells were lysed and run on

a 4~20% Mini-PROTEAN TGX Stain-Free Precast Gel. The blot was probed for phospho-Thr172-

TBK1 antibody (CST, clone D52C2), phosphor-Ser 396-IRF3 (CST, clone 4D4G), STING

(Proteintech, #19851-1-AP) and IκBα (CST, clone 44D4) antibody. **B.** BMDM from *WT/WT,*

WT/SAVI, HAQ/SAVI and *AQ/SAVI* were treated with 100ng/ml diABZi in culture for 24 hrs. Cells

were lysed and run on a 4~20% Mini-PROTEAN TGX Stain-Free Precast Gel. The blot was

probed for STING antibody (Proteintech, 19851-1-AP). **D.** BMDM from *WT/WT, WT/SAVI,*

HAQ/SAVI and *AQ/SAVI* were treated with 400μM cleavable chemical crosslinker DSP (Pierce™,

cat no: PG82081) in PBS for 1 hr at 4°C. Cells were washed with PBS and lysed in RIPA buffer.

Whole cell lysate was mixed with 4x Laemmli Sample Buffer (BioRad, cat no 1610747) containing

703   5% 2-mercaptoethanol, heated at 95°C for 10 min and, run on a 4~20% Mini-PROTEAN TGX

704   Stain-Free Precast Gel. The blot was probed for STING antibody (Proteintech, 19851-1-AP). Data

705   are representative of three independent experiments.

Figure 6. *HAQ/SAVI(N153S)* and *AQ/SAVI(N153S)* cells had 10-fold and 20-fold increased spleen T-regs compared to *WT/SAVI* mice. A. Flow cytometry analysis of IFNγ producing spleen CD4$^+$ T cells from *WT/HAQ, WT/AQ, WT/SAVI, HAQ/SAVI* and *AQ/SAVI* mice. **B**. Flow cytometry analysis of CD4$^+$ FoxP3$^+$ spleen T-regs. n=3~5 mice/group. Data are representative of at least 2 independent experiments. Graphs represent the mean with error bars indication s.e.m. *P* values are determined by one-way ANOVA Tukey's multiple comparison test. * *p*<0.05, **p<0.01, ***p<0.001. n.s.: not significant.