The spatiotemporal distribution of human pathogens in ancient Eurasia and the emergence of zoonotic diseases

Martin Sikora[1,*], Elisabetta Canteri[2], Antonio Fernandez-Guerra[1], Nikolay Oskolkov[3], Rasmus Ågren[4], Lena Hansson[5], Evan K. Irving-Pease[2], Barbara Mühlemann[6,7], Sofie Holtsmark Nielsen[8], Gabriele Scorrano[1,9], Morten E. Allentoft[1,10], Frederik Valeur Seersholm[1], Hannes Schroeder[2], Charleen Gaunitz[1], Jesper Stenderup[1], Lasse Vinner[1], Terry C. Jones[5,6,9], Björn Nystedt[12], Karl-Göran Sjögren[13], Julian Parkhill[14], Lars Fugger[15,16,17], Fernando Racimo[2], Kristian Kristiansen[1,13], Astrid K. N. Iversen[1, 17,*], Eske Willerslev[1,18,19,*]

[1]Centre for Ancient Environmental Genomics & The Lundbeck Foundation GeoGenetics Centre, Globe Institute, University of Copenhagen, Copenhagen, Denmark
[2]Section for Molecular Ecology and Evolution, Globe Institute, University of Copenhagen, Copenhagen, Denmark
[3]Department of Biology, National Bioinformatics Infrastructure Sweden, Science for Life Laboratory, Lund University, Lund, Sweden
[4]Department of Biology and Biological Engineering, National Bioinformatics Infrastructure Sweden, Science for Life Laboratory, Chalmers University of Technology, Göteborg, Sweden
[5]Definitive Healthcare, Gothenburg, Sweden.
[6]Institute of Virology, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, 10117 Berlin, Germany.
[7]German Centre for Infection Research (DZIF), partner site Charité, 10117 Berlin, Germany
[8]Department of Bacteria, Parasites and Fungi, Statens Serum Institut, Copenhagen, Denmark
[9]Center for Molecular Anthropology for the study of ancient DNA, Department of Biology, University of Rome "Tor Vergata", 00173 Rome, Italy.
[10]Trace and Environmental DNA (TrEnD) Laboratory, School of Molecular and Life Sciences, Curtin University, Perth, Australia
[11]Centre for Pathogen Evolution, Department of Zoology, University of Cambridge, Cambridge, UK
[12]Department of Cell and Molecular Biology, National Bioinformatics Infrastructure Sweden, Science for Life Laboratory, Uppsala University, Uppsala, Sweden
[13]Department of Historical Studies, University of Gothenburg, Gothenburg, Sweden.
[14]Department of Veterinary Medicine, University of Cambridge, Cambridge, UK
[15]Oxford Centre for Neuroinflammation, Nuffield Department of Clinical Neurosciences, John Radcliffe Hospital, University of Oxford, Oxford, UK
[16]MRC Human Immunology Unit, John Radcliffe Hospital, University of Oxford, Oxford, UK
[17]Nuffield Department of Clinical Neurosciences, Weatherall Institute of Molecular Medicine, University of Oxford, Oxford, UK
[18]Department of Genetics, University of Cambridge, Cambridge, UK
[19]MARUM Center for Marine Environmental Sciences and Faculty of Geosciences, University of Bremen, Bremen, Germany

* Corresponding authors; email: martin.sikora@sund.ku.dk;
astrid.iversen@ndcn.ox.ac.uk;ewillerslev@sund.ku.dk

Summary

Infectious diseases have had devastating impacts on human populations throughout history, but important questions about their origins and past dynamics remain[1]. To create the first archaeogenetic-based spatiotemporal map of human pathogens, we screened shotgun sequencing data from 1,313 ancient humans covering 37,000 years of Eurasian history. We demonstrate the widespread presence of ancient bacterial, viral and parasite DNA, identifying 5,486 individual hits against 492 species from 136 genera. Among those hits, 3,384 involve known human pathogens[2], many of which were detected for the first time in ancient human remains. Grouping the ancient microbial species according to their likely reservoir and type of transmission, we find that most groups are identified throughout the entire sampling period. Intriguingly, zoonotic pathogens are only detected ~6,500 years ago, peaking ~5,000 years ago, coinciding with the widespread domestication of livestock[3]. Our findings provide the first direct evidence that this lifestyle change resulted in an increased infectious disease burden. Importantly, they also suggest that the spread of these pathogens increased substantially during subsequent millenia, coinciding with the pastoralist migrations from the Eurasian Steppe[4,5].

Introduction

Pathogens have been a constant threat to human health throughout our evolutionary history. Until approximately 1850, at least a quarter of all children died before age one, and around another quarter before turning 15. Infectious diseases are estimated to have been responsible for over half of these deaths[6]. Larger disease outbreaks have profoundly impacted human societies, sometimes devastatingly affecting entire civilizations[7]. Infectious diseases have left lasting impressions on human genomes, as selective pressures from pathogens have continuously shaped human genetic variation[8–10]. Where and when different human pathogens first emerged, how and why they spread, and how they affected human populations are important but largely unresolved questions.

During the Holocene (beginning ~12,000 years ago), the agricultural transition created larger and more sedentary communities, facilitating pathogen transmission and persistence within populations[11]. Simultaneously, the rise of animal husbandry and pastoralism are thought to have increased the risk of zoonoses[3]. Technological advances, such as horses and carts, increased both mobility and the risk of disease transmission between populations[12]. It has been hypothesised that these changes led to the so-called "first epidemiological transition" characterised by increased infectious disease mortality[3]. However, direct evidence remains scarce and the idea is debated[13]. Paleopathological examinations of ancient skeletons offer insights into past infectious disease burden[14], but are limited to few diseases identifiable from the available tissue. Recent advances in ancient DNA (aDNA) techniques allow for the retrieval of direct genomic evidence of past microbial infections, which can enable the reconstruction of complete ancient pathogen genomes. These studies have typically concentrated on specific pathogens and have provided surprising insights into the evolutionary history of the causative agents of some of the most historically important infectious diseases affecting humans, including plague (*Yersinia pestis*)[12,15–23], tuberculosis (*Mycobacterium tuberculosis*)[24,25], smallpox (*Variola virus*)[26,27], Hepatitis B (*Hepatitis B virus*)[28–30] and others[31–36]. However, there is an unmet need to investigate the combined landscape of ancient bacteria, viruses, and parasites that impacted our ancestors across various regions and time periods. Here we use a new high-throughput computational workflow to screen for ancient microbial DNA and use our data to investigate long-standing questi no ons in paleoepidemiology: When and where did important human pathogens arise? And what factors influenced their spatiotemporal distribution?

To understand the distribution of ancient pathogenic challenges, we developed an accurate and
scalable workflow to identify ancient microbial DNA in shotgun-sequenced aDNA data (Extended
Data Figs. 1-4; Supplementary information 1). The data (~405 billion sequencing reads) derived from
1,313 ancient individuals from Western Eurasia (n=1,015; 77%), Central and North Asia (n=265;
20%) and Southeast Asia (n=33; 3%), spanning a ~37,000 year period, from the Upper Paleolithic to
historical times (Fig. 1b; Supplementary table S1; Supplementary information 2). As burial practices
varied across cultures and time, these samples represent a subset of groups within past societies.
Nevertheless, the identified pathogens likely affected the broader population, as diseases spread easily
in communities with poor sanitation and hygiene[37]. Initial metagenomic classification showed a large
fraction of reads classified as soil-dwelling taxa including genera such as *Streptomyces* or
Pseudomonas, reflecting a predominantly environmental source of microbial DNA. Further
characterization using topic-model however suggested that microbial DNA in ancient tooth samples
often derives from genera commonly associated with the human oral microbiome such *Actinomyces*
or *Streptococcus* (Extended Data Fig. 1d-g).
106

Fig. 1. Dataset overview. Spatiotemporal distribution and site contexts of the study samples. White squares in the
geographic maps indicate locations of the full set of 1,313 study samples, whereas coloured circles highlight location and
age of samples from the time period and region indicated in the respective panel. Bar plots show numbers of samples for
different site type contexts in each region (nd - not determined).

112
We selected a set of 136 bacterial and protozoan genera (11,553 species total) containing human
pathogenic species[2] as well as 1,356 viral genera (259,979 species total) for further authentication and
detection of ancient taxa. We found that ancient microbial DNA was widely detected, with 5,486
authenticated individual hits identified across 1,005 samples (Z-score for aDNA damage rate from
metaDMG $\geq$ 1.5; Fig. 2a; Supplementary table S2; Extended Data Fig. 4). Of those, 3,384 hits were
found among 214 known human pathogen species[2], with the remaining 2,104 hits involving 278 other
species. The highest numbers were observed in bacterial genera associated with the human oral

microbiome, such as *Actinomyces* (380; 28.5% of samples) and *Streptococcus* (242; 18.1% of
samples) or those commonly found in soil environments, such as *Clostridium* (252; 18.9% of
samples) and Pseudomonas (111; 8.3% of samples).
123
We observed marked differences in the distributions of the genetic similarity of the ancient microbial
sequences to their reference assemblies, both among genera and between species within a genus (Fig.
2b; Extended Data Fig. 5).. High average nucleotide identity (ANI) indicates that ancient microbial
sequences are closely related to a reference assembly in the modern database, and was observed in
hits across all species from some genera (e.g. *Yersinia,* Fig. 2b). In other genera, only a few hits had a
closely related database reference assembly match. An example is the genus *Mycobacterium,* where
only hits of the leprosy-causing bacterium *Mycobacterium leprae* were highly similar to their
reference assembly (ANI > 99%; Fig. 2b). Low ANI indicates that the ancient microbial DNA is only
distantly related to the reference assembly, for example, due to aDNA damage, poor representation of
the diversity of the genus in the database or false-positive classification of ancient microbial reads
deriving from a related genus (Extended Data Fig. 3). Alternatively, ANI can also be reduced when
reads mapped to a particular reference assembly originate from multiple closely related strains or
species in a sample. To test for such mixtures, we quantified the rate of observing different alleles at
two randomly sampled reads at nucleotide positions across the genomes of hits with read depth ≥1X.
We found a high rate of multiple alleles in many species associated with the human oral microbiome,
such as *Streptococcus sanguinis* or *Treponema denticola*. Hits for these species also showed lower
ANI, consistent with the expectation for mixtures of ancient microbial DNA (Extended Data Fig. 6b,
c).
142

143
144 **Fig. 2. Overview and characteristics of detected ancient microbial DNA. a,** Barplot showing total number of putative
145 ancient microbial hits (overall detection rate in brackets) for bacterial, eukaryotic and viral (n ≥ 4) genera. Bar colour and
146 shading distinguishes counts in the different aDNA damage categories. **b, c** Distributions of ANI (**b**) and log10-fold change
147 of mapped reads over median of reads classified at taxonomic rank of genus per sample (**c**) for individual species hits in
148 selected example genera. Symbol colour indicates aDNA damage category.

149
The rate of read mapping varied by orders of magnitude between species, from hits in species with
high read recruitment, such as *Mycobacterium leprae* (> 100-fold enrichment over the median number
of classified reads across target genera) to hits at the lower limits of detection, e.g., for the louse-
borne pathogen *Borrelia recurrentis* (lowest read recruitment ~100-fold less than the median number
of classified reads across target genera; Fig. 2c; Extended Data Fig. 5b). Ancient microbial DNA from
species commonly found in soil, such as *Clostridium botulinum,* was detected at similar rates in tooth
and bone samples. Conversely, species associated with the human oral microbiome (e.g.,
Fusobacterium nucleatum, Streptococcus mutans and *Porphyromonas gingivalis*) or pathogenic
infections (e.g., *Yersinia pestis* and *Hepatitis B virus*) were significantly more frequently identified in
tooth samples (Extended Data Fig. 6a). To further verify hits with low read numbers, we performed a
BLASTn search for all reads of each hit with N ≤ 100 final reads (n=712 hits total; Supplementary
table S3). Most hits showed a high proportion (≥ 80%) of reads assigned to the same species using
BLASTn, and the species with the most top-ranked BLASTn hits generally matched the inferred hit
species (Extended Data Fig. 7a, b).

164
165 Our results show that ancient microbial DNA isolated from human remains originates from complex
166 mixtures of distinct endogenous and exogenous sources. The high detection rate, high read

recruitment, lower ANI, and evidence of mixtures in genera such as *Clostridium* or *Pseudomonas* (Fig 2.; Extended Data Fig. 5, 6) suggest that a substantial fraction of this ancient microbial metagenome derives from environmental sources, possibly associated with the "necrobiome" involved in post-mortem putrefaction processes (Supplementary information 3)[38,39]. By contrast, species from other frequently observed genera, including *Actinomyces* or *Streptococcus,* were predominantly identified from teeth and likely originated from the endogenous oral microbiome[38]. Species representing likely cases of pathogenic infections (e.g., *Yersinia pestis* and *Mycobacterium leprae*) were often characterised by higher ANI and/or low multi-allele rate, consistent with pathogen load predominantly originating from a single dominant strain.

The landscape of ancient pathogens across Eurasia

Our dataset provides a unique opportunity to investigate the origins and spatiotemporal distribution of major human pathogens in Eurasia, expanding the known range of some ancient pathogenic species and identifying others for the first time using paleogenomic data (Supplementary tables S3, S5).

Considering bacterial pathogens, we found widespread distribution of the plague-causing bacterium *Yersinia pestis*, consistent with previous studies[12,16,17,19,22,41,42]. We identified 42 putative cases of *Y. pestis* (35 newly reported; Extended Data Fig. 6e), corresponding to a detection rate of ~3% in our samples. These newly identified cases expand the spatial and temporal extent of ancient plague over previous results (Fig. 3). The earliest three cases were dated between approximately 5,700-5,300 calibrated years before present (cal. BP), across a broad geographic area ranging from Western Russia (NEO168, 5,583-5,322 cal. BP), to Central Asia (BOT2016, 5,582-5,318 cal. BP), and to Lake Baikal in Siberia[43] (DA342, 5,745-5,474 cal. BP). This broad range of detection among individuals pre-dating 5,000 cal. BP challenges previous interpretations that early plague strains represent only isolated zoonotic spillovers[20]. We replicated previously identified cases of plague in Late Neolithic and Bronze Age (LNBA) contexts across the Eurasian Steppe[16] and identified many instances where multiple individuals from the same burial context were infected (Afanasievo Gora, Russia; Kytmanovo, Russia; Kapan, Armenia; Arban 1, Russia) (Supplementary table S2). These results indicate that the transmissibility and potential for local epidemic outbreaks for strains at those sites were likely higher than previously assumed[20]. Finally, 11 out of 42 cases were identified in late mediaeval and early modern period individuals (800-200 BP) from two cemeteries in Denmark (Aalborg, Randers), highlighting the high burden of plague during this time in Europe. All but one hit (NEO627, n=84 reads total) showed expected coverage for the virulence plasmids pCD1 and pMT1, with hits before 2,500 years BP characterized by the previously reported absence of a 19 kilobase region on pMT1 containing the *ymt* gene[16] (Extended Data Fig. 7c; Supplementary information 4).

202
203 **Fig. 3. Spatiotemporal distribution of selected ancient pathogens.** Each panel shows geographic distribution (top) and
204 timeline (bottom) for identified cases of the respective pathogen (indicated by coloured circle). Geographic locations and age
205 distributions of all 1,313 study samples are shown in each panel using white squares. The panel for *Plasmodium* combines
206 the three species detected (*P. vivax* n=5; *P. malariae* n=3; *P. falciparum* n=1).

207
Another bacterial pathogen frequently detected was the spirochaete bacterium *Borrelia recurrentis*,
causative agent of louse-borne relapsing fever (LBRF), a disease with a mortality of 10-40%
(Supplementary information 5)[44]. While previous paleogenomic evidence for LBRF is limited to a
few cases from Scandinavia and Britain[31,45], we report 34 new putative cases (2.5% detection rate;
Extended Data Fig. 6e), with wide geographic distribution across Europe, Central Asia, and Siberia
(Fig. 3). We detected the earliest case in a Neolithic farmer individual from Scandinavia (NEO29,
Lohals, 5,647-5,471 cal. BP), suggesting that human body lice were already vectors for infectious
disease during the Neolithic period, supported by phylogenetic analyses of *B. recurrentis* in a recent

preprint[45]. The highest detection rates were found during the Iron and Viking Ages. LBRF outbreaks
were historically associated with crowded living conditions, poor personal hygiene, and wet and cold
seasons, but are rare today in most regions (Supplementary information 5)[46]. Our results suggest that
B. recurrentis infections exerted a substantial disease burden on past populations.
220
We also report novel cases of other bacterial pathogens previously detected in paleogenomic data.
The leprosy-causing bacterium *Mycobacterium leprae* was identified in seven individuals (0.5%
detection rate) from Scandinavia and only appeared from the Late Iron Age onwards (earliest case
RISE174, 1,523-1,339 cal. BP). Because *M. leprae* can infect both red squirrels and humans[47], and
archaeological evidence demonstrates that fur trade from Scandinavia, including squirrel fur,
increased substantially during the late Iron and Viking Ages[48], our results support the suggestion that
squirrel fur trade could have facilitated transmission[49]. Our findings are also consistent with the
widespread distribution of leprosy in mediaeval Europe[50]. We further detected three putative cases of
Treponema pallidum - subspecies of which are the causative agents of treponematoses such as yaws,
and endemic and venereal syphilis - in three individuals from recent time periods (earliest case
101809T, Denmark, 600-500 BP; Fig. 3). Two cases were identified in individuals from Borneo in
Southeast Asia (approximately 500-300 years BP); to our knowledge the first paleogenomic evidence
for treponemal disease from this region.
234
Among the species reported for the first time using paleogenomic data, we identified twelve putative
cases of *Yersinia enterocolitica,* the causative agent of yersiniosis, commonly contracted through
consuming contaminated raw or undercooked meat (Fig. 3). The animal reservoirs for *Y.*
enterocolitica include boars, deer, horses, cattle, and sheep. As *Y. enterocolitica* rarely enters the
bloodstream, our results likely underestimate the disease burden. Interestingly, this species includes
some of the only identified putative zoonotic infections in individuals from Mesolithic hunter-gatherer
contexts (NEO941, Denmark, 6,446-6,302 cal. BP). We also detected other members of the order
Enterobacterales, transmitted via the fecal-oral route, including members of the genera *Shigella*,
Salmonella, and *Escherichia* (Supplementary table S2). We report the first evidence for ancient
leptospirosis (genus *Leptospira*) dating back to the Neolithic, 5,650-5,477 cal. BP (NEO46, Sweden;
Leptospira borgpetersenii). While earlier cases predominantly involved *Leptospira borgpetersenii*
(n=5, 0.4% detection rate), the majority of hits were *Leptospira interrogans* (n=20, 1.5% detection
rate), almost exclusively in Scandinavian contexts from the Viking Age onwards (Fig. 3). *Leptospira*
borgpetersenii is today primarily found in cattle, while *Leptospira interrogans* is detected more
broadly in both domestic and wild animals. Although the clinical manifestations are similar, with an
untreated fatality rate of 1% today, transmission routes vary[51]. While host-to-host transmission
predominates for *Leptospira borgpetersenii*, transmission via urine-contaminated environments
dominates for *Leptospira interrogans* transmission. We also report two putative cases of
Corynebacterium diphtheriae, the causative agent of diphtheria; the oldest of which dates back to the
Mesolithic (Sidelkino, 11,336-11,181 cal. BP) (Supplementary table S2).
255
Other diseases associated with animals and livestock, such as listeriosis (*Listeria monocytogenes*) and
brucellosis (genus *Brucella*), could not be reliably identified. Another major human pathogen not
identified in our dataset is *Mycobacterium tuberculosis,* which causes tuberculosis (TB). However, as
the *Mycobacterium tuberculosis* load in blood is typically low in immunocompetent patients without
advanced disease[52] and latent TB develops in 60% of cases and can persist for decades, it is, based on
current knowledge, unlikely to be readily identified using aDNA data from tooth and bone remains
sampled for ancient human DNA.
263

Identifying eukaryotic pathogens is challenging as sequence contamination from other organisms
frequently occurs in their large and often fragmented reference genomes[53]. An illustrative example in
our dataset is the protozoan parasite *Toxoplasma gondii*, which we readily identified in hits with high
ANI and aDNA damage but low support from coverage evenness statistics, due to reads mapping to
short contigs representing human contamination (Extended Data Fig. 4a,b; Supplementary Data 1).
Despite these challenges, we identified nine putative malaria infections across three different human-
infecting species (*P. vivax* n=5; *P. malariae* n=3; *P. falciparum* n=1; Fig. 3; Supplementary table S2).
The most widely detected parasite species was *P. vivax,* with the earliest evidence in a Bronze Age
individual from Central Europe (RISE564, 4,750-3,750 BP based on archaeological context). Other
cases include a mediaeval individual from Central Asia (DA204, Kazakhstan; 1,053-1,025 cal. BP)
and two Viking Age individuals from Eastern Europe (VK224, 950-750 BP and VK253, 950-850 BP;
Russia). The *P. vivax* malaria vector *Anopheles atroparvus* is currently widespread in Europe and
nearby regions, including the Pontic Steppe, and our cases suggest this was also true in the past[54,55].
The single case of *P. falciparum* malaria was found in a sample from Armenia (NEO111; 463-0 cal.
BP), where malaria was eliminated in the 1960s[56].
279
Among DNA viral species, we found widespread infections with *Hepatitis B virus* (HBV; 28 cases,
2.1% detection rate), consistent with previous studies[28–30] (Extended Data Fig. 6e). Our newly
reported HBV cases include individuals from Mesolithic (Kolyma River, n=1) and Neolithic (Lake
Baikal, n=3) contexts in Siberia dating back to 9,906-9,665 cal. BP, providing the first evidence for
ancient HBV from those regions (Fig. 3). We also report the first putative ancient case (n=1) of
Torque teno virus (TTV) dating back ~7,000 years (NEO498, Ukraine; 7,161-6,950 cal. BP). TTV
infects approximately 80% of the human population today, and while it is not associated with any
particular disease, it replicates rapidly in immunocompromised individuals[57]. Other ancient virus hits
included viruses not known to infect humans, such as ancient phage DNA (e.g,. *Escherichia phage*
T4, Proteus virus Isfahan; Supplementary table S2) and one putative case of an ancient insect virus
(*Invertebrate iridescent virus 31 (IIV-31))* in a tooth sample of a Viking Age individual from Sweden
(VK30, Varnhem; 950-650 BP)[58]. The virus source is likely exogenous, potentially originating from
aDNA of food sources in the tooth remains.
293
Co-infections with multiple pathogens can worsen disease progression and outcomes[59] and they were
likely an important morbidity factor in ancient human populations. Searching for individuals showing
co-occurrence of distinct ancient microbial species, we identified 15 cases of putative co-infections in
our dataset (Supplementary table S2). A striking case was a Viking Age individual from Norway
(VK388), where we replicated previous results of infection with a likely smallpox-causing variola
virus[27] and additionally found evidence of infection with the leprosy-causing bacterium
Mycobacterium leprae. Another case of possible co-infection with *Mycobacterium leprae* was found
in VK366, a Viking Age individual from Denmark, who also showed evidence for leptospirosis
(*Leptospira interrogan*s). Interestingly, among the 15 cases, six involved co-infections of HBV with
non-viral pathogens (*Yersinia pestis* n=3; *Borrelia recurrentis* n=2; *Plasmodium malariae* n=1;
Supplementary table S2). This suggests that some of these cases involved chronic hepatitis, possibly
reflecting HBV infection during infancy, when hepatitis becomes chronic in 90-95% of modern cases,
compared to only 2-6% in adult infections. An intriguing early case of a possible co-infection was
found in a Mesolithic hunter-gatherer from Russia (Sidelkino, 11,336-11,181 cal. BP). This individual
showed evidence of the respiratory pathogen *Corynebacterium diphtheriae*, and *Helicobacter pylori*,
usually restricted to gastric infections; however, rare contemporary examples of bacteremia have been
reported for both[60,61]. Overall, our results show that co-infections can be detected using ancient
metagenomic screening, but are likely underestimated given methodological limitations such as

differences in pathogen load, tissue availability, and other factors impacting detectability of ancient
microbial DNA.

Temporal dynamics and drivers of epidemic pathogens

Understanding the factors affecting the dynamics of past epidemics is a major aim of
paleoepidemiology. Our dataset allows us to address this question using direct molecular evidence for
ancient pathogens across prehistory. To investigate changes in pathogen incidence over time, we
performed Bayesian change-point detection and time series decomposition[62] on two pathogens with
high detection rates, *Yersinia pestis* (plague) and *Borrelia recurrentis* (LBRF), using the detection
rate of the respective pathogen as a proxy for its incidence (Methods). For plague, we inferred a
gradual rise in detection rate starting from ~6,000 BP, about 1,000 years after the estimated time to
the most recent common ancestor of currently known ancient strains (7,100 cal. BP)[20]. It reached a
first peak around ~5,000 BP across Europe and the Eurasian Steppe, coinciding with the emergence
and early spread of the LNBA- strains, believed to have had limited flea-borne transmissibility[16,17,22]
(Fig. 4). Detection remained high with additional peaks for a ~3,000 year period, until an abrupt
change ~2,800 BP led to a ~800 year period where plague was only detected in one sample (VK522,
Oland, Sweden 2,343-2,154 cal. BP). Starting at ~2,000 BP, plague reappeared in three samples from
Central Asia (DA92, DA101, DA104, Kazakhstan and Kyrgyzstan; Fig. 3; Supplementary table S2),
just before the first historically documented plague pandemic (Fig. 4). Another hiatus of ~600 years
led to a rise and peak associated with the second plague pandemic ~600 BP (European late mediaeval
cases, Denmark and previously published cases[63]; Fig. 4). This pattern of change coincides with the
extinction of the LNBA- strains ~2,700 BP[22] and the second *Yersinia pestis* diversification event
starting ~3,700 BP, which gave rise to an extinct Bronze Age lineage (RT5, LNBA+)[18] and present-
day lineages; these had increased flea-mediated transmission adaptations favouring bubonic plague
and led to all known later plague pandemics[64]. The adaptations included acquiring two plasmids: one
with the *ymt* gene for survival in the flea midgut and another with the *pla* gene for invasiveness after
transmission[65]. The lack of detection during both periods is also seen in publicly available ancient
Yersinia pestis genomes from other Eurasian sites[63], suggesting that sampling bias is unlikely to
significantly influence the observed dynamics.
340

341
342 **Fig. 4. Bayesian time series decomposition of major epidemic pathogens.** Panels show estimated trendlines and 95%
343 credible interval for detection rates (top), probability distributions and locations (dotted lines) for change points (middle) and
344 probability of trend slope (bottom) being positive (red), negative (blue) or zero (white), inferred using Bayesian change-
345 point detection and time series decomposition. Top of panels show temporal distributions of newly reported pathogen hits
346 (blue circles) as well as previously published ancient pathogens (grey circles) from the respective species.
347

The inferred temporal dynamics of LBRF show a first peak in detection around 5,500 BP, slightly
more recent than for plague, but with more sporadic occurrences and sharper peaks during the first
~2,000 years (Fig. 4). The geographic extent during the early period ranges from Scandinavia
(NEO29, Denmark, 5,647-5,471 cal. BP) to the Altai mountains (RISE503, Russia, 3,677-3,461 cal.
BP) (Fig. 3; Supplementary table S2). From ~2,800 BP, LBRF was detected more consistently,
peaking approximately 2,000 years ago, predominantly in the Eurasian Steppe region (Fig. 3). This
change from epidemic outbreaks to endemicity overlaps in time with the estimated emergence of a
distinct *Borrelia recurrentis* Iron Age clade[45] (Supplementary information 5). The period of high
LBRF detection coincided with a time without detectable plague activity (Fig. 4), reinforcing that the
absence of plague is not due to sample size limitations or poor DNA preservation. This opposing
pattern is unlikely to result from any cross-immunity between *Yersinia pestis* and *Borrelia recurrentis*
but could plausibly, in part, be caused by population size decreases and behavioural and societal
adjustments during plague epidemics. LBRF remained detectable until the end of the time series,
particularly in Europe; the continued presence might have facilitated the emergence of a Medieval *B.*
recurrentis clade ~600 years BP[45] (Supplementary information 5) (Fig. 3, 4).
363
A striking feature shared in the temporal dynamics of plague and LBRF was the absence of detectable
cases before ~6,000 BP, coinciding with a transition of individuals in predominantly hunter-gatherer
contexts to those in farming or pastoralist cultural contexts. It has been hypothesized that this
transition led to a higher risk of zoonotic disease transmission, and facilitated the spread of both old
and new pathogens[3]. Our dataset allows us to test this hypothesis using molecular evidence for
infectious disease burden. To increase power to detect changes in the load of different pathogen types,
we focused on grouped ancient microbial hit categories (Supplementary table S4).
371
We found that species associated with the ancient oral microbiome showed the highest relative
detection rate, accounting for up to 50% of ancient hits across various periods (Fig. 5a; Extended Data
Fig. 8). Species in the "environmental" classes of likely exogenous origins were also detected at
consistent rates throughout time. Species in the "infection" classes occurred at low detection rates
throughout (mostly <10%). Strikingly, we found that species in the "zoonotic" reservoir classes were
not detected until approximately 6,500 BP (Fig. 5a). Using Bayesian time series decomposition[62], we
inferred an overall increase in the detection rates of the "zoonotic" reservoir classes from ~6,000 BP,
thereafter remaining at elevated levels until the mediaeval period (Fig. 5b; Extended Data Fig. 8, 9a).
While species in the "anthroponotic" reservoir classes also occur earlier (predominantly species with
human-to-human transmission, Extended Data Fig. 9a), we observe increased detection rates from
~2,500 BP onwards (Fig 5b, Extended Data Fig. 8). Our results provide the first direct evidence for an
epidemiological transition of increased infectious disease burden after the onset of agriculture through
to historical times.
385

386
387 **Fig. 5. Time series of ancient microbes by microbial source. a,** Timeline of relative detection rates in sliding windows of
388 21 temporally consecutive samples, for different ancient microbial species classes. Coloured horizontal lines indicate the
389 expected rates if species in all classes would be detected at equal rates, based on the total number of distinct species in each
390 class. **b,** Trendlines for detection rates inferred using Bayesian change-point detection and time series decomposition, for
391 ancient microbial species in the "zoonotic" (top) and "anthroponotic" (bottom) reservoir class.
392
We used Bayesian spatiotemporal modelling[67] to investigate possible drivers of the observed ancient
microbial incidences. We modelled the presence/absence of either individual microbial species or
combined species groups using sets of putative covariates, including spatiotemporal variables
(longitude, latitude, and sample age), paleoclimatic variables (mean annual temperature and
precipitation), human mobility and ancestry, sample material (tooth or other), and a proxy for
"detectability" (number of human-classified reads). In the models for the "zoonotic" or
"anthroponotic" infection species classes, sample age was an important predictor (Fig. 6; Extended
Data Fig.10; Supplementary table S6), consistently negatively associated with incidence, and high
effect sizes in the individual species models for *Borrelia recurrentis* and *Leptospira interrogans* (Fig.
6, Extended Data Fig. 10). Longitude was another important factor in the "infection" classes; it was
positively associated with incidence rates for the combined "anthroponotic" class, and in individual
models for *Yersinia pestis* and *Borrelia recurrentis*. The positive effect of longitude suggests a higher
incidence in the eastern part of our spatiotemporal range, where samples from the Eurasian Steppe
predominate.
407
The increased infection incidence in Steppe populations could reflect an increased genetic
susceptibility or a higher risk of acquiring diseases associated with the pastoralist lifestyle. The latter
suggestion seems more plausible as continued exposure to selective pressures from certain infectious
diseases likely would reduce susceptibility in these populations. Human ancestry showed small but
consistent positive effects in some models, particularly the infection classes, for the Caucasus hunter-
gatherers (CHG). Across all models, the incidence of ancient microbes was positively associated with
teeth as sample material; the highest effect sizes were found in the "oral microbiome" and "infection"

classes (Fig. 6, Extended Data Fig. 10). Teeth preserved ancient oral microbiome and pathogen DNA better than petrous bones (the source of 86% of our samples), likely due to oral cavity exposure and better access to microbial DNA in the bloodstream[68]. These results support the notion that species detected in those classes are predominantly of endogenous origin.

Fig. 6. Predictors of ancient microbial species incidence. Matrix showing effect sizes and of 12 potential predictors (columns) for presence of selected combined ancient microbial species groups inferred from spatiotemporal modelling. For each class, the model with lowest Watanabe–Akaike information criterion is shown. Symbols indicate the predictors included in the respective model. Predictors with positive effect (2.5% and 97.5% posterior quantiles both positive) are shown as red triangles, whereas predictors with negative effect (2.5% and 97.5% posterior quantiles both negative) are shown as blue inverted triangles. Predictors included in the best-fitting model but without effect (posterior quantile range spanning zero) are indicated using white circles. Posterior standard error of effect sizes is indicated by error bars.

Conclusions

During the Holocene, human lifestyles changed significantly as agriculture, animal husbandry, and pastoralism became key practices but the impact on infectious disease incidence is debated. Our study represents the first large-scale characterization of ancient pathogens across Eurasia, providing clear evidence that identifiable zoonotic pathogens emerged around 6,500 years ago and were consistently detected after 6,000 years ago. While zoonotic cases likely existed before 6,500 years ago, the risk and extent of zoonotic transmission probably increased with the widespread adoption of husbandry

435 practices and pastoralism. Today, zoonoses account for over 60% of newly emerging infectious
436 diseases[70].
437
438
Strikingly, we observed some of the highest detection rates at ~5,000 BP, a time of significant
demographic changes in Europe due to the migration of Steppe pastoralists and the displacement of
earlier populations[4,5]. Steppe pastoralists, through their long-term continuous exposure to animals,
likely developed some immunity to certain zoonoses and their dispersals may have carried these
diseases westward and eastwards. Consequently, the genetic upheaval in Europe could have been
facilitated by epidemic waves of zoonotic diseases causing population declines, with depopulated
areas subsequently being repopulated by opportunistic settlers who intermixed with the remaining
original population. This scenario would mirror the population decline of Indigenous people in the
Americas following their exposure to diseases introduced by European colonists[71,72]. Our findings
support the interpretation of increased pathogen pressure as a likely driver of positive selection on
immune genes associated with the risk of multiple sclerosis in Steppe populations ~5,000 years ago[73],
and immune gene adaptations having occurred predominantly after the onset of the Bronze Age in
Europe[10].
452
Expanding our analyses to the broader pathogen landscape allowed us to infer and contrast incidence
patterns between different species and types of pathogens to a greater extent than previously possible.
If ancient pathogen DNA of a single species is not detected in a particular region or period, asserting
whether this is due to low disease incidence or confounding factors such as differential DNA
preservation between different periods and environments is challenging. Our analyses counter these
limitations; we demonstrate that pathogens with known epidemic potential and high detection rates,
such as *Yersinia pestis* (plague) and *Borrelia recurrentis* (LBRF)*,* show striking differences in their
detection rate over time, suggesting that low detection rate in these cases represent an actual reduction
in incidence. During the early period (~5,700-2,700 years ago), the continuous detection of *Yersinia*
pestis is suggestive of endemic disease. The succeeding pattern of distinct waves and periods without
detection indicate epidemic outbreaks; these detection peaks match the historically described plague
pandemics. This shift from endemic to epidemic is concurrent with significant changes in the *Yersinia*
*pesti*s genome, particularly increased flea-transmissibility and pathogenicity[16,18]. The pattern for
Borrelia recurrentis is almost entirely the opposite, with narrow peaks and long periods without
detection, suggesting local epidemics before ~2,700 years ago and consistent detection afterwards.
This later endemicity of LBRF could be driven by changes in the bacterial genome and by human and
environmental factors known to increase the risk of louse infestation[45,74,66]. Experimental studies have
demonstrated that *Yersinia pestis,* like *Borrelia recurrentis,* can infect body lice in the midgut, and
sometimes, also the Pawlowsky glands (PG), a putative salivary gland[66]. Body lice infected in the PG
can transmit *Yersinia pestis* in concentrations sufficient to initiate disease in humans, possibly
contributing to transmission during plague outbreaks. Infected body lice have higher mortality than
uninfected lice, and it remains unknown whether co-infection of body lice with *Yersinia pestis* and
Borrelia recurrentis is possible.
476
Our study has some important limitations. While ancient shotgun metagenomic data offers direct
evidence of past infections, its usefulness depends on having a high pathogen load and the right tissue
samples. Our ancient tooth and bone samples are well suited to detect high-load bloodstream
infections like *Yersinia pestis* and *Borrelia recurrentis*, but pathogens with lower loads or different
tissue preferences are underrepresented. Moreover, differentiating ancient infections from those
arising from environmental sources, like the "necrobiome," is challenging. Finally, our dataset lacks

information on RNA viruses, therefore underestimating the zoonotic disease burden. However, the
timing is probably accurate as the conditions favouring zoonotic transmission of RNA viruses are
similar to those of other zoonotic pathogens[69].
486
Our findings represent the first example of how the nascent field of genomic paleoepidemiology can
create a map of the spatial and temporal distribution of diverse human pathogens over millennia. This
map will develop as more ancient specimens are investigated, as will our abilities to match their
distribution with genetic, archaeological, and environmental data. Our current map shows clear
evidence that lifestyle changes in the Holocene led to an epidemiological transition, resulting in a
greater burden of zoonotic infectious diseases. This transition profoundly affected human health and
history throughout the millennia and continues to do so today.
494

Methods

We compiled a dataset of aDNA shotgun-sequencing data from 1,313 ancient individuals previously
sequenced for studies of human population history (references for previous publications describing
laboratory procedures and sample/site descriptions in Supplementary table S1). To facilitate ancient
microbial DNA authentication, we excluded sequencing libraries subjected to UDG treatment which
removes characteristic aDNA damage patterns from further analyses. Samples sequenced across
multiple libraries were combined into single analysis units to maximise sensitivity for detection of
ancient microbial DNA present in low abundance.

Ancient microbial DNA screening

We carried out screening for ancient microbial DNA using a computational workflow combining *k*-
mer-based taxonomic classification, read mapping and aDNA authentication. We first performed
taxonomic classification of the sequencing reads (minimum read length 30 bp) using *KrakenUniq*[75],
against a comprehensive database of complete bacterial, archaeal, viral, protozoan genomes in the
RefSeq database (built with default parameters of *k*-mer size 31 and low-complexity sequences
masked). To increase sensitivity for ancient viral DNA, we re-ran the classification on a viral-specific
database of complete viral genomes and neighbour assemblies from RefSeq
(https://www.ncbi.nlm.nih.gov/genome/viruses/about/assemblies/), using all reads classified as non-
human from the previous run.
514
Following this initial metagenomic classification, a subset of genera was further processed in the
genus-level read mapping and authentication stages. For bacterial pathogens, we selected genera with
two or more established species of human pathogens from a recent survey of human bacterial
pathogens[2] (n=125 genera). Genera with a single pathogenic species were not included in order to
balance between including genera responsible for substantial human pathogenic burden and
computational feasibility. We further included genera including human protozoan pathogens (n=11
genera), as well as all viral genera (n=1,356).
522
For each genus of interest showing $\geq$ 50 unique *k*-mers assigned, all sequencing reads classified were
collected and aligned in parallel against a representative reference assembly for each individual
species within the genus. We selected the assembly with the most unique *k*-mers assigned as the
representative reference genome for each species in a particular sample. If no reads were assigned to
any assembly of the species in *KrakenUniq*, we selected the first assembly for mapping. Read
mapping against the selected assembly was carried out using *bowtie2*[76], using the 'very sensitive'
preset and allowing one mismatch in the seed ('-N 1' option). Mapped BAM files were subjected to
duplicate marking using '*samtools markdup*'[77], and filtered for mapping quality MAPQ$\geq$20. aDNA
damage rates were estimated using *metaDMG*[78].

Authentication of ancient microbial DNA

To authenticate ancient microbial DNA, we calculated sets of summary statistics quantifying expected
molecular characteristics of true positive ancient microbial DNA hits[79]:

Similarity to the reference assembly

Summary statistics in this category measure how similar sequencing reads are to a particular reference
assembly, with true positive hits expected to show higher similarity than false positive hits. Summary
statistics used include:

Average edit distance
The average number of mismatches in sequencing reads mapped to a particular reference (lower -
more similar to reference).

Average nuclear identity (ANI)
The average number of bases in a mapped sequencing read matching the reference assembly,
normalised by the read length (higher - more similar to reference).

Number of unique *k*-mers assigned
The number of unique *k*-mers assigned to a particular reference assembly from *KrakenUniq*
classification (higher - more similar to reference).

Ancient DNA characteristics

Summary statistics in this category measure the evidence for sequencing reads deriving from an
aDNA source. Summary statistics used include:

Average read length
The average length in base pairs of sequencing reads mapped to a particular reference (shorter - more
likely ancient).

Terminal aDNA substitution rates
The frequency of C>T (G>A) substitutions observed at the 5' (3') terminal base across all sequencing
reads mapped to a particular reference (higher - more likely ancient).

Bayesian D_{max}
Bayesian estimator of aDNA damage rate from *metaDMG* (higher - more likely ancient).

Bayesian Z_x
Bayesian estimator of significance of evidence for aDNA damage rate from *metaDMG* (higher - more
likely ancient).

Evenness of genomic coverage

Summary statistics in this category measure how evenly mapped sequencing reads are distributed
across a reference assembly. Summary statistics used include:

Average read depth
The average number of reads covering a base in the reference assembly.

Breadth of coverage
The fraction of the reference assembly that is covered by one or more sequencing reads.

Expected breadth of coverage

Breadth of coverage expected for a particular average read depth, calculated[80] as

$1 - e^{-(\text{average read depth})}$

<u>Ratio of observed over expected breadth of coverage</u>
The ratio of breadth of coverage observed in mapping over breadth of coverage expected given
observed average read depth (higher - more even coverage).

<u>Relative entropy of read start positions</u>
A measure for the information content of the genomic positions of mapped reads. To obtain this
statistic, we calculate the frequency of read alignments with their start positions falling within
windows along the reference assembly, using two different window sizes (100bp and 1000bp). The
obtained frequency vector is converted into Shannon information entropy, and normalised using the
maximum entropy attainable if the same total number of reads were evenly distributed across the
windows (higher - more even coverage).

Filtering of putative ancient microbial hits

From this initial screening, we then selected a subset of putative microbial "hits" (sample/species
combinations) for further downstream analysis based on a set of aDNA authentication summary
statistics:

-    Number of mapped reads ≥ 20
-    5' C>T deamination rate ≥ 0.01
-    3' G>A deamination rate ≥ 0.01
-    Ratio of observed/expected breadth of coverage ≥ 0.8
-    Relative entropy of read start positions ≥ 0.9
-    ANI > 0.965
-    Rank of number of unique k-mers assigned ≤ 2

For this initial filtered list of putative microbial hits, we ran *metaDMG* using the full Bayesian
inference method to obtain Z-scores measuring the strength of evidence for observing aDNA damage
(Supplementary Data 2).

The final list of putative individual ancient microbial hits was then obtained using the filtering cutoffs

-    *metaDMG* Bayesian $D_{max} \geq 0.05$
-    *metaDMG* Bayesian $Z \geq 1.5$
-    Rank of number of unique k-mers assigned $== 1$

For authentication of viral species, we used the same filtering cutoffs described above, except for a
lower ANI cutoff (> 0.95), as well as a lower cutoff for relative entropy of read start positions (>0.7)
for short viral genomes (< 10kb).

The result of this filtering is a single best-matching species hit for each sample and genus of interest
Supplementary table S2. We note that this approach will miss potential cases where aDNA from
multiple species of the same genus are present in the sample. However, due to the considerable
challenges involved in distinguishing this scenario from false positives due to cross-mapping of

ancient reads from a single source of DNA to reference assemblies of a closely related species (e.g.,
Yersinia pestis / Yersinia pseudotuberculosis), we opted for the conservative option of retaining only
the best hit for each genus.
629
To further authenticate putative hits with low read counts (N ≤ 100 final reads), we carried out a
BLASTn analysis. We extracted the reads for a species hit from the final filtered BAM files, and
queried them against the 'nt' database (downloaded 20240828) using `blastn -task blastn -max_hsps
1`. For the reads of each putative ancient microbial hit, we then tabulated the number of times and
proportion of the highest scoring BLAST hits matched either the genus or species inferred from our
workflow Supplementary table S3).

Simulations of ancient microbial DNA

We simulated aDNA fragments from microbial reference genomes *in silico* using *gargammel*[81]. We
chose nine species representing pathogens of interest, and for each selected an assembly not present in
the pathogen screening workflow database:
640
-   *Brucella melitensis* (GCF_027625455.1)
-   *Helicobacter pylori* (NZ_CP134396.1)
-   *Mycobacterium tuberculosis* (NZ_CP097110.1)
-   *Salmonella enterica* (NZ_CP103966.1)
-   *Yersinia pestis* (NZ_CP064125.2)
-   *Yersinia pseudotuberculosis* (NZ_CP130901.1)
-   *Plasmodium vivax* (GCA_900093555.2)
-   *Variola virus* (GCA_037113635.1)
-   *Human betaherpesvirus 5* (GCA_027927465.1)
650
For each reference genome, we simulated 5 million single-end sequencing reads (100 bp read length)
with adapter sequences, with read length distribution and damage patterns from a *mapDamage2*
results of a previously published ancient pathogen genome (RISE509, *Yersinia pestis*[16]). The full-
length simulated reads were then adapter-trimmed using *AdapterRemoval*[82]. To investigate the ability
of the workflow to detect low abundance ancient microbes, we randomly down-sampled the full read
set for each reference genome using *seqtk* (https://github.com/lh3/seqtk) (50, 100, 200, 500 reads; 10
replicates each).

Topic model analysis

We carried out topic model analysis on taxonomic classification profiles for each sample using the R
package *fastTopics*[83] (https://github.com/stephenslab/fastTopics). We used the number of unique *k*-
mers assigned to non-human genera from *KrakenUniq* as the observed count data for each sample,
excluding genera with less than 50 unique *k*-mers assigned. The analysis was carried out for L=2 and
L=3 topics, to capture broad structure in the classification profiles.
664
665 Ancient microbial groups

666 For combined analyses, we grouped the ancient microbial hits into three categories, based on the
667 likely source of the microbial DNA (Supplementary table S4):
668

1) Environmental, to capture all hits derived from environmental sources including the necrobiome
(labelled environment_background, environment_pathogen, to distinguish potential pathogenic
species from non-pathogenic ones);
2) Oral microbiome, including both commensal and pathogenic species (microbiome_oral)
3) likely pathogenic infections, further distinguished into different modes of transmission
(infection_anthroponotic; infection_vector_borne; infection_zoonotic).
675
676   We define zoonotic pathogens here as those transmitted from animals to humans or which made such
677   a host jump in our sampling time frame[40].

Time series

To infer temporal dynamics of ancient microbial species, we calculated detection rates in a sliding
window of k=21 temporally consecutive samples across the entire timeline of the 1,266 samples with
dating information. For individual species, the detection rate for each window corresponds to the
proportion of the 21 samples in each window that were positive for the species of interest. For
analyses of species combined in classes, we calculated the detection rate as the ratio of the total
number of hits within a class in the window over the total number of possible hits across all species in
a window (21 samples x 258 species across all classes). For individual species with $n \geq 20$ hits or
combined species classes, we further performed Bayesian change-point detection and time series
decomposition (BEAST) [62] implemented in the R package *Rbeast* (https://github.com/zhaokg/Rbeast),
using the detection frequencies described above as response variables.

Spatiotemporal models of species incidence

To identify possible drivers of the observed spatiotemporal ancient microbial incidence, we combined
the individual microbial species and the combined species groups with palaeoclimatic variables,
human mobility estimates and kriged estimates of ancestry composition for Holocene West Eurasia.
Palaeoclimatic reconstructions were accessed using the CHELSA-Trace21k data, which provides
global monthly climatologies for temperature and precipitation at 30 arcsec spatial resolution in 100-
year time steps for the last 21,000 years[84]. To pair the microbial species/groups to the palaeoclimatic
reconstructions, we took the average climatic value across all the time steps that fall within the
microbial species/group age ± sd at each of the sampling locations. Palaeoclimatic variables
considered were annual mean temperature (BIO01) and annual precipitation (BIO12). Human
mobility values were accessed from Schmid & Schiffels[85] and approximately represent the distance in
kilometres between the burial location of the ancient human individual and its putative ancestral
origin, based on patterns of genetic similarity derived from a MDS analysis. Microbial species/groups
were paired to the mobility estimate of the ancient human individual that occurs closest in space and
time. Kriged ancestry estimates were extracted from Allentoft *et al.*[86], using the spatiotemporal
ancestry kriging method from Racimo et al.[87], and paired to the closest spatiotemporal location of the
ancient human remain where the corresponding microbial species/groups were sampled.
706
To determine the influence of the covariates on the microbial incidence, we used a hierarchical
Bayesian model implemented in the *inlabru* R package[67,88], where ancient microbial presence/absence
follows a binomial distribution and the spatiotemporal variables (latitude, longitude and sample age),
number of human-classified reads, sample material, palaeoclimatic variables, human mobility and
human ancestry constitute the linear predictors. The sample material is a categorical variable
indicating whether the material used for sequencing was a tooth or not (bone), which *inlabru* treats as
a random effect variable. We followed the default *inlabru* priors, where distributions are distributed as

a Gaussian variable with mean μ and precision τ. The prior on the precision τ is a Gamma with parameters 1 and 0.00005. The mean is a linear combination of the covariates. By default, the prior on the intercept of the linear combination is a uniform distribution, while the priors on the coefficients are Gaussian with zero mean and precision 0.001. All covariates were normalised before the analyses. For each microbial species and group, we tested multiple models with different sets of covariates: 1) palaeoclimate + mobility + ancestry, 2) palaeoclimate + mobility, 3) palaeoclimate + ancestry, 4) only climate, 5) mobility + ancestry, 6) only mobility, 7) only ancestry, 8) no climate, nor mobility, nor ancestry. Spatiotemporal variables, number of human-classified reads, and sample material were included in all models. Because covariates were normalised, results indicate deviations from the mean. The effect size is interpreted in units of standard deviation. We used the deviance information criterion (DIC) to assess the model fit to each set of covariates, and prevent overfitting. The results shown in the main text are for the best-performing models (i.e., models with the lowest DIC score for each microbial species or combined species group). DIC scores as well as Watanabe–Akaike information criterion (WAIC) for each model, and results for all the other models we tested can be found in the Supplementary table S6.

Data availability

All sequencing data used in this study is available as trimmed read files (FASTQ) at the European Nucleotide Archive under accession PRJEB65256. Processed analysis files including *KrakenUniq* database file and metagenomic profiling results, microbial species read alignments (BAM format) as well as per-sample summary tables and plots from screening pipeline are available at Zenodo under accession XX.

Code availability

A *Snakemake* workflow implementing the computational screening pipeline is available at https://github.com/martinsikora/pathopipe.

Acknowledgements

The Lundbeck Foundation GeoGenetics Centre is supported by the the Lundbeck Foundation (R302-2018-2155, R155-2013-16338), the Novo Nordisk Foundation (NNF18SA0035006), the Wellcome Trust (UNS69906), Carlsberg Foundation (CF18-0024), the Danish National Research Foundation (DNRF94, DNRF174), the University of Copenhagen (KU2016 programme) and Ferring Pharmaceuticals A/S to E.W. Additional support was provided by Germany's Excellence Strategy (EXC- 2077), project 390741603 "The Ocean Floor – Earth's Uncharted Interface". We thank A. Razeto, P. Selmer Olsen, for administrative and technical assistance. We thankfully acknowledge Illumina Inc. for collaboration. EW thanks St. John's College, Cambridge, for providing a stimulating environment of discussion and learning. This work was further supported by the Swedish Foundation for Humanities and Social Sciences grant (Riksbankens Jubileumsfond M16-0455:1) to KK. M.E.A. was supported by Marie Skłodowska-Curie Actions of the EU (grant no. 300554), The Villum Foundation (grant no. 10120) and Independent Research Fund Denmark (grant no. 7027-00147B). G.S. is funded by Marie Skłodowska-Curie Individual Fellowship 'PALAEO-ENEO' (grant agreement number 751349). H.S. was supported by a Carlsberg Semper Ardens Grant (grant no. CF19-0601) and an ERC Consolidator Grant (grant no. 101045643). F. R. is supported by a Villum Young Investigator Grant (project no. 00025300), a Novo Nordisk Fonden Data Science Ascending Investigator Award (NNF22OC0076816) and by the European Research Council (ERC) under the

European Union's Horizon Europe programme (grant agreements No. 101077592 and 951385). N.O., R.Å., L.H. and B.N. are financially supported by Knut and Alice Wallenberg Foundation as part of the National Bioinformatics Infrastructure Sweden at SciLifeLab. A.K.N.I. and L. F. thank the OAK Foundation.

Author contributions

M.S. and E.W. conceptualized the study. M.S., E. C., A. F. G., S. H. N., A.K.N.I, and F. V. S analysed data. M.S., E.C. A. F. G., N. O., R. A., L. H., E. K. I.-P., B. M., S. H. N. and H.S. were involved in method development and implementation. G. S., M. E. A., F. V. S., H.S., C. G. J. S. and L. V. were involved in data generation. M. S., M. E. A., K. G., and K. K. curated bioarchaeological data. M.S. T. C. J., B. N., J. P., L. F., F. R. and E.W. supervised the research. M.S., A. K. N. I. and E.W. wrote the first draft of the paper. M.S, A. K. N. I., L. F. J. P. and E.W. were involved in reviewing drafts and editing. All co-authors read, commented on and agreed on the submitted manuscript.

References

1. Lewis, C. M., Jr, Akinyi, M. Y., DeWitte, S. N. & Stone, A. C. Ancient pathogens provide a window into health and well-being. *Proc. Natl. Acad. Sci. U. S. A.* **120**, e2209476119 (2023).
2. Bartlett, A., Padfield, D., Lear, L., Bendall, R. & Vos, M. A comprehensive list of bacterial pathogens infecting humans. *Microbiology* **168**, (2022).
3. Barrett, R., Kuzawa, C. W., McDade, T. & Armelagos, G. J. EMERGING AND RE-EMERGING INFECTIOUS DISEASES: The third epidemiologic transition. *Annu. Rev. Anthropol.* **27**, 247–271 (1998).
4. Haak, W. *et al.* Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature* **522**, 207–211 (2015).
5. Allentoft, M. E. *et al.* Population genomics of Bronze Age Eurasia. *Nature* **522**, 167–172 (2015).
6. Infant and child death in the human environment of evolutionary adaptation. *Evol. Hum. Behav.* **34**, 182–192 (2013).
7. Harper, K. *Plagues Upon the Earth: Disease and the Course of Human History*. (Princeton University Press, 2021).
8. Karlsson, E. K., Kwiatkowski, D. P. & Sabeti, P. C. Natural selection and infectious disease in human populations. *Nat. Rev. Genet.* **15**, 379–393 (2014).
9. Quintana-Murci, L. Human Immunology through the Lens of Evolutionary Genetics. *Cell* **177**, 184–199 (2019).
10. Kerner, G. *et al.* Genetic adaptation to pathogens and increased risk of inflammatory disorders in post-Neolithic Europe. *Cell Genom.* **3**, 100248 (2023).
11. Page, A. E. *et al.* Reproductive trade-offs in extant hunter-gatherers suggest adaptive mechanism for the Neolithic expansion. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 4694–4699 (2016).
12. Rascovan, N. *et al.* Emergence and Spread of Basal Lineages of Yersinia pestis during the Neolithic Decline. *Cell* **176**, 295–305.e10 (2019).
13. Fuchs, K. *et al.* Infectious diseases and Neolithic transformations: Evaluating biological and archaeological proxies in the German loess zone between 5500 and 2500 BCE. *Holocene* **29**, 1545–1557 (2019).
14. Abegg, C., Desideri, J., Dutour, O. & Besse, M. More than the sum of their parts: reconstituting

the paleopathological profile of the individual and commingled Neolithic populations of Western
Switzerland. *Archaeol. Anthropol. Sci.* **13**, (2021).

15. Bos, K. I. *et al.* A draft genome of Yersinia pestis from victims of the Black Death. *Nature* **478**,
506–510 (2011).

16. Rasmussen, S. *et al.* Early Divergent Strains of Yersinia pestis in Eurasia 5,000 Years Ago. *Cell*
163, 571–582 (2015).

17. Andrades Valtueña, A. *et al.* The Stone Age Plague and Its Persistence in Eurasia. *Curr. Biol.* **27**,
3683–3691.e8 (2017).

18. Spyrou, M. A. *et al.* Analysis of 3800-year-old Yersinia pestis genomes suggests Bronze Age
origin for bubonic plague. *Nat. Commun.* **9**, 2234 (2018).

19. Keller, M. *et al.* Ancient genomes from across Western Europe reveal early diversification
during the First Pandemic (541-750). *Proc. Natl. Acad. Sci. U. S. A.* **116**, 12363–12372 (2019).

20. Susat, J. *et al.* A 5,000-year-old hunter-gatherer already plagued by Yersinia pestis. *Cell Rep.* **35**,
109278 (2021).

21. Feldman, M. *et al.* A High-Coverage Yersinia pestis Genome from a Sixth-Century Justinianic
Plague Victim. *Mol. Biol. Evol.* **33**, 2911–2923 (2016).

22. Andrades Valtueña, A. *et al.* Stone Age genomes shed light on the early evolution, diversity, and
ecology of plague. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2116722119 (2022).

23. Guellil, M. *et al.* A genomic and historical synthesis of plague in 18th century Eurasia. *Proc.*
Natl. Acad. Sci. U. S. A. **117**, 28328–28335 (2020).

24. Bos, K. I. *et al.* Pre-Columbian mycobacterial genomes reveal seals as a source of New World
human tuberculosis. *Nature* **514**, 494–497 (2014).

25. Vågene, Å. J. *et al.* Geographically dispersed zoonotic tuberculosis in pre-contact South
American human populations. *Nat. Commun.* **13**, 1195 (2022).

26. Duggan, A. T. *et al.* 17th Century Variola Virus Reveals the Recent History of Smallpox. *Curr.*
Biol. **26**, 3407–3412 (2016).

27. Mühlemann, B. *et al.* Diverse variola virus (smallpox) strains were widespread in northern
Europe in the Viking Age. *Science* **369**, (2020).

28. Mühlemann, B. *et al.* Ancient Hepatitis B viruses from the Bronze Age to the Medieval.

29. Kocher, A. *et al.* Ten millennia of hepatitis B virus evolution. *Science* **374**, 182–188 (2021).

30. Krause-Kyora, B. *et al.* Neolithic and medieval virus genomes reveal complex evolution of
hepatitis B. *Elife* **7**, (2018).

31. Guellil, M. *et al.* Genomic blueprint of a relapsing fever pathogen in 15th century Scandinavia.
Proc. Natl. Acad. Sci. U. S. A. **115**, 10422–10427 (2018).

32. Schuenemann, V. J. *et al.* Genome-Wide Comparison of Medieval and Modern Mycobacterium
leprae. *Science* **341**, 179–183 (2013).

33. Guellil, M. *et al.* Ancient herpes simplex 1 genomes reveal recent viral structure in Eurasia. *Sci*
Adv **8**, eabo4435 (2022).

34. van Dorp, L. *et al.* Plasmodium vivax Malaria Viewed through the Lens of an Eradicated
European Strain. *Mol. Biol. Evol.* **37**, 773–785 (2020).

35. Key, F. M. *et al.* Emergence of human-adapted Salmonella enterica is linked to the
Neolithization process. *Nat Ecol Evol* **4**, 324–333 (2020).

36. Mühlemann, B. *et al.* Ancient human parvovirus B19 in Eurasia reveals its long-term association
with humans. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 7557–7562 (2018).

37. Bonczarowska, J. H. *et al.* Pathogen genomics study of an early medieval community in
Germany reveals extensive co-infections. *Genome Biol.* **23**, 250 (2022).

38. Abdoun, A., Amir, N. & Fatima, M. Thanatomicrobiome in forensic medicine. *New Microbiol.*
46, 236–245 (2023).

39.   Burcham, Z. M. *et al.* A conserved interdomain microbial network underpins cadaver
850   decomposition despite environmental variables. *Nat Microbiol* **9**, 595–613 (2024).
851   40.   WHO. Zoonoses fact sheet. https://www.who.int/news-room/fact-sheets/detail/zoonoses.
852   41.   Spyrou, M. A. *et al.* Phylogeography of the second plague pandemic revealed through analysis of
853   historical Yersinia pestis genomes. *Nat. Commun.* **10**, 4470 (2019).
854   42.   Neumann, G. U. *et al.* Ancient Yersinia pestis and Salmonella enterica genomes from Bronze
855   Age Crete. *Curr. Biol.* **32**, 3641–3649.e8 (2022).
856   43.   Macleod, R. *et al.* Lethal plague outbreaks in Lake Baikal Hunter-gatherers 5500 years ago.
857   *bioRxiv* (2024) doi:10.1101/2024.11.13.623490.
858   44.   ECDC. Louse-borne relapsing fever in the EU.
859   *https://www.ecdc.europa.eu/sites/default/files/media/en/publications/Publications/louse-borne-*
860   *relapsing-fever-in-eu-rapid-risk-assessment-17-nov-15.pdf*
861   https://www.ecdc.europa.eu/sites/default/files/media/en/publications/Publications/louse-borne-
862   relapsing-fever-in-eu-rapid-risk-assessment-17-nov-15.pdf (2015).
863   45.   Swali, P. *et al.* Ancient Borrelia genomes document the evolutionary history of louse-borne
864   relapsing fever. *bioRxiv* 2024.07.18.603748 (2024) doi:10.1101/2024.07.18.603748.
865   46.   Warrell, D. A. Louse-borne relapsing fever (Borrelia recurrentis infection). *Epidemiol. Infect.*
866   **147**, e106 (2019).
867   47.   Avanzi, C. *et al.* Red squirrels in the British Isles are infected with leprosy bacilli. *Science* **354**,
868   744–747 (2016).
869   48.   Hennius, A. Outlanders? : Resource colonisation, raw material exploitation and networks in
870   Middle Iron Age Sweden. (Uppsala University, Archaeology, 2021).
871   49.   Urban, C. *et al.* Ancient Mycobacterium leprae genome reveals medieval English red squirrels as
872   animal leprosy host. *Curr. Biol.* **34**, 2221–2230.e8 (2024).
873   50.   Pfrengle, S. *et al.* Mycobacterium leprae diversity and population dynamics in medieval Europe
874   from novel ancient genomes. *BMC Biol.* **19**, 220 (2021).
875   51.   Bulach, D. M. *et al.* Genome reduction in Leptospira borgpetersenii reflects limited transmission
876   potential. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 14560–14565 (2006).
877   52.   Rees, C. E., Swift, B. M. & Haldar, P. State-of-the-art detection of Mycobacterium tuberculosis
878   in blood during tuberculosis infection using phage technology. *Int. J. Infect. Dis.* **141S**, 106991
879   (2024).
880   53.   Steinegger, M. & Salzberg, S. L. Terminating contamination: large-scale search identifies more
881   than 2,000,000 contaminated entries in GenBank. *Genome Biol.* **21**, 115 (2020).
882   54.   Hay, S. I., Guerra, C. A., Tatem, A. J., Noor, A. M. & Snow, R. W. The global distribution and
883   population at risk of malaria: past, present, and future. *Lancet Infect. Dis.* **4**, 327–336 (2004).
884   55.   Hulden, L. & Hulden, L. The decline of malaria in Finland--the impact of the vector and social
885   variables. *Malar. J.* **8**, 94 (2009).
886   56.   Davidyants, V. A. *et al.* Role of malaria partners in malaria elimination in Armenia. *Malar. J.* **18**,
887   178 (2019).
888   57.   Roberto, P. *et al.* Torque teno virus (TTV): A gentle spy virus of immune status, predictive
889   marker of seroconversion to COVID-19 vaccine in kidney and lung transplant recipients. *J. Med.*
890   *Virol.* **95**, e28512 (2023).
891   58.   İnce, İ. A., Özcan, O., Ilter-Akulke, A. Z., Scully, E. D. & Özgen, A. Invertebrate Iridoviruses: A
892   Glance over the Last Decade. *Viruses* **10**, (2018).
893   59.   Singer, M., Bulled, N., Ostrach, B. & Mendenhall, E. Syndemics and the biosocial conception of
894   health. *Lancet* **389**, 941–950 (2017).
895   60.   Zasada, A. A., Zaleska, M., Podlasin, R. B. & Seferynska, I. The first case of septicemia due to
896   nontoxigenic Corynebacterium diphtheriae in Poland: case report. *Ann. Clin. Microbiol.*

Antimicrob. **4**, 8 (2005).

61. Han, X. Y., Tarrand, J. J., Dickey, B. F. & Esteva, F. J. Helicobacter pylori bacteremia with
sepsis syndrome. *J. Clin. Microbiol.* **48**, 4661–4663 (2010).

62. Zhao, K. *et al.* Detecting change-point, trend, and seasonality in satellite time series data to track
abrupt changes and nonlinear dynamics: A Bayesian ensemble algorithm. *Remote Sens. Environ.*
232, 111181 (2019).

63. Fellows Yates, J. A. *et al.* Community-curated and standardised metadata of published ancient
metagenomic samples with AncientMetagenomeDir. *Sci Data* **8**, 31 (2021).

64. Demeure, C. *et al.* Yersinia pestis and plague: an updated view on evolution, virulence
determinants, immune subversion, vaccination and diagnostics. *Microbes Infect.* **21**, 202–212
(2019).

65. Sun, Y.-C., Jarrett, C. O., Bosio, C. F. & Hinnebusch, B. J. Retracing the evolutionary path that
led to flea-borne transmission of Yersinia pestis. *Cell Host Microbe* **15**, 578–586 (2014).

66. Bland, D. M., Long, D., Rosenke, R. & Hinnebusch, B. J. Yersinia pestis can infect the
Pawlowsky glands of human body lice and be transmitted by louse bite. *PLoS Biol.* **22**, e3002625
(2024).

67. doi:10.1111/2041-210X.13168.

68. Margaryan, A. *et al.* Ancient pathogen DNA in human teeth and petrous bones. *Ecol. Evol.* **8**,
3534–3542 (2018).

69. Ellwanger, J. H. & Chies, J. A. B. Zoonotic spillover: Understanding basic aspects for better
prevention. *Genet. Mol. Biol.* **44**, e20200355 (2021).

70. Jones, K. E. *et al.* Global trends in emerging infectious diseases. *Nature* **451**, 990–993 (2008).

71. Meltzer, D. J. *First Peoples in a New World: Colonizing Ice Age America.* (Univ of California
Press, 2009).

72. Collen, E. J., Johar, A. S., Teixeira, J. C. & Llamas, B. The immunogenetic impact of European
colonization in the Americas. *Front. Genet.* **13**, 918227 (2022).

73. Barrie, W. *et al.* Genetic risk for Multiple Sclerosis originated in Pastoralist Steppe populations.
(2022) doi:10.1101/2022.09.23.509097.

74. Cutler, S. J. Relapsing fever--a forgotten disease revealed. *J. Appl. Microbiol.* **108**, 1115–1122
(2010).

75. Breitwieser, F. P., Baker, D. N. & Salzberg, S. L. KrakenUniq: confident and fast metagenomics
classification using unique k-mer counts. *Genome Biol.* **19**, 198 (2018).

76. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
357–359 (2012).

77. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

78. Michelsen, C. *et al.* MetaDMG – A fast and accurate ancient DNA damage toolkit for
metagenomic data. *bioRxiv* (2022) doi:10.1101/2022.12.06.519264.

79. Warinner, C. *et al.* A Robust Framework for Microbial Archaeology. *Annu. Rev. Genomics Hum.*
Genet. **18**, 321–356 (2017).

80. Lander, E. S. & Waterman, M. S. Genomic mapping by fingerprinting random clones: a
mathematical analysis. *Genomics* **2**, 231–239 (1988).

81. Renaud, G., Hanghøj, K., Willerslev, E. & Orlando, L. gargammel: a sequence simulator for
ancient DNA. *Bioinformatics* **33**, 577–579 (2017).

82. Schubert, M., Lindgreen, S. & Orlando, L. AdapterRemoval v2: rapid adapter trimming,
identification, and read merging. *BMC Res. Notes* **9**, (2016).

83. Carbonetto, P., Sarkar, A., Wang, Z. & Stephens, M. Non-negative matrix factorization
algorithms greatly improve topic model fits. (2021) doi:10.48550/ARXIV.2105.13440.

84. Karger, D. N., Nobis, M. P., Normand, S., Graham, C. H. & Zimmermann, N. E. CHELSA-

TraCE21k -- high-resolution (1 km) downscaled transient temperature and precipitation data
since the Last Glacial Maximum. *Clim. Past* **19**, 439–456 (2023).

85. Schmid, C. & Schiffels, S. Estimating human mobility in Holocene Western Eurasia with large-
scale ancient genomic data. *Proc. Natl. Acad. Sci. U. S. A.* **120**, e2218375120 (2023).

86. Allentoft, M. E. *et al.* Population genomics of Stone Age Eurasia. *bioRxiv* (2022)
doi:10.1101/2022.05.04.490594.

87. Racimo, F. *et al.* The spatiotemporal spread of human migrations during the European Holocene.
Proc. Natl. Acad. Sci. U. S. A. **117**, 8989–9000 (2020).

88. Lindgren, F. & Rue, H. Bayesian Spatial Modelling withR-INLA. *J. Stat. Softw.* **63**, (2015).

Extended Data Figures

955
956 **Extended Data Fig. 1. Workflow overview and metagenome composition. b,** Distribution of total number of sequencing
957 reads screened across the 1,313 study samples. **c,** Violin plots showing distributions of proportions of reads classified as
958 human, non-human or not classified for the study samples. Median values for each genus are indicated by horizontal lines. **d,**
959 Violin plots showing fraction of reads classified on the taxonomic level of genus, for the top 20 most abundant genera. **e,**

Barplots showing inferred proportions for L=3 topics (indicated by fill colour) from topic model analysis for 1,272 study samples with sample material information. **f,** Factor loadings for the 10 highest loading genera for each of the L=3 topics from the topic model analysis. **g,** Boxplots showing distributions of proportions for topic K3 (associated with oral microbiome taxa) in different sample materials.

Extended Data Fig. 2. Reference genome similarity in simulated ancient microbial data. a, Illustration showing phylogenetic context and expected average nucleotide identity (ANI) for a hypothetical sampled microbial species X and four genomes (A1, A2; B1, B2) of two genera (A, B) present in the reference database. **b,** Number of unique *k*-mers classified at the level of genus using *KrakenUniq* for replicates of different read numbers across all simulated species.

969 Dashed line indicates cutoff used in analysis of real data (150 unique k-mers). **c**, Number of unique k-mers classified at the
970 level of species as a function of average nucleotide identity for mappings against all individual species reference genomes in
971 the genus of reads simulated for a particular species. Blue diamonds indicate results for the mapping against a reference
972 genome from the same species as the simulated read data, whereas grey circles indicate reference genomes of other species.
973 Selected individual species results are highlighted by species name. Dashed line indicates ANI ≥ 0.97 cutoff value. **d**,
974 Barplots showing number of replicates where the true positive species reference genome was highest ranking in numbers of
975 unique k-mers classified at level of species.

Extended Data Fig. 3. Read mappings across genera in simulated ancient microbial data. a, Observed breadth of genomic coverage as a function of average read depth for distinct species hits (i.e., mappings with highest number of unique k-mers at species level for a genus; n ≥ 20 reads mapped). Each panel shows results for reads simulated from species indicated. Results for mappings against the simulated species are indicated by diamond shape, whereas mappings against species from other genera are indicated with circles. Symbol fill colour indicates average nucleotide identity for mapped reads (grey symbols ANI < 0.97). Solid black line shows theoretical expected breadth of coverage for a given average read depth[80]. Vertical dashed line indicates 1X average read depth. **b,** Relative entropy statistic (1000 bp window size) as a function of average nucleotide identity. Blue diamonds indicate results for the mapping against reference genome from the same species as the simulated read data, whereas grey circles indicate reference genomes for species hits in other genera.

Dashed lines indicate cutoffs used in analyses of real data (ANI ≥ 0.97, entropy ≥ 0.9). False positive hits of reads mapped to
a reference genome from a different genome passing cutoffs and their final number of mapped reads (out of 5 million total
simulated reads) are labelled. **c**, Illustration showing potential sources of false positive hits and expected results for
authentication summary statistics. **d**, Matrix plot showing all microbial hits with n ≥ 20 reads mapped and their
authentication statistics, for all simulated species and read numbers. Symbol colour and size indicates the number of
replicates passing the cutoff for each of three summary statistics shown (ANI ≥ 0.97, ratio of observed / expected coverage
breadth ≥ 0.8, entropy ≥ 0.9). Hits passing cutoffs for all three statistics are indicated with coloured outline and background
lines (black - true positives; grey - cross-genus false positive mappings).

995
Extended Data Fig. 4. Examples of authentication for microbial hits. a, Observed breadth of genomic coverage as a
function of average read depth. Coloured symbols indicate hits in species *Toxoplasma gondii* (left panel) and *Yersinia pestis*
(right panel), with symbol colour indicating relative entropy of read start positions. Solid black line shows theoretical
expected breadth of coverage for a given average read depth[80]. **b**, Lengths of contigs in the reference genome of *Toxoplasma*
gondii and number of samples showing n ≥ 20 reads mapped. Symbol colour indicates the average number of reads mapped

to a specific contig across samples. **c,** Bayesian estimator of aDNA damage (D max) and significance (Z-score) obtained
from *metaDMG,* for hits in species *Clostridium botulinum* (left) and *Yersinia pestis* (right). Error bars indicate ± 1 standard
deviation, and symbol fill colour indicates average read depth for mapped reads. Samples used as examples in aDNA
damage curves (d) are labelled and indicated with black circles. **d**, aDNA damage patterns for four example hits in species
Clostridium botulinum and *Yersinia pestis*. Plots show observed nucleotide misincorporation frequencies (red symbols and
line) and *metaDMG* fit (black line) and 68% credible intervals (shaded region) for C>T transitions as a function of distance
from the 5' read end.

Extended Data Fig. 5. Ancient microbial hit ANI and read recruitment. a, b, Distributions of ANI (**a**) and log10-fold change of mapped reads over median of reads classified at taxonomic rank of genus per sample (**b**) for individual species hits detected in n ≥ 5 samples. Symbol colour indicates species hit category.

Extended Data Fig. 6. Ancient microbial hit characteristics. a, Odds ratios for association of ancient hits with sample
material (tooth or bone) across 61 species with ≥ 20 ancient hits. Symbols indicate significance of association (p ≤ 0.01,
Fisher's exact test; white triangles - more frequently identified in tooth; grey circles - no significant association). Error bars
indicate 95% confidence interval of odds ratio **b, c,** Rates of observing multiple alleles in 2 randomly sampled sequencing
reads at genomic sites in 190 ancient hits (average read depth ≥ 1X) across 120 samples. **b,** Multi-allele rate as a function of
ANI. Symbol colour indicates average read depth. **c,** Distribution of multi-allele rate across species hits. Symbol colour
indicates ANI. **d, e** Barplots showing number of hits identified in each microbial species group (**d**) or each species within

groups of likely infections (**e**). Novel and previously reported ancient pathogen hits are distinguished by bar colour, with

total number in each category labelled.

Extended Data Fig. 7. Additional ancient microbial hit authentication. a, Bar plots showing proportion of reads assigned
to same species (dark blue) or genus (light blue) using BLASTn for all hits with N ≤ 100 final reads (N=712), stratified by
genus and microbial source groups **b,** Bar plots showing the proportion of ancient microbial hits with N ≤ 100 final reads
matching the species with most reads assigned using BLASTn, stratified by microbial source group. **c,** Heatmap showing
number of reads mapped to *Yersinia pestis* CO92 chromosome and plasmids, for N=42 *Yersinia pestis* hits. Cell color
indicates ratio of observed over expected breadth of coverage. Results for plasmid pMT1 are shown for full plasmid, as well
as separately for the 19 kb region contaiinng the *ymt* gene absent in the LNBA- strains. Samples are ordered by decreasing
age from top to bottom.

1034
1035 **Extended Data Fig. 8. Time series of detection rates for ancient microbial groups. a-h,** Panels show estimated trendlines
1036 and 95% credible interval for detection rates (top), probability distributions and locations (dotted lines) for change points
1037 (middle) and probability of trend slope (bottom) being positive (red), negative (blue) or zero (white), inferred using Bayesian
1038 change-point detection and time series decomposition.
1039

1040
1041
Extended Data Fig. 9. Spatiotemporal distribution and host genetic structure for ancient microbial groups. a, Panels
showing geographic distributions (top) and timelines (bottom) for identified cases of ancient microbial hits in the oral
microbiome and infection groups classes (indicated by coloured circle). Geographic locations and age distributions of all
1,313 study samples are shown in each panel using white squares. **b,** Principal component analyses showing ancient and
modern human genetic population structure in non-African (left panels) and west Eurasian (right panels) individuals. Grey
crosses indicate present-day individuals, whereas coloured symbols indicate ancient individuals (coloured by sample age).
Diamonds with black outlines indicate position in PCA space for samples with hits in combined infection groups. Major
clines of known ancient and modern human ancestry groups are indicated with labels.

1050
Extended Data Fig. 10. Predictors of ancient microbial species incidence. a, Watanabe–Akaike information criterion
values for each model and response variable. **b,** Matrix showing effect sizes and of 12 potential predictors (columns) for
presence of selected combined ancient microbial species and combined groups inferred from spatiotemporal modelling. For
each class, the model with lowest Watanabe–Akaike information criterion is shown. Symbols indicate the predictors
included in the respective model. Predictors with positive effect (2.5% and 97.5% posterior quantiles both positive) are
shown as red triangles, whereas predictors with negative effect (2.5% and 97.5% posterior quantiles both negative) are
shown as blue inverted triangles. Predictors included in the best-fitting model but without effect (posterior quantile range
spanning zero) are indicated using white circles. Posterior standard error of effect sizes is indicated by error bars.