Synergistic, Multi-level Understanding of Psychedelics: Three Systematic Reviews and Meta-analyses of Their Pharmacology, Neuroimaging and Phenomenology

Kenneth Shinozuka[1,2,3*], Katarina Jerotic[1,2], Pedro Mediano[1,4], Alex T. Zhao[5], Katrin H. Preller[6,7], Robin Carhart-Harris[8,9,10] & Morten L. Kringelbach[1,2,11]

1. Centre for Eudaimonia and Human Flourishing, Linacre College, University of Oxford, Oxford, OX3 9BX, UK
2. Department of Psychiatry, University of Oxford, Oxford, OX3 7JX, UK
3. Oxford Mathematics of Consciousness and Applications Network (OMCAN), University of Oxford, Oxford, OX2 6GG, UK
4. Department of Computing, Imperial College London, London, SW7 2RH, UK
5. Department of Statistics and Data Science (Alumnus), The Wharton School, University of Pennsylvania, Philadelphia, PA 19104, USA
6. Departments of Psychiatry, Neuroscience, and Psychology, Yale University, New Haven, CT 06511, USA
7. Department of Psychiatry, Psychotherapy and Psychosomatics, Psychiatric University Hospital Zurich, University of Zurich, Zurich, 8032, Switzerland
8. Centre for Psychedelic Research, Imperial College London, London, SW7 2AZ, UK
9. Department of Neurology, University of California, San Francisco, San Francisco, CA 94158, USA
10. Department of Neurology, Psychiatry and Behavioral Sciences, University of California, San Francisco, CA 94143, USA
11. Center for Music in the Brain, Department of Clinical Medicine, Aarhus University, Aarhus, 8000, Denmark

*Corresponding email: kenneth.shinozuka@psych.ox.ac.uk
*Corresponding address: Centre for Eudaimonia and Human Flourishing, 7 Stoke Place, Headington, Oxford, UK OX3 9BX

Abstract:

Serotonergic psychedelics induce altered states of consciousness and have shown potential for treating a variety of neuropsychiatric disorders, including depression and addiction. Yet their modes of action are not fully understood. Here, we provide a novel, synergistic understanding of psychedelics arising from systematic reviews and meta-analyses of three hierarchical levels of analysis: 1) subjective experience (phenomenology), 2) neuroimaging and 3) molecular pharmacology. Phenomenologically, medium and high doses of LSD yield significantly higher ratings of visionary restructuralisation than psilocybin on the 5-dimensional Altered States of Consciousness Scale. Our neuroimaging results reveal that, in general, psychedelics significantly strengthen between-network functional connectivity (FC) while significantly diminishing within-network FC. Pharmacologically, LSD induces significantly more inositol phosphate formation at the 5-HT_{2A} receptor than DMT and psilocin, yet there are no significant between-drug differences in the selectivity of psychedelics for the 5-HT_{2A}, 5-HT_{2C}, or D_2 receptors, relative to the 5-HT_{1A} receptor. Our meta-analyses link DMT, LSD, and psilocybin to specific neural fingerprints at each level of analysis. The results show a highly non-linear relationship between these fingerprints. Overall, the results point to the need for standardising experimental procedures and analysis techniques, as well as for more research on the emergence between different levels of psychedelic effects.

Keywords:

psychedelics, DMT, ayahuasca, LSD, psilocybin, meta-analysis, fMRI, functional connectivity, pharmacology, receptor agonism, GPCR, phenomenology, subjective experience, consciousness

Table of Contents

Section 1. Introduction

Psychedelics have been utilised by early cultures for millennia within diverse sociocultural contexts, as well as spiritual and healing rituals (Garcia & Maia, 2022; Nichols, 2016; Tupper et al., 2015). Western research into psychedelics, having blossomed in the '50s and '60s following the first study of lysergic acid diethylamide (LSD) in 1947 (Stoll, 1947), came to a halt in the late 1960s due to changes in government regulations and the classification of these substances as "drugs of abuse" with "no recognised medical value" (Tupper et al., 2015). Revitalisation of psychedelic research commenced in the 1990s (Hermle et al., 1992; Strassman et al., 1994; Strassman & Qualls, 1994), with three key factors inspiring the current interest: scientific advances, evidence of therapeutic potential, and increased public interest (Aday et al., 2021). Researchers have come to recognise that psychedelics may be effective tools in the treatment of psychiatric disorders such as depression (Muttoni et al., 2019; Nutt et al., 2020) and addiction (Tófoli & de Araujo, 2016).

The effects of psychedelic substances have been examined at several levels. Pharmacological research has measured their interaction with various receptors in the brain (e.g. Glennon et al., 1984; Watts et al., 1995), while other research domains have explored the subjective experience, or phenomenology, of a psychedelic 'trip'. Neuroimaging research has examined changes in brain activity and functional connectivity under the influence of psychedelics, yet there are still relatively few studies and a plethora of different analysis techniques. Here, we report the findings of three systematic meta-analyses focused on synthesising the evidence about three classical psychedelics: DMT, LSD, and psilocybin. The main aim is to analyse the literature across three levels of description: (1) phenomenology—to discuss key differences in the states of consciousness elicited by psychedelics; (2) functional neuroimaging—to compare the changes in brain activity, functional connectivity, and entropy that are induced by psychedelics; and (3) pharmacology—to discuss the binding affinities and functional activity of psychedelics with respect to various serotonergic and dopaminergic receptors.

Section 1.1. Phenomenology

Psychedelics produce altered states of consciousness that are characterised by ego death, or a breakdown in one's sense of self, and "mystical" experiences (Griffiths et al., 2018; Yaden et al., 2021). Mystical experiences on psychedelics can be defined by four components: a sense of unity and sacredness; transcendence of time and space; positive mood; and ineffability (Barrett et al., 2015; Maclean et al., 2012). Individuals who have taken one of the classic psychedelics are significantly more likely to have had a mystical experience with all four components than people who had other kinds of meaningful experiences (Griffiths et al., 2019).

However, perhaps the strongest predictor of therapeutic outcomes from psychedelic research is so-called 'emotional breakthrough' experiences, as assessed by the emotional breakthrough inventory or EBI (Roseman et al., 2019). Additionally, psychological insight, which includes awareness of previously repressed emotions, correlates with the therapeutic effects of psychedelics (Davis et al., 2021; Peill et al., 2022).

The Altered States of Consciousness (ASC) scale is one popular questionnaire that captures several of the aforementioned subjective effects. The 5D-ASC measures altered states of consciousness along five different dimensions: oceanic boundlessness (e.g. a sense of interconnectedness), anxious ego dissolution, visionary restructuralisation (e.g. visual hallucinations), auditory alterations, and reduction of vigilance. Subsequent factor analyses led to the extraction of 11 lower-order dimensions from the 5D-ASC, resulting in the development of the 11D-ASC: experience of unity, spiritual experience, blissful state, insightfulness, disembodiment, impaired control and cognition, anxiety, elementary imagery, complex imagery, audio-visual

synesthesia, and changed meaning of percepts (Studerus et al., 2010). Because the ASC is the most common subjective questionnaire in the literature on the neuroimaging of psychedelics, we performed a meta-analysis on both the 5D- and 11D-ASC scores of DMT, LSD, and psilocybin.

However, the ASC is just one of many scales for rating the subjective effects of psychedelics. The multiplicity of scales makes it difficult to comprehensively evaluate the literature on the phenomenology of psychedelics. To date, only two studies have used a single set of scales to compare the subjective effects of LSD and psilocybin in the same group of participants (Holze et al., 2022; Ley et al., 2023). More recent research has attempted to draw direct comparisons by qualitatively examining larger datasets, such as the Erowid database of trip reports (Ballentine et al., 2022; Coyle et al., 2012; Qiu & Minda, 2021; Sanz et al., 2018; Zamberlan et al., 2018). The key findings from these comparative studies are discussed in *Section S3*.

Section 1.2. Neuroimaging

After a long pause in psychedelic research on humans, the pioneering positron emission tomography (PET) scans by Vollenweider and Gouzoulis-Mayfrank in the late 1990s ushered in a new era of research on the human brain under psychedelics (Gouzoulis-Mayfrank et al., 1999; Vollenweider et al., 1997, 1999). Forty to fifty years after the initial spate of electroencephalography (EEG) studies on psychedelics (see Tagliazucchi (2020) for a review), several studies used EEG to measure human brain activity under psychedelics, especially ayahuasca (Hoffmann, 2001; Riba, Anderer, et al., 2002; Riba et al., 2004; Riba, Rodríguez-Fornells, et al., 2002; Stuckey et al., 2005) and psilocybin (Umbricht et al., 2002, 2003). However, many of these studies were predicated on the notion that psychedelics are principally useful as a model for psychosis and schizophrenia (Vollenweider et al., 1997). However, the last decade has seen a huge rise in more exploratory, data-driven or hypothesis-free neuroimaging studies on psychedelics, as well as studies on their therapeutic benefits for clinical populations, e.g. patients with depression. The first functional neuroimaging study of psychedelics, published in 2012 (Carhart-Harris, Erritzoe, et al., 2012), was an exploratory analysis of psilocybin-induced changes in cerebral blood flow and BOLD activity in healthy human participants. Since then, dozens of fMRI studies have been performed on human participants under the influence of ayahuasca, LSD, psilocybin and DMT, including several studies on depressed patients (Carhart-Harris et al., 2017; Daws et al., 2022; Doss et al., 2021; Mertens et al., 2020; Roseman et al., 2018; Wall et al., 2022). Additionally, in the last decade, two studies collected primary magnetoencephalography (MEG) data on healthy human participants who had been administered LSD and psilocybin (Carhart-Harris et al., 2016; Muthukumaraswamy et al., 2013).

We chose to perform a meta-analysis exclusively on the fMRI data and not on any of the other modalities used to measure brain activity. There were too few primary PET (4), SPECT (4), or MEG (2) datasets to merit a meta-analysis; by comparison, 30 fMRI studies have collected original data. While there are many more primary EEG datasets (25) from the post-1960s era (and even more from the 1950s-1960s era) than PET, SPECT, or MEG, most of the EEG studies do not spatially localise the brain activity recorded at EEG electrodes. Without the raw data, which may be difficult to access when some of the studies are nearly 20 years old, we cannot source-localise the EEG activity ourselves. On the other hand, the published fMRI data on psychedelics tends to report the specific coordinates of brain regions that become more or less active under psychedelics. In general, EEG has higher temporal resolution than fMRI (Haufe et al., 2018), so a future meta-analysis that focuses on the temporal, rather than spatial, characteristics of brain activity on psychedelics could consider the EEG rather than fMRI data.

The fMRI studies assess three different metrics of brain activity: BOLD activation, connectivity, and
entropy. Studies on BOLD activation analyse changes in the trajectory of BOLD timeseries with a rapidly-
acting psychedelic, such as intravenously-administered psilocybin. Connectivity studies can examine
undirected, instantaneous correlations between different regions (a type of functional connectivity) or model
the experimental factors that modulate directed connections between regions, using dynamic causal modeling
(a tool for measuring effective connectivity). The entropy of spontaneous brain activity is best assessed with
MEG and EEG, but efforts to compute entropy on the spontaneous BOLD signal, as well as the entropy
rate (Lempel-Ziv complexity) and indices of criticality, have also been attempted (see *Section S1.2* for a full
list of citations). While studies on BOLD activation and functional connectivity are common for most subject
areas within cognitive neuroscience, entropy is a rather unique feature of the psychedelic neuroimaging
literature. The studies on entropy were motivated by Robin Carhart-Harris' influential Entropic Brain
Hypothesis (EBH) (Carhart-Harris et al., 2014; Carhart-Harris, 2018), which states that psychedelics alter
consciousness by elevating the entropy of spontaneous brain activity across time.

We performed a quantitative meta-analysis of the pairwise functional connectivity data using a novel
algorithm. We initially attempted to conduct a quantitative meta-analysis of the BOLD activation data with
the GingerALE method, which examines common clusters of BOLD activity across studies. However, we
found that the data was too heterogeneous for the results to be valid. From the outset, we decided to provide
a qualitative discussion of the studies on entropy, due to the wide range of methods in that section of the
literature.

Section 1.3. Pharmacology

Chemically, psychedelics fall into three categories: tryptamines (e.g. DMT, psilocybin, ayahuasca), ergolines
(e.g. LSD), and phenethylamines (e.g. mescaline, 2C-B) (Nichols, 2018). The molecular structure of
tryptamine psychedelics strongly resembles that of serotonin; in fact, serotonin is itself a tryptamine, as both
are synthesised from the same amino acid, tryptophan (Brandt et al., 2004). Ergolines are rigid and more
complex analogues of tryptamines (Nichols, 2004), whereas the chemical backbone of phenethylamines is
not the same as that of tryptamines; it is instead constituted from the amino acid phenylalanine (Fantegrossi
et al., 2008).

Despite their chemical differences, both tryptamines and phenethylamines – indeed, all classic
psychedelics – activate serotonin receptors. Phenethylamines are very selective for the serotonin-2 (5-HT$_2$)
receptor subtype, including 5-HT$_{2A}$, 5-HT$_{2B}$, and 5-HT$_{2C}$, whereas tryptamines exhibit a more "promiscuous"
binding profile, as they also have affinity for the 5-HT$_1$ receptor subtype (Araújo et al., 2015). In spite of the
broad range of serotonin receptors that psychedelics, especially tryptamines, bind to, their interaction with
the 5-HT$_{2A}$ receptor is their primary mechanism of action (González-Maeso et al., 2007; López-Giménez &
González-Maeso, 2018). Multiple studies have demonstrated that blocking the 5-HT$_{2A}$ receptor with
antagonists such as ketanserin eliminates their subjective effects (Preller et al., 2017; Quednow et al., 2012;
Vollenweider et al., 1998). 5-HT$_{2A}$ is expressed postsynaptically in the apical dendrites of layer V pyramidal
neurons (Aleksandrova & Phillips, 2021; Hamada et al., 1998; Jakab & Goldman-Rakic, 1998), which are the
primary excitatory neurons in mammalian cortex (Spruston, 2008). Binding to 5-HT$_{2A}$ amplifies the firing
rate of pyramidal neurons (Puig et al., 2003) and upregulates the release of glutamate by increasing the
amplitude and frequency of excitatory postsynaptic potentials (Aghajanian & Marek, 1997, 1999). 5-HT$_{2A}$
expression is profuse in human cortex, with one study finding highest expression in the posterior cingulate

cortex and V1 (Beliveau et al., 2017). In both humans and monkeys, 86-100% of glutamatergic cells in layers II-V of cortex express mRNA encoding 5-HT_{2A} receptors (de Almeida & Mengod, 2007). 5-HT_{2A} has also been identified in many subcortical regions (Nichols, 2016), though human PET imaging implies lower expression than in the cortex (Beliveau et al., 2017).

Although the consciousness-altering effects of psychedelics do primarily arise from their action at the 5-HT_{2A} receptor, different psychedelics display different binding profiles. For instance, unlike DMT and psilocybin, LSD has moderate affinity for dopamine receptors (de Vos et al., 2021). Differences in binding affinities could partially account for the distinctive phenomenology of each psychedelic. There is also some evidence that the 5-HT_{1A} and 5-HT_{2C} receptors may mediate the effects of psychedelics (Canal & Murnane, 2017; Pokorny et al., 2016).

However, binding affinity paints an incomplete picture of the pharmacology of psychedelics. Another key aspect is their functional activity at receptors. Serotonin and dopamine receptors are G protein-coupled receptors (GPCRs) (Gurevich et al., 2016; Shah et al., 2020), which perform essential roles in neurophysiological processes like smell, taste, light perception, and more (de Oliveira et al., 2019). GPCRs regulate these processes through G protein signaling pathways, which are initiated when agonists bind to GPCRs and activate G proteins (Tuteja, 2009). Different families of G proteins are selective for particular signaling pathways (Giulietti et al., 2014). In a phenomenon known as biased agonism, agonists can induce certain conformations in GPCRs, which then selectively activate certain pathways and not others (Andresen, 2011). 5-HT_2 receptors couple preferentially to a family of G proteins known as $G\alpha_{q/11}$, which activates the enzyme phospholipase C (PLC). This enzyme then catalyses the synthesis of the secondary messenger inositol triphosphate (IP_3, a species of inositol phosphate [IP]) , which subsequently leads to the release or "mobilisation" of calcium from the endoplasmic reticulum (López-Giménez & González-Maeso, 2018). Psychedelics are also known to recruit β-arrestin proteins (Pottie, Cannaert, et al., 2020; Pottie, Dedecker, et al., 2020; Rodriguiz et al., 2021; Schmitz et al., 2022). These proteins block the interaction between G proteins and the GPCR and remove the GPCR from the cell membrane, while also coupling GPCRs to signaling proteins without activating G proteins; thus, β-arrestin promotes alternative signaling pathways (Bohn & Schmid, 2010). Functional GPCR assays on psychedelics have tended to focus on the three aspects of GPCR signaling described above: (1) IP formation, (2) calcium mobilisation, and (3) β-arrestin (specifically β-arrestin2) recruitment.

We performed a meta-analysis of the selective affinity of DMT, LSD, and psilocin for the 5-HT_{2A}, 5-HT_{2C}, and D_2 receptors, relative to the 5-HT_{1A} receptor. Additionally, we conducted a meta-analysis of their functional activity at the 5-HT_{2A} receptor, as measured by the three aforementioned assays.

Section 2. Results

We present the results of our meta-analysis of the phenomenology, neuroimaging, and pharmacology of three classical psychedelics: DMT, LSD, and psilocybin. At each level, we measured the alignment between the corresponding results and the Yeo networks, which facilitated comparisons between the three hierarchical levels.

Section 2.1. Phenomenology

We performed a meta-analysis on the 5D- and 11D-ASC scores of DMT, LSD, and psilocybin. Our literature search identified $n = 44$ phenomenology studies that met the inclusion criteria, including 5 studies on DMT (5D-ASC: $n = 3$; 11D-ASC: $n = 3$), 14 studies on LSD (5D-ASC: $n = 9$; 11D-ASC: $n = 12$), and 25 studies on psilocybin (5D-ASC: $n = 12$; 11D-ASC: $n = 17$). The eleven dimensions of the 11D-ASC are subscales of three dimensions in the 5D-ASC: oceanic boundlessness (OB), anxious ego dissolution (AED), and visionary restructuralisation (VR). Nevertheless, the 11D-ASC questionnaire contains fewer items than the 5D-ASC questionnaire, so 5D-ASC data cannot be directly compared to 11D-ASC data. Hence, we ran separate meta-analyses for the 5D-ASC and 11D-ASC data. In order to account for the measurement of multiple subjective dimensions, sometimes with more than one dose or drug, in individual studies, we performed a multilevel random-effects meta-analysis (Assink & Wibbelink, 2016; Harrer et al., 2021).

DMT was administered intravenously (IV) in the phenomenological studies, whereas LSD and psilocybin were administered orally (we excluded phenomenological studies that used IV LSD and psilocybin). IV administration yields different pharmacokinetics, which are known to influence the subjective experience of the drug (Luan et al., 2023; Vogt et al., 2023). Therefore, while we conducted meta-analyses of all three drugs, we only examined significant between-drug differences for ASC ratings of LSD and psilocybin. Additionally, we only compared similar doses of LSD and psilocybin. We assumed 20 mg psilocybin is equivalent to 0.01 mg LSD (Ley et al., 2023), and we defined low, medium, and high doses based on the literature (Carbonaro et al., 2018; Hasler et al., 2004; Vollenweider et al., 2007).

Pooled 5D-ASC scores for DMT, LSD, and psilocybin are shown in **Figure 1** (numerical data are given in **Table S1**). Between-drug differences for LSD and psilocybin are displayed in the left column of **Figure 1a-c**, and within-drug differences are displayed in the right column. At medium doses (LSD: 0.075-0.109 mg, psilocybin: 15-21 mg), 5D-ASC scores were significantly greater for LSD than psilocybin in the OB dimension ($p = 0.0283$), which corresponds to feelings of interconnectedness, and the VR dimension ($p = 0.0468$), which measures the quality and intensity of visual hallucinations. We also observed a significant difference between LSD and psilocybin in the VR dimension ($p = 0.0417$) at high doses (LSD: ≥0.010 mg, psilocybin: ≥22 mg). There were no significant differences at low doses (LSD: 0.050-0.074 mg, psilocybin: 8-14 mg).

Within-drug differences between the five subjective dimensions were quite similar. A clear separation can be seen between two groups of subjective categories: (1) OB, VR, RV, and (2) AA and AED. At medium doses, VR and OB scores were significantly higher than both AED and AA scores for both LSD and psilocybin. For DMT, there were no significant differences between dimensions at high doses (≥16 mg bolus injection, followed by continuous infusion of 1 mg/min), but for low doses (10-15 mg bolus + 0.6-0.99 mg/min continuous infusion), AED ranked significantly lower than both OB and VR. (Note that only one study measured AA and RV scores for DMT (Vogt et al., 2023), so we did not perform a meta-analysis on these dimensions for DMT.)

The results of the 11D-ASC meta-analysis were not very consistent with those of the 5D-ASC meta-analysis (**Figure S3**; **Table S2**). Note that we could not include DMT in the 11D-ASC meta-analysis because two of the three studies on DMT did not report standard errors, which are necessary for pooling the subjective data. At high doses, there were no significant differences between LSD and psilocybin in any of the dimensions, and standard errors were very large for psilocybin in some dimensions, particularly in some OB and VR subscales. At medium doses, where standard errors tended to be lower, we observed that two

of the OB subscales (experience of unity and insightfulness), one of the AED subscales (impaired control and cognition), and three of the VR subscales (complex imagery, audio-visual synaesthesia, and changed meaning of percepts) ranked significantly higher for LSD than psilocybin. In line with the 5D-ASC results, we did not find any significant differences between LSD and psilocybin at low doses. However, unlike the 5D-ASC analysis, the 11D-ASC analysis showed that psilocybin rated higher than LSD in most of the dimensions; that being said, standard errors were very high for both LSD and psilocybin, so our estimates are not reliable. For medium doses, within-drug comparisons for the 11D-ASC analysis generally aligned with those of the 5D-ASC analysis.

We did not find any significant relationship between the pooled 5D-ASC scores and three methodological covariates that were defined *a priori*: prior psychedelic use, the time that the questionnaire was administered relative to the drug, and the presence or absence of a task in the experiment. Additionally, there was significant residual heterogeneity in the pooled 5D- and 11D-ASC scores ($p < 0.0001$ in both cases). Unfortunately, we were unable to obtain reliable estimates of heterogeneity specifically attributable to within-study and between-study differences. For three of the 5D-ASC scales, we found evidence of significant publication bias with Egger's regression test, which determines whether small studies report disproportionately high effect sizes (OB: $p = 0.0187$, AED: $p < 0.0001$, VR: $p = 0.6279$, AA: $p < 0.0001$, RV: $p = 0.6532$). The results of our risk-of-bias assessments for individual studies are shown in **Table S3**. Due to the moderate-to-serious risk of bias, significant residual heterogeneity, high probability of publication bias, and indirectness (some studies measured subjective effects while participants were performing a specific task, as opposed to during resting-state), our certainty in the body of phenomenological evidence is low.

In order to directly compare the phenomenology of psychedelics to their neuroimaging and pharmacological profiles (*Section 2.2* and *Section 2.3*), we sought to determine the "phenomenology profiles" of DMT, LSD, and psilocybin based on the neural correlates of their subjective effects. We performed a separate literature search of studies that measured correlations between ASC ratings and fMRI activity. We determined the Yeo network that contained each brain region that exhibited a significant correlation. The phenomenology profiles were defined by a weighted combination of the mean correlations between each ASC scale and each Yeo network, in which the weights were the pooled ASC ratings of high doses of each psychedelic for the corresponding scale.

Because there were few significant differences in the pooled ASC scores, the phenomenology profiles of the psychedelics look very similar (**Figure 2**). However, these profiles may not reflect the true neural correlates of the subjective effects of psychedelics. Many studies performed correlations between ASC ratings and brain regions that were active only in specific tasks, but the tasks varied strongly in the literature. The ventral attention network (VAN) appears to be more strongly associated with the subjective effects than any of the other Yeo networks, but this result may be an artifact of one study's decision to correlate the ASC ratings with a specific seed in the VAN. There need to be more studies that link whole-brain fMRI activity to ASC ratings before we can estimate the phenomenology profiles with confidence.

Finally, we also aimed to relate the phenomenology of different psychedelics by qualitatively synthesising the findings of papers that directly compared the subjective experiences of DMT/ayahuasca, LSD, and psilocybin. *Section S3* contains a comprehensive summary of their major aims, analysis approaches and findings.

Section 2.2. Neuroimaging

As stated above, the fMRI studies on psychedelics measure three different characteristics of brain activity: BOLD activation, entropy, and connectivity. We identified $n = 17$ studies on BOLD activation, including $n = 3$ studies on ayahuasca or DMT, $n = 7$ studies on LSD, and $n = 4$ studies on psilocybin. Initially, we used the GingerALE method to determine common clusters of BOLD activation across studies. We observed that psilocybin was associated with a cluster of activity in visual cortices, whereas LSD tended to affect more frontal areas. However, three of the four studies on psilocybin displayed visual stimuli to participants. Therefore, our results were likely confounded by the tasks that were used in the literature. We chose not to report the results of our GingerALE analysis because they are likely misleading.

The $n = 12$ fMRI studies on entropy and criticality in the psychedelic literature are very heterogeneous, employing a wide variety of measures on dissimilar variables of interest. A quantitative meta-analysis was extremely challenging, so we opted instead to perform a qualitative review of this literature, which is reported in *Section S4*.

The studies on connectivity were the only segment of the fMRI literature for which we determined that we could conduct a valid meta-analysis. These studies tended to use the same resting-state experimental procedure, and many of them employed similar methods for measuring connectivity. However, one major obstacle was the variety of parcellations in the literature. Since each study defined regions of interest (ROIs) in a different way, it was not possible to simply perform a weighted average of connectivity values across studies. (This weighted average method is essentially the core technique of the phenomenology and pharmacology meta-analyses.) An alternative approach is to "re-reference" the data to a coarse-grained parcellation: the Yeo networks, of which there are only seven in the brain (Thomas Yeo et al., 2011). That is, for every pair of functionally connected ROIs in the literature, we determine the corresponding Yeo networks that contain them. If the connection is positive (it increases under psychedelics), then the aggregate FC between those Yeo networks is incremented; otherwise, it is decremented. Our method for weighting each connection in the literature is described in *Section S2.2.2*. We determined significance by repeatedly applying our method to an independent resting-state fMRI dataset of sober, healthy individuals (Van Essen et al., 2013), which resulted in a null distribution.

This method is capable of synthesising studies on *pairwise* FC, such as seed-to-seed, seed-to-voxel, and within- and between-network FC analyses using independent components analysis (see **Table S5** for a full description of the inputs). The algorithm cannot accommodate global, structural metrics of connectivity such as global brain connectivity (GBC) or graph-theoretic modularity. Additionally, the algorithm does not capture any information about the direction of connectivity between regions; therefore, we did not incorporate studies of effective connectivity or directed FC. (However, the vast majority of FC studies measured undirected FC.)

Many FC studies are secondary analyses of the same primary dataset. For instance, the primary FC data from Carhart-Harris et al. (2016) was re-analysed at least seven times with different methods in subsequent studies. To mitigate bias, we only inputted the most informative analysis of each unique dataset into the meta-analysis algorithm, such that each dataset was only represented once in the meta-analysis. Our method for selecting the most informative analysis is described in *Section S2.2.2*.

Based on these criteria and others, we ultimately identified $n = 12$ studies on connectivity that were eligible for our meta-analysis, including $n = 3$ studies on ayahuasca or DMT, $n = 3$ studies on LSD, and $n = 6$ studies on psilocybin. (There are 22 other studies on connectivity that we qualitatively review in *Section S5*.) Our

estimates of aggregate FC between Yeo networks, across all psychedelics, are shown in **Figure 3**. All pairwise
changes in aggregate FC were deemed to be significant. In general, psychedelics significantly strengthened
between-network connectivity, except between three pairs of networks: the visual network & somatomotor
network (SMN), SMN & limbic network, and the ventral attention network (VAN) & limbic network.
Within-network connectivity significantly decreased in all networks except for the SMN and frontoparietal
network (FPN).

In addition to aggregating FC across all psychedelics, we performed our analysis separately on each
individual psychedelic. All three of the psychedelics lowered FC in the visual network and default mode
network (DMN) and generally elevated between-network FC. Intriguingly, ayahuasca/DMT
disproportionately augmented FC between two "transmodal" or associative networks (FPN and DMN) &
the two "unimodal" or sensory networks (visual network and SMN) while strongly reducing FC within and
between the two sensory networks. LSD dramatically increased FC between the visual network and all other
networks, yet sharply decreased FC within the visual network. Summing across the rows of each aggregate
FC matrix, we find that DMT and psilocybin tend to affect the FC of the DMN the most, whereas LSD
displays a more uniform profile of changes in FC across the Yeo networks (**Figure 6**).

Finally, there was no significant association between our results and any of the methodological covariates
in the literature that we identified *a priori*: the width of the Gaussian kernel used to smooth the data, the use
of FSL or SPM, the technique used to regress out white matter, and the use of scrubbing to correct head
motion. The results of our risk-of-bias assessment for individual studies is shown in **Table S6**.
379

Section 2.3. Pharmacology

381 We performed two separate meta-analyses on the pharmacology of psychedelics. The first assessed the
382 selective affinity of DMT, LSD, and psilocin for the 5-HT_{2A}, 5-HT_{2C}, and D_2 receptors, relative to 5-HT_{1A}.
383 The second examined the relative functional activity of the psychedelics at the 5-HT_{2A} receptor, as captured
384 in three different assays of GPCR signaling.
385

Section 2.3.1. Affinity

We first present our meta-analysis of the selective affinity data. Selectivity is defined as the ratio between the
K_i of each psychedelic for a receptor of interest, relative to a reference receptor. K_i refers to the inhibition
constant, which reflects the concentration of a drug that is needed to inhibit the binding of another ligand,
for instance a radioactively-labeled ligand (radioligand), to the receptor of interest; it is inversely related to
binding affinity (Sykes et al., 2019). We chose 5-HT_{1A} to be the reference receptor. We decided to analyse
selectivity rather than absolute K_i because the latter is biased by the potency of the drug. Since LSD is much
more potent than DMT and psilocin (Rickli et al., 2016), the meta-analysis would have simply revealed that
LSD has higher affinity for all receptors if we were to examine the absolute K_i instead.

Our literature search identified $n = 14$ studies on selectivity, including $n = 6$ studies on DMT, $n = 9$ studies
on LSD, and $n = 5$ studies on psilocin. We performed a random-effects meta-analysis, the results of which
are displayed in **Figure 5a** (numerical data are given in **Table S7**). We found no significant between-drug
differences in selectivity for any of the receptors – 5-HT_{2A}, 5-HT_{2C}, and D_2 – relative to the 5-HT_{1A} receptor.
Standard errors were very large for all of the pooled selectivity values, and heterogeneity was very high for
the 5-HT_{2A} selectivity data ($I^2 = 93.69\%$) and for the 5-HT_{2C} selectivity data ($I^2 = 99.14\%$). Nevertheless, it
is clear that all drugs display less selectivity for D_2 than they do for 5-HT_{2C} and 5-HT_{2A}. Each individual drug,

especially LSD and psilocin, is more selective for 5-HT$_{2A}$ than 5-HT$_{1A}$. DMT and psilocin are about as equally
selective for 5-HT$_{2C}$ as they are for 5-HT$_{1A}$, but LSD is much less selective for 5-HT$_{2C}$ than for 5-HT$_{1A}$.
We constructed another random-effects model in which the radioligand for the receptor of interest was
included as a covariate; we refer to this as the "full" model and the previous one, in which the influence of
the radioligand was not modeled, as the "reduced" model. For 5-HT$_{2A}$, the use of the radioligands [^{3}H]-
ketanserin ($p = 0.0141$) and [^{3}H]-spiperone ($p = 0.0271$) significantly influenced the pooled selectivity
estimates. Use of [^{3}H]-mesulgerine ($p = 0.0292$), but not [^{3}H]-ketanserin ($p = 0.0872$), significantly affected
our estimates of selectivity for 5-HT$_{2C}$. According to the corrected Akaike's information criterion, the full
model performed significantly better than the reduced model for 5-HT$_{2A}$ selectivity ($p < 0.0001$), whereas
the reverse was true for 5-HT$_{2C}$ selectivity ($p = 0.0024$).
We assessed publication bias due to small study effects and found significant evidence for bias in the 5-
HT$_{2A}$ ($p = 0.0001$) and 5-HT$_{2C}$ ($p = 0.0004$) affinity data. (There were not enough studies on D$_2$ affinity to
measure bias.) Because the affinity studies are not interventional, we could not measure risk of bias in
individual studies with the same tool that we used in the phenomenology and neuroimaging meta-analyses.
Instead, in *Section S6.3*, we qualitatively describe some major confounders in the affinity literature. Due to
the large imprecision of the results and high probability of publication bias, our certainty in the body of
evidence about selectivity is low.
To relate the selective affinities to the neuroimaging and phenomenology of psychedelics, we first
determined the expression of the 5-HT$_{2A}$ and D$_2$ receptors in the Yeo networks, based on an atlas of PET
maps (Hansen et al., 2022) (unfortunately, this did not include maps of the 5-HT$_{2C}$ receptor). Then, we
defined the "pharmacology profile" as the weighted combination of the expression patterns, in which the
weights were the selectivity of each psychedelic for the corresponding receptor (**Figure 6**). The PET atlas
indicates that 5-HT$_{2A}$ receptor density is around two orders of magnitude higher than that of D$_2$ receptor
density, so the pharmacology profile predominantly reflects selectivity for 5-HT$_{2A}$. Because psilocin's
selectivity for 5-HT$_{2A}$ is greater than that of LSD and DMT, psilocin has the "largest" pharmacology profile.
5-HT$_{2A}$ is expressed more in transmodal networks than the visual network; thus, selectivity for 5-HT$_{2A}$ may
affect brain activity the most in these networks.
429

Section 2.3.2. Functional activity
Binding affinity is only one facet of the pharmacology of psychedelics. Another key aspect is their functional
activity: the responses that they elicit in receptors after binding to them. While there are many ways to
measure functional activity, we focused on GPCR signaling through three different pathways: (1) IP
formation, (2) calcium mobilisation, and (3) β-arrestin2 recruitment. We limited our analysis to functional
activity at specifically the 5-HT$_{2A}$ receptor, as there was much more data about this receptor in humans. The
variable of interest in our analysis was $\Delta\log(E_{max}/EC_{50})$.
We found $n = 18$ relevant studies on functional activity, including $n = 6$ studies on DMT, $n = 13$ studies
on LSD, and $n = 6$ studies on psilocin. We performed a random-effects meta-analysis. For the IP formation
and calcium mobilisation assays, the reference ligand was serotonin; for the β-arrestin2 recruitment assays,
the reference ligand was chosen to be mescaline, as this was the only reference ligand used in experiments
on all three psychedelics. The pooled relative activity values indicate that LSD induces significantly more IP
formation than DMT ($p < 0.0001$) and psilocin ($p = 0.0002$) (**Figure 5b**; numerical data given in **Table S8**).
Additionally, DMT elicits significantly more IP formation than psilocin ($p = 0.0203$). There were no

significant between-drug differences for the other two pathways. Heterogeneity was very high for all three pathways (IP formation: $I^2 = 95.35\%$, calcium mobilisation: $I^2 = 92.66\%$, β-arrestin2 recruitment: $I^2 = 91.05\%$). There was no significant publication bias due to small study effects for the literature on IP formation ($p = 0.3460$). (There were too few studies on the other pathways to assess publication bias.) Because of the large standard errors of the pooled estimates and high unexplained heterogeneity, our certainty in the body of evidence about functional activity is low, despite the lack of evidence for publication bias.

Section 3. Discussion

Here we examine the effects of psychedelics at three levels: 1) subjective experience (phenomenology), 2) functional connectivity (neuroimaging), and 3) the interaction of psychedelics with serotonin and dopamine receptors (pharmacology). At each level, we performed a quantitative meta-analysis and computed the alignment between the results and the seven Yeo networks. The latter network analysis enabled us to directly compare the effects of three psychedelics – ayahuasca/DMT, LSD, and psilocybin – within and between levels.

Section 3.1. Unifying the levels of analysis

For the phenomenology literature, we conducted a meta-analysis of both the 5-dimensional and 11-dimensional versions of the Altered States of Consciousness (ASC) scale, a common questionnaire for measuring changes in subjective experience on psychedelics. At both medium and high doses, LSD ranked significantly higher than psilocybin in "visionary restructuralisation," a dimension capturing the quality and intensity of visual hallucinations. Additionally, at medium doses, LSD was associated with significantly higher scores in the "oceanic boundlessness" dimension, which captures feelings of interconnectedness. The VAN appears to be correlated with the phenomenology of all three psychedelics, but this result may be spurious due to the small number of studies that measured correlations between ASC ratings and fMRI activity. Nevertheless, there are reasons to think that the VAN may be uniquely implicated in the phenomenology of psychedelics. The VAN is responsible for orienting attention toward surprising objects or events (Vossel et al., 2014), and it is engaged during mismatch negativity (MMN), an event-related potential that is triggered when an unexpected stimulus is shown (Lee et al., 2017). Psychedelics diminish the MMN response while enhancing the neural response to familiar stimuli (Duerler et al., 2021; Timmermann et al., 2018), perhaps because these drugs engender profound feelings of childlike awe and wonder (Hendricks, 2018). This heightened predisposition towards subjective experiences of novelty may increase activity in the VAN.

The whole-brain neuroimaging data consisted of three subsets of data: BOLD activation, entropy, and FC. We did not report a meta-analysis of the BOLD activation or entropy literature due to the heterogeneity of experimental procedures and analysis methods, respectively; however, we did conduct a qualitative review of the entropy literature (*Section S4*). To perform a meta-analysis on the FC data, we essentially re-parcellated the published data into the Yeo networks and then calculated a weighted sum of the connections between Yeo networks. LSD strongly elevated FC between the visual network and the other networks, whereas ayahuasca/DMT increased FC most between transmodal networks (specifically, the FPN and DMN) and unimodal networks (i.e. SMN and visual).

For pharmacology, we conducted two meta-analyses: one of selectivity and another of relative functional activity. There were no significant differences between psychedelics in selectivity for the 5-HT$_{2A}$, 5-HT$_{2C}$, or

D$_2$ receptors relative to the 5-HT$_{1A}$ receptor. Compared to both DMT and psilocin, LSD elicited significantly
more relative activity in IP formation assays. We did not observe any significant between-drug differences
for the calcium mobilisation and β-arrestin2 recruitment assays.

Directly comparing the pharmacology, neuroimaging, and phenomenology profiles of psychedelics
(**Figure 7**) reveals a weak one-to-one relationship between the three levels of analysis, which is unsurprising
given the highly non-linear, complex, and reciprocal (Kringelbach et al., 2020) interactions between them.
However, the results of the meta-analyses at each level may reveal some insights into the nature of these
interactions.

On a superficial level, there do appear to be some direct correspondences between the phenomenology,
neuroimaging, and pharmacology. Phenomenologically, we only found one significant difference between
psychedelics that was consistent across medium and high doses: LSD induces more visionary
restructuralisation than psilocybin does. Neurobiologically, LSD disproportionately enhanced connectivity
between the visual network & all other networks, while very strongly reducing connectivity within the visual
network. LSD's effect on FC between other networks was much weaker.

The relationship between the pharmacology results and the above findings is more difficult to establish.
Our results show that LSD increases IP formation, whereas DMT and psilocin decrease it. IP formation
occurs whenever Gα$_q$ proteins are recruited to initiate cellular signaling. Furthermore, there is evidence that
Gα$_q$ activation is necessary for 5-HT$_{2A}$ agonists to bring about hallucinogenic effects. When administered the
psychedelic (±)1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane (DOI), mice that lack the gene for Gα$_q$
proteins do not exhibit behaviours (head twitches) that typically indicate hallucinations (Garcia et al., 2007).
More recently, activation of Gα$_q$ pathways by 5-HT$_{2A}$ agonists has been shown to predict head-twitch
behaviour, whereas 5-HT$_{2A}$ agonists that are biased for the β-arrestin2 pathway do not induce head twitches
(Wallach et al., 2023). There appears to be a threshold level of Gα$_q$ activation that must be surpassed in order
for a 5-HT$_{2A}$ agonist to induce hallucinogenic effects (Wallach et al., 2023). Intriguingly, the pro-psychotic
properties of drugs can be predicted from the extent to which they increase Gα$_q$ activation (relative to a
different G-protein pathway) when these drugs interact with a complex formed between the 5-HT$_{2A}$ receptor
and a metabotropic glutamate receptor (Fribourg et al., 2011). In other words, the more that a drug upsets
the normal balance of Gα$_q$ signaling, the more likely it is to engender hallucinations. However, these studies
only establish the presence of hallucinations elicited by Gα$_q$ signaling. They do not indicate whether Gα$_q$
signaling induces *visual* hallucinations in particular. Thus, based on the available evidence, we cannot yet
conclude that the pharmacology, neuroimaging, and phenomenology literature all convergently indicate that
LSD has a uniquely strong effect on visual awareness, compared to the other psychedelics.

There do not appear to be any tools at the moment for relating the pharmacology of drugs to the precise
nature of the subjective experiences that they produce. Methods for examining the relationship between
neuroimaging and phenomenology are currently limited to correlations and linear regressions (Burt et al.,
2021; Carhart-Harris et al., 2016), which do not capture causality. There is a need to develop techniques for
elucidating the causal relationships between phenomenology, neuroimaging, and pharmacology.

Section 3.2. Relating the results of the meta-analysis to the literature

Section 3.2.1. Phenomenology

We sought to verify the results of our phenomenology meta-analysis by comparing them to the two
experimental studies that measured ASC ratings of LSD and psilocybin in the same group of participants.

Holze et al. (2022) gave a medium (15 mg) and a high (30 mg) dose of psilocybin, as well as a medium (0.01
mg) and a high (0.02 mg) dose of LSD, and then compared both 5D- and 11D-ASC scores. It is worth noting
that the doses of psilocybin and LSD in this study are not equivalent, since 0.01 mg LSD = 20 mg psilocybin
(Ley et al., 2023). When comparing 15 mg psilocybin to 0.01 mg LSD, mean 5D-ASC and 11D-ASC scores
tended to be higher for LSD than psilocybin. However, the only differences that reached significance were
in the OB and VR scales for the 5D-ASC analysis, and in the complex imagery and audio-visual synaesthesia
subscales for the 11D-ASC analysis. In our meta-analysis, LSD and psilocybin were significantly different in
both OB and VR ratings, as well as in complex imagery and audio-visual synaesthesia ratings, at medium
doses. Ley et al. (2023) compared the subjective effects of medium doses of LSD (0.01 mg) and psilocybin
(20 mg). Here, the doses are equivalent, unlike in the previous study. The study found no significant
differences between LSD and psilocybin in any of the dimensions, both in the 5D and 11D analyses.
Therefore, the results of our meta-analysis are not entirely supported by studies that conducted direct
comparisons between psychedelics.
541
Section 3.2.2. Neuroimaging
One major finding in the neuroimaging literature is that psychedelics desegregate and disintegrate brain
networks. Segregation is defined as the lack of FC *between* brain regions, while integration denotes FC *within*
a network. Carhart-Harris et al. (2016) found that LSD led to disintegration and desegregation for most
RSNs. This finding has been independently confirmed in several other studies (Barrett, Doss, et al., 2020;
Dai et al., 2023; Girn et al., 2022; Lebedev et al., 2015; Madsen et al., 2021; F. Müller, Dolder, et al., 2018;
Roseman et al., 2014). In a review of three studies, Müller, Liechti, et al. (2018) found good convergence in
reports of desegregation.
Our meta-analysis generally corroborates the desegregation and disintegration hypothesis. FC significantly
increased between most, though not all, non-identical pairs of brain networks. Within-network FC
significantly decreased for all networks except for the SMN and FPN. In this meta-analysis, we determined
statistical significance by comparing the psychedelic literature to resting-state FC data collected by the
Human Connectome Project in sober human participants. However, the HCP data does not reflect the
heterogeneity of analysis methods in the psychedelic literature. Whereas we used a whole-brain seed-to-seed
FC analysis of the HCP data, many of the studies in the psychedelic literature isolated specific seeds rather
than examining the entire brain. Additionally, we applied a single parcellation (DK-80) to the HCP data,
whereas multiple parcellations were used in the psychedelic literature. Thus, the insignificant result may
reflect inconsistencies between our methods and the literature. Nevertheless, it does raise the intriguing
possibility that reductions in within-network FC can sometimes be greater in sober resting-state than on
psychedelics.
While there is a correlation between desegregation/disintegration and the subjective experience of ego
dissolution (Tagliazucchi et al., 2016), there is no evidence for a *causal* link. There appear to be parallels
between the conscious experience of interconnectedness and the "interconnectedness" of the brain on
psychedelics, but further research is needed to demonstrate that the connection between the two is more
than merely semantic.
Another common finding in the literature is that psychedelics decrease activity and connectivity within
the DMN, a region that is generally implicated in self-awareness, self-reflection, and other self-referential
cognitive processes (Davey et al., 2016; Davey & Harrison, 2018; Molnar-Szakacs & Uddin, 2013; Moran et

al., 2013). This popular 'meme' likely has its origin in the first resting-state fMRI study of a psychedelic, where decreased blood flow and BOLD signal were observed in a pattern of regions resembling the DMN (Carhart-Harris et al., 2012). Eleven years later, evidence continues to indicate that psychedelics potently affect the regularity of population brain activity in DMN regions (Timmermann et al., 2023). Since a commonly reported experience on psychedelics is the dissolution of the self, a phenomenon known as "ego death" (Moreton et al., 2020; Nour et al., 2016; Pahnke, 1969), it would superficially make sense that psychedelics dysregulate the DMN. Many studies have found significant correlations between subjective ratings of ego dissolution and DMN disintegration (Atasoy et al., 2017; Barrett, Krimmel, et al., 2020; Luppi et al., 2021; Madsen et al., 2021; Palhano-Fontes et al., 2015; Pasquini et al., 2020; Roseman et al., 2014; Smigielski et al., 2019; Tagliazucchi et al., 2014, 2016), though other analyses did not find evidence for a relationship between the two (Lebedev et al., 2015; F. Müller, Dolder, et al., 2018).

However, claims of "de-activation" or worse, "shutting-off" of the DMN as signatures of psychedelic action are wrong and misleading. Other neuroimaging studies by Carhart-Harris et al. (2012, 2013, 2016) and Roseman et al. (2014) discovered that psychedelics affect the *connectivity* of the DMN, decreasing within-DMN FC ("disintegration") while increasing DMN coupling with other resting-state networks (RSNs) ("desegregation"). Our neuroimaging meta-analysis confirms that psychedelics significantly decrease connectivity within the DMN while significantly elevating its connectivity with all other networks. However, it is worth noting that the reduction in the within-network FC of the DMN is smaller than that of the VAN, limbic network, and visual network. In fact, the drop in within-network FC is nearly 58 times greater in the visual network than the DMN. Our phenomenology meta-analysis revealed that all psychedelics induced positive experiences of depersonalisation and derealisation, as captured by the oceanic boundlessness dimension of the 5D-ASC scale; these experiences are related to the feeling of ego death.

Nevertheless, there are many cases in which within-DMN connectivity may decrease without giving rise to the subjective effects that characterise psychedelics. Several other drugs such as amphetamines and alcohol, as well as drug addiction in general, reduce within-DMN connectivity in humans (Schrantee et al., 2016; Weber et al., 2014; Zhang & Volkow, 2019), even though they typically do not lead to ego dissolution (Doss et al., 2020). Certain psychological states that exacerbate self-rumination, such as maladaptive self-focused attention, are *not* associated with hyperconnectivity in the DMN (Fang et al., 2022), and one study found that some tasks requiring self-related judgments were associated with a *suppression* of within-DMN connectivity (van Buuren et al., 2010). That being said, a recent meta-analysis of 14 studies did endorse an association between rumination and the DMN (Zhou et al., 2020).

Section 3.2.3. Pharmacology
The results of our affinity meta-analysis aligned well with studies that performed direct comparisons of selectivity between different psychedelics (Eshleman et al., 2014; Janowsky et al., 2014; Kozell et al., 2023; Pierce & Peroutka, 1989; Rickli et al., 2016). In line with our findings, all of these studies showed that LSD is more selective than DMT for 5-HT$_{2A}$, relative to 5-HT$_{1A}$. Two studies also observed that psilocin is more selective than LSD for 5-HT$_{2A}$ (Kozell et al., 2023; Rickli et al., 2016). Our meta-analysis demonstrated that DMT is more selective than LSD for 5-HT$_{2C}$, a conclusion that is supported by three of four studies (Eshleman et al., 2014; Janowsky et al., 2014; Kozell et al., 2023). Note that none of these studies measured the statistical significance of between-drug differences in selectivity.

Our findings about functional activity were also congruent with studies that measured the effect of multiple psychedelics on IP formation at the 5-HT$_{2A}$ receptor (Braden & Nichols, 2007; Eshleman et al., 2014; Janowsky et al., 2014; Kozell et al., 2023). We found that LSD's relative activity at this pathway was significantly greater than that of psilocin, which was in turn significantly larger than that of DMT. All four studies showed that LSD had higher relative activity at this pathway than DMT. Two studies also reported that psilocin's relative activity was lower than that of LSD, but larger than that of DMT (Braden & Nichols, 2007; Kozell et al., 2023).

Section 3.3. Recommendations for future research

We encourage researchers to develop tools for modelling the nonlinear relationships between the pharmacology, neuroimaging, and phenomenology of psychedelics, the three hierarchical levels of analysis that are at the heart of this meta-analysis. While some researchers have attempted to examine the associations between each level (Ballentine et al., 2022; Lawn et al., 2022), their techniques are model-free and only measure *correlations* between the effects of psychedelics, rather than elucidating the *causal mechanisms* that underpin them.

Recent whole-brain models have made significant advances in discovering these causal mechanisms (see Kringelbach & Deco (2020) for a general review). Whole-brain models consist of nodes that approximate local or mesoscopic neuronal dynamics through mean field models of coupled excitatory-inhibitory interactions (Deco et al., 2014; Honey et al., 2007) or Hopf models of bifurcations into and out of sustained oscillatory activity (Deco et al., 2017; Freyer et al., 2012). Crucially, whole-brain models have enabled researchers to identify the specific regions that should be stimulated in order to force transitions between different states of consciousness, such as from sleep to wakefulness (Deco et al., 2019). Whole-brain models have also been applied to psychedelics; in particular, the 5-HT$_{2A}$ receptor density in each area of the brain was used to fit a mean field model of fMRI data on LSD, capturing the nonlinear interactions between connectivity and the distribution of 5-HT$_{2A}$ receptors (Deco et al., 2018). This model was recently extended to determine the associations between global brain connectivity on LSD and categories of subjective experience (Burt et al., 2021). The model has also been applied to the relationship between neurotransmitters (namely, serotonin) and neuronal regions on psilocybin; in particular, Kringelbach et al. (2020) coupled a mean field model of each neuronal region to the release-and-reuptake dynamics of serotonin concentration. Importantly, this coupled neuronal-neurotransmitter model exhibited a better fit to the empirical FC than a model of solely the neuronal activity. Overall, whole-brain models display great promise for linking neurotransmitter release, which is partly determined by pharmacology, to the brain regions that causally mediate, rather than merely correlate with, the effects of psychedelics. Most recently, whole-brain models have also been combined with another cutting-edge method, turbulence, to determine the susceptibility of the brain to external stimuli under psychedelics (Cruzat et al., 2022).

Section 4. Conclusion

This is the first meta-analysis of the literature on the phenomenology, neuroimaging, and pharmacology of psychedelics. We assess and compare these three hierarchical levels of analysis across three different classical psychedelics: DMT, LSD, and psilocybin. Thus, we performed comparisons not only between psychedelics, but also between the levels of the hierarchy.

652   We found that the different levels of the hierarchy exhibited a weak one-to-one correspondence with one
653 another. This is not surprising, as the interactions between phenomenology, neuroimaging, and
654 pharmacology are highly non-linear and complex. We encourage future research to develop tools for
655 modelling these relationships in order to improve our scientific understanding of psychedelics.
656

Acknowledgments

K.S. is funded by the Clarendon Fund, a Department of Psychiatry studentship, the John Henry Jones
Scholarship from Balliol College, and the Research Council U.K. P.M. is funded by the Wellcome Trust
(grant no. 210920/Z/18/Z). K.P. is funded by the Swiss National Science Foundation (P2ZHP1\161626).
R.C-H is funded by a Ralph Metzner endowment. M.K. is supported by the Centre for Eudaimonia and
Human Flourishing (funded by the Pettit and Carlsberg Foundations) and Center for Music in the Brain
(funded by the Danish National Research Foundation, DNRF117).

Code and Data Availability
Code and data used for the phenomenology and pharmacology analyses are available upon request by
emailing the corresponding author: kenneth.shinozuka@psych.ox.ac.uk
668

669 **Competing Interests Statement**
670 The authors have no conflicts of interest to declare.
671

672 **Study Registration and Protocol**
673 This study was not registered. A review protocol was not prepared.
674

References

Aday, J. S., Davis, A. K., Mitzkovitz, C. M., Bloesch, E. K., & Davoli, C. C. (2021). Predicting Reactions to Psychedelic Drugs: A Systematic Review of States and Traits Related to Acute Drug Effects. *ACS Pharmacology & Translational Science*, *4*(2), 424–435. https://doi.org/10.1021/acsptsci.1c00014

Aghajanian, G. K., & Marek, G. J. (1997). Serotonin induces excitatory postsynaptic potentials in apical dendrites of neocortical pyramidal cells. *Neuropharmacology*, *36*(4–5), 589–599. https://doi.org/10.1016/s0028-3908(97)00051-8

Aghajanian, G. K., & Marek, G. J. (1999). Serotonin, via 5-HT2A receptors, increases EPSCs in layer V pyramidal cells of prefrontal cortex by an asynchronous mode of glutamate release. *Brain Research*, *825*(1–2), 161–171. https://doi.org/10.1016/s0006-8993(99)01224-x

Aleksandrova, L. R., & Phillips, A. G. (2021). Neuroplasticity as a convergent mechanism of ketamine and classical psychedelics. *Trends in Pharmacological Sciences*, *42*(11), 929–942. https://doi.org/10.1016/j.tips.2021.08.003

Andresen, B. T. (2011). A Pharmacological Primer of Biased Agonism. *Endocrine, Metabolic & Immune Disorders Drug Targets*, *11*(2), 92–98. https://doi.org/10.2174/187153011795564179

Araújo, A. M., Carvalho, F., Bastos, M. de L., Guedes de Pinho, P., & Carvalho, M. (2015). The hallucinogenic world of tryptamines: An updated review. *Archives of Toxicology*, *89*(8), 1151–1173. https://doi.org/10.1007/s00204-015-1513-x

Atasoy, S., Roseman, L., Kaelen, M., Kringelbach, M. L., Deco, G., & Carhart-Harris, R. L. (2017). Connectome-harmonic decomposition of human brain activity reveals dynamical repertoire re-organization under LSD. *Scientific Reports*, *7*(1), Article 1. https://doi.org/10.1038/s41598-017-17546-0

Ballentine, G., Friedman, S. F., & Bzdok, D. (2022). Trips and neurotransmitters: Discovering principled patterns across 6850 hallucinogenic experiences. *Science Advances*, *8*(11), eabl6989. https://doi.org/10.1126/sciadv.abl6989

Barrett, F. S., Doss, M. K., Sepeda, N. D., Pekar, J. J., & Griffiths, R. R. (2020). Emotions and brain function are altered up to one month after a single high dose of psilocybin. *Scientific Reports*, *10*(1), 2214. https://doi.org/10.1038/s41598-020-59282-y

Barrett, F. S., Johnson, M. W., & Griffiths, R. R. (2015). Validation of the revised Mystical Experience Questionnaire in experimental sessions with psilocybin. *Journal of Psychopharmacology*, *29*(11), 1182–1190. https://doi.org/10.1177/0269881115609019

Barrett, F. S., Krimmel, S. R., Griffiths, R. R., Seminowicz, D. A., & Mathur, B. N. (2020). Psilocybin acutely alters the functional connectivity of the claustrum with brain networks that support perception, memory, and attention. *NeuroImage*, *218*, 116980. https://doi.org/10.1016/j.neuroimage.2020.116980

Beliveau, V., Ganz, M., Feng, L., Ozenne, B., Højgaard, L., Fisher, P. M., Svarer, C., Greve, D. N., & Knudsen, G. M. (2017). A High-Resolution In Vivo Atlas of the Human Brain's Serotonin System. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *37*(1), 120–128. https://doi.org/10.1523/JNEUROSCI.2830-16.2016

Bogenschutz, M. P., Ross, S., Bhatt, S., Baron, T., Forcehimes, A. A., Laska, E., Mennenga, S. E., O'Donnell, K., Owens, L. T., Podrebarac, S., Rotrosen, J., Tonigan, J. S., & Worth, L. (2022). Percentage of Heavy Drinking Days Following Psilocybin-Assisted Psychotherapy vs Placebo in the Treatment of Adult Patients With Alcohol Use Disorder: A Randomized Clinical Trial. *JAMA Psychiatry*, *79*(10), 953–962. https://doi.org/10.1001/jamapsychiatry.2022.2096

Bohn, L. M., & Schmid, C. L. (2010). Serotonin receptor signaling and regulation via β-arrestins. *Critical Reviews in Biochemistry and Molecular Biology*, *45*(6), 555. https://doi.org/10.3109/10409238.2010.516741

Bosma, R., Stoddart, L. A., Georgi, V., Bouzo-Lorenzo, M., Bushby, N., Inkoom, L., Waring, M. J., Briddon, S. J., Vischer, H. F., Sheppard, R. J., Fernández-Montalván, A., Hill, S. J., & Leurs, R. (2019). Probe dependency in the determination of ligand binding kinetics at a prototypical G protein-coupled receptor. *Scientific Reports*, *9*(1), Article 1. https://doi.org/10.1038/s41598-019-44025-5

Braden, M. R., & Nichols, D. E. (2007). Assessment of the roles of serines 5.43(239) and 5.46(242) for binding and potency of agonist ligands at the human serotonin 5-HT2A receptor. *Molecular Pharmacology*, *72*(5), 1200–1209. https://doi.org/10.1124/mol.107.039255

Brandt, S. D., Freeman, S., McGagh, P., Abdul-Halim, N., & Alder, J. F. (2004). An analytical perspective on favoured synthetic routes to the psychoactive tryptamines. *Journal of Pharmaceutical and Biomedical Analysis*, *36*(4), 675–691. https://doi.org/10.1016/j.jpba.2004.08.022

Burt, J. B., Preller, K. H., Demirtas, M., Ji, J. L., Krystal, J. H., Vollenweider, F. X., Anticevic, A., & Murray, J. D. (2021). Transcriptomics-informed large-scale cortical model captures topography of pharmacological neuroimaging effects of LSD. *eLife*, *10*, e69320. https://doi.org/10.7554/eLife.69320

Butler, M., Jelen, L., & Rucker, J. (2022). Expectancy in placebo-controlled trials of psychedelics: If so, so what? *Psychopharmacology*, *239*(10), 3047–3055. https://doi.org/10.1007/s00213-022-06221-6

Canal, C. E., & Murnane, K. S. (2017). The serotonin 5-HT2C receptor and the non-addictive nature of classic hallucinogens. *Journal of Psychopharmacology*, *31*(1), 127–143. https://doi.org/10.1177/0269881116677104

Carbonaro, T. M., Johnson, M. W., Hurwitz, E., & Griffiths, R. R. (2018). Double-blind comparison of the two hallucinogens psilocybin and dextromethorphan: Similarities and differences in subjective experiences. *Psychopharmacology*, *235*(2), 521–534. https://doi.org/10.1007/s00213-017-4769-4

Carhart-Harris, R. L. (2018). The entropic brain—Revisited. *Neuropharmacology*, *142*, 167–178. https://doi.org/10.1016/j.neuropharm.2018.03.010

Carhart-Harris, R. L., Erritzoe, D., Williams, T., Stone, J. M., Reed, L. J., Colasanti, A., Tyacke, R. J., Leech, R., Malizia, A. L., Murphy, K., Hobden, P., Evans, J., Feilding, A., Wise, R. G., & Nutt, D. J. (2012). Neural correlates of the psychedelic state as determined by fMRI studies with psilocybin. *Proceedings of the National Academy of Sciences*, *109*(6), 2138–2143. https://doi.org/10.1073/pnas.1119598109

Carhart-Harris, R. L., Leech, R., Erritzoe, D., Williams, T. M., Stone, J. M., Evans, J., Sharp, D. J., Feilding, A., Wise, R. G., & Nutt, D. J. (2013). Functional connectivity measures after psilocybin inform a novel hypothesis of early psychosis. *Schizophrenia Bulletin*, *39*(6), 1343–1351. https://doi.org/10.1093/schbul/sbs117

Carhart-Harris, R. L., Muthukumaraswamy, S., Roseman, L., Kaelen, M., Droog, W., Murphy, K., Tagliazucchi, E., Schenberg, E. E., Nest, T., Orban, C., Leech, R., Williams, L. T., Williams, T. M., Bolstridge, M., Sessa, B., McGonigle, J., Sereno, M. I., Nichols, D., Hellyer, P. J., … Nutt, D. J. (2016). Neural correlates of the LSD experience revealed by multimodal neuroimaging. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(17), 4853–4858. https://doi.org/10.1073/pnas.1518377113

Carhart-Harris, R. L., Roseman, L., Bolstridge, M., Demetriou, L., Pannekoek, J. N., Wall, M. B., Tanner, M., Kaelen, M., McGonigle, J., Murphy, K., Leech, R., Curran, H. V., & Nutt, D. J. (2017). Psilocybin for treatment-resistant depression: fMRI-measured brain mechanisms. *Scientific Reports*, *7*(1), Article 1. https://doi.org/10.1038/s41598-017-13282-7

Carhart-Harris, R., Leech, R., Hellyer, P., Shanahan, M., Feilding, A., Tagliazucchi, E., Chialvo, D., & Nutt, D. (2014). The entropic brain: A theory of conscious states informed by neuroimaging research with psychedelic drugs. *Frontiers in Human Neuroscience*, *8*. https://www.frontiersin.org/article/10.3389/fnhum.2014.00020

Colquhoun, D. (1998). Binding, gating, affinity and efficacy: The interpretation of structure-activity relationships for agonists and of the effects of mutating receptors. *British Journal of Pharmacology*, *125*(5), 923–947. https://doi.org/10.1038/sj.bjp.0702164

Coyle, J. R., Presti, D. E., & Baggott, M. J. (2012). *Quantitative Analysis of Narrative Reports of Psychedelic Drugs* (arXiv:1206.0312). arXiv. https://doi.org/10.48550/arXiv.1206.0312

Cruzat, J., Perl, Y. S., Escrichs, A., Vohryzek, J., Timmermann, C., Roseman, L., Luppi, A. I., Ibañez, A., Nutt, D., Carhart-Harris, R., Tagliazucchi, E., Deco, G., & Kringelbach, M. L. (2022). Effects of classic psychedelic drugs on turbulent signatures in brain dynamics. *Network Neuroscience*, *6*(4), 1104–1124. https://doi.org/10.1162/netn_a_00250

Dai, R., Larkin, T. E., Huang, Z., Tarnal, V., Picton, P., Vlisides, P. E., Janke, E., McKinney, A., Hudetz, A. G., Harris, R. E., & Mashour, G. A. (2023). Classical and non-classical psychedelic drugs induce common network changes in human cortex. *NeuroImage*, *273*, 120097. https://doi.org/10.1016/j.neuroimage.2023.120097

Davey, C. G., & Harrison, B. J. (2018). The brain's center of gravity: How the default mode network helps us to understand the self. *World Psychiatry*, *17*(3), 278–279. https://doi.org/10.1002/wps.20553

Davey, C. G., Pujol, J., & Harrison, B. J. (2016). Mapping the self in the brain's default mode network. *NeuroImage*, *132*, 390–397. https://doi.org/10.1016/j.neuroimage.2016.02.022

Davis, A. K., Barrett, F. S., So, S., Gukasyan, N., Swift, T. C., & Griffiths, R. R. (2021). Development of the Psychological Insight Questionnaire among a sample of people who have consumed psilocybin or LSD. *Journal of Psychopharmacology (Oxford, England)*, *35*(4), 437–446. https://doi.org/10.1177/0269881120967878

Daws, R. E., Timmermann, C., Giribaldi, B., Sexton, J. D., Wall, M. B., Erritzoe, D., Roseman, L., Nutt, D., & Carhart-Harris, R. (2022). Increased global integration in the brain after psilocybin therapy for depression. *Nature Medicine*, *28*(4), Article 4. https://doi.org/10.1038/s41591-022-01744-z

de Almeida, J., & Mengod, G. (2007). Quantitative analysis of glutamatergic and GABAergic neurons expressing 5-HT(2A) receptors in human and monkey prefrontal cortex. *Journal of Neurochemistry*, *103*(2), 475–486. https://doi.org/10.1111/j.1471-4159.2007.04768.x

De Lean, A., Stadel, J. M., & Lefkowitz, R. J. (1980). A ternary complex model explains the agonist-specific binding properties of the adenylate cyclase-coupled beta-adrenergic receptor. *The Journal of Biological Chemistry*, *255*(15), 7108–7117.

de Oliveira, P. G., Ramos, M. L. S., Amaro, A. J., Dias, R. A., & Vieira, S. I. (2019). Gi/o-Protein Coupled Receptors in the Aging Brain. *Frontiers in Aging Neuroscience*, *11*. https://www.frontiersin.org/articles/10.3389/fnagi.2019.00089

de Vos, C. M. H., Mason, N. L., & Kuypers, K. P. C. (2021). Psychedelics and Neuroplasticity: A Systematic Review Unraveling the Biological Underpinnings of Psychedelics. *Frontiers in Psychiatry*, *12*. https://www.frontiersin.org/article/10.3389/fpsyt.2021.724606

Deco, G., Cruzat, J., Cabral, J., Knudsen, G. M., Carhart-Harris, R. L., Whybrow, P. C., Logothetis, N. K., & Kringelbach, M. L. (2018). Whole-Brain Multimodal Neuroimaging Model Using Serotonin Receptor Maps Explains Non-linear Functional Effects of LSD. *Current Biology*, *28*(19), 3065-3074.e6. https://doi.org/10.1016/j.cub.2018.07.083

Deco, G., Cruzat, J., Cabral, J., Tagliazucchi, E., Laufs, H., Logothetis, N. K., & Kringelbach, M. L. (2019). Awakening: Predicting external stimulation to force transitions between different brain states. *Proceedings of the National Academy of Sciences*, *116*(36), 18088–18097. https://doi.org/10.1073/pnas.1905534116

Deco, G., Kringelbach, M. L., Jirsa, V. K., & Ritter, P. (2017). The dynamics of resting fluctuations in the brain: Metastability and its dynamical cortical core. *Scientific Reports*, *7*(1), 3095. https://doi.org/10.1038/s41598-017-03073-5

Deco, G., Ponce-Alvarez, A., Hagmann, P., Romani, G. L., Mantini, D., & Corbetta, M. (2014). How Local Excitation–Inhibition Ratio Impacts the Whole Brain Dynamics. *Journal of Neuroscience*, *34*(23), 7886–7898. https://doi.org/10.1523/JNEUROSCI.5068-13.2014

Doss, M. K., May, D. G., Johnson, M. W., Clifton, J. M., Hedrick, S. L., Prisinzano, T. E., Griffiths, R. R., & Barrett, F. S. (2020). The Acute Effects of the Atypical Dissociative Hallucinogen Salvinorin A on

Functional Connectivity in the Human Brain. *Scientific Reports*, *10*(1), Article 1. https://doi.org/10.1038/s41598-020-73216-8

Doss, M. K., Považan, M., Rosenberg, M. D., Sepeda, N. D., Davis, A. K., Finan, P. H., Smith, G. S., Pekar, J. J., Barker, P. B., Griffiths, R. R., & Barrett, F. S. (2021). Psilocybin therapy increases cognitive and neural flexibility in patients with major depressive disorder. *Translational Psychiatry*, *11*(1), Article 1. https://doi.org/10.1038/s41398-021-01706-y

Duerler, P., Brem, S., Fraga-González, G., Neef, T., Allen, M., Zeidman, P., Stämpfli, P., Vollenweider, F. X., & Preller, K. H. (2021). Psilocybin Induces Aberrant Prediction Error Processing of Tactile Mismatch Responses-A Simultaneous EEG-FMRI Study. *Cerebral Cortex (New York, N.Y.: 1991)*, *32*(1), 186–196. https://doi.org/10.1093/cercor/bhab202

Dyer, D. C., & Gant, D. W. (1973). Vasoconstriction Produced by Hallucinogens on Isolated Human and Sheep Umbilical Vasculature. *Journal of Pharmacology and Experimental Therapeutics*, *184*(2), 366–375.

Eshleman, A. J., Forster, M. J., Wolfrum, K. M., Johnson, R. A., Janowsky, A., & Gatch, M. B. (2014). Behavioral and neurochemical pharmacology of six psychoactive substituted phenethylamines: Mouse locomotion, rat drug discrimination and in vitro receptor and transporter binding and function. *Psychopharmacology*, *231*(5), 875–888. https://doi.org/10.1007/s00213-013-3303-6

Fang, A., Baran, B., Beatty, C. C., Mosley, J., Feusner, J. D., Phan, K. L., Wilhelm, S., & Manoach, D. S. (2022). Maladaptive self-focused attention and default mode network connectivity: A transdiagnostic investigation across social anxiety and body dysmorphic disorders. *Social Cognitive and Affective Neuroscience*, *17*(7), 645–654. https://doi.org/10.1093/scan/nsab130

Fantegrossi, W. E., Murnane, K. S., & Reissig, C. J. (2008). The behavioral pharmacology of hallucinogens. *Biochemical Pharmacology*, *75*(1), 17–33. https://doi.org/10.1016/j.bcp.2007.07.018

Freyer, F., Roberts, J. A., Ritter, P., & Breakspear, M. (2012). A Canonical Model of Multistability and Scale-Invariance in Biological Systems. *PLOS Computational Biology*, *8*(8), e1002634. https://doi.org/10.1371/journal.pcbi.1002634

Fribourg, M., Moreno, J. L., Holloway, T., Provasi, D., Baki, L., Mahajan, R., Park, G., Adney, S. K., Hatcher, C., Eltit, J. M., Ruta, J. D., Albizu, L., Li, Z., Umali, A., Shim, J., Fabiato, A., MacKerell, A. D., Brezina, V., Sealfon, S. C., … Logothetis, D. E. (2011). Decoding the Signaling of a GPCR Heteromeric Complex Reveals a Unifying Mechanism of Action of Antipsychotic Drugs. *Cell*, *147*(5), 1011–1023. https://doi.org/10.1016/j.cell.2011.09.055

Garcia, A. C. M., & Maia, L. de O. (2022). The therapeutic potential of psychedelic substances in Hospice and Palliative Care. *Progress in Palliative Care*, *30*(1), 1–3. https://doi.org/10.1080/09699260.2022.2001140

Garcia, E. E., Smith, R. L., & Sanders-Bush, E. (2007). Role of G(q) protein in behavioral effects of the hallucinogenic drug 1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane. *Neuropharmacology*, *52*(8), 1671–1677. https://doi.org/10.1016/j.neuropharm.2007.03.013

Girn, M., Roseman, L., Bernhardt, B., Smallwood, J., Carhart-Harris, R., & Nathan Spreng, R. (2022). Serotonergic psychedelic drugs LSD and psilocybin reduce the hierarchical differentiation of unimodal and transmodal cortex. *NeuroImage*, *256*, 119220. https://doi.org/10.1016/j.neuroimage.2022.119220

Giulietti, M., Vivenzio, V., Piva, F., Principato, G., Bellantuono, C., & Nardi, B. (2014). How much do we know about the coupling of G-proteins to serotonin receptors? *Molecular Brain*, *7*(1), 49. https://doi.org/10.1186/s13041-014-0049-y

Glennon, R. A., Titeler, M., & McKenney, J. D. (1984). Evidence for 5-HT2 involvement in the mechanism of action of hallucinogenic agents. *Life Sciences*, *35*(25), 2505–2511. https://doi.org/10.1016/0024-3205(84)90436-3

González-Maeso, J., Weisstaub, N. V., Zhou, M., Chan, P., Ivic, L., Ang, R., Lira, A., Bradley-Moore, M., Ge, Y., Zhou, Q., Sealfon, S. C., & Gingrich, J. A. (2007). Hallucinogens recruit specific cortical 5-

HT(2A) receptor-mediated signaling pathways to affect behavior. *Neuron*, *53*(3), 439–452. https://doi.org/10.1016/j.neuron.2007.01.008

Goodwin, G. M., Croal, M., Marwood, L., & Malievskaia, E. (2023). Unblinding and demand characteristics in the treatment of depression. *Journal of Affective Disorders*, *328*, 1–5. https://doi.org/10.1016/j.jad.2023.02.030

Gouzoulis-Mayfrank, E., Schreckenberger, M., Sabri, O., Arning, C., Thelen, B., Spitzer, M., Kovar, K.-A., Hermle, L., Büll, U., & Sass, H. (1999). Neurometabolic Effects of Psilocybin, 3,4-Methylenedioxyethylamphetamine (MDE) and d-Methamphetamine in Healthy Volunteers A Double-Blind, Placebo-Controlled PET Study with [18F]FDG. *Neuropsychopharmacology*, *20*(6), Article 6. https://doi.org/10.1016/S0893-133X(98)00089-X

Griffiths, R. R., Hurwitz, E. S., Davis, A. K., Johnson, M. W., & Jesse, R. (2019). Survey of subjective 'God encounter experiences': Comparisons among naturally occurring experiences and those occasioned by the classic psychedelics psilocybin, LSD, ayahuasca, or DMT. *PLOS ONE*, *14*(4), e0214377. https://doi.org/10.1371/journal.pone.0214377

Griffiths, R. R., Johnson, M. W., Richards, W. A., Richards, B. D., Jesse, R., MacLean, K. A., Barrett, F. S., Cosimano, M. P., & Klinedinst, M. A. (2018). Psilocybin-occasioned mystical-type experience in combination with meditation and other spiritual practices produces enduring positive changes in psychological functioning and in trait measures of prosocial attitudes and behaviors. *Journal of Psychopharmacology (Oxford, England)*, *32*(1), 49–69. https://doi.org/10.1177/0269881117731279

Gurevich, E. V., Gainetdinov, R. R., & Gurevich, V. V. (2016). G protein-coupled receptor kinases as regulators of dopamine receptor functions. *Pharmacological Research*, *111*, 1–16. https://doi.org/10.1016/j.phrs.2016.05.010

Hamada, S., Senzaki, K., Hamaguchi-Hamada, K., Tabuchi, K., Yamamoto, H., Yamamoto, T., Yoshikawa, S., Okano, H., & Okado, N. (1998). Localization of 5-HT2A receptor in rat cerebral cortex and olfactory system revealed by immunohistochemistry using two antibodies raised in rabbit and chicken. *Brain Research. Molecular Brain Research*, *54*(2), 199–211. https://doi.org/10.1016/s0169-328x(97)00322-7

Hansen, J. Y., Shafiei, G., Markello, R. D., Smart, K., Cox, S. M. L., Nørgaard, M., Beliveau, V., Wu, Y., Gallezot, J.-D., Aumont, É., Servaes, S., Scala, S. G., DuBois, J. M., Wainstein, G., Bezgin, G., Funck, T., Schmitz, T. W., Spreng, R. N., Galovic, M., … Misic, B. (2022). Mapping neurotransmitter systems to the structural and functional organization of the human neocortex. *Nature Neuroscience*, *25*(11), Article 11. https://doi.org/10.1038/s41593-022-01186-3

Hasler, F., Grimberg, U., Benz, M. A., Huber, T., & Vollenweider, F. X. (2004). Acute psychological and physiological effects of psilocybin in healthy humans: A double-blind, placebo-controlled dose-effect study. *Psychopharmacology*, *172*(2), 145–156. https://doi.org/10.1007/s00213-003-1640-6

Haufe, S., DeGuzman, P., Henin, S., Arcaro, M., Honey, C. J., Hasson, U., & Parra, L. C. (2018). Elucidating relations between fMRI, ECoG, and EEG through a common natural stimulus. *NeuroImage*, *179*, 79–91. https://doi.org/10.1016/j.neuroimage.2018.06.016

Hendricks, P. S. (2018). Awe: A putative mechanism underlying the effects of classic psychedelic-assisted psychotherapy. *International Review of Psychiatry*, *30*(4), 331–342. https://doi.org/10.1080/09540261.2018.1474185

Hermle, L., Fünfgeld, M., Oepen, G., Botsch, H., Borchardt, D., Gouzoulis, E., Fehrenbach, R. A., & Spitzer, M. (1992). Mescaline-induced psychopathological, neuropsychological, and neurometabolic effects in normal subjects: Experimental psychosis as a tool for psychiatric research. *Biological Psychiatry*, *32*(11), 976–991. https://doi.org/10.1016/0006-3223(92)90059-9

Hillman, E. M. C. (2014). Coupling Mechanism and Significance of the BOLD Signal: A Status Report. *Annual Review of Neuroscience*, *37*, 161–181. https://doi.org/10.1146/annurev-neuro-071013-014111

Hoffmann, E. (2001). Effects of a psychedelic, tropical tea, ayahuasca, on the electroencephalographic (EEG) activity of the human brain during a shamanistic ritual. *MAPS Bulletin.*, *11*, 25–30.

Holze, F., Ley, L., Müller, F., Becker, A. M., Straumann, I., Vizeli, P., Kuehne, S. S., Roder, M. A., Duthaler, U., Kolaczynska, K. E., Varghese, N., Eckert, A., & Liechti, M. E. (2022). Direct comparison of the acute effects of lysergic acid diethylamide and psilocybin in a double-blind placebo-controlled study in healthy subjects. *Neuropsychopharmacology*, *47*(6), 1180–1187. https://doi.org/10.1038/s41386-022-01297-2

Honey, C. J., Kötter, R., Breakspear, M., & Sporns, O. (2007). Network structure of cerebral cortex shapes functional connectivity on multiple time scales. *Proceedings of the National Academy of Sciences*, *104*(24), 10240–10245. https://doi.org/10.1073/pnas.0701519104

Hulme, E. C., & Trevethick, M. A. (2010). Ligand binding assays at equilibrium: Validation and interpretation. *British Journal of Pharmacology*, *161*(6), 1219–1237. https://doi.org/10.1111/j.1476-5381.2009.00604.x

Jakab, R. L., & Goldman-Rakic, P. S. (1998). 5-Hydroxytryptamine2A serotonin receptors in the primate cerebral cortex: Possible site of action of hallucinogenic and antipsychotic drugs in pyramidal cell apical dendrites. *Proceedings of the National Academy of Sciences of the United States of America*, *95*(2), 735–740. https://doi.org/10.1073/pnas.95.2.735

Janowsky, A., Eshleman, A. J., Johnson, R. A., Wolfrum, K. M., Hinrichs, D. J., Yang, J., Zabriskie, T. M., Smilkstein, M. J., & Riscoe, M. K. (2014). Mefloquine and Psychotomimetics Share Neurotransmitter Receptor and Transporter Interactions In Vitro. *Psychopharmacology*, *231*(14), 2771–2783. https://doi.org/10.1007/s00213-014-3446-0

Kenakin, T. (2017). A Scale of Agonism and Allosteric Modulation for Assessment of Selectivity, Bias, and Receptor Mutation. *Molecular Pharmacology*, *92*(4), 414–424. https://doi.org/10.1124/mol.117.108787

Kenakin, T., Watson, C., Muniz-Medina, V., Christopoulos, A., & Novick, S. (2011). A Simple Method for Quantifying Functional Selectivity and Agonist Bias. *ACS Chemical Neuroscience*, *3*(3), 193–203. https://doi.org/10.1021/cn200111m

Kenakin, T., Watson, C., Muniz-Medina, V., Christopoulos, A., & Novick, S. (2012). A Simple Method for Quantifying Functional Selectivity and Agonist Bias. *ACS Chemical Neuroscience*, *3*(3), 193–203. https://doi.org/10.1021/cn200111m

Knudsen, G. M. (2023). Sustained effects of single doses of classical psychedelics in humans. *Neuropsychopharmacology*, *48*(1), 145–150. https://doi.org/10.1038/s41386-022-01361-x

Kozell, L. B., Eshleman, A. J., Swanson, T. L., Bloom, S. H., Wolfrum, K. M., Schmachtenberg, J. L., Olson, R. J., Janowsky, A., & Abbas, A. I. (2023). Pharmacologic Activity of Substituted Tryptamines at 5-Hydroxytryptamine (5-HT)2A Receptor (5-HT2AR), 5-HT2CR, 5-HT1AR, and Serotonin Transporter. *The Journal of Pharmacology and Experimental Therapeutics*, *385*(1), 62–75. https://doi.org/10.1124/jpet.122.001454

Kringelbach, M. L., Cruzat, J., Cabral, J., Knudsen, G. M., Carhart-Harris, R., Whybrow, P. C., Logothetis, N. K., & Deco, G. (2020). Dynamic coupling of whole-brain neuronal and neurotransmitter systems. *Proceedings of the National Academy of Sciences*, *117*(17), 9566–9576. https://doi.org/10.1073/pnas.1921475117

Kringelbach, M. L., & Deco, G. (2020). Brain States and Transitions: Insights from Computational Neuroscience. *Cell Reports*, *32*(10), 108128. https://doi.org/10.1016/j.celrep.2020.108128

Kumral, D., Şansal, F., Cesnaite, E., Mahjoory, K., Al, E., Gaebler, M., Nikulin, V. V., & Villringer, A. (2020). BOLD and EEG signal variability at rest differently relate to aging in the human brain. *NeuroImage*, *207*, 116373. https://doi.org/10.1016/j.neuroimage.2019.116373

Lawn, T., Dipasquale, O., Vamvakas, A., Tsougos, I., Mehta, M. A., & Howard, M. A. (2022). Differential contributions of serotonergic and dopaminergic functional connectivity to the phenomenology of LSD. *Psychopharmacology*, *239*(6), 1797–1808. https://doi.org/10.1007/s00213-022-06117-5

Lebedev, A. V., Lövdén, M., Rosenthal, G., Feilding, A., Nutt, D. J., & Carhart-Harris, R. L. (2015). Finding the self by losing the self: Neural correlates of ego-dissolution under psilocybin. *Human Brain Mapping*, *36*(8), 3137–3153. https://doi.org/10.1002/hbm.22833

Lee, M., Sehatpour, P., Hoptman, M. J., Lakatos, P., Dias, E. C., Kantrowitz, J. T., Martinez, A. M., & Javitt, D. C. (2017). Neural mechanisms of mismatch negativity (MMN) dysfunction in schizophrenia. *Molecular Psychiatry*, *22*(11), 1585–1593. https://doi.org/10.1038/mp.2017.3

Ley, L., Holze, F., Arikci, D., Becker, A. M., Straumann, I., Klaiber, A., Coviello, F., Dierbach, S., Thomann, J., Duthaler, U., Luethi, D., Varghese, N., Eckert, A., & Liechti, M. E. (2023). Comparative acute effects of mescaline, lysergic acid diethylamide, and psilocybin in a randomized, double-blind, placebo-controlled cross-over study in healthy participants. *Neuropsychopharmacology*, 1–9. https://doi.org/10.1038/s41386-023-01607-2

López-Giménez, J. F., & González-Maeso, J. (2018). Hallucinogens and Serotonin 5-HT2A Receptor-Mediated Signaling Pathways. *Current Topics in Behavioral Neurosciences*, *36*, 45–73. https://doi.org/10.1007/7854_2017_478

Luan, L., Eckernäs, E., Ashton, M., Rosas, F., Uthaug, M., Bartha, A., Jagger, S., Gascon-Perai, K., Gomes, L., Nutt, D., Erritzoe, D., Carhart-Harris, R., & Timmermann, C. (2023). *Psychological and physiological effects of extended DMT*. PsyArXiv. https://doi.org/10.31234/osf.io/vg4dp

Luppi, A. I., Carhart-Harris, R. L., Roseman, L., Pappas, I., Menon, D. K., & Stamatakis, E. A. (2021). LSD alters dynamic integration and segregation in the human brain. *NeuroImage*, *227*, 117653. https://doi.org/10.1016/j.neuroimage.2020.117653

Maclean, K. A., Leoutsakos, J.-M. S., Johnson, M. W., & Griffiths, R. R. (2012). Factor Analysis of the Mystical Experience Questionnaire: A Study of Experiences Occasioned by the Hallucinogen Psilocybin. *Journal for the Scientific Study of Religion*, *51*(4), 721–737. https://doi.org/10.1111/j.1468-5906.2012.01685.x

Madsen, M. K., Stenbæk, D. S., Arvidsson, A., Armand, S., Marstrand-Joergensen, M. R., Johansen, S. S., Linnet, K., Ozenne, B., Knudsen, G. M., & Fisher, P. M. (2021). Psilocybin-induced changes in brain network integrity and segregation correlate with plasma psilocin level and psychedelic experience. *European Neuropsychopharmacology*, *50*, 121–132. https://doi.org/10.1016/j.euroneuro.2021.06.001

Makowski, C., Lepage, M., & Evans, A. C. (2019). Head motion: The dirty little secret of neuroimaging in psychiatry. *Journal of Psychiatry & Neuroscience : JPN*, *44*(1), 62–68. https://doi.org/10.1503/jpn.180022

Mason, N. L., Kuypers, K. P. C., Müller, F., Reckweg, J., Tse, D. H. Y., Toennes, S. W., Hutten, N. R. P. W., Jansen, J. F. A., Stiers, P., Feilding, A., & Ramaekers, J. G. (2020). Me, myself, bye: Regional alterations in glutamate and the experience of ego dissolution with psilocybin. *Neuropsychopharmacology*, *45*(12), Article 12. https://doi.org/10.1038/s41386-020-0718-8

McCulloch, D. E.-W., Olsen, A. S., Ozenne, B., Stenbæk, D. S., Armand, S., Madsen, M. K., Knudsen, G. M., & Fisher, P. M. (2023). *Navigating the chaos of psychedelic neuroimaging: A multi-metric evaluation of acute psilocybin effects on brain entropy* (p. 2023.07.03.23292164). medRxiv. https://doi.org/10.1101/2023.07.03.23292164

Mertens, L. J., Wall, M. B., Roseman, L., Demetriou, L., Nutt, D. J., & Carhart-Harris, R. L. (2020). Therapeutic mechanisms of psilocybin: Changes in amygdala and prefrontal functional connectivity during emotional processing after psilocybin for treatment-resistant depression. *Journal of Psychopharmacology (Oxford, England)*, *34*(2), 167–180. https://doi.org/10.1177/0269881119895520

Molnar-Szakacs, I., & Uddin, L. (2013). Self-Processing and the Default Mode Network: Interactions with the Mirror Neuron System. *Frontiers in Human Neuroscience*, *7*. https://www.frontiersin.org/article/10.3389/fnhum.2013.00571

Moran, J. M., Kelley, W. M., & Heatherton, T. F. (2013). What Can the Organization of the Brain's Default Mode Network Tell us About Self-Knowledge? *Frontiers in Human Neuroscience*, *7*, 391. https://doi.org/10.3389/fnhum.2013.00391

Moreton, S. G., Szalla, L., Menzies, R. E., & Arena, A. F. (2020). Embedding existential psychology within psychedelic science: Reduced death anxiety as a mediator of the therapeutic effects of psychedelics. *Psychopharmacology*, *237*(1), 21–32. https://doi.org/10.1007/s00213-019-05391-0

Moujaes, F., Preller, K. H., Ji, J. L., Murray, J. D., Berkovitch, L., Vollenweider, F. X., & Anticevic, A. (2023). Toward Mapping Neurobehavioral Heterogeneity of Psychedelic Neurobiology in Humans. *Biological Psychiatry*, *93*(12), 1061–1070. https://doi.org/10.1016/j.biopsych.2022.10.021

Müller, F., Dolder, P. C., Schmidt, A., Liechti, M. E., & Borgwardt, S. (2018). Altered network hub connectivity after acute LSD administration. *NeuroImage. Clinical*, *18*, 694–701. https://doi.org/10.1016/j.nicl.2018.03.005

Müller, F., Liechti, M. E., Lang, U. E., & Borgwardt, S. (2018). Chapter 6 - Advances and challenges in neuroimaging studies on the effects of serotonergic hallucinogens: Contributions of the resting brain. In T. Calvey (Ed.), *Progress in Brain Research* (Vol. 242, pp. 159–177). Elsevier. https://doi.org/10.1016/bs.pbr.2018.08.004

Muthukumaraswamy, S. D., Carhart-Harris, R. L., Moran, R. J., Brookes, M. J., Williams, T. M., Errtizoe, D., Sessa, B., Papadopoulos, A., Bolstridge, M., Singh, K. D., Feilding, A., Friston, K. J., & Nutt, D. J. (2013). Broadband Cortical Desynchronization Underlies the Human Psychedelic State. *Journal of Neuroscience*, *33*(38), 15171–15183. https://doi.org/10.1523/JNEUROSCI.2063-13.2013

Muttoni, S., Ardissino, M., & John, C. (2019). Classical psychedelics for the treatment of depression and anxiety: A systematic review. *Journal of Affective Disorders*, *258*, 11–24. https://doi.org/10.1016/j.jad.2019.07.076

Nichols, D. E. (2004). Hallucinogens. *Pharmacology & Therapeutics*, *101*(2), 131–181. https://doi.org/10.1016/j.pharmthera.2003.11.002

Nichols, D. E. (2016). Psychedelics. *Pharmacological Reviews*, *68*(2), 264–355. https://doi.org/10.1124/pr.115.011478

Nichols, D. E. (2018). Chemistry and Structure–Activity Relationships of Psychedelics. In A. L. Halberstadt, F. X. Vollenweider, & D. E. Nichols (Eds.), *Behavioral Neurobiology of Psychedelic Drugs* (pp. 1–43). Springer. https://doi.org/10.1007/7854_2017_475

Nour, M. M., Evans, L., Nutt, D., & Carhart-Harris, R. L. (2016). Ego-Dissolution and Psychedelics: Validation of the Ego-Dissolution Inventory (EDI). *Frontiers in Human Neuroscience*, *10*. https://www.frontiersin.org/article/10.3389/fnhum.2016.00269

Nutt, D., Erritzoe, D., & Carhart-Harris, R. (2020). Psychedelic Psychiatry's Brave New World. *Cell*, *181*(1), 24–28. https://doi.org/10.1016/j.cell.2020.03.020

Ort, A., Smallridge, J. W., Sarasso, S., Casarotto, S., von Rotz, R., Casanova, A., Seifritz, E., Preller, K. H., Tononi, G., & Vollenweider, F. X. (2023). TMS-EEG and resting-state EEG applied to altered states of consciousness: Oscillations, complexity, and phenomenology. *iScience*, *26*(5), 106589. https://doi.org/10.1016/j.isci.2023.106589

Özbay, P. S., Chang, C., Picchioni, D., Mandelkow, H., Chappel-Farley, M. G., van Gelderen, P., de Zwart, J. A., & Duyn, J. (2019). Sympathetic activity contributes to the fMRI signal. *Communications Biology*, *2*(1), Article 1. https://doi.org/10.1038/s42003-019-0659-0

Özbay, P. S., Chang, C., Picchioni, D., Mandelkow, H., Moehlman, T. M., Chappel-Farley, M. G., van Gelderen, P., de Zwart, J. A., & Duyn, J. H. (2018). Contribution of systemic vascular effects to fMRI activity in white matter. *NeuroImage*, *176*, 541–549. https://doi.org/10.1016/j.neuroimage.2018.04.045

Pahnke, W. N. (1969). The Psychedelic Mystical Experience in the Human Encounter with Death*. *Harvard Theological Review*, *62*(1), 1–21. https://doi.org/10.1017/S0017816000027577

Palhano-Fontes, F., Andrade, K. C., Tofoli, L. F., Santos, A. C., Crippa, J. A. S., Hallak, J. E. C., Ribeiro, S., & Araujo, D. B. de. (2015). The Psychedelic State Induced by Ayahuasca Modulates the Activity and Connectivity of the Default Mode Network. *PLOS ONE*, *10*(2), e0118143. https://doi.org/10.1371/journal.pone.0118143

Pasquini, L., Palhano-Fontes, F., & Araujo, D. B. (2020). Subacute effects of the psychedelic ayahuasca on the salience and default mode networks. *Journal of Psychopharmacology*, *34*(6), 623–635. https://doi.org/10.1177/0269881120909409

Peill, J. M., Trinci, K. E., Kettner, H., Mertens, L. J., Roseman, L., Timmermann, C., Rosas, F. E., Lyons, T., & Carhart-Harris, R. L. (2022). Validation of the Psychological Insight Scale: A new scale to assess psychological insight following a psychedelic experience. *Journal of Psychopharmacology (Oxford, England)*, *36*(1), 31–45. https://doi.org/10.1177/02698811211066709

Pierce, P. A., & Peroutka, S. J. (1989). Hallucinogenic drug interactions with neurotransmitter receptor binding sites in human cortex. *Psychopharmacology*, *97*(1), 118–122. https://doi.org/10.1007/BF00443425

Pokorny, T., Preller, K. H., Kraehenmann, R., & Vollenweider, F. X. (2016). Modulatory effect of the 5-HT1A agonist buspirone and the mixed non-hallucinogenic 5-HT1A/2A agonist ergotamine on psilocybin-induced psychedelic experience. *European Neuropsychopharmacology*, *26*(4), 756–766. https://doi.org/10.1016/j.euroneuro.2016.01.005

Pottie, E., Cannaert, A., & Stove, C. P. (2020). In vitro structure-activity relationship determination of 30 psychedelic new psychoactive substances by means of β-arrestin 2 recruitment to the serotonin 2A receptor. *Archives of Toxicology*, *94*(10), 3449–3460. https://doi.org/10.1007/s00204-020-02836-w

Pottie, E., Dedecker, P., & Stove, C. P. (2020). Identification of psychedelic new psychoactive substances (NPS) showing biased agonism at the 5-HT2AR through simultaneous use of β-arrestin 2 and miniGαq bioassays. *Biochemical Pharmacology*, *182*, 114251. https://doi.org/10.1016/j.bcp.2020.114251

Preller, K. H., Herdener, M., Pokorny, T., Planzer, A., Kraehenmann, R., Stämpfli, P., Liechti, M. E., Seifritz, E., & Vollenweider, F. X. (2017). The Fabric of Meaning and Subjective Effects in LSD-Induced States Depend on Serotonin 2A Receptor Activation. *Current Biology: CB*, *27*(3), 451–457. https://doi.org/10.1016/j.cub.2016.12.030

Puig, M. V., Celada, P., Díaz-Mataix, L., & Artigas, F. (2003). In Vivo Modulation of the Activity of Pyramidal Neurons in the Rat Medial Prefrontal Cortex by 5-HT2A Receptors: Relationship to Thalamocortical Afferents. *Cerebral Cortex*, *13*(8), 870–882. https://doi.org/10.1093/cercor/13.8.870

Qiu, T. (Tim), & Minda, J. P. (2021). *Recreational Psychedelic Users Frequently Encounter Complete Mystical Experiences: Trip Content and Implications for Wellbeing*. PsyArXiv. https://doi.org/10.31234/osf.io/xrbzs

Quednow, B. B., Kometer, M., Geyer, M. A., & Vollenweider, F. X. (2012). Psilocybin-Induced Deficits in Automatic and Controlled Inhibition are Attenuated by Ketanserin in Healthy Human Volunteers. *Neuropsychopharmacology*, *37*(3), Article 3. https://doi.org/10.1038/npp.2011.228

Riba, J., Anderer, P., Jané, F., Saletu, B., & Barbanoj, M. J. (2004). Effects of the South American Psychoactive Beverage Ayahuasca on Regional Brain Electrical Activity in Humans: A Functional Neuroimaging Study Using Low-Resolution Electromagnetic Tomography. *Neuropsychobiology*, *50*(1), 89–101. https://doi.org/10.1159/000077946

Riba, J., Anderer, P., Morte, A., Urbano, G., Jané, F., Saletu, B., & Barbanoj, M. J. (2002). Topographic pharmaco-EEG mapping of the effects of the South American psychoactive beverage ayahuasca in healthy volunteers. *British Journal of Clinical Pharmacology*, *53*(6), 613–628. https://doi.org/10.1046/j.1365-2125.2002.01609.x

Riba, J., Rodríguez-Fornells, A., & Barbanoj, M. J. (2002). Effects of ayahuasca on sensory and sensorimotor gating in humans as measured by P50 suppression and prepulse inhibition of the startle reflex, respectively. *Psychopharmacology*, *165*(1), 18–28. https://doi.org/10.1007/s00213-002-1237-5

Rickli, A., Moning, O. D., Hoener, M. C., & Liechti, M. E. (2016). Receptor interaction profiles of novel psychoactive tryptamines compared with classic hallucinogens. *European Neuropsychopharmacology: The Journal of the European College of Neuropsychopharmacology*, *26*(8), 1327–1337. https://doi.org/10.1016/j.euroneuro.2016.05.001

Rodriguiz, R. M., Nadkarni, V., Means, C. R., Pogorelov, V. M., Chiu, Y.-T., Roth, B. L., & Wetsel, W. C. (2021). LSD-stimulated behaviors in mice require β-arrestin 2 but not β-arrestin 1. *Scientific Reports*, *11*(1), Article 1. https://doi.org/10.1038/s41598-021-96736-3

Roseman, L., Demetriou, L., Wall, M. B., Nutt, D. J., & Carhart-Harris, R. L. (2018). Increased amygdala responses to emotional faces after psilocybin for treatment-resistant depression. *Neuropharmacology*, *142*, 263–269. https://doi.org/10.1016/j.neuropharm.2017.12.041

Roseman, L., Haijen, E., Idialu-Ikato, K., Kaelen, M., Watts, R., & Carhart-Harris, R. (2019). Emotional breakthrough and psychedelics: Validation of the Emotional Breakthrough Inventory. *Journal of Psychopharmacology (Oxford, England)*, *33*(9), 1076–1087. https://doi.org/10.1177/0269881119855974

Roseman, L., Leech, R., Feilding, A., Nutt, D. J., & Carhart-Harris, R. L. (2014). The effects of psilocybin and MDMA on between-network resting state functional connectivity in healthy volunteers. *Frontiers in Human Neuroscience*, *8*. https://www.frontiersin.org/article/10.3389/fnhum.2014.00204

Sanz, C., Zamberlan, F., Erowid, E., Erowid, F., & Tagliazucchi, E. (2018). The Experience Elicited by Hallucinogens Presents the Highest Similarity to Dreaming within a Large Database of Psychoactive Substance Reports. *Frontiers in Neuroscience*, *12*. https://doi.org/10.3389/fnins.2018.00007

Schmitz, G. P., Jain, M. K., Slocum, S. T., & Roth, B. L. (2022). 5-HT2A SNPs Alter the Pharmacological Signaling of Potentially Therapeutic Psychedelics. *ACS Chemical Neuroscience*, *13*(16), 2386–2398. https://doi.org/10.1021/acschemneuro.1c00815

Schrantee, A., Ferguson, B., Stoffers, D., Booij, J., Rombouts, S., & Reneman, L. (2016). Effects of dexamphetamine-induced dopamine release on resting-state network connectivity in recreational amphetamine users and healthy controls. *Brain Imaging and Behavior*, *10*(2), 548–558. https://doi.org/10.1007/s11682-015-9419-z

Shah, U., Pincas, H., Sealfon, S. C., & González-Maeso, J. (2020). Chapter 11—Structure and function of serotonin GPCR heteromers. In C. P. Müller & K. A. Cunningham (Eds.), *Handbook of Behavioral Neuroscience* (Vol. 31, pp. 217–238). Elsevier. https://doi.org/10.1016/B978-0-444-64125-0.00011-6

Smigielski, L., Scheidegger, M., Kometer, M., & Vollenweider, F. X. (2019). Psilocybin-assisted mindfulness training modulates self-consciousness and brain default mode network connectivity with lasting effects. *NeuroImage*, *196*, 207–215. https://doi.org/10.1016/j.neuroimage.2019.04.009

Spruston, N. (2008). Pyramidal neurons: Dendritic structure and synaptic integration. *Nature Reviews Neuroscience*, *9*(3), Article 3. https://doi.org/10.1038/nrn2286

Stoll, W. (1947). Lysergsaure-diathylamid, ein Phantastikum aus der Mutterkorngruppe. *Royal Society of Medicine, Microfilm Unit*.

Strange, P. G. (1999). G-protein coupled receptors: Conformations and states. *Biochemical Pharmacology*, *58*(7), 1081–1088. https://doi.org/10.1016/s0006-2952(99)00144-6

Strange, P. G. (2008). Agonist binding, agonist affinity and agonist efficacy at G protein-coupled receptors. *British Journal of Pharmacology*, *153*(7), 1353–1363. https://doi.org/10.1038/sj.bjp.0707672

Strassman, R. J., & Qualls, C. R. (1994). Dose-response study of N,N-dimethyltryptamine in humans. I. Neuroendocrine, autonomic, and cardiovascular effects. *Archives of General Psychiatry*, *51*(2), 85–97. https://doi.org/10.1001/archpsyc.1994.03950020009001

Strassman, R. J., Qualls, C. R., Uhlenhuth, E. H., & Kellner, R. (1994). Dose-response study of N,N-dimethyltryptamine in humans. II. Subjective effects and preliminary results of a new rating scale. *Archives of General Psychiatry*, *51*(2), 98–108. https://doi.org/10.1001/archpsyc.1994.03950020022002

Stuckey, D. E., Lawson, R., & Luna, L. E. (2005). EEG gamma coherence and other correlates of subjective reports during ayahuasca experiences. *Journal of Psychoactive Drugs*, *37*(2), 163–178. https://doi.org/10.1080/02791072.2005.10399798

Studerus, E., Gamma, A., & Vollenweider, F. X. (2010). Psychometric Evaluation of the Altered States of Consciousness Rating Scale (OAV). *PLOS ONE*, *5*(8), e12412. https://doi.org/10.1371/journal.pone.0012412

Sykes, D. A., Stoddart, L. A., Kilpatrick, L. E., & Hill, S. J. (2019). Binding kinetics of ligands acting at GPCRs. *Molecular and Cellular Endocrinology*, *485*, 9–19. https://doi.org/10.1016/j.mce.2019.01.018

Tagliazucchi, E. (2020). *Early and Contemporary Human Neuroimaging Studies of Serotonergic Psychedelics*. https://doi.org/10.20944/preprints202005.0510.v1

Tagliazucchi, E., Carhart-Harris, R., Leech, R., Nutt, D., & Chialvo, D. R. (2014). Enhanced repertoire of brain dynamical states during the psychedelic experience. *Human Brain Mapping*, *35*(11), 5442–5456. https://doi.org/10.1002/hbm.22562

Tagliazucchi, E., Roseman, L., Kaelen, M., Orban, C., Muthukumaraswamy, S. D., Murphy, K., Laufs, H., Leech, R., McGonigle, J., Crossley, N., Bullmore, E., Williams, T., Bolstridge, M., Feilding, A., Nutt, D. J., & Carhart-Harris, R. (2016). Increased Global Functional Connectivity Correlates with LSD-Induced Ego Dissolution. *Current Biology: CB*, *26*(8), 1043–1050. https://doi.org/10.1016/j.cub.2016.02.010

Thomas Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125–1165. https://doi.org/10.1152/jn.00338.2011

Timmermann, C., Roseman, L., Haridas, S., Rosas, F. E., Luan, L., Kettner, H., Martell, J., Erritzoe, D., Tagliazucchi, E., Pallavicini, C., Girn, M., Alamia, A., Leech, R., Nutt, D. J., & Carhart-Harris, R. L. (2023). Human brain effects of DMT assessed via EEG-fMRI. *Proceedings of the National Academy of Sciences of the United States of America*, *120*(13), e2218949120. https://doi.org/10.1073/pnas.2218949120

Timmermann, C., Spriggs, M. J., Kaelen, M., Leech, R., Nutt, D. J., Moran, R. J., Carhart-Harris, R. L., & Muthukumaraswamy, S. D. (2018). LSD modulates effective connectivity and neural adaptation mechanisms in an auditory oddball paradigm. *Neuropharmacology*, *142*, 251–262. https://doi.org/10.1016/j.neuropharm.2017.10.039

Tófoli, L. F., & de Araujo, D. B. (2016). Treating Addiction: Perspectives from EEG and Imaging Studies on Psychedelics. *International Review of Neurobiology*, *129*, 157–185. https://doi.org/10.1016/bs.irn.2016.06.005

Tupper, K. W., Wood, E., Yensen, R., & Johnson, M. W. (2015). Psychedelic medicine: A re-emerging therapeutic paradigm. *CMAJ: Canadian Medical Association Journal*, *187*(14), 1054–1059. https://doi.org/10.1503/cmaj.141124

Tuteja, N. (2009). Signaling through G protein coupled receptors. *Plant Signaling & Behavior*, *4*(10), 942–947.

Umbricht, D., Koller, R., Vollenweider, F. X., & Schmid, L. (2002). Mismatch negativity predicts psychotic experiences induced by NMDA receptor antagonist in healthy volunteers. *Biological Psychiatry*, *51*(5), 400–406. https://doi.org/10.1016/s0006-3223(01)01242-2

Umbricht, D., Vollenweider, F. X., Schmid, L., Grübel, C., Skrabo, A., Huber, T., & Koller, R. (2003). Effects of the 5-HT2A Agonist Psilocybin on Mismatch Negativity Generation and AX-Continuous Performance Task: Implications for the Neuropharmacology of Cognitive Deficits in Schizophrenia. *Neuropsychopharmacology*, *28*(1), Article 1. https://doi.org/10.1038/sj.npp.1300005

van Buuren, M., Gladwin, T. E., Zandbelt, B. B., Kahn, R. S., & Vink, M. (2010). Reduced functional coupling in the default-mode network during self-referential processing. *Human Brain Mapping*, *31*(8), 1117–1127. https://doi.org/10.1002/hbm.20920

Van Essen, D. C., Smith, S. M., Barch, D. M., Behrens, T. E. J., Yacoub, E., & Ugurbil, K. (2013). The WU-Minn Human Connectome Project: An overview. *NeuroImage*, *80*, 62–79. https://doi.org/10.1016/j.neuroimage.2013.05.041

Vogt, S. B., Ley, L., Erne, L., Straumann, I., Becker, A. M., Klaiber, A., Holze, F., Vandersmissen, A., Mueller, L., Duthaler, U., Rudin, D., Luethi, D., Varghese, N., Eckert, A., & Liechti, M. E. (2023). Acute effects of intravenous DMT in a randomized placebo-controlled study in healthy participants. *Translational Psychiatry*, *13*(1), 172. https://doi.org/10.1038/s41398-023-02477-4

Vollenweider, F. X., Csomor, P. A., Knappe, B., Geyer, M. A., & Quednow, B. B. (2007). The effects of the preferential 5-HT2A agonist psilocybin on prepulse inhibition of startle in healthy human volunteers depend on interstimulus interval. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *32*(9), 1876–1887. https://doi.org/10.1038/sj.npp.1301324

Vollenweider, F. X., Leenders, K. L., Scharfetter, C., Maguire, P., Stadelmann, O., & Angst, J. (1997). Positron emission tomography and fluorodeoxyglucose studies of metabolic hyperfrontality and psychopathology in the psilocybin model of psychosis. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *16*(5), 357–372. https://doi.org/10.1016/S0893-133X(96)00246-1

Vollenweider, F. X., Vollenweider-Scherpenhuyzen, M. F., Bäbler, A., Vogel, H., & Hell, D. (1998). Psilocybin induces schizophrenia-like psychosis in humans via a serotonin-2 agonist action. *Neuroreport*, *9*(17), 3897–3902. https://doi.org/10.1097/00001756-199812010-00024

Vollenweider, F. X., Vontobel, P., Hell, D., & Leenders, K. L. (1999). 5-HT Modulation of Dopamine Release in Basal Ganglia in Psilocybin-Induced Psychosis in Man—A PET Study with [11C]raclopride. *Neuropsychopharmacology*, *20*(5), Article 5. https://doi.org/10.1016/S0893-133X(98)00108-0

Vossel, S., Geng, J. J., & Fink, G. R. (2014). Dorsal and Ventral Attention Systems. *The Neuroscientist*, *20*(2), 150–159. https://doi.org/10.1177/1073858413494269

Wall, M. B., Lam, C., Ertl, N., Kaelen, M., Roseman, L., Nutt, D. J., & Carhart-Harris, R. L. (2022). *Increased low-frequency brain responses to music after psilocybin therapy for depression* (p. 2022.02.13.480302). bioRxiv. https://doi.org/10.1101/2022.02.13.480302

Wallach, J., Cao, A. B., Calkins, M. M., Heim, A. J., Lanham, J. K., Bonniwell, E. M., Hennessey, J. J., Bock, H. A., Anderson, E. I., Sherwood, A. M., Morris, H., Klein, R. de, Klein, A. K., Cuccurazzu, B., Gamrat, J., Fannana, T., Zauhar, R., Halberstadt, A. L., & McCorvy, J. D. (2023). *Identification of 5-HT2A Receptor Signaling Pathways Responsible for Psychedelic Potential* (p. 2023.07.29.551106). bioRxiv. https://doi.org/10.1101/2023.07.29.551106

Watts, V. J., Lawler, C. P., Fox, D. R., Neve, K. A., Nichols, D. E., & Mailman, R. B. (1995). LSD and structural analogs: Pharmacological evaluation at D1 dopamine receptors. *Psychopharmacology*, *118*(4), 401–409. https://doi.org/10.1007/BF02245940

Weber, A. M., Soreni, N., & Noseworthy, M. D. (2014). A preliminary study on the effects of acute ethanol ingestion on default mode network and temporal fractal properties of the brain. *Magma (New York, N.Y.)*, *27*(4), 291–301. https://doi.org/10.1007/s10334-013-0420-5

Wen, A., Singhal, N., Jones, B. D. M., Zeifman, R. J., Mehta, S., Shenasa, M. A., Blumberger, D. M., Daskalakis, Z. J., & Weissman, C. R. (2023). A Systematic Review of Study Design and Placebo Controls in Psychedelic Research. *Psychedelic Medicine*. https://doi.org/10.1089/psymed.2023.0028

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: A Network Visualization Tool for Human Brain Connectomics. *PLOS ONE*, *8*(7), e68910. https://doi.org/10.1371/journal.pone.0068910

Yaden, D. B., Johnson, M. W., Griffiths, R. R., Doss, M. K., Garcia-Romeu, A., Nayak, S., Gukasyan, N., Mathur, B. N., & Barrett, F. S. (2021). Psychedelics and consciousness: Distinctions, demarcations, and opportunities. *International Journal of Neuropsychopharmacology*, *24*(8), 615–623. https://doi.org/10.1093/ijnp/pyab026

Zamberlan, F., Sanz, C., Martínez Vivot, R., Pallavicini, C., Erowid, F., Erowid, E., & Tagliazucchi, E. (2018). The Varieties of the Psychedelic Experience: A Preliminary Study of the Association Between the Reported Subjective Effects and the Binding Affinity Profiles of Substituted Phenethylamines and Tryptamines. *Frontiers in Integrative Neuroscience*, *12*, 54. https://doi.org/10.3389/fnint.2018.00054

Zhang, R., & Volkow, N. D. (2019). Brain default-mode network dysfunction in addiction. *NeuroImage*, *200*, 313–331. https://doi.org/10.1016/j.neuroimage.2019.06.036

Zhou, H.-X., Chen, X., Shen, Y.-Q., Li, L., Chen, N.-X., Zhu, Z.-C., Castellanos, F. X., & Yan, C.-G. (2020). Rumination and the default mode network: Meta-analysis of brain imaging studies and implications for depression. *NeuroImage*, *206*, 116287. https://doi.org/10.1016/j.neuroimage.2019.116287

Figures

1262
1263

Figure 1. Meta-analysis of the 5-Dimensional Altered States of Consciousness (5D-ASC) data reveals few significant differences between psychedelics, but many more significant differences within psychedelics. *The 5D-ASC is one of the most common scales for assessing the subjective effects of psychedelics. It measures altered states of consciousness along five different dimensions: oceanic boundlessness (OB; a feeling of interconnectedness), anxious ego dissolution (AED), visionary restructuralisation (VR; the quality and intensity of visual hallucinations), auditory alterations (AA), and reduction of vigilance (RV). Our literature search identified 23 studies that reported 5D-ASC data. We performed a multilevel random-effects meta-analysis in order to account for the lack of statistical*

independence between measurements of different dimensions within the same group of participants. Within-study and between-study heterogeneity were estimated with the restricted maximum likelihood procedure. Using a Correlated and Hierarchical Effects model to account for within-study correlations in sampling error, we analysed the effect of dose and drug (DMT, LSD, or psilocybin) on pooled 5D-ASC scores. DMT was administered intravenously in phenomenological studies, whereas LSD and psilocybin were administered orally. Thus, DMT has different pharmacokinetics from LSD and psilocybin, which impacts the subjective experience, so we only compared significant between-drug differences for LSD and psilocybin. We only compared the two drugs for similar doses, assuming that 0.1 mg LSD = 20 mg psilocybin (Ley et al., 2023). (a-c, left column) Between-drug comparisons of pooled 5D-ASC scores for LSD and psilocybin. LSD almost always ranked higher than psilocybin, but differences only reached significance in the VR dimension for high and medium doses and in the OB dimension for medium dose. (a-c, right column) Within-drug comparisons for LSD and psilocybin. VR and OB received significantly higher scores than AED and AA, for both LSD and psilocybin. (d-e) Within-drug comparisons for DMT. At low doses, OB and VR ranked significantly higher than AED. Only one study measured ASC scores for the AA and RV dimensions on DMT, so we did not perform a meta-analysis on these dimensions for DMT. Numerical data for the above figure are given in **Table S1**. *We also conducted a meta-analysis of the 11-dimensional ASC scale, which is also widely used in the literature; the results are displayed in* **Figure S3** *and* **Table S2**. ** < 0.05, ** < 0.01, *** < 0.001.*

Figure 2. Phenomenology profiles of the psychedelics demonstrate broad similarity between the neural correlates of their subjective effects. *We identified 14 studies that measured correlations between ASC ratings and fMRI activity or pairwise connectivity, including three studies on ketamine. We determined the Yeo network that contained each brain region that exhibited a significant correlation, and then we averaged across the correlations associated with each Yeo network, resulting in the mean correlation between each Yeo network and each ASC scale. The phenomenology profiles were defined by a weighted combination of the mean correlations for each ASC scale, in which the weights were the pooled ASC ratings of high doses of each psychedelic for the corresponding scale. Because the pooled ASC ratings are similar across psychedelics, the phenomenology profiles are very alike one another as well. However, the lack of data on the neural correlates of the ASC ratings limits our confidence in the validity of these profiles.*

(a)

(b)

Figure 3. Meta-analysis of the functional connectivity (FC) data indicates that psychedelics increase between-network FC and decrease within-network FC. *To perform a meta-analysis on the FC data, we determined the Yeo networks that contained each pair of functionally connected regions of interest (ROIs) in the literature and then performed a weighted sum of the number of significant connections between Yeo networks. We included n = 12 studies in this meta-analysis, after excluding studies that did not measure pairwise FC and secondary analyses of identical datasets. (a) Aggregate FC matrix, in which each entry contains the overall connectivity between a pair of Yeo networks. All connections were deemed to be significant according to our method, which compared the results to an independent resting-state fMRI dataset from the Human Connectome Project. (b) A rendering of the FC results on the surface of the brain. Positive edges are yellow, and negative edges are red. Images produced through the BrainNet Viewer (Xia et al., 2013).*

Figure 4. Functional connectivity (FC) profiles show unique FC patterns for each psychedelic. *Out of the n = 12 studies that we examined in our quantitative FC meta-analysis, n = 3 were studies on ayahuasca/DMT (ayahuasca: n = 2; DMT: n = 1), n = 3 on LSD, and n = 6 on psilocybin. Each FC profile contains the total FC of each network, which was obtained by taking the normalised sum of the rows of the corresponding pooled FC matrices, including insignificant values. (The units of the profiles are arbitrary.) The psychedelics display distinct FC profiles; for instance, ayahuasca, DMT, and psilocybin increase total FC for the DMN more strongly than for other networks, whereas LSD elevates FC more for the SMN and DAN than for other networks.*

(a) Selectivity

(b) Relative activity at 5-HT$_{2A}$ receptor

Figure 5. Pharmacology meta-analysis reveals that there are no significant differences in selectivity between psychedelics and that LSD induces significantly higher relative activity at the inositol phosphate (IP) formation pathway. For both the selectivity and relative activity data, we created random-effects models that modeled between-study variance. (a) Selectivity is the ratio between the binding affinity (measured as K_i) for a receptor of interest and the affinity for a reference receptor. We measured selectivity for three different receptors – 5-HT$_{2A}$, 5-HT$_{2C}$, and D$_2$ – relative to 5-HT$_{1A}$. Our literature search identified n = 14 studies on selectivity, including n = 6 studies on DMT, n = 9 studies on LSD, and n = 5 studies on psilocin. We did not find any significant between-drug differences in selectivity for any of the three receptors. (b) Relative activity is $\Delta\log(E_{max}/EC_{50})$, where E_{max} is the maximal effect of the drug relative to a reference ligand and EC_{50} is the concentration needed to elicit 50% of the maximal effect. The log of the ratio is measured for both the drug and the reference ligand, and then the difference is taken (Kenakin, 2017). We found n = 18 studies on functional activity, including n = 6 studies on DMT, n = 13 studies on LSD, and n = 6 studies on psilocin. We measured relative activity at three different signaling pathways: calcium mobilisation, IP formation, and β-arrestin2 recruitment. IP formation was the only pathway that exhibited any significant between-drug differences; LSD elicited significantly higher activity than both DMT and psilocin. * < 0.05, ** < 0.01, *** < 0.001.

1344

1345
1346

Figure 6. Pharmacology profiles primarily reveal the distribution of 5-HT$_{2A}$ receptors in the Yeo
networks, as well as the (insignificantly) higher selectivity of psilocin for the 5-HT$_{2A}$ receptor
compared to DMT and LSD. Based on an available PET atlas of the 5-HT$_{2A}$ and D$_2$ receptors (Hansen et al.,
2022), we created the pharmacology profile of each psychedelic. The profiles show the expression of those two receptors in the
Yeo networks, weighted by the selectivity of the corresponding psychedelics for those receptors. The PET atlas reveals that the
5-HT$_{2A}$ receptor is expressed much more in the human brain than the D$_2$ receptor. Therefore, the profiles are dominated by 5-
HT$_{2A}$ receptor expression. Since psilocin has the most specificity for 5-HT$_{2A}$ (relative to 5-HT$_{1A}$), followed by LSD then
DMT, psilocin's pharmacology profile is the most extensive.

1355

1356

1357
1358
Figure 7. Our multi-level analysis of psychedelic effects highlights the nonlinear relationship
between their pharmacology, neuroimaging, and phenomenology. *The effects of psychedelics form a*
tripartite hierarchy, consisting of subjective experience (a, phenomenology), functional connectivity (b, neuroimaging), and their
selective affinity for receptors (c, pharmacology). Here, we show the neural correlates of each level of the hierarchy in the seven
Yeo networks. In brief, each profile was derived from (a) neural correlates of the subjective dimensions of the ASC scale, (b)
summing the aggregate FC between each network and other networks, and (c) the distribution of each receptor, weighted by
each psychedelic's selectivity for that receptor. Clearly, there is a very weak correspondence between the different levels of the
hierarchy, revealing the highly non-linear relationship between phenomenology, neuroimaging, and pharmacology. Note: we
analysed data on both DMT and ayahuasca in our FC meta-analysis, but only on DMT for the other meta-analyses. We
examined data on psilocin in our pharmacology meta-analysis and on psilocybin in both of the other meta-analyses.

(a) High Dose (LSD: ≥0.110 mg, Psilocybin: ≥22 mg)

(d) High Dose DMT (≥16 mg bolus + ≥1 mg/min)

(b) Medium Dose (LSD: 0.075-0.109 mg, Psilocybin: 15-21 mg)

(e) Low Dose DMT (10-15 mg bolus + 0.6-0.99 mg/min)

(c) Low Dose (LSD: 0.05-0.074 mg, Psilocybin: 8-14 mg)

(a)

	Visual	SMN	DAN	VAN	Limbic	FPN	DMN
Visual	-20.01	-6.404	5.047	6.609	10.62	12.51	11.02
SMN	-6.404	1.3	7.08	1.423	-1.589	11.96	13.55
DAN	5.047	7.08	-0.05905	0.437	2.653	2.34	8.235
VAN	6.609	1.423	0.437	-0.7903	-0.2201	3.564	13.4
Limbic	10.62	-1.589	2.653	-0.2201	-3.176	0.488	0.3069
FPN	12.51	11.96	2.34	3.564	0.488	0.529	6.747
DMN	11.02	13.55	8.235	13.4	0.3069	6.747	-0.3453

(b)

(a)

Selectivity

(b)

Relative activity at 5-HT$_{2A}$ receptor

DMT:
Pharmacology Profile

LSD:
Pharmacology Profile

Psilocin:
Pharmacology Profile