Chemotherapy activates inflammasomes to cause inflammation-associated bone loss

Chun Wang[1], Khushpreet Kaur[1], Canxin Xu[2], Yousef Abu-Amer[3,4], and Gabriel Mbalaviele[1*]

[1]Division of Bone and Mineral Diseases, Washington University School of Medicine, St. Louis,

MO 63110, USA; [2]Aclaris Therapeutics, Inc., St. Louis, MO 63108, USA; [3]Department of

Orthopaedic Surgery, Washington University School of Medicine, St. Louis, Missouri, USA,

[4]Shriners Hospital for Children, St. Louis, Missouri, USA

Running title: Chemotherapy causes inflammasome-mediated bone loss

*To whom correspondence should be addressed. Division of Bone and Mineral Diseases,

Washington University School of Medicine, 660 South Euclid Avenue, Campus Box 8301, St.

Louis, MO 63110, Phone: (314) 286-1114; Fax: (314) 454-5047; e-mail: gmbalaviele@wustl.edu

Abstract

Chemotherapy is a widely used treatment for a variety of solid and hematological malignancies. Despite its success in improving the survival rate of cancer patients, chemotherapy causes significant toxicity to multiple organs, including the skeleton, but the underlying mechanisms have yet to be elucidated. Using tumor-free mouse models, which are commonly used to assess direct off-target effects of anti-neoplastic therapies, we found that doxorubicin caused massive bone loss in wild-type mice, a phenotype associated with increased number of osteoclasts, leukopenia, elevated serum levels of danger-associated molecular patterns (DAMPs; e.g., cell-free DNA and ATP) and cytokines (e.g., IL-1β and IL-18). Accordingly, doxorubicin activated the absent in melanoma (AIM2) and NLR family pyrin domain containing 3 (NLRP3) inflammasomes in macrophages and neutrophils, causing inflammatory cell death pyroptosis and NETosis, which correlated with its leukopenic effects. Moreover, the effects of this chemotherapeutic agent on cytokine secretion, cell demise, and bone loss were attenuated to various extent in conditions of AIM2 and/or NLRP3 insufficiency. Thus, we found that inflammasomes are key players in bone loss caused by doxorubicin, a finding that may inspire the development of a tailored adjuvant therapy that preserves the quality of this tissue in patients treated with this class of drugs.

Introduction

The chemotherapeutic drug, doxorubicin, is widely used for the treatment of breast cancer, bladder cancer, lymphomas, and acute lymphocytic leukemia (Smith et al., 2010, Jang et al., 2023, Geffen and Man, 2002). Despite its success in improving the survival rate of cancer patients, doxorubicin causes serious adverse effects, including cardiomyopathy, bone marrow suppression, hair loss, and skeletal manifestations (Cardinale et al., 2015, Tacar et al., 2013, Coleman et al., 2013). Bone complications include osteoporosis, a metabolic disease that is characterized by decreased bone mass and deteriorated microarchitecture, and associated with increased risks for the development of late fractures, and morbidities in the elderly populations (Cauley et al., 2009, Cawthon et al., 2009, Cawthon et al., 2012). In fact, it was reported that 20-50% of geriatric patients (≥65 years) with a hip fracture die within one year of fracture (Coleman et al., 2013). Consistent with the dogma that bone resorption by osteoclasts (OCs) and bone formation by osteoblasts is uncoupled in osteoporotic patients, doxorubicin causes bone loss by promoting osteoclastogenesis while suppressing osteoblastogenesis (Yao et al., 2020, Chai et al., 2014, Rana et al., 2013, Zhou et al., 2020). Increased production of senescence-associated secretory phenotype (SASP), enhanced generation of reactive oxygen species (ROS), and dysregulated autophagy and mitochondrial metabolism are proposed mechanisms of doxorubicin-induced bone pathology (Yao et al., 2020, Park et al., 2022).

Doxorubicin intercalates into DNA thereby impeding the activity of DNA repair enzymes such as topoisomerase II and impairing DNA replication (Robson et al., 1987, Lawrence, 1988, Bonner and Lawrence, 1990, Abe et al., 2022, Tewey et al., 1984). Defective DNA repair ultimately culminates in genomic instability and cell demise, events that can provoke uncontrollable release of intracellular contents such as DNA and various danger-associated molecular patterns (DAMPs). DNA-enriched entities include neutrophil extracellular traps (NETs),

web-like structures in which DNA is decorated with peptides, some of which have anti-microbial

and inflammatory properties (Komada et al., 2018, Apel et al., 2021). NETs can also propagate

inflammation following their engulfment by phagocytes (Boccia et al., 2022, Blayney and

Schwartzberg, 2022, Nakazawa et al., 2016, Jeong et al., 2021). Since DNA normally resides in

the nucleus and mitochondria, its presence in the cytoplasm is detected by DNA sensors,

including absent in melanoma 2 (AIM2), and cyclic guanosine monophosphate-adenosine

monophosphate synthase (cGAS), which can trigger immune responses aimed at eliminating the

mislocated DNA (Sun et al., 2013, Zhang et al., 2014, Rathinam et al., 2010). Oxidized DNA and

various DAMPs can also be sensed by NLRP3 (Shimada et al., 2012, Lu et al., 2014, Xian et al.,

2022). Upon recognition of DAMPs or pathogen-associated molecular patterns (PAMPs), AIM2

and NLRP3 assemble protein platforms comprising the adaptor protein apoptosis-associated

speck-like protein containing a CARD (ASC) and caspase-1. These protein complexes known as

inflammasomes are responsible for the maturation of pro-interleukin-1β (pro-IL-1β) and pro-IL-18

to IL-1β and IL-18, respectively (Guo et al., 2015, Sharma and Kanneganti, 2021, Wang et al.,

2021). Inflammasome-comprising caspase-1 also cleaves gasdermin D (GSDMD), generating N-

terminal fragments, which form IL-1β- and IL-18-secreting conduits, and cause the inflammatory

cell death, pyroptosis (Guo et al., 2015, Sharma and Kanneganti, 2021, Wang et al., 2021). While

acute activation of inflammasomes is important for the clearance of the perceived danger and

restoration of homeostasis, chronic or excessive stimulation of these safeguard mechanisms can

cause disease.

Inflammasomes are involved in skeletal pathophysiology. Gain-of-function mutations of

NLRP3 cause skeletal abnormalities in humans (Aksentijevich et al., 2002, Feldmann et al., 2002),

which are reproduced to a great extent in knockin mice expressing NLRP3 harboring mutations

found in these patients (Bonar et al., 2012, Qu et al., 2015, Snouwaert et al., 2016, Wang et al.,

2017). In normal mice, degraded bone matrix components, which are released during bone

resorption promote infammasome activation and OC differentiation (Alippe et al., 2017). Age-

associated bone loss has also been linked to chronic low grade inflammation mediated by the

NLRP3 inflammasome (Youm et al., 2013). More relevant to this study, radiation, which is also

used as an anti-neoplastic therapy, causes bone loss through GSDMD downstream of AIM2 and

NLRP3 inflammasomes, but not NLR family caspase recruitment domain containing protein 4

(NLRC4) inflammasome (Xiao et al., 2020). Collectively, the bone phenotypes of genetically or

pharmacologically activated inflammasome sensors suggest that the fate of bone cells can be

influenced by inflammation-driven by inflammasomes, which are mainly activated in myeloid cells

(Brydges et al., 2009, Meng et al., 2009). This view provides a strong rationale for exploring the

role of inflammasome pathways in bone loss induced by off-target actions of doxorubicin as this

agent causes the death of cancer and bystander normal cells, thereby releasing DAMPs such as

ATP and DNA, which activate these pathways.

We used non-tumor-bearing mouse models, which are commonly used to assess off-

target outcomes of anti-neoplastic therapies (Park et al., 2022, Borniger et al., 2015, Harrison et

al., 1980, Yao et al., 2020) to study bone adverse effects of doxorubicin. We found that

doxorubicin caused massive bone loss in wild-type (WT) mice, a phenotype associated with

increased number of OCs, leukopenia, and cytokinemia. These outcomes implicated the AIM2

and NLRP3 inflammasomes as they were attenuated upon genetic inactivation of these sensors.

Thus, our results show that inflammasomes are key players in bone loss caused by doxorubicin,

a finding that may enable the implementation of novel strategies for chemotherapy-related bone

complications.

Results

Doxorubicin causes bone loss

To determine the effects of doxorubicin on bone mass, femurs of 10 week-old WT female

mice were analyzed by VivaCT before (baseline) and 4 weeks after a single intraperitoneal

injection of 5 mg/kg doxorubicin or vehicle. Doxorubicin, but not the vehicle, caused bone loss

(S1A-D Fig), findings that were consistent with the recently reported stimulatory and suppressive

effects of this drug on bone resorption and formation in WT female mice, respectively (Yao et al.,

2020). Doxorubicin also caused a low bone mass phenotype attended by an increased OC

number and surface in 10 week-old WT male mice (Fig 1A-E and S1E Fig). Since doxorubicin

inflicted bone damage independently of the sex, afterward mechanistic studies focused mainly on

male mice and revolved around innate immune responses, which regulate OC-mediated bone

resorption in pathological conditions.

Doxorubicin causes cytokinemia, leukopenia, release of DAMPs, and NETosis in vivo

Doxorubicin induces inflammatory responses in patients and experimental models

(Wittenburg et al., 2019). Accordingly, WT mice exposed to doxorubicin for 3 days exhibited

higher serum levels of IL-1β, IL-18, IL-6, and TNF-α compared to vehicle-injected counterparts

(Fig 2A-D). Levels of these inflammatory cytokines inversely correlated with the abundance of

white blood cells (WBC; Fig 2E). While doxorubicin lowered the number of circulating

lymphocytes and monocytes, the number of neutrophils, the most abundant immune cells in the

blood, increased 2 hours post-drug exposure before progressively returning to baseline levels.

Consistent with the leukopenic outcome, levels of ATP, which is released by dead cells, were

higher in doxorubicin-treated mice compared to vehicle-treated cohorts (Fig 2F). To gain further

insight into the mechanisms of leukopenia, we focused on neutrophils, the most abundant

leukocytes in blood. These cells exhibit morphological changes such as NET extrusion upon

exposure to PAMPs or sterile DAMPs, and eventually undergo NETosis (Komada et al., 2018,

Apel et al., 2021). To determine the effects of doxorubicin on NET formation, we measured serum

levels of NET components in mice treated with vehicle or this drug for 48 hours. We found that

doxorubicin induced NET formation as evidenced by increased levels of citrullinated histone 3

(cit-H3; Fig 2G), myeloperoxidase (MPO; Fig 2H), and cell-free DNA (cfDNA) (Fig 2I, J). Thus,

doxorubicin causes cytokinemia, a response that is associated with increased cell death and

decreased number of WBCs.

139 Doxorubicin activates inflammasome –dependent and –
140 independent pathways, and caused macrophage pyroptosis

The effects of doxorubicin on macrophages have been reported (Chen et al., 2023, Saleh

et al., 2021).   To test the hypothesis that these cells were implicated in the inflammatory

phenotype of mice treated with doxorubicin, we treated bone marrow-derived macrophages

(BMMs) with lipopolysaccharide (LPS) for 3 hours to induce the expression of inflammasome

components (Wang et al., 2021), then with various concentrations of this drug for 16 hours. Within

its reported potent concentrations (1.5 µM – 12 µM) (Saleh et al., 2021, Eljack et al., 2022),

doxorubicin did not induce IL-1β secretion, but it significantly caused the release of lactate

dehydrogenase (LDH; Fig 3A, B), a marker of cell death as it is released only upon plasma

membrane rupture (Wang et al., 2021). Since LDH release was not induced by doxorubicin in a

dose-dependent manner, this response may be the result of nonselective cytotoxic actions of this

drug. By contrast, doxorubicin promoted IL-1β and LDH release by LPS-primed BMMs in a dose-

dependent fashion, with the maximal effect on IL-1β secretion achieved at 6 µM (Fig 3A).

Unexpectedly, LPS attenuated LDH release induced by low doxorubicin concentrations (1. 5 and

3 µM) (Fig 3B). To further gain insight into the mechanism of action of doxorubicin, we measured

the levels of ATP, which is released by dead cells and activates multiple pathways, including the

NLRP3 inflammasome (Karmakar et al., 2016, Carta et al., 2015). ATP levels were higher in the

supernatants of doxorubicin-treated BMMs compared to controls, a response that was further

enhanced by LPS (Fig 3C). Collectively, these results suggest that BMMs underwent pyroptosis

in the presence of LPS and doxorubicin, releasing DAMPs such as ATP.

To reinforce the proposition that doxorubicin activates inflammasomes, we assessed the

effects of this drug on the formation of ASC specks, a readout of inflammasome-activated states

(Stutz et al., 2013). As anticipated, LPS induced ASC speck formation only in the presence of

nigericin, a well-known trigger of NLRP3 inflammasome assembly signals (Fig 3D, E, H). Likewise,

doxorubicin induced ASC speck formation only in LPS-primed BMMs (Fig 3F, G, H). Next, we

performed immunoblotting to analyze the expression of NLRP3 and AIM2 since these sensors

assemble inflammasomes in response to DAMPs such as ATP and DNA, which were released

by doxorubicin-damaged cells. We also determined the expression of other key components of

these pathways such as caspase-1 and gasdermins. LPS induced the expression of NLRP3, but

not AIM2, caspase-1, caspase-3, GSDMD, and GSDME (Fig 4). The active fragments of

caspase-1 and GSDMD, p10 and p30, respectively, were mainly detected in cells treated with

LPS and doxorubicin (Fig 4 and S2A Fig). GSDMD p10 and GSDME p35, which are

proteolytically generated by caspase-3 (S2A, B Fig), were also detected, but their abundance was

not affected by LPS (Fig 4). Together, these results suggest that doxorubicin activates both

caspase-1 and caspase-3, which cleave GSDMD and GSDME, ultimately, causing pyroptosis and

IL-1β release.

Doxorubicin activates inflammasome –dependent and –

independent pathways, and causes NETosis in vitro

To further support the conclusion that doxorubicin induced the formation of NETs, first, we

analyzed the expression of some key players directly or indirectly involved in this process.

Incubation of mouse bone marrow neutrophils with LPS for 19 hours resulted in increased NLRP3

expression (Fig 5A). GSDMD, GSDME, caspase-1, and caspase-3 were readily detected in

neutrophil lysates (Fig 5A). The abundance of these proteins and their cleaved fragments were

unaltered upon cell exposure to LPS, but markedly reduced in doxorubicin-treated cells, likely

because of cell death as the levels of β-actin used as loading control were markedly reduced in

samples from these cells (Fig 5A). GSDMD p20 and p10 fragments were detected only in

neutrophils treated with LPS and doxorubicin. LPS also induced IL-1β secretion, a response that

was enhanced by doxorubicin in a dose-dependent manner (Fig 5B). LPS reduced baseline as

well as doxorubicin-induced LDH release (Fig 5C). In sum, unchallenged neutrophils released

LDH, a response that aligned with the presence of functional caspase-3, GSDMD, and GSDME

fragments in these cells. However, LPS was required for optimal NLRP3 expression, caspase-1

activation, and IL-1β production by neutrophils. Although neutrophils secreted higher levels of IL-

1β in response to doxorubicin, they were highly sensitive to the cytotoxic effects of this drug.

Next, we performed immunofluorescence to visualize NET components in neutrophils

treated with vehicle or this drug for 16 hours. Consistent with in vivo results, Cit-H3 and MPO

were detected in cells treated with doxorubicin but not with vehicle (Fig 6A). Accordingly, levels

of cfDNA were higher in doxorubicin-exposed cultures compared to untreated or cultures treated

with LPS (Fig 6B). To determine the biological relevance of cfDNA while modeling the in vivo

bone microenvironment, we assessed the impact of degrading cfDNA with DNAse I on IL-1β

release by cultured whole bone marrow cells, which comprised various cell types, including

neutrophils and macrophages. IL-1β levels were higher in cells treated with LPS and doxorubicin

compared to LPS, an outcome that was inhibited by DNAse I (Fig 6C). Thus, doxorubicin

promoted IL-1β release, a response that correlated with its effects on NET formation and

abundance of cfDNA.

207 AIM2 and NLRP3 inflammasomes are involved in bone-
208 damaging effects of doxorubicin

We hypothesized that doxorubicin induced IL-1β and IL-18 release by activating the AIM2

and NLRP3 inflammasomes as blood levels of their activators (DNA, ATP) were increased n

response doxorubicin administration (Fig 2F, I, J).   To test this idea, we measured the effects of

this drug on IL-1β and LDH release by WT, $Aim2^{-/-}$, $Nlrp3^{-/-}$, or $Aim2^{-/-};Nlrp3^{-/-}$ BMMs and

neutrophils. LPS induction of IL-1β and LDH release by BMMs required doxorubicin, a response

that was significantly reduced in $Aim2$ or $Nlrp3$ deficient cells (Fig 7A, B).   LPS induced IL-1β

secretion by neutrophils, an outcome that was enhanced by doxorubicin, and comparably

attenuated in all mutants (Fig 7C, D).   Differences in LDH release were marginal, perhaps

because baseline levels of this readout were high, consistent with the short lifespan of these cells

in vitro. Because inflammation leads to bone loss, we assessed bone outcomes of 10 weeks old

WT or mice globally lacking AIM2 and/or NLRP3, 4 weeks after receiving a single dose of 5 mg/kg

doxorubicin or vehicle. Administration of doxorubicin to WT mice caused bone loss associated

with increased OC number and surface (Fig 7E-I and S3A-C Fig), consistent with the results

shown above (Fig 1).   These responses were reduced slightly in $Nlrp3$ deficient mice, but

significantly in $Aim2$ null mice. $Aim2^{-/-}$ male mice lost bone comparably to $Aim2^{-/-};Nlrp3^{-/-}$ and

$caspase-1^{-/-}$ counterparts (Fig. 7E-G). Similar trends in bone changes were observed in female

mice, though $Aim2^{-/-}$ mice were more osteopenic than $caspase-1^{-/-}$ mice (S4A-D Fig). Collectively,

our results suggest that the AIM2 and NLRP3 inflammasomes participate to various extent in

doxorubicin-bone-damaging effects. They also suggest that inflammasome-independent actions

of this drug on bone are not negligible.

230 Discussion

We found that the AIM2 inflammasome and the NLRP3 inflammasome to a lesser extent,

played an important role in bone-damaging effects of doxorubicin. The comparable bone

phenotype of *Aim2^{-/-};Nlrp3^{-/-}* and *caspase-1^{-/-}* mice suggested that the AIM2 and NLRP3

inflammasomes were the primary mediators of doxorubicin actions. Because doxorubicin

activates several pathways, some of which interact or overlap with inflammasome functions (e.g.,

senescence factors), the remaining bone loss in compound mutant mice was expected. The

interplay among these pathways may have accounted for the sex differences in the bone

outcomes of inflammasome insufficiency as residual doxorubicin-induced bone loss was higher

in *Aim2^{-/-}* and *Aim2^{-/-};Nlrp3^{-/-}* female mice than in male counterparts. Sexual dimorphic actions of

inflammasomes were not unprecedented as uneven severity of atherosclerosis was found in male

and female *Nlrp3*-deficient mice with gonadal insufficiency (Chen et al., 2020) and sex-dependent

differential activation of AIM2 and NLRP3 inflammasomes in macrophages from systemic lupus

erythematous had been reported (Yang et al., 2015). Although the impacts of doxorubicin on

bone pathology are complex, including its direct actions on bone cells (Yao et al., 2020, Chai et

al., 2014, Rana et al., 2013, Zhou et al., 2020), our investigation focused on immune cells, and

found that the inactivation of inflammasomes is sufficient to attenuate the drug's bone-damaging

effects.

Doxorubicin causes neutropenia, lymphopenia, and anemia in patients (Gatti et al., 2018,

Boccia et al., 2022). Consistent with the clinical situation, administration of doxorubicin to mice

caused leukopenia, which correlated with lymphocytopenia and monocytopenia, and was

associated with fluctuations in neutrophil counts. Since neutrophils were highly sensitive to the

cytotoxic effects of doxorubicin, the transient neutrophilic effects of this drug in mice may be the

result of emergency granulopoiesis, a physiological response that is rapidly triggered to restore

adequate neutrophil number. This leukopenic outcome was consistent with increased serum

levels of cell death-associated DAMPS (ATP and cfDNA) and our results showing that doxorubicin

activated the effectors of apoptosis, pyroptosis, and NETosis, including caspase-1, caspase-3,

GSDMD, and GSDME. Since pyroptosis and NETosis promote inflammation and immune

responses, we argued that they accounted for the cytokinemic effects of doxorubicin. Other

studies, however, found that doxorubicin reduced NET formation in cancer models and by human

neutrophils in vitro (Lu et al., 2021, Khan et al., 2019). This discrepancy may be due to differences

in the experimental models and cell context-dependent actions of doxorubicin. Other limitations

of our study include its focus on: i) macrophages and neutrophils while oversighting lymphocytes

or even other cells bone microenvironment such as mesenchymal and adipocytes whose fate is

affected by this drug (Fan et al., 2018, Fan et al., 2017, Wang et al., 2012, Buttiglieri et al., 2011),

which were also targeted by doxorubicin; ii) immune cells without assessing the direct effects of

doxorubicin on bone cells, as noted above, and iii) the use of the tumor-free model as immune

responses can differ significantly in the absence or presence of cancer cells. Despite these

limitations, our findings point to a novel mechanism of action for doxorubicin in bone**.**

DNA accumulates in the cytoplasm as the result of genomic instability, damaged

mitochondria, or lysed intracellular pathogens. Extracellular DNA from pathogens, NETotic, or

pyroptotic cells can be internalized and culminate in the cytoplasm. In either case, sensors such

as AIM2 detect mislocated DNA in the cytoplasm and triggers inflammatory responses (Boccia et

al., 2022, Blayney and Schwartzberg, 2022, Nakazawa et al., 2016, Jeong et al., 2021). This

view was consistent with our data showing a correlation between the levels of extracellular DNA

and IL-1β as well as by the inhibition of IL-1β secretion by DNAse I. We also found that

doxorubicin activated the NLRP3 inflammasome and induced the release of ATP, a well-known

activator of the NLRP3 inflammasome. Whether doxorubicin activated the NLRP3 inflammasome

directly by perturbing the plasma membrane or indirectly via generation of secondary signals such

as ATP remained unclear.

By showing that inflammasomes are key players in bone loss caused by doxorubicin, this

work advances our knowledge on potential mechanisms of action of this drug on this tissue (S5

Fig). This insight is difficult to get in clinical situations because this chemotherapeutic is employed

not alone but in conjunction with other medications (Zimny, 1988, Svendsen et al., 2017, Müller

et al., 2020).

Materials and Methods

Animals

Casp1$^{-/-}$ were kindly provided by Dr. Thirumala-Devi Kanneganti (St. Jude Children's

Research Hospital)*.* WT*, Asc-citrine, Aim2*$^{-/-}$, and *Nlrp3*$^{-/-}$ mice were purchased from The Jackson

Laboratory (Sacramento, CA). *Aim2*$^{-/-}$ mice and *Nlrp3*$^{-/-}$ mice were intercrossed to generate *Aim2*$^{-}$

$^{/-}$;*Nlrp3*$^{-/-}$ mice*.* All mice were on the C57BL6J background and mouse genotyping was performed

by PCR. All procedures were approved by the Institutional Animal Care and Use Committee

(IACUC) of Washington University School of Medicine in St. Louis. All experiments were

performed in accordance with the relevant guidelines and regulations described in the IACUC-

approved protocol 22-0335.

Doxorubicin administration and VivaCT analysis

The femurs of 10 week-old mice were analyzed by VivaCT 2 weeks before (baseline) and

4 weeks after a single intraperitoneal injection (i.p.) of 5 mg/kg doxorubicin (Sigma-Aldrich, MO)

formulated in H_2O at 1 mg/ml or vehicle. For bone analysis, mice were anesthetized with

isofluorane and trabecular volume in the distal femoral metaphysis of the right leg was measured

using VivaCT 40 (Scanco Medical AG, Zurich, Switzerland) set at 70 kVp, 114 µA, and 20 µm

resolution as previously described (Yao et al., 2020). For the trabecular bone compartment,

contours were traced on the inside of the cortical shell using 2D images of the femoral metaphysis.

The end of the growth plate region was used as a landmark to establish a consistent location for

starting analysis, and the next 50 slices were analyzed. The following trabecular parameters are

reported for all Viva CT experiments: bone volume over total volume (BV/TV), trabecular number

(Tb.N), trabecular thickness (Tb.Th), trabecular separation (Tb.Sp), and volumetric bone mineral

density (vBMD).

Histomorphometry

For static histomorphometry, the femurs were fixed in 10% neutral buffered formalin

overnight and decalcified in 14% (wt/vol) EDTA, pH 7.2, for 10–14 days at room temperature.

Fixed femurs were embedded in paraffin, sectioned at 5-μm thicknesses, and mounted on glass

slides. The sections were stained with tartrate-resistant acidic phosphatase (TRAP) as described

previously (Wang et al., 2018). For dynamic histomorphometry, mice were i.p. injected with 10

mg/kg calcein green (Sigma-Aldrich, MO) and 6 days later with 50 mg/kg alizarin red (Sigma-

Aldrich, MO). Mice were euthanized 2 days after the second injection. The tibias were collected

and fixed in 10% neutral buffered formalin overnight, embedded in methyl methacrylate, and

sectioned at 7–10 μm. Images were taken using a Nanozoomer 2.0 HT whole slide scanner

(Hamamatsu Photonics, Shizuoka, Japan) at 20X magnification. Bioquant Osteo software

(v18.2.6; Bioquant Image Analysis Corp, TN) was used for image analysis. Measurements of

dynamic bone histomorphometry were calculated from fluorochrome double labels at the

endocortical surfaces as previously described (Xiao et al., 2020).

Serum assays

Blood was collected by cardiac puncture and was allowed to clot at room temperature.

Serum obtained after centrifugation at 2,000 x g for 10 minutes was used for various assays.

Cytokine and chemokine levels were measured by V-PLEX Plus Proinflammatory Panel 1 Mouse

Kit (Meso Scale Diagnostics, MD), except IL-18, which was analyzed by enzyme linked

immunosorbent assay (ELISA) kit (Sigma-Aldrich, MO). The levels of Citrine-H3 (Cit-H3) and

myeloperoxidase (MPO) were determined by ELISA kits (Abcam, MA & Cayman, MI).

337   Peripheral blood analysis

338   Mouse blood was collected by cardiac puncture in the EDTA-containing tubes. Complete

339   blood counts were performed by the Washington University School of Medicine as previously

340   described (Wang et al., 2017).

342   Cell cultures

Murine primary bone marrow-derived macrophages (BMMs) were obtained by culturing

mouse bone marrow cells in culture media containing a 1:10 dilution of supernatant from the

fibroblastic cell line CMG 14-12 as a source of macrophage colony-stimulating factor, a mitogenic

factor for BMMs, for 4-5 days in a 15-cm dish as previously described (Takeshita et al., 2000,

Wang et al., 2020). Briefly, nonadherent cells were removed by vigorous washes with PBS, and

adherent BMMs were detached with trypsin-EDTA and cultured in culture media containing a 1:10

dilution of CMG for various experiments. Murine primary neutrophils were isolated by collecting

bone marrow cells and subsequently over a discontinuous Percoll (Sigma Aldrich, MO) gradient

as described previously (Sun et al., 2022). Briefly, all bone marrow cells from femurs and tibias

were washed by PBS and then resuspended in 2 ml PBS. Cell suspension was gently layered

on top of gradient (72% Percoll, 64% Percoll, 52% Percoll) and centrifuged at 1545x g for 30

minutes at room temperature. After carefully discarding the top two cell layers, the third layer

containing neutrophils was transferred to a clean 15 ml tube. Cells were washed and counted,

then plated at a density of 1×10^5 cells/well in 96-well plate or 5×10^6 cells/well in 6-well plate for

1 hour followed by various experiments. For all *in vitro* experiments except otherwise specified,

BMMs were plated at 2×10^4 cells per well on a 96-well plate or 10^6 cells per well on a 6-well plate

overnight. Neutrophils were plated at 10^5 cells per well on a 96-well plate or 5×10^6 cells per well

on a 6-well plate for one hour prior to treatment. BMMs and neutrophils were primed with 100

ng/ml LPS (Sigma-Aldrich, MO) for 3 hours, then with different concentrations of doxorubicin

(Sigma-Aldrich, MO) as indicated for 16 hours. Conditioned media were collected for the analysis

of IL-1β and lactate dehydrogenase (LDH). Cell lysates were collected for protein expression

analysis by Western blot as described below.

366   Western blot analysis

Cell extracts were prepared by lysing cells with RIPA buffer (50 mM Tris, 150 mM NaCl, 1

mM EDTA, 0.5% NaDOAc, 0.1% SDS, and 1.0% NP-40) plus phosphatase and protease

inhibitors (GenDEPOT, TX). Protein concentrations were determined by the Bio-Rad Laboratories

method (Bio-Rad, CA), and equal amounts of proteins were subjected to SDS-PAGE gels (12%

or 15%) as previously described (Wang et al., 2021). Proteins were transferred onto nitrocellulose

membranes and incubated with antibodies against GSDMD (1;1,000; Abcam, MA), GSDME

(1;1,000; Abcam, MA), caspase-1 (1;1,000; Abcam, MA), caspase-3 (1:1,000; Cell Signaling

Technologies, MA), NLRP3 (1:1,000; AdipoGen,CA), AIM2 (1:1000; Cell Signaling Technologies,

MA) or β-actin (1:2,000; Santa Cruz Biotechnology, TX) overnight at 4°C followed by incubation

for 1 hour with secondary goat anti-mouse IRDye 800 (Thermo Fisher Scientific, MA) or goat anti–

rabbit Alexa Fluor 680 (Thermo Fisher Scientific, MA), respectively. The results were visualized

using the Odyssey infrared imaging system (LI-COR Biosciences, NE).

LDH assay and IL-1β ELISA

380   Cell death was assessed by the release of LDH in conditioned medium using LDH

381 Cytotoxicity Detection Kit (TaKaRa, CA). IL-1β levels in conditioned media were measured by an

383 ELISA kit (eBiosciences, NY).

ASC specks assay

386   *Asc-citrine*-WT BMMs were plated at 10^4 cells per well on a 16-well glass plate overnight.

387 Cells were primed with LPS for 3 hours followed by 15 µM nigericin (AdipoGen, CA) for 30 minutes

388 or the indicated doxorubicin concentrations for 16 hours. Cells were washed with PBS, fixed with

389 4% paraformaldehyde buffer for 10 minutes at room temperature, then counterstained with

390 Fluoro-gel II containing DAPI (Fluoro-Gel, Fisher Scientific Intl INC, PA). Asc-citrine photographs

391 were taken using ZEISS microscopy (Carl Zeiss Industrial Metrology, MN). Quantification of ASC

392 specks was carried out using ImageJ.

Immunofluorescence

395   Isolated neutrophils were plated at 10^5 cells per well on a 16-well glass plate for 1 hour.

396 Cells were primed with LPS for 3 hours, treated with doxorubicin for 16 hours, washed with PBS,

397 and fixed with 4% paraformaldehyde buffer for 10 minutes at room temperature. Cells were

398 permeabilized with 0.2% Triton in PBS for 20 minutes, blocked with 0.2% Triton, 1% BSA, and

399 CD61 antibody (1:1000; Alexa Fluor 647, BD, NJ) in PBS for 30 minutes, and were incubated with

400 Cit-H3 antibody (1:1,000; Cayman, MI) and MPO (1;1,000; Abcam, MA) overnight at 4°C in

401 blocking buffer, followed by incubation with secondary antibody (Alexa Fluor 594, 1:2,000; Life

402 Technologies, CA) for 30 minutes. Cells were counterstained with Fluoro-gel II containing DAPI

403 (Fluoro-Gel, Fisher Scientific Intl INC, PA). Immunostaining images were taken using a Leica

404 inverted microscope with a TCS SPEII confocal module and processed using LAS X software

405 (Leica Microsystems Inc, IL).

cfDNA assay

cfDNA in the conditioned cell culture medium was extracted using NucleoSpin Gel and

PCR Clean-up kit (Takara, Duren, Germany), and quantified with Nanodrop (Thermo Fisher

Scientific, MA). cfDNA in the serum was purified and measured using Qubit by Washington

University School of Medicine Hope Center DNA/RNA purification Core or using SYBR Green

(Thermo Fisher Scientific, MA) as previously described (Goldshtein et al., 2009, Villalba-Campos

et al., 2016).

ATP assay

416 ATP levels in conditioned media and serum were measured by RealTime-Glo Extracellular

417 ATP Assay kit (Promega, Madison, WI).

Statistical analysis

Statistical analysis was performed using the Student's t test, one-way ANOVA with Tukey's

multiple comparisons test, or two-way ANOVA with Tukey's multiple comparisons test, Dunnett's

multiple comparisons test, or Sidak's multiple comparisons test using the GraphPad Prism 9.0

Software. Values are expressed as mean ± SEM. *$p < 0.05$ was considered statistically significant.

Acknowledgements

We want to thank Dr. Deborah J. Veis for reading this manuscript.

References

ABE, K., IKEDA, M., IDE, T., TADOKORO, T., MIYAMOTO, H. D., FURUSAWA, S., TSUTSUI, Y., MIYAKE, R., ISHIMARU, K., WATANABE, M., MATSUSHIMA, S., KOUMURA, T., YAMADA, K. I., IMAI, H. & TSUTSUI, H. 2022. Doxorubicin causes ferroptosis and cardiotoxicity by intercalating into mitochondrial DNA and disrupting Alas1-dependent heme synthesis. *Sci Signal,* 15**,** eabn8017.

AKSENTIJEVICH, I., NOWAK, M., MALLAH, M., CHAE, J. J., WATFORD, W. T., HOFMANN, S. R., STEIN, L., RUSSO, R., GOLDSMITH, D., DENT, P., ROSENBERG, H. F., AUSTIN, F., REMMERS, E. F., BALOW, J. E., JR., ROSENZWEIG, S., KOMAROW, H., SHOHAM, N. G., WOOD, G., JONES, J., MANGRA, N., CARRERO, H., ADAMS, B. S., MOORE, T. L., SCHIKLER, K., HOFFMAN, H., LOVELL, D. J., LIPNICK, R., BARRON, K., O'SHEA, J. J., KASTNER, D. L. & GOLDBACH-MANSKY, R. 2002. De novo CIAS1 mutations, cytokine activation, and evidence for genetic heterogeneity in patients with neonatal-onset multisystem inflammatory disease (NOMID): a new member of the expanding family of pyrin-associated autoinflammatory diseases. *Arthritis Rheum,* 46**,** 3340-8.

ALIPPE, Y., WANG, C., RICCI, B., XIAO, J., QU, C., ZOU, W., NOVACK, D. V., ABU-AMER, Y., CIVITELLI, R. & MBALAVIELE, G. 2017. Bone matrix components activate the NLRP3 inflammasome and promote osteoclast differentiation. *Sci Rep,* 7**,** 6630.

APEL, F., ANDREEVA, L., KNACKSTEDT, L. S., STREECK, R., FRESE, C. K., GOOSMANN, C., HOPFNER, K. P. & ZYCHLINSKY, A. 2021. The cytosolic DNA sensor cGAS recognizes neutrophil extracellular traps. *Sci Signal,* 14.

BLAYNEY, D. W. & SCHWARTZBERG, L. 2022. Chemotherapy-induced neutropenia and emerging agents for prevention and treatment: A review. *Cancer Treat Rev,* 109**,** 102427.

BOCCIA, R., GLASPY, J., CRAWFORD, J. & AAPRO, M. 2022. Chemotherapy-Induced Neutropenia and Febrile Neutropenia in the US: A Beast of Burden That Needs to Be Tamed? *Oncologist,* 27**,** 625-636.

BONAR, S. L., BRYDGES, S. D., MUELLER, J. L., MCGEOUGH, M. D., PENA, C., CHEN, D., GRIMSTON, S. K., HICKMAN-BRECKS, C. L., RAVINDRAN, S., MCALINDEN, A., NOVACK, D. V., KASTNER, D. L., CIVITELLI, R., HOFFMAN, H. M. & MBALAVIELE, G. 2012. Constitutively activated NLRP3 inflammasome causes inflammation and abnormal skeletal development in mice. *PLoS One,* 7**,** e35979.

BONNER, J. A. & LAWRENCE, T. S. 1990. Doxorubicin decreases the repair of radiation-induced DNA damage. *Int J Radiat Biol,* 57**,** 55-64.

BORNIGER, J. C., GAUDIER-DIAZ, M. M., ZHANG, N., NELSON, R. J. & DEVRIES, A. C. 2015. Cytotoxic chemotherapy increases sleep and sleep fragmentation in non-tumor-bearing mice. *Brain Behav Immun,* 47**,** 218-27.

BRYDGES, S. D., MUELLER, J. L., MCGEOUGH, M. D., PENA, C. A., MISAGHI, A., GANDHI, C., PUTNAM, C. D., BOYLE, D. L., FIRESTEIN, G. S., HORNER, A. A., SOROOSH, P., WATFORD, W. T., O'SHEA, J. J., KASTNER, D. L. & HOFFMAN, H. M. 2009. Inflammasome-mediated disease animal models reveal roles for innate but not adaptive immunity. *Immunity,* 30**,** 875-87.

BUTTIGLIERI, S., RUELLA, M., RISSO, A., SPATOLA, T., SILENGO, L., AVVEDIMENTO, E. V. & TARELLA, C. 2011. The aging effect of chemotherapy on cultured human mesenchymal stem cells. *Exp Hematol,* 39**,** 1171-81.

CARDINALE, D., COLOMBO, A., BACCHIANI, G., TEDESCHI, I., MERONI, C. A., VEGLIA, F., CIVELLI, M., LAMANTIA, G., COLOMBO, N., CURIGLIANO, G., FIORENTINI, C. & CIPOLLA, C. M. 2015. Early detection of anthracycline cardiotoxicity and improvement with heart failure therapy. *Circulation,* 131**,** 1981-8.

CARTA, S., PENCO, F., LAVIERI, R., MARTINI, A., DINARELLO, C. A., GATTORNO, M. & RUBARTELLI, A. 2015. Cell stress increases ATP release in NLRP3 inflammasome-mediated autoinflammatory diseases, resulting in cytokine imbalance. *Proc Natl Acad Sci U S A,* 112**,** 2835-40.

CAULEY, J. A., LUI, L. Y., BARNES, D., ENSRUD, K. E., ZMUDA, J. M., HILLIER, T. A., HOCHBERG, M. C., SCHWARTZ, A. V., YAFFE, K., CUMMINGS, S. R. & NEWMAN, A. B. 2009. Successful skeletal aging: a marker of low fracture risk and longevity. The Study of Osteoporotic Fractures (SOF). *J Bone Miner Res,* 24**,** 134-43.

CAWTHON, P. M., EWING, S. K., MACKEY, D. C., FINK, H. A., CUMMINGS, S. R., ENSRUD, K. E., STEFANICK, M. L., BAUER, D. C., CAULEY, J. A. & ORWOLL, E. S. 2012. Change in hip bone mineral density and risk of subsequent fractures in older men. *J Bone Miner Res,* 27**,** 2179-88.

CAWTHON, P. M., EWING, S. K., MCCULLOCH, C. E., ENSRUD, K. E., CAULEY, J. A., CUMMINGS, S. R. & ORWOLL, E. S. 2009. Loss of hip BMD in older men: the osteoporotic fractures in men (MrOS) study. *J Bone Miner Res,* 24**,** 1728-35.

CHAI, R. C., KOUSPOU, M. M., LANG, B. J., NGUYEN, C. H., VAN DER KRAAN, A. G., VIEUSSEUX, J. L., LIM, R. C., GILLESPIE, M. T., BENJAMIN, I. J., QUINN, J. M. & PRICE, J. T. 2014. Molecular stress-inducing compounds increase osteoclast formation in a heat shock factor 1 protein-dependent manner. *J Biol Chem,* 289**,** 13602-14.

CHEN, S., HUANG, Y., HUANG, S., ZHOU, Z., LIU, K., PAN, J. & WU, Z. 2023. M2b macrophages protect against doxorubicin induced cardiotoxicity via alternating autophagy in cardiomyocytes. *PLoS One,* 18**,** e0288422.

CHEN, S., MARKMAN, J. L., SHIMADA, K., CROTHER, T. R., LANE, M., ABOLHESN, A., SHAH, P. K. & ARDITI, M. 2020. Sex-Specific Effects of the Nlrp3 Inflammasome on Atherogenesis in LDL Receptor-Deficient Mice. *JACC Basic Transl Sci,* 5**,** 582-598.

COLEMAN, R. E., RATHBONE, E. & BROWN, J. E. 2013. Management of cancer treatment-induced bone loss. *Nat Rev Rheumatol,* 9**,** 365-74.

ELJACK, S., ALLARD-VANNIER, E., MISERICORDIA, Y., HERVÉ-AUBERT, K., AUBREY, N., CHOURPA, I., FAGGAD, A. & DAVID, S. 2022. Combination of Nanovectorized siRNA Directed against Survivin with Doxorubicin for Efficient Anti-Cancer Activity in HER2+ Breast Cancer Cells. *Pharmaceutics,* 14.

FAN, C., GEORGIOU, K. R., MORRIS, H. A., MCKINNON, R. A., KEEFE, D. M. K., HOWE, P. R. & XIAN, C. J. 2017. Combination breast cancer chemotherapy with doxorubicin and cyclophosphamide damages bone and bone marrow in a female rat model. *Breast Cancer Res Treat,* 165**,** 41-51.

FAN, C. M., SU, Y. W., HOWE, P. R. & XIAN, C. J. 2018. Long Chain Omega-3 Polyunsaturated Fatty Acid Supplementation Protects Against Adriamycin and Cyclophosphamide Chemotherapy-Induced Bone Marrow Damage in Female Rats. *Int J Mol Sci,* 19.

FELDMANN, J., PRIEUR, A. M., QUARTIER, P., BERQUIN, P., CERTAIN, S., CORTIS, E., TEILLAC-HAMEL, D., FISCHER, A. & DE SAINT BASILE, G. 2002. Chronic infantile neurological cutaneous and articular syndrome is caused by mutations in CIAS1, a gene highly expressed in polymorphonuclear cells and chondrocytes. *Am J Hum Genet,* 71**,** 198-203.

GATTI, D. M., WEBER, S. N., GOODWIN, N. C., LAMMERT, F. & CHURCHILL, G. A. 2018. Genetic background influences susceptibility to chemotherapy-induced hematotoxicity. *Pharmacogenomics J,* 18**,** 319-330.

GEFFEN, D. B. & MAN, S. 2002. New drugs for the treatment of cancer, 1990-2001. *Isr Med Assoc J,* 4**,** 1124-31.

GOLDSHTEIN, H., HAUSMANN, M. J. & DOUVDEVANI, A. 2009. A rapid direct fluorescent assay for cell-free DNA quantification in biological fluids. *Ann Clin Biochem,* 46**,** 488-94.

GUO, H., CALLAWAY, J. B. & TING, J. P. 2015. Inflammasomes: mechanism of action, role in disease, and therapeutics. *Nat Med,* 21**,** 677-87.

HARRISON, S. D., JR., GILES, H. D. & DENINE, E. P. 1980. Antitumor drug toxicity in tumor-free and tumor-
bearing mice. *Cancer Chemother Pharmacol,* 4**,** 199-204.
JANG, J. Y., KIM, D. & KIM, N. D. 2023. Recent Developments in Combination Chemotherapy for Colorectal
and Breast Cancers with Topoisomerase Inhibitors. *Int J Mol Sci,* 24.
JEONG, J. H., CHOI, S. J., AHN, S. M., OH, J. S., KIM, Y. G., LEE, C. K., YOO, B. & HONG, S. 2021. Neutrophil
extracellular trap clearance by synovial macrophages in gout. *Arthritis Res Ther,* 23**,** 88.
KARMAKAR, M., KATSNELSON, M. A., DUBYAK, G. R. & PEARLMAN, E. 2016. Neutrophil P2X7 receptors
mediate NLRP3 inflammasome-dependent IL-1β secretion in response to ATP. *Nat Commun,* 7**,**
10555.
KHAN, M. A., D'OVIDIO, A., TRAN, H. & PALANIYAR, N. 2019. Anthracyclines Suppress Both NADPH Oxidase-
Dependent and -Independent NETosis in Human Neutrophils. *Cancers (Basel),* 11.
KOMADA, T., CHUNG, H., LAU, A., PLATNICH, J. M., BECK, P. L., BENEDIKTSSON, H., DUFF, H. J., JENNE, C. N.
& MURUVE, D. A. 2018. Macrophage Uptake of Necrotic Cell DNA Activates the AIM2
Inflammasome to Regulate a Proinflammatory Phenotype in CKD. *J Am Soc Nephrol,* 29**,** 1165-
1181.
LAWRENCE, T. S. 1988. Reduction of doxorubicin cytotoxicity by ouabain: correlation with topoisomerase-
induced DNA strand breakage in human and hamster cells. *Cancer Res,* 48**,** 725-30.
LU, B., KWAN, K., LEVINE, Y. A., OLOFSSON, P. S., YANG, H., LI, J., JOSHI, S., WANG, H., ANDERSSON, U.,
CHAVAN, S. S. & TRACEY, K. J. 2014. α7 nicotinic acetylcholine receptor signaling inhibits
inflammasome activation by preventing mitochondrial DNA release. *Mol Med,* 20**,** 350-8.
LU, Z., LONG, Y., LI, J., LI, J., REN, K., ZHAO, W., WANG, X., XIA, C., WANG, Y., LI, M., ZHANG, Z. & HE, Q. 2021.
Simultaneous inhibition of breast cancer and its liver and lung metastasis by blocking
inflammatory feed-forward loops. *J Control Release,* 338**,** 662-679.
MENG, G., ZHANG, F., FUSS, I., KITANI, A. & STROBER, W. 2009. A mutation in the Nlrp3 gene causing
inflammasome hyperactivation potentiates Th17 cell-dominant immune responses. *Immunity,* 30**,**
860-74.
MÜLLER, A., DREYLING, M., ROEDER, F., BAUR-MELNYK, A., KNÖSEL, T., KLEIN, A., BIRKENMAIER, C.,
JANSSON, V. & DÜRR, H. R. 2020. Primary bone lymphoma: Clinical presentation and therapeutic
considerations. *J Bone Oncol,* 25**,** 100326.
NAKAZAWA, D., SHIDA, H., KUSUNOKI, Y., MIYOSHI, A., NISHIO, S., TOMARU, U., ATSUMI, T. & ISHIZU, A.
2016. The responses of macrophages in interaction with neutrophils that undergo NETosis. *J*
Autoimmun, 67**,** 19-28.
PARK, H. J., YOON, S. Y., PARK, J. N., SUH, J. H. & CHOI, H. S. 2022. Doxorubicin Induces Bone Loss by
Increasing Autophagy through a Mitochondrial ROS/TRPML1/TFEB Axis in Osteoclasts.
Antioxidants (Basel), 11.
QU, C., BONAR, S. L., HICKMAN-BRECKS, C. L., ABU-AMER, S., MCGEOUGH, M. D., PEÑA, C. A., BRODERICK,
L., YANG, C., GRIMSTON, S. K., KADING, J., ABU-AMER, Y., NOVACK, D. V., HOFFMAN, H. M.,
CIVITELLI, R. & MBALAVIELE, G. 2015. NLRP3 mediates osteolysis through inflammation-
dependent and -independent mechanisms. *Faseb j,* 29**,** 1269-79.
RANA, T., CHAKRABARTI, A., FREEMAN, M. & BISWAS, S. 2013. Doxorubicin-mediated bone loss in breast
cancer bone metastases is driven by an interplay between oxidative stress and induction of TGFβ.
PLoS One, 8**,** e78043.
RATHINAM, V. A., JIANG, Z., WAGGONER, S. N., SHARMA, S., COLE, L. E., WAGGONER, L., VANAJA, S. K.,
MONKS, B. G., GANESAN, S., LATZ, E., HORNUNG, V., VOGEL, S. N., SZOMOLANYI-TSUDA, E. &
FITZGERALD, K. A. 2010. The AIM2 inflammasome is essential for host defense against cytosolic
bacteria and DNA viruses. *Nat Immunol,* 11**,** 395-402.

ROBSON, C. N., HOBAN, P. R., HARRIS, A. L. & HICKSON, I. D. 1987. Cross-sensitivity to topoisomerase II inhibitors in cytotoxic drug-hypersensitive Chinese hamster ovary cell lines. *Cancer Res,* 47**,** 1560-5.

SALEH, H. A., RAMDAN, E., ELMAZAR, M. M., AZZAZY, H. M. E. & ABDELNASER, A. 2021. Comparing the protective effects of resveratrol, curcumin and sulforaphane against LPS/IFN-γ-mediated inflammation in doxorubicin-treated macrophages. *Sci Rep,* 11**,** 545.

SHARMA, B. R. & KANNEGANTI, T. D. 2021. NLRP3 inflammasome in cancer and metabolic diseases. *Nat Immunol,* 22**,** 550-559.

SHIMADA, K., CROTHER, T. R., KARLIN, J., DAGVADORJ, J., CHIBA, N., CHEN, S., RAMANUJAN, V. K., WOLF, A. J., VERGNES, L., OJCIUS, D. M., RENTSENDORJ, A., VARGAS, M., GUERRERO, C., WANG, Y., FITZGERALD, K. A., UNDERHILL, D. M., TOWN, T. & ARDITI, M. 2012. Oxidized mitochondrial DNA activates the NLRP3 inflammasome during apoptosis. *Immunity,* 36**,** 401-14.

SMITH, L. A., CORNELIUS, V. R., PLUMMER, C. J., LEVITT, G., VERRILL, M., CANNEY, P. & JONES, A. 2010. Cardiotoxicity of anthracycline agents for the treatment of cancer: systematic review and meta-analysis of randomised controlled trials. *BMC Cancer,* 10**,** 337.

SNOUWAERT, J. N., NGUYEN, M., REPENNING, P. W., DYE, R., LIVINGSTON, E. W., KOVAROVA, M., MOY, S. S., BRIGMAN, B. E., BATEMAN, T. A., TING, J. P. & KOLLER, B. H. 2016. An NLRP3 Mutation Causes Arthropathy and Osteoporosis in Humanized Mice. *Cell Rep,* 17**,** 3077-3088.

STUTZ, A., HORVATH, G. L., MONKS, B. G. & LATZ, E. 2013. ASC speck formation as a readout for inflammasome activation. *Methods Mol Biol,* 1040**,** 91-101.

SUN, K., WANG, C., XIAO, J., BRODT, M. D., YUAN, L., YANG, T., ALIPPE, Y., HU, H., HAO, D., ABU-AMER, Y., SILVA, M. J., SHEN, J. & MBALAVIELE, G. 2022. Fracture healing is delayed in the absence of gasdermin-interleukin-1 signaling. *Elife,* 11.

SUN, L., WU, J., DU, F., CHEN, X. & CHEN, Z. J. 2013. Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science,* 339**,** 786-91.

SVENDSEN, P., SHEKHRAJKA, N., NIELSEN, K. L., VESTERGAARD, P., POULSEN, M., VISTISEN, A. K., MUNKSGAARD, P. S., SEVERINSEN, M. T., JENSEN, P., JOHNSEN, H. E., JAKOBSEN, L. H., BØGSTED, M., FRØKJÆR, J. B. & EL-GALALY, T. C. 2017. R-CHOP(-like) treatment of diffuse large B-cell lymphoma significantly reduces CT-assessed vertebral bone density: a single center study of 111 patients. *Leuk Lymphoma,* 58**,** 1105-1113.

TACAR, O., SRIAMORNSAK, P. & DASS, C. R. 2013. Doxorubicin: an update on anticancer molecular action, toxicity and novel drug delivery systems. *J Pharm Pharmacol,* 65**,** 157-70.

TAKESHITA, S., KAJI, K. & KUDO, A. 2000. Identification and characterization of the new osteoclast progenitor with macrophage phenotypes being able to differentiate into mature osteoclasts. *J Bone Miner Res,* 15**,** 1477-88.

TEWEY, K. M., ROWE, T. C., YANG, L., HALLIGAN, B. D. & LIU, L. F. 1984. Adriamycin-induced DNA damage mediated by mammalian DNA topoisomerase II. *Science,* 226**,** 466-8.

VILLALBA-CAMPOS, M., RAMÍREZ-CLAVIJO, S. R., SÁNCHEZ-CORREDOR, M. C., RONDÓN-LAGOS, M., IBÁÑEZ-PINILLA, M., PALMA, R. M., VARONA-URIBE, M. E. & CHUAIRE-NOACK, L. 2016. Quantification of cell-free DNA for evaluating genotoxic damage from occupational exposure to car paints. *J Occup Med Toxicol,* 11**,** 33.

WANG, C., HOCKERMAN, S., JACOBSEN, E. J., ALIPPE, Y., SELNESS, S. R., HOPE, H. R., HIRSCH, J. L., MNICH, S. J., SAABYE, M. J., HOOD, W. F., BONAR, S. L., ABU-AMER, Y., HAIMOVICH, A., HOFFMAN, H. M., MONAHAN, J. B. & MBALAVIELE, G. 2018. Selective inhibition of the p38α MAPK-MK2 axis inhibits inflammatory cues including inflammasome priming signals. *J Exp Med,* 215**,** 1315-1325.

WANG, C., XIAO, J., NOWAK, K., GUNASEKERA, K., ALIPPE, Y., SPECKMAN, S., YANG, T., KRESS, D., ABU-AMER, Y., HOTTIGER, M. O. & MBALAVIELE, G. 2020. PARP1 Hinders Histone H2B Occupancy at the NFATc1 Promoter to Restrain Osteoclast Differentiation. *J Bone Miner Res,* 35**,** 776-788.

WANG, C., XU, C. X., ALIPPE, Y., QU, C., XIAO, J., SCHIPANI, E., CIVITELLI, R., ABU-AMER, Y. & MBALAVIELE,
G. 2017. Chronic inflammation triggered by the NLRP3 inflammasome in myeloid cells promotes
growth plate dysplasia by mesenchymal cells. *Sci Rep,* 7**,** 4880.
WANG, C., YANG, T., XIAO, J., XU, C., ALIPPE, Y., SUN, K., KANNEGANTI, T. D., MONAHAN, J. B., ABU-AMER,
Y., LIEBERMAN, J. & MBALAVIELE, G. 2021. NLRP3 inflammasome activation triggers gasdermin D-
independent inflammation. *Sci Immunol,* 6**,** eabj3859.
WANG, W. L., KATZ, D., ARAUJO, D. M., RAVI, V., LUDWIG, J. A., TRENT, J. C., PATEL, S. R., LIN, P. P.,
GUADAGNOLO, A., LÒPEZ-TERRADA, D., DEI TOS, A. P., LEWIS, V. O., LEV, D., POLLOCK, R. E.,
ZAGARS, G. K., BENJAMIN, R. S., MADEWELL, J. E. & LAZAR, A. J. 2012. Extensive adipocytic
maturation can be seen in myxoid liposarcomas treated with neoadjuvant doxorubicin and
ifosfamide and pre-operative radiation therapy. *Clin Sarcoma Res,* 2**,** 25.
WITTENBURG, L. A., WEISHAAR, K., RAMIREZ, D. & GUSTAFSON, D. L. 2019. Doxorubicin area under the
curve is an important predictor of neutropenia in dogs with naturally occurring cancers. *Vet Comp*
Oncol, 17**,** 147-154.
XIAN, H., WATARI, K., SANCHEZ-LOPEZ, E., OFFENBERGER, J., ONYURU, J., SAMPATH, H., YING, W.,
HOFFMAN, H. M., SHADEL, G. S. & KARIN, M. 2022. Oxidized DNA fragments exit mitochondria via
mPTP- and VDAC-dependent channels to activate NLRP3 inflammasome and interferon signaling.
Immunity, 55**,** 1370-1385.e8.
XIAO, J., WANG, C., YAO, J. C., ALIPPE, Y., YANG, T., KRESS, D., SUN, K., KOSTECKI, K. L., MONAHAN, J. B.,
VEIS, D. J., ABU-AMER, Y., LINK, D. C. & MBALAVIELE, G. 2020. Radiation causes tissue damage by
dysregulating inflammasome-gasdermin D signaling in both host and transplanted cells. *PLoS Biol,*
18**,** e3000807.
YANG, C. A., HUANG, S. T. & CHIANG, B. L. 2015. Sex-dependent differential activation of NLRP3 and AIM2
inflammasomes in SLE macrophages. *Rheumatology (Oxford),* 54**,** 324-31.
YAO, Z., MURALI, B., REN, Q., LUO, X., FAGET, D. V., COLE, T., RICCI, B., THOTALA, D., MONAHAN, J., VAN
DEURSEN, J. M., BAKER, D., FACCIO, R., SCHWARZ, J. K. & STEWART, S. A. 2020. Therapy-Induced
Senescence Drives Bone Loss. *Cancer Res,* 80**,** 1171-1182.
YOUM, Y. H., GRANT, R. W., MCCABE, L. R., ALBARADO, D. C., NGUYEN, K. Y., RAVUSSIN, A., PISTELL, P.,
NEWMAN, S., CARTER, R., LAQUE, A., MÜNZBERG, H., ROSEN, C. J., INGRAM, D. K., SALBAUM, J.
M. & DIXIT, V. D. 2013. Canonical Nlrp3 inflammasome links systemic low-grade inflammation to
functional decline in aging. *Cell Metab,* 18**,** 519-32.
ZHANG, X., WU, J., DU, F., XU, H., SUN, L., CHEN, Z., BRAUTIGAM, C. A., ZHANG, X. & CHEN, Z. J. 2014. The
cytosolic DNA sensor cGAS forms an oligomeric complex with DNA and undergoes switch-like
conformational changes in the activation loop. *Cell Rep,* 6**,** 421-30.
ZHOU, L., KUAI, F., SHI, Q. & YANG, H. 2020. Doxorubicin restrains osteogenesis and promotes
osteoclastogenesis in vitro. *Am J Transl Res,* 12**,** 5640-5654.
ZIMNY, M. L. 1988. Mechanoreceptors in articular tissues. *Am J Anat,* 182**,** 16-32.

Figures

Figure 1. Doxorubicin causes bone loss in male mice. (A, B) Femurs from WT male mice
were analyzed by VivaCT before (baseline) and 4 weeks after a single intraperitoneal injection of
5 mg/kg doxorubicin. (A) Cross sections of 3D reconstructions. Scale bar: 500 μm. (B) Bone
parameters. (C-E) Femurs from WT male mice, analyzed 4 weeks after a single intraperitoneal
injection of vehicle or doxorubicin. Specimens were stained for TRAP activity. (C) Representative
images. Scale bar: 500 μm. (D) N.Oc/BS. (E) Oc.S/BS. N=5 mice/group. Data are mean ± SEM.
Student t test was used. *P < 0.05; **P < 0.01; ***P < 0.001. BMD, bone mineral density; BV/TV,
bone volume/total volume; Dox, doxorubicin; N.Oc/BS, OC number/bone surface; Oc.S/BS, OC
surface/bone surface; OC, osteoclast; ns, not significant; Tb.N, trabecular number; Tb.Th,
trabecular thickness; Tb.Sp, trabecular separation; WT, wild-type.

Figure 2. Doxorubicin causes cytokinemia, leukopenia, release of DAMPs, and NETosis in vivo. Twelve-week-old WT mice were exposed to a single dose of vehicle or 5 mg/kg doxorubicin. Serum samples were harvested 3 days (A-D) or 2 days (F-J) later and analyzed by MSD (IL-1β, IL-6, and TNF-α) or ELISA (IL-18). (E) Blood was collected for cell counts at the indicated time-points after a single dose of 5 mg/kg doxorubicin injection. cfDNA was measured using Qubit (I) or SYBR green (J). Data are mean ± SEM. N=5-12 mice/group. *P < 0.05; **P < 0.01; ***P < 0.001 vs. 0 hour; ##P < 0.01 vs. 2 or 24 hours. Student t test (A-D, F-J) and One-way ANOVA (E) were used. cfDNA, cell free DNA; Cit-H3, citrullinated histone H3; Dox, doxorubicin; IL, interleukin; MPO, myeloperoxidase; WBC, white blood cells.

Figure 3. Doxorubicin activates inflammasomes and causes macrophage pyroptosis. WT
BMMs were left untreated or primed with LPS for 3 hours, then treated with various doxorubicin
concentrations for 16 hours. IL-1β (A), LDH (B), and ATP (C) in the conditioned media were
measured by ELISA, the cytotoxicity detection Kit or ATP detection Kit, respectively. (D-F) WT
BMMs from Asc-citrine mice were primed with 100 ng/ml LPS for 3 hours and treated or not with
15 μM nigericin for 30 minutes or 10 μM doxorubicin for 16 hours. (G) Non-primed cells were
also treated with 10 μM doxorubicin for 16 hours. Scale bar: 50 μm. ASC specks were visualized
under fluorescence microscopy and quantified using imageJ. (H) Quantitative data. Data are
mean ± SEM from experimental triplicates and are representative of at least 2 independent
experiments. ***P < 0.001 vs. untreated- or LPS-treated cultures. One-way ANOVA. ASC,
apoptosis-associated speck-like protein containing a CARD; ATP, adenosine triphosphate; Dox,
doxorubicin; IL-1β, interleukin-1β; LDH, lactate dehydrogenase; LPS, lipopolysaccharide; WT,
wild-type.

Figure 4. Doxorubicin activates inflammasome –dependent and –independent pathways in macrophages. WT BMMs were left untreated or primed with LPS for 3 hours, then treated with various doxorubicin concentrations for 16 hours. Whole cell lysates were analyzed by immunoblotting. Blots are representative of at least 3 independent experiments. AIM2, absent in melanoma 2; cCasp, cleaved caspase; cGSDM, cleaved gasdermin; LPS, lipopolysaccharide; Dox, doxorubicin; WT, wild-type.

Figure 5. Doxorubicin activates inflammasome –dependent and –independent pathways in neutrophils. WT mouse bone marrow neutrophils were left untreated or primed with LPS for 3 hours, then treated with various doxorubicin concentrations for 16 hours. (A) Whole cell lysates were analyzed by immunoblotting. Blots are representative of at least 3 independent experiments. IL-1β (B) and LDH (C) in the conditioned media were measured by ELISA and by the cytotoxicity detection Kit, respectively. Data are mean ± SEM from experimental triplicates and are representative of at least 2 independent experiments. ***P < 0.001 vs. LPS; ##P < 0.01, ###P < 0.001 vs. untreated cultures. One-way ANOVA was used. cCasp, cleaved caspase; cGSDM, cleaved gasdermin; IL-1β, interleukin-1β LDH, lactate dehydrogenase; LPS, lipopolysaccharide; Dox, doxorubicin.

Figure 6. Doxorubicin causes NETosis in vitro. (A) WT mouse bone marrow neutrophils were left untreated or treated with 10 μM doxorubicin for 16 hours. Cit-H3 and MPO were analyzed by immunofluorescence. Scale bar: 50 μm. Images are representative of at least 3 independent experiments. (B) Neutrophils were left untreated or primed with LPS for 3 hours, then treated with 10 μM doxorubicin for 16 hours. cfDNA in the conditioned medium was extracted and quantified. (C) Neutrophils were left untreated or primed with LPS for 3 hours, then treated with 10 μM doxorubicin and/or DNase I for 16 hours. IL-1β in the conditioned media were measured by ELISA. Data are mean ± SEM from experimental triplicates and are representative of at least 2 independent experiments. *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$ vs. LPS; #$P < 0.05$, ###$P < 0.001$ vs. LPS + Dox. One-way ANOVA was used. Dox, doxorubicin; cfDNA, cell free DNA; Cit-H3, citrullinated histone H3; MPO, myeloperoxidase.

Figure 7. AIM2 and NLRP3 inflammasomes are involved in bone-damaging effects of doxorubicin. WT, *Aim2⁻/⁻*, *Nlrp3⁻/⁻*, *Aim2⁻/⁻;Nlrp3⁻/⁻* or *Casp1⁻/⁻* BMMs (A, B) and neutrophils (C, D) were left untreated or primed with LPS for 3 hours, then exposed or not to 10 µM doxorubicin for 16 hours. IL-1β (A, C) and LDH (B, D) in the conditioned media were measured by ELISA and by the cytotoxicity detection Kit, respectively. (E-G) Femurs from male mice were analyzed by VivaCT before (baseline) and 4 weeks after a single intraperitoneal injection of 5 mg/kg doxorubicin. (E) Cross sections of 3D reconstructions. Scale bar: 500 µm. (F) BV/TV changes. (G) BMD changes. (H, I) Femurs harvested from different genotypes male mice were analyzed 4 weeks after a single intraperitoneal injection of vehicle or doxorubicin. Specimens were stained for TRAP activity. (H) N.Oc/BS. (I) Oc.S/BS. Data are mean ± SEM from experimental triplicates and are representative of at least 2 independent experiments (A-D); n=5-8 mice/group (E-I). Data are mean ± SEM. *P < 0.05; **P < 0.01; ***P < 0.001 vs. control, LPS or vehicle; ##P < 0.01, ###P < 0.001 vs. other genotypes or WT treated with Dox. Two-way ANOVA (A-D, H-I) and One-way ANOVA (F-G) were used. AIM2, absent in melanoma 2; BMD bone mineral density; BV/TV, bone volume/total volume; casp1, caspase-1; Cont, control; Dox, doxorubicin; IL-1β, interleukin-1β;

745   LDH, lactate dehydrogenase; LPS, lipopolysaccharide; N.Oc/BS, OC number/bone surface;
746   Oc.S/BS; OC, osteoclast; WT, wild-type.