Insertion sequence driven evolution of the microbiota revealed by a novel metagenomics

pipeline

Joshua M. Kirsch[1], Andrew J. Hryckowian[2, 3], and Breck A. Duerkop[1]*

[1]Department of Immunology and Microbiology, University of Colorado - Anschutz Medical

Campus, School of Medicine, Aurora, Colorado, USA

[2]Department of Medicine, Division of Gastroenterology and Hepatology, University of Wisconsin

School of Medicine and Public Health, Madison, Wisconsin, USA

[3]Department of Medical Microbiology & Immunology, University of Wisconsin School of

Medicine and Public Health, Madison, Wisconsin, USA

*Correspondence: Breck A. Duerkop breck.duerkop@cuanschutz.edu

Abstract

Insertion sequence (IS) elements are mobile genetic elements in bacterial genomes that support

adaptation. We developed a database of IS elements coupled to a computational pipeline that

identifies IS element insertions in the microbiota. We discovered that diverse IS elements insert

into the genomes of intestinal bacteria regardless of human host lifestyle. These insertions

targeted bacterial accessory genes that aid in their adaptation to unique environmental

conditions. Using IS element expansion in *Bacteroides*, we show that IS element activity leads

to insertion "hot spots" in accessory genes. We show that IS element insertions are stable and

can be transferred between humans. Extreme environmental perturbations force IS element

insertions to fall out of the microbiota and many fail to rebound following homeostasis. Our work

shows that IS elements drive bacterial genome diversification within the microbiota and

establishes a framework for understanding how strain level variation within the microbiota

impacts human health.

Introduction

Bacteria in the intestinal microbiota are under constant evolutionary pressure to adapt.

One way that bacteria can rapidly evolve their genomes is through the activity of mobile genetic

elements, including bacteriophages, plasmids, DNA inversions, and transposable elements (1-

3). Genetic rearrangements have a strong impact on bacterial fitness, influencing diverse

phenotypes including virulence, antibiotic resistance, interbacterial competition, and secondary

metabolism. For example, pathogenicity islands can distribute virulence traits to previously non-

pathogenic bacteria and DNA inversions can alter the expression of antibiotic resistance genes

in the microbiota (4, 5). Additionally, novel genetic traits such as metabolism and virulence

genes are acquired by intestinal bacteria through plasmid exchange, bacteriophage lysogeny,

and conjugative transposition (6, 7). Thus, mobile elements actively drive bacterial adaptation

and evolution in the intestine.

Insertion sequence (IS) elements are small (~1 kB) simple transposons that are

common to bacterial genomes (8). IS elements are capable of mobilizing themselves by the

action of an encoded transposase that recognizes sequence-specific inverted repeats at the

boundaries of the element. IS element transposases are diverse (9), and include enzymes with

a DD(E/D) motif that support "copy-and-paste" and "cut-and-paste" mechanisms or those with

HUH nuclease chemistry which transpose as ssDNA molecules at the replication fork (10-12).

Gene expression and genome fidelity can be modulated by IS element activity in a variety of

ways, including IS element insertions into protein coding regions inactivating genes, insertions

into intergenic regions forming strong hybrid promoters that increase gene expression, or

recombination with other IS element sequences resulting in large deletions (13-16).

Additionally, IS elements can rapidly expand in bacterial genomes when commensal bacteria

transition to become pathogens (15, 17, 18). Thus, IS elements have a profound impact on the

evolution of bacterial genomes and the physiological traits of bacteria (19).

Despite a deep knowledge of fundamental IS element biology from diverse bacteria, we

have a rather rudimentary understanding of how these elements function in polymicrobial

communities. We believe this knowledge is critical for deciphering how bacterial communities

evolve ecosystems such as the human microbiota. A barrier to studying IS elements in such

complex environments results from imperfect methodologies for measuring *in situ* IS element

dynamics. This stems from the poor recovery of multi-copy genes with repetitive sequences by

short-read assemblers, leading to fragmented assemblies where IS elements are either absent

or become break-points between contigs (20). Therefore, assembly-level analyses of IS

elements from metagenomic datasets are often underpowered. Despite these limitations,

previous culture-based studies show that IS elements are active in the human intestine.

Longitudinal sampling of *Bacteroides fragilis* isolates revealed the gain and loss of IS element

transposases (21). Furthermore, IS elements were shown to rapidly expand in copy number in

the genome of *Escherichia coli* during intestinal colonization of mice, driving increased virulence

in a mouse model of Chron's disease (22, 23). Recently, longitudinal intestinal metagenomic

samples from a single individual were sequenced using long DNA fragment partitioning,

producing higher quality assemblies compared to short-read methodologies. This analysis

revealed that *Bacteroides caccae* gained and lost numerous IS insertions over the course of

sampling (20). Finally, we previously utilized a targeted IS element sequencing approach to

identify novel insertions of the IS element IS256 in intestinal *Enterococcus faecium* populations

isolated from an individual undergoing multiple antibiotic treatment (19). Specific antibiotics

treatment were associated with increased IS256 abundances and enrichment of IS256

insertions in genes related to antibiotic resistance and virulence, suggesting that antibiotic

treatment drives IS-mediated pathoadaptation in the human intestine.

In this work, we developed an open-source IS element database that greatly expands

the diversity of IS elements compared to current databases. We built a computational pipeline

that utilizes this database to find IS element insertions in a variety of public metagenomic

datasets. Our analysis reveals widespread IS element insertions and expansions within the

human microbiota. We show that IS element activity is transferable between individuals and are

stable over extended timescales. Distinct families of IS elements are favored for insertional

activity within the microbiota and these IS elements preferentially insert into specific classes of

genes. These genes are specific to distinct microbial taxa with an overrepresentation within the

Bacteroidia and *Clostridia* and are reproducible across discrete metagenomic datasets. Gene

classes targeted by IS elements are primarily metabolic, cell surface, and mobile genetic

element accessory genes. Using an *in vitro* ISOSDB412 (IS4351) expansion model, we confirm

that identical and related accessory genes are preferentially targeted within both *Bacteroides*

thetaiotaomicron and *Bacteroides fragilis*. Finally, we show that the stability of IS element

insertions is lost following antibiotic perturbation and diet intervention, and following microbiota

recovery after such alterations, new IS element abundances and insertion site locations arise.

Together, this work establishes a framework for studying IS elements within the context of the

microbiota and is the first step toward understanding how these mobile elements contribute to

the function of the microbiota during human health.

Results

ISOSDB: a comprehensive open-source IS element database

To help facilitate the study of IS elements in bacterial genomes and metagenomic

datasets, we sought to build an updated and exhaustive IS element database that can serve as

an open-source tool for the scientific community. Prior to our development of this database and

the analysis workflow, the main repository for IS elements was the ISFinder database (24).

While ISFinder has served as a gold-standard for the systematic naming of IS elements and

their identification in single isolate genomes (25), it is underpowered for high-throughput

genomics. Namely, ISFinder lacks integration into downstream tools that can facilitate in-depth

analyses of IS elements, has accessibility issues in that the entirety of the database cannot be

manually downloaded for personal use, and it relies solely on manual curation and submission

of new IS elements, decreasing the speed at which new IS elements can be disseminated to the

scientific community. These are serious drawbacks in the current genomic age, where the

identification of IS elements from complex datasets such as metagenomes would greatly

improve our understanding of their function in affecting complex bacterial traits from diverse

environments.

To address the need for an unrestricted open-access IS element database to support

genomic research, we built the Insertion Sequence Open-Source Database (ISOSDB), which

consists of IS elements identified from 39,878 complete bacterial genomes and 4,497

metagenome assembled genomes (MAGs) (26).These IS elements were identified using

OASIS, a rigorously tested IS identification tool that allows for the high-throughput analysis of

multiple genomes and contigs (27). IS elements were considered valid and included in ISOSDB

if there were at least two copies of the IS element in a single genome and it was flanked by

terminal inverted repeats (IRs). Redundant IS elements were deduplicated at 95% nucleotide

identity. The resulting set of IS elements totaled 22,804 distinct IS elements, an almost five-fold

excess to the ISFinder database (Fig. 1A). We identified transposase amino acid homologs, in

addition to nucleotide homology over the whole sequence of the IS elements, in the ISFinder

database (Fig. 1B). 92.6% of the transposases included in ISOSDB had homologs in ISFinder,

but only 37.7% of IS elements in the ISOSDB had nucleotide homologs in ISFinder. This

demonstrates that the ISOSDB has an abundance of novel IS element nucleotide sequence

diversity. The ISOSDB also has a wide range of transposases representing multiple IS families

(Fig. 1C). In summary, the ISOSDB is an expansive freely available database of IS elements

that contains substantially improved IS element diversity compared to current IS element

databases.

Development of an IS element detection pipeline that uses ISOSDB

IS elements are known to drive genomic diversity in almost every bacterial species (9,

28, 29). Considering the importance of IS elements to bacterial genome evolution, a systematic

method to identify novel IS insertions in complex microbial communities has not been previously

developed. This has precluded the study of IS element biology in the microbiota at the

population scale. To obtain a deeper understanding of IS element biology in the microbiota, we

developed the pseudoR pipeline that utilizes ISOSDB to identify IS element insertions in

previously assembled genomic sequences. pseudoR is named for its ability to find IS element

inactivated pseudogenes and is implemented in the R programming language. This pipeline is

built for fragmented, incomplete assemblies such as fecal metagenomic assemblies, and has

ample multi-threading support to speed up analyses of deeply sequenced samples. This

approach was inspired by previous tools developed for the analysis of transposon insertions

from eukaryotic (30) and bacterial genomes (31, 32). These tools either require *a priori*

knowledge of the target site duplications which is often not consistent for IS element insertions

(33), use read pairing to infer IS element insertions which limits the data available to verify IS

insertions, or require a matched reference assembly.

The backbone of the pseudoR pipeline is a split read approach. First, reads are mapped

against assembled contigs and unmapped reads are binned as separate read files. These

unmapped reads are aligned against a database of IS element terminal ends (150 bp on either

end of an IS element) compiled from ISOSDB (Fig. S1). If an unmapped read has an IS element

termini, the termini is trimmed and the remaining read is remapped against the assembled

contigs. To ensure that insertions are genuine, a minimum depth of four reads is required and

these reads must include at least one left and one right termini.

To compare between samples and multiple disparate assemblies, we built two modes

(multi and single) into pseudoR. Multi mode takes each sample and maps its reads against its

own assembly. These reads are then subsequently mapped against a deduplicated gene

database built from all assemblies in the dataset. Single mode was developed to better analyze

time series datasets, where a single assembly can be used for multiple samples (such as one

from the start of the time series) which is key for analyzing complex assemblies from multiple

individuals and timepoints.

IS element insertions are widespread in the healthy human intestinal microbiota

We first utilized the pseudoR pipeline to identify IS element insertions in healthy human

fecal metagenomes (Fig. 2). To gain a comprehensive view of IS insertions in this context, we

analyzed data from three geographically disparate studies: a survey of healthy Italian adults

ranging in age from 30-105 years old (ITA) (34), healthy individuals from a Japanese colorectal

cancer study (JPN) (35), and healthy rural Madagascan adults (MDG) (36). Each study showed

evidence of widespread IS insertion heterogeneity (Fig. 2A-B). Some individuals had few

detectable IS element insertions while others had over 100 unique IS element insertions (Fig.

2B). We used relative IS depth (calculated by dividing IS depth by the sum of the IS depth and

the depth of the reads that map to the insertion site without the insertion present) as a metric to

report the abundance of IS alleles. Most insertions had a relative IS depth above 10% and

below 100%, demonstrating that the inserted allele exists in equilibrium with the wild type allele

(Fig. 2A). To determine the extent of IS element insertions within the microbiota, we determined

the bacterial taxa that underwent extensive IS element insertional activity (Fig. 2C, Fig. S2A). IS

element insertions were abundant in the *Bacteroidia* and *Clostridia* and insertions in these

classes were found in almost every individual. These two classes make up the majority of the

healthy intestinal microbiota (37) and the genera *Bacteroides*, *Phocaeicola*, *Ruminococcus*, and

Blautia accounted for the majority of IS element insertion diversity (Fig. S2A). While *Bacteroidia*

IS element insertions were present across all studies, the IS elements underlying these

insertions varied between studies (Fig. 2D). Unique IS element types for the *Bacteroidia*

included ISOSDB412 (IS30 family) insertions for ITA individuals, ISOSDB18121 (IS1380 family)

insertions for JPN individuals, and ISOSDB33 (IS1182 family) insertions for MDG individuals

(Fig. 2D). Insertions in the *Bacteroidia* from JPN and ITA individuals were formed from shared

IS elements, including ISOSDB4584 (ISL3 family), ISOSDB634 (IS982 family), and ISOSDB445

(IS66 family), whereas the IS element diversity in the *Bacteroidia* from MDG individuals was

mostly unique consisting of ISOSDB250 (IS66 family), ISOSDB426 (IS630 family), and

ISOSDB45 (IS256 family) insertions. This suggests that regional differences in genomic makeup

of the *Bacteroidia* contributes to differences in IS element content or activity that drives insertion

events.

IS element insertion types among the *Clostridia* were found across all three geographic

locations, with many more IS elements shared between JPN and ITA individuals (Fig. 2D).

Interestingly, the Clostridia shared three IS family types (IS982, IS66, and IS30) with the

Bacteroidia, yet the specific IS family members active were assigned to different ISOSBD

groups indicating they are unique IS elements (Fig 2D). We conclude from this data that

although the IS element sequence space of these two classes of bacteria are unique, they

share IS element features at the IS family level. This suggests that certain IS element families

may be more promiscuous across diverse intestinal bacteria.

MDG individuals, despite having similar numbers of IS element insertions among their

bacterial communities (Fig. 2B), had relatively low abundances of IS elements from *Bacteroides*,

Phocaeicola, Blautia, and *Bifidobacterium* and much higher abundance of IS elements

associated with pathobionts including *Escherichia* and *Prevotella* species (Fig. S2A) (38-41).

This is supported by the presence of ISOSDB45 (IS256 family) insertions that are associated

with bacteria of pathogenic potential (Fig. 2D) (19, 42-44).

IS elements can insert into coding sequences or intergenic regions of the genome which

can inactivate genes or influence the expression of adjacent genes, respectively (28). To

measure IS element insertions in intragenic versus intergenic sites we analyzed the precise

positions of all IS elements identified across ITA, JPN, and MDG individuals by comparing the

number of intragenic insertions to intergenic insertions on a per individual basis. We found that

specific IS elements had preferences for intragenic or intergenic insertions (Fig. 2E).

ISOSDB1159 (IS3 family), ISOSDB805 (IS3 family), ISOSDB178 (ISAS1 family), and

ISOSDB237 (IS4 family) preferentially inserted into intergenic loci, while ISOSDB18121 (IS1380

family), ISOSDB445 (IS66 family), and ISOSDB412 (IS30 family) inserted more readily into

intragenic loci. These data indicate that some IS elements are suited to diversify genomes

through mutation whereas others may preferentially insert adjacent to coding sequences to

drive phenotypic diversity by altering gene expression.

IS elements frequently insert into accessory genes important for bacterial adaptation

Having identified numerous IS element insertions within predicted coding sequences, we

analyzed ORFs carrying IS element insertions, from here on called iORFs, in the ITA, JPN, and

MDG individuals (Fig. 3). A deduplicated gene database of all predicted ORFs from the

metagenomes from each study was assessed for iORF's. iORF's with insertions from only a

single individual were about 10-fold more abundant as iORFs with insertions from multiple

individuals (Fig. 3A). This demonstrates that IS element insertions per individual have a high

degree of inter-individual insertion site variation. We next performed functional annotation of the

iORF's and found five broad functional categories that were shared between individuals and

encompassed 20.7% of iORFs, on average (Fig. 3B). An abundance of iORFs were annotated

as *susC, susD*, or *tonB* receptor genes in ITA and JPN individuals. These genes encode high-

affinity substrate-uptake receptors for carbohydrates and cofactors (45, 46). *susC-D* homologs

are frequently involved in the uptake of polysaccharides used for metabolism, with *Bacteroidia*

species containing many different *susC-D* homologs (45). Mobilome genes, including

transposases, integrases, prophage elements, and mobile element defense genes, such as

restriction modification genes, contained insertions in almost every individual across all studies.

IS element insertions in transposase and integrase genes suggests mobile element competition

within bacterial genomes. Genes involved in exopolysaccharide biogenesis, such as LPS

modifying enzymes and glycosyltransferases involved in cell wall synthesis were consistent IS

element insertion targets in all three studies. Cell wall and membrane modifications are crucial

for *in vivo* survival to avoid host immunity and during competition with other bacteria (47).

Finally, antibiotic resistance genes frequently contained IS element insertions, including a high

prevalence of insertions in *tetQ* and *ermF* in JPN and ITA individuals. This indicates that the

carriage of antibiotic resistance genes in the microbiota is metabolically unfavorable and IS

elements may modulate their activity in the absence of antibiotic pressure. Insertion

abundances were enriched in these functional categories when comparing insertion

abundances within all predicted protein coding sequences (Fig. 3C, Fig. S2B). Together, these

data show that accessory metabolic genes and genes involved in antibiotic resistance and

genomic plasticity are common IS insertion targets and indicates that the intestinal microbiota

modulates the function of these genes to adapt to their environment.

We next asked if certain iORFs were shared between individuals from the three studies.

Although infrequent, a few iORFs were shared. 14% and 37% of ITA and JPN individuals,

respectively, had IS insertions in the gene *ermF*, a macrolide resistance gene, and 3.2% and

9.8% of ITA and JPN individuals, respectively, had IS insertions in *KAP*, a P-loop NTPase

predicted to be involved in phage defense (Fig. 3D) (48, 49). 7.9% and 6.1% ITA and MDG

individuals, respectively, shared IS insertions in *shufflon*, a gene encoding an invertase that

regulates pilin phase variation (Fig. 3D) (50). We found that iORF functional classes were

conserved across diverse intestinal bacteria (Fig. 3E) yet the IS element types responsible for

these insertions differed among the various classes of bacteria (Fig. 3F). This demonstrates that

although IS elements target similar genes within disparate bacteria, the types of IS elements

that promote these diversification events are specific to certain classes of bacteria.

Multiple iORF's were classified as transposases (Fig. 3B) and examples of IS elements

inserting in other IS elements has been previously reported (51). In order to understand the

dynamics of IS insertions in IS elements, we classified both the transposase iORF and the IS

element forming the insertion by IS family (Fig. 3G). We found multiple examples of both self-

targeting (an IS element inserting into a closely-related IS element) and orthologous-targeting

(an IS element inserting into an unrelated IS element). We found that some IS families such as

ISL3, IS4, IS110, and IS256 experienced promiscuous insertion events. Interestingly, IS

elements from some of these families create composite transposons capable of mobilizing

cargo genes including metabolic and antibiotic resistance genes (52-55). This observation is

consistent with an enrichment of iORFs belonging to both metabolic and antibiotic resistance

accessory genes and suggests IS element self- and orthologous-targeting could be a

mechanism to control IS element mobilization of genetic traits when these events are

unfavorable to the bacterial host.

IS elements provide mutational diversity to Bacteroides species

Recently a genome wide mutagenesis study of the *Bacteroides thetaiotaomicron*

chromosome identified genes that support adaptation to environmental pressures, including

antibiotics, bile acids, and carbon sources (56). We used this dataset to evaluate whether we

could identify similar fitness determinants as iORFs within the natural *Bacteroides* population

within the intestine. We searched our iORF database for homologs to these fitness-associated

ORFs from *B. thetaiotaomicron* and found multiple closely related iORFs that likely influence

Bacteroides fitness both positively and negatively (Fig. S3A). Examples include iORFs that

provide fitness advantages during growth in the presence of the antibiotics doxycycline and

vancomycin and during hyaluronic acid and glucosamine consumption (Fig. S3A). Interestingly

we found iORFs associated with bacterial fitness during exposure to the antipsychotic

medication chlorpromazine (Fig S3A). Antipsychotic compounds are connected to alterations in

the intestinal microbiota and directly inhibit bacterial growth as well as potentiate antibiotic

resistance (57-60). Additionally, we found multiple IS element insertions in homologs of the *B.*

thetaiotaomicron susC gene BT1119 (Fig. S3B). Inactivation of this *susC* gene decreases

fitness during growth on galacturonic acid, a downstream product of pectin degradation (56).

Our *in silico* analysis shows that the intestinal *Bacteroidia* experience large-scale IS

element mobilization into discrete groups of genes including *susC-D/tonB* carbohydrate

utilization genes as well as genes involved in exopolysaccharide synthesis and mobile genetic

element interactions (Fig 2, Fig. 3). These data indicate that these genes are "hotspots" for IS

element insertion. To test if the activation of an IS element leads to predictable insertion into

these specific classes of genes, we transformed a multi-copy plasmid carrying an *ermF* cassette

disrupted with a copy of ISOSDB412 (IS30 family member IS4351) into *B. thetaiotaomicron* VPI-

5482 (*Bt*) and *Bacteroides fragilis* NCTC 9343 (*Bf*), both which lack native ISOSDB412.

Following the growth of these transformants, we isolated their genomic DNA and performed IS-

Seq, a technique that enriches for ISOSDB412 amplicons during Illumina DNA sequencing

library construction and read mapping to identify specific ISOSDB412 insertion locations (19).

We found that both *Bt* and *Bf* strains carrying the ISOSDB412 plasmid obtained numerous

ISOSDB412 insertions throughout their genomes compared to controls that lacked the

ISOSDB412 plasmid construct (Fig. 4A, Fig. 4D). As predicted, ISOSDB412 insertions were

enriched in gene categories reflected from our findings from insertions from the human

microbiota (Fig. 4B, Fig. 4E). Specifically, for *Bt* there was an enrichment of ISOSDB412

insertions in *susC-D/tonB* and EPS biogenesis genes, whereas *Bf* acquired an abundance of

insertions in *susC-D/tonB* genes despite a significantly lower number of I ISOSDB412 insertion

events compared to *Bt* (Fig. S4A). Additionally, insertions were more frequently found in

intragenic loci compared to intergenic loci, confirming our findings from the intestinal microbiota

(Fig. 4C, Fig. 4F). These results demonstrate that IS elements can rapidly diversify *Bacteroidia*

genomes.

Our discovery that IS insertions target bacterial accessory genes suggests that these IS

insertions inactivate genes that are currently dispensable for survival in the intestine. To explore

this, we measured IS insertional dynamics in a nutrient-deplete environment with readily-

digestible glucose as the sole carbon source. We hypothesized that a lack of fiber and complex

carbon sources, high levels of glucose, and single sources of iron and nitrogen, would

incentivize higher rates of IS insertions to inactivate costly metabolic machinery needed for

competition in complex nutritional environments. We measured ISOSDB412 insertion events

from three *Bt* colonies after three sequential passages in either the complex media (BHIS) or

minimal media (MM) (Fig. 4G, Fig. 4H, Fig. 4I). We found that colonies passaged in MM had

substantially more IS4351 insertions compared to BHIS-passaged colonies. Numerous

insertions were found in the open reading frame BT3642, a Na^+-dependent transporter (Fig.

S4B) in MM colonies, but not in BHIS colonies. This gene has been shown to be detrimental for

growth in glucose minimal media (56). Next, we measured ISOSDB412 abundance in *Bt* cells

chronically infected with the *Crassvirales* DAC15 or DAC17 (Fig. 4J, 4K, 4L). *Crassvirales* are

bacteriophages that selectively infect *Bacteroides* species and are highly prevalent in the

human intestinal microbiota (61). These strains shed high levels of infectious phage particles

following chronic phage infection (Fig. S4C). While DAC15 and DAC17 share 99.24%

nucleotide identity over 98% of their genomes, DAC17 infection is associated with an increased

number of insertions genome-wide (Fig. 4J, Fig. 4K) and preferentially inserted into genes

involved in EPS biogenesis (Fig. 4L). Furthermore, almost all of the infected strains carried IS

insertions in the CPS3 locus (Fig. S4D), which is crucial for productive phage infection and likely

leads to phage resistance (62). Together, these results show that IS element diversification of

the *Bt* genome is more prevalent under nutrient-limited conditions and during infection with

intestinal-resident phages. This indicates that IS element activity supports the ability of *Bt* to

adapt when faced with fitness constraints.

Temporal monitoring of IS element activity shows that IS insertions are maintained over

time.

Our assessment of IS element insertions in members of the intestinal microbiota

revealed that IS insertions in specific genetic loci coexist with the wild type allele lacking the IS

element (Fig. 2). This indicates that IS element insertions in target loci represent only a

subpopulation of bacterial cells within the community and that these insertions are maintained in

equilibrium with bacterial cells harboring uninserted versions of the loci. To understand the

temporal longevity of IS element insertions in the microbiota, we analyzed longitudinal fecal

metagenomic samples from healthy individuals (63, 64). Using a set of two individuals whose

fecal samples were collected over the course of 76 and 91 weeks (63), we measured the IS

element landscape of their microbiotas using pseudoR (Fig. 5A & Fig. S5). We also analyzed

longitudinal samples from an additional 10 individuals from a separate study (64) (Fig. S6). For

each individual, we used the assemblies from the first timepoint (indicated as week 0 in Fig. 5A)

as the reference for the pseudoR pipeline which was then compared to all other time points to

assess 1) the maintenance of the IS elements, and 2) the acquisition of new IS elements that

were not found at the starting timepoint. We found that most IS element insertions are

maintained for periods extending to up the endpoint of the study for each individual and that

some of these insertions frequently went in and out of detection (Fig. 5A, Fig. S5-6). New

insertions not present at week 0 arose at almost every timepoint, with a variable number of

insertions per timepoint. Some new insertions were maintained for extended periods of time

while others were only detected at the initial timepoint of week 0. These results demonstrate

that IS elements are both stably carried in the microbiota and are actively forming new

insertions.

Next, we analyzed whether the host bacterium's abundance could account for the

detection of new IS elements (Fig. S7A). Increasing levels of an IS element's host bacterium

would generate higher sequencing read depth and more detection power for new IS insertions.

For every insertion not present at week 0, we compared the insertion site's read depth (a

marker of host abundance) to the maximum insertion site depth of all earlier timepoints (prior to

the detection of the insertion). Insertions with insertion site depths equal to or less than 200% of

the previous maximum insertion site depth were considered to arise from new insertional activity

(Fig. S7B). We found that the majority of new insertions could be accounted for as active IS

element insertions and not increased bacterial host abundance (Fig. S7C).

After week 0, multiple insertions were never detected again. To confirm that these

insertions were lost from the population rather than being undetectable due to low sequencing

depth, we compared the WT allele depth of insertions that were present at week 0 and absent at

every other timepoint. Insertions where the WT allele depth reached at least 50% of the week 0

WT allele depth at a later timepoint were classified as "rebounders" and those that did not were

classified as "non-rebounders". We found that, for every insertion that fit the above criteria, the

WT allele depth rebounded to at least 50% of the week 0 WT allele depth (S7D). This suggests

that these insertions were lost due to competition from the WT allele, rather than a change in

read depth.

Insertions across the time series included diverse IS element families and were

frequently associated with the *Clostridia* and *Bacilli* (Fig. 5B, S5-6). Additionally, we observed

that the rate of new insertions per week for each individual ranged on average from around 1 to

10 new insertions and that the maintenance, loss, and gain of IS elements tracked with common

bacterial accessory genes that we established to be "hot spots" for IS element insertions (Fig.

5C, Fig. 5D).

IS elements are efficiently transferred and maintained within new individuals

Having discovered that individual microbiotas have unique patterns of IS element

insertions, we wanted to understand how the human host influences the maintenance, loss, and

gain of IS elements within the microbiota. To test this, we used pseudoR to profile the IS

element insertions during a fecal microbiota transplantation (FMT) where the microbiotas of

FMT donors and recipients were longitudinally sampled before and after fecal transplantation

(63). We compared the rates of IS element insertions and their maintenance in both the donors

and recipients using the assembly of the donor's transplanted microbiota as a reference. Donor-

derived communities harbored both insertion site DNA with and without an IS element insertions

and were stably maintained over one year in the recipients that received the FMT (Fig. 6A & Fig.

S8). Insertions present prior to transplantation in recipients were lost, but a minority returned at

later timepoints. Recipients maintained significantly less IS element insertions that were present

at week 0 compared to the donors (Fig. 6C), but both donor and recipients had similar rates of

new insertions (Fig. 6D). To understand if IS element insertional activity was similar between

host and recipient, we compared the pattern of new shared insertions in donors and recipients

at all timepoints post transplantation (Fig. 6B & Fig. S9). We found multiple instances of the

same newly detected insertion arising in both donor and recipients at the same timepoint. These

results demonstrate that complex IS element insertions can be stably maintained in new hosts

for extended periods of time and that IS element mobilization occurs at similar rates and

patterns in new hosts.

Antibiotics drive widespread loss of IS insertions within the microbiota

Considering IS element insertions were found to persist for prolonged periods of time

within the microbiota and were portable between individuals, we wanted to know what would

happen to IS element diversity if a stable community was disrupted. To test this, we analyzed a

microbiota dataset for IS element composition before, during, and after broad spectrum

antibiotic treatment (65). The metagenome of the microbiota preceding antibiotic treatment was

used as the reference for IS element comparison using pseudoR. To account for IS element

activity within the microbiota prior to antibiotic treatment and to exclude the possibility that new

insertions were caused by large shifts in rebounding bacterial abundances following antibiotic

treatment, new insertions had to meet the selection criteria established where the insertion site

lacking an IS element must have a sequencing depth less than or equal to 200% compared to

the reference insertion site prior to antibiotic treatment. As expected, antibiotic treatment caused

a substantial loss of IS element diversity, which was not fully restored after treatment (Fig. 7A &

Fig. S10A). The starkest example was sample ERAS10. Extensive IS element diversity was

present at week 0 in ERAS10 before antibiotic treatment and by at week 0.6 there were no

detectable IS element signatures nor did any of the original IS elements return by week 26 post

antibiotic therapy. In contrast, samples ERAS1 and ERAS12 both had a complete loss of pre-

antibiotic treatment IS element insertions immediately following treatment, yet a considerable

number of pre-antibiotic treatment IS element insertions returned at later timepoints. Across all

samples, IS element insertions in *Clostridia* genomes showed a dramatic loss of original IS

element signatures, similar to the pattern seen in ERAS10 (Fig. 7C). To better understand the

mechanism behind the loss of week 0 insertions, the wild type allele depth of week 0 insertions

that were never detected again was plotted against time (Fig. 7B & Fig. S10B). Again, lost

insertions were classified as either non-rebounders, where the insertion site DNA sequencing

depth at later timepoints never reached 50% of the pre-antibiotic treatment insertion site read

depth, or rebounders, where the insertion site DNA sequencing depth at later timepoints was

greater than or equal to 50% of the pre-antibiotic treatment insertion site read depth.

Rebounders are indicative of the wild type allele out-competing the IS inserted allele, while non-

rebounders are indicative of complete loss of the host bacterium. Examples of both non-

rebounding and rebounding alleles were found for every individual. Finally, we compared the

rate of maintenance of pre-antibiotic treatment insertions in individuals treated with antibiotics

and those who were not. Antibiotic treatment was associated with a significant decrease in the

ability of the microbiota to maintain IS insertions (Fig. 7D), due to both competition from wild

type alleles and loss of host bacteria.

IS element insertion into susC-D/tonB loci is associated with diet intervention.

Our analysis has identified *susC-D/tonB* genes as major targets for IS element

insertions, suggesting that IS elements can modulate carbohydrate metabolism in the

Bacteroidia (Fig. 3), and we showed in *B. thetaiotaomicron* that nutrient stress can promote

these insertions events into predictable metabolic loci (Fig. 4). With these data in hand, we

suspected that a radical diet intervention in humans would lead to a shift in IS insertions in

metabolic-associated genes and particularly *susC-D/tonB*. To test this hypothesis, we analyzed

the IS element insertions among the intestinal microbiota of obese individuals before, during,

and after a rigorous, one year-long diet intervention (66). Participants in this study received a

low-calorie formula-based diet for 12 weeks, followed by a 6-week period of phasing out the

formula diet for solid food without changing total caloric intake, followed by a 7-week phase

where caloric intake was increased while preventing weight gain (67). Similar to antibiotic

treatment (Fig. 7), we found that diet change was associated with frequent loss of the starting

set of IS insertions at week 0 (Fig. 8A, 8E & Fig. S11A). Loss of the target insertion site without

the IS element was assessed in addition to the loss of IS element from these sites. In contrast to

antibiotic treatment, read depth at target insertion sites lacking an IS element usually

rebounded, suggesting that loss of week 0 insertions is due to competition from wild type alleles

rather than loss of the host bacterium (Fig. 8B & Fig. S11B). *susC-D/tonB* genes import specific

carbohydrates and we suspected that these genes were the targets for IS-mediated insertion

events during diet intervention, thus we reasoned that the presence of certain carbohydrates

would increase the fitness burden of an insertion in that carbohydrate's cognate *susC-D/tonB*

gene, promoting loss of the IS element from this location. Alternatively, low abundance of a

carbohydrate would favor insertions in the carbohydrate's cognate *susC-D/tonB* genes to

reduce the fitness burden of maintaining the gene's expression. We found that ancestral *susC-*

D/tonB insertions were frequently lost during the course of the diet study, new insertions in

susC-D/tonB genes arose after the diet intervention ended (Fig. 8C). In many cases, insertions

arose in the *susC-D/tonB* homolog BT1119 which we identified as an insertion site of preference

(Fig. 8C, 8D). Together, these results demonstrate that diet is associated with IS insertion

patterns that specify alterations in insertion events in *susC-D/tonB* genes.

Discussion

IS elements are fundamental to the evolution of bacterial genomes. They likely play a

key role in the assembly of polymicrobial communities and influence how bacterial members of

such communities interact. Despite this, IS element diversity and function within polymicrobial

communities like the intestinal microbiota are poorly defined and understudied. Here, we built a

novel computational pipeline for identifying IS element insertions within complex metagenomic

DNA sequences. This pipeline, which we call pseudoR, was built using a custom open-source

IS element database (ISOSDB), which vastly improves nucleotide diversity compared to current

state-of-the-art databases for IS element analysis. Using pseudoR, we found that IS elements

contribute to high levels of bacterial genomic diversification within the microbiota, are

maintained as integrated elements at specific loci for extended periods of time, can be efficiently

transferred between individuals, and experience both losses and gains within bacterial

populations following perturbation (i.e., antibiotics treatment or diet alteration). Together, these

results demonstrate that IS elements are important genetic elements that can dictate genotype

diversity within the microbiota. The functional consequences of IS element activity within the

microbiota are just beginning to be understood and should prove to have a strong impact on the

microbiota and its relationship to human health.

Using previously published metagenomic datasets, we showed that the majority of open

reading frames containing IS element insertions (iORFs) are unique and not shared between

different individuals (Fig. 3A). This is similar to the observation that single nucleotide

polymorphism (SNP) diversity in the human intestinal microbiota is more likely to be unique than

shared (68). Our work expands upon this idea and suggests that each person's "fingerprint" of

microbial diversity as it relates to genotype is also strongly influenced by IS element activity.

While iORFs are unique within individuals, the functional classes of genes that constitute

these iORFs are consistent between individuals (Fig. 3B). iORFs were frequently annotated

within five broad categories: *susC-D/tonB*, mobilome primally composed of prophage elements

and transposase genes, EPS genes, antibiotic resistance genes, and genes implicated in anti-

MGE systems. Genes belonging to these categories are frequently found as accessory genes

and are often non-essential. Interestingly, genes regulated by DNA inversions including those

regulated by phase variation fall into similar functional categories compared to those targeted by

IS element insertions (69, 70). This suggests that insertions and inversions control gene

expression of the microbiota to position bacteria for optimal adaptation to changing

environmental conditions. Additionally, *susC-D/tonB* receptors are often targets of nucleotide

evolution in the human intestine possibly to help gain new abilities based on substrate

availability (21). In support of this, a SusC epitope was shown to active T cells to promote an

anti-inflammatory IL-10 response in healthy individuals and a proinflammatory IL-17A response

in people with Chron's disease (71). This suggests that IS elements could influence the

immunomodulatory capacity of the microbiota depending on whether they are inserted within

genetic loci such as *susC* homologs. Carbohydrate and other metabolite availability can have a

strong influence on the virulence of intestinal of pathogens (40, 72, 73). Since *susC-D*/*tonB* are

most prevalent among the *Bacteroidia* which dominate the intestinal microbiota and participate

in the utilization of carbohydrate substrates, IS element insertions in *susC-D*/*tonB* genes could

effectively shape the nutrient landscape of the intestine promoting or subverting pathogen

fitness. In support of this theory, we showed that IS elements frequently target *susC* genes

homologous to BT1119 from *B. thetaiotaomicron* implicated in the transport galacturonic acid.

Galacturonic acid inhibits the activation of the type III secretion system of enterohemorrhagic *E.*

coli and reduces its virulence (74), thus the inactivation of BT1119 by IS element insertion may

result in regulated control of intestinal galacturonic acid concentrations that can provide

protection against invading pathogenic *E. coli*.

Often, the microbiome of urban individuals is less-well suited to degrade diverse

polysaccharides compared to rural individuals (75). Here, we found that urban individuals (those

in the ITA and JPN cohorts) have more extensive modulation of the microbiome through higher

rates of IS insertions in *susC-D/tonB* genes, compared to individuals living in rural settings

(MDG cohort) (Fig. 3E). This could partly be explained by the higher ITA and JPN individuals

have almost exclusively *Bacteroidia* IS insertions in *Bacteroides* and *Phocaeicola* species, while

MDG individuals have *Bacteroidia* insertions in *Prevotella* species. However, we expect that the

varied diet of MDG individuals, which heavily includes wild meat and plants (76), partially

selects for a diverse set of functioning *susC-D/tonB* receptors. In support of this finding, diet

intervention modulated IS insertions in *susC-D/tonB* genes (Fig. 8). Further work to better

characterize the IS insertional landscape in individuals with diverse lifestyles, such as the Hadza

hunter gatherers (77), can better demonstrate how IS elements impact nutrient acquisition.

Frequent IS element inactivation of diverse MGE systems has been observed and was

hypothesized to help protect host bacteria from the toxic effects of these systems or to allow

incoming MGEs to subvert these systems (33). We found that IS element insertions into other

MGEs was frequent within the microbiota and that these insertions often co-exist with the

insertion site lacking the IS element. This suggests that IS elements exist in an equilibrium

between the inserted and non-inserted state at the population scale to maintain effective host

defense against MGEs and when dispensable, relax the cost of maintaining these systems

through inactivation.

In addition to metabolic and MGE system targeting by IS elements, another common

iORF was *ermF*, a macrolide resistance gene that acts by methylating the 23S rRNA subunit to

inhibit protein translation (78). We previously found that gentamicin resistance operons are

frequently interrupted by IS elements during enterococcal infection (19). We presume that in the

absence of antibiotic pressure, bacteria can dampen their effort toward maintaining high-level

antibiotic resistance at the population level to favor their fitness. However, since these are

heterogenous populations that co-exist between antibiotic resistance genes carrying or lacking

IS element insertions, we hypothesize that IS element insertions in *ermF* and other antibiotic

resistance genes serves as a mechanism to maintain a low level of population-wide antibiotic

resistance, while reducing the burden of expressing the resistance gene. Such a "hedge betting"

strategy would be advantageous upon antibiotic exposure where the population would not

require genetic mutation to overcome antibiotic pressure but instead regain antibiotic resistance

thorough selection of variants within the population.

Prior to this study, it was not known whether IS element insertions could be stably

maintained in bacterial genomes within host associated microbiotas. Having this knowledge

provides new insights into the genomic composition of the microbiota and their capacity to

harbor stable subpopulations of genetic variants that are prime for selection and expansion

under changing environmental conditions. This could account for the rapid adaptability of the

microbiota and its resiliency to recover from perturbation. We found that specific IS element

insertions were detectable for just under two years, demonstrating that IS insertions within

specific genetic loci are often not sanitized from the population. Additionally, new previously

undetected IS elements were detectable and maintained for extended periods of time. Such

insertions were likely due to new transpositions, rather than bacterial host immigration within the

microbiota or changes in preexisting host bacterial abundances. These findings are analogous

to the discovery of SNP conservation within the microbiota (79). Together, our results show that

in addition to active and widespread transposition, IS element insertions in the microbiota are

maintained for prolonged periods of time which may support rapid responses to bacterial

community assembly under selective pressure and/or fluctuations in environmental resources.

In addition to longevity, we also established that IS elements can be stably transferred

between individuals. Donor-derived bacterial communities have been shown to persist in

recipients following FMT (80-82). Bacterial diversity frequently rebounds after antibiotic

treatment (65, 83), but how bacterial communities assemble at the strain level is more varied

and includes widespread loss of mutations and enrichment of others (84). We found that IS

insertions were efficiently transplanted between individuals (Fig. 6), but were readily lost after

antibiotic treatment and diet intervention (Fig. 7-8). We suspect that IS element insertions impart

only a minor impact on bacterial fitness under steady state conditions when selective pressure

is low, but become burdensome during strong selective pressure when environmental

conditions are dramatically altered. Furthermore, the dysbiosis that is caused by antibiotic

treatment and diet fluctuations can eventually resolve to the previous taxonomic composition

(65, 66, 85, 86) . We show that in contrast to higher level taxonomic restoration of the

microbiota following perturbation, IS element insertions are irreversibly altered following strong

selective pressure (Fig. 7-8). This indicates that many IS element insertions within the

population come with fitness tradeoffs that can be detrimental when the appropriate selection is

applied. It would also be fair to speculate that some IS element insertions would be beneficial

and become dominate and fixed within the population depending on the type of environmental

pressure applied. Our results suggest that IS elements can be viewed as strain-level features

that are more variable than taxonomic-level diversity following perturbation.

Antibiotic exposure enriches IS insertions in *E. faecium* during a chronic human infection

and antibiotics can stimulate IS element activity in *Staphylococcus aureus* (19, 87). In each of

these scenarios this activity was linked to the IS element IS256, which is strongly associated

with pathogenic and hospital-associated bacteria (18). In contrast, we found that during

antibiotic treatment the intestinal microbiota undergoes extreme reductions in IS element

diversity among the commensal bacterial population. This suggests that IS element activation

by antibiotic pressure could be dependent on the specific family of IS element or certain

pathogenic bacteria have developed IS elements as a tool to mutagenize their genomes in

response to antibiotic stress. The latter strategy could help support pathogen fitness in a context

dependent manner (i.e. infection), for which there is little to no benefit to commensal bacteria

during intestinal colonization.

In summary, the development of a contemporary and stringently curated IS element

database, combined with a streamlined computational approach to identify IS elements from

metagenomic data, has revealed novel insights into how IS elements shape the genomes of the

microbiota and sets the stage for exploring how specific IS element variants in the microbiota

influence human health. However, a few limitations to the tools and study exist. First, delineation

of IS element insertion sties between closely related strains is imperfect, thus insertions

between very closely related strains may group together and fail to resolve. Work from the

intestinal microbiota has found that most species are represented by one dominant strain and

that these strains are stable over long periods (88), suggesting that the insertion sites that we

measured are from subpopulations originating from single strains. Second, we do not

understand how IS element insertions influences the fitness of the microbiota. We can begin to

overcome this problem by assembling and assessing Tn-Seq libraries of diverse intestinal

bacteria under conditions relevant to the intestinal environment and in animal colonization

experiments. Coupling such studies with pseudoR analyses of microbiota datasets would begin

to help determine what IS element insertion sites are beneficial or detrimental to commensal

bacterial genome evolution and would aid in the identification of IS element families that are

important for such evolutionary events in the intestine. Third, the pseudoR pipeline relies on

assemblies containing some level of wild type allele being measured in comparison to an IS

element insertion at that same allele. This limits the detection of IS elements that assemble as

part of the reference contig. Conceivably, an inverse method to find wild type alleles in IS-allele

assemblies can be built and be implemented into a future release of pseudoR. Fourth, the single

reference mode of pseudoR excludes assemblies from later timepoints to prevent over-

assembly and focus the analysis on the initial timepoint. This hinders our understanding of how

IS elements are impacting the microbiota not present at the initial timepoints.

Materials and Methods

Creation of the ISOSDB

All genomes classified as "complete" in NCBI Assembly were downloaded on

05/08/2023 using bit v1.8.57 (89). IS elements were identified using OASIS (27). This program

uses annotated transposases as points for sequence extension and comparison to find

complete IS elements. Identified IS elements were filtered using the following criteria: 1) multiple

(minimum of 2 copies) complete copies of the element must be present in at least one genome

and 2) the elements must contain identified inverted repeat sequences. To supplement these

genomes, we also identified IS elements in hybrid-assembled MAGs from a large human

intestinal microbiota study (26). These metagenomes were annotated with Prokka v1.14.6 (90),

processed as described above, and then combined with the IS elements identified from the

genomes from NCBI Assembly. Redundant IS elements in the final database were deduplicated

using CD-HIT-EST (91) with a 95% sequence cutoff and a word length of 9. IS element

homologs in ISFinder were found using either blastn (complete elements) with a minimum e-

value of 0.000001 and a minimum length of alignment of 224 bp or blastp (transposases) with a

minimum e-value of 0.000001 (92). Phylogenetic analysis of transposase genes was performed

by multiple sequence alignment using MAFFT v7.520 (93). A phylogenetic tree was created

using FastTree with default settings (94) and visualized using iTOL (95). The ISOSDB and

pseudoR pipeline is freely available at https://github.com/joshuakirsch/pseudoR. Each IS

element in this database is given a unique numeric identifier.

pseudoR pipeline

To characterize IS elements and their insertion sites in metagenomic DNA sequences

we built the pseudoR pipeline. This tool relies on deeply-sequenced short read Illumina data

and can be used for both metagenomic and single isolate genome data. We found a minimum

of ~20 million paired-end reads is sufficient for identifying IS elements using pseudoR. Data

input into the pipeline are deduplicated reads and assembled contigs. For this study, reads were

downloaded from NCBI SRA (96) using fasterq-dump v2.11.0 from the studies previously

mentioned (34-36, 63, 65, 66). Study accession numbers are provided in Table S1.

Duplications, quality trimming, and read decontamination was performed using programs that

are part of the BBTools software suite (97). PCR duplicated reads were removed with

clumpify.sh with the flag "subs=0". Adapter and quality trimming was performed with bbduk.sh

and human, mouse, and phi29 contamination was removed using bbsplit.sh. Cleaned and

deduplicated reads were assembled into contigs using MEGAHIT v1.2.7 using the "meta-large"

preset (-k-max 127 -k-min 27 -k-step 10) (98). Contigs input into the pipeline include those equal

to and greater than 1 kB. When using the multiple reference mode, assemblies were combined

and ORFs were predicted using pprodigal.py (a parallelizable wrapper of prodigal (99)) and

deduplicated using dedupe.sh. If using the single reference mode, only the reference assembly

is annotated as described above. Nucleotide sequences of the predicted ORFs were used for

read mapping using Bowtie2 v2.4.5 (100). To identify IS insertions within the microbiota from

longitudinal samples from the same individual, we built the single reference mode, where the

individual's starting timepoint assembly is used to find novel IS element insertions. This

approach reduces the chances that meaningful information is lost during clustering of multiple

assemblies from multiple timepoints. In addition to time-series, the single reference mode can

be used for evolution studies where a single population is compared between different

treatments. Multiple reference mode is used for comparing the insertional patterns between

multiple individuals. To identity IS elements in multiple reference mode, reads were aligned to

each sample's own assembly using Bowtie2. Any mapped read, including discordantly mapped

reads, were considered positive hits. Unmapped reads were collected and aligned to a

database of IS termini using blastn and then filtered using a custom R script. This filtering script

ensures that the IS termini is properly positioned on the read such that the remainder of the

read is outside of the IS element. The IS termini are removed and the trimmed reads were then

re-mapped to the sample's assembly as unpaired reads using Bowtie2. Following this, the

original mapped reads and IS-termini trimmed reads are mapped against the deduplicated gene

database. For single reference mode, reads from every sample are mapped to the provided

reference assembly and the remainder of the steps are the same. Following read mapping

completion, a custom R script is used to identify DNA insertion sites with read mapping from

both the left and right terminus of the IS element at most 20 bp away from one another and this

script compiles read abundances for these regions. Mosdepth v0.3.3 (101) is used to determine

insertion site sequencing depth for the loci described above. Seqkit v2.2.0 (102), samtools v1.6

(103), R, ggplot2, tidyr, and MetBrewer were used for various computational tasks. Scripts to

reproduce all data analysis presented here are available in the GitHub repository above.

iORFs assoated with B. thetaiotaomicron

B. thetaiotaomicron VPI-5482 coding sequences were downloaded from

fit.genomics.lbl.gov. Homologs to iORFs were found using blastp with a maximum e-value of

0.00005 and a minimum identity of 75%. Hits to multiple related fitness determinants were

filtered to include the representative with the strongest fitness value.

Bacteria

Bacteroides thetaiotaomicron VPI-5482, *Bacteroides thetaiotaomicron* VPI-5482 CPS3,

and *Bacteroides fragilis* NCTC 9343 were cultured as previously described (104). Briefly, these

strains were grown in brain-heart infusion (BHI) broth (Becton Dickinson, Franklin Lakes, NJ)

supplemented with 1 g/L cysteine, 5% w/v $NaHCO_3$, and 5 mg/L hemin (BHIS), Varel-Bryant

minimal media (105) supplemented with 28 mM glucose, or *Bacteroides* phage recovery

medium (BPRM) (106) at 37^o C in a Coy Type A vinyl anaerobic chamber in an atmosphere of

5% hydrogen, 20% carbon dioxide, and balanced nitrogen (Coy Lab Products, Grass Lake, MI).

For growth on solid-media we used BHI agar supplemented with 10% defibrinated calf blood

(Colorado Serum Company, Denver, CO). *Escherichia coli* S17-1 was grown in Lennox L broth

(Fisher Scientific, Hampton, NH) with aeration at 37^o C. Tetracycline was used at 2 µg/mL,

gentamicin was used at 200 µg/mL, and ampicillin was used at 100 µg/mL.

DNA manipulation and cloning

Primers and plasmids used in this study are listed in Table S2. All plasmid constructs

were made using Gibson assembly master mix (New England Biolabs, Ipswich, MA).

Assembled constructs were electroporated into electrocompetent *E. coli* S17-1. These cells

were prepared by washing cell pellets three times in ice-cold electroporation buffer (0.5 M

sucrose, 10% glycerol). The resulting transformants were mated with *B. thetaiotaomicron* or *B.*

fragilis by mixing 500 µL of stationary phase *E. coli* S17-1 and 500 µL of stationary phase

Bacteroides on BHI blood agar overnight at 37^o C under aerobic conditions. Transconjugants

were selected by scraping the bacterial lawns into BHI and plating serial dilutions on BHI blood

agar supplemented with tetracycline and gentamicin, and incubating at 37°C anaerobically for

48 hours. The ISOSDB412 (IS4351) DNA sequence construct was purchased from Twist

Biosciences (San Francisco, CA). pB006 (Addgene plasmid #182320), a gift from Dr. Lei Dai,

was used as a shuttle vector and purchased from Addgene (Watertown, MA) (107). The *tetM*

cassette was cloned from pCIE-*tetM* (108) and the *ermF* cassette was cloned from pG10K, a

gift from Dr. Janina Lewis (Addgene plasmid #191377) and purchased from Addgene (109).

DNA extraction and IS-Seq

Genomic DNA was extracted from 1.5 mL cultures of *Bacteroides* strains using the

Zymobionics DNA Miniprep kit (Zymo Research, Irvine, CA) and used as input for IS-Seq as

described previously (19). Briefly, NGS libraries were produced using the Illumina DNA Prep Kit

(Illumina, San Diego, CA) and amplified using ISOSDB412 IS-Seq Step1 and p7 primers for 13

cycles using Q5 Master Mix (New England Biolabs). The products of this reaction were

amplified with IS-Seq Step2 and p7 primers for 9 cycles using Q5 Master Mix and sequenced on

a Novaseq 6000 at Novogene (Sacramento, CA). R1 reads were binned and trimmed of the

ISOSDB412 terminus and adapters using cutadapt v1.1.18 (110). To remove PCR duplicates,

R2 pairs of R1 reads that contained the ISOSDB412 terminus were binned and deduplicated

using dedupe.sh from the BBTools suite with the flag minidentity=100. Deduplication of R2

reads ensures that each amplified read pair is a unique molecule, while maintaining read depth

information of the R1 (IS-amplified) read. This set of deduplicated R2 reads was re-paired to

trimmed R1 reads with reformat.sh from the BBTools suite and these pairs were aligned to the

host genome using Bowtie2. R1 read depth at each genomic position was calculated using

bedtools v2.30.0 (111) IS seq reads can be found at the European Nucleotide Archive under

study accession number PRJEB66483.

B. thetaiotaomicron passaging experiment

B. thetaiotaomicron was grown overnight in BHIS and then subcultured 1:100 into either

Varel-Bryant minimal media or BHIS containing antibiotic and grown overnight. The next day,

bacteria cultured in BHIS were subcultured 1:100 into BHIS and bacteria cultured in Varel-

Bryant minimal media cultured were subcultured 1:100 into Varel-Bryant minimal media with

antibiotic. After overnight growth, this passaging was repeated once more. Genomic DNA was

isolated and the final passage was sequenced as described above.

B. thetaiotamicron phage infection

B. thetaiotamicron CPS3 (a mutant of *B. thetaiotamicron* VPI-5482 expressing only the

CPS3 capsule) (62) (112) was transformed with pB6T-ermF-IS4351 and grown overnight in

Bacteroides phage recovery medium (BPRM) with antibiotic (106). Overnight cultures were

diluted to OD_{600} of 1.0 in phage buffer (62) and was plated on BPRM agar containing

tetracycline and 1×10^8 PFU/mL of either DAC15 or DAC17. Dilutions of resuspended overnight

culture were also plated on BPRM containing tetracycline. Plated cultures on DAC15 agar

media were incubated at 37^o C for 48 hours and individual colonies were picked and patched

onto fresh BPRM plates with DAC15. This process was repeated. Cultures plated on DAC17

agar and agar without phage were similarly passaged. Passaged colonies were cultured into

BPRM broth without phage and incubated for 48 hours at 37^o C. Genomic DNA was isolated

and sequenced as described above.

Enumeration of infectious phage particles

90 µL of supernatant from liquid cultures of passaged colonies were mixed with 10 µL

chloroform (Fisher Scientific), centrifuged for 1 minute at 21,000 RCF at 25^o C, and the

supernatant was collected. Dilutions of this supernatant were made in phage buffer and 5 µL

spots of the dilutions were plated on a BPRM plate containing *B. thetaiotamicron* CPS3

embedded in 0.35% BPRM top agar, and incubated at 37^o C overnight.

Functional and taxonomic classification of contigs and ORFs

Metagenomic sequences with IS insertions were functionally annotated using eggNOG

mapper v2 (113). Orthologous groups were cross-validated using the COG database (114).

Antibiotic resistance genes were identified using CARD RGI 6.0.0 (115). ORFs and contigs

containing IS insertions were classified using Kraken2 v2.0.7 (116).

Acknowledgements

We would like to thank the members of the Duerkop lab for helpful discussions related to this

manuscript. We would like to thank A. Murat Eren and Alejandro Reyes Muñoz for critically

reading this manuscript and providing useful insights.

Funding

This work was supported by the National Institutes of Health grants R01AI141479 (B.A.D.),

T32AI052066 (J.M.K), F31AI169976 (J.M.K.), and R35GM150996 (A.J.H). The funders had no

role in the study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

Figure Legends

Figure 1. The ISOSDB: a database of diverse and complete IS elements. (A) Number of

unique IS element sequences in ISFinder and the ISOSDB. (B) Comparison of transposase

(Tnp amino acid) and nucleotide sequences of the full IS elements from the ISOSDB to the IS

elements from ISFinder. (C) Phylogenetic tree of the Tnp genes of IS elements in the ISOSDB.

Leaves are colored by Tnp family.

Figure 2. IS element insertions are abundant in intestinal bacteria. (A) Abundance of IS

elements from ITA, JPN, and MDG individuals. Each data point represents one new insertion

not found in the reference assembly. The y-axis is the relative IS depth percentage (IS depth/

(WT allele depth + IS depth)). (B) Number of insertions per sample shown in A. (C) IS element

insertions per bacterial class. The intensity of each bar is proportional to the number of IS

element insertions. (D). IS element insertions organized by IS family that from insertions in

Bacteroidia and *Clostridia* bacteria. The intensity of each bar is proportional to the number of IS

element insertions. (E) Preferential IS element insertion in either intergenic (left side, red) or

intragenic (right side, blue) loci for highly abundant IS elements (paired T test with FDR multiple

testing correction, maximum adjusted *p* value = 0.05).

Figure 3. IS element insertions are commonly found in bacterial accessory genes. (A)

Number of shared and unique iORFs within ITA, JPN, and MDG individuals. (B) Heatmap of IS

element insertions in different gene functional categories. (C) Statistically significant enrichment

of IS elements in the functional categories from B (Fisher's exact test with FDR multiple

comparison correction, ***$p < 10^{-25}$, **$p < 10^{-4}$, *$p < 0.05$). (D) Location and relative IS element

depth in shared iORFs. The location of the insertion (either above or below the gene

representation) is representative of the study. (E-F) Number of IS element insertions in iORFs in

the functional categories in B from different bacterial classes (E) and IS families (F). (G) Rate of

IS insertions in transposases. Rectangles in the red dashed line are self-targets and rectangles

outside of the red dashed line are non-synonymous-targets.

Figure 4. ISOSDB412 insertion in *Bacteroides* species replicates findings from intestinal

metagenomic data. (A and D) IS-Seq deep sequencing of ISOSDB412 insertions in *Bt* (A) and

Bf (D). (B and E) Functional enrichment of ISOSDB412 insertions in *Bt* (B) and *Bf* (E) (Fisher's

exact test, *p* < 0.06, ***p < 0.001). (C and F) ISOSDB412 insertions were counted in intergenic

and intragenic loci in *Bt* (C) and *Bf* (F) (paired T-test, *p* < 0.05). (G) IS-Seq of ISOSDB412

insertions of *Bt* passaged in either BHIS or MM. (H) Number of ISOSDB412 insertions in each

condition in G (paired T test). (I) Number of ISOSDB412 insertions in EPS biogenesis genes in

G (paired T test). (J) IS-Seq of ISOSDB412 insertions of *Bt* chronically infected with DAC15 or

DAC17. (K) Number of ISOSDB412 insertions in each condition in J (unpaired T test, *p* < 0.05).

(L) Number of ISOSDB412 insertions in EPS biogenesis genes in J (unpaired T test, *p* < 0.05).

Figure 5. IS element insertions are maintained in the intestinal microbiota for extended

time periods. (A) IS element insertions are depicted as rectangles, with the color intensity

proportional to the relative IS depth percentage. Each insertion is unique based on insertion

position and the same insertions are aligned vertically. Orange-colored IS element insertions

are present at the first timepoint (week 0), green IS element insertions are new insertions not

present at the initial timepoint, and blue IS element insertions are new insertions at timepoints

after their first appearance. (B) Number of IS element insertions per timepoint for IS families

(red) or taxonomic classes (blue). (C) Rate of new Is element insertions per week. (D) Dynamics

of IS element insertions over time for gene functional category iORFs. Data from every

individual was used for this figure.

Figure 6. FMT transfer of IS element insertions between donors and recipients. (A) Blue

rectangles are IS element insertions in donor samples and red rectangles are IS element

insertions in recipient samples. The same insertions in donor and recipients are aligned

vertically. Week 0 is the transplanted fecal material. (B) Heatmap showing new IS element

insertions that were detected 2 weeks after transplantation and are detected in both donor and

recipient samples. (C) Loss of IS element insertions detected in the transplanted fecal material

(week 0) from recipients (paired T test, **$p < 0.01$). (D) Similar rates of new IS element

insertions are found in both donor and recipients (unpaired T test).

Figure 7. Antibiotic treatment sanitizes IS element insertions from intestinal bacteria. (A)

Heatmap showing IS element insertion dynamics during antibiotic treatment. Only insertions in

alleles with less than or equal to 200% of the week 0 read depth are shown in this panel.

Horizontal black bars at the 0.6 Week timepoint are present in individuals who did not have an

available sample for this timepoint. (B) Rebounding and non-rebounding wild type alleles of IS

element insertions lost from week 0 during antibiotic treatment. (C) Clostridia bacteria rapidly

lose IS element insertions. Data from every individual was used for this figure panel. (D)

Antibiotic treatment significantly decreases the number of IS element insertions present

following week 0 (unpaired T test, *** $p < 0.005$).

Figure 8. Diet intervention alters IS element mediated *susCD/tonB* gene inactivation. (A)

Heatmap showing IS element insertion dynamics before, during, and after diet intervention. Only

insertions in alleles with less than or equal to 200% of the week 0 read depth are shown in this

panel. (B) Rebounding and non-rebounding wild type alleles of IS element insertions lost from

week 0 during diet intervention. (C) Dynamics of IS element insertions in *susCD/tonB* genes

during the timecourse. Data from every individual was used for this figure panel. (D) New

BT1119 IS element insertions arise after the diet intervention has ended. Wild type allele depth

is shown as a black line and IS element insertion allele depth is shown as a red bar. (E) Diet

intervention significantly decreases the number of IS element insertions present following week

0 (unpaired T test, ***$p < 0.005$).

Figure S1. Schematic of the pseudoR computational pipeline. Bacterial genomes are

sequenced using Illumina short-read DNA sequencing. These reads are assembled into contigs

and the reads (black rectangles) are mapped against the assembly contigs. Unmapped reads

(black and blue lines) are compared against the ISOSDB containing IS element termini. IS

element termini are trimmed from the reads and the reads are then are re-mapped to the

assembly contigs.

Figure S2. IS element insertions in diverse genera and functional classes. (A) Abundance

of IS element insertions for intestinal bacteria at the genera level. (B) Replotting of the data in

Fig. 3C to include the "other" category which encompasses all other genes containing IS

element insertions that were not the focus of this study.

Figure S3. IS element insertions in genes that affect *Bt* fitness. (A) Table of IS element hits

in genes involved in fitness in *Bt*. Genes whose fitness values < 0 are beneficial for fitness in the

specified condition and genes whose fitness > 0 are detrimental to fitness in the specified

condition (+++ fitness > 4, ++ fitness > 2, + fitness > 1, - - - fitness < -4, - - fitness < -2, - fitness

< -1). (B) Schematic of IS element insertions in *BT1119*.

Figure S4. ISOSDB412 drives IS element insertions during selective pressure. (A) IS

element insertions in *Bt* and *Bf* strains following passage in nutrient limited conditions (unpaired

T test, **$p < 0.01$) (B) Schematic of IS element insertions in BT3642. (C) Quantification of

infectious phage particles in the supernatant of *Bt* CPS3 strains chronically infected with DAC15

or DAC17. (D) Schematic of IS element insertions in the CPS3 locus.

Figure S5-6. Remainder of healthy individuals followed longitudinally (Related to Fig. 5A-

B). The same scale in 5A-B is used for these figures.

Figure S7. New IS element insertions are not associated with increased wild type allele

depth. (A) The percentage of new IS element insertions is shown on the Y axis and the X axis

is the percentage of the wild type allele depth from the first detection of each new insertion

divided by the maximum detected wild type allele depth of the timepoints prior to the first

detection timepoint. The red dashed line intersects the X axis at 200%. (B) Representation of

abundance-driven or transposition-driven allele measurements. (C) Significantly more IS

element insertions are predicted to arise from transposition compared to changes in host

abundance (***$p < 0.01$, paired T-test). (D) Wild type allele depth of IS element insertions that

were present at week 0 and were not detected at another timepoint. Each IS element insertion

is shown as a line. Red lines ("rebounders") represent IS element insertions where the wild type

allele depth at later timepoints reached at least 50% of the week 0 wild type allele depth. Black

lines ("non-rebounders") represent IS element insertions where the wild type allele depth at later

timepoints did not reached at least 50% of the week 0 wild type allele depth.

Figure S8. Remainder of individuals followed longitudinally after FMT (Related to Fig.

6A). Same scale in Fig. 6A is used in these figures.

Figure S9. Remainder of shared insertion plots (Related to Fig. 6B).

Figure S10. Remainder of antibiotic treated individual plots (Related to Fig. 7A and 7B).

Figure S11. Remainder of antibiotic treated individual plots (Related to Fig. 8A and 8B).

References

1.   Shkoporov AN, Clooney AG, Sutton TDS, Ryan FJ, Daly KM, Nolan JA, McDonnell SA,

Khokhlova EV, Draper LA, Forde A, Guerin E, Velayudhan V, Ross RP, Hill C. 2019.

The human gut virome is highly diverse, stable, and individual specific. Cell Host

Microbe 26:527-541.e5. 10.1016/j.chom.2019.09.009

2.   Forster SC, Liu J, Kumar N, Gulliver EL, Gould JA, Escobar-Zepeda A, Mkandawire T,

Pike LJ, Shao Y, Stares MD, Browne HP, Neville BA, Lawley TD. 2022. Strain-level

characterization of broad host range mobile genetic elements transferring antibiotic

resistance from the human microbiome. Nat Commun 13:1445. 10.1038/s41467-022-

29096-9

3.   Jiang X, Hall AB, Xavier RJ, Alm EJ. 2019. Comprehensive analysis of chromosomal

mobile genetic elements in the gut microbiome reveals phylum-level niche-adaptive

gene pools. PLoS One 14:e0223680. 10.1371/journal.pone.0223680

4.   Jiang X, Hall AB, Arthur TD, Plichta DR, Covington CT, Poyet M, Crothers J, Moses PL,

Tolonen AC, Vlamakis H, Alm EJ, Xavier RJ. 2019. Invertible promoters mediate

bacterial phase variation, antibiotic resistance, and host adaptation in the gut. Science

363:181-187. 10.1126/science.aau5238

5.   Desvaux M, Dalmasso G, Beyrouthy R, Barnich N, Delmas J, Bonnet R. 2020.

Pathogenicity factors of genomic islands in intestinal and extraintestinal *Escherichia coli*.

Front Microbiol 11:2065. 10.3389/fmicb.2020.02065

6.   Groussin M, Poyet M, Sistiaga A, Kearney SM, Moniz K, Noel M, Hooker J, Gibbons SM,

Segurel L, Froment A, Mohamed RS, Fezeu A, Juimo VA, Lafosse S, Tabe FE, Girard

C, Iqaluk D, Nguyen LTT, Shapiro BJ, Lehtimäki J, Ruokolainen L, Kettunen PP,

Vatanen T, Sigwazi S, Mabulla A, Domínguez-Rodrigo M, Nartey YA, Agyei-Nkansah A,

Duah A, Awuku YA, Valles KA, Asibey SO, Afihene MY, Roberts LR, Plymoth A,

Onyekwere CA, Summons RE, Xavier RJ, Alm EJ. 2021. Elevated rates of horizontal

gene transfer in the industrialized human microbiome. Cell 184:2053-2067.e18.

10.1016/j.cell.2021.02.052

7.   Brito IL, Yilmaz S, Huang K, Xu L, Jupiter SD, Jenkins AP, Naisilisili W, Tamminen M,

Smillie CS, Wortman JR, Birren BW, Xavier RJ, Blainey PC, Singh AK, Gevers D, Alm

EJ. 2016. Mobile genes in the human microbiome are structured from global to individual

scales. Nature 535:435-439. 10.1038/nature18927

8.   Aziz RK, Breitbart M, Edwards RA. 2010. Transposases are the most abundant, most

ubiquitous genes in nature. Nucleic Acids Res 38:4207-17. 10.1093/nar/gkq140

9.   Siguier P, Gourbeyre E, Chandler M. 2014. Bacterial insertion sequences: their genomic

impact and diversity. FEMS Microbiol Rev 38:865-91. 10.1111/1574-6976.12067

10.   He S, Corneloup A, Guynet C, Lavatine L, Caumont-Sarcos A, Siguier P, Marty B, Dyda

F, Chandler M, Ton Hoang B. 2015. The IS200/IS605 family and "peel and paste" single-

strand transposition mechanism. Microbiol Spectr 3. 10.1128/microbiolspec.MDNA3-

0039-2014

11.   Piegu B, Bire S, Arensburger P, Bigot Y. 2015. A survey of transposable element

classification systems--a call for a fundamental update to meet the challenge of their

diversity and complexity. Mol Phylogenet Evol 86:90-109. 10.1016/j.ympev.2015.03.009

12.   Kosek D, Hickman AB, Ghirlando R, He S, Dyda F. 2021. Structures of ISCth4

transpososomes reveal the role of asymmetry in copy-out/paste-in DNA transposition.

EMBO J 40:e105666. 10.15252/embj.2020105666

13.   Jellen-Ritter AS, Kern WV. 2001. Enhanced expression of the multidrug efflux pumps

AcrAB and AcrEF associated with insertion element transposition in *Escherichia coli*

mutants selected with a fluoroquinolone. Antimicrob Agents Chemother 45:1467-72.

10.1128/aac.45.5.1467-1472.2001

14.   Shankar N, Baghdayan AS, Gilmore MS. 2002. Modulation of virulence within a

pathogenicity island in vancomycin-resistant *Enterococcus faecalis*. Nature 417:746-50.

10.1038/nature00802

15.   Chain PS, Carniel E, Larimer FW, Lamerdin J, Stoutland PO, Regala WM, Georgescu

AM, Vergez LM, Land ML, Motin VL, Brubaker RR, Fowler J, Hinnebusch J, Marceau M,

Medigue C, Simonet M, Chenal-Francisque V, Souza B, Dacheux D, Elliott JM, Derbise

A, Hauser LJ, Garcia E. 2004. Insights into the evolution of *Yersinia pestis* through

whole-genome comparison with *Yersinia pseudotuberculosis*. Proc Natl Acad Sci U S A

101:13826-31. 10.1073/pnas.0404012101

16.   Manson JM, Hancock LE, Gilmore MS. 2010. Mechanism of chromosomal transfer of

Enterococcus faecalis pathogenicity island, capsule, antimicrobial resistance, and other

traits. Proc Natl Acad Sci U S A 107:12269-74. 10.1073/pnas.1000139107

17.   Hawkey J, Monk JM, Billman-Jacobe H, Palsson B, Holt KE. 2020. Impact of insertion

sequences on convergent evolution of *Shigella* species. PLoS Genet 16:e1008931.

10.1371/journal.pgen.1008931

18.   Leavis HL, Willems RJ, van Wamel WJ, Schuren FH, Caspers MP, Bonten MJ. 2007.

Insertion sequence-driven diversification creates a globally dispersed emerging

multiresistant subspecies of *E. faecium*. PLoS Pathog 3:e7.

10.1371/journal.ppat.0030007

19.   Kirsch JM, Ely S, Stellfox ME, Hullahalli K, Luong P, Palmer KL, Van Tyne D, Duerkop

BA. 2023. Targeted IS-element sequencing uncovers transposition dynamics during

selective pressure in enterococci. PLoS Pathog 19:e1011424.

10.1371/journal.ppat.1011424

20. Zlitni S, Bishara A, Moss EL, Tkachenko E, Kang JB, Culver RN, Andermann TM, Weng

Z, Wood C, Handy C, Ji HP, Batzoglou S, Bhatt AS. 2020. Strain-resolved microbiome

sequencing reveals mobile elements that drive bacterial competition on a clinical

timescale. Genome Med 12:50. 10.1186/s13073-020-00747-0

21. Zhao S, Lieberman TD, Poyet M, Kauffman KM, Gibbons SM, Groussin M, Xavier RJ,

Alm EJ. 2019. Adaptive evolution within gut microbiomes of healthy people. Cell Host

Microbe 25:656-667.e8. 10.1016/j.chom.2019.03.007

22. Elhenawy W, Tsai CN, Coombes BK. 2019. Host-specific adaptive diversification of

Crohn's disease-associated adherent-invasive *Escherichia coli*. Cell Host Microbe

25:301-312.e5. 10.1016/j.chom.2018.12.010

23. Barreto HC, Frazão N, Sousa A, Konrad A, Gordo I. 2020. Mutation accumulation and

horizontal gene transfer in *Escherichia coli* colonizing the gut of old mice. Commun

Integr Biol 13:89-96. 10.1080/19420889.2020.1783059

24. Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. 2006. ISfinder: the reference

centre for bacterial insertion sequences. Nucleic Acids Res 34:D32-6.

10.1093/nar/gkj014

25. Kichenaradja P, Siguier P, Pérochon J, Chandler M. 2010. ISbrowser: an extension of

ISfinder for visualizing insertion sequences in prokaryotic genomes. Nucleic Acids Res

38:D62-8. 10.1093/nar/gkp947

26.   Gounot JS, Chia M, Bertrand D, Saw WY, Ravikrishnan A, Low A, Ding Y, Ng AHQ, Tan

1021   LWL, Teo YY, Seedorf H, Nagarajan N. 2022. Genome-centric analysis of short and long

1022   read metagenomes reveals uncharacterized microbiome diversity in Southeast Asians.

1023   Nat Commun 13:6044. 10.1038/s41467-022-33782-z

1024   27.   Robinson DG, Lee MC, Marx CJ. 2012. OASIS: an automated program for global

1025   investigation of bacterial and archaeal insertion sequences. Nucleic Acids Res 40:e174.

1026   10.1093/nar/gks778

1027   28.   Tempel S, Bedo J, Talla E. 2022. From a large-scale genomic analysis of insertion

1028   sequences to insights into their regulatory roles in prokaryotes. BMC Genomics 23:451.

1029   10.1186/s12864-022-08678-3

1030   29.   Nzabarushimana E, Tang H. 2018. Insertion sequence elements-mediated structural

1031   variations in bacterial genomes. Mob DNA 9:29. 10.1186/s13100-018-0134-3

1032   30.   Quadrana L, Bortolini Silveira A, Mayhew GF, LeBlanc C, Martienssen RA, Jeddeloh JA,

1033   Colot V. 2016. The *Arabidopsis thaliana* mobilome and its impact at the species level.

1034   Elife 5:e15716. 10.7554/eLife.15716

1035   31.   Hawkey J, Hamidian M, Wick RR, Edwards DJ, Billman-Jacobe H, Hall RM, Holt KE.

1036   2015. ISMapper: identifying transposase insertion sites in bacterial genomes from short

1037   read sequence data. BMC Genomics 16:667. 10.1186/s12864-015-1860-2

1038   32.   Durrant MG, Li MM, Siranosian BA, Montgomery SB, Bhatt AS. 2020. A bioinformatic

1039   analysis of integrative mobile genetic elements highlights their role in bacterial

1040   adaptation. Cell Host Microbe 27:140-153.e9. 10.1016/j.chom.2019.10.022

33. Sheng Y, Wang H, Ou Y, Wu Y, Ding W, Tao M, Lin S, Deng Z, Bai L, Kang Q. 2023.

Insertion sequence transposition inactivates CRISPR-Cas immunity. Nat Commun

14:4366. 10.1038/s41467-023-39964-7

34. Rampelli S, Soverini M, D'Amico F, Barone M, Tavella T, Monti D, Capri M, Astolfi A,

Brigidi P, Biagi E, Franceschi C, Turroni S, Candela M. 2020. Shotgun metagenomics of

gut microbiota in humans with up to extreme longevity and the increasing role of

xenobiotic degradation. mSystems 5. 10.1128/mSystems.00124-20

35. Yachida S, Mizutani S, Shiroma H, Shiba S, Nakajima T, Sakamoto T, Watanabe H,

Masuda K, Nishimoto Y, Kubo M, Hosoda F, Rokutan H, Matsumoto M, Takamaru H,

Yamada M, Matsuda T, Iwasaki M, Yamaji T, Yachida T, Soga T, Kurokawa K, Toyoda

A, Ogura Y, Hayashi T, Hatakeyama M, Nakagama H, Saito Y, Fukuda S, Shibata T,

Yamada T. 2019. Metagenomic and metabolomic analyses reveal distinct stage-specific

phenotypes of the gut microbiota in colorectal cancer. Nat Med 25:968-976.

10.1038/s41591-019-0458-7

36. Pasolli E, Asnicar F, Manara S, Zolfo M, Karcher N, Armanini F, Beghini F, Manghi P,

Tett A, Ghensi P, Collado MC, Rice BL, DuLong C, Morgan XC, Golden CD, Quince C,

Huttenhower C, Segata N. 2019. Extensive unexplored human microbiome diversity

revealed by over 150,000 genomes from metagenomes spanning age, geography, and

lifestyle. Cell 176:649-662.e20. 10.1016/j.cell.2019.01.001

37. Rinninella E, Raoul P, Cintoni M, Franceschi F, Miggiano GAD, Gasbarrini A, Mele MC.

2019. What is the healthy gut microbiota composition? A changing ecosystem across

age, environment, diet, and diseases. Microorganisms 7.

10.3390/microorganisms7010014

38.   Alpizar-Rodriguez D, Lesker TR, Gronow A, Gilbert B, Raemy E, Lamacchia C, Gabay

1065   C, Finckh A, Strowig T. 2019. *Prevotella copri* in individuals at risk for rheumatoid

1066   arthritis. Ann Rheum Dis 78:590-593. 10.1136/annrheumdis-2018-214514

1067   39.   Scher JU, Sczesnak A, Longman RS, Segata N, Ubeda C, Bielski C, Rostron T,

1068   Cerundolo V, Pamer EG, Abramson SB, Huttenhower C, Littman DR. 2013. Expansion

1069   of intestinal *Prevotella copri* correlates with enhanced susceptibility to arthritis. Elife

1070   2:e01202. 10.7554/eLife.01202

1071   40.   Hughes ER, Winter MG, Duerkop BA, Spiga L, Furtado de Carvalho T, Zhu W, Gillis CC,

1072   Buttner L, Smoot MP, Behrendt CL, Cherry S, Santos RL, Hooper LV, Winter SE. 2017.

1073   Microbial respiration and formate oxidation as metabolic signatures of inflammation-

1074   associated dysbiosis. Cell Host Microbe 21:208-219. 10.1016/j.chom.2017.01.005

1075   41.   Conway T, Cohen PS. 2015. Commensal and pathogenic *Escherichia coli* metabolism in

1076   the gut. Microbiol Spectr 3. 10.1128/microbiolspec.MBP-0006-2014

1077   42.   Lebreton F, van Schaik W, McGuire AM, Godfrey P, Griggs A, Mazumdar V, Corander J,

1078   Cheng L, Saif S, Young S, Zeng Q, Wortman J, Birren B, Willems RJ, Earl AM, Gilmore

1079   MS. 2013. Emergence of epidemic multidrug-resistant *Enterococcus faecium* from

1080   animal and commensal strains. mBio 4. 10.1128/mBio.00534-13

1081   43.   McEvoy CR, Tsuji B, Gao W, Seemann T, Porter JL, Doig K, Ngo D, Howden BP,

1082   Stinear TP. 2013. Decreased vancomycin susceptibility in *Staphylococcus aureus*

1083   caused by IS256 tempering of WalKR expression. Antimicrob Agents Chemother

1084   57:3240-9. 10.1128/AAC.00279-13

1085   44.   Kozitskaya S, Cho SH, Dietrich K, Marre R, Naber K, Ziebuhr W. 2004. The bacterial

1086   insertion sequence element IS256 occurs preferentially in nosocomial *Staphylococcus*

epidermidis isolates: association with biofilm formation and resistance to

aminoglycosides. Infect Immun 72:1210-5. 10.1128/IAI.72.2.1210-1215.2004

45.   Martens EC, Koropatkin NM, Smith TJ, Gordon JI. 2009. Complex glycan catabolism by

the human gut microbiota: the Bacteroidetes Sus-like paradigm. J Biol Chem 284:24673-

7. 10.1074/jbc.R109.022848

46.   Noinaj N, Guillier M, Barnard TJ, Buchanan SK. 2010. TonB-dependent transporters:

regulation, structure, and function. Annu Rev Microbiol 64:43-60.

10.1146/annurev.micro.112408.134247

47.   Yadav AK, Espaillat A, Cava F. 2018. Bacterial Strategies to Preserve Cell Wall Integrity

Against Environmental Threats. Front Microbiol 9:2064. 10.3389/fmicb.2018.02064

48.   Vassallo CN, Doering CR, Littlehale ML, Teodoro GIC, Laub MT. 2022. A functional

selection reveals previously undetected anti-phage defence systems in the *E. coli*

pangenome. Nat Microbiol 7:1568-1579. 10.1038/s41564-022-01219-4

49.   Aravind L, Iyer LM, Leipe DD, Koonin EV. 2004. A novel family of P-loop NTPases with

an unusual phyletic distribution and transmembrane segments inserted within the

NTPase domain. Genome Biol 5:R30. 10.1186/gb-2004-5-5-r30

50.   Komano T. 1999. Shufflons: multiple inversion systems and integrons. Annu Rev Genet

33:171-91. 10.1146/annurev.genet.33.1.171

51.   Blattner FR, Plunkett G, 3rd, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J,

Glasner JD, Rode CK, Mayhew GF, Gregor J, Davis NW, Kirkpatrick HA, Goeden MA,

Rose DJ, Mau B, Shao Y. 1997. The complete genome sequence of *Escherichia coli* K-

12. Science 277:1453-62. 10.1126/science.277.5331.1453

52.   Lyon BR, Gillespie MT, Skurray RA. 1987. Detection and characterization of IS256, an

insertion sequence in *Staphylococcus aureus*. J Gen Microbiol 133:3031-8.

10.1099/00221287-133-11-3031

53.   Quintiliani R, Jr., Courvalin P. 1996. Characterization of Tn1547, a composite

transposon flanked by the IS16 and IS256-like elements, that confers vancomycin

resistance in *Enterococcus faecalis* BM4281. Gene 172:1-8. 10.1016/0378-

1119(96)00110-2

54.   Dufour A, Rince A, Uguen P, Le Pennec JP. 2000. IS1675, a novel lactococcal insertion

element, forms a transposon-like structure including the lacticin 481 lantibiotic operon. J

Bacteriol 182:5600-5. 10.1128/JB.182.19.5600-5605.2000

55.   Udaondo Z, Abram KZ, Kothari A, Jun S-R. 2021. Insertion sequences associated with

antibiotic resistance genes in *Enterococcus* isolates from an inpatient with prolonged

bacteremia. bioRxiv doi:10.1101/2021.07.22.453447. 10.1101/2021.07.22.453447

56.   Liu H, Shiver AL, Price MN, Carlson HK, Trotter VV, Chen Y, Escalante V, Ray J, Hern

KE, Petzold CJ, Turnbaugh PJ, Huang KC, Arkin AP, Deutschbauer AM. 2021.

Functional genetics of human gut commensal *Bacteroides thetaiotaomicron* reveals

metabolic requirements for growth across environments. Cell Rep 34:108789.

10.1016/j.celrep.2021.108789

57.   Flowers SA, Evans SJ, Ward KM, McInnis MG, Ellingrod VL. 2017. Interaction between

atypical antipsychotics and the gut microbiome in a bipolar disease cohort.

Pharmacotherapy 37:261-267. 10.1002/phar.1890

58. Maier L, Pruteanu M, Kuhn M, Zeller G, Telzerow A, Anderson EE, Brochado AR,

Fernandez KC, Dose H, Mori H, Patil KR, Bork P, Typas A. 2018. Extensive impact of

non-antibiotic drugs on human gut bacteria. Nature 555:623-628. 10.1038/nature25979

59. Ait Chait Y, Mottawea W, Tompkins TA, Hammami R. 2020. Unravelling the

antimicrobial action of antidepressants on gut commensal microbes. Sci Rep 10:17878.

10.1038/s41598-020-74934-9

60. Kyono Y, Ellezian L, Hu Y, Eliadis K, Moy J, Hirsch EB, Federle MJ, Flowers SA. 2022.

The atypical antipsychotic quetiapine promotes multiple antibiotic resistance in

Escherichia coli. J Bacteriol 204:e0010222. 10.1128/jb.00102-22

61. Dutilh BE, Cassman N, McNair K, Sanchez SE, Silva GG, Boling L, Barr JJ, Speth DR,

Seguritan V, Aziz RK, Felts B, Dinsdale EA, Mokili JL, Edwards RA. 2014. A highly

abundant bacteriophage discovered in the unknown sequences of human faecal

metagenomes. Nat Commun 5:4498. 10.1038/ncomms5498

62. Hryckowian AJ, Merrill BD, Porter NT, Van Treuren W, Nelson EJ, Garlena RA, Russell

DA, Martens EC, Sonnenburg JL. 2020. Bacteroides thetaiotaomicron-Infecting

Bacteriophage Isolates Inform Sequence-Based Host Range Predictions. Cell Host

Microbe 28:371-379.e5. 10.1016/j.chom.2020.06.011

63. Watson AR, Füssel J, Veseli I, DeLongchamp JZ, Silva M, Trigodet F, Lolans K, Shaiber

A, Fogarty E, Runde JM, Quince C, Yu MK, Söylev A, Morrison HG, Lee STM, Kao D,

Rubin DT, Jabri B, Louie T, Eren AM. 2023. Metabolic independence drives gut

microbial colonization and resilience in health and disease. Genome Biol 24:78.

10.1186/s13059-023-02924-x

64.   Lloyd-Price J, Arze C, Ananthakrishnan AN, Schirmer M, Avila-Pacheco J, Poon TW,

Andrews E, Ajami NJ, Bonham KS, Brislawn CJ, Casero D, Courtney H, Gonzalez A,

Graeber TG, Hall AB, Lake K, Landers CJ, Mallick H, Plichta DR, Prasad M, Rahnavard

G, Sauk J, Shungin D, Vázquez-Baeza Y, White RA, 3rd, Braun J, Denson LA, Jansson

JK, Knight R, Kugathasan S, McGovern DPB, Petrosino JF, Stappenbeck TS, Winter

HS, Clish CB, Franzosa EA, Vlamakis H, Xavier RJ, Huttenhower C. 2019. Multi-omics

of the gut microbial ecosystem in inflammatory bowel diseases. Nature 569:655-662.

10.1038/s41586-019-1237-9

65.   Palleja A, Mikkelsen KH, Forslund SK, Kashani A, Allin KH, Nielsen T, Hansen TH,

Liang S, Feng Q, Zhang C, Pyl PT, Coelho LP, Yang H, Wang J, Typas A, Nielsen MF,

Nielsen HB, Bork P, Wang J, Vilsbøll T, Hansen T, Knop FK, Arumugam M, Pedersen O.

2018. Recovery of gut microbiota of healthy adults following antibiotic exposure. Nat

Microbiol 3:1255-1265. 10.1038/s41564-018-0257-9

66.   Louis S, Tappu RM, Damms-Machado A, Huson DH, Bischoff SC. 2016.

Characterization of the gut microbial community of obese patients following a weight-

loss intervention using whole metagenome shotgun sequencing. PLoS One

11:e0149564. 10.1371/journal.pone.0149564

67.   Bischoff SC, Damms-Machado A, Betz C, Herpertz S, Legenbauer T, Löw T, Wechsler

JG, Bischoff G, Austel A, Ellrott T. 2012. Multicenter evaluation of an interdisciplinary 52-

week weight loss program for obesity with regard to body weight, comorbidities and

quality of life--a prospective study. Int J Obes (Lond) 36:614-24. 10.1038/ijo.2011.107

68.   Schloissnig S, Arumugam M, Sunagawa S, Mitreva M, Tap J, Zhu A, Waller A, Mende

DR, Kultima JR, Martin J, Kota K, Sunyaev SR, Weinstock GM, Bork P. 2013. Genomic

1175   variation landscape of the human gut microbiome. Nature 493:45-50.

1176   10.1038/nature11711

1177   69.   Chanin RB, West PT, Park RM, Wirbel J, Green GZM, Miklos AM, Gill MO, Hickey AS,

1178   Brooks EF, Bhatt AS. 2023. Intragenic DNA inversions expand bacterial coding capacity.

1179   bioRxiv doi:10.1101/2023.03.11.532203. 10.1101/2023.03.11.532203

1180   70.   Milman O, Yelin I, Kishony R. 2023. Systematic identification of gene-altering

1181   programmed inversions across the bacterial domain. Nucleic Acids Res 51:553-573.

1182   10.1093/nar/gkac1166

1183   71.   Pedersen TK, Brown EM, Plichta DR, Johansen J, Twardus SW, Delorey TM, Lau H,

1184   Vlamakis H, Moon JJ, Xavier RJ, Graham DB. 2022. The CD4(+) T cell response to a

1185   commensal-derived epitope transitions from a tolerant to an inflammatory state in

1186   Crohn's disease. Immunity 55:1909-1923.e6. 10.1016/j.immuni.2022.08.016

1187   72.   Curtis MM, Hu Z, Klimko C, Narayanan S, Deberardinis R, Sperandio V. 2014. The gut

1188   commensal *Bacteroides thetaiotaomicron* exacerbates enteric infection through

1189   modification of the metabolic landscape. Cell Host Microbe 16:759-69.

1190   10.1016/j.chom.2014.11.005

1191   73.   Spiga L, Winter MG, Furtado de Carvalho T, Zhu W, Hughes ER, Gillis CC, Behrendt

1192   CL, Kim J, Chessa D, Andrews-Polymenis HL, Beiting DP, Santos RL, Hooper LV,

1193   Winter SE. 2017. An oxidative central metabolism enables *Salmonella* to utilize

1194   microbiota-derived succinate. Cell Host Microbe 22:291-301 e6.

1195   10.1016/j.chom.2017.07.018

74.   Jimenez AG, Ellermann M, Abbott W, Sperandio V. 2020. Diet-derived galacturonic acid

regulates virulence and intestinal colonization in enterohaemorrhagic *Escherichia coli*

and *Citrobacter rodentium*. Nat Microbiol 5:368-378. 10.1038/s41564-019-0641-0

75.   De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, Collini S,

Pieraccini G, Lionetti P. 2010. Impact of diet in shaping gut microbiota revealed by a

comparative study in children from Europe and rural Africa. Proc Natl Acad Sci U S A

107:14691-6. 10.1073/pnas.1005963107

76.   Golden CD, Anjaranirina EJG, Fernald LCH, Hartl DL, Kremen C, Milner DA, Jr.,

Ralalason DH, Ramihantaniarivo H, Randriamady H, Rice BL, Vaitla B, Volkman SK,

Vonona MA, Myers SS. 2017. Cohort Profile: The Madagascar Health and

Environmental Research (MAHERY) study in north-eastern Madagascar. Int J Epidemiol

46:1747-1748d. 10.1093/ije/dyx071

77.   Carter MM, Olm MR, Merrill BD, Dahan D, Tripathi S, Spencer SP, Yu FB, Jain S, Neff

N, Jha AR, Sonnenburg ED, Sonnenburg JL. 2023. Ultra-deep sequencing of Hadza

hunter-gatherers recovers vanishing gut microbes. Cell 186:3111-3124.e13.

10.1016/j.cell.2023.05.046

78.   Rasmussen JL, Odelson DA, Macrina FL. 1986. Complete nucleotide sequence and

transcription of *ermF*, a macrolide-lincosamide-streptogramin B resistance determinant

from *Bacteroides fragilis*. J Bacteriol 168:523-33. 10.1128/jb.168.2.523-533.1986

79.   Garud NR, Good BH, Hallatschek O, Pollard KS. 2019. Evolutionary dynamics of

bacteria in the gut microbiome within and across hosts. PLoS Biol 17:e3000102.

10.1371/journal.pbio.3000102

1218 80. Wilson BC, Vatanen T, Jayasinghe TN, Leong KSW, Derraik JGB, Albert BB, Chiavaroli

1219   V, Svirskis DM, Beck KL, Conlon CA, Jiang Y, Schierding W, Holland DJ, Cutfield WS,

1220   O'Sullivan JM. 2021. Strain engraftment competition and functional augmentation in a

1221   multi-donor fecal microbiota transplantation trial for obesity. Microbiome 9:107.

1222   10.1186/s40168-021-01060-7

1223 81. Li SS, Zhu A, Benes V, Costea PI, Hercog R, Hildebrand F, Huerta-Cepas J, Nieuwdorp

1224   M, Salojärvi J, Voigt AY, Zeller G, Sunagawa S, de Vos WM, Bork P. 2016. Durable

1225   coexistence of donor and recipient strains after fecal microbiota transplantation. Science

1226   352:586-9. 10.1126/science.aad8852

1227 82. Smillie CS, Sauk J, Gevers D, Friedman J, Sung J, Youngster I, Hohmann EL, Staley C,

1228   Khoruts A, Sadowsky MJ, Allegretti JR, Smith MB, Xavier RJ, Alm EJ. 2018. Strain

1229   tracking reveals the determinants of bacterial engraftment in the human gut following

1230   fecal microbiota transplantation. Cell Host Microbe 23:229-240.e5.

1231   10.1016/j.chom.2018.01.003

1232 83. Ng KM, Aranda-Díaz A, Tropini C, Frankel MR, Van Treuren W, O'Loughlin CT, Merrill

1233   BD, Yu FB, Pruss KM, Oliveira RA, Higginbottom SK, Neff NF, Fischbach MA, Xavier

1234   KB, Sonnenburg JL, Huang KC. 2019. Recovery of the gut microbiota after antibiotics

1235   depends on host diet, community context, and environmental reservoirs. Cell Host

1236   Microbe 26:650-665.e4. 10.1016/j.chom.2019.10.011

1237 84. Roodgar M, Good BH, Garud NR, Martis S, Avula M, Zhou W, Lancaster SM, Lee H,

1238   Babveyh A, Nesamoney S, Pollard KS, Snyder MP. 2021. Longitudinal linked-read

1239   sequencing reveals ecological and evolutionary responses of a human gut microbiome

1240   during antibiotic treatment. Genome Res 31:1433-1446. 10.1101/gr.265058.120

1241   85.   David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV,

1242   Devlin AS, Varma Y, Fischbach MA, Biddinger SB, Dutton RJ, Turnbaugh PJ. 2014. Diet

1243   rapidly and reproducibly alters the human gut microbiome. Nature 505:559-63.

1244   10.1038/nature12820

1245   86.   Leeming ER, Johnson AJ, Spector TD, Le Roy CI. 2019. Effect of diet on the gut

1246   microbiota: Rethinking intervention duration. Nutrients 11. 10.3390/nu11122862

1247   87.   Nagel M, Reuter T, Jansen A, Szekat C, Bierbaum G. 2011. Influence of ciprofloxacin

1248   and vancomycin on mutation rate and transposition of IS256 in *Staphylococcus aureus*.

1249   Int J Med Microbiol 301:229-36. 10.1016/j.ijmm.2010.08.021

1250   88.   Truong DT, Tett A, Pasolli E, Huttenhower C, Segata N. 2017. Microbial strain-level

1251   population structure and genetic diversity from metagenomes. Genome Res 27:626-638.

1252   10.1101/gr.216242.116

1253   89.   Lee M. 2022. bit: a multipurpose collection of bioinformatics tools. F1000Research

1254   11:122. 10.5281/zenodo.3383647

1255   90.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics

1256   30:2068-9. 10.1093/bioinformatics/btu153

1257   91.   Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of

1258   protein or nucleotide sequences. Bioinformatics 22:1658-9.

1259   10.1093/bioinformatics/btl158

1260   92.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment

1261   search tool. J Mol Biol 215:403-10. 10.1016/s0022-2836(05)80360-2

1262   93.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

1263   improvements in performance and usability. Mol Biol Evol 30:772-80.

1264   10.1093/molbev/mst010

1265   94.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood

1266   trees for large alignments. PLoS One 5:e9490. 10.1371/journal.pone.0009490

1267   95.   Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic

1268   tree display and annotation. Nucleic Acids Res 49:W293-w296. 10.1093/nar/gkab301

1269   96.   Leinonen R, Sugawara H, Shumway M. 2011. The sequence read archive. Nucleic Acids

1270   Res 39:D19-21. 10.1093/nar/gkq1019

1271   97.   Bushnell B. BBTools, SourceForge. https://sourceforge.net/projects/bbmap/.

1272   98.   Li D, Liu CM, Luo R, Sadakane K, Lam TW. 2015. MEGAHIT: an ultra-fast single-node

1273   solution for large and complex metagenomics assembly via succinct de Bruijn graph.

1274   Bioinformatics 31:1674-6. 10.1093/bioinformatics/btv033

1275   99.   Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:

1276   prokaryotic gene recognition and translation initiation site identification. BMC

1277   Bioinformatics 11:119. 10.1186/1471-2105-11-119

1278   100.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat

1279   Methods 9:357-9. 10.1038/nmeth.1923

1280   101.   Pedersen BS, Quinlan AR. 2018. Mosdepth: quick coverage calculation for genomes

1281   and exomes. Bioinformatics 34:867-868. 10.1093/bioinformatics/btx699

1282   102.   Shen W, Le S, Li Y, Hu F. 2016. SeqKit: A cross-platform and ultrafast toolkit for

1283   FASTA/Q file manipulation. PLoS One 11:e0163962. 10.1371/journal.pone.0163962

1284   103.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,

1285   Durbin R. 2009. The sequence alignment/map format and SAMtools. Bioinformatics

1286   25:2078-9. 10.1093/bioinformatics/btp352

1287   104.   Bacic MK, Smith CJ. 2008. Laboratory maintenance and cultivation of *Bacteroides*

1288   species. Curr Protoc Microbiol Chapter 13:Unit 13C.1.

1289   10.1002/9780471729259.mc13c01s9

1290   105.   Varel VH, Bryant MP. 1974. Nutritional features of *Bacteroides fragilis* subsp. *fragilis*.

1291   Appl Microbiol 28:251-7. 10.1128/am.28.2.251-257.1974

1292   106.   Tartera C, Araujo R, Michel T, Jofre J. 1992. Culture and decontamination methods

1293   affecting enumeration of phages infecting Bacteroides fragilis in sewage. Appl Environ

1294   Microbiol 58:2670-3. 10.1128/aem.58.8.2670-2673.1992

1295   107.   Zheng L, Tan Y, Hu Y, Shen J, Qu Z, Chen X, Ho CL, Leung EL, Zhao W, Dai L. 2022.

1296   CRISPR/Cas-Based Genome Editing for Human Gut Commensal Bacteroides Species.

1297   ACS Synth Biol 11:464-472. 10.1021/acssynbio.1c00543

1298   108.   Willett JLE, Ji MM, Dunny GM. 2019. Exploiting biofilm phenotypes for functional

1299   characterization of hypothetical genes in Enterococcus faecalis. NPJ Biofilms

1300   Microbiomes 5:23. 10.1038/s41522-019-0099-0

1301   109.   Jones KR, Belvin BR, Macrina FL, Lewis JP. 2020. Sequence and characterization of

1302   shuttle vectors for molecular cloning in Porphyromonas, Bacteroides and related

1303   bacteria. Mol Oral Microbiol 35:181-191. 10.1111/omi.12304

110. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnetjournal 17:3. 10.14806/ej.17.1.200

111. Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics 26:841-2. 10.1093/bioinformatics/btq033

112. Hickey CA, Kuhn KA, Donermeyer DL, Porter NT, Jin C, Cameron EA, Jung H, Kaiko

GE, Wegorzewska M, Malvin NP, Glowacki RW, Hansson GC, Allen PM, Martens EC,

Stappenbeck TS. 2015. Colitogenic Bacteroides thetaiotaomicron Antigens Access Host

Immune Cells in a Sulfatase-Dependent Manner via Outer Membrane Vesicles. Cell

Host Microbe 17:672-80. 10.1016/j.chom.2015.04.002

113. Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021.

eggNOG-mapper v2: Functional annotation, orthology assignments, and domain

prediction at the metagenomic scale. Mol Biol Evol 38:5825-5829.

10.1093/molbev/msab293

114. Tatusov RL, Galperin MY, Natale DA, Koonin EV. 2000. The COG database: a tool for

genome-scale analysis of protein functions and evolution. Nucleic Acids Res 28:33-6.

10.1093/nar/28.1.33

115. Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, Edalatmand A, Huynh W,

Nguyen AV, Cheng AA, Liu S, Min SY, Miroshnichenko A, Tran HK, Werfalli RE, Nasir

JA, Oloni M, Speicher DJ, Florescu A, Singh B, Faltyn M, Hernandez-Koutoucheva A,

Sharma AN, Bordeleau E, Pawlowski AC, Zubyk HL, Dooley D, Griffiths E, Maguire F,

Winsor GL, Beiko RG, Brinkman FSL, Hsiao WWL, Domselaar GV, McArthur AG. 2020.

CARD 2020: antibiotic resistome surveillance with the comprehensive antibiotic

resistance database. Nucleic Acids Res 48:D517-d525. 10.1093/nar/gkz935

116.   Wood DE, Lu J, Langmead B. 2019. Improved metagenomic analysis with Kraken 2.

1328   Genome Biol 20:257. 10.1186/s13059-019-1891-0

1329