Title

NEDD4-binding protein 1 suppresses HBV replication by degrading HBV RNAs

Authors

Nobuhiro Kobayashi[a,b,*], Saori Suzuki[b,c,*], Yuki Sakamoto[b], Rigel Suzuki[b,c], Kento Mori[b], Yume

Kosugi[b], Lihan Liang[b], Tomoya Saito[a,b], Takuma Izumi[d], Kisho Noda[e], Daisuke Okuzaki[f], Yumi

Kanegae[g], Sanae Hayashi[h], Yasuhito Tanaka[h], Yoshiharu Matsuura[i,j], Osamu Takeuchi[k], Tomokazu

Tamura[b,c,1], Akinobu Taketomi[a], Takasuke Fukuhara #[b,c,j,l, m, n]

Affiliations

a. Department of Gastroenterological Surgery 1, Graduate School of Medicine, Hokkaido

12   University, Hokkaido, Japan

13   b. Department of Microbiology and Immunology, Graduate School of Medicine, Hokkaido

14   University, Hokkaido, Japan

15   c. Institute for Vaccine Research and Development: HU-IVReD, Hokkaido University,

16   Hokkaido, Japan

17   d. Department of Surgery, Matsuyama Red Cross Hospital, Ehime, Japan

18   e. Tokyo Bay Urayasu Ichikawa Medical Center, Chiba, Japan

19   f. Genome Information Research Center, Research Institute for Microbial Diseases, Osaka

20   University, Osaka, Japan

21   g. Core Research Facilities, Research Center for Medical Sciences, The Jikei University

22   School of Medicine, Tokyo, Japan

23   h. Department of Gastroenterology and Hepatology, Faculty of Life Sciences, Kumamoto

24   University, Kumamoto, Japan

25   i. Department of Molecular Virology, Research Institute for Microbial Diseases, Osaka

26   University, Osaka, Japan

27   j. Laboratory of Virus Control, Center for Infectious Disease Education and Research, Osaka

28   University, Osaka, Japan

29   k. Department of Medical Chemistry, Graduate School of Medicine, Kyoto University, Kyoto,

30   Japan

31   l. One Health Research Center, Hokkaido University, Hokkaido, Japan

32   m. AMED-CREST, Japan Agency for Medical Research and Development (AMED), Tokyo,

33   Japan

n. Department of Virology, Faculty of Medicine, Kyushu University, Fukuoka, Japan

Running Head: Effect of NEDD4-binding protein 1 on HBV replication

*These authors contributed equally

#Address correspondence to Fukuhara Takasuke, fukut@pop.med.hokudai.ac.jp.

Author order was determined by the basis of contribution (see Acknowledgement for detail).

Conflict of Interest statement

Nothing to report.

Abbreviations

cccDNA: covalently closed circular DNA

d.p.i.: days post-infection

d.p.t.: days post-transfection

ETV: entecavir

HBV: hepatitis B virus

HBcrAg: hepatitis B core-related antigens

HCC: hepatocellular carcinoma

IFN: interferon

RIP assay: RNA Immunoprecipitation assay

ISG: interferon-stimulated gene

KO: knockout

MALT1: mucosa-associated lymphoid tissue lymphoma translocation 1

NA: nucleos(t)ide analog

NTCP: sodium-taurocholate cotransporting polypeptide

N4BP1: NEDD4-binding protein 1

pgRNA: pregenomic RNA

PHH: primary human hepatocyte

qPCR: quantitative PCR

RBP: RNA-binding protein

rcDNA: relaxed circular DNA

WB: western blotting

WT: wild type

<u>Abstract</u>

Chronic infection with hepatitis B virus (HBV) places patients at increased risk for liver

cirrhosis and hepatocellular carcinoma. Although nucleos(t)ide analogs are mainly used for the

treatment of HBV, they require long-term administration and may lead to the emergence of drug

resistant mutants. Therefore, to identify targets for the development of novel anti-HBV drugs, we

screened for HBV-suppressive host factors using a plasmid expression library of RNA-binding

proteins (RBPs). We screened 132 RBPs using an expression plasmid library by measuring HBV

relaxed circular DNA (rcDNA) levels in hepatocellular carcinoma. Our screen identified

NEDD4-binding protein 1 (N4BP1) as having an anti-HBV effect. In hepatocellular carcinoma cell

lines transfected or infected with HBV, overexpression of N4BP1 decreased rcDNA levels while

knockdown or knockout of the gene encoding N4BP1 rescued rcDNA levels. N4BP1 possesses the

KH-like and RNase domains and both were required for the anti-HBV effect of N4BP1. Additionally,

we measured levels of HBV pregenomic RNA (pgRNA) and covalently closed circular DNA

(cccDNA) in the RBP-transfected cells and confirmed that N4BP1 binds pgRNA directly and

degraded both the 3.5 kb and 2.4/2.1 kb HBV RNA. In summary, N4BP1 is a newly identified host

factor able to counteract HBV production by promoting the degradation of 3.5 kb and 2.1/2.4 kb

HBV RNA.

Importance

There is still a large number of HBV-infected people in the world today because of no

curative treatment for HBV infection. In this study, we focused on and screened RNA-binding

proteins to identify new host factors which inhibit HBV replication. As a result, we found that

NEDD4-binding protein 1 (N4BP1) expression suppresses rcDNA production by promoting the

degradation of pregenomic RNA, 2.4kb and 2.1kb HBV RNA. Furthermore, KH-like domain or

RNase domain of N4BP1 were involved in this anti-HBV effect. In addition, the N4BP1 levels were

lower in HCC resection samples of exacerbated patients, suggesting that individual N4BP1 levels

might be related to HCC progression. This novel factor can potentially become a key to new HBV

treatments.

Introduction

Hepatitis B virus (HBV) infection causes either acute or chronic hepatitis, with approximately

296 million people worldwide suffering from chronic hepatitis B and an estimated 1.5 million new

cases arising each year. Chronically infected individuals are at increased risk for liver disease, and

annually ~820,000 die from cirrhosis and hepatocellular carcinoma (HCC) caused by HBV infection

(1). Although there is a prophylactic vaccine for HBV, there are still many new cases of HBV

infection due to the large number of chronic HBV infected patients still in Africa, Asia, and Eastern

Europe and the low vaccination rates in these highly endemic areas (2).

HBV is an enveloped DNA virus with a small, partially double-stranded genome of 3.2 kbp.

Entry into hepatocytes is mediated by binding to the bile acid transporter sodium-taurocholate

cotransporting polypeptide (NTCP). After being released into the nucleus, the HBV genome is

converted from its relaxed circular DNA (rcDNA) form to covalently closed circular DNA

(cccDNA) using the host DNA repair system (3-6). The cccDNA serves as the transcription template

for viral mRNAs, including the HBV pregenomic RNA (pgRNA). pgRNA is used to synthesize the

negative-strand DNA first and then the positive-strand is synthesized to generate rcDNA.

Interferon (IFN) and nucleos(t)ide analogs (NAs) are currently used as treatments for HBV

infection, and both have shown efficacy in controlling viremia (7, 8). IFNs are involved in the

inhibition of HBV replication, transcription, and other important processes by inducing the

expression of various IFN-stimulated genes (ISGs) in host cells (9). However, IFN treatment can

cause considerable side effects, including liver damage, and is thus unsuitable for patients with

severe liver dysfunction (10).

In contrast, NAs impair further HBV intrahepatic spreading by blocking production of new

virions. But the HBV antigens HBs or HBe are not eliminated with treatment as NAs inhibit the

reverse transcription step that occurs after these antigens have already been translated. Furthermore,

NAs do not directly target cccDNA, allowing the virus to persist in hepatocytes (11, 12).

Consequently, to prevent relapse due to reactivation of HBV after discontinuing NAs, long-term

treatment is recommended (13).

In efforts to overcome the limitations of these current therapeutic options, research has looked

to identify host factors with an antiviral role. Understanding the mechanism of action of such factors

could aid in the development of new antiviral treatments. Several host factors are known to inhibit

HBV replication, including RNA-binding proteins (RBPs). For example, the CCCH zinc finger

antiviral protein ZAP (ZC3HAV1) binds to HBV RNA with its four zinc finger motifs at the

N-terminus, promoting pgRNA degradation (14). Similarly, MCPIP1 (ZC3H12A, regnase-1) has

also been reported to inhibit HBV replication by binding to HBV pgRNA through its zinc finger

motif, directly degrading pgRNA via its RNase activity (15). Furthermore, PUF60 is a host factor

that not only positively regulates HBV 3.5 kb precore plus pregenomic RNA expression by binding

to the HBV enhancer region II with the transcription factor TCF7L2 but can also promote

degradation of the HBV 3.5 kb RNA by binding via its N-terminal RNA recognition motif (16).

Despite these descriptions, HBV produces high levels of HBV RNAs in infected primary human

hepatocytes (PHHs) and i*n vivo*, so the antiviral effects of these host factors appear limited.

To determine if other RBPs may have an antiviral role, we performed a comprehensive screen

using an expression library of RBPs. We identified NEDD4-binding protein 1 (N4BP1) as a host

factor with anti-HBV activity. We subsequently elucidated the mechanism of how N4BP1 suppresses

HBV replication by promoting the degradation of HBV RNAs.

Results

Screen for RNA-binding proteins that inhibit HBV propagation

To identify novel RBPs involved in HBV propagation, we screened 132 RBPs using an

expression plasmid library. Huh7 cells were co-transfected with an RBP expression plasmid and a

1.3-fold-overlength HBV genome (genotype C). At 3 days post-transfection (d.p.t.), cells were

collected to measure intracellular rcDNA in core particles. As shown in Fig 1A, 23 out of 132 RBPs

suppressed rcDNA levels by at least 50% compared to the control.

As a secondary screen, we then examined the effect of these 23 RBPs on HBV entry and

replication. HepG2-hNTCP C4 cells expressing each of the 23 RBPs were infected with HBV

(genotype D), and intracellular rcDNA was measured by quantitative PCR (qPCR) at 14 days

post-infection (d.p.i.). Seven of the 23 RBPs (N4BP1, TIA-1, ZC3H12B, PJA1, HNRNPC,

ZCCHC10, and KHNYN) reduced rcDNA levels in comparison with the control (Fig 1B). These

results were consistent with previous reports that found TIA-1 and PJA1 inhibit expression of HBs

antigen and HBV transcription and replication, respectively (17, 18).

Of the five remaining factors, we found by western blot that three (N4BP1, HNRNPC, and

ZCCHC10) are endogenously expressed in Huh7 and HepG2-hNTCP C4 cells (Fig 1C). To test

whether these proteins affect HBV replication, we knocked down each one and assessed rcDNA

levels following transfection with the 1.3-fold-overlength genome of HBV genotype C. The antiviral

drug entecavir, a nucleoside analog, was used as a positive control. Knockdown of each factor was

confirmed by western blotting (WB), but unlike knockdown of N4BP1, knockdown of HNRNPC or

ZCCHC10 did not consistently increase the levels of rcDNA depending on the siRNA used (Fig

2A-2C). Thus, N4BP1 became the focus of our study.

N4BP1 is a suppressive host factor of HBV replication

Next, to determine whether N4BP1 also had an antiviral effect against other HBV genotypes,

we co-transfected Huh7 cells with a plasmid expressing N4BP1 and a plasmid containing

1.3-fold-overlength genomes of HBV genotypes A, B, and C. At 3 d.p.t., rcDNA levels were

decreased in cells transfected with each genotype (Fig 3A).

To further elucidate the role of N4BP1 in HBV replication, we constructed Huh7 N4BP1

knockout (KO) cells using CRISPR/Cas9. We generated two clones of N4BP1 KO Huh7 cells (Fig

3B) and once more assessed whether intracellular rcDNA levels changed at 3 d.p.t. with a plasmid

containing the 1.3-fold-overlength genome of HBV genotype C. In both KO lines, rcDNA levels

increased approximately 3-fold compared to control cells (Fig 3B). After rescuing N4BP1 expression

in our KO line, rcDNA levels became significantly decreased again (Fig 3C, 3D), suggesting that

N4BP1 is a suppressive host factor of HBV replication.

In addition, we performed a transcriptomic analysis of N4BP1-overexpressing PHHs vs control

PHHs to assess whether N4BP1 affects regulatory transcription factors that might affect HBV

replication (Fig. 4). We set a cutoff value of gene expression at absolute fold change greater than 2

and an adjusted P-value less than 0.1 compared to control cells and found 178 genes upregulated in

N4BP1-overexpressing vs control PHHs and 230 genes downregulated (Fig 4A-B, S1 Table). None

of these factors were regulatory transcription factors. In addition, pathway enrichment analysis did

not show significant coordinated changes in genes associated with translational pathways (Fig 4C).

These findings suggest that N4BP1 overexpression was not associated with overarching changes in

the expression of regulatory transcription factors that would also affect HBV RNA production.

N4BP1 expression is not induced by HBV

We then evaluated whether N4BP1 expression level was increased by HBV infection. A

plasmid containing the 1.3-fold-overlength genome of HBV genotype C was transfected into Huh7

cells, and N4BP1 expression assessed by WB and qPCR at 3 d.p.t. Transfection of the HBV genome

did not significantly affect N4BP1 expression at either the transcriptional or protein level (Fig 5A,

4B). N4BP1 expression also did not significantly change in HepG2-hNTCP C4 cells at 3 and 10 d.p.i.

(Fig 5A, 5C). Finally, we measured N4BP1 expression in HepAD38.7 cells that stably secrete HBV

particles unless cultured in the presence of tetracycline. In this system, removal of tetracycline and

induction of HBV particle production also did not result in any change in N4BP1 expression (Fig 5A,

5D). Together, these results indicate that N4BP1 expression is not affected by HBV infection.

The KH-like and RNase domains of N4BP1 are critical for the anti-HBV effect of N4BP1

As shown in Fig 6A, N4BP1 is composed of 896 amino acids and contains a KH-like domain at

its N terminus side and an RNase domain at its C terminus (19). To examine which N4BP1 domains

are needed to suppress HBV replication, we constructed deletion mutants lacking either the KH-like

(ΔKH; deletion of amino acids 59-143) or the RNase domain (ΔRNase; deletion of amino acids

617-769).

The plasmid encoding wild-type or mutant N4BP1 was respectively co-transfected into naïve

Huh7 cells with the plasmid containing the 1.3-fold-overlength genome of HBV genotype C and the

rcDNA levels were measured. The expression of the deletion mutants was confirmed by WB using

an HA-tagged antibody (Fig 6B, upper). The rcDNA level in N4BP1 overexpressing cells was

significantly reduced, while the rcDNA levels in the ΔKH- or ΔRNase-expressing cells were

comparable to those in the control cells (Fig 6B, lower).

Residue D623 in the RNase domain of N4BP1 (Fig 6A) has been shown to be important for the

protein's RNase activity (20). A D623N substitution causes loss of RNase activity without changing

the structure of N4BP1. Co-transfecting the Huh7 cells with a plasmid expressing this N4BP1 D623

mutant and the HBV overlength genome also did not lead to a decrease in rcDNA levels (Fig 6C).

These results suggest that both the KH-like and RNase domains are critical for the anti-HBV effect

of N4BP1.

N4BP1 promotes pgRNA degradation

Having shown the effect of N4BP1 on rcDNA, we then sought to determine more specifically

which step of HBV replication N4BP1 suppresses. To that end, we assessed the impact of N4BP1

overexpression on cccDNA and pgRNA levels. HepG2-hNTCP C4 cells were infected with HBV

and cells were harvested at 14 d.p.i. While cccDNA levels were not affected by overexpression of

N4BP1, pgRNA and rcDNA levels were both significantly reduced (Fig 7A-7C). The HBV progeny

secreted from infected HepG2-NTCP C4 cells is not infectious, and thus only single-round infection

is observed in this system (21, 22). Thus, these results indicate that N4BP1 affects the steps of HBV

replication after cccDNA was formed and augmented pgRNA degradation.

To elucidate the mechanism of pgRNA suppression by N4BP1, we turned to HepAD38.7 cells.

First, the cells were cultured for 2 days in media without tetracycline to induce HBV replication.

Then N4BP1 or the domain deletion mutant plasmids were transfected, and tetracycline added at 2

d.p.t. to stop viral replication. Cells were collected starting 2 days after adding tetracycline to

measure the pgRNA levels (Fig 7D). As shown in Fig 7E, the expression of wild-type N4BP1

augmented the rate of pgRNA degradation. In contrast, both domain deficient mutants resulted in

pgRNA degradation at a rate similar to the control, suggesting that N4BP1 affects the degradation

rate of pgRNA. Next, we examined whether N4BP1 expression also suppresses HBV RNAs other

than pgRNA. We co-transfected Huh7 cells with a plasmid expressing N4BP1 and a plasmid

containing 1.3-fold-overlength genomes of HBV genotype C. At 3 d.p.t., cells were collected and

HBV RNA expression was measured by Northern blotting. Results showed that control Huh7 cells

showed intense expression of 3.5kb, 2.4/2.1kb HBV RNA, whereas N4BP1 overexpression Huh7

cells showed weak expression of all these RNAs (Fig 7F).

Next, to test that N4BP1 binds HBV RNA directly, we performed an RNA immunoprecipitation

(RIP) assay. Huh7 cells were co-transfected with a plasmid containing the 1.3-fold-overlength

genome of HBV genotype C and HA-tagged N4BP1 wildtype or domain deficient mutants or HA

control vector. Cells were collected at 3 d.p.t. and immunoprecipitated with anti-HA-tag monoclonal

antibody magnetic beads. Wildtype and N4BP1 domain-deficient mutants were confirmed to be

precipitated by an anti-HA-tag monoclonal antibody by western blot (Fig. 7G). The RNA-protein

interaction was expected to be most stable in the ΔRNase domain mutant, and our RIP-quantitative

PCR (qPCR) assay confirmed this (Fig. 7H). In contrast, the amount of pgRNA recovered with

wildtype N4BP1 and ΔKH-like domain mutants was far less, presumably due to the intact RNase

activity.

Furthermore, to investigate whether N4BP1 also led to reduced expression of viral proteins, we

assessed levels of HBeAg, HBsAg and HB core-related antigen (HBcrAg) in the supernatant of

HBV-infected PHHs at 14d.p.i., the timepoint when we had observed decreased pgRNA (Fig. 7I). All

3 HBV proteins we evaluated were decreased in the N4BP1-overexpressing PHHs compared to

control, indicating that N4BP1 inhibits pcRNA, PreS2/S RNA and potentially pgRNA, consistent

with our northern blot data.

N4BP1 expression is induced by IFN-α/λ in PHHs

It has been reported that N4BP1 expression is increased 3- to 5-fold with IFN-α treatment in

Jurkat cells and THP-1 cells (20). To see if this was the case in PHHs and hepatoma cell lines, we

treated Huh7, HepG2-hNTCP C4, and PHHs with 1000 U/ml of IFN-α and 100 ng/ml of IFN-λ,

which are known to also inhibit HBV replication. Cells were collected at 6, 24 and 48 hours

post-drug treatment without HBV infection, and expression of *N4BP1* mRNA and protein was

examined by qPCR and WB, respectively (Fig 8A). *ISG15* mRNA was included as a positive control.

As shown in Fig 8B, levels of *N4BP1* mRNA were increased 2-fold in both Huh7 and

HepG2-hNTCP C4 cells by IFN-α at 6 hours and 24 hours, respectively, which is small, albeit

significant in expression. IFN-λ stimulation increased levels of *N4BP1* mRNA in PHHs at 24 hours

and 48 hours significantly and N4BP1 protein levels were prompted in Huh7 cells and PHHs (Fig

8C).

IFN-α and IFN-λ treatment in PHHs resulted in an approximately 2-fold increase in *N4BP1*

mRNA expression and protein level was also increased. Collectively, these results indicate that

IFN-α treatment induces N4BP1 expression in Huh7 and HepG2-hNTCP C4 cells, while both IFN-α

and IFN-λ induces N4BP1 expression in PHHs. As primary cells, PHHs are more physiologically

relevant than hepatoma lines such as Huh7 and HepG2-hNTCP C4 cells in which the response to

IFN is disrupted. To build upon our in vitro findings, we then examined samples from patients with

HBV-HCC to assess whether there was an association between *N4BP1* mRNA expression in the liver

and HBV reactivation status. All the patients showed HBs-Ag (+), HBe-Ag (-) and HBe-Ab (+), and

categorized as either HBV DNA positive or negative per HBV DNA levels in the serum, and *ISG15*

and *N4BP1* mRNA expression were assayed in non-cancerous liver tissue from these patients (Fig.

9). *ISG15* mRNA did not significantly differ between the patient groups, but *N4BP1* mRNA was

significantly higher in patients who were HBV DNA negative vs positive.

<u>Discussion</u>

NAs are generally well tolerated, very effective in inhibiting HBV replication, and are currently

the most commonly used treatment for chronic hepatitis B. However, since NAs inhibit reverse

transcription, HBV cccDNA remains in the nucleus of hepatocytes so the patient is never completely

cured. Indeed, patients who discontinue NA treatment can experience virologic relapse and thus

must continue NA therapy for extended periods. Long-term treatment with NA is associated with the

development of antiviral drug resistance; for example, mutations in the HBV polymerase gene,

which is a target of NAs, have been reported (23-25). Resistance to NA therapy can lead to hepatitis

relapse and even death. Thus, we sought to identify host factors involved in HBV proliferation that

would be distinct from the viral sites targeted by NAs. In a library of 132 RBPs containing proteins

with known RNA-binding domains such as RNA recognition motif domains or zinc finger domains,

we found that N4BP1 inhibited HBV replication, was negatively correlated with HBV rcDNA levels

in hepatoma cell lines and suppressed HBV replication by promoting pgRNA degradation.

We found that both the KH-like domain and the RNase domain of N4BP1 were essential for the

protein's anti-HBV effect (Fig 6B). It has been reported that the KH-like domain binds to RNA or

single-strand DNA and is involved in splicing, transcriptional regulation, and translational regulation

(19, 26, 27). On the other hand, the RNase domain of N4BP1 is also involved in regulating genome

stability and repressing the expression of target genes through mRNA strand breaks (28). Based on

our RIP assay data, it is possible that N4BP1 binds HBV RNA via its KH-like domain and degrades

HBV RNA by the RNase activity of its RNase domain. However, this would need to be confirmed

with a RIP assay using a doubly deficient RNase and KH-like domain mutant. In fact, in this study,

N4BP1 suppressed HBV replication at the step of pgRNA production in the HBV life cycle (Fig 7).

We also found that N4BP1 suppresses production of not only the 3.5 kb RNA of HBV, but also the

2.4/2.1 kb RNA by northern blot, which implies N4BP1 promotes degradation of pgRNA, PreS1 and

PreS2/S. However, it is still unclear and remains to be assessed whether HBx is also affected by

N4BP1.

N4BP1 is known to be an inhibitor of innate immunity and inflammation-mediated cytokine

production. NF-κB is an important transcription factor for the elicitation of cytokine responses, and

its activation is necessary for many immune responses. N4BP1 interacts with the NF-κB signal

essential modulator (NEMO, also known as IκB kinase γ) to suppress Toll-like receptor (TLR)–

dependent NF-κB activation (29). Furthermore, even in vivo, it has been reported that N4BP1-/-

mice develop mild inflammation and have exacerbated TLR-dependent inflammatory responses (30).

Recently, N4BP1 was found as a new hotspot gene for HBV integration (31). Although the

significance of HBV integration to N4BP1 remains unknown, it is interesting that the virus

integration occurred to N4BP1 which has anti-HBV activity. Regarding virus replication, N4BP1 is

known to directly recognize and degrade retroviruses such as HIV-1 (20) and to be involved in

replication of the Porcine Reproductive and Respiratory Syndrome Virus (32). In both cases, the

RNase activity of N4BP1 was found to inhibit viral replication by directly degrading viral RNA,

consistent with our data. In addition, the RNase domain of N4BP1 has a similar structure to that of

MCPIP1(33), which has been reported to recognize and degrade HBV pgRNA directly (15).

However, MCPIP1 is a cytoplasmic RNase, whereas N4BP1 is mainly localized on the nucleus.

Furthermore, MCPIP1 has not been reported to suppress 2.4 kb and 2.1 kb HBV RNAs, whereas we

have demonstrated that N4BP1 suppresses these RNAs. These results suggest that MCPIP1 and

N4BP1 may regulate HBV degrading mRNAs through different processes. The mechanism behind

our finding that the KH-like domain of N4BP1 was also important for the anti-HBV effect of this

protein is presumably due to its binding to HBV RNA but further work is needed to elucidate the

mechanism more precisely.

Although N4BP1 expression has been reported to be induced by type I IFN (20), stimulation of

N4BP1 expression by IFN-α (type I IFN) and IFN-λ (type III IFN) treatment varied between the

PHHs and hepatoma cell lines we tested. This could be due to the disrupted IFN signaling reported

in the cell lines Huh7 and HepG2 that result in them being less sensitive to IFN (34, 35). Also,

ISG15 is an inducible factor by IFN-α and IFN-λ (36, 37). In characterizing N4BP1 expression, we

found that N4BP1 expression was not induced by HBV infection or transfection in hepatoma cell

lines (Fig 5). Furthermore, in clinical liver tissue samples, N4BP1 expression was significantly lower

in HBV-DNA positive vs HBV-DNA negative samples (Fig 9B). These results suggest that N4BP1

expression may differ according to each individual and that expression of N4BP1 in the liver may

impact HBV reactivation. Otherwise, it is thought that factors upstream of N4BP1 may cause

differences in N4BP1 expression and affect HBV reactivation. In addition, research has shown that

N4BP1 can be inactivated by caspase-8 (30) or mucosa-associated lymphoid tissue lymphoma

translocation 1 (MALT1) (20) in immune cells. In the latter instance, N4BP1 inactivation by

MALT-1 induced HIV-1 reactivation. This does not appear to be the case in our study, as our

RNA-seq data did not indicate MALT-1 induction by N4BP1 overexpression in PHH, but other

signaling proteins may have a similar effect, resulting in greater HBV replication in hepatocytes.

This will need to be studied further.

A limitation of our current study is the small number of patient samples used to evaluate the

relationship between N4BP1 and HBV reactivation. Studies with a greater number of samples are

needed to further characterize this possible relationship. In addition, although transcriptomic analysis

implies that N4BP1 was not associated with transcriptional regulation, it is required to strengthen

this result by other methods such as ActD chase assay. Also, identification of the element(s) of HBV

RNA that bind N4BP1 remains to be determined. As HBV RNAs overlap each other, explicit

identification of the binding region(s) was difficult. A previous report suggested that sequence

motif(s) other than GC dinucleotides may be recognized by N4BP1 as N4BP1 can degrade HIV-1

transcripts, which have a markedly reduced frequency of GC dinucleotides (20). However, based on

our data, it is conceivable that an overlapping area or areas of the 3.5kb and 2.4kb/2.1kb HBV RNAs

may bind to N4BP1.

In conclusion, to investigate different treatments against HBV infection from currently

approved HBV therapeutics, we focused on RBPs. We found that N4BP1 is a RBP with novel

anti-HBV activity, inhibiting HBV replication via degradation of HBV RNAs. Although the detailed

molecular mechanism behind the anti-HBV effect of N4BP1 remains to be uncovered, understanding

the association between N4BP1 and HBV could contribute to the development of new HBV

treatments.

Materials and methods

Cell culture

All cells were grown at 37°C in a 5% (v/v) CO2 environment. Huh7 cells and HEK293T cells

were obtained from the Japanese Collection of Research Bioresources Cell Bank (JCRB, Osaka,

Japan, JCRB0403 and JCRB9068). HepG2-hNTCP C4 cells and HepAD38.7 cells were kindly

provided by Takaji Wakita (NIID, Tokyo, Japan). PHHs were obtained from Phoenix Bio (Hiroshima,

Japan). Huh7 cells and HEK293T cells were maintained in DMEM (Nacalai Tesque, Kyoto, Japan)

supplemented with 100 U/ml of penicillin, 100 μg/ml of streptomycin (Nacalai Tesque) and 10%

fetal bovine serum (FBS, Sigma, MO, USA). HepG2-hNTCP C4 cells were maintained in the

aforementioned media containing 400 μg/ml G418 (Nacalai Tesque). HepAD38.7 cells were

maintained in DMEM/Ham's F-12 media (Nacalai Tesque) supplemented with 10% FBS, 100 U/ml

of penicillin, 100 μg/ml of streptomycin, 400 μg/ml of G418, 5 μg/ml of insulin (Nacalai Tesque),

and 400 ng/ml of tetracycline (Nacalai Tesque). PHHs were maintained in PHH-specific media

provided by the manufacture (Phoenix Bio).

Generation of RBP overexpressing cells

RBP-overexpressing Huh7 cells were generated by transfection of RBP expression plasmids

using TransIT LT-1 transfection reagent (Mirus, WI, USA). Briefly, Huh7 cells were seeded in

24-well plates (Greiner, Bad Haller, Austria) 1 day prior to the transfection. The following day, 1 μl

of TransIT LT-1 transfection reagent, 0.3 μg of RBP expression plasmid, and 0.1 μg of a plasmid

containing a 1.3-fold-overlength genome of HBV (genotype C) were mixed in 50 μl of Opti-MEM

(Thermo Fisher Scientific, MA, USA). After 15 minutes incubation at room temperature, the mixture

was added into the supernatant of the cultured Huh7 cells. Entecavir (Fujifilm, Tokyo, Japan) was

added at 5 μM the day after transfection as a positive control.

RBP-overexpressing HepG2-hNTCP C4 cells were made by lentiviral transduction.

Lentiviruses were made by transfecting HIV gag/pol, VSV-G, REV, and pcs-II EF RBP expression

plasmids into HEK293T cells using TransIT LT-1 transfection reagent. At 2 d.p.t., the supernatant of

HEK293T was collected and spun at 4,200 rpm for 2 minutes at room temperature and the

392 supernatant added to HepG2-hNTCP C4 cells. Entecavir or heparin was used as a positive control.

393 Entecavir was added at 5 μM the day after infection. Heparin (Mochida Pharmaceutical Co., Tokyo,

394 Japan) was added at 5 U/ml 30 min before infection.

395  To express N4BP1 in PHHs, we used an adenovirus vector (AdV). The N4BP1expressing

396 adenovirus vector (N4BP1 AdV) was constructed using the cosmid cassette pAxEFwit2, which has a

397 cloning site *SwaI* flanked by the EF1α promoter and rabbit β-globin polyA signal in the E1-deleted

398 region of the adenovirus genome. The N4BP1 cDNA was prepared from the pCS2-EF-N4BP1-HA

399 plasmid and inserted into the *SwaI* cloning site. The AdV was prepared as previously described (38,

400 39). The AdV was titrated using the methods described by Pei et al (40). Briefly, the copy numbers

401 of viral genome successfully transduced into target cells were measured by qPCR. For generating

402 N4BP1-overexpressing PHHs, cells were infected with N4BP1 AdV at a multiplicity of infection

403 (MOI) of 5 or 30 two days before HBV infection.

405 **Generation of RBP knockdown cells**

406  RBP knockdown cell lines were generated using Silencer Select siRNA (Thermo Fisher

407 Scientific) and Lipofectamine RNAiMAX (Thermo Fisher Scientific) following the manufacturer's

408 protocol. Briefly, 1.5 μl of Lipofectamine RNAiMAX and 5 pg of siRNA were mixed in 50 μl of

409 Opti-MEM. After incubating for 5 minutes at room temperature, the mixture was gently added to the

cell supernatant. The following siRNA reagents were used in this study: control siRNA (Silencer

Select Negative Control No.1 siRNA, 4390843), ZCCHC10 siRNA#1 (s29507), ZCCHC10

siRNA#2 (s29508), ZCCHC10 siRNA#3 (s29509), HNRNPC siRNA#1 (s6719), HNRNPC

siRNA#2 (s6720), HNRNPC siRNA#3 (s6721), N4BP1 siRNA#1 (s18638), N4BP1 siRNA#2

(s18639).

Generation of N4BP1 knockout cells

N4BP1 KO Huh7 cells were generated by CRISPR/Cas9 using

pX330-U6-Chimeric_BB-CBh-hSpCas9 (pX330, Addgene, MA, USA) and HR110PA-1 (System

Biosciences, CA, USA). The pX330-puro vector was constructed by integrating the puromycin

resistance gene sequence from the HR110PA-1 vector into the pX330 vector. The pX330-puro vector

was digested using *BbsI* (Biolabs, MA, USA) and the target sequence was subcloned. The plasmid

was transfected into Huh7 cells using TransIT LT-1 transfection reagent and after selection with

puromycin dihydrochloride (Invivogen, CA, USA), a single colony was isolated using cloning

cylinders (Merck, Darmstadt, Germany) and propagated. KO of the target gene was confirmed by

WB and DNA sequencing.

Plasmid Constructions

We used the following plasmids containing 1.3-fold-overlength genomes of different HBV

genotypes: pUC19-Ae_us (genotype A, accession no. AB246337), pGEM-Bj-JPN56 (genotype B,

accession no. AB246342), and pUC19-C_JPNAT (genotype C, accession no. AB246345).

Each RBP expression plasmid had a C-terminus HA tag and was inserted between the *XhoI* and

XbaI sites of the pcs-II EF vector. The KH-like domain deletion mutant (ΔKH), RNase domain

deletion mutant (ΔRNase), and D623N mutant were generated from the N4BP1 expression plasmid.

ΔKH plasmid was generated by deleting amino acids 59-143 from N4BP1; ΔRNase was generated

by deleting amino acids 617-769 N4BP1; and D623N was generated by changing the 623rd amino

acid from asparagine acid to asparagine. To insert these mutated N4BP1 genes into the pcs-II EF

vector between the *XhoI* and *XbaI* sites, we used the following primer sets and In-Fusion Snap

Assembly Master Mix (Takara Bio, Shiga, Japan) per the manufacturer's directions. Each N4BP1

mutant plasmid had a C-terminus HA tag.

ΔKH: 5'- CGCTACCGGTCTCGAGATGGCGGCCCGGGCGGTGCT -3' (forward)

5'- GGGTAGGTTCTCTTTCGCCCCGCAGAGCTGCAGCC -3' (reverse) , and

5'- CAGCTCTGCGGGGCGAAAGAGAACCTACCCAGTAG -3' (forward)

5'- GGTACATGGTCTCGAGATCCAACACCATGGCAGAAA -3' (reverse)

ΔRNase: 5'- CGCTACCGGTCTCGAGATGGCGGCCCGGGCGGTGCT -3' (forward)

5'- TTCCTTCTGAAGAAAATCCGTTCTCCCTGGTTCAT -3' (reverse) , and

5'- CCAGGGAGAACGGATTTTCTTCAGAAGGAAGTCTG -3' (forward)

5'- GGGTAGGTTCTCTTTCGCCCCGCAGAGCTGCAGCC -3' (reverse)

D623N: 5'- CGCTACCGGTCTCGAGATGGCGGCCCGGGCGGTGCT -3' (forward)

5'- AACATTGCTCCCATTTATAACAATGTGTTTCAAAT -3' (reverse) , and

5'- AATGGGAGCAATGTTGCAATTACCCATGGTCTGAA -3' (forward)

5'- GGTACATGGTCTCGAGATCCAACACCATGGCAGAAA -3' (reverse)

Viruses

The production of HBV particles (genotype D) was adapted from a previous report (41). Briefly,

HepAD38.7 cells were cultured under tetracycline (400 ng/mL) for maintenance. To produce HBV

particles, tetracycline was removed and the supernatant collected after 6 and 9 days. The collected

supernatant was mixed with 30% polyethylene glycol (PEG) 8000 (Sigma) and incubated overnight

at 4°C. Then it was spun at 3,000 rpm for 20 minutes at 4°C for concentration. Before HBV infection,

HepG2-hNTCP C4 cells were seeded in collagen type I-coated 24-well plates (Iwaki, Tokyo, Japan)

and transduced with lentivirus to express RBP (see "Generation of RBP-overexpressing cells"

above). At 2 days post-transduction, the cells were infected with 5,000 genome equivalents

(GEq)/cell of HBV in DMEM containing 3% DMSO (Nacalai Tesque) and 4% PEG 8000, and the

media was changed every 2-3 days until 14 d.p.i..

HBV genotype C for PHHs infection was purchased from Phoenix Bio. The infection into

PHHs was performed at 5 Geq/cell in PHH-specific media containing 40% PEG 8000 following the

manufacturer's protocol. The culture media was replaced every 4 days. Cells were harvested at 14

d.p.i. for WB and reverse transcription quantitative PCR (RT-qPCR).

Western blotting

Cells were lysed with $2 \times$ Laemmli sample buffer (Bio Rad Laboratories, CA, USA) containing

10% 2-mercaptoethanol (Sigma) and boiled for 10 minutes at 100°C. Then lysates were loaded on a

10-20% gradient SDS-PAGE gel (Atto, Tokyo, Japan) and transferred onto a polyvinylidene fluoride

transfer membrane (Millipore, MA, USA). Membranes were immersed in Block Ace (Kac, Kyoto,

Japan) containing 1% bovine serum albumin (Sigma) and incubated for 1 hour at room temperature.

The blocking solution was then decanted, and replaced with Can Get Signal Solution 1 (Toyobo,

Osaka, Japan) containing the appropriate antibodies described below and incubated for 1 hour. The

membranes were washed with $1 \times$ PBS (Fujifilm) containing 0.1% polyoxyethylene (20) sorbitan

monolaurate (Fujifilm) and incubated with Can Get Signal Solution 2 (Toyobo) containing

horseradish peroxidase-conjugated antibody against rabbit or mouse immunoglobulins for 1 hour.

After addition of Amersham ECL Prime Western Blotting Detection Reagents (Cytiva, Tokyo, Japan),

the membranes were subsequently imaged via LuminoGraph I (Atto).

The following antibodies were used in this study: anti-N4BP1 (Bethyl Laboratories, TX, USA),

anti-ZC3H12B (Gene Tex, CA, USA), anti-HNRNPC (abcam, Cambridgeshire, UK),

anti-ZCCHC10 (Sigma), anti-KHNYN (Fujifilm), anti-HA tag (Biolegend, CA, USA), anti-rabbit

IgG (H+L) HRP (Jackson Immuno Research Laboratories, MD, USA), anti-mouse IgG (H+L) HRP

(Thermo Fisher Scientific), and anti-GAPDH (Fujifilm). The antibodies used in this study are listed

in S2 Table.

Purification of intracellular HBV rcDNA

To extract intracellular HBV DNA, cells were washed twice with PBS, and cells were lysed in

lysis buffer (100 mM Tris-HCl [pH 8.0], 0.2% NP-40) for 15 minutes at 4°C. After spinning at

13,000 rpm for 1 minute at room temperature, the supernatant was collected and incubated with 1 M

MgCl2, 0.2 mg/ml of DNase I (Sigma), and 10 mg/ml of RNase A (Nacalai Tesque) for at least 3

hours at 37°C. After adding 500 mM EDTA and 5 M NaCl, the lysates were digested with 20 mg/ml

of proteinase K (Kanto Chemical, Tokyo, Japan) and 10% SDS for 5 hours at 55°C. The extracted

HBV DNA was purified using phenol-chloroform-isoamyl alcohol (Sigma), then precipitated with

isopropanol, 0.5 mg/ml of glycogen, and 3M sodium acetate. After washing with 70% ethanol, the

purified HBV DNA was resolved in 50 μl pure water. HBV rcDNA was detected by qPCR using

Power SYBR Green PCR Master Mix (Applied Biosystems, CA, USA). As a standard for rcDNA,

pUC19-C_JPNAT was used. qPCR procedures were following the manufacturer's protocol. Briefly,

the samples were incubated at 50℃ for 2 minutes and at 95℃ for 10 minutes to initial denaturation.

Then, the samples were incubated at 95℃ for 15 seconds and at 60℃ for 1 minute for 40 cycles to

polymerase reaction. The following primer pair was used in this study.

rcDNA: 5'- GGAGGGATACATAGAGGTTCCTTGA -3' (forward)

5'- GTTGCCCGTTTGTCCTCTAATTC -3' (reverse)

Quantification of intracellular HBV cccDNA level

To extract intracellular HBV cccDNA, the cells were lysed by lysis buffer (10mM Tris-HCl [pH

7.4], 10mM EDTA, 0.5% SDS) and 5M NaCl was added. After incubation overnight at 4℃, the

samples were spun at 15,000 rpm for 30 minutes at 4℃ and the supernatant was collected to isolate

protein-free DNA. The DNA was purified a total of three times by phenol (Nacalai Tesque),

phenol-chloroform-isoamyl, and chloroform (Fujifilm). After precipitation with isopropanol and

ethanol, Plasmid-Safe ATP-Dependent DNase (Lucigen, WI, USA) was added to the purified DNA

to remove double-stranded and circular DNA. The Purified DNA was extracted in 30 μl of 10mM

Tris-Cl (pH 8.5) using the FastGene Gel/PCR Extraction Kit (NIPPON Genetics, Tokyo, Japan).

HBV cccDNA was measured by qPCR using TaqMan Fast Advanced Master Mix (Applied

Biosystems, CA, USA). As a standard for cccDNA, pUC19-C_JPNAT was used. qPCR procedures

were done per the manufacturer's protocol. In brief, the samples were incubated at 50℃ for 2

minutes and at 95℃ for 10 minutes to initiate denaturation. Then, the samples were incubated at

95℃ for 15 seconds and at 60℃ for 1 minute for 50 cycles for the polymerase reaction. The

following primer pair and probe were used in this study.

cccDNA: 5'- CGTCTGTGCCTTCTCATCTGC -3' (forward)

523   5'- GCACAGCTTGGAGGCTTGAA -3' (reverse)

524   5'- CTGTAGGCATAAATTGGT -3' (Probe)

Measurement of intracellular HBV pgRNA

Total RNA including pgRNA was extracted using the RNeasy Mini Kit (Qiagen, Hilden,

Germany) following the manufacturer's protocol. The extracted RNA was resolved in 50 μl pure

water and measured by RT-qPCR using Power SYBR Green RNA-to-Ct 1-step Kit (Applied

Biosystems). The pgRNA expression level was normalized against the expression level of GAPDH

as a housekeeping gene and calculated by the ΔΔCT method. RT-qPCR procedures were done

following manufacturer's protocol. First, the samples were incubated at 50℃ for 2 minutes and at

95℃ for 10 minutes to intiate denaturation. Then, the samples were incubated at 95℃ for 15

seconds and at 60℃ for 1 minute for 40 cycles for the polymerase reaction. Samples were then

incubated at 95℃ for 15 seconds, 60℃ for 1 minute, and 95℃ for 15 seconds for the melting curve

measurement. The following primer pairs were used in this study.

pgRNA: 5'- TCCCTCGCCTCGCAGACG -3' (forward)

5'- GTTTCCCACCTTATGAGTC -3' (reverse)

GAPDH: 5'- AGAAGGCTGGGGCTCATTTG -3' (forward)

5'- AGGGGCCATCCACAGTCTTC -3' (reverse)

pgRNA degradation assay

HepAD38.7 cells were seeded in collagen type I-coated 12-well plates (Iwaki) and cultured

overnight in media containing tetracycline hydrochloride. After washing twice with PBS, cells were

cultured without tetracycline for 48 hours. Then, the cells were transfected with N4BP1 wild-type

expression plasmid, ΔKH expression plasmid, ΔRNase expression plasmid, or control vector. These

plasmids were transfected as follows: 100 μl of Opti-MEM, 6 μl of TransIT LT-1, and 3μg of

expression plasmid. After transfection of plasmids, cells were cultured for 48 hours and changed to a

tetracycline-containing media in order to stop HBV production. Cells were collected at 2, 3, and 4

days after changing to tetracycline-containing media. Total RNA, including pgRNA, was extracted

as described above for RT-qPCR.

Drug treatment assay

Huh7 cells, HepG2-hNTCP C4 cells, and PHHs were stimulated with 1000 U/ml of IFN-α

(PBL Assay Science, NJ, USA) or 100 ng/ml of IFN-λ (Pepro Tech, NJ, USA). At 6, 24 and 48 hours

post-drug stimulation, total RNA in cells was extracted for qPCR using the RNeasy Mini Kit, and

cDNA was synthesized using SuperScript IV VILO Master Mix (Invitrogen). For WB, the cells were

lysed with $2 \times$ Laemmli sample buffer containing 10% 2-mercaptoethanol and processed as

described above. qPCR was performed using Power SYBR Green PCR Master Mix. The mRNA

expression levels of N4BP1 and ISG15 were normalized against the expression level of GAPDH as a

housekeeping gene and calculated by the $\Delta\Delta$CT method. The following primer pairs were used in

this study.

N4BP1: 5'- CCCGATGATCCTCTGGGAAG -3' (forward)

5'- TTTGGCAGGGCACTGAGTAG -3' (reverse)

ISG15: 5'- ACTCATCTTTGCCAGTACAGGAG -3' (forward)

5'- CAGCATCTTCACCGTCAGGTC -3' (reverse)

GAPDH: see above

Analysis of N4BP1 expression in clinical liver samples

To assess the N4BP1 expression of HBV-HCC resection samples, we confirmed that all

specimens were HBs-Ag (+), HBe-Ag (-), and HBe-Ab (+) as documented in the patient medical

records. Based on HBV DNA data in the serum, the samples were divided into an HBV DNA

positive group (n=9) and an HBV DNA negative group (n=8). In all cases, RNA was extracted from

the non-cancerous areas of resected tissues using the RNeasy Mini Kit and *N4BP1* and *ISG15*

mRNA expression were measured by qPCR. This study was approved by the Institutional Review

Board of the Graduate School of Medicine, Hokkaido University (No. 14-015, 022-0195). All

patients participating in this study provided written informed consent for the use of their samples and

clinical data and they were informed of the opportunity to opt out.

Northern blotting

Huh7 cells were co-transfected with a plasmid containing the 1.3-fold-overlength genome of

HBV genotype C and N4BP1 using TransIT LT-1 transfection reagent, and cells were collected at 3

d.p.t.. Total RNA was extracted using Isogen (Nippon Gene, Tokyo, Japan) and subjected to

electrophoresis in 1% agarose-formaldehyde gel and morpholinepropanesulfonic acid buffer

(Nacalai Tesque). Ribosomal RNA was visualized by ethidium bromide staining, and

electrophoresed RNA was transferred onto a positively charged nylon membrane (Roche,

Basel-Stadt, Switzerland). An HBV RNA probe and GAPDH RNA probe were linearized and

synthesized by digoxigenin RNA labeling kit (Roche). Hybridization and detection were performed

using the DIG Northern Starter kit (Roche) following the manufacturer's protocol. The signals were

detected by LuminoGraph I (Atto).

RNA Immunoprecipitation (RIP) assay

Huh7 cells were co-transfected with a plasmid containing the 1.3-fold-overlength genome

of HBV genotype C and HA-tagged N4BP1 wildtype or domain deficient mutants or HA control

vector using TransIT LT-1 transfection reagent, and cells were collected at 3 d.p.t. Cells were

collected and immunoprecipitated with anti-HA-tag monoclonal antibody magnetic beads (Medical

& Biological Laboratories Co., Ltd., Tokyo, Japan). Immunoprecipitation and RNA extraction were

performed using RiboCluster Profiler RIP-Assay Kit (Medical & Biological Laboratories Co., Ltd.,

Tokyo, Japan) following the manufacture's protocol. Western blotting and pgRNA qPCR were

measured as describe above.

HBeAg CLIA, HBs Ag ELISA and HBcrAg ELISA

N4BP1-overexpressing PHHs were infected with HBV genotype C (5 Geq/cell). At 14

d.p.i., RNA was extracted from the cells and pgRNA was measured as described above. HBV

proteins in the supernatant were assessed as follows: HBeAg CLIA (Ig Biotechnology, Burlingame,

USA); HBs S Antigen Quantitative ELISA Kit, Rapid-II (Takara Bio) following manufacturers'

protocol; and HBcr Ag assay (42) by SRL, Inc. (Tokyo, Japan).

RNA sequencing

PHHs were analyzed for RNA sequencing. N4BP1-overexpressing PHHs were infected

with HBV genotype C (5 Geq/cell) as described above. At 3 d.p.i., total RNA in PHHs was extracted

using RNeasy Mini Kit following the manufacturer's protocol. RNA sequencing was performed by

Osaka University (Genome Information Research Center, Research Institute for Microbial Diseases,

Osaka University, Osaka, Japan). We ran 2 well (N4BP1-overexpressing) vs 2 well (control) samples.

Whole transcriptome sequencing was applied to RNA samples using the Illumina NovaSeq 6000

platform in a 101 bp single-end mode. Sequenced reads were mapped to the human reference genome

sequence (hg19) using TopHat version 2.2.1 in combination with Bowtie 2 version 2.2.3 and

SAMtools version 1.0. The number of fragments per kilobase of exon per million mapped fragments

was calculated using Cufflinks version 2.2.1. Following the exclusion of genes with expression

counts below 50, differential expression analysis was conducted utilizing the DESeq2 package

(version 1.42.0) (43). Genes were considered significantly differentially expressed if they satisfied

the criteria of an adjusted P-value less than 0.1 and an absolute fold change greater than 2.

Subsequently, pathway enrichment analysis of the differentially expressed genes was performed

using the Kyoto Encyclopedia of Genes and Genomes database (44) with the clusterProfiler package

(version 4.10.0) (45). Plots were generated using the ggplot2 package (version 3.5.1) (46). To review

GEO data, access https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE225322. The accession

No. is GSE225322.

Statistical analysis

Data are averages of three or more independent experiments. All data in figures are presented as

the mean ± standard deviation (SD). Evaluation of data was performed by Mann-Whitney *U* test.

P-value of < 0.05 was considered statistically significant. Statistical analysis was performed using

JMP Pro version 16.

<u>**Data Availability**</u>

The GEO data that support the findings of this study are openly available in

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE225322. The accession No. is

GSE225322.

<u>**Acknowledgements**</u>

We thank Haruko Kubo, Mizuho Tetsuka, and Saya Shimamura for their secretarial work and

Mikiko Ishibashi, Minako Hommura and Hiroko Azuma for their technical assistance. We are

grateful to Takaji Wakita (NIID) for providing HepG2-hNTCP C4 cells and HepAD38.7 cells. We

also thank Jenna Gaska for English proofreading.

This work was supported in part by the Japan Agency for Medical Research and Development

(AMED) under Grant Numbers 22fk0310501h0001, 22fk0310506h0001, 22fk0310513h0001,

22fk0310524h0001, and 22fk0310523h0001; AMED CREST under Grant Number JP22gm1610008;

JSPS KAKENHI Fund for the Promotion of Joint International Research (International Leading

Research) under grant number JP23K20041.

Conceptualization, Y. M., O. T., A. T., and T. F.; Data curation, N. K., S. S., R. S., T. T., D. O., and T.

F.; Formal analysis, N. K., S. S., R. S., T. T., D. O., K. M., and T. F.; Funding acquisition, T. F.;

Investigation, N. K., S. S., Y. S., T. I., K. N., D. O., K. M., L. L., Y. K., and Y. K.; Methodology, N.

K., S. S., R. S., T. T., S. H., Y. T., A. T., and T. F.; Project administration, A. T. and T. F.; Resources,

D. O., Y. K., Y. M., O. T., A. T., and T. F.; Software, D. O.; Supervision, A. T., T. F.; Validation, N. K.,

S. S., Y. S., T. S., T. I., and K. N.; Visualization, N. K., S. S., and T. F.; Writing - original draft, N. K.,

S. S., D. O., Y. K., K. M., and T. F.; Writing - review & editing, N. K., S. S., R. S., T. T., Y. S., T. S.,

T. I., K. N., D. O., Y. K., Y. M., O. T., A. T., and T. F. All authors have approved the submitted

version of the manuscript and agreed to be accountable for any part of the work.

<u>References</u>

1.   World Health Organisation. Hepatitis B Fact Sheet 2021 [updated 24 June 202213 Feb

2023]. Available from: https://www.who.int/en/news-room/fact-sheets/detail/hepatitis-b.

2.   World Health Organisation. Immunization Coverage 2021 [updated 14 July 202213 Feb

2023]. Available from: https://www.who.int/news-room/fact-sheets/detail/immunization-coverage

3.   Koniger C, Wingert I, Marsmann M, Rosler C, Beck J, Nassal M. 2014. Involvement of

the host DNA-repair enzyme TDP2 in formation of the covalently closed circular DNA persistence

reservoir of hepatitis B viruses. Proc Natl Acad Sci U S A. 2014;111(40):E4244-53.

4.   Qi Y, Gao Z, Xu G, Peng B, Liu C, Yan H, Yao Q, Sun G, Liu Y, Tang D, Song Z, He W,

Sun Y, Guo JT, Li W. 2016. DNA Polymerase kappa Is a Key Cellular Factor for the Formation of

Covalently Closed Circular DNA of Hepatitis B Virus. PLoS Pathog. 2016;12(10):e1005893.

5.   Sheraz M, Cheng J, Tang L, Chang J, Guo JT. 2019. Cellular DNA Topoisomerases Are

Required for the Synthesis of Hepatitis B Virus Covalently Closed Circular DNA. J Virol.

2019;93(11).

6.   Kitamura K, Que L, Shimadu M, Koura M, Ishihara Y, Wakfae K, Nakamura T, Watashi K,

Wakita T, Muramatsu M. 2018. Flap endonuclease 1 is involved in cccDNA formation in the

hepatitis B virus. PLoS Pathog. 2018;14(6):e1007124.

7.   Rijckborst V, Janssen HL. 2010. The Role of Interferon in Hepatitis B Therapy. Curr Hepat

Rep. 2010;9(4):231-8.

8.   Doong SL, Tsai CH, Schinazi RF, Liotta DC, Cheng YC. 1991. Inhibition of the

replication of hepatitis B virus in vitro by 2',3'-dideoxy-3'-thiacytidine and related analogues. Proc

Natl Acad Sci U S A. 1991;88(19):8495-9.

9.   Li Q, Sun B, Zhuo Y, Jiang Z, Li R, Lin C, Jin Y, Gao Y, Wang D. 2022. Interferon and

interferon-stimulated genes in HBV treatment. Front Immunol. 2022;13:1034968.

10.   Tang LSY, Covert E, Wilson E, Kottilil S. Chronic Hepatitis B Infection: A Review. 2018.

JAMA. 2018;319(17):1802-13.

11.   Dienstag JL, Perrillo RP, Schiff ER, Bartholomew M, Vicary C, Rubin M. 1995. A

preliminary trial of lamivudine for chronic hepatitis B infection. N Engl J Med.

1995;333(25):1657-61.

12.   Nowak MA, Bonhoeffer S, Hill AM, Boehme R, Thomas HC, McDade H. 1996. Viral

dynamics in hepatitis B virus infection. Proc Natl Acad Sci U S A. 1996;93(9):4398-402.

13.   Suzuki F, Akuta N, Suzuki Y, Sezaki H, Arase Y, Hosaka T, Someya T, Kobayashi M,

Saitoh S, Ikeda K, Kobayashi M, Matsuda M, Satoh J, Watahiki S, Kumada H. Clinical and

virological features of non-breakthrough and severe exacerbation due to lamivudine-resistant

hepatitis B virus mutants. J Med Virol. 2006;78(3):341-52.

14.   Mao R, Nie H, Cai D, Zhang J, Liu H, Yan R, Cuconati A, Block T, Guo J, Guo H. 2013.

Inhibition of hepatitis B virus replication by the host zinc finger antiviral protein. PLoS Pathog.

2013;9(7):e1003494.

15.   Li M, Yang J, Zhao Y, Song Y, Yin S, Guo J, Zhang H, Wang K, Wei L, Li S, Xu W. 2020.

MCPIP1 inhibits Hepatitis B virus replication by destabilizing viral RNA and negatively regulates

the virus-induced innate inflammatory responses. Antiviral Res. 2020;174:104705.

16.   Sun S, Nakashima K, Ito M, Li Y, Chida T, Takahashi H, Watashi K, Sawasaki T, Wakita T,

Suzuki T. 2017. Involvement of PUF60 in Transcriptional and Post-transcriptional Regulation of

Hepatitis B Virus Pregenomic RNA Expression. Sci Rep. 2017;7(1):12874.

17.   Tang H, Huang Y, Chen J, Yu C, Huang AL. 2008. Cellular protein TIA-1 regulates the

expression of HBV surface antigen by binding the HBV posttranscriptional regulatory element.

Intervirology. 2008;51(3):203-9.

18.   Xu W, Ma C, Zhang Q, Zhao R, Hu D, Zhang X, Chen J, Liu F, Wu K, Liu Y, Wu J. 2018.

PJA1 Coordinates with the SMC5/6 Complex To Restrict DNA Viruses and Episomal Genes in an

Interferon-Independent Manner. J Virol. 2018;92(22).

19.   Cai J, Yang L, Wang B, Huang Y, Tang J, Lu Y, Wu Z, Jian J. 2014. Identification of a

novel N4BP1-like gene from grass carp (Ctenopharyngodon idella) in response to GCRV infection.

Fish Shellfish Immunol. 2014;36(1):223-8.

20.   Yamasoba D, Sato K, Ichinose T, Imamura T, Koepke L, Joas S, Reith E, Hotter D,

Misawa N, Akaki K, Uehata T, Mino T, Miyamoto S, Noda T, Yamashita A, Standley D, Kirchhoff F,

Sauter D, Koyanagi Y, Takeuchi O. 2019. N4BP1 restricts HIV-1 and its inactivation by MALT1

promotes viral reactivation. Nat Microbiol. 2019;4(9):1532-44.

21.   Iwamoto M, Watashi K, Tsukuda S, Aly H. H, Fukasawa M, Fujimoto A, Suzuki R, Aizaki

H, Ito T, Koiwai O, Kusuhara H, Wakita T. 2014. Evaluation and identification of hepatitis B virus

entry inhibitors using HepG2 cells overexpressing a membrane transporter NTCP. BioChem Biophys

Res Commun. 2014;443(3):808-13.

22.   Köing A, Yang J, Jo E, Park HPK, Kim H, Than TT, Song X, Qi X, Dai X, Park S, Shum D,

Ryu WS, Kim JH, Yoon KS, Park YJ, Ahn HS, Han KH, Gerlich HW, Windisch PM. 2019. Efficient

long-term amplification of hepatitis B virus isolates after infection of slow proliferating

HepG2-NTCP cells. J Hepatol. 2019; 71(2):289-300.

23.   Honkoop P, Niesters HG, de Man RA, Osterhaus AD, Schalm SW. 1997. Lamivudine

resistance in immunocompetent chronic hepatitis B. Incidence and patterns. J Hepatol.

1997;26(6):1393-5.

24.   Tipples GA, Ma MM, Fischer KP, Bain VG, Kneteman NM, Tyrrell DL. 1996. Mutation in

HBV RNA-dependent DNA polymerase confers resistance to lamivudine in vivo. 1996. Hepatology.

1996;24(3):714-7.

25.   Bartholomew MM, Jansen RW, Jeffers LJ, Reddy KR, Johnson LC, Bunzendahl H,

Condreay L, Tzakis A, Schiff E, Brown N. 1997. Hepatitis-B-virus resistance to lamivudine given

for recurrent infection after orthotopic liver transplantation. Lancet. 1997;349(9044):20-2.

26.   Valverde R, Edwards L, Regan L. 2008. Structure and function of KH domains. FEBS J.

2008;275(11):2712-26.

27.   Hollingworth D, Candel AM, Nicastro G, Martin SR, Briata P, Gherzi R, Ramos A. 2012.

KH domains with impaired nucleic acid binding as a tool for functional analysis. Nucleic Acids Res.

2012;40(14):6873-86.

28.   Anantharaman V, Aravind L. 2006. The NYN domains: novel predicted RNAses with a

PIN domain-like fold. RNA Biol. 2006;3(1):18-27.

29.   Shi H, Sun L, Wang Y, Liu A, Zhan X, Li X, Tang M, Anderton P, Hildebrand S, Quan J,

Ludwig S, Moresco EM, Beutler B. 2021. N4BP1 negatively regulates NF-kappaB by binding and

inhibiting NEMO oligomerization. Nat Commun. 2021;12(1):1379.

30.   Gitlin AD, Heger K, Schubert AF, Reja R, Yan D, Pham VC, Suto E, Zhang J, Kwon YC,

Freund EC, Kang J, Pham A, Caothien R, Bacarro N, Hinkle T, Xu M, McKenzie BS, Haley B, Lee

WP, Lill JR, Roose-Girma M, Dohse M, Webster JD, Newton K, Dixit VM. 2020. Integration of

innate immune signalling by caspase-8 cleavage of N4BP1. Nature. 2020;587(7833):275-80.

31.   Cui X, Li Y, Xu H, Sun Y, Jiang S, Li W. 2023. Characteristics of Hepatitis B virus

integration and mechanism of inducing chromosome translocation. NPJ Genom Med. 2023;8(1):11.

32.   Gu H, Zheng S, Han G, Yang H, Deng Z, Liu Z, He F. 2022. Porcine Reproductive and

Respiratory Syndrome Virus Adapts Antiviral Innate Immunity via Manipulating MALT1. mBio.

2022;13(3):e0066422.

33.   Mino T, Takeuchi O. 2021. Regnase-1-related endoribonucleases in health and

immunological diseases. Immunol Rev. 2021;304(1):97-110.

34.   Melén K, Keskinen P, Lehtonen A, Julkunen I. 2000. Interferon-induced gene expression

and signaling in human hepatoma cell lines. J Hepatol. 2000;33(5):764-72.

35.   Marozin S, Altomonte J, Stadler F, Thasler WE, Schmid RM, Ebert O. 2008. Inhibition of

the IFN-β Response in Hepatocellular Carcinoma by Alternative Spliced Isoform of IFN Regulatory

Factor-3. Molecular Therapy. 2008;16(11):1789-97.

36.   Speer SD, Li Z, Buta S, Payelle-Brogard B, Qian L, Vigant F, Rubino E, Gardner T,

Wedeking T, Hermann M, Duehr J, Sanal O, Tezcan I, Mansouri N, Tabarsi P, Mansouri D,

Francois-Newton V, Daussy CF, Rodriguez MR, Lenschow DJ, Freiberg AN, Tortorella D, Piehler J,

Lee B, Garcia-Sastre A, Pellegrini S, Bogunovic D. 2016. ISG15 deficiency and increased viral

resistance in humans but not mice. Nat Commun. 2016;7:11496.

37.   Dickensheets H, Sheikh F, Park O, Gao B, Donnelly RP. 2013. Interferon-lambda (IFN-λ)

induces signal transduction and gene expression in human hepatocytes, but not in lymphocytes or

monocytes. J Leukoc Biol. 2013;93(3):377-85.

38.   Nakai M, Komiya K, Murata M, Kimura T, Kanaoka M, Kanegae Y, Saito I. 2007.

Expression of pIX gene induced by transgene promoter: possible cause of host immune response in

first-generation adenoviral vectors. Hum Gene Ther. 2007;18(10):925-36.

39.   Suzuki M, Kondo S, Pei Z, Maekawa A, Saito I, Kanegae Y. Preferable sites and

orientations of transgene inserted in the adenovirus vector genome: The E3 site may be unfavorable

for transgene position. Gene Ther. 2015;22(5):421-9.

40.   Pei Z, Kondo S, Kanegae Y, Saito I. Copy number of adenoviral vector genome transduced

into target cells can be measured using quantitative PCR: application to vector titration. Biochem

Biophys Res Commun. 2012;417(3):945-50.

41.   Watashi K, Liang G, Iwamoto M, Marusawa H, Uchida N, Daito T, Kitamura K,

Muramatsu M, Ohashi H, Kiyohara T, Suzuki R, Li J, Tong S, Tanaka Y, Murata K, Aizaki H, Wakita

T. 2013. Interleukin-1 and tumor necrosis factor-alpha trigger restriction of hepatitis B virus

infection via a cytidine deaminase activation-induced cytidine deaminase (AID). J Biol Chem.

2013;288(44):31715-27.

42.   Inoue T, Kusumoto S, Iio E, Ogawa S, Suzuki T, Yagi S, Kaneko A, Matsuura K, Aoyagi

K, Tanaka Y. 2021. Clinical efficacy of a novel, high-sensitivity HBcrAg assay in the management of

chronic hepatitis B and HBV reactivation. J Hepatol. 2021;75(2):302-10.

43.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol 15:550.

44.   Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. 1999. KEGG: Kyoto

Encyclopedia of genes and genomes. Nucleic Acids Res 27:29–34.

45.   Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L, Fu X, Liu S,

Bo X, Yu G. 2021. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.

Innovation (Camb) 2:100141.

46.   Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New

York. ISBN 978-3-319-24277-4, https://ggplot2.tidyverse.org.

Figure legends

Fig 1. Screening for RNA binding proteins (RBPs) that inhibit HBV propagation.

(A) A plasmid containing the 1.3-fold-overlength genome of HBV genotype C and 132 RBP

expression plasmids were co-transfected into Huh7 cells. Intracellular rcDNA levels were assessed at

3 d.p.t.. Using the 50% of rcDNA level of the control as a cutoff, 23 out of the 132 RBPs suppressed

HBV rcDNA levels. (B) HepG2-hNTCP C4 cells overexpressing the 23 RBPs were infected with

HBV genotype D. At 14 d.p.i., cells were harvested and intracellular rcDNA levels were measured

by qPCR. Seven out of the 23 RBPs (N4BP1, TIA-1, ZC3H12B, PJA1, HNRNPC, ZCCHC10, and

KHNYN) reduced rcDNA levels in comparison to the control. (C) WB analyses in naïve Huh7 cells,

HepG2-hNTCP C4 cells of the five factors (N4BP1, ZC3H12B, HNRNPC, ZCCHC10, and

KHNYN) identified in the secondary screening. As a positive control, WB were performed on Huh7

cells transfected to overexpress each factor. Error bars represent the SD. Abbreviations: RBP (RNA

binding protein), ETV (entecavir), HepG2 C4 (HepG2-hNTCP C4), OE (overexpression)

Fig 2. Measurement of rcDNA levels after RBP knockdown.

(A) Huh7 cells were transfected with siRNA, then co-transfected with the 1.3-fold-overlength

genome of HBV genotype C and an RBP expression plasmid. WB analyses and qPCR of rcDNA

were performed at 3 d.p.t.. ZCCHC10 (A), HNRNPC (B), and N4BP1 (C). GAPDH was used as a

loading control and ETV was used as a positive control. Error bars represent the SD. *P < 0.05, **P

< 0.01, as determined by Mann-Whitney U test. All qPCR experiments were independently

performed at least three times. Abbreviations: NS (not significant)

Fig 3. N4BP1 is a suppressive host factor of HBV replication.

(A) Plasmids containing 1.3-fold-overlength genomes of HBV genotypes A, B, or C were

co-transfected with N4BP1 expression plasmids into Huh7 cells. At 3 d.p.t., the rcDNA levels were

measured by qPCR. Genotype A (left), genotype B (middle), and genotype C (right). (B) WB

analysis and rcDNA levels of N4BP1 KO Huh7 cells. N4BP1 KO Huh7 cells were transfected with

the plasmid containing 1.3-fold-overlength genomes C and WB and qPCR were performed at 3 d.p.t..

N4BP1 expression by WB (upper, anti-N4BP1). GAPDH was used as a loading control (lower,

anti-GAPDH). rcDNA levels were increased relative to control in both N4BP1-KO lines that were

generated. N4BP1 KO Huh7 cell lines #1 (C) and #2 (D) were transfected with the N4BP1 plasmid

to restore N4BP1 expression (upper, anti-N4BP1). GAPDH was used as a loading control (lower,

anti-GAPDH). In the N4BP1-rescued KO Huh7 cells, the level of rcDNA was significantly lower

than in the non-rescued N4BP1 KO Huh7 cells. Error bars represent the SD. *$P < 0.05$, **$P < 0.01$,

as determined by Mann-Whitney U test. All qPCR experiments were independently performed at

least three times.

Fig 4. N4BP1 overexpression affected expression of several host factors

(A) Heat map of the transcriptomic profiles of N4BP1-expressing and control PHHs infected with

HBV. (B) In the volcano plot, we set a cutoff value of gene expression at absolute fold change

greater than 2 and an adjusted P-value less than 0.1 compared to control cells. Relative to the control

cells, 178 factors were upregulated, and 230 factors downregulated in the N4BP1-overexpressing

PHHs. (C) Pathway enrichment analysis of the differentially expressed genes in

N4BP1-overexpressing PHHs versus the control PHHs demonstrates the distinct differences in

pathways induced by N4BP1-overexpression. The graph depicts the number of genes corresponding

to each pathway through the size of its dots. The color gradient, which transitions from red to purple,

illustrates the adjusted p-value indicating significance of enrichment. The Gene Ratio on the

horizontal axis indicates the proportion of input genes that are annotated to each term, thereby

highlighting the distinct impact of N4BP1 over-expression. The raw data sets are shown in

Supporting Information S1 table.

Fig 5. N4BP1 expression is not induced by HBV.

(A) We evaluated whether N4BP1 expression is induced by HBV infection or transfection in

hepatocyte cell lines. In each system, we confirmed by qPCR whether pgRNA was induced. (B)

Huh7 cells were transfected with a plasmid containing the 1.3-fold-overlength genome of HBV

genotype C and cells were collected at 3 d.p.t. (upper). *N4BP1* mRNA expression was assessed by

qPCR (middle) and protein expression by WB (lower). (C) HepG2-hNTCP C4 cells were infected

with HBV genotype D and cells were collected at 3 d.p.i. and 10 d.p.i. (upper). N4BP1 expression

level by qPCR (middle). N4BP1 expression levels by WB (lower). (D) Tetracycline was removed

from the medium of HepAD38.7 cells and the cells were collected 3 days later (upper). N4BP1

expression levels by qPCR (middle). N4BP1 expression levels by WB (lower). Error bars represent

the SD. *P < 0.05, **P < 0.01, as determined by Mann-Whitney U test. All qPCR experiments were

independently performed at least three times.

Fig 6. KH-like domain and RNase domain are involved in the anti-HBV effect of N4BP1.

(A) Schematic of N4BP1. N4BP1 is composed of 896 amino acids and has both a KH-like domain

and an RNase domain. We constructed a KH-like domain deficient mutant (deletion of amino acids

59-143) and an RNase domain deficient mutant (deletion of amino acids 617-769). Each plasmid had

a HA tag at its C-terminus. (B) Huh7 cells were transfected with the plasmid containing the

1.3-fold-overlength genome of HBV genotype C and each expression plasmid (control vector,

N4BP1 WT, ΔKH, and ΔRNase) and cells were collected at 3 d.p.t. WB was used to confirm the

expression of each N4BP1 mutant (upper, anti-HA tag). GAPDH was used as a loading control

(lower, anti-GAPDH). HBV levels were assessed by measuring intracellular rcDNA levels at 3 d.p.t.

(C) Huh7 cells were transfected with the plasmid containing the 1.3-fold-overlength genome of HBV

genotype C and each expression plasmid (control vector, WT, and D623N) and cells were collected

at 3 d.p.t. WB was used to confirm the expression of each N4BP1 mutant (upper, anti-HA tag).

GAPDH was used as a loading control (lower, anti-GAPDH). HBV levels were assessed by

measuring intracellular rcDNA levels at 3 d.p.t. ETV was used as a positive control. Error bars

represent the SD. *$P < 0.05$ as determined by Mann-Whitney U test. All qPCR experiments were

independently performed at least three times.

Fig 7. N4BP1 degrades HBV RNAs by binding HBV RNAs

HepG2-hNTCP C4 cells were infected with HBV genotype D and cells were collected at 14 d.p.i.

Comparison of cccDNA (A), pgRNA (B), and rcDNA (C) levels in N4BP1-overexpressing

HepG2-hNTCP C4 cells and control at 14 d.p.i. Heparin was used as a positive control. (D) Time

course of pgRNA degradation assay in HepAD38.7 cells. (E) pgRNA levels over time in HepAD38.7

cells overexpressing N4BP1 WT, ΔKH, ΔRNase, or control. (F) Northern blotting analyses in control,

N4BP1-overexpressing Huh7 cells transfected with the 1.3-fold-overlength genome of HBV and

naïve Huh7 cells (mock). 28S and 18S ribosomal RNA was used as a loading control (lower panel).

(G) Huh7 cells were co-transfected with the 1.3-fold-overlength genome of HBV genotype C with

HA control vector, HA-tagged N4BP1 or ΔRNase. Cells were collected at 3 d.p.t. and RIP assay was

performed. WB was used to confirm the expression of each N4BP1 mutant (Blot: HA) and IP was

performed. GAPDH was used as a loading control (lower, anti-GAPDH). (H) RNA was extracted

from total cell lysate and the immunoprecipitated protein, and then pgRNA was measured by qPCR.

(I) PHHs were infected with HBV at 5 Geq/cell and supernatant was collected at 14 d.p.i. RNA was

extracted from the cells and pgRNA was measured and N4BP1 overexpression was confirmed by

western blot (left). The viral protein levels in supernatant were measured by HBeAg CLIA (second

from left), HBsAg ELISA (second from right) and HBcrAg assay (right). Error bars represent the SD.

*P < 0.05, **P < 0.01, as determined by Mann-Whitney U test. All experiments were independently

performed at least three times.

Fig 8. N4BP1 expression is induced by IFN-α and IFN-λ in PHHs.

(A) Time course of the drug treatment assay. Cells were harvested at 6, 24 or 48 hours post-IFN-α/λ

treatment. (B) N4BP1 expression levels by qPCR in Huh7 cells, HepG2-hNTCP C4 cells, and PHHs

treated with IFN-α (upper). ISG15 expression levels by qPCR (middle). N4BP1 protein expression

levels by WB. GAPDH was used as a loading control (lower). (C) N4BP1 expression levels by

qPCR in Huh7 cells, HepG2-hNTCP C4 cells, and PHHs treated with IFN-λ (upper). ISG15

expression levels by qPCR (middle). N4BP1 protein expression levels by WB. GAPDH was

measured as loading control (lower).

Fig 9. ISG15 and N4BP1 expression levels in HBV-infected human liver tissues.

The mRNA levels of *ISG15* (A) and *N4BP1* (B) from non-cancerous tissues of HBV-HCC patients

with HBs-Ag (+), HBe-Ag (-), HBe-Ab (+) were examined by qPCR. The samples were divided into

HBV DNA positive group (n=9) and HBV DNA negative group (n=8). Error bars represent the SD.

*$P < 0.05$, as determined by Mann-Whitney U test.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

A

seed   IFN-α/λ   Harvest   Harvest   Harvest

-1   0   6h   24h   48h   (day)

B

IFN-α

Huh7   C4   PHH

N4BP1 mRNA expression

ISG15 mRNA expression

(kDa)
100-
37-

Blot :
N4BP1
GAPDH

C

IFN-λ

Huh7   C4   PHH

N4BP1 mRNA expression

ISG15 mRNA expression

(kDa)
100-
37-

Blot :
N4BP1
GAPDH

Fig. 9