# Depletion of SMN Protein in Mesenchymal Progenitors Impairs the Development of Bone and Neuromuscular Junction in Spinal Muscular Atrophy

**Sang-Hyeon Hann[1]\*, Seon-Yong Kim[2]\*, Ye-Lynne Kim[1], Young-Woo Jo[1], Jong-Seol Kang[1], Hyerim Park[1], Se-Young Choi[2#] and Young-Yun Kong[1#]**

[1]School of Biological Sciences, Seoul National University, Seoul 08826, Republic of Korea

[2]Department of Physiology, Dental Research Institute, Seoul National University School of Dentistry, Seoul, 03080, Republic of Korea

[3]

\*These authors contributed equally to this work

#Correspondence and requests for materials should be addressed to Y.Y.K. (ykong@snu.ac.kr) and S.Y.C. (sychoi@snu.ac.kr)

**Abstract**

Spinal Muscular Atrophy (SMA) is a neuromuscular disorder characterized by the deficiency

of the survival motor neuron (SMN) protein, which leads to motor neuron dysfunction and

muscle atrophy. In addition to the requirement for SMN in motor neurons, recent studies

suggest that SMN deficiency in peripheral tissues plays a key role in the pathogenesis of

SMA. Using limb mesenchymal progenitor cells (MPCs)-specific SMN-depleted mouse

models, we reveal that SMN reduction in chondrocytes and fibro-adipogenic progenitors

(FAPs) derived from limb MPCs causes defects in the development of bone and

neuromuscular junction (NMJ), respectively. We show that impaired growth plate

homeostasis, which causes skeletal growth defects in SMA, is cell-autonomous due to SMN

ablation in chondrocytes. Furthermore, the reduction of SMN in FAPs resulted in abnormal

NMJ maturation, altered release of neurotransmitters, and NMJ morphological defects.

Transplantation of healthy FAPs rescued the morphological deterioration. Our findings

highlight the significance of mesenchymal SMN in neuromusculoskeletal pathogenesis in

SMA and provide insights into potential therapeutic strategies targeting mesenchymal cells

for the treatment of SMA.

## Introduction

The survival motor neuron (SMN) protein is a crucial component of the spliceosome complex and is essential for the proper function of all cell types (Mercuri et al., 2022). Deficiency in SMN protein disrupts the formation of spliceosome complexes, ultimately causing splicing defects in multiple genes. Mutations in the SMN1 gene, which encodes the SMN protein, give rise to the neuromuscular disorder Spinal Muscular Atrophy (SMA). SMA is characterized by neuromuscular junctions (NMJ) disruption, muscular atrophy, and alpha motor neuron loss (Mercuri et al., 2022; Burghes et al., 2009). The severity of the disease in humans correlates with the copy number of SMN2, which is a paralog of SMN1 in humans. SMN2 primarily produces less functional exon 7-deleted SMN protein and rarely generates a limited quantity of functional full-length SMN protein via alternative splicing. SMA patients are classified into types 0 through 4 based on the severity and timing of disease onset, predominantly determined by the number of copies of the SMN2 gene they possess. More than 50% of SMA patients are categorized as type 1, characterized by muscle defects with proximal muscle atrophy during infancy, eventually resulting in death within a few years (Mercuri et al., 2022).

Previous studies suggest that the onset of SMA is mainly attributed to SMN loss in motor neurons (Monani et al., 2000; Burghes et al., 2009). Nevertheless, motor neuron-specific SMN deficiency in the SMA mouse model exhibits relatively mild phenotypes compared to whole-body SMA mouse models (Park et al., 2010; McGovern et al., 2015). Furthermore, restoring SMN to motor neurons in SMA mouse models result in only partial rescue in lifespan and neuromuscular defects (Passini et al., 2010; Martinez et al., 2012; McGovern et al., 2015; Besse et al., 2020). Systemic administration of antisense oligonucleotide (ASO),

which corrects SMN2 splicing to restore SMN expression, significantly prolongs survival compared to CNS administration (Hua et al., 2011). In the mouse treated with a systemically delivered ASO, blocking the effect of ASO in the CNS by a complementary decoy did not have any detrimental effect on survival, motor function, or NMJ integrity (Hua et al., 2015). These studies suggest that peripheral SMN plays a crucial part in SMA pathology. Overall, investigating the impacts of SMN depletion in peripheral tissues is critical for alleviating neuromuscular impairments and increasing life expectancy in SMA.

In SMA patients, bone growth retardation has been observed (Khatri et al., 2008; Vai et al., 2015; Wasserman et al., 2017; Baranello et al., 2019; Hensel et al., 2020). Studies using whole-body SMA mouse models have revealed that this is caused by diminished growth plate chondrocyte proliferation and extracellular matrix (ECM) remodeling defects, independent of muscle atrophy (Hensel et al., 2020). However, it is still unclear whether these defects result from an internal factor such as SMN ablation in chondrocytes, or an external factor such as IGF1 decline in SMA patients and mice (Hua et al., 2011; Yesbek-Kaymaz et al., 2016).

Mesenchymal progenitor cells (MPCs) derived from the lateral plate mesoderm (LPM) differentiate into various types of limb mesenchymal cells, including bone, cartilage, and intramuscular mesenchymal cells like fibro-adipogenic progenitors (FAPs) (Nassari et al., 2017). Recent studies have revealed the role of FAPs in skeletal muscle homeostasis, as reducing the number of FAPs resulted in diminished muscle regeneration capacity, long-term muscle atrophy, and NMJ denervation (Wosczyna et al., 2019; Uezumi et al., 2021). Our latest research demonstrated that FAPs-specific deficiency of Bap1, one of the deubiquitinases, leads to NMJ defects (Kim et al., 2022). These recent findings raise the possibility that FAPs may have a specific role in the pathogenesis of neuromuscular diseases

such as SMA. However, it has not been studied if depletion of SMN in FAPs can lead to

SMA-like neuromuscular pathology.

In this study, we crossed a limb MPCs-specific Cre mouse with a floxed Smn exon7 mouse

carrying multiple copies of the human SMN2 gene, allowing us to examine the impact of

mesenchymal SMN reduction on SMA pathogenesis. Our findings indicate that the SMN

reduction in FAPs, similar to the extent of severe SMA, causes altered NMJ development like

SMA. Additionally, in chondrocytes, it leads to skeletal growth abnormalities due to growth

plate cell-autonomous defects.

**Results**

**Bone growth restriction due to growth plate cell-autonomous defects caused by MPCs-**

**specific SMN depletion**

To investigate the effects of SMN reduction within MPCs in SMA pathogenesis, we crossed

$Smn^{f7/f7}$ mice, which possess loxP sites flanking exon 7 of the Smn gene (Frugier et al., 2000),

with $Prrx1^{Cre}$ mice. This produced $Smn^{\Delta MPC}$ mice ($Prrx1^{Cre}; Smn^{f7/f7}$) that lacked the Smn

gene specifically in $Prrx1^{Cre}$-expressed limb MPCs that give rise to bone, cartilage, and FAPs

(Logan et al., 2002; Leinroth et al., 2022). To ascertain whether mutant mice carrying the

SMN2 gene, like SMA patients, present pathological phenotypes, we additionally generated

SMN2 2-copy $Smn^{\Delta MPC}$ ($Prrx1^{Cre}; Smn^{f7/f7}; SMN2^{+/+}$) and SMN2 1-copy $Smn^{\Delta MPC}$ ($Prrx1^{Cre};$

$Smn^{f7/f7}; SMN2^{+/0}$) mice. Control littermates that lacked $Prrx1^{Cre}$ were used as controls for

comparison.

SMN2 0-copy $Smn^{\Delta MPC}$ mice died within 24 hours after birth. Regions where limbs should

have formed at E18.5 only had rudimentary limb structures (Figure 1—figure supplement 1A). Furthermore, since the upper head bone did not envelop the brain, it was directly attached to the skin and protruded. These observations can be attributed to Prrx1[Cre]-mediated SMN deletion in the LPM-derived limb MPCs and the craniofacial mesenchyme, which is accountable for the formation of calvarial bone (Wilk et al., 2017). To investigate whether the lack of SMN proteins in MPCs is responsible for bone development abnormalities, we performed alcian blue and alizarin red staining on E18.5 SMN2 0-copy Smn[ΔMPC] mutants to analyze the structure of bones and cartilage. The appendages displayed restricted bone and cartilage formations, with scarcely discernible femur and tibia (Figure 1—figure supplement 1B-C). In the cranial region, there was an absence of both cartilage and bone at the location of the calvarial bone, with the parietal bone entirely missing and partially absent frontal bone (Figure 1—figure supplement 1D). Additionally, the sternum, which is one of the bones originating from the LPM (Sheng et al., 2015), was shorter than the control (Figure 1—figure supplement 1E).

The SMN2 2-copy Smn[ΔMPC] mice, carrying two homologous SMN2 genes, did not show any discernible differences from the Prrx1[Cre]-negative control littermates into adulthood. However, the SMN2 1-copy Smn[ΔMPC] mice exhibited reduced body size and shorter limb length compared to the SMN2 1-copy control (Smn[f7/f7]; SMN2[+/0]). To assess postnatal bone growth defects observed in SMA patients and mouse models, we conducted micro-CT analysis on femurs obtained from postnatal day 14 (P14) SMN2 1-copy Smn[ΔMPC] and SMN2 2-copy Smn[ΔMPC] mice (Figure 1A). The data showed that SMN2 1-copy mutants exhibited reduced femur diaphysis length, diameter, and trabecular bone volume compared to the control group, indicating bone formation defects (Figure 1B-D). We then examined femoral

bone thickness and diaphyseal bone mineral density (BMD) to determine whether mineralization was normal after bone formation. The thickness of the bone in the mutants did not differ significantly from the control, suggesting that bone mineralization was intact (Figure 1—figure supplement 1E-F). Unexpectedly, BMD slightly increased in SMN2 1-copy mutants compared to the control group (Figure 1—figure supplement 1G). The higher BMD may be attributed to greater mechanical stress caused by the shorter femur supporting the weight of the body, consistent with prior research indicating that elevated mechanical force leads to higher BMD in the femur (Hoxha et al., 2014; Ike et al., 2015). Nevertheless, the decrease in bone growth without apparent deterioration in bone mineralization of the femur of SMN2 1-copy Smn$^{\Delta MPC}$ mutants is consistent with findings from the whole-body SMA mouse model (Hensel et al., 2020). Collectively, these results suggest that mice carrying low copies of SMN2, with the SMN gene specifically deleted in MPCs, exhibit bone growth defects.

To determine whether bone growth defects in SMN2 1-copy Smn$^{\Delta MPC}$ mutants arise from disrupted chondrocyte homeostasis at growth plates, we stained the femur growth plate of P14 mice with hematoxylin and eosin (Figure 1—figure supplement 2A). In line with earlier findings in whole-body SMA mice (Hensel et al., 2020), SMN2 1-copy Smn$^{\Delta MPC}$ mice exhibited shorter proliferative and hypertrophic zones compared to control mice (Figure 1—figure supplement 2B-D). Additionally, there was a significant reduction in the number of chondrocytes in the hypertrophic zone (Figure 1—figure supplement 1J). Our data indicate that adequate expression of SMN is essential for the homeostasis of chondrocytes at growth plates.

**Mesenchymal cell-specific SMN reduction similar to severe SMA mouse model in SMN2 1-copy Smn$^{\Delta MPC}$ mutants**

The proliferation and differentiation of growth plate chondrocytes are regulated by extrinsic factors such as GH-IGF-I, glucocorticoids, and estrogen (Shim et al., 2015; Karimian et al., 2012; Shantanam et al., 2018). Previous research indicates that reduced expression of Igf-I/Igfbp3/Igfals in the liver leads to impaired chondrocyte proliferation and shorter femur length, emphasizing the importance of these factors in postnatal bone development (Yakar et al., 2002; Yakar et al., 2009). As the IGF pathway proteins are downregulated in whole-body SMA mice, the bone growth defects observed in the mice have sparked debate as it remains unclear whether they are due to intrinsic factors, such as chondrocyte SMN reduction, or the low IGF axis level (Hua et al., 2011; Tsai et al., 2014; Deguise et al., 2021; Hensel et al., 2015).

To clarify this issue, we used Smn$^{\Delta MPC}$ mutant mice, which enabled us to investigate the effect of SMN depletion in chondrocytes on bone growth, while circumventing the impact of external factors. To confirm the specific deletion of Smn in limb mesenchymal cells, including chondrocytes and FAPs, we performed qRT-PCR for full-length Smn mRNA expression in each tissue. Our findings indicate that full-length Smn mRNA expression in the brain, liver, skeletal muscle, and spinal cord of SMN2 1-copy Smn$^{\Delta MPC}$ mice at postnatal day 21 (P21) was similar to that of control mice, while it was significantly reduced in isolated FAPs and chondrocytes (Figure 1I). Additionally, we confirmed the presence of the Smn$\Delta 7$ variant, which is the exon 7-deleted form of Smn created by cre-lox mediated recombination, in both FAPs and chondrocytes by conducting genomic PCR on the SMN2 1-copy Smn$^{\Delta MPC}$ mutant (Figure 1H). Since SMN was not down-regulated in the tissues other than limb

mesenchymal cells, non-mesenchymal cells were ruled out from being responsible for the

phenotype observed in Smn$^{\Delta MPC}$ mutants. Indeed, the liver's mRNA expression of IGF

pathway genes, reported to be decreased in SMA mouse models and patients (Hua et al., 2011;

Deguise., 2021; Sahashi et al., 2013), showed no difference when comparing controls to

SMN2 1-copy Smn$^{\Delta MPC}$ mutants (Figure 1—figure supplement 2E). Based on these findings,

we concluded that the bone growth defects observed in the whole-body SMA mouse model

are mainly due to a depletion of SMN in chondrocytes.

Among our Smn$^{\Delta MPC}$ models, mice with two copies of SMN2 exhibit similar bone

development parameters as control mice without SMN deletion. However, mice with one

copy of SMN2 display bone pathological defects akin to SMA mouse models. This may be

because the quantity of human full-length SMN protein produced by SMN2 2-copy, was

sufficient to sustain SMN complex function in the SMN2 2-copy Smn$^{\Delta MPC}$ mutants, despite

the absence of functional mouse SMN protein in limb mesenchymal cells. On the contrary,

due to the lower expression of full-length SMN compared to SMN2 2-copy mutants, the

SMN complex may not function properly in SMN2 1-copy mutant cells. We confirmed this

by comparing the amount of full-length SMN protein in isolated FAPs from the hindlimbs of

control, Smn$^{\Delta MPC}$ mutants, and a severe SMA mouse model (*Smn$^{-/-}$; SMN2$^{+/+}$; SMNΔ7$^{+/+}$*; -

SMAΔ7 mutants -) (Figure 1I). The data show that SMN2 1-copy Smn$^{\Delta MPC}$ mutants exhibited

~80% reduction in SMN levels compared to the control group. The level of SMN protein in

SMN2 1-copy Smn$^{\Delta MPC}$ mutants was similar to that in SMAΔ7 mutants. Conversely, SMN2

2-copy mutants display a decrease of approximately 40% in SMN protein levels compared to

the control (Figure 1J). The moderately reduced expression of SMN is adequate to support

regular bone development in SMN2 2-copy mutants. Taken together, our findings indicate

that the reduction of mesenchymal SMN to levels comparable to that of the severe SMA

mouse model causes SMA-like bone pathology in the SMN2 1-copy mutant.

**Abnormal NMJ maturation in SMN2 1-copy Smn$^{\Delta MPC}$ mutants**

To investigate whether disabling SMN in FAPs results in SMA-like neuromuscular

impairments, we assessed if NMJ phenotypes observed in SMA mouse models also occur in

SMN2 1-copy Smn$^{\Delta MPC}$ mutants. Both severe and mild SMA mouse models exhibit impaired

NMJ maturation markers, including plaque-like morphology of acetylcholine receptor (AChR)

clusters, neurofilament (NF) varicosities, and poor terminal arborization (Kong et al., 2009;

Martinez, 2012; Monani et al., 2003; Kariya et al., 2008). To evaluate the impact of SMN

deficiency in FAPs on NMJ maturation, we evaluated the NMJ maturation markers in the TA

muscles of control and SMN2 1-copy Smn$^{\Delta MPC}$ mutant mice at P21, a time when NMJ

maturation is in progress (Figure 2A). Our examination revealed the presence of NF

varicosities in SMN2 1-copy mutants as compared with control mice (Figure 2A and 2D).

Additionally, the number of nerve branches was decreased and half of the total NMJs were

poorly arborized in SMN2 1-copy mutants (Figure 2B-C). These presynaptic alterations are

specific phenotypes in neurogenic atrophy like SMA. Unlike neurogenic atrophy, physiologic

atrophy shows no differences in presynaptic morphology, such as nerve branching

(Deschenes et al., 2006). This suggests that the NMJ phenotypes observed in SMN2 1-copy

Smn$^{\Delta MPC}$ mutant mice are not caused by decreased muscle size and activity resulting from

bone growth abnormalities. The morphology of AChR clusters shows that the mutants have

more immature plaque-like NMJs than the controls' pretzel-like structure (Figure 2E).

Therefore, these findings indicate that SMN2 1-copy Smn$^{\Delta MPC}$ mutants exhibit NMJ

maturation abnormalities common in SMA mouse models.

**Undisturbed NMJ formation in neonatal SMN2 1-copy Smn$^{\Delta MPC}$ mutants**

To determine whether any NMJ defects were present prior to juvenile NMJ maturation in

SMN2 1-copy Smn$^{\Delta MPC}$, we examined NMJ formation in SMN2 1-copy Smn$^{\Delta MPC}$ mice at the

neonatal stage on postnatal day 3. We evaluated the AChR and nerve terminal areas to assess

postsynaptic and presynaptic development, respectively (Figure 2F). Measurements of AChR

cluster size indicated no differences between control and SMN2 1-copy Smn$^{\Delta MPC}$ mice

(Figure 2G). However, the area of AChR covered by nerve terminals was slightly larger in

SMN2 1-copy Smn$^{\Delta MPC}$ (Figure 2H). We have no reasonable explanation for why the

coverage is higher in the mutant. However, there does not appear to be abnormal

development of the NMJ in the mutant, at least until the neonatal period. Therefore, we

reasoned that SMN2 1-copy Smn$^{\Delta MPC}$ mutants began to exhibit deterioration in the NMJ

maturation during the juvenile stage, following the intact neonatal development of NMJ.

**Aberrant NMJ morphology in the adult SMN2 1-copy Smn$^{\Delta MPC}$ mice**

To evaluate the organization of NMJ after the conclusion of postnatal NMJ development,

considering mesenchymal SMN expression, we examined NMJ morphology in the TA muscle

of control, SMN2 1-copy, and 2-copy Smn$^{\Delta MPC}$ mice at postnatal day 56 (P56). Our analysis

revealed that presynapses were fragmented in SMN2 1-copy mutants, resulting in a bouton-

like morphology, in contrast to the control and SMN2 2-copy Smn$^{\Delta MPC}$ mice (Figure 3A). In

SMN2 1-copy Smn$^{\Delta MPC}$ mice, a two-fold presynaptic fragmentation compared to the control

was quantified, demonstrating nerve terminal shrinkage (Figure 3B). Additionally, NF ends

displayed more severe varicosities than at P21 and were only connected to the proximal nerve

by very thin NF, unlike control and SMN2 2-copy mice (Figure 3C). Remarkably, numerous

presynaptic islands formed in SMN2 1-copy Smn$^{\Delta MPC}$ mice through the merging of

fragmented presynapses and NF varicosity. In SMN2 1-copy mutants, AChR clusters

displayed fragmented grape-shaped morphology that overlapped with nerve terminals,

whereas control and SMN2 2-copy mice displayed pretzel-like structures (Figure 3D). These

results suggest that defects in adult NMJ morphology occur when mesenchymal SMN protein

is reduced to the extent of the SMN2 1-copy Smn$^{\Delta MPC}$ mutants.

**Presynaptic neurotransmission alteration in SMN2 1-copy Smn$^{\Delta MPC}$ mutants**

To investigate whether the morphologically aberrant NMJs of SMN2 1-copy Smn$^{\Delta MPC}$ mice

have functional impairments, we isolated hindlimb extensor digitorum longus (EDL) muscles

from P56 mice and conducted electrophysiological recording. We incubated the muscles with

μ-conotoxin, which selectively inhibits muscle voltage-gated Na+ channels, prevented the

induction of muscle action potential (Ling et al., 2010; Zanetti 2018), and recorded the

Miniature endplate potential (mEPP) and evoked endplate potential (eEPP). (Ling et al., 2010;

Zanetti 2018). mEPP, a response that occurs when spontaneously released acetylcholine binds

to nicotinic AChR without nerve stimulation, was measured ex vivo near the NMJs of the

EDL muscles in the control and SMN2 1-copy Smn$^{\Delta MPC}$ mice (Figure 4A). The mEPP

amplitude was increased in SMN2 1-copy Smn$^{\Delta MPC}$ mice (Figure 4B), whereas mEPP

frequency was comparable between the controls and mutants (Figure 4C). The results

indicate that the NMJ synapses of SMN2 1-copy Smn$^{\Delta MPC}$ mice are functional and more

sensitive to acetylcholine compared to the controls. Next, we measured the eEPP by

stimulating an action potential at the peroneal nerve (Figure 4D). Despite the increased mEPP

amplitude, the amplitude of eEPPs was significantly decreased in SMN2 1-copy Smn$^{\Delta MPC}$

mice (Figure 4E). These results suggest that the nerve terminals in SMN2 1-copy Smn$^{\Delta MPC}$

mice exhibit decreased quantal content. This could be due to a decrease in vesicle release

probability or a reduced readily releasable pool (RRP). Notably, there was no difference in

the paired-pulse response, indicating normal neurotransmitter release probability (Figure 4F).

Taken together, these findings suggest that the presynaptic neurotransmission ability of the

NMJ is reduced in SMN2 1-copy Smn$^{\Delta MPC}$ mutants.

**Disturbed nerve terminal structure in SMN2 1-copy Smn$^{\Delta MPC}$ mice**

To examine the NMJ ultrastructure of SMN2 1-copy Smn$^{\Delta MPC}$ mutants, we utilized

transmission electron microscopy (TEM) (Figure 5A). The density of junctional folds in

SMN2 1-copy Smn$^{\Delta MPC}$ mutant specimens was comparable to that of the control (Figure 5B).

However, the density of synaptic vesicles was substantially elevated in the SMN2 1-copy

mutants (Figure 5C). Since previous electrophysiological results suggested a decrease in

presynaptic neurotransmission capacity in SMN2 1-copy mutants, this could be due to

synaptic vesicles failing to fuse with the membrane, leading to the accumulation of vesicles

in the terminal. Additionally, the detachment of the nerve terminal is more frequent at the

NMJ of mutants (Figure 5D). The detachment of nerve terminals observed in SMN2 1-copy

Smn$^{\Delta MPC}$ mutants could have resulted in diminished presynaptic neurotransmission capacity.

Collectively, these findings indicate that SMN2 1-copy Smn$^{\Delta MPC}$ mutants have nerve

terminal-specific pathological defects at the NMJ ultrastructural level.

**FAPs transplantation rescues NMJ morphology in limb mesenchymal SMN mutants**

SMN-deleted limb mesenchymal tissues in SMN2 1-copy $Smn^{\Delta MPC}$ mutants comprise not

only FAPs, but also bone, cartilage, pericytes, and tendon, among others (Leinroth et al.,

2022; Nassari et al., 2017). To evaluate the critical role of FAPs in the postnatal development

of the NMJ, we isolated fluorescent protein-labeled wild-type FAPs from $Prrx1^{Cre}$;

$Rosa26^{YFP/+}$ or $Prrx1^{Cre}$; $Rosa26^{tdTomato/+}$ mice and transplanted them into the TA muscles of

SMN2 1-copy $Smn^{\Delta MPC}$ mice at postnatal day 10. The TA muscle on the contralateral side

was treated with a vehicle as a control. In SMN2 1-copy $Smn^{\Delta MPC}$ mice at P56, the muscle

that received FAPs showed decreased presynaptic fragmentation, NF varicosities, and

postsynaptic fragmentation compared to the contralateral muscle (Figure 6A-D). These data

demonstrate that the transplantation of wild-type FAPs rescues the abnormal NMJ

development in SMN2 1-copy $Smn^{\Delta MPC}$ mice. Overall, our findings indicate that SMN

depletion in FAPs leads to the neuronal SMN-independent NMJ pathology in severe SMA,

which is rescued by the transplantation of healthy FAPs.

**Discussion**

303    In this paper, we elucidate the contribution of SMN depletion in mesenchymal progenitors for

304    the pathogenesis of SMA. To test this hypothesis, we generated conditional knockout mouse

305    strains to delete the Smn allele specifically in limb mesenchymal cells and carry human

306    SMN2 copies. Our research using these mouse models resulted in three major discoveries.

307First, SMN deficiency in FAPs contributes to NMJ pathological defects in SMA. We

308observed delayed NMJ maturation and varicosities in juvenile SMN2 1-copy Smn$^{\Delta MPC}$

309mutant. The pathogenic NMJ phenotypes were also observed in the SMAΔ7 mutant, which is

310one of the severe SMA mouse models (Kong et al., 2009; Martinez, 2012; Kariya et al., 2008).

311As the SMAΔ7 mutant typically lives for approximately 12 days, the fragmentation of the

312NMJ in adult SMN2 1-copy mutant was not evaluated in the severe SMA mutant.

313Nevertheless, models that induce a conditional adult SMN deficiency through either Cre$^{ER}$

314allele or oligonucleotide administration resulting in SMN reduction, demonstrated the

315depletion of SMN throughout the body caused fragmentation of NMJ (Sahashi et al., 2013;

316Kariya et al., 2014). Thus, we demonstrate that the SMN2 1-copy Smn$^{\Delta MPC}$ mutant model

317mimics whole-body SMA mouse models in NMJ morphology. In the electrophysiological test,

318the SMAΔ7 mutant exhibited reduced quantal content, readily releasable pool (RRP), and

319vesicle release probability (Torres-Benito et al., 2011). In our study, we did not assess RRP in

320SMN2 1-copy Smn$^{\Delta MPC}$ mutant, but instead reduced quantal content and normal vesicle

321release probability, indicating a reduced RRP size. The decrease in RRP size in the SMAΔ7

322mutant was attributed to abnormal axonal transport, which led to a decline in synaptic vesicle

323density in the nerve terminal (Dale et al., 2011; Kong et al., 2009). However, we found that

324synaptic vesicle density was increased in the SMN2 1-copy Smn$^{\Delta MPC}$ mutant. It is possible

325that the alteration of active zones, which were observed in the SMA motor terminals,

326contributed to the reduction in synaptic vesicle fusion and the higher synaptic vesicle density.

327Indeed, nerve terminal detachment in SMN2 1-copy mutant mice was also found in active

328zone complex protein integrin-α3 knockout mice (Ross et al., 2017). Based on the rescue data

329of transplanted healthy FAPs, we can report that FAPs-specific SMN depletion is involved in

330NMJ pathology of SMA.

Second, we demonstrated that skeletal growth defects, a phenotype observed in SMA (Khatri

et al., 2008; Vai et al., 2015; Wasserman et al., 2017; Baranello et al., 2019; Hensel et al.,

2020), are a cell-autonomous pathological effect of the depletion of SMN in chondrocytes.

We observed reduced bone size and volume in juvenile SMN2 1-copy $Smn^{\Delta MPC}$ mutant.

Depletion of SMN from chondrocytes is sufficient to cause growth problems without altering

the hepatic IGF axis, which affects growth plate homeostasis. Recent studies have reported

that IGF1 overexpression improves biochemical and behavioral manifestations in SMA mice,

suggesting potential therapies for SMA (Bosch-Marce et al., 2011; Tsai et al., 2014). Our

study demonstrated that these IGF1 therapies for SMA without mesenchymal SMN

supplementation have a limited ability to rescue bone abnormalities. In addition, the skeletal

growth defect may affect physiologic muscle atrophy through imbalanced muscle contraction

and reduced tension. This physiologic atrophy may be part of SMA-associated muscle

weakness independent of neurogenic atrophy.

Third, it was demonstrated that adequate levels of SMN protein are essential for MPCs to

contribute to limb neuromusculoskeletal development. The mutants with only 1 copy of

SMN2 exhibit problematic symptoms observed in both SMA patients and mouse models,

while SMN2 2-copy mutants display a typical phenotype in bone and NMJ. The lack of SMN

protein in MPCs by insufficient SMN2 copies, similar to the deficiency seen in severe SMA,

is responsible for the onset of SMA pathology. Thus, we propose that restoration of deficient

SMN in MPCs is crucial for rehabilitating their function. Based on these discoveries, SMN

replenishment treatments for MPCs, specifically FAPs, and chondrocytes, are necessary to

provide a complete solution for neuromusculoskeletal defects in severe SMA patients.

The initial focus for the treatment of SMA was on motor neurons located in the spinal cord,

and pharmaceuticals were created to address the deficiency of SMN in these neurons

(Mercuri et al., 2020). For example, intrathecal injections of Spinraza can efficiently boost

SMN in the CNS, including the spinal cord (Passini et al., 2011; Claborn et al., 2018).

However, Spinraza does not address the lack of SMN in peripheral tissues, including

mesenchymal cells, highlighted in recent studies. The drug Zolgensma, which employs AAV9

to express SMN through systemic delivery, appears to resolve this issue (Foust et al., 2009;

Foust et al., 2022; Valori et al., 2010; Mattar et al., 2013). Nevertheless, a prior study

indicates that chondrocytes within the growth plate and articular cartilage do not get infected

with AAV9 (Yang et al., 2019). Furthermore, it is well-established that DNA vectors

delivered via AAV9 undergo dilution during cell proliferation (Penaud-Budloo et al., 2008;

Colella et al., 2018; Van-Alstyne et al., 2021; Heller et al., 2021). Given the infection issue

and the dilution issue by the active cell population changes of chondrocytes and FAPs during

postnatal development (Petrany., 2020; Bachman et al., 2022), SMN supplementation through

Zolgensma alone would not be sufficient for severe SMA patients. Thus, there is an urgent

need to research and develop therapeutic strategies that target mesenchymal progenitors.

While this study is the first to demonstrate the impact of SMN depletion in FAPs on the NMJ

development, it does not elucidate the specific mechanism by which FAPs influence the NMJ.

Our study observed NMJ recovery in the muscles transplanted with healthy FAPs, but not in

the contralateral muscles, indicating that FAPs are likely involved in NMJ development

through juxtacrine or paracrine signaling. Since FAPs interact with surrounding tissues

through a variety of signaling factors, such as extracellular matrix (Contreras et al., 2021;

Scott et al 2019), Wnt-related protein (Lukjanenko et al., 2019), and Bmp signaling protein

(Uezumi et al., 2021; Camps et al., 2020), mis-splicing of the signaling factors due to SMN

reduction could disturb the homeostasis of neighboring tissues (Zhang et al., 2008).

Furthermore, a population of FAPs associated with NMJ was discovered through single-cell

RNA sequencing in a recent study (Leinroth et al., 2022). This population is located adjacent

to the NMJ and responds to denervation, indicating an increased possibility of interaction

with the NMJ organization. Therefore, it is necessary to conduct additional investigations into

the expression of various signaling factors by diverse FAP populations in future studies.

**Methods**

**Animals**

$Prrx1^{Cre}$ (stock 005584), $Smn^{f7/+}$ (stock 006138), $Rosa26^{YFP/+}$ (stock 006148), $Rosa26^{tdTomato/+}$ (stock 007914), $SMN2^{+/+}$, $Smn^{+/-}$ (stock 005024) and $Smn^{+/-}$; $SMN2^{+/+}$; $SMN\Delta7^{+/+}$ (stock 005025) mice were acquired from The Jackson Laboratory (Bar Harbor, ME, USA). SMN2 0-copy $Smn^{\Delta MPC}$ mice ($Prrx1^{Cre}$; $Smn^{f7/f7}$), SMN2 1-copy $Smn^{\Delta MPC}$ mice ($Prrx1^{Cre}$; $Smn^{f7/f7}$; $SMN2^{+/0}$), and SMN2 2-copy $Smn^{\Delta MPC}$ mice ($Prrx1^{Cre}$; $Smn^{f7/f7}$; $SMN2^{+/+}$) were generated by crossing $Prrx1^{Cre}$ mice with $Smn^{f7/+}$ and $Smn^{+/-}$; $SMN2^{+/+}$ mice. To utilize fibro-adipogenic progenitors (FAPs) transplantation, we generated $Prrx1^{Cre}$; $Rosa26^{YFP/+}$ mice and $Prrx1^{Cre}$; $Rosa26^{tdTomato/+}$ mice by breeding $Prrx1^{Cre}$ with $Rosa26^{YFP/+}$ and $Rosa26^{tdTomato/+}$ mice, respectively. To avoid deletion of the floxed allele by $Prrx1^{Cre}$ expression in female germline, we bred females without $Prrx1^{Cre}$ line to $Prrx1^{Cre}$ transgenic males. SMAΔ7 mutants ($Smn^{-/-}$; $SMN2^{+/+}$; $SMN\Delta7^{+/+}$) were produced by mating $Smn^{+/-}$; $SMN2^{+/+}$; $SMN\Delta7^{+/+}$ mice. Both male and female mice were used in the experiments, and no sex-specific differences were observed. Control littermates lacking $Prrx1^{Cre}$ were utilized for analysis. All mouse lines were housed under controlled conditions with specific pathogen-free and handled according to the guidelines of the Seoul National University Institutional Animal Care and Use Committee (IACUC).

**Micro CT**

The femurs from three groups - control, SMN2 2-copy $Smn^{\Delta MPC}$, and SMN2 1-copy $Smn^{\Delta MPC}$ mice at postnatal day 14 (P14) - were isolated and cleaned of muscles and skin. Subsequently,

the femurs were preserved in 4% paraformaldehyde (PFA) in PBS overnight at 4°C before

micro-computed tomography (micro CT). The femurs were then imaged through a Quantum

GX II micro-CT imaging system (PerkinElmer, Hopkinton, MA, USA). The X-ray source for

scanning was set at 90 kV and 88 mA with a field of view of 10 mm (voxel size, 20 μm;

scanning time, 14 min). The 3D imaging was viewed using the 3D Viewer software of the

Quantum GX II. The size and volume of the femur bone were measured via AccuCT analysis

software within the ossified diaphysis and metaphysis of the femur, excluding the epiphysis.

Bone mineral density (BMD) calibration was performed using a 4.5 mm BMD phantom and

BMD measurements were taken at the center of the diaphysis.

**Histology**

Alcian blue & alizarin red staining was performed on control and SMN2 0-copy Smn[ΔMPC]

mice at E18.5 using a previously reported protocol (Ovchinnikov, 2009). For hematoxylin &

eosin staining of the growth plate in the control and SMN2 1-copy Smn[ΔMPC] mice at P14, the

femurs were fixed in 4% paraformaldehyde for 24 hours, rinsed in running tap water for 24

hours, and incubated with 10% EDTA (pH 7.4) at 4°C with shaking for 2-3 days.

Subsequently, the samples were rinsed in running tap water for 24 hours and then dehydrated

through ethanol/xylene and embedded in paraffin. The embedded samples were then

sectioned to a thickness of 5 μm, rehydrated, and stained with Gill No. 3 formula

hematoxylin and eosin Y (H&E, Sigma Aldrich, St. Louis, MO, USA). Stained slides were

analyzed using 10× and 20x objectives in the EVOS M7000 imaging system (Thermo Fisher

Scientific, Waltham, MA, USA). The growth plate's proliferative and hypertrophic zones

were defined by their respective cell sizes.

**Hindlimb fibro-adipogenic progenitors isolation**

Isolation of limb muscle FAPs was performed according to a previously reported protocol (Kim et al., 2021) with modifications. Limb muscles were dissected and mechanically dissociated in Dulbecco's modified Eagle's medium (DMEM, Hyclone) containing 10% horse serum (Hyclone, Logan, UT, USA), collagenase II (800 units/mL; Worthington, Lakewood, NJ, USA), and dispase (1.1 units/mL; ThermoFisher Scientific, Waltham, MA, USA) at 37°C for 40 min. Digested suspensions were subsequently triturated by sterilized syringes with 20G 1/2 needle (BD Biosciences, Franklin Lakes, NJ, USA) and washed with DMEM to harvest mononuclear cells. Mononuclear cells were stained with corresponding antibodies. All antibodies used in FACS analysis were listed in Supplemental Table 1. To exclude dead cells, 7AAD (Sigma Aldrich; St. Louis, MO, USA) was used. Stained cells were analyzed and 7AAD$^-$Lin$^-$Vcam$^-$Sca1$^+$ (FAPs) were isolated using FACS Aria III cell sorter (BD Biosciences) with 4-way-purity precision. For western blot, freshly isolated FAPs were cultured at 37 °C in alpha-MEM (Hyclone) supplemented with Antibiotic-Antimycotic (anti-anti, Gibco) and 20% FBS (Hyclone). For transplantation, YFP$^+$ FAPs and tdTomato$^+$ FAPs were isolated from postnatal day 10~21 *Prrx1$^{Cre}$; Rosa26$^{YFP/+}$* mice and *Prrx1$^{Cre}$; Rosa26$^{tdTomato/+}$* mice, respectively. Isotype control density plots were used as a reference for positive gating.

**Chondrocytes isolation**

Isolation of chondrocytes followed a modified protocol previously published (Jonason et al.,

2015). Using blunt forceps, cartilage caps were removed from P21 mouse femoral heads and

dissected into ~1mm fragments in a petri dish with 10X anti-anti in PBS. The cartilage

fragments were washed twice with PBS and then incubated in 5 mL collagenase II solution

(800 units/mL collagenase II in DMEM with 1X anti-anti, sterilized by 0.2 µm filtration) in a

60-mm culture dish at 37 °C in a 5% CO2 incubator overnight. Chondrocytes were released

by pipetting the remaining cartilage fragments ten times and then filtering them through a 70

µm cell strainer to a 50 ml conical tube. The cells were then washed twice with PBS and

pelleted by centrifugation at 500g for 5 min. Subsequently, the cells were cultured overnight

at 37 °C in a 5% CO2 incubator, in complete culture medium (DMEM with 10% FBS and 1X

anti-anti) overnight at 37 °C in a 5% CO2 incubator.

**PCR reaction**

To detect the presence of Smn exon 7 floxed and deleted allele, 20 mg of each tissue were

dissolved in direct PCR buffer (VIAGEN) with proteinase K overnight at 65°C. After

inactivation at 95°C for 30 minutes, PCR was performed using previously reported primer

sets to verify the existence of Cre, Smn$^{f7}$, and Smn$^{\Delta 7}$ alleles (Frugier et al., 2000). Primers are

listed in Supplemental Table 1.

**RNA extraction and measurement of mRNA expression**

Total RNA was extracted from the brain, liver, spinal cord L4, tibialis anterior muscle, freshly

isolated FAPs, and chondrocytes using a TRIzol Reagent (Life Technologies, Carlsbad, CA,

USA) and analyzed by qRT-PCR. First-strand complementary DNA was synthesized from 1

µg of RNA using ReverTra Ace (Toyobo, Osaka, Japan) containing random oligomer

according to the manufacturer's instructions. qRT-PCR (Qiagen) was performed with SYBR

Green technology (SYBR Premix Ex Taq, Qiagen) using specific primers against indicated

genes. Relative mRNA levels were determined using the $2^{-\Delta\Delta Ct}$ method and normalized to

*Gapdh* (Figure 1H, Figure 1—figure supplement 1K). Primers are listed in Supplemental

Table 1.

## Western blot

Cultured FAPs at passage 3 were homogenized in RIPA buffer (50 mM Tris-HCl, pH 7.5, 0.5%

SDS, 20 µg/ mL aprotinin, 20 µg/ mL leupeptin, 10 µg/ mL phenylmethylsulfonyl fluoride, 1

mM sodium orthovanadate, 10 mM sodium pyrophosphate, 10 mM sodium fluoride, and 1

mM dithiothreitol). Cell lysates were centrifuged at 13,000 rpm for 15 min. Supernatants

were collected and subjected to immunoblot. BCA protein assay (Thermo Fisher Scientific)

was used for estimating total protein concentrations. Normalized total proteins were analyzed

by electrophoresis in 10% polyacrylamide gels and transferred to PVDF membranes

(Millipore, Billerica, MA, USA). Membranes were blocked in 5% skim milk (BD

Biosciences) in TBS with 0.1% Tween-20 and incubated with primary antibodies overnight at

4°C. After incubation with the corresponding HRP-conjugated secondary antibodies, the

membranes were developed using a Fusion solo chemiluminescence imaging system (Vilber,

Marne-la-Vallée, France). α-tubulin was used as a loading control. Band intensities were

quantified using ImageJ software. Antibodies used in this study were listed in Supplemental

Table 1. Primary and secondary antibodies were diluted 1:1000 and 1:10000 with PBS

containing 0.1% Tween-20 and 3% BSA, respectively.

## FAPs transplantation

498     YFP$^+$ or tdTomato$^+$ FAPs were isolated by FACS from the limb muscles of indicated mice.

499     $1\times10^5$ FAPs were suspended in 0.1% gelatin (Sigma Aldrich; St. Louis, MO, USA) in PBS

500     and then transplanted into one side of the TA muscles of SMN2 1-copy Smn$^{\Delta MPC}$ mice. The

501     contralateral muscle received an equivalent volume of 0.1% gelatin in PBS (Vehicle).

## Electrophysiology

The extensor digitorum longus (EDL) muscle was dissected from control and SMN2 1-copy

Smn$^{\Delta MPC}$ mice, along with the peroneal nerve, and then pinned to a Sylgard-coated recording

chamber. Intracellular recording was conducted in oxygenated Ringer's solution, which

comprised 138.8 mM NaCl, 4 mM KCl, 12 mM NaHCO3, 1 mM KH2PO4, 1 mM MgCl2

and 2 mM CaCl2 with a pH of 7.4. Action potential of the muscle was prevented by preparing

the muscle in 2.5 uM μ-conotoxin GIIB (Alomone, Jerusalem, Israel) for 10 minutes

beforehand. The recording was performed in toxin-free Ringer's solution. Miniature endplate

potentials (mEPPs) were recorded from a junction, followed by recordings of evoked

endplate potentials (eEPPs) by stimulating the attached peroneal nerve. The eEPPs were

elicited using evoked stimulation. Paired-pulse stimulation (10 ms inter-stimulus interval)

was utilized to assess synaptic transmission. The data was obtained and analyzed with

Axoclamp 900A and Clampfit version 10.7 software.

## Transmission electron microscopy

NMJ transmission electron microscopy followed a modified protocol previously reported (Modla et al., 2010). Mouse EDL muscle was swiftly excised and fixed in 4% paraformaldehyde (PFA) dissolved in Sorensen's phosphate buffer (0.1M, pH 7.2), followed by washing in 0.1 M phosphate buffer. The EDL was then gradually infiltrated on a rotator at room temperature with sucrose: 0.1M phosphate buffer solutions of 30% and 50%, for one hour each, followed by an overnight incubation in 70% sucrose. Excess sucrose was then eliminated using filter paper, and the muscle was embedded in an optimal cutting temperature compound (O.C.T.; Sakura Fineteck, Torrance, CA, USA), followed by being frozen in a cryostat (Leica, Wetzlar, Germany). 10-µm thick longitudinal sections were washed in PBS and treated with Alexa fluor 555-conjugated α-bungarotoxin (1:500, Invitrogen) for an hour. Imaging was conducted with the EVOS M7000 imaging system, and we selected 4-5 NMJ-rich regions for processing with TEM. The sections were fixed with 2% glutaraldehyde and 2% PFA in 0.1M cacodylate buffer (pH 7.2) for two hours at room temperature, with an additional overnight incubation at 4°C. After washing with 0.1M cacodylate buffer they were post-fixed with 1% osmium tetraoxide in 0.1M cacodylate buffer (pH 7.2)- for two hours at 4°C. The sections were then stained en bloc with 0.5% uranyl acetate overnight, washed with distilled water, and dehydrated using serial ethanol and propylene oxide. The sections were embedded in epoxy resin (Embed-812, Electron Microscopy Sciences) and detached from the slides by dipping them in liquid nitrogen. Ultra-thin sections (70nm) were prepared with a diamond knife on an ultramicrotome (ULTRACUT UC7, Leica) and mounted on 100 mesh copper grids. Sections were stained with 2% uranyl acetate for 10 minutes and lead citrate for 3 minutes, then observed using a transmission electron microscope (80kV, JEM1010, JEOL or 120kV, Talos L120C, FEI). Synaptic vesicle density was quantified within a distance of 500 nm from the presynaptic membrane.

**Immunohistochemistry**

Freshly dissected TA muscles were fixed in 4% PFA for 30 min at room temperature. Subsequently, the muscles were cryoprotected in 30% sucrose overnight, embedded in O.C.T., snap-frozen in liquid nitrogen, and stored at -80°C prior to sectioning. Longitudinal 40 μm-thick sections were obtained from the embedded muscles using a cryostat. The sections were blocked for 2 hours at room temperature using 5% goat serum and 5% bovine serum albumin in PBS/0.4% Triton X-100. Then, the sections were incubated with primary antibodies in the blocking buffer for two days at 4 °C. After washing the sections three times with PBS/0.4% Triton X-100, the sections were stained with secondary antibodies overnight at 4 °C, and then incubated with Alexa fluor 488 or 555-conjugated α-bungarotoxin (1:500, Invitrogen) for 2hr at RT, washed three times with PBS/0.4% Triton X-100 and mounted in VECTASHIELD. Z-serial images were collected at 40X with a Leica SP8 confocal laser scanning microscope. To analyze NMJ morphology, LasX software was used to obtain maximal projections. Neurofilament (NF) varicosity refers to the varicose NF end connected to the rest of the nerve terminal. To quantify NMJ size and synaptophysin coverage, the Btx area and synaptophysin area were measured by ImageJ analysis software.

**Statistical analysis**

All statistical analyses were performed using GraphPad Prism 9 (GraphPad Software). For comparison of significant differences in multiple groups for normally distributed data, statistical analysis was performed by one-way or two-way ANOVA followed by Tukey's

pairwise comparison post hoc test. For non-normally distributed data, Brown–Forsythe and

Welch ANOVA followed by Games-Howell multiple comparisons test was used. For the

comparison of two groups, Student's unpaired t-test assuming a two-tailed distribution with

Welch's correction was used. Unless otherwise noted, all error bars represent s.e.m. The

number of biological replicates and statistical analyses for each experiment were indicated in

the figure legends. Independent experiments were performed at least in triplicates. $P < 0.05$

was considered statistically significant at the 95% confidence level. *$p < 0.05$, **$p < 0.01$,

***$p < 0.001$, ****$p < 0.0001$.

**Supplemental information**

Supplemental information includes one Supplementary Table.

**Ethics**

The care and treatment of animals were approved by the IACUC of Seoul National University.

**Acknowledgments**

We express our gratitude to the Kong and Choi laboratory members for their valuable

feedback during the project. This work was supported by the National Research Foundation

of Korea (NRF-2022R1A2C3007621 and NRF-2020R1A5A1018081).

**References**

Bachman, J. F., & Chakkalakal, J. V. (2022). Insights into muscle stem cell dynamics during
postnatal development. *FEBS Journal*, *289*(10), 2710–2722.
https://doi.org/10.1111/febs.15856

Baranello, G., Vai, S., Broggi, F., Masson, R., Arnoldi, M. T., Zanin, R., Mastella, C., &
Bianchi, M. L. (2019). Evolution of bone mineral density, bone metabolism and fragility
fractures in Spinal Muscular Atrophy (SMA) types 2 and 3. *Neuromuscular Disorders*,
*29*(7), 525–532. https://doi.org/10.1016/j.nmd.2019.06.001

Besse, A., Astord, S., Marais, T., Roda, M., Giroux, B., Lejeune, F. X., Relaix, F., Smeriglio,
P., Barkats, M., & Biferi, M. G. (2020). AAV9-Mediated Expression of SMN Restricted
to Neurons Does Not Rescue the Spinal Muscular Atrophy Phenotype in Mice.
*Molecular Therapy*, *28*(8), 1887–1901. https://doi.org/10.1016/j.ymthe.2020.05.011

Bosch-Marcé, M., Wee, C. D., Martinez, T. L., Lipkes, C. E., Choe, D. W., Kong, L., Van
Meerbeke, J. P., Musarò, A., & Sumner, C. J. (2011). Increased IGF-1 in muscle
modulates the phenotype of severe SMA mice. Human Molecular Genetics, 20(9),
1844–1853. https://doi.org/10.1093/hmg/ddr067

Camps, J., Breuls, N., Sifrim, A., Giarratana, N., Corvelyn, M., Danti, L., Grosemans, H.,
Vanuytven, S., Thiry, I., Belicchi, M., Meregalli, M., Platko, K., MacDonald, M. E.,
Austin, R. C., Gijsbers, R., Cossu, G., Torrente, Y., Voet, T., & Sampaolesi, M. (2020).
Interstitial Cell Remodeling Promotes Aberrant Adipogenesis in Dystrophic Muscles.
*Cell Reports*, *31*(5), 107597. https://doi.org/10.1016/j.celrep.2020.107597

Cifuentes-Diaz, C., Frugier, T., Tiziano, F. D., Lacène, E., Roblot, N., Joshi, V., Moreau, M.
H., & Melki, J. (2001). Deletion of murine SMN exon 7 directed to skeletal muscle leads
to severe muscular dystrophy. Journal of Cell Biology, 152(5), 1107–1114.
https://doi.org/10.1083/jcb.152.5.1107

Claborn, M. K., Stevens, D. L., Walker, C. K., & Gildon, B. L. (2019). Nusinersen: A
Treatment for Spinal Muscular Atrophy. *Annals of Pharmacotherapy*, *53*(1), 61–69.
https://doi.org/10.1177/1060028018789956

Colella, P., Ronzitti, G., & Mingozzi, F. (2018). Emerging Issues in AAV-Mediated In Vivo
Gene Therapy. *Molecular Therapy - Methods and Clinical Development*, *8*(March), 87–
104. https://doi.org/10.1016/j.omtm.2017.11.007

Contreras, O., Rossi, F. M. V., & Theret, M. (2021). Origins, potency, and heterogeneity of
skeletal muscle fibro-adipogenic progenitors—time for new definitions. *Skeletal Muscle*,
*11*(1), 1–25. https://doi.org/10.1186/s13395-021-00265-6

Currie, J. D., Grosser, L., Murawala, P., Schuez, M., Michel, M., Tanaka, E. M., & Sandoval-
Guzman, T. (2019). The Prrx1 limb enhancer marks an adult subpopulation of injury-

responsive dermal fibroblasts. *Biology Open*, *8*(7), 1–25.
https://doi.org/10.1242/bio.043711

Dale, J. M., Shen, H., Barry, D. M., Garcia, V. B., Rose, F. F., Lorson, C. L., & Garcia, M. L.
(2011). The spinal muscular atrophy mouse model, SMAΔ7, displays altered axonal
transport without global neurofilament alterations. Acta Neuropathologica, 122(3), 331–
341. https://doi.org/10.1007/s00401-011-0848-5

Deguise, M., Pileggi, C., Repentigny, Y. De, Beauvais, A., Tierney, A., Chehade, L.,
Michaud, J., Llavero-hurtado, M., Lamont, D., Atrih, A., Wishart, T. M., Gillingwater, T.
H., Schneider, B. L., Harper, M., Parson, S. H., & Kothary, R. (2021). SMN Depleted
Mice Offer a Robust and Rapid Onset Model of. *Cellular and Molecular*
*Gastroenterology and Hepatology*, *12*(1), 354-377.e3.
https://doi.org/10.1016/j.jcmgh.2021.01.019

Deschenes, M. R., Tenny, K. A., & Wilson, M. H. (2006). Increased and decreased activity
elicits specific morphological adaptations of the neuromuscular junction. *Neuroscience*,
*137*(4), 1277–1283. https://doi.org/10.1016/j.neuroscience.2005.10.042

Foust, K. D., Nurre, E., Montgomery, C. L., Hernandez, A., Chan, C. M., & Kaspar, B. K.
(2009). Intravascular AAV9 preferentially targets neonatal neurons and adult astrocytes.
*Nature Biotechnology*, *27*(1), 59–65. https://doi.org/10.1038/nbt.1515

Foust, K. D., Wang, X., McGovern, V. L., Braun, L., Bevan, A. K., Haidet, A. M., Le, T. T.,
Morales, P. R., Rich, M. M., Burghes, A. H. M., & Kaspar, B. K. (2022). Retraction
Note: Rescue of the spinal muscular atrophy phenotype in a mouse model by early
postnatal delivery of SMN. *Nature Biotechnology*, *40*(11), 1692–1692.
https://doi.org/10.1038/s41587-022-01497-7

Frugier, T., Tiziano, F. D., Cifuentes-Diaz, C., Miniou, P., Roblot, N., Dierich, A., Le Meur,
M., & Melki, J. (2000). Nuclear targeting defect of SMN lacking the C-terminus in a
mouse model of spinal muscular atrophy. *Human Molecular Genetics*, *9*(5), 849–858.
https://doi.org/10.1093/hmg/9.5.849

Heller, G. J., Marshall, M. S., Issa, Y., Marshall, J. N., Nguyen, D., Rue, E., Pathmasiri, K. C.,
Domowicz, M. S., van Breemen, R. B., Tai, L. M., Cologna, S. M., Crocker, S. J.,
Givogri, M. I., Sands, M. S., & Bongarzone, E. R. (2021). Waning efficacy in a long-
term AAV-mediated gene therapy study in the murine model of Krabbe disease.
*Molecular Therapy*, *29*(5), 1883–1902. https://doi.org/10.1016/j.ymthe.2021.01.026

Hensel, N., Brickwedde, H., Tsaknakis, K., Grages, A., Braunschweig, L., Lüders, K. A.,
Lorenz, H. M., Lippross, S., Walter, L. M., Tavassol, F., Lienenklaus, S., Neunaber, C.,
Claus, P., & Hell, A. K. (2020). Altered bone development with impaired cartilage
formation precedes neuromuscular symptoms in spinal muscular atrophy. *Human*
*Molecular Genetics*, *29*(16), 2662–2673. https://doi.org/10.1093/hmg/ddaa145

Hoxha, R., Islami, H., QorrajBytyqi, H., Thaci, S., & Bahtiri, E. (2014). Relationship of Weight and Body Mass Index with Bone Mineral Density in Adult Men from Kosovo. *Materia Socio Medica*, *26*(5), 306. https://doi.org/10.5455/msm.2014.26.306-308

Hua, Y., Sahashi, K., Rigo, F., Hung, G., Horev, G., Bennett, C. F., & Krainer, A. R. (2011). Peripheral SMN restoration is essential for long-term rescue of a severe spinal muscular atrophy mouse model. *Nature*, *478*(7367), 123–126. https://doi.org/10.1038/nature10485

Passini, M. A., Bu, J., Roskelley, E. M., Richards, A. M., Sardi, S. P., O'Riordan, C. R., Klinger, K. W., Shihabuddin, L. S., & Cheng, S. H. (2010). CNS-targeted gene therapy improves survival and motor function in a mouse model of spinal muscular atrophy. Journal of Clinical Investigation, 120(4), 1253–1264. https://doi.org/10.1172/JCI41615

Torres-Benito, L., Neher, M. F., Cano, R., Ruiz, R., & Tabares, L. (2011). SMN requirement for synaptic vesicle, active zone and microtubule postnatal organization in motor nerve terminals. PLoS ONE, 6(10). https://doi.org/10.1371/journal.pone.0026164

Ike, H., Inaba, Y., Kobayashi, N., Hirata, Y., Yukizawa, Y., Aoki, C., Choe, H., & Saito, T. (2015). Comparison between mechanical stress and bone mineral density in the femur after total hip arthroplasty by using subject-specific finite element analyses. *Computer Methods in Biomechanics and Biomedical Engineering*, *18*(10), 1056–1065. https://doi.org/10.1080/10255842.2013.869320

Jonason, J. H., Hoak, D., & O'Keefe, R. J. (2015). Primary Murine Growth Plate and Articular Chondrocyte Isolation and Cell Culture. In *Osteoporosis and Osteoarthritis* (Vol. 1226, pp. 11–18). https://doi.org/10.1007/978-1-4939-1619-1_2

Kariya, S., Park, G. H., Maeno-Hikichi, Y., Leykekhman, O., Lutz, C., Arkovitz, M. S., Landmesser, L. T., & Monani, U. R. (2008). Reduced SMN protein impairs maturation of the neuromuscular junctions in mouse models of spinal muscular atrophy. *Human Molecular Genetics*, *17*(16), 2552–2569. https://doi.org/10.1093/hmg/ddn156

Kariya, S., Obis, T., Garone, C., Akay, T., Sera, F., Iwata, S., Homma, S., & Monani, U. R. (2014). Requirement of enhanced Survival Motoneuron protein imposed during neuromuscular junction maturation. Journal of Clinical Investigation, 124(2), 785–800. https://doi.org/10.1172/JCI72017

Khatri, I. A., Chaudhry, U. S., Seikaly, M. G., Browne, R. H., & Iannaccone, S. T. (2008). Low bone mineral density in spinal muscular atrophy. *Journal of Clinical Neuromuscular Disease*, *10*(1), 11–17. https://doi.org/10.1097/CND.0b013e318183e0fa

Kim, J. K., Jha, N. N., Feng, Z., Faleiro, M. R., Chiriboga, C. A., Wei-Lapierre, L., Dirksen, R. T., Ko, C. P., & Monani, U. R. (2020). Muscle-specific SMN reduction reveals motor neuron–independent disease in spinal muscular atrophy models. Journal of Clinical Investigation, 130(3), 1271–1287. https://doi.org/10.1172/JCI131989

Kim, J. H., Kang, J. S., Yoo, K., Jeong, J., Park, I., Park, J. H., Rhee, J., Jeon, S., Jo, Y. W., Hann, S. H., Seo, M., Moon, S., Um, S. J., Seong, R. H., & Kong, Y. Y. (2022). Bap1/SMN axis in Dpp4+ skeletal muscle mesenchymal cells regulates the neuromuscular system. *JCI Insight*, *7*(10). https://doi.org/10.1172/jci.insight.158380

Kong, L., Wang, X., Choe, D. W., Polley, M., Burnett, B. G., Bosch-Marcé, M., Griffin, J. W., Rich, M. M., & Sumner, C. J. (2009). Impaired synaptic vesicle release and immaturity of neuromuscular junctions in spinal muscular atrophy mice. *Journal of Neuroscience*, *29*(3), 842–851. https://doi.org/10.1523/JNEUROSCI.4434-08.2009

Leinroth, A. P., Mirando, A. J., Rouse, D., Kobayahsi, Y., Tata, P. R., Rueckert, H. E., Liao, Y., Long, J. T., Chakkalakal, J. V., & Hilton, M. J. (2022). Identification of distinct non-myogenic skeletal-muscle-resident mesenchymal cell populations. *Cell Reports*, *39*(6), 110785. https://doi.org/10.1016/j.celrep.2022.110785

Ling, K. K. Y., Lin, M. Y., Zingg, B., Feng, Z., & Ko, C. P. (2010). Synaptic defects in the spinal and neuromuscular circuitry in a mouse model of spinal muscular atrophy. *PLoS ONE*, *5*(11). https://doi.org/10.1371/journal.pone.0015457

Martinez, T. L., Kong, L., Wang, X., Osborne, M. A., Crowder, M. E., Van Meerbeke, J. P., Xu, X., Davis, C., Wooley, J., Goldhamer, D. J., Lutz, C. M., Rich, M. M., & Sumner, C. J. (2012). Survival Motor Neuron Protein in Motor Neurons Determines Synaptic Integrity in Spinal Muscular Atrophy. *Journal of Neuroscience*, *32*(25), 8703–8715. https://doi.org/10.1523/JNEUROSCI.0204-12.2012

Mattar, C. N., Waddington, S. N., Biswas, A., Johana, N., Ng, X. W., Fisk, A. S., Fisk, N. M., Tan, L. G., Rahim, A. A., Buckley, S. M. K., Tan, M. H., Lu, J., Choolani, M., & Chan, J. K. Y. (2013). Systemic delivery of scAAV9 in fetal macaques facilitates neuronal transduction of the central and peripheral nervous systems. *Gene Therapy*, *20*(1), 69–83. https://doi.org/10.1038/gt.2011.216

McGovern, V. L., Iyer, C. C., David Arnold, W., Gombash, S. E., Zaworski, P. G., Blatnik, A. J., Foust, K. D., & Burghes, A. H. M. (2015). SMN expression is required in motor neurons to rescue electrophysiological deficits in the SMNΔ7 mouse model of SMA. *Human Molecular Genetics*, *24*(19), 5524–5541. https://doi.org/10.1093/hmg/ddv283

Mercuri, E., Pera, M. C., Scoto, M., Finkel, R., & Muntoni, F. (2020). Spinal muscular atrophy — insights and challenges in the treatment era. *Nature Reviews Neurology*, *16*(12), 706–715. https://doi.org/10.1038/s41582-020-00413-4

Modla, S., Mendonca, J., Czymmek, K. J., & Akins, R. E. (2010). Identification of neuromuscular junctions by correlative confocal and transmission electron microscopy. *Journal of Neuroscience Methods*, *191*(2), 158–165. https://doi.org/10.1016/j.jneumeth.2010.06.015

Monani, U. R., Pastore, M. T., Gavrilina, T. O., Jablonka, S., Le, T. T., Andreassi, C., DiCocco, J. M., Lorson, C., Androphy, E. J., Sendtner, M., Podell, M., & Burghes, A. H.

M. (2003). A transgene carrying an A2G missense mutation in the SMN gene modulates phenotypic severity in mice with severe (type I) spinal muscular atrophy. *Journal of Cell Biology*, *160*(1), 41–52. https://doi.org/10.1083/jcb.200208079

Monani, U. R., Sendtner, M., Coovert, D. D., Parsons, D. W., Andreassi, C., Le, T. T., Jablonka, S., Schrank, B., Rossol, W., Prior, T. W., Morris, G. E., & Burghes, A. H. M. (2000). The human centromeric survival motor neuron gene (SMN2) rescues embryonic lethality in Smn(-/-) mice and results in a mouse with spinal muscular atrophy. *Human Molecular Genetics*, *9*(3), 333–339. https://doi.org/10.1093/hmg/9.3.333

Nassari, S., Duprez, D., & Fournier-Thibault, C. (2017). Non-myogenic Contribution to Muscle Development and Homeostasis: The Role of Connective Tissues. *Frontiers in Cell and Developmental Biology*, *5*(March), 1–17. https://doi.org/10.3389/fcell.2017.00022

Ovchinnikov, D. (2009). Alcian blue/alizarin red staining of cartilage and bone in mouse. *Cold Spring Harbor Protocols*, *4*(3), 5170–5172. https://doi.org/10.1101/pdb.prot5170

Passini, M. A., Bu, J., Roskelley, E. M., Richards, A. M., Sardi, S. P., O'Riordan, C. R., Klinger, K. W., Shihabuddin, L. S., & Cheng, S. H. (2010). CNS-targeted gene therapy improves survival and motor function in a mouse model of spinal muscular atrophy. Journal of Clinical Investigation, 120(4), 1253–1264. https://doi.org/10.1172/JCI41615

Passini, M. A., Bu, J., Richards, A. M., Kinnecom, C., Sardi, S. P., Stanek, L. M., Hua, Y., Rigo, F., Matson, J., Hung, G., Kaye, E. M., Shihabuddin, L. S., Krainer, A. R., Bennett, C. F., & Cheng, S. H. (2011). Antisense oligonucleotides delivered to the mouse CNS ameliorate symptoms of severe spinal muscular atrophy. *Science Translational Medicine*, *3*(72), 1–12. https://doi.org/10.1126/scitranslmed.3001777

Penaud-Budloo, M., Le Guiner, C., Nowrouzi, A., Toromanoff, A., Chérel, Y., Chenuaud, P., Schmidt, M., von Kalle, C., Rolling, F., Moullier, P., & Snyder, R. O. (2008). Adeno-Associated Virus Vector Genomes Persist as Episomal Chromatin in Primate Muscle. *Journal of Virology*, *82*(16), 7875–7885. https://doi.org/10.1128/jvi.00649-08

Petrany, M. J., Swoboda, C. O., Sun, C., Chetal, K., Chen, X., Weirauch, M. T., Salomonis, N., & Millay, D. P. (2020). Single-nucleus RNA-seq identifies transcriptional heterogeneity in multinucleated skeletal myofibers. *Nature Communications*, *11*(1). https://doi.org/10.1038/s41467-020-20063-w

Ross, J. A., Webster, R. G., Lechertier, T., Reynolds, L. E., Turmaine, M., Bencze, M., Jamshidi, Y., Cetin, H., Muntoni, F., Beeson, D., Hodilvala-Dilke, K., & Conti, F. J. (2017). Multiple roles of integrin-α3 at the neuromuscular junction. Journal of Cell Science, 130(10), 1772–1784. https://doi.org/10.1242/jcs.201103

Sahashi, K., Ling, K. K. Y., Hua, Y., Wilkinson, J. E., Nomakuchi, T., Rigo, F., Hung, G., Xu, D., Jiang, Y. P., Lin, R. Z., Ko, C. P., Bennett, C. F., & Krainer, A. R. (2013).

Pathological impact of SMN2 mis-splicing in adult SMA mice. *EMBO Molecular*
*Medicine*, *5*(10), 1586–1601. https://doi.org/10.1002/emmm.201302567

Scott, R. W., Arostegui, M., Schweitzer, R., Rossi, F. M. V., & Underhill, T. M. (2019). Hic1
Defines Quiescent Mesenchymal Progenitor Subpopulations with Distinct Functions and
Fates in Skeletal Muscle Regeneration. *Cell Stem Cell*, *25*(6), 797-813.e9.
https://doi.org/10.1016/j.stem.2019.11.004

Shababi, M., Lorson, C. L., & Rudnik-Schöneborn, S. S. (2014). Spinal muscular atrophy: A
motor neuron disorder or a multi-organ disease? *Journal of Anatomy*, *224*(1), 15–28.
https://doi.org/10.1111/joa.12083

Sheng, G. (2015). The developmental basis of mesenchymal stem/stromal cells (MSCs).
*BMC Developmental Biology*, *15*(1), 26–31. https://doi.org/10.1186/s12861-015-0094-5

Torres-Benito, L., Neher, M. F., Cano, R., Ruiz, R., & Tabares, L. (2011). SMN requirement
for synaptic vesicle, active zone and microtubule postnatal organization in motor nerve
terminals. PLoS ONE, 6(10). https://doi.org/10.1371/journal.pone.0026164

Tsai, L. K., Chen, C. L., Ting, C. H., Lin-Chao, S., Hwu, W. L., Dodge, J. C., Passini, M. A.,
& Cheng, S. H. (2014). Systemic administration of a recombinant AAV1 vector
encoding IGF-1 improves disease manifestations in SMA mice. Molecular Therapy,
22(8), 1450–1459. https://doi.org/10.1038/mt.2014.84

Uezumi, A., Ikemoto-Uezumi, M., Zhou, H., Kurosawa, T., Yoshimoto, Y., Nakatani, M.,
Hitachi, K., Yamaguchi, H., Wakatsuki, S., Araki, T., Morita, M., Yamada, H., Toyoda,
M., Kanazawa, N., Nakazawa, T., Hino, J., Fukada, S. I., & Tsuchida, K. (2021).
Mesenchymal Bmp3b expression maintains skeletal muscle integrity and decreases in
age-related sarcopenia. *Journal of Clinical Investigation*, *131*(1).
https://doi.org/10.1172/JCI139617

Vai, S., Bianchi, M. L., Moroni, I., Mastella, C., Broggi, F., Morandi, L., Arnoldi, M. T.,
Bussolino, C., & Baranello, G. (2015). Bone and Spinal Muscular Atrophy. *Bone*, *79*,
116–120. https://doi.org/10.1016/j.bone.2015.05.039

Valori, C. F., Ning, K., Wyles, M., Mead, R. J., Grierson, A. J., Shaw, P. J., & Azzouz, M.
(2010). Systemic delivery of scAAV9 expressing SMN prolongs survival in a model of
spinal muscular atrophy. *Science Translational Medicine*, *2*(35), 1–9.
https://doi.org/10.1126/scitranslmed.3000830

Van Alstyne, M., Tattoli, I., Delestrée, N., Recinos, Y., Workman, E., Shihabuddin, L. S.,
Zhang, C., Mentis, G. Z., & Pellizzoni, L. (2021). Gain of toxic function by long-term
AAV9-mediated SMN overexpression in the sensorimotor circuit. *Nature Neuroscience*.
https://doi.org/10.1038/s41593-021-00827-3

Wasserman, H. M., Hornung, L. N., Stenger, P. J., Rutter, M. M., Wong, B. L., Rybalsky, I.,
Khoury, J. C., & Kalkwarf, H. J. (2017). Low bone mineral density and fractures are

highly prevalent in pediatric patients with spinal muscular atrophy regardless of disease
severity. *Neuromuscular Disorders*, *27*(4), 331–337.
https://doi.org/10.1016/j.nmd.2017.01.019

Wilk, K., Yeh, S. C. A., Mortensen, L. J., Ghaffarigarakani, S., Lombardo, C. M., Bassir, S.
H., Aldawood, Z. A., Lin, C. P., & Intini, G. (2017). Postnatal Calvarial Skeletal Stem
Cells Expressing PRX1 Reside Exclusively in the Calvarial Sutures and Are Required
for Bone Regeneration. *Stem Cell Reports*, *8*(4), 933–946.
https://doi.org/10.1016/j.stemcr.2017.03.002

Woszczyna, M. N., Konishi, C. T., Perez Carbajal, E. E., Wang, T. T., Walsh, R. A., Gan, Q.,
Wagner, M. W., & Rando, T. A. (2019). Mesenchymal Stromal Cells Are Required for
Regeneration and Homeostatic Maintenance of Skeletal Muscle. *Cell Reports*, *27*(7),
2029-2035.e5. https://doi.org/10.1016/j.celrep.2019.04.074

Yakar, S., Rosen, C. J., Beamer, W. G., Ackert-Bicknell, C. L., Wu, Y., Liu, J. L., Ooi, G. T.,
Setser, J., Frystyk, J., Boisclair, Y. R., & LeRoith, D. (2002). Circulating levels of IGF-1
directly regulate bone growth and density. *Journal of Clinical Investigation*, *110*(6),
771–781. https://doi.org/10.1172/JCI0215463

Yakar, S., Rosen, C. J., Bouxsein, M. L., Sun, H., Mejia, W., Kawashima, Y., Wu, Y.,
Emerton, K., Williams, V., Jepsen, K., Schaffler, M. B., Majeska, R. J., Gavrilova, O.,
Gutierrez, M., Hwang, D., Pennisi, P., Frystyk, J., Boisclair, Y., Pintar, J., … LeRoith, D.
(2009). Serum complexes of insulin-like growth factor-1 modulate skeletal integrity and
carbohydrate metabolism. *The FASEB Journal*, *23*(3), 709–719.
https://doi.org/10.1096/fj.08-118976

Yang, Y. S., Xie, J., Wang, D., Kim, J. M., Tai, P. W. L., Gravallese, E., Gao, G., & Shim, J.
H. (2019). Bone-targeting AAV-mediated silencing of Schnurri-3 prevents bone loss in
osteoporosis. *Nature Communications*, *10*(1), 1–13. https://doi.org/10.1038/s41467-019-
10809-6

Yeo, C. J. J., & Darras, B. T. (2020). Overturning the Paradigm of Spinal Muscular Atrophy
as Just a Motor Neuron Disease. *Pediatric Neurology*, *109*, 12–19.
https://doi.org/10.1016/j.pediatrneurol.2020.01.003

Yesbek Kaymaz, A., Bal, A., Bora-Tatar, G., Ozon, A., Alikasifoglu, A., Topaloglu, H., &
Erdem-Yurter, H. (2016). Serum IGF1 and IGFBP3 levels in SMA patients.
*Neuromuscular Disorders*, *26*, S105. https://doi.org/10.1016/j.nmd.2016.06.075

Zanetti, G., Negro, S., Megighian, A., & Pirazzini, M. (2018). Electrophysiological
Recordings of Evoked End-Plate Potential on Murine Neuro-muscular Synapse
Preparations. *Bio-Protocol*, *8*(8), 1–16. https://doi.org/10.21769/bioprotoc.2803

**Figure 1**

Skeletal growth abnormalities by limb mesenchymal cell-specific SMN depletion in SMN2 1-copy Smn$^{\Delta MPC}$ mice. (**A**) Representative 3D images of ossified femur bone. Scale bars, 1 mm. (**B-C**) SMN2 1-copy mutant's femurs showed reduced growth in diaphysis length and diameter, and (**D**) decreased trabecular bone volume. (**E-F**) Trabecular and cortical bone thicknesses were not significantly different between the control and mutant groups. (**G**) Bone mineral density was slightly increased in the diaphysis of SMN2 1-copy Smn$^{\Delta MPC}$ mice. The micro CT analysis was performed in femur diaphysis and metaphysis from SMN2 1-copy Smn$^{\Delta WT}$, SMN2 2-copy, and 1-copy Smn$^{\Delta MPC}$ mice at P14. 1-way ANOVA with Tukey's post hoc test, n = 3-5 mice in each genotype (**B-G**). (**H**) Representative images of genomic PCR analysis from SMN2 1-copy Smn$^{\Delta MPC}$ mice tissues at P21. (**I**) qRT PCR analysis from tissues of SMN2 1-copy Smn$^{WT}$ and Smn$^{\Delta MPC}$ mice at P21 (n = 3 mice in each genotype; Unpaired t-test with Welch's correction). Deletion of Smn exon 7 was detected only in limb mesenchymal cells using genomic PCR (**H**) and mRNA expression (**I**). (**J**) Representative images of western blot analysis in cultured FAPs. SMN protein in FAPs of SMN2 1-copy Smn$^{\Delta MPC}$ mice exhibited a decrease comparable to that observed in the SMA$\Delta$7 mice. (**K**) Relative SMN levels in cultured FAPs of the controls and mutants (n = 3 mice in each genotype; 1-way ANOVA with Tukey's post hoc test). ns; not significantly different. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$. Error bars show s.e.m.

**Figure 1 -figure supplement 1**

Growth defects in the Prrx1-lineage bone of SMN2 0-copy Smn$^{\Delta MPC}$ mice. **(A)** Representative control and mutant mice with 0 copies of SMN2 were photographed at E18.5. **(B-C)** Alcian blue & alizarine red staining in the SMN2 0-copy Smn$^{\Delta MPC}$ mutant showed abnormal skeletal development of the limbs, **(D)** calvaria (shown by the dashed line), and **(E)** sternum (indicated by a white arrow). Scale bars, 5 mm **(B-C)** and 2 mm **(D-E)**.

**Figure 1 -figure supplement 2**

Altered growth plate homeostasis independent of the IGF axis from the liver in SMN2 1-copy Smn$^{\Delta MPC}$ mice. (**A**) Representative images of H&E staining in the femur growth plate of control and mutant mice with 1 copy of SMN2 at P14. Scale bars, 200 μm (Left) and 100 μm (Right). (**B-D**) Hypertrophic zone (HZ) and proliferative zone (PZ), indicated by black arrows and dashed lines, lengths were reduced in SMN2 1-copy Smn$^{\Delta MPC}$ mice, and the hypertrophic cell number in a section of the 1-copy mutant was decreased (n = 4 mice in each genotype; Unpaired t-test with Welch's correction). (**E**) Relative IGF axis mRNA expression in the livers of SMN2 1-copy control and mutant mice. The IGF pathway genes showed no difference when comparing controls to SMN2 1-copy Smn$^{\Delta MPC}$ mutants (n = 3 mice in each genotype; Unpaired t-test with Welch's correction). ns; not significantly different. *$p < 0.05$; **$p < 0.01$. Error bars show s.e.m.

**A**    Synaptophysin/AChR/Neurofilament

SMN2 1cp Smn^WT

SMN2 1cp Smn^ΔMPC

**B** Presynaptic terminal arborization (average terminals/NMJ) ****

**C** Poor arborization % (terminals<4) ***

**D** Nerve terminals with NF varicosities (% total) *

☐ SMN2 1-copy Smn^WT
■ SMN2 1-copy Smn^ΔMPC

**E**
● SMN2 1-copy Smn^WT
■ SMN2 1-copy Smn^ΔMPC

Percentage(%)

Plaque  Small perforated  Large perforated  Open  Branched

**F**    SYN/AChR/NF

SMN2 1cp Smn^WT

SMN2 1cp Smn^ΔMPC

**G** AChR cluster size(μm²) ns

**H** Terminal covered by synaptophysion (%) ***

**Figure 2**

Aberrant postnatal NMJ maturation in SMN2 1-copy $Smn^{\Delta MPC}$ mice. (**A**) Immunostaining of NMJs in TA muscle of SMN2 1-copy $Smn^{WT}$ and $Smn^{\Delta MPC}$ mice at P21 with anti-NF (Green), anti-synaptophysin (Blue), α-Btx staining AChR (Red). Scale bars, 20 μm. The confocal images of NMJs showed decreased presynaptic terminal branching and the existence of nerve terminal varicosities that were enlarged with NF (indicated by arrowheads) in the mutant. (**B**) The NMJs of the SMN2 1-copy mutant exhibited a significant decrease in presynaptic terminal arborization and (**C**) an increased percentage of poorly arborized NMJs (n = 3 mice in each genotype; Unpaired t-test with Welch's correction). (**D**) The percentage of NMJs exhibiting NF varicosities was higher in the SMN2 1-copy mutant group than in the control group (n = 3-4 mice in each genotype; Unpaired t-test with Welch's correction). (**E**) For quantification of the NMJ maturation stage, we classified NMJs into five distinct developmental stages (Plaque: Plaque-shaped endplate without any perforation; Small perforated: Plaque-shaped endplate with small perforations; Large perforated: Plaque-shaped endplate with large perforations; Open: C-shaped endplate; Branched: Pretzel-like branched -shaped endplate) and then compared the frequency patterns of SMN2 1-copy control and mutant mice (n = 3 mice in each genotype; 2-way ANOVA with Tukey's post hoc test). The NMJs of SMN2 1-copy mutants displayed plaque-like shapes, indicating that they were in the immature stage. (**F**) Immunostaining of NMJs in TA muscle of SMN2 1-copy $Smn^{WT}$ and $Smn^{\Delta MPC}$ mice at P3 with anti-NF (Green), anti-synaptophysin (Blue), α-Btx staining AChR (Red). Scale bars, 10 μm. (**G**) There were no significant differences in AChR cluster size between the SMN2 1-copy control and mutant at P3 (n = 3-4 mice in each genotype; Unpaired t-test with Welch's correction). (**H**) The ratio of the Synaptophysin area to the AChR area in NMJ was slightly higher in the SMN2 1-copy mutant at P3 (n = 3-4 mice in each genotype; Unpaired t-test with

Welch's correction). ns; not significantly different. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$. All box-and-whisker plots show the median, interquartile range, minimum, and maximum. For the bar and line graph, error bars show s.e.m.

**A**

Synaptophysin/AChR/Neurofilament

SMN2 1cp Smn^WT

SMN2 2cp Smn^ΔMPC

SMN2 1cp Smn^ΔMPC

**B**

Presynaptic fragments/NMJ

****
****
ns

**C**

Nerve terminals with NF varicosities (% total)

****
****
ns

**D**

AChR fragments/NMJ

****
****
ns

☐ SMN2 1-copy Smn^WT
▨ SMN2 2-copy Smn^ΔMPC
■ SMN2 1-copy Smn^ΔMPC

**Figure 3**

Morphological deterioration in NMJs of adult SMN2 1-copy $Smn^{\Delta MPC}$ mice. (**A**) Immunostaining of NMJs in TA muscle of SMN2 1-copy $Smn^{WT}$, SMN 2-copy, and SMN2 1-copy $Smn^{\Delta MPC}$ mice at P56 with anti-NF (Green), anti-synaptophysin (Blue), α-Btx staining AChR (Red). Scale bars, 20 μm. The confocal images of NMJs showed fragmentation and bouton-like NF varicosities (indicated by arrowheads) in the SMN2 1-copy $Smn^{\Delta MPC}$ mice. (**B-D**) The NMJs of the SMN2 1-copy mutant displayed fragmented presynapse (**B**), endplate (**D**), and NF varicosities (**C**) compared to SMN2 1-copy $Smn^{WT}$ and SMN2 2-copy $Smn^{\Delta MPC}$ mice (n = 3-5 mice in each genotype; Presynaptic fragments & AChR fragments: Brown–Forsythe and Welch ANOVA with Games-Howell's test; NF varicosities: 1-way ANOVA with Tukey's post hoc test). ns; not significantly different. ****$p < 0.0001$. All box-and-whisker plots show the median, interquartile range, minimum, and maximum. For the bar graph, error bars show s.e.m.

**Figure 4**

Reduced presynaptic neurotransmission ability in the NMJs of SMN2 1-copy Smn$^{\Delta MPC}$ mice. **(A)** Representative traces of mEPP from SMN2 1-copy Smn$^{\Delta WT}$ (top) and SMN2 1-copy Smn$^{\Delta MPC}$ (bottom) mice. **(B-C)** SMN2 1-copy mutant's NMJs showed an increase in mEPP amplitude and no differences in mEPP frequency (1-copy control, n = 25, 9 mice; 1-copy mutant, n = 21, 8 mice; Unpaired t-test with Welch's correction). **(D)** Representative traces of eEPP from SMN2 1-copy Smn$^{\Delta WT}$ (top) and SMN2 1-copy Smn$^{\Delta MPC}$ (bottom) mice. **(E)** The mutant's NMJs showed a stronger amplitude of eEPPs (1-copy control, n = 12, 4 mice; 1-copy mutant, n = 12, 3 mice; Unpaired t-test with Welch's correction). **(F)** Representative traces of paired-pulse response from SMN2 1-copy Smn$^{\Delta WT}$ (top) and SMN2 1-copy Smn$^{\Delta MPC}$ (bottom) mice. **(G)** Paired-pulse response was not different between SMN2 1-copy control and mutant NMJs, indicating a comparable neurotransmitter release probability (1-copy control, n = 8, 3 mice; 1-copy mutant, n = 6, 3 mice; Unpaired t-test with Welch's correction). The electrophysiological recording was performed in the EDL muscle at P56. ns; not significantly different. *$p < 0.05$. All box-and-whisker plots show the median, interquartile range, minimum, and maximum.

**A**

SMN2 1cp Smn^WT          SMN2 1cp Smn^ΔMPC

**B**
ns

Junctional folds density (#/µm)

**C**
***

Synaptic vesicle density (#/µm²)

**D**
*

Detached nerve terminal %

☐ SMN2 1-copy Smn^WT
■ SMN2 1-copy Smn^ΔMPC

**Figure 5**

Abnormal nerve terminal ultrastructure in SMN2 1-copy mutant. **(A)** Representative TEM images from NMJs of SMN2 1-copy Smn$^{\Delta WT}$ and SMN2 1-copy Smn$^{\Delta MPC}$ mice at P56. Scale bars, 500 nm. Nerve terminal (NT; indicated by the blue zone). Synaptic vesicles (SVs). Muscle fiber (MS; indicated by the red zone). Endplate junctional folds (JF). Nerve terminal detachment (indicated by arrow) was observed in SMN2 1-copy Smn$^{\Delta MPC}$ mice. **(B)** The density of junctional folds in the NMJ of SMN2 1-copy Smn$^{\Delta MPC}$ mice no significant change compared to the control, whereas **(C)** the density of synaptic vesicles was increased (n = 3-4 mice in each genotype; Unpaired t-test with Welch's correction). **(D)** The detachment of the nerve terminal occurs more frequently at the NMJ of mutants (n = 3-4 mice in each genotype; Unpaired t-test with Welch's correction). ns; not significantly different. *$p < 0.05$; ***$p < 0.001$. All box-and-whisker plots show the median, interquartile range, minimum, and maximum. For the bar graph, error bars show s.e.m.

**A**

**B**

Presynaptic fragments/NMJ

****

**C**

Nerve terminals with
NF varicosities (% total)

***

**D**

AChR fragments/NMJ

**

■ SMN2 1-copy Smn$^{\Delta MPC}$+Veh
■ SMN2 1-copy Smn$^{\Delta MPC}$+FAP

**Figure 6**

Improved postnatal NMJ development in the TA muscle of SMN2 1-copy Smn$^{\Delta MPC}$ mice following healthy FAPs transplantation. (**A**) Immunostaining of NMJs in tdTomato$^+$ FAPs-transplanted TA muscle (+FAP) and vehicle-treated contralateral muscle (+Veh) in SMN2 1-copy Smn$^{\Delta MPC}$ mice at P56 with anti-NF (Yellow), anti-synaptophysin (Blue), α-Btx staining AChR (Green), and tdTomato fluorescence (Red). Scale bars, 40 μm. The images revealed NF varicosities (indicated by arrowheads) in the +Veh NMJs. The tdTomato+ FAPs (marked by asterisks) were transplanted into +FAP NMJs, which exhibited (**B**) decreased presynaptic fragmentation, (**C**) NF varicosities, and (**D**) AChR fragmentation compared to +Veh NMJs. (n = 4 mice in each group; Unpaired t-test with Welch's correction). ns; not significantly different. **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. All box-and-whisker plots show the median, interquartile range, minimum, and maximum. For the bar graph, error bars show s.e.m.