A wave of minor *de novo* DNA methylation initiates in mouse 8-cell embryos and co-regulates imprinted X- chromosome inactivation with H3K27me3

Yuan Yue[1], Wei Fu[1], Qianying Yang[1], Chao Zhang[1], Wenjuan Wang[1], Meiqiang Chu[1], Qingji Lyu[1], Yawen Tang[1], Jian Cui[1], Xiaodong Wang[1], Zhenni Zhang[1], Jianhui Tian[1], Lei An[1*]

[1] Frontiers Science Center for Molecular Design Breeding of the Ministry of Education, State Key Laboratory of Animal Biotech Breeding, National Engineering Laboratory for Animal Breeding; Key Laboratory of Animal Genetics, Breeding and Reproduction of the Ministry of Agriculture and Rural Affairs, College of Animal Science and Technology, China Agricultural University, Beijng, China.

*Authors for correspondence (anleim@cau.edu.cn)

Keywords: preimplantation embryos, *de novo* DNA methylation, DNA methyltransferase 3B, imprinted X chromosome inactivation, polycomb repressive complexes 2

Summary statement

A minor wave of *de novo* DNA methylation has been initiated prior to blastocyst formation, but not during the implantation period, and co-regulates imprinted X-chromosome inactivation.

ABSTRACT

DNA methylation is extensively reprogrammed during early stage of mammalian

development and is essential for normal embryogenesis. It is well established that mouse

embryos acquire genome-wide DNA methylation during implantation, referred to as *de novo*

DNA methylation, from globally hypomethylated blastocysts. However, the fact that the main

de novo DNA methyltransferase 3B (DNMT3B) is initially expressed as early as the 8-cell

stage, contradicts the current knowledge about timing of initiation of *de novo* DNA

methylation. Here, we reported that a previously overlooked minor wave of *de novo* DNA

methylation initially occurs during the transition from the 8-cell to blastocyst stage, before the

well-known large-scale *de novo* DNA methylation during implantation. Functional analyses

indicated that minor *de novo* DNA methylation regulates proliferation, lineage differentiation

and metabolic homeostasis of preimplantation embryos, and is critical for embryonic

developmental potential and pregnancy outcomes. Furthermore, bioinformatic and functional

analyses indicated that minor *de novo* DNA methylation preferentially occurs on the X

chromosome and co-regulates imprinted X-chromosome inactivation via the interaction

between DNMT3B and polycomb repressive complexes 2 core components during blastocyst

formation. Thus, our study updates the current knowledge of embryonic *de novo* DNA

methylation, thereby providing a novel insight of early embryonic epigenetic reprogramming.

INTRODUCTION

Mammalian early embryos are co-regulated by a series of spatiotemporally restricted

epigenomic events that are critical for acquiring developmental potential. In mouse, it has

been widely accepted that a wave of genome-wide *de novo* DNA methylation occurs around

the time of implantation, thus re-establishing DNA methylation patterns from a globally

hypomethylated blastocysts to relatively static DNA methylation patterns in post-implantation

embryos (Borgel et al., 2010; Smith et al., 2012a). DNA methyltransferase 3B (DNMT3B) is

the main enzyme that catalyzes embryonic *de novo* DNA methylation. Genetic depletion of

Dnmt3b results in hypomethylation and dysregulation of pluripotency genes, gastrulation

genes, germline-specific genes, *etc* (Auclair et al., 2014; Borgel et al., 2010), thus causing

severe developmental defects and lethality after implantation (Li et al., 1992; Okano et al.,

1999). However, despite its developmental importance, the exact timing of initiation of *de*

novo DNA methylation remains largely elusive.

Although earlier results using high-throughput sequencing methods suggested that *de novo*

DNA methylation occurs during the transition from blastocysts to early post-implantation

embryos (Auclair et al., 2014; Borgel et al., 2010; Guo et al., 2014; Smith et al., 2012a), our

detection assays (Figure 1A, B) and other previous studies (Guenatri et al., 2013; Hirasawa et

al., 2008; Li et al., 2016) indicated that *Dnmt3b,* the main *de novo* DNA methyltransferase, is

initially expressed as early as the 4-cell stage, and shows clear nuclear localization at 8-cell

stage. This fact challenges current knowledge about timing of initiation of embryonic *de novo*

DNA methylation, and therefore raising the possibility that embryonic *de novo* DNA

methylation may precede, but not follows, the blastocyst stage.

To test this possibility, we reanalyzed well-established bisulfite sequencing data that

profiles DNA methylation maps from the mouse preimplantation to post-implantation

embryos (Smith et al., 2012a), and found approximately one-third of genes gained promoter

DNA methylation during the transition from the 8-cell to blastocyst stage. It means that

despite global DNA demethylation throughout the preimplantation development, a small

proportion of regions have initiated *de novo* DNA methylation prior to the blastocyst stage,

which we designated as minor *de novo* DNA methylation. We next ask its developmental

functions for embryogenesis. The results showed that minor *de novo* methylation also

regulates proliferation, lineage differentiation and metabolism of preimplantation embryos,

and thus affecting the developmental potential and final pregnancy outcomes. Of interest,

minor *de novo* DNA methylation is more prone to occur on the X chromosome. We next

confirmed that minor *de novo* DNA methylation plays an important role in establishing

imprinted X chromosome inactivation (iXCI) via the interaction between DNMT3B and the

polycomb repressive complexes 2 (PRC2) core components, which are critical for X-linked

heterochromatin. Thus, our study provides a novel understanding of *de novo* DNA

methylation during early development, which updates the current knowledge of embryonic

epigenomic hallmark events and their interactions.

RESULTS AND DISCUSSION

A small proportion of promoters initiate *de novo* DNA methylation by the 8-cell stage

To investigate the earliest initiation of *de novo* DNA methylation, we first examined the expression patterns of *Dnmt3s* in mouse preimplantation embryos. Results from previously published transcriptome (Fan et al., 2015) and our quantitative RT-PCR (qRT-PCR) detection showed that expression of *Dnmt3b*, the main enzyme for embryonic *de novo* DNA methylation (Borgel et al., 2010; Okano et al., 1999), as well as its non-catalytic partner *Dnmt3l* (Gowher et al., 2005), initiated by the 8-cell stage, and were further increased upon preimplantation development (Figure 1A). By contrast, the expression of *Dnmt3a*, which is mainly required for gametic methylation (Kaneda et al., 2004a), was maintained at low levels concurrently (Figure 1A). The initial expression of *Dnmt3b* was also confirmed at the protein level. Nuclear location of DNMT3B can be clearly detected from the 8-cell stage onward (Figure 1B). Given the essential role of DNMT3B in catalyzing *de novo* DNA methylation, we postulated that embryonic *de novo* DNA methylation may initiate as early as the 8-cell stage, which challenges current knowledge that *de novo* DNA methylation initiates during implantation.

To test the hypothesis, we next reanalyzed a well-established reduced representation bisulfite sequencing (RRBS) data that profiles DNA methylation maps from the mouse preimplantation to post-implantation stage (Smith et al., 2012a). Promoter regions are important 's target sites of DNMT3B (Choi et al., 2011). The acquisition of DNA methylation in promoters, especially in intermediate and low CpG promoters, during implantation is largely dependent on DNMT3B, and plays an important role in regulating developmental genes (Auclair et al., 2014; Borgel et al., 2010; Dahlet et al., 2020). Thus, among genomic regions that may undergo *de novo* DNA methylation, we initially focused our analysis on DNA methylation dynamics of promoters from the 8-cell stage onward. Despite the global loss of promoter DNA methylation (Figure 1—figure supplement 1A), more detailed characterization showed that a considerable proportion of promoters (approximately 30%) gained DNA methylation during the transition from the 8-cell to blastocyst stage, even though most promoters underwent demethylation (Figure 1C). Similar observation was also obtained

by reanalyzing another published whole-genome bisulfite sequencing (WGBS) data (Smith et al., 2017) (Figure 1—figure supplement 1B). In addition, a reanalysis based on a publicly available single cell multi-omics sequencing (COOL-seq) data of mouse early embryos (Guo et al., 2017) showed that both male and female embryos gain DNA methylation during the transition from the 8-cell to blastocyst stage (Figure 1—figure supplement 1C, D).

Many previous analyses using different methodologies have showed the promoter regions are relatively hypomethylated in general compared with other genome elements (Auclair et al., 2014; Borgel et al., 2010; Dahlet et al., 2020; Smith et al., 2012b). Despite this, by tracking DNA methylation dynamics of two subsets of promoters that gained or lost DNA methylation from the 8-cell to blastocyst stage respectively, we found promoters that underwent demethylation showed a relatively higher DNA methylation levels following fertilization, and then were gradually demethylated until reaching their lowest value in inner cell mass (ICM) (Figure 1D). In contrast, promoters that gained DNA methylation from the 8-cell to blastocyst stage, showed a relatively lower DNA methylation levels, following fertilization, and reached their lowest DNA methylation levels at the 8-cell stage (Figure 1E). Based on the low basal levels, these promoters gain DNA methylation from the 8-cell to ICM, and then continued to gain DNA methylation during implantation (Figure 1E). To distinguish two waves of *de novo* DNA methylation before and after the blastocyst stage, we designated them as minor and major *de novo* DNA methylation, respectively. By performing gene ontology enrichment analysis of genes with promoter regions that undergo minor or major *de novo* DNA methylation respectively, we noticed that besides of many important basic processes common to two waves of *de novo* DNA methylation, genes subject to minor *de novo* DNA methylation have been enriched in processes such as organic substance transport, chromosome organization, and cell fate specification (Figure 1F). Compared with conventionally accepted major *de novo* DNA methylation, minor *de novo* DNA methylation showed a smaller increase in promoter DNA methylation that tended to occur in low CpG-containing promoters (Figure 1—figure supplement 1E).

Minor *de novo* DNA methylation participates in embryonic proliferation, differentiation and affects developmental potential

It has been known that *de novo* DNA methylation during implantation stage plays an essential

role in transcriptional regulation of developmental genes that are critical for embryonic

differentiation and survival (Auclair et al., 2014; Borgel et al., 2010), thus we next attempted

to test the short-and long-term developmental consequences of minor *de novo* DNA

methylation. To this end, we performed an integrated analysis using the publicly available

bisulfite sequencing data (Smith et al., 2012a) and transcriptome data (Wu et al., 2016) that

profiles DNA methylation and gene expression maps from the 8-cell to ICM, respectively.

Putative genes transcriptionally regulated by minor *de novo* DNA methylation (Figure

2—figure supplement 1A) were functionally related to many basic processes, such as cell

differentiation and metabolic regulation. Phenotype annotations in Mouse Genome

Informatics (MGI) showed that genes enriched in these basic processes are critical for normal

embryonic survival, development and postnatal growth (Figure 2—figure supplement 1B, C).

It should be noted that genes subject to DNA demethylation also participate in similar

processes (Figure 2—figure supplement 1D). In line with our results, previous studies using

Tet1-deficient or *Tet1/3*-deficient embryos, indicated the critical role of TET-mediated DNA

demethylation in these processes, as well as embryonic and postnatal development (Dawlaty

et al., 2011; Ito et al., 2010; Kang et al., 2015). Thus, it can be presumed that minor *de novo*

DNA methylation, and TET-mediated demethylation, may synergistically co-regulate DNA

methylation homeostasis before blastocyst formation, thus fine-tuning embryonic

developmental potential and long-term outcomes, enriching the knowledge of developmental

significance of DNA methylation homeostasis in early embryos, as proposed previously

(Iurlaro et al., 2017; Smith and Meissner, 2013).

We next wanted to examine whether minor *de novo* methylation is functionally associated

with these basic biological processes before blastocyst formation and in turn affects

developmental potential and pregnancy outcomes. Given identification of long-term

developmental consequence of minor *de novo* DNA methylation requires that only minor, but

not major *de novo* methylation, is functionally inactivated before blastocyst formation, we

knocked down *Dnmt3b* in preimplantation embryos by microinjecting *Dnmt3b* siRNA into

zygotes (Figure 2—figure supplement 2A-E). As expected, *Dnmt3b* mRNA and protein levels

were significantly reduced in morulae, and tended to be lower in blastocysts compared to

those of the negative control (NC) group (Figure 2—figure supplement 2A-C). The inhibition

of *Dnmt3b* resulted in a significant decrease in the global level of 5mC staining before

blastocyst formation (Figure 2—figure supplement 2D, E). *Dnmt3b* deficiency not only

inhibited proliferation (Figure 2B), but also changed lineage differentiation of blastocysts

(Figure 2C). The decreased EdU-positive signals, as well as the slight lower total cell number,

suggested an inhibited proliferation due to *Dnmt3b* knockdown (KD). Since energy

metabolism is crucial for supporting embryo differentiation and development, and oxidative

phosphorylation (OXPHOS) is activated during the blastocyst stage (Zhao et al., 2021), we

next examined the energy metabolism, in particular OXPHOS activity, of *Dnmt3b*-KD

embryos. Despite the unchanged total ATP production, *Dnmt3b* deficiency led to a

significantly disruption in OXPHOS-dependent-ATP production (Figure 2D). These

observations were further confirmed via the transient treatment of 5-aza-dC specifically

during the window of minor *de novo* DNA methylation (Figure 2—figure supplement 3A-D).

5-aza-dC is a well-established global DNA hypomethylating agent that efficiently inhibit the

activity of all DNMTs and thus has been frequently used to study the maintenance of DNA

methylation and *de novo* DNA methylation, despite its inhibitory effect common to various

DNMTs (Maslov et al., 2012; Oka et al., 2005). Transient treatment of embryos with 5-aza-dC

specifically during the window of minor *de novo* DNA methylation allows us to test its

function in a temporally specific manner. Thus, it can be presumed that the disrupted

proliferation and energy metabolism, appears not to be due to defective major *de novo* DNA

methylation. Finally, we transferred *Dnmt3b*-KD embryos into pseudopregnant recipient

females. Compared with that in NC group, *Dnmt3b*-KD embryos resulted in a significantly

reduced implantation rate and live birth rate (Figure 2E-F), suggesting the important role of

minor *de novo* DNA methylation in influencing embryonic developmental potential.

Minor *de novo* DNA methylation preferentially occurs on the X chromosome

and plays an important role in iXCI

To obtain a global view of minor *de novo* DNA methylation, we next detected the

chromosome-wide distribution of *de novo* methylated promoters from the 8-cell to ICM, and

found that they were globally distributed across autosomes, with the percentage ranging from

23% to 33% (Figure 3A-B). Of note, approximately half of X-linked promoters (48%)

underwent minor *de novo* DNA methylation throughout the X chromosome (Figure 3—figure

supplement 1A), much higher than those of autosomal promoters (Figure 3B). This result was

similar with that of major *de novo* DNA methylation during implantation (Figure 3—figure

supplement 1B, C). Using the single cell COOL-seq data, we also specifically reanalyzed the

DNA methylation changes on the X chromosome in female embryos. The X chromosome

showed a more notable increase than that on autosomes, and the female X chromosome

showed a higher DNA methylation level than that of the male (Figure 3—figure supplement

2A, B). The X chromosome-biased prevalence of *de novo* DNA methylation is reminiscent of

XCI, a female-specific epigenetic event that occurs during early development for balancing

the X-linked gene dosage between males and females. In mice, early embryos undergo two

waves of XCI: iXCI initiates and occurs specifically on the paternal X chromosome in

preimplantation embryos, and then persists in extraembryonic tissues; while random X

chromosome inactivation (rXCI) occurs randomly on either the maternal or paternal X

chromosome in the embryonic cells by implantation stage (Augui et al., 2011; Chow and

Heard, 2009; Lee and Bartolomei, 2013). Spatiotemporal co-occurrence of minor *de novo*

DNA methylation and iXCI on the X chromosome, led us to ask if these two epigenetic events

are functional linked during preimplantation development.

Earlier studies have focused on whether the DNA methylation imprinting established

during oogenesis as the imprinting mark responsible for the monoallelic expression of *Xist*

(the initiation event of iXCI) in preimplantation embryos and demonstrated that oocyte DNA

methylation is dispensable for *Xist* imprinting (Chiba et al., 2008; Kaneda et al., 2004b; Sado

et al., 2004). However, the role of DNA methylation in achieving X chromosomal

heterochromatinization (the late event of XCI) remains controversial, although it has been

well-accepted DNA methylation is tightly associated with maintaining X chromosome-wide

transcriptional silence of rXCI during implantation (Blewitt et al., 2008; Sado et al., 2000).

Recently, targeted DNA demethylation of genes on the X chromosome, using inactive Cas9

protein (dCas9) fused with the catalytic domain of ten-eleven translocation dioxygenase 1

(TET1), supported the causal role between DNA methylation and XCI status (Halmai et al.,

2020). However, the conclusion is based on rXCI, the role of DNA methylation in the

heterochromatinization of iXCI remains unclear, and this issue has been documented as a

long-standing open question of epigenetic programming during preimplantation development

(Csankovszki et al., 2001; Kaneda et al., 2004b; Sado et al., 2000; Sado et al., 2004). Actually,

it is even plausibly thought that DNA methylation is not required for achieving

heterochromatinization of iXCI because preimplantation embryos is undergoing global and

massive DNA demethylation. To identify the biological function of minor *de novo* DNA

methylation in iXCI, we knocked down *Dnmt3b* in preimplantation embryos by

microinjecting *Dnmt3b* siRNA into zygotes (Figure 3C). Next, we detected the H3K27me3, a

classic marker for establishment of XCI achieving X chromosome-wide

heterochromatinization of transcriptional depression (Chow and Heard, 2009; Huynh and Lee,

2005; Wutz, 2011). In line with the decreased DNA methylation levels (Figure 2—figure

supplement 2D, E), we found nearly all detected *Dnmt3b*-KD female blastocysts contained

trophoblast cells lacking the H3K27me3 domain, with a variable frequency ranging from 10.7%

to 42.4%; whereas nearly all NC female blastocysts showed a single H3K27me3 domain in

each trophoblast (Figure 3D-E). The proportion H3K27me3-positive cells in total trophoblast

cells were significantly lower due to *Dnmt3b*-KD (Figure 3F), indicating *Dnmt3b* deficiency

leads to an impaired iXCI in preimplantation female embryos. This result was confirmed

using *Dnmt3b*-knockout (KO) embryos (Figure 3—figure supplement 3A-D). To further

validate the function of minor *de novo* DNA methylation in iXCI, we take advantage of

chemical-induced DNMT inhibition using 5-aza-dC (Figure 3—figure supplement 3E-H). In

addition, the involvement of minor *de novo* DNA methylation in iXCI establishment was also

supported by the finding that many representative X-linked non-escaping genes tended to be

de novo methylated during the transition from the 8-cell to ICM (Figure 3—figure supplement

3I-J).

To test whether DNA methylation loss leads to a prolonged effect on iXCI, we further

reanalyzed the RNA-seq and H3K27me3 CHIP-seq data in extraembryonic ectoderm (ExE)

of E6.5 single embryos that underwent *Dnm3a/3b* knockout because preimplantation iXCI

status maintains in extraembryonic cells (Chen et al., 2019; Galupa and Heard, 2015; Schulz

and Heard, 2013). The results showed that chromosome-wide loss of DNA methylation led to

a nearly complete loss of H3K27me3 on paternal X chromosome (specifically inactivated in

iXCI), along with a notable transcriptional upregulation cross the chromosome, including the

locus of some non-escaping genes. By contrast, these changes cannot be not observed on

maternal X chromosome (Figure 3—figure supplement 4A, B).

Taken together, these data suggest that a wave of minor *de novo* DNA methylation

preceding the blastocyst stage, participate in the establishment of iXCI. Although a recent

study reported that *Dnmt3b* has a dominant role on the X chromosome in mouse embryonic

fibroblasts (Yagi et al., 2020), the detectable DNMT3A/*Dnmt3a* in preimplantation embryos,

albeit at low levels (Uysal et al., 2017; Zhang et al., 2015), led us to speculate that DNMT3B

and DNMT3A synergistically act on minor *de novo* DNA methylation, similar with that

reported in subsequent peri-implantation embryos (Borgel et al., 2010; Chen et al., 2019).

269   Minor *de novo* DNA methylation co-regulates iXCI via the interaction between

270   DNMT3B and PRC2 core components

Having confirmed the functional role of minor *de novo* methylation in iXCI, we next explored

the underlying mechanism. Because iXCI is initiated by *Xist* RNA upregulation and coating

during the 8-cell and morular stage (Penny et al., 1996), we first detected *Xist* mRNA

expression. No significant difference in the degrees of absence of the *Xist* domain in each

blastomere nucleus between NC and *Dnmt3b*-KD female morulae (Figure 4A, B), which were

also confirmed by detecting *Dnmt3b*-KO female morulae (Figure 4—figure supplement 1A,

B). In line with this, single female embryo qRT-PCR analysis of *Xist* and its essential

upstream activator *Rnf12* during initial stage of iXCI, showed that the expression levels of

Xist and *Rnf12* were not affected by *Dnmt3b* deficiency (Figure 4C). Our results demonstrate

that minor *de novo* DNA methylation may participate in establishment, but not in initiation of

iXCI.

In attempt to further understand the mechanism underlying the role of minor *de novo* DNA

methylation in establishing iXCI, we next asked whether the *Dnmt3b* deficiency-induced loss

of H3K27me3 domains was related to abnormal expression of histone-modifiers. We detected

the expression of H3K27 histone methyltransferases or the partner (*Ezh2*, *Suz12, Eed*) and

demethylase (*Kdm6a*, *Kdm6b*). The results showed *Dnmt3b* deficiency has no impact on the

expression levels of these chromatin modifiers (Figure 4D).

DNA methylation and H3K27me3 have been shown to have complex relationship, possibly

in a cell-type and/or genomic region-specific manner (King et al., 2016). Under certain

circumstances, DNA methylation shows antagonistic effect to H3K27me3 at promoters, via

excluding the binding of PRC2 components to their targets (Bartke et al., 2010; Jermann et al., 2014), while other studies have presented alternative evidence that PRC2 and DNMT cooperate to achieve silencing (Vire et al., 2006), and has a synergistic relationship in the process of global heterochromatin formation (Hagarman et al., 2013; King et al., 2016). A recent work reported that maintenance of noncanonical imprinting requires maternal allele-specific *de novo* DNA methylation in extraembryonic cells (Chen et al., 2019). It has been thought that during the late phase of rXCI in fully differentiated cells, gene silencing is initially achieved by PRC2 complex-induced H3K27me3, and further stably maintained by the redundant action of multiple layers of epigenetic modifications, including DNA methylation, to reach the maximum level of chromatin compaction (Chow and Heard, 2009; Heard et al., 2004; Pintacuda and Cerase, 2015). In line with this, a recent multifaceted analysis showed that DNA methylation and H3K27me3 are concurrently enriched in genes subject to XCI (Balaton and Brown, 2021). Thus, we next reanalyzed H3K27me3 dynamics of promoters that underwent minor *de novo* DNA methylation during the transition from the 8-cell to ICM, using a well-established H3K27me3 ChIP-seq data (Liu et al., 2016). As expected, concurrent increase in H3K27me3 enrichment can be clearly observed throughout the X chromosome and many *de novo* methylated X-linked promoters (Figure 4—figure supplement 2A, B). Next, we used *in situ* proximity ligation assays (PLA), a method that is suitable for visualizing protein interactions by detecting the close proximity of two molecules (Soderberg et al., 2006) to test interactions between DNMT3B and PRC2 core components, *i.e.,* EZH2, SUZ12, EED, which are essential for X-linked heterochromatin. It is clearly visible that DNMT3B-EZH2, DNMT3B-SUZ12 and DNMT3B-EED interactions can be detected in the nuclei by the blastocyst stage (Figure 4E). In line with this, the function of *de novo* DNA methylation in XCI, as well as direct interaction between DNMT3B and PRC2 core components, were also confirmed in differentiated female ES cells (Figure 4—figure supplement 3A-H). Collectively, our results support the idea that minor *de novo* DNA methylation plays an important role in iXCI via synergistic interaction with heterochromatic histone modifications, rather than transcriptional regulation of iXCI-regulating genes.

It is also worth mentioning that minor *de novo* DNA methylation is definitely distinct from previously reported DNA re-methylation in monkey and human preimplantation embryos,

which occurs during the 2- to 8-cell and 4- to 8-cell transition respectively (Gao et al., 2017; Guo et al., 2014; Zhu et al., 2018). The so-called DNA re-methylation does not occur in naturally conceived mouse embryos. However, an aberrant wave of DNA re-methylation has been identified in cloned mouse preimplantation embryos as an epigenetic barrier that impairs full-term developmental potential (Gao et al., 2018).

Collectively, our study presents an update on the current knowledge of embryonic *de novo* DNA methylation. We identified that a wave of minor *de novo* DNA methylation has initially occurred during the transition from the mouse 8-cell to blastocyst stage, but not during the implantation stage. Of interest, we provide the evidence that DNA methylation is important for establishing iXCI in preimplantation female embryos. Our data, together with previous results, also support a model where minor *de novo* DNA methylation and demethylation co-orchestrate DNA methylation homeostasis before blastocyst formation, and serve as epigenetic mechanisms affecting embryonic developmental potential by fine-tuning the processes of lineage commitment, proliferation and metabolic homeostasis (Figure 4F). Thus, our work provides a novel insight for understanding epigenetic programming and reprogramming during early embryonic development.

MATERIALS AND METHODS

Mouse embryo preparation

All experimental procedures were approved by and performed in accordance with the guidelines of the Institutional Animal Care and Use Committee of China Agricultural University. Mice were fed ad libitum and housed under controlled lighting (12 light:12 dark) and specific pathogen-free conditions. ICR female mice were superovulated with 5 IU pregnant mare serum gonadotrophin (PMSG; Ningbo Hormone Product Co., Ltd, Ningbo, China) and a further 5 IU human chorionic gonadotropin (hCG; Ningbo Hormone Product CO., Ltd) 46-48 h later. Then, superovulated ICR females were cogaged individually with ICR males. *In vivo* zygote, 2-cell embryos, 4-cell embryos, 8-cell embryos, morulae, or blastocysts were flushed from oviducts and uterus at 16 h, 42 h, 54 h, 66 h, 78 h, or 93 h post-hCG, respectively.

The *Dnmt3b*[tm2Enl]/+ mouse line(Okano et al., 1999) was purchased from Mutant Mouse

351   Resource & Research Centers (MMRRC, 029886-UNC). For collection of *Dnmt3b*[+/+] (WT)

352   and *Dnmt3b* [tm2Enl/ tm2Enl] (KO) embryos, *Dnmt3b*[tm2Enl/+] female mice were superovulated as

353   above, and mated with *Dnmt3b*[tm2Enl/+] male mice. Morulae and blastocysts were collected at

354   78 h and 93 h, respectively.

Quantitative real-time PCR analysis

Total RNA was extracted from embryos using TRIzol[TM] (Invitrogen, Carlsbad, CA, USA).

RNA was treated with DNase, and reverse-transcribed into cDNA using the HiScript II Q

Select RT Kit (Vazyme, Nanjing, China). qRT-PCR was performed using SsoFast Eva Green

Supermix (Bio-Rad, Hercules, CA, USA). The relative expression data were calculated using

the $2^{-\Delta Ct}$ method. Relative expression level was normalized to the housekeeping gene *H2afz*,

and each experiment was repeated at least three times, as previously reported. Primers used

are listed in Table S1.

For qRT-PCR on single female or male embryo, a blastomere was isolated and treated with

2 μL Cells-to-cDNA™ II Cell Lysis Buffer (Invitrogen) at 75°C for 15 min. Next, the lysate

was digested with proteinase K (Merck, Kenilworth, NJ, USA) at 55°C for 2 h, and used for

sex determination according to the manufacturer's instruction of Ex Taq Hot Start Version

(Takara Bio, Shiga, Japan). The amplification was conducted as follows: 95 °C for 3 min; 40

cycles of 95 °C for 10 s, 52 °C for 15 s, 72 °C for 15 s; and extension at 72 °C for 5 min. The

remaining part of embryo was treated with 5 μL Cells-to-cDNA™ II Cell Lysis Buffer

(Invitrogen), and then reverse-transcribed into cDNA with the HiScript II Q Select RT Kit

(Vazyme).

Immunofluorescence analysis

Embryos were fixed with 4% Paraformaldehyde (PFA) in PBS-0.1% PVA at 4 °C overnight.

After permeabilizing with 0.5% Triton X-100 in PBS-0.1%PVA (PBST-PVA) for 1 h at room

temperature, embryos were blocked with 1% BSA in PBST-PVA at 4 °C for 6 h. Embryos

were incubated with primary anti-DNMT3B antibody (1:200; GeneTex, Irvine, CA, USA),

anti-H3K27me3 antibody (1:1000; MilliporeSigma, Burlington, MA, USA), or anti CDX2

antibody (1:200; BioGenex, Fremont, CA, USA) overnight at 4 °C, followed by Alexa

Fluor-488 (anti-mouse; Invitrogen) and Alexa Fluor-594 (anti-rabbit; Invitrogen) labeled

secondary antibodies for 1 h at room temperature. Finally, embryos were counterstained with

DAPI. Fluorescence images were obtained under an upright microscope (BX51; Olympus)

using an attached digital microscope camera (DP72; Olympus). Quantification of the

immunofluorescence results were performed using Image J (National Institutes of Health,

Bethesda, MD, USA).

For 5-methylcytosine (5-mC) staining, permeabilized embryos were treated with 4 M HCl

for 10 min and 100 mM Tris-HCl for 10 min at room temperature, and then blocked overnight

at 4 °C. Next, embryos were incubated with primary anti-5mC antibody (1:200; Active Motif,

Carlsbad, CA, USA) at room temperature for 2 h, followed by secondary antibody for 1 h.

Furthermore, embryos for sex determination and genotyping were removed from the slides

and digested with Cells-to-cDNA™ II Cell Lysis Buffer (Invitrogen) and proteinase K (Merck)

as described above, followed by sex determination and genotyping using PCR. For

genotyping, the amplification was conducted as follows: 94 °C for 3 min; 40 cycles of 94 °C

for 30 s, 65 °C for 30 s, 72 °C for 45 s; and extension at 72 °C for 7 min. The primer

sequences are listed in Table S1.

Bisulfite sequencing

The DNA methylation profile was analyzed by PCR-based bisulfite sequencing. 40 IVO

blastocysts or 80 8-cell embryos were digested and bisulfite converted according to the

manufacturer's instruction of EZ DNA Methylation-Direct Kit (Zymo, Irvine, CA, USA).

Bisulfite-specific primers were designed with the online Methyl Primer Express, v1.0

(http://www.urogene.org/methprimer/ Table S1). The converted DNA was amplified by PCR

using Ex Taq Hot Start Version (Takara). And the PCR products were subcloned into

pEASY-T5 Zero Cloning Kit (TransGen, Beijing, China). At least 10 clones per group were

sequenced. The sequencing results were analyzed with QUMA (http://quma.cdb.riken.jp/).

Microinjection of siRNA

siRNA oligo sequences were synthesized by GenePharma (Shanghai, China). The sequences

of *Dnmt3b* siRNA used in the present study were as follows: sense

5'-CCUCAAGACAAAUAGCUAUTT-3', antisense

5'-AUAGCUAUUUGUCUUGAGGTT-3'. siRNA was diluted to 100 μM and stored at

-80 °C. For siRNA microinjection, 5 pL siRNA was microinjected into the cytoplasm of

zygotes. Then embryos were cultured in KSOM at 37 °C under 5% CO_2 and collected at the

8-cell, morula or blastocyst stage for further analyses.

Treatment of 5-aza-dC

Zygotes were collected as described above, and 10 nM 5-aza-2'-deoxycytidine (5-aza-dC) was

supplemented into *in vitro* culture medium from the 8-cell stage onward. Blastocysts were

collected at 110 h post-hCG administration.

RNA-FISH

RNA-FISH was performed using ViewRNA™ ISH Cell Assay Kit (Invitrogen) according to

the manufacturer's instructions. Fixed embryos were treated with Detergent Solution QC at

room temperature for 5 min, and digested with protease QS at room temperature for 10 min.

Next, the embryos were hybridized with probes against *Xist* (Invitrogen) at 40 °C for 3 h.

Then the embryos were hybridized with PreAmplifier and Amplifier Mix at 40 °C for 30 min,

respectively. Label Probe Mix was used to produce a signal. The embryos were

counterstained with DAPI, and imaged using an attached digital microscope camera (DP72;

Olympus).

Proximity ligation assays

The in situ proximity ligation assays (PLA) was performed using the Duolink® In Situ Red

Starter Kit according to the manufacturer's instruction (MilliporeSigma). Embryos were fixed

and permeabilized as described for immunofluorescence. After blocking, embryos were

incubated with primary antibody. Embryos were then incubated with PLA probes, followed

by ligation and amplification reaction. Finally, embryos were counterstained with Duolink®

In Situ Mounting Medium with DAPI (MilliporeSigma). Furthermore, DNMT3B, EZH2,

SUZ12, or EED single primary antibody, and secondary antibodies alone were used as

technical negative controls.

Generation of Dnmt3a/Dnmt3b KO ESC

Mouse ESCs line PGK12.1 was used in the present study. Cells were cultured in KnockOut™

DMEM medium (Invitrogen) containing 10% fetal bovine serum (VISTECH, Sydney,

Australia), 2mM GlutaMAX (Invitrogen), 1×non-essential amino acid (MilliporeSigma), 55

μM β-mercaptoethanol (Invitrogen), 1000 units/ml mouse recombinant leukemia inhibitor

factor (MilliporeSigma), 100 units/ml penicillin and 100 µg/ml streptomycin (both

Invitrogen). To induce ESC differentiation, ESCs were cultured in DMEM/F12 (Invitrogen)

and Neurobasal medium supplement with B27 and N2 supplement (Invitrogen), 100 units/ml

penicillin-streptomycin (Invitrogen).

Knockout of *Dnmt3b* or *Dnmt3a* for ES cells was performed using CRISPR/Cas9n. sgRNA

was designed using Benchling (https://www.benchling.com/). sgRNAs were constructed to

pSpCas9n(BB)-2A-Puro vector (PX462, Addgene plasmid no. 62987). 4 μg of the plasmid

was transfected using Lipofectamine 2000 (Invitrogen) according to the manufacturer's

protocol. 2 μg/ml puromycin (Solarbio, Beijing, China) was used to selected the positive

colonies. After selection, colonies were picked out and expanded. Cells were verified by

genotyping and Sanger sequencing. The sgRNA sequences and knockdown cell detection

primers are listed in Table S1.

Co-immunoprecipitation

Co-immunoprecipitation (Co-IP) was performed with Dynabeads™ Protein G

Immunoprecipitation Kit (Invitrogen) according to the manufacturer's protocol. Briefly, 3 μg

DNMT3B antibody (GeneTex) or mouse IgG (MilliporeSigma) was incubated to Protein G

beads with rotation for 10 minutes at room temperature. The beads were washed with

Washing Buffer. Then cell lysates were then incubated with the beads-antibody complex with

rotation for 4h at 4 °C. Then the beads were washed 3 times with Washing Buffer, and were

eluted with loading buffer by heating at 98 °C for 10min. Co-IP products were analyzed by

western blot using the indicated primary antibodies and secondary antibodies. And protein

bands were detected with Tanon 5200 detection systems (Tanon, Shanghai, China).

Bioinformatics analysis

RRBS data of mouse embryos (2-cell embryos, 4-cell embryos, 8-cell embryos, ICM, and

E6.5 embryos) were downloaded from the published article by Smith et al (Smith et al.,

2012b) (accession number: GSE34864). The methylation level was calculated as the number

of "methylated" reads (reporting as C), divided by the total number of "methylated" and

"unmethylated" read, which reporting as C or T. The genomic region information was

downloaded from the mm9 Repeat Masker. As described in the published article, promoters

were defined as 1 kb up- and downstream of the TSS and classified into high-density CpG

promoter (HCP), intermediate-density CpG promoter (ICP) and low-density CpG promoter

(LCP). Only CpG sites with at least fivefold coverage were included in the methylation

analysis. WGBS data of 8-cell and ICM were from GSE84236. Cool-seq data of female and

male embryos were from GSE78140, and analyzed as described in the article (Guo et al.,

2017). Briefly, every covered WCG site (W includes A or T) with at least one-fold coverage

was summed for the analysis of DNA methylation levels. The bigwig files of RRBS,

RNA-seq and H3K27me3 CUT&RUN data of WT and Dnmt3a/3b DKO E6.5 ExE were

downloaded from the published article by Chen et al (Chen et al., 2019) (accession number:

GSE130115). The bigwig files of H3K27me3 ChIP-seq data of 8-cell embryos and ICM were

downloaded from the published article by Liu et al. (Liu et al., 2016) (accession number:

GSE73952).

Gene ontology (GO) annotation analysis was performed using Database for Annotation,

Visualization and Integrated Discovery (DAVID 6.8: https://david.ncifcrf.gov/). Heatmaps

were plotted using the R software "pheatmap" package (version 1.0.12). The integrative

genomics viewer (IGV) was used to visualize the ChIP-seq results. The phenotype of genes

was analyzed on the Mouse Genome Informatics (MGI; http://www.informatics.jax.org/)

database.

EdU assay

The proliferation of embryos was detected with a BeyoClick™ EdU-594 Cell Proliferation

Kit (Beyotime, Shanghai, China) according to the manufacturer's instruction. Embryos were

incubated in EmbryoMax® KSOM Mouse Embryo Media (MilliporeSigma) containing 10

μM EdU at 37 °C for 2 h, and then fixed and permeabilized. Afterwards, embryos were

incubated with the Click Reaction mixture for 2 min at room temperature before stained with

DAPI. Images were acquired under an upright microscope (BX51; Olympus) using an

attached digital microscope camera (DP72; Olympus).

ATP measurement

ATP level was measured using an ATP determination kit according to the manufacturer's

instruction (Beyotime). Briefly, blastocysts were incubated with M2 medium, and 1 μM

oligomycin (Abmore, Houston, USA) or 1 mM sodium oxamate (MilliporeSigma) in M2

medium at 37 °C for 15 min, respectively. Next, embryos were digested with 20 μL ATP lysis

buffer on ice, and the lysate was added to 96-well plates containing 100 μL ATP detection

working dilution. The luminescence was detected by a multifunctional microplate reader

(Infinite F200; TECAN, Mannedorf, Switzerland).

Embryo transfer

503 Pseudo-pregnant ICR females were mated with ICR males 3.5 d before embryo transfer.

504 Blastocysts derives from NC or *Dnmt3b*-KD group were transferred to the uterine horn of

505 pseudo-pregnant female mice.

Statistical analysis

507 All data are presented as mean ± SD and analyzed with the student's *t* test or correct χ^2

508 procedure by using SPSS v.25.0 (IBM, Armonk, NY, USA). Values of $P < 0.05$ were

509 considered statistically significant.

Data availability

512 All data are available in the article and/or supporting information.

Acknowledgements

We thank Neil Brockdorff (University of Oxford) and Ingolf Bach (University of

Massachusetts Medical School) for the gift and delivery of PGK12.1 cells. We thank Zihuan

Du for the assistance in the analysis of sequencing data. This work was supported by grants

from the National Natural Science Foundation of China (31930103), National Agriculture

Key Science & Technology Project (NK20221201), Ningbo Major Science and Technology

Project (2021Z112), the National Key R&D Program (2022YFD1300301).

Author contributions

523 Y.Y., J.T. and L.A. conceived and designed the experiments. Y.Y. C.Z. and W.W. were

524 responsible for animal care and management. Y.Y., W.F., Q.Y., C.Z., W.W., M.C., Q.L., Y.T.,

525 J.C., X.W. and Z.Z. performed the experiments. Y.Y., W.F., J.T. and L.A. analyzed the data.

526 Y.Y., W.F., J.T. and L.A wrote, and finalized the manuscript. All authors approved the paper.

Competing interests

529 The authors declare no competing interests.

References

Auclair, G., Guibert, S., Bender, A. and Weber, M. (2014). Ontogeny of CpG island methylation and specificity of DNMT3 methyltransferases during embryonic development in the mouse. *Genome Biol.* **15**, 545.

Augui, S., Nora, E. P. and Heard, E. (2011). Regulation of X-chromosome inactivation by the X-inactivation centre. *Nat. Rev. Genet.* **12**, 429-442.

Balaton, B. P. and Brown, C. J. (2021). Contribution of genetic and epigenetic changes to escape from X-chromosome inactivation. *Epigenetics Chromatin* **14**, 30.

Bartke, T., Vermeulen, M., Xhemalce, B., Robson, S. C., Mann, M. and Kouzarides, T. (2010). Nucleosome-interacting proteins regulated by DNA and histone methylation. *Cell* **143**, 470-484.

Blewitt, M. E., Gendrel, A. V., Pang, Z., Sparrow, D. B., Whitelaw, N., Craig, J. M., Apedaile, A., Hilton, D. J., Dunwoodie, S. L., Brockdorff, N., et al. (2008). SmcHD1, containing a structural-maintenance-of-chromosomes hinge domain, has a critical role in X inactivation. *Nat. Genet.* **40**, 663-669.

Borgel, J., Guibert, S., Li, Y., Chiba, H., Schubeler, D., Sasaki, H., Forne, T. and Weber, M. (2010). Targets and dynamics of promoter DNA methylation during early mouse development. *Nat. Genet.* **42**, 1093-1100.

Chen, Z., Yin, Q., Inoue, A., Zhang, C. and Zhang, Y. (2019). Allelic H3K27me3 to allelic DNA methylation switch maintains noncanonical imprinting in extraembryonic cells. *Sci Adv* **5**, eaay7246.

Chiba, H., Hirasawa, R., Kaneda, M., Amakawa, Y., Li, E., Sado, T. and Sasaki, H. (2008). De novo DNA methylation independent establishment of maternal imprint on X chromosome in mouse oocytes. *Genesis* **46**, 768-774.

Choi, S. H., Heo, K., Byun, H. M., An, W., Lu, W. and Yang, A. S. (2011). Identification of preferential target sites for human DNA methyltransferases. *Nucleic Acids Res.* **39**, 104-118.

Chow, J. and Heard, E. (2009). X inactivation and the complexities of silencing a sex chromosome. *Curr. Opin. Cell Biol.* **21**, 359-366.

Csankovszki, G., Nagy, A. and Jaenisch, R. (2001). Synergism of Xist RNA, DNA methylation, and histone hypoacetylation in maintaining X chromosome inactivation. *J. Cell Biol.* **153**, 773-784.

Dahlet, T., Argueso Lleida, A., Al Adhami, H., Dumas, M., Bender, A., Ngondo, R. P., Tanguy, M., Vallet, J., Auclair, G., Bardet, A. F., et al. (2020). Genome-wide analysis in the mouse embryo reveals the importance of DNA methylation for transcription integrity. *Nat Commun* **11**, 3153.

Dawlaty, M. M., Ganz, K., Powell, B. E., Hu, Y. C., Markoulaki, S., Cheng, A. W., Gao, Q., Kim, J., Choi, S. W., Page, D. C., et al. (2011). Tet1 is dispensable for maintaining pluripotency and its loss is compatible with embryonic and postnatal development. *Cell Stem Cell* **9**, 166-175.

Fan, X., Zhang, X., Wu, X., Guo, H., Hu, Y., Tang, F. and Huang, Y. (2015). Single-cell

RNA-seq transcriptome analysis of linear and circular RNAs in mouse
preimplantation embryos. *Genome Biol.* **16**, 148.

Galupa, R. and Heard, E. (2015). X-chromosome inactivation: new insights into cis and
trans regulation. *Curr. Opin. Genet. Dev.* **31**, 57-66.

Gao, F., Niu, Y., Sun, Y. E., Lu, H., Chen, Y., Li, S., Kang, Y., Luo, Y., Si, C., Yu, J., et al.
(2017). De novo DNA methylation during monkey pre-implantation embryogenesis.
Cell Res. **27**, 526-539.

Gao, R., Wang, C., Gao, Y., Xiu, W., Chen, J., Kou, X., Zhao, Y., Liao, Y., Bai, D., Qiao,
Z., et al. (2018). Inhibition of Aberrant DNA Re-methylation Improves
Post-implantation Development of Somatic Cell Nuclear Transfer Embryos. *Cell*
Stem Cell **23**, 426-435 e425.

Gowher, H., Liebert, K., Hermann, A., Xu, G. and Jeltsch, A. (2005). Mechanism of
stimulation of catalytic activity of Dnmt3A and Dnmt3B
DNA-(cytosine-C5)-methyltransferases by Dnmt3L. *J. Biol. Chem.* **280**,
13341-13348.

Guenatri, M., Duffie, R., Iranzo, J., Fauque, P. and Bourc'his, D. (2013). Plasticity in
Dnmt3L-dependent and -independent modes of de novo methylation in the
developing mouse embryo. *Development* **140**, 562-572.

Guo, F., Li, L., Li, J., Wu, X., Hu, B., Zhu, P., Wen, L. and Tang, F. (2017). Single-cell
multi-omics sequencing of mouse early embryos and embryonic stem cells. *Cell Res.*
27, 967-988.

Guo, H., Zhu, P., Yan, L., Li, R., Hu, B., Lian, Y., Yan, J., Ren, X., Lin, S., Li, J., et al.
(2014). The DNA methylation landscape of human early embryos. *Nature* **511**,
606-610.

Hagarman, J. A., Motley, M. P., Kristjansdottir, K. and Soloway, P. D. (2013). Coordinate
regulation of DNA methylation and H3K27me3 in mouse embryonic stem cells. *PLoS*
One **8**, e53880.

Halmai, J., Deng, P., Gonzalez, C. E., Coggins, N. B., Cameron, D., Carter, J. L.,
Buchanan, F. K. B., Waldo, J. J., Lock, S. R., Anderson, J. D., et al. (2020).
Artificial escape from XCI by DNA methylation editing of the CDKL5 gene. *Nucleic*
Acids Res. **48**, 2372-2387.

Heard, E., Chaumeil, J., Masui, O. and Okamoto, I. (2004). Mammalian X-chromosome
inactivation: an epigenetics paradigm. *Cold Spring Harb. Symp. Quant. Biol.* **69**,
89-102.

Hirasawa, R., Chiba, H., Kaneda, M., Tajima, S., Li, E., Jaenisch, R. and Sasaki, H.
(2008). Maternal and zygotic Dnmt1 are necessary and sufficient for the maintenance
of DNA methylation imprints during preimplantation development. *Genes Dev.* **22**,
1607-1616.

Huynh, K. D. and Lee, J. T. (2005). X-chromosome inactivation: a hypothesis linking
ontogeny and phylogeny. *Nat. Rev. Genet.* **6**, 410-418.

Ito, S., D'Alessio, A. C., Taranova, O. V., Hong, K., Sowers, L. C. and Zhang, Y. (2010).
Role of Tet proteins in 5mC to 5hmC conversion, ES-cell self-renewal and inner cell
mass specification. *Nature* **466**, 1129-1133.

Iurlaro, M., von Meyenn, F. and Reik, W. (2017). DNA methylation homeostasis in human

and mouse development. *Curr. Opin. Genet. Dev.* **43**, 101-109.

Jermann, P., Hoerner, L., Burger, L. and Schubeler, D. (2014). Short sequences can efficiently recruit histone H3 lysine 27 trimethylation in the absence of enhancer activity and DNA methylation. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E3415-3421.

Kaneda, M., Okano, M., Hata, K., Sado, T., Tsujimoto, N., Li, E. and Sasaki, H. (2004a). Essential role for de novo DNA methyltransferase Dnmt3a in paternal and maternal imprinting. *Nature* **429**, 900-903.

Kaneda, M., Sado, T., Hata, K., Okano, M., Tsujimoto, N., Li, E. and Sasaki, H. (2004b). Role of de novo DNA methyltransferases in initiation of genomic imprinting and X-chromosome inactivation. *Cold Spring Harb. Symp. Quant. Biol.* **69**, 125-129.

Kang, J., Lienhard, M., Pastor, W. A., Chawla, A., Novotny, M., Tsagaratou, A., Lasken, R. S., Thompson, E. C., Surani, M. A., Koralov, S. B., et al. (2015). Simultaneous deletion of the methylcytosine oxidases Tet1 and Tet3 increases transcriptome variability in early embryogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E4236-4245.

King, A. D., Huang, K., Rubbi, L., Liu, S., Wang, C. Y., Wang, Y., Pellegrini, M. and Fan, G. (2016). Reversible Regulation of Promoter and Enhancer Histone Landscape by DNA Methylation in Mouse Embryonic Stem Cells. *Cell Rep.* **17**, 289-302.

Lee, J. T. and Bartolomei, M. S. (2013). X-inactivation, imprinting, and long noncoding RNAs in health and disease. *Cell* **152**, 1308-1323.

Li, E., Bestor, T. H. and Jaenisch, R. (1992). Targeted mutation of the DNA methyltransferase gene results in embryonic lethality. *Cell* **69**, 915-926.

Li, Y., Seah, M. K. and O'Neill, C. (2016). Mapping global changes in nuclear cytosine base modifications in the early mouse embryo. *Reproduction* **151**, 83-95.

Liu, X. Y., Wang, C. F., Liu, W. Q., Li, J. Y., Li, C., Kou, X. C., Chen, J. Y., Zhao, Y. H., Gao, H. B., Wang, H., et al. (2016). Distinct features of H3K4me3 and H3K27me3 chromatin domains in pre-implantation embryos. *Nature* **537**, 558-562.

Maslov, A. Y., Lee, M., Gundry, M., Gravina, S., Strogonova, N., Tazearslan, C., Bendebury, A., Suh, Y. and Vijg, J. (2012). 5-aza-2'-deoxycytidine-induced genome rearrangements are mediated by DNMT1. *Oncogene* **31**, 5172-5179.

Oka, M., Meacham, A. M., Hamazaki, T., Rodic, N., Chang, L. J. and Terada, N. (2005). De novo DNA methyltransferases Dnmt3a and Dnmt3b primarily mediate the cytotoxic effect of 5-aza-2'-deoxycytidine. *Oncogene* **24**, 3091-3099.

Okano, M., Bell, D. W., Haber, D. A. and Li, E. (1999). DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell* **99**, 247-257.

Penny, G. D., Kay, G. F., Sheardown, S. A., Rastan, S. and Brockdorff, N. (1996). Requirement for Xist in X chromosome inactivation. *Nature* **379**, 131-137.

Pintacuda, G. and Cerase, A. (2015). X Inactivation Lessons from Differentiating Mouse Embryonic Stem Cells. *Stem Cell Rev Rep* **11**, 699-705.

Sado, T., Fenner, M. H., Tan, S. S., Tam, P., Shioda, T. and Li, E. (2000). X inactivation in the mouse embryo deficient for Dnmt1: distinct effect of hypomethylation on imprinted and random X inactivation. *Dev. Biol.* **225**, 294-303.

Sado, T., Okano, M., Li, E. and Sasaki, H. (2004). De novo DNA methylation is dispensable for the initiation and propagation of X chromosome inactivation.

Development **131**, 975-982.

Schulz, E. G. and Heard, E. (2013). Role and control of X chromosome dosage in
mammalian development. *Curr. Opin. Genet. Dev.* **23**, 109-115.

Smith, Z. D., Chan, M. M., Mikkelsen, T. S., Gu, H., Gnirke, A., Regev, A. and Meissner,
A. (2012a). A unique regulatory phase of DNA methylation in the early mammalian
embryo. *Nature* **484**, 339-344.

Smith, Z. D., Chan, M. M., Mikkelsen, T. S., Gu, H. C., Gnirke, A., Regev, A. and
Meissner, A. (2012b). A unique regulatory phase of DNA methylation in the early
mammalian embryo. *Nature* **484**, 339-344.

Smith, Z. D. and Meissner, A. (2013). DNA methylation: roles in mammalian development.
Nat. Rev. Genet. **14**, 204-220.

Smith, Z. D., Shi, J., Gu, H., Donaghey, J., Clement, K., Cacchiarelli, D., Gnirke, A.,
Michor, F. and Meissner, A. (2017). Epigenetic restriction of extraembryonic
lineages mirrors the somatic transition to cancer. *Nature* **549**, 543-547.

Soderberg, O., Gullberg, M., Jarvius, M., Ridderstrale, K., Leuchowius, K. J., Jarvius,
J., Wester, K., Hydbring, P., Bahram, F., Larsson, L. G., et al. (2006). Direct
observation of individual endogenous protein complexes in situ by proximity ligation.
Nat. Methods **3**, 995-1000.

Uysal, F., Ozturk, S. and Akkoyunlu, G. (2017). DNMT1, DNMT3A and DNMT3B
proteins are differently expressed in mouse oocytes and early embryos. *J. Mol. Histol.*
48, 417-426.

Vire, E., Brenner, C., Deplus, R., Blanchon, L., Fraga, M., Didelot, C., Morey, L., Van
Eynde, A., Bernard, D., Vanderwinden, J. M., et al. (2006). The Polycomb group
protein EZH2 directly controls DNA methylation. *Nature* **439**, 871-874.

Wu, J. Y., Huang, B., Chen, H., Yin, Q. Z., Liu, Y., Xiang, Y. L., Zhang, B. J., Liu, B. F.,
Wang, Q. J., Xia, W. K., et al. (2016). The landscape of accessible chromatin in
mammalian preimplantation embryos. *Nature* **534**, 652-+.

Wutz, A. (2011). Gene silencing in X-chromosome inactivation: advances in understanding
facultative heterochromatin formation. *Nat. Rev. Genet.* **12**, 542-553.

Yagi, M., Kabata, M., Tanaka, A., Ukai, T., Ohta, S., Nakabayashi, K., Shimizu, M.,
Hata, K., Meissner, A., Yamamoto, T., et al. (2020). Identification of distinct loci
for de novo DNA methylation by DNMT3A and DNMT3B during mammalian
development. *Nat Commun* **11**, 3199.

Zhang, J., Wang, Y., Liu, X., Jiang, S., Zhao, C., Shen, R., Guo, X., Ling, X. and Liu, C.
(2015). Expression and potential role of microRNA-29b in mouse early embryo
development. *Cell. Physiol. Biochem.* **35**, 1178-1187.

Zhao, J., Yao, K., Yu, H., Zhang, L., Xu, Y., Chen, L., Sun, Z., Zhu, Y., Zhang, C., Qian,
Y., et al. (2021). Metabolic remodelling during early mouse embryo development.
Nat Metab **3**, 1372-1384.

Zhu, P., Guo, H., Ren, Y., Hou, Y., Dong, J., Li, R., Lian, Y., Fan, X., Hu, B., Gao, Y., et
al. (2018). Single-cell DNA methylome sequencing of human preimplantation
embryos. *Nat. Genet.* **50**, 12-19.

Figure legends

Figure 1. A small proportion of promoters initiate *de novo* DNA methylation by 8-cell

stage. (A) The mRNA expression levels of *Dnmt3s* from the 2-cell to blastocyst stage. The

left ordinate represents FPKM values of *Dnmt3s* from published transcriptome data (Fan et al.,

2015), and the right ordinate represents relative expression levels detected by qRT–PCR. (B)

Immunofluorescent staining of DNMT3B in mouse preimplantation embryos. Scale bars: 50

µm. (C) DNA methylation changes of promoters during the transition from the 8-cell to

blastocyst stage. Left panel: heat map of DNA methylation levels at promoters. Right panel:

general trends of promoters undergoing *de novo* methylation or demethylation, and their

relative contribution to differentially methylated regions. (D-E) DNA methylation dynamics

of promoters subject to demethylation (D) or *de novo* methylation (E) during the transition

from the 8-cell to blastocyst stage (red and green lines, average methylation levels; colored

space, 10th/ 90th percentile). (F) A model illustrating two waves of *de novo* DNA methylation

before and after the blastocyst stage, and their potential functions.

Figure 2. Minor *de novo* DNA methylation participates in embryonic proliferation,

differentiation and affects developmental potential. (A) Schematic diagram of the

experimental design. (B) EdU staining of NC and *Dnmt3b*-KD blastocysts. Right panel: the

percentage of EdU-positive cells. (C) The total cell number (left panel) and the ratio of

ICM/TE (right panel) in NC and *Dnmt3b*-KD blastocysts. (D) The total ATP production (left

panel) and OXPHOS-dependent ATP production (right panel) in NC and *Dnmt3b*-KD

blastocysts. (E-F) Implantation rate (E) and live birth rate (F) in NC and *Dnmt3b*-KD groups.

All data are presented as the mean ± SD of at least three independent experiments. *$P < 0.05$,

**$P < 0.01$. ns, not significant. Scale bars: 50 µm (A, B).

Figure 3. Minor *de novo* methylation preferentially occurs on the X chromosome and

plays an important role in iXCI. (A) The chromosome-wide distribution of *de novo*

methylated promoters during the transition from the 8-cell to blastocyst stage. (B) Percentage

of *de novo* methylated promotors on each chromosome during the transition from the 8-cell to

blastocyst stage. (C) Schematic diagram of the experimental design and the molecular

processes of iXCI initiation and establishment. *Dnmt3b* knockdown was performed by

microinjecting *Dnmt3b* siRNA into zygotes. Immunostaining assay for H3K27me3 was

performed and quantified in trophoblast cells in each female blastocyst. (D) Representative immunostaining for H3K27me3 (red) in the nuclei (DAPI) of NC and *Dnmt3b*-KD female blastocysts colabeled with CDX2 (green)-positive trophoblast cells. Scale bars: 50 µm. The arrowhead indicates the H3k27me3 domain and the blank arrowhead indicates the blastomere without the H3k27me3 domain. (E) The ratio of trophoblast cells classified by the number of H3K27me3 domains in each NC and *Dnmt3b*-KD female blastocysts. Each bar represents one female embryo. (F) The percentage of H3K27me3-positive and H3K27me3-negative trophoblast cells (TCs) to total trophoblast cells in female blastocysts. All data are presented as the mean ± SD of at least three independent experiments. **P < 0.01. ns, not significant.

Figure 4. Minor *de novo* DNA methylation co-regulates iXCI via the interaction between DNMT3B and PRC2 core components. (A) Representative localization of *Xist* expression detected by RNA-FISH in the nuclei (DAPI) of NC and *Dnmt3b*-KD female morulae. (B) The ratio of blastomere with *Xist* signal to the total number of blastomeres in each NC and *Dnmt3b*-KD female morulae. Each bar represents one female embryo. The arrowhead indicates *Xist* RNA domain and the blank arrowhead indicates the blastomere without *Xist* RNA domain. (C) *Rnf12* and *Xist* expression levels in individual *Dnmt3b*-KD female 8-cell embryos and morulae. (D) Expression of H3K27me3 modifiers in individual *Dnmt3b*-KD female morulae. (E) Validation of the interaction between DNMT3B and PRC2 core components, *i.e.,* EZH2, SUZ12, EED, via *in situ* PLA. All data are presented as the mean ± SD of at least three independent experiments. **P < 0.01. ns, not significant. Scale bars: 50 µm (A, E). (F) A model illustrating a wave of minor *de novo* DNA methylation that initiates during the transition from the mouse 8-cell to blastocyst stage. Minor *de novo* DNA methylation co-regulates iXCI via the interaction between DNMT3B and PRC2 core components, and fine-tunes the processes of lineage commitment, proliferation and metabolic homeostasis before blastocyst formation.

Figure 1—figure supplement 1. (A) Dynamics of global methylation during preimplantation
embryos (blue line, average methylation levels; colored space, 10th/ 90th percentile). (B)
General trends of CpG islands undergoing *de novo* methylation or demethylation, and their
relative contribution to differentially methylated regions. (C) Number of promoters that gain
or loss DNA methylation in female and male embryos during the transition from the 8-cell to
blastocyst stage. (D) DNA methylation changes of promoters that gain DNA methylation
during the transition from the 8-cell to blastocyst stage in female or male embryos. (E) DNA
methylation dynamics of promoters with different CpG-densities during consecutive
transitions.

Figure 2—figure supplement 1. (A) A scatter plots of changes in promoter DNA methylation
and gene expression from the 8-cell embryos to ICM. Red plots indicate putative genes that
undergo minor *de novo* DNA methylation (ΔDNA methylation > 0.01) and are downregulated
(ΔDNA methylation < -0.01). (B-C) Gene ontology analysis (B) and phenotype annotation (C)
of genes transcriptionally regulated by minor *de novo* DNA methylation. Blue squares
indicate genes that are related to developmental or lethal phenotypes. (D) Gene ontology
analysis of genes transcriptionally regulated by DNA demethylation from the 8-cell embryos
to ICM.

Figure 2—figure supplement 2. (A) Relative expression levels of *Dnmt3b* in NC and
Dnmt3b-KD morulae and blastocysts. (B-C) Immunofluorescent staining (C) and
quantification (D) of DNMT3B in NC and *Dnmt3b*-KD morulae and blastocysts. (D-E)
Immunofluorescent staining (D) and quantification (E) of 5-mC in NC and *Dnmt3b*-KD
morulae and blastocysts. All data are presented as the mean ± SD of at least three independent
experiments. *$P < 0.05$, **$P < 0.01$. ns, not significant. Scale bars: 50 µm (B, D).

Figure 2—figure supplement 3. (A) Schematic diagram of the experimental design. (B)
EdU staining of blastocysts exposed to 5-aza-dC from the 8-cell to blastocyst stage. Right
panel: the percentage of EdU-positive cells. (C) The total cell number (left panel) and the
ratio of ICM/TE (right panel) of blastocysts treated with or without 5-aza-dC. (D) The total
ATP production (left panel) and OXPHOS-dependent ATP production (right panel) of
blastocysts treated with or without 5-aza-dC. All data are presented as the mean ± SD of at

least three independent experiments. *P < 0.05, **P < 0.01. ns, not significant. Scale bars: 50

µm (B, C).

Figure 3—figure supplement 1. (A) The distribution of *de novo* methylated and

demethylated promoters cross the X chromosome during the transition from the 8-cell to

blastocyst stage. (B) The chromosome-wide distribution of *de novo* methylated promoters

during the transition from the blastocyst to post-implantation stage. (C) Percentage of *de novo*

methylated promoters on each chromosome during the transition from the blastocyst to

post-implantation stage.

Figure 3—figure supplement 2. (A) DNA methylation changes of promoters that gain DNA

methylation during the transition from the 8-cell to blastocyst stage in chromosome X and

autosomes of female embryos. (B) DNA methylation changes of promoters that gain DNA

methylation during the transition from the 8-cell to blastocyst stage in chromosome X and

autosome of male embryos.

Figure 3—figure supplement 3. (A) Schematic diagram of the experimental design.

Dnmt3b-KO female embryos were collected by intercrossing *Dnmt3b*[+/-] male and female mice.

Immunostaining assay for H3K27me3 was performed and quantified in trophoblast cells in

each female homozygous blastocyst. (B) Representative immunostaining for H3K27me3 (red)

in the nuclei (DAPI) of wild-type and *Dnmt3b*-KO female blastocysts colabeled with CDX2

(green)-positive trophoblast cells. The arrowhead indicates the H3k27me3 domain and the

blank arrowhead indicates the blastomere without the H3k27me3 domain. (C) The ratio of

trophoblast cells classified by the number of H3K27me3 domains in each wild-type and

Dnmt3b-KO female blastocysts. Each bar represents one female embryo. (D) The percentage

of H3K27me3-positive and H3K27me3-negative trophoblast cells (TCs) to total trophoblast

cells in female blastocysts. (E) Schematic diagram of the experimental design. Zygotes were

cultured in vitro and transiently exposed to 5-aza-dC during the window of minor *de novo*

DNA methylation (the 8-cell to blastocyst stage) or not. Immunostaining assay for

H3K27me3 was performed and quantified in trophoblast cells in each female blastocyst. (F)

Representative immunostaining for H3K27me3 (red) in the nuclei (DAPI) of 5-aza-dC treated

or untreated female blastocysts colabeled with CDX2 (green)-positive trophoblast cells. Scale

bars: 50 µm. The arrowhead indicates the H3k27me3 domain and the blank arrowhead

indicates the blastomere without the H3k27me3 domain. (G) The ratio of trophoblast cells

classified by the number of H3K27me3 domains in female blastocysts exposed or not exposed

to 5-aza-dC from the 8-cell to blastocyst stage. Each bar represents one female embryo. (H)

The percentage of H3K27me3-positive and H3K27me3-negative trophoblast cells (TCs) to

total trophoblast cells in female blastocysts. (I) The changes of promoter methylation levels of

X-linked non-escaping genes from the 8-cell to blastocyst stage. (J) The expression level

changes of *de novo* methylated X-linked non-escaping genes during the transition from 8-cell

to blastocyst stage. All data are presented as the mean ± SD of at least three independent

experiments. *$P < 0.05$, **$P < 0.01$. ns, not significant. Scale bars: 50 µm (B, E).

Figure 3—figure supplement 4. (A) Genome browser views of the allelic expression, DNA

methylation and H3K27me3 enrichment cross the X chromosome in WT and DKO female

ExE based on the previously published data (Chen et al., 2019). (B) Genome browser views

of the paternal expression of selected X-linked non-escaping genes that undergo minor *de*

novo DNA methylation based on the previously published data (Chen et al., 2019).

Figure 4—figure supplement 1. (A) Representative localization of *Xist* expression detected

by RNA-FISH in the nuclei (DAPI) of wild-type and *Dnmt3b*-KO female morulae. The

arrowhead indicates *Xist* RNA domain and the blank arrowhead indicates the blastomere

without *Xist* RNA domain. Scale bars: 50 µm. (B) The ratio of blastomere with Xist signal to

the total number of blastomeres in each wild-type and *Dnmt3b*-KO female morulae. Each bar

represents one female embryo. All data are presented as the mean ± SD of at least three

independent experiments. ns, not significant.

Figure 4—figure supplement 2. (A) H3K27me3 enrichment cross the X chromosome from

the 8-cell to blastocyst stage based on the previously published data (Liu et al., 2016). (B)

Concurrent increase in H3K27me3 enrichment (left panel) and DNA methylation levels (right

panel) at selected X-linked promoters that undergo minor *de novo* DNA methylation.

Figure 4—figure supplement 3. (A) Schematic illustration of generation of *Dnmt3b*-KO

female ES cells via the CRISPR/Cas9n system. (B) Detection of deletion introduced by

sgRNA-Cas9n targeting *Dnmt3b* via PCR with genomic DNA from wild-type and

Dnmt3b-KO female ES cells. (C) Relative expression levels of *Dnmt3b* in wild-type and

Dnmt3b-KO female ES cells. (D) Schematic illustration of generation of DKO female ES

cells via the CRISPR/Cas9n system. (E) Detection of deletion introduced by sgRNA-Cas9n

targeting *Dnmt3a/3b* via PCR with genomic DNA from wild-type and DKO female ES cells.

(F) Relative expression levels of *Dnmt3a* and *Dnmt3b* in wild-type and DKO female ES cells.

(G) Representative immunostaining for H3K27me3 (red) in the nuclei (DAPI) of wild-type,

Dnmt3b-KO, and DKO female ES cells differentiated for 5, 7 or 9 days, as well as the ratio of

ES cells classified by the number of H3K27me3 domains. Scale bars: 20 µm. (H) Western

blot showing immunoprecipitation of DNMT3B and EZH2, SUZ12, EED from differentiated

female ES cells. All data are presented as the mean ± SD of at least three independent

experiments. **$P < 0.01$.

Supplemental Tables (separate files)

Supplemental Table S1. Primers sequences used in this study.

Figure 1

Figure 2

A

Microinjection of Control or *Dnmt3b* siRNA

Blastocyst transfer

E7.5

E19

• Proliferation assay
• Total cell number
• ICM/TE ratio
• ATP production

• Implantation rate

• Live birth rate

B

DAPI   EdU

NC

Dnmt3b-KD

% of EdU-positive cells

**

NC   *Dnmt3b*-KD

C

DAPI / CDX2

NC

Dnmt3b-KD

Total cell number

ns

NC   *Dnmt3b*-KD

Ratio of ICM / TE

*

NC   *Dnmt3b*-KD

D

Total

ATP production / embryo

ns

NC   *Dnmt3b*-KD

OXPHOS

ATP production / embryo

**

NC   *Dnmt3b*-KD

E

Implantation rate (%)

*

N = 60   N = 72

NC   *Dnmt3b*-KD

F

Live birth rate (%)

*

N = 60   N = 60

NC   *Dnmt3b*-KD

Figure 3

Figure 4

Figure 1—figure supplement 1

Figure 2—figure supplement 1

A

- Δ DNA methylation>0.01 & mRNA level FC<0.5
- Δ DNA methylation<-0.01 & mRNA level FC>2

1093

ΔDNA Methylation

$Log_{10}(FPKM\ FC)$

B

-log10 (P value)

regulation of metabolic process
cellular component organization
cell differentiation
multicellular organism development
system development
transport
regulation of developmental process
cell fate commitment

D

-log10 (P value)

cell-substrate adhesion
glycoprotein metabolic process
embryo development
regulation of developmental process
transport
cell differentiation
system development
cellular component organization

C

	cell differentiation							cellular metabolism						
	Shh	*Pax6*	*Cxcl12*	*Arih2*	*Eng*	*Adm*	*Pnpla1*	*Wtap*	*Rnf7*	*Tshz2*	*Sumo2*	*Cdan1*	*Atp7a*	*Otub1*
abnormal embryo development	■							■						
decreased embryo size	■	■			■		■				■	■	■	
embryonic growth retardation	■			■	■				■		■	■		
abnormal placenta morphology		■			■	■								
embryonic lethality	■				■				■	■	■	■		
embryonic tissue cell apoptosis									■					
prenatal lethality	■	■	■		■						■	■		
preweaning lethality	■		■	■		■	■					■	■	
postnatal growth retardation	■	■	■					■			■			

Figure 2—figure supplement 2

Figure 2—figure supplement 3

Figure 3—figure supplement 1

A

De novo methylated promoters Demethylated promoters

8-cell→ICM

Δ DNA methylation

ChrX (M bp)

B

ICM→E6.5

Number of promoters

Gain Loss

C

% of methylated promoters compared to all promoters

ChrX 9% 91% Chr10 25% 75% Chr17 27% 73% Chr19 27% 73%

Gain Loss

Figure 3—figure supplement 2

A

B

Figure 3—figure supplement 3

Figure 3—figure supplement 4

A

B

Pat | WT / DKO — *Chic1*

Pat | WT / DKO — *Atp7a*

Pat | WT / DKO — *Kif4*

Figure 4—figure supplement 1

A

B

Figure 4—figure supplement 2