Cell autonomous role of leucine-rich repeat kinase in protection of dopaminergic neuron survival

Jongkyun Kang[1,¶], Guodong Huang[1,¶], Long Ma[1], Youren Tong[1], Phoenix Chen[1], and Jie Shen[1,2*]

[1]Department of Neurology, Brigham and Women's Hospital, [2]Program in Neuroscience, Harvard Medical School, Boston, MA 02445, United States of America

*Corresponding author

E-mail: jshen@bwh.harvard.edu

[¶]These authors contributed equally to this work.

Abstract

Mutations in leucine-rich repeat kinase 2 (LRRK2) are the most common genetic cause of Parkinson's disease (PD), which is the leading neurodegenerative movement disorder characterized by the progressive loss of dopaminergic (DA) neurons in the substantia nigra pars compacta (SNpc). However, whether LRRK2 mutations cause PD and degeneration of DA neurons *via* a toxic gain-of-function or a loss-of-function mechanism is unresolved and has pivotal implications in LRRK2 based PD therapy. In this study, we investigate whether LRRK2 and its functional homologue LRRK1 play an essential, intrinsic role in DA neuron survival through the development of DA neuron-specific *LRRK* conditional double knockout (cDKO) mice. We first generated and characterized floxed *LRRK1* and *LRRK2* mice and then confirmed that germline deletions of the floxed *LRRK1* and *LRRK2* alleles result in null alleles, as evidenced by the absence of *LRRK1* and *LRRK2* mRNA and protein in the respective homozygous deleted mutant mice. We further examined the specificity of Cre-mediated recombination driven by the *dopamine transporter-Cre* (*DAT-Cre*) knockin (KI) allele using a GFP reporter line and confirmed that *DAT-Cre*-mediated recombination is restricted to DA neurons in the SNpc. Crossing these validated floxed *LRRK1* and *LRRK2* mice with *DAT-Cre* KI mice, we then generated DA neuron-restricted *LRRK* cDKO mice and further showed reduced levels of LRRK1 and LRRK2 in dissected ventral midbrains of *LRRK* cDKO mice. While DA neuron-restricted *LRRK* cDKO mice of both sexes exhibit normal mortality and body weight, they develop age-dependent loss of DA neurons in the SNpc, as demonstrated by the progressive reduction of DA neurons in the SNpc of cDKO mice at 20 and 24 months of age. Moreover, DA neurodegeneration is accompanied with increases of apoptosis and elevated microgliosis in the SNpc of *LRRK* cDKO mice. These findings provide unequivocal evidence for the importance of LRRK in DA neurons and raise the possibility that LRRK2 mutations may impair its protection of DA neurons, leading to the loss of DA neurons in Parkinson's disease.

Introduction

Parkinson's disease (PD) is the most common movement disorder and is characterized by the progressive loss of dopaminergic (DA) neurons in the substantia nigra pars compacta (SNpc). Dominantly inherited missense mutations in the *leucine-rich repeat kinase 2* (*LRRK2*) gene are the most common cause of both familial and sporadic PD, highlighting the importance of LRRK2 in PD pathogenesis (1-16). LRRK2 is a large protein of 2,527 amino acid residues containing several functional domains including several leucine-rich repeats (LRRs), a GTPase-like domain of Ras-of-complex (Roc), a C-terminal of Roc (COR) domain, and a serine/threonine MAPKKK-like kinase domain. LRRK1, a homologue of LRRK2, belongs to the evolutionarily conserved Roco protein family and contains similar LRRs, Roc, COR, and kinase domains (17-19). LRRK proteins are broadly expressed with LRRK2 being most abundant in the kidney (20). While most PD mutations are found in LRRK2, rare variants in the Roc, COR, and kinase domains of LRRK1 have been reported and may be associated with PD (21). Furthermore, the Roc-COR domain of LRRK2 forms dimers and exhibits conventional Ras-like GTPase properties, and the R1441/C/G/H and I1371V mutations destabilize dimer formation and decrease GTPase activity (22, 23). Recent high-resolution cryoEM structural studies of full-length LRRK2 demonstrated its existence as dimers and pathogenic mutations such as R1441/C/G/H and Y1699I at the Roc-COR interface, whereas the G2910S mutant and wild-type LRRK2 share similar structure (24).

Previous genetic studies demonstrated that LRRK2 plays essential roles in the autophagy-lysosomal pathway (25-28). Consistent with abundant expression of LRRK2 in kidneys, *LRRK2*-/- mice develop age-dependent phenotypes in the kidney, including autophagy-lysosomal impairments and increases of α-synuclein, apoptosis, and inflammatory responses (25, 26, 29). It has also been reported that LRRK2 is important for maintaining lung homeostasis, and *LRRK2* deficiency results in impaired autophagy in alveolar type II epithelial cells (28). It was proposed that the lack of brain phenotypes in *LRRK2*-/- mice might be due to the presence of LRRK1 and potential functional compensation mechanisms (25, 26). Indeed, *LRRK* double knockout (DKO) mice develop age-dependent, progressive loss of DA neurons in the SNpc, beginning at 14 months of age (30). However, *LRRK* DKO mice also

exhibit lower body weight and increased mortality, raising the possibility that DA neurodegeneration in

LRRK DKO mice may be secondary to poor health.

In this study, we investigate whether LRRK2 and its functional homologue LRRK1 play an

essential, intrinsic role in DA neuron survival through the development of DA neuron-specific *LRRK*

conditional DKO (cDKO) mice. We first generated and confirmed floxed *LRRK1* and *LRRK2* mice, and

then crossed them with the deleter *CMV-Cre* to create germline deletions of the floxed *LRRK1* and

LRRK2 alleles, followed by Northern and Western analyses to confirm the absence of *LRRK1* and *LRRK2*

mRNA and protein in the respective homozygous deleted mutant mice. We also crossed a GFP reporter

mouse with *DAT-Cre* KI mice and found that Cre-mediated recombination occurs in most, if not all, DA

neurons and is restricted to DA neurons in the SNpc. We then crossed these thoroughly validated floxed

LRRK1 and *LRRK2* mice with *DAT-Cre* KI mice to generate DA neuron-restricted *LRRK* cDKO mice and

further confirmed the reduction of LRRK1 and LRRK2 in dissected ventral midbrains of *LRRK* cDKO

mice. While DA neuron-restricted *LRRK* cDKO mice of both sexes exhibit normal mortality and body

weight, they develop age-dependent loss of DA neurons in the SNpc, as demonstrated by the progressive

reduction of TH+ DA neurons or NeuN+ neurons in the SNpc of cDKO mice at 20 and 24 months of age.

Interestingly, quantitative electron microscopy analysis showed a similar number of electron-dense

vacuoles in SNpc neurons of *LRRK* cDKO mice relative to controls, in contrast to age-dependent,

dramatic increases of vacuoles in SNpc neurons of germline *LRRK* DKO mice (30, 31). Moreover, DA

neurodegeneration is accompanied with increases of apoptosis and elevated microgliosis in the SNpc of

LRRK cDKO mice. These findings provide unequivocal evidence for the importance of LRRK in DA

neurons and raise the possibility that LRRK2 mutations may impair this crucial physiological function,

leading to the loss of DA neurons in Parkinson's disease.

Results

Generation and molecular characterization of the floxed and deleted *LRRK1* and *LRRK2* alleles

LRRK2 and its homologue LRRK1 share several functional (LRRs, GTPase Roc, COR, and

kinase) domains (Figure 1A). To investigate the intrinsic role of LRRK in DA neurons, we generated

floxed *LRRK1* (*fLRRK1*) and floxed *LRRK2* (*fLRRK2*) mice, which permit deletions of *LRRK1* and

LRRK2 selectively in DA neurons by *DAT-Cre*, which expresses Cre recombinase under the control of the

endogenous *DAT* promoter (32). We introduced two *loxP* sites in introns 26 and 29 of *LRRK1* through

homologous recombination and site-specific recombination by FLP recombinase to remove the positive

selection *PGK-neo* cassette, which is flanked by two *FRT* sites (Figure 1B, C; Figure 1-figure supplement

1, 2). The embryonic stem (ES) cells carrying the targeted allele or the floxed allele were identified and

validated by Southern analysis using the 5' and 3' external probes as well as the *neo* probe (Figure 1D;

Figure 1-figure supplement 3). The validated ES cells carrying the floxed *LRRK1* allele were injected into

mouse blastocysts to develop floxed *LRRK1* mice, which were further confirmed by Southern using the 5'

and 3' external probes (Figure 1D). In the presence of Cre recombinase, the floxed *LRRK1* region

containing part of intron 26 (1,288 bp), exons 27-29, which encode the kinase domain, and part of intron

29 (1,023 bp) is deleted (Figure 1C), and the removal of exons 27-29 (625 bp) results in a frameshift of

downstream codons.

The *LRRK2* targeting vector contains the 5' homologous region, a *loxP* site 1,768 bp upstream of

the transcription initiation site, exons 1-2, the *PGK-neo* cassette flanked by two *loxP* sites and two *FRT*

sequences, and the 3' homologous region (Figure 1B, E; Figure 1-figure supplement 4, 5). The ES cell

clones carrying the targeted allele were identified and validated by Southern analysis using the 5' external

probe (Figure 1F; Figure 1-figure supplement 6) and genomic PCR followed by sequencing. Mice

carrying the *LRRK2* targeted allele were further verified by Southern analysis using the 5' and 3' external

probes (Figure 1-figure supplement 6) as well as the *neo* probe and were then crossed with the *Actin-FLP*

mice (33) to remove the *PGK-neo* cassette flanked by two *FRT* sites to generate floxed *LRRK2* mice

(Figure 1E). The resulting floxed *LRRK2* mice were confirmed by Southern analysis using the 5' and 3'

external probes (Figure 1F). In the presence of Cre recombinase, the floxed *LRRK2* region containing the

promoter and exons 1-2 are deleted, likely resulting in a null allele, as we previously targeted a very

similar region (~2.5 kb upstream of the transcription initiation site and exons 1-2) to generate germline

deletion of *LRRK2* (25).

Dopaminergic neurons in the SNpc are a small neuronal population embedded in the ventral

midbrain, making it difficult to confirm whether DA neuron-specific deletions of the floxed *LRRK1* and

LRRK2 regions result in null alleles. We previously generated three independent *LRRK1* knockout (KO)

mice, and only one KO line (Line 2) represents a *LRRK1* null allele (30), whereas deletion of *LRRK2*

promoter region and exons 1-2 resulted in a *LRRK2* null allele (25). We therefore generated germline

deleted *LRRK1* and *LRRK2* mice from floxed *LRRK1* and *LRRK2* mice, respectively, by crossing them to

germline deleter *CMV-Cre* transgenic mice (34).

We then performed Northern analysis of *LRRK1* using both an upstream probe specific for exons

2-3 and a downstream probe specific for exons 27-29 (Figure 1G; Figure 1-figure supplement 7, 8).

Because of the low expression level of *LRRK1* and the relative abundance of *LRRK1* in the lung (20), we

enriched polyA+ RNA from the lung of the mice carrying homozygous germline deleted (Δ/Δ) *LRRK1*

alleles derived from the floxed *LRRK1* alleles. Using a probe specific for exons 2-3, the *LRRK1*

transcripts in wild-type mice are the expected size of ~7.4 kb, whereas the detected *LRRK1* transcripts in

LRRK1 Δ/Δ mice are smaller, consistent with the deletion of exons 27-29 (625 bp), resulting in a

frameshift of downstream codons and the likely degradation of the truncated *LRRK1* mRNA (Figure. 1G;

Figure 1-figure supplement 8). Using a probe specific for exons 27-29, there is no *LRRK1* transcript in

LRRK1 Δ/Δ mice (Figure. 1G; Figure 1-figure supplement 8). Extensive RT-PCR analysis of total RNA

isolated from the kidney, brain, and lung of *LRRK1* Δ/Δ mice using an exon 32-specific primer for RT

and exon-specific primer sets for PCR (*e.g.* exons 4-8, 11-17, 20-25, and 25-31), followed by sequencing

confirmation of the PCR products, indicated normal *LRRK1* splicing in *fLRRK1/fLRRK1* mice and the

lack of *LRRK1* exons 27-29 in *LRRK1* Δ/Δ mice (Figure 1-figure supplement 9).

Similarly, Northern analysis of *LRRK2* using a probe specific for exons 1-5 and RT-PCR

followed by sequencing confirmed the absence of *LRRK2* mRNA in *LRRK2* Δ/Δ brains and normal

LRRK2 transcripts in *fLRRK2/fLRRK2* brains (Figure 1H; Figure 1-figure supplement 10, 11).

Furthermore, Western analysis confirmed the absence of LRRK1 and LRRK2 proteins in the brain of

LRRK1 Δ/Δ and *LRRK2* Δ/Δ mice, respectively (Fig. 1I). Taken together, our Northern, RT-PCR followed

by sequencing, and Western analyses demonstrated that deletion of the floxed *LRRK1* and *LRRK2* alleles

results in null mutations. Thus, floxed *LRRK1* and *LRRK2* alleles can be used to generate DA neuron-

specific *LRRK* cDKO mice.

Generation and molecular characterization of DA neuron-specific *LRRK* cDKO mice

To generate DA neuron-specific *LRRK* cDKO mice, we used *DAT-Cre* KI mice, which express

Cre recombinase under the control of the endogenous *DAT* promoter (32). To confirm if Cre-mediated

recombination occurs broadly and specifically in DA neurons of the SNpc, we crossed *DAT-Cre* mice

with the *Rosa26-CAG-LSL-ZsGreen1* reporter mouse (35). Upon Cre expression, Cre recombinase

removes the floxed "stop" cassette, resulting in the expression of EGFP. We found that Cre-mediated

recombination (GFP+) occurs in TH+ cells in the SNpc (Figure 2A). Quantification of GFP+ and/or TH+

cells in the SNpc showed that 99% of TH+ dopaminergic neurons are also GFP+, demonstrating that Cre-

mediated recombination takes place in essentially all DA neurons in the SNpc (Figure 2B).

Having confirmed *fLRRK1* and *fLRRK2* mice as well as *DAT-Cre* mice, we then bred them

together to generate *LRRK* cDKO mice (*fLRRK1/fLRRK1; fLRRK2/fLRRK2; DAT-Cre/+*), which were

further bred with *fLRRK1/fLRRK1; fLRRK2/fLRRK2* mice to generate cDKO and littermate controls

(*fLRRK1/fLRRK1; fLRRK2/fLRRK2*). It was previously reported that germline *LRRK* DKO mice failed to

gain weight as they aged (30). However, DA neuron-specific *LRRK* cDKO and littermate control mice

have similar body and brain weights at the ages of 2-24 months (Figure 2C, D). Western analysis showed

a significant reduction of LRRK1 and LRRK2 proteins in the dissected ventral midbrain but not in the

cerebral cortex of DA neuron-specific *LRRK* cDKO, relative to littermate controls at 2-3 months of age

(Figure 2E, F).

Age-dependent loss of DA neurons in the SNpc of *LRRK* cDKO mice

To determine whether the inactivation of LRRK selectively in DA neurons of the SNpc impacts

their survival, we performed TH immunostaining and quantified TH+ DA neurons in the SNpc of *LRRK*

cDKO mice and littermate controls. The morphology of TH+ DA neurons in *LRRK* cDKO mice at the

ages of 15, 20, and 24 months appears normal (Figure 3A). TH+ neuron counting revealed that the

number of DA neurons in the SNpc at the age of 15 months is similar between cDKO mice (10,000 ± 141)

and littermate controls (10,077 ± 310, $F_{1,46}$ = 16.59, p = 0.0002; p > 0.9999, two-way ANOVA with

Bonferroni's post hoc multiple comparisons; Figure 3B). However, at the age of 20 months, the number

of TH+ neurons in the SNpc of cDKO mice (8,948 ± 273) is significantly reduced, compared to controls

(10,244 ± 220; p = 0.0041), and is further decreased at the age of 24 months (Control: 9,675 ± 232, cDKO:

8,188 ± 452; p = 0.0010).

We further performed TH and NeuN double immunostaining of *LRRK* cDKO and control mice at

24 months of age (Figure 3C), and found that the number of NeuN+ neurons is also significantly reduced

in the SNpc of cDKO mice (17,923 ± 813), compared to controls (21,907 ± 469, p = 0.0006, Student's *t*-

test; Figure 3D). The number of TH+/NeuN+ cells in the SNpc of *LRRK* cDKO mice (10,500 ± 644) is

also lower, compared to control mice (14,102 ± 310, p = 0.0001; Figure 3D). These data indicate that the

decreases in the number of TH+ cells are not due to decreases in TH expression in DA neurons, but rather

a result of the loss of DA neuron cell bodies in the SNpc of *LRRK* cDKO mice.

We further evaluated apoptosis in the SNpc of *LRRK* cDKO and littermate controls at the age of

24 months using an antibody specific for active Caspase-3 to label apoptotic cells. We observed increases

in apoptotic DA neurons, labeled by active Caspase-3+ and TH+ immunoreactivity, in the SNpc of *LRRK*

cDKO mice (Figure 4A). Quantification of active Caspase-3+/TH+ apoptotic DA neurons shows a

significant increase in the SNpc of *LRRK* cDKO mice (323 ± 38), compared to controls (157 ± 8, p =

0.0004, Student's *t*-test; Figure 4B). These results further support that LRRK plays an intrinsic role in the

survival of DA neurons in the SNpc during aging.

Previously, we reported the reduction of noradrenergic neurons in the locus coeruleus (LC) of

LRRK DKO mice (30). We therefore quantified TH+ noradrenergic neurons in the LC of *LRRK* cDKO

mice at the age of 24 months, and found a similar number of TH+ cells in the LC of *LRRK* cDKO and

littermate controls (Control: 3,419 ± 86, cDKO: 3,350 ± 99, p = 0.6057, Student's *t*-test; Figure 5A, B).

Further examination of the GFP reporter line crossed with *DAT-Cre* showed the lack of Cre-mediated

recombination in the LC (Figure 5C).

203 Quantitative EM analysis of the SNpc in *LRRK* cDKO mice

We previously reported striking age-dependent increases of electron-dense autophagic and

autolysosomal vacuoles as well as the presence of large lipofuscin granules in the surviving SNpc neurons

of *LRRK* DKO mice beginning at 10 months of age (30, 31). To determine whether selective inactivation

of LRRK in DA neurons similarly results in the accumulation of electron-dense autophagic and

autolysosomal vacuoles in the SNpc, we performed EM analysis in the SNpc of *LRRK* cDKO mice and

littermate controls at the ages of 25 months (Figure 6). We observed various electron-dense double

membrane autophagosomes and single membrane autolysosomes as well as lipofuscin granules composed

of lipid-containing residues of lysosomal digestion in the SNpc of *LRRK* cDKO and littermate control

mice (Figure 6C-F). Interestingly, there are similar numbers of electron-dense vacuoles in the SNpc of

LRRK cDKO mice and littermate controls at the age of 25 months (Control: 6.72 ± 0.43, cDKO: 6.99 ±

0.52, p = 0.6839, Student's *t*-test; Figure 6G). We also found no significant difference in the area of

electron-dense vacuoles in the SNpc between *LRRK* cDKO and littermate controls at the age of 25 months

(Control: 4.43 ± 0.44 μm^2; cDKO: 4.60 ± 0.49 μm^2, p = 0.8048; Figure 6G). The difference in age-

dependent accumulation of electron-dense vacuoles in the SNpc between germline DKO mice and DA

neuron-restricted cDKO suggests that LRRK in non-DA neurons such as microglia, which is lacking in

germline DKO but is expressed normally in cDKO mice, may play a more prominent role.

220 Enhanced microgliosis in the SNpc of *LRRK* cDKO mice

We further evaluated whether elevated microgliosis accompanies DA neurodegeneration in the

SNpc of *LRRK* cDKO mice. We performed immunohistochemical analysis of Iba1, which labels

microglia, and TH, which marks DA neurons and processes, thus showing the boundary of the SNpc

(Figure 7A-C). We found that the number of Iba1+ immunoreactive microglia is significantly increased in

the SNpc of *LRRK* cDKO mice at 15 months of age (2541.3 ± 193.0), compared to controls (1736.7 ±

82.9; p = 0.0017, two-way ANOVA with Bonferroni's post hoc multiple comparisons; Figure 7A, D).

The number of Iba1+ immunoreactive microglia in the SNpc of *LRRK* cDKO is further increased

compared to controls at the age of 20 (Control: 2426.0 ± 67.7, cDKO: 3639.0 ± 127.1, p < 0.0001; Figure

7B, D) and 25 months (Control: 2640.0 ± 187.1, cDKO: 4088.8 ± 100.3, p < 0.0001; Figure 7C, D).

These results show that despite the selective inactivation of LRRK in DA neurons of *LRRK* cDKO mice,

microgliosis accompanies DA neuronal loss in the SNpc.

Discussion

LRRK2 mutations represent ~5% familial and 1-2% of sporadic PD cases, and the G2019S

mutation is the most common mutation but with a higher age of onset and incomplete penetrance (74% at

age 79) (36). Despite the importance of LRRK2 in PD pathogenesis, whether LRRK2 mutations cause PD

via a loss-of-function or a gain-of-function mechanism remains highly debated and unresolved. Neither

LRRK2-deficient mice nor *LRRK2* R1441C and G2019S KI mice develop dopaminergic

neurodegeneration (25, 37, 38). It was proposed that the lack of brain phenotypes in *LRRK2*-/- mice might

be due to the presence of LRRK1 (25, 26), which is broadly expressed in the brain including the midbrain

(https://www.proteinatlas.org/ENSG00000154237-LRRK1/brain). Indeed, germline deletions of LRRK2

and LRRK1 result in age-dependent loss of DA neurons and increases of apoptosis and microgliosis in the

SNpc of *LRRK* DKO mice (30, 31). However, the pleiotropic roles of LRRK1 and LRRK2 in the kidney

(25, 26, 29), lung (28), and bone (39, 40), and the observed earlier mortality and lower body weight in

LRRK DKO mice raised the possibility that DA neurodegeneration in *LRRK* DKO mice may be due to

poor health.

In this study, we address this important question through the development of floxed *LRRK1* and

LRRK2 mice (**Figure 1; Figure 1-figure supplement 1-11**), which permit the dissection of cell intrinsic

function of LRRK in DA neurons. We generated DA neuron-specific *LRRK* cDKO mice using the *DAT-*

Cre KI allele (32) to delete floxed *LRRK1* and *LRRK2* regions selectively in DA neurons. Indeed, using an EGFP reporter line, we found that *DAT-Cre* drives Cre-mediated recombination in almost all DA neurons in the SNpc (**Figure 2**). Unlike the *LRRK* germline DKO mice (30), DA neuron-specific *LRRK* cDKO mice exhibit normal body weight and mortality during mouse lifespan (**Figure 2**). Importantly, *LRRK* cDKO mice develop age-dependent DA neurodegeneration, as evidenced by progressive reductions of DA neurons in the SNpc at the age of 20-24 months (**Figure 3**) and increases of apoptotic DA neurons (**Figure 4**), whereas the number of noradrenergic neurons in the LC is uncaged, consistent with the lack of Cre-mediated recombination in the LC by *DAT-Cre* (**Figure 5**). Interestingly, quantitative EM analysis showed similar numbers of electron-dense vacuoles in the SNpc of *LRRK* cDKO and control mice even at 25 months of age (**Figure 6**), in contrast to dramatic, age-dependent increases of vacuoles in the SNpc of *LRRK* DKO mice (30, 31). Lastly, DA neurodegeneration in the SNpc of *LRRK* cDKO mice is accompanied with elevated microgliosis (**Figure 7**).

The toxic gain-of-function pathogenic mechanism was supported by earlier biochemical reports showing that LRRK2 mutations (*e.g.* R1441C/G, G2019S) enhance kinase activity, leading to elevated levels of pSer1292-LRRK2 (41), pSer935- LRRK2 (42, 43), pT73-Rab10, and pS106-Rab12 (44). However, most transgenic mice overexpressing mutant LRRK2 did not produce neurodegeneration or DA neuron loss (45-53), suggesting that enhanced kinase activity does not correlate with compromised DA neuronal survival, though two reports showed that overexpression of LRRK2 G2019S but not R1441C resulted in DA neurodegeneration (54, 55). Thus, further studies are needed to elucidate the differences of these seemingly conflicting transgenic findings and to understand the differences between the R1441C and G2019S mutations.

The loss-of-function pathogenic mechanism was supported by mouse knockout findings showing that germline inactivation of LRRK results in age-dependent loss of DA neurons in the SNpc and DA terminals in the striatum (30, 31). The current study demonstrated a cell intrinsic role of LRRK in support of DA neuron survival in the SNpc of the aging brain, providing further genetic evidence of an essential, cell autonomous requirement of LRRK in DA neuron survival. However, highly penetrant PD-linked

mutations are dominantly-inherited missense mutations (6) in LRRK2, and there have been no reported deletion mutations that are linked to or causal of familial PD, despite various variants were reported in LRRK2 and LRRK1 that are associated with sporadic PD (but not linked to familial PD). One plausible explanation is that LRRK2 mutant may act in a dominant negative manner, inhibiting the activity of the wild-type LRRK2. This notion is supported by structural studies showing that the GTPase domain forms a dimer, and that the R1441/C/G/H and I1371V mutations destabilize dimer formation and decrease GTPase activity (22). LRRK2 may function as a dimer, which is further supported by the crystal structure of LRRK2's WD40 domain dimer at 2.6 Å resolution (56). Additional studies also suggested that LRRK2 functions as a dimer (57-59), and that LRRK2 and LRRK1 can form a heterodimer (60). Furthermore, LRRK2 G2385R variant, a risk factor for PD (61-64), has been reported as a partial loss-of-function mutation (65, 66). Future studies are needed to differentiate the possibility between a toxic gain-of-function pathogenic mechanism vs. loss-of-function mutations exerting a dominant negative effect further impairing LRRK2 activity of the wild-type protein encoded by the LRRK2 wild-type allele, as the gain-of-function or loss-of-function pathogenic mechanism underlying LRRK2 mutations pivots opposing LRRK2 based therapeutic development of PD therapies.

Finally, it was surprising to find no accumulation of autophagic and autolysosomal vacuoles in SNpc neurons of *LRRK* cDKO mice, even at the age of 25 months. This suggests that non-dopaminergic cells (*e.g.* microglia), in which LRRK expression is unaffected in cDKO mice but is absent in germline DKO mice, may contribute to the dramatic phenotypic differences. Development of cell type-specific cDKO mice, such as microglia-specific *LRRK* cDKO mice, would permit further dissection of cell autonomous and non-cell autonomous role of LRRK in dopaminergic neuronal survival and whether glial LRRK plays a more important role in the regulation of the autophagy-lysosomal pathway.

Materials and methods

Key Resource Table

Antibodies				
Reagent Name	**Source**	**Catalog #**	**Identifiers**	**Titer**
Rabbit anti-LRRK1	Alomone Lab	ANR-101	AB_2756700	WB: 1:1000
Rabbit anti-LRRK2	abcam	Ab133474	AB_2713963	WB: 1:1000
Mouse anti-α-Vinculin	Millipore	05-386	AB_309711	WB: 1:2000
Rabbit anti-GFP	abcam	ab290	AB_303395	IF: 1:1000
Mouse anti-TH	SantaCruz	Sc-25269	AB_628422	IF: 1:50
Rabbit anti-TH	abcam	Ab112	AB_297840	IHC: 1:750
Rabbit anti-NeuN	Cell Signaling Technology	12943S	AB_2630395	IF: 1:400
Rabbit anti-cleaved caspase-3	Cell Signaling Technology	9661	AB_2341188	IF: 1:150
Rabbit anti-Iba1	Wako	019-19741	AB_839504	IF: 1:500
Goat anti-rabbit IRdye800	LI-COR Biosciences	926-32211	AB_2651127	WB: 1:20000
Goat anti-mouse IRdye680	LI-COR Biosciences	926-68020	AB_2651128	WB: 1:20000
Alexa Fluor 488 goat anti-mouse IgG	ThermoFisher Scientific	A-11001	AB_2534069	IF: 1:250
Alexa Fluor 555 goat anti-rabbit IgG	ThermoFisher Scientific	A-32732	AB_2633281	IF: 1:250
Alexa Fluor 488 goat anti-rabbit IgG	ThermoFisher Scientific	A-11034	AB_2576217	IF: 1:250
Alexa Fluor 555 goat anti-mouse IgG	ThermoFisher Scientific	A-21424	AB_141780	IF: 1:250
Goat Biotinylated anti-Rabbit IgG (H&L)	Vector Laboratories	BA-1000	AB_2313606	IHC: 1:250
Oligonucleotides				
LRRK1 PCR Forward Primer for Left arm amplification: P1 5'-GACAT<u>CCGCGG</u>CACCATGTGAGTGGCAGCTGTGGTGAGAAC-3' *Sac*II sequence is underlined.				
LRRK1 PCR Reverse Primer for Left arm amplification: P2 5'-GACAT<u>GCGGCCGC</u><u>AAGCTT</u>TTTAATAGCCGTTCTTTCTTAGAGAAGGCAG-3' *Not*I and *Hind*III sequences are underlined.				
LRRK1 PCR Forward Primer for Left-Middle arm amplification: P3 5'-CCAGTCACTTCTCCACCTCAGGGAAAATGG-3'				
LRRK1 PCR Reverse Primer for Left-Middle arm amplification: P4 5'-CCTTGT<u>GGTACC</u>CGGACCTTCTATCACCTTTATCC-3' *Kpn*I sequence is underlined.				
LRRK1 PCR Forward Primer for Right-Middle arm amplification: P5 5'-GGTCCG<u>GGTACC</u>ACAAGGTGCTGGTTAAGTGCC-3' *Kpn*I sequence is underlined.				

LRRK1 PCR Reverse Primer for Right-Middle arm amplification: P6 5'-AGCAGACCTCTTGCCTTCTACTACTGACTG-3'
LRRK1 PCR Forward Primer for Right arm amplification: P7 5'- GACAT<u>GTCGAC</u>GGATCCGTAGGGAAGACCCACTAGGAGGAAGAAAG-3' *Sal*I sequence is underlined.
LRRK1 PCR Reverse Primer for Right arm amplification: P8 5'- GACAT<u>AAGCTT</u>TGGTACCTTTCTAAAGGCAGCATTTTGCTTGC-3' *Hind*III sequence is underlined.
LRRK1 PCR Forward Primer for 5' Southern Probe: P15 5'-CAGAATACCCCATGCTGGGGAATTGC-3'
LRRK1 PCR Reverse Primer for 5' Southern Probe: P16 5'-CCGTTTCTAGGATTCTAATTTTTC-3'
LRRK1 PCR Forward Primer for 3' Southern Probe: P17 5'-AACTCCTTCCTGGTGCTGGCAGGCCTGGCTG-3'
LRRK1 PCR Reverse Primer for 3' Southern Probe: P18 5'-ACAAGTGACGTGACCATGGACGGAGCTGCG-3'
Neo PCR Forward Primer for Southern Probe: P23 5'-ATTCGGCTATGACTGGGCAC-3'
Neo PCR Reverse Primer for Southern Probe: P24 5'-GACCACCAAGCGAAACATCG-3'
LRRK1 PCR Forward Primer for exons 2-3 Northern Probe: P31 5'-CAGGATGAGCGTGTGTCTGCAG-3'
LRRK1 PCR Reverse Primer for exons 2-3 Northern Probe: P32 5'-CCTTCTCCTGTGAGGATTCGCTCT-3'
LRRK1 PCR Forward Primer for exons 27-29 Northern Probe: P33 5'-CTGGCCTACCTGCACAAGAA-3'
LRRK1 PCR Reverse Primer for exons 27-29 Northern Probe: P34 5'-CCTTCCCATCCCAGAACACC-3'
GADPH PCR Forward Primer for Northern Probe: P37 5'-ACCACAGTCCATGCCATCAC-3'
GADPH PCR Reverse Primer for Northern Probe: P38 5'-TCCACCACCCTGTTGCTGTA-3'
LRRK1 RT Primer in exons 32: P47 5'-GGCTCAGGTCATGCTCAGTT-3'
LRRK1 PCR Forward Primer for exons 4-8 RT-PCR: P53 5'-TTTTGGACACGCCGAAGTAGT-3'
LRRK1 PCR Reverse Primer for exons 4-8 RT-PCR: P54 5'-AGCCGCTCCAGGTAGTTTTT-3'
LRRK1 PCR Forward Primer for exons 11-17 RT-PCR: P57 5'-GGACCTCTCCAGAAACCAGC-3'
LRRK1 PCR Reverse Primer for exons 11-17 RT-PCR: P58 5'-GCAGGGTTGCTATCCTCTCC-3'
LRRK1 PCR Forward Primer for exons 20-25 RT-PCR: P63 5'-GCGGTCAGTGGCAAAGAATG-3'
LRRK1 PCR Reverse Primer for exons 20-25 RT-PCR: P64 5'-AATGCTGTTCTCACCCTCCG-3'
LRRK1 PCR Forward Primer for exons 25-31 RT-PCR: P41 5'-GAATTCTGCTAATGCCCCAGC-3'
LRRK1 PCR Reverse Primer for exons 25-31 RT-PCR: P42 5'-AGGCTGTAGATATAGATTTTCTGGT-3'

LRRK2 PCR Forward Primer for Left arm amplification: P9 5'-GAACACACAAGGCTATGGCTATTGTC-3'
LRRK2 PCR Reverse Primer for Left arm amplification: P10 5'-GTAGGACTATCATCCACCTGTAGGACTCC-3'
loxP site introducing oligo for *LRRK2*: P39 5'-<u>GATCC</u>ATAACTTCGTATAATGTATGCTATACGAAGTTAT<u>GCTAGC</u>A-3' *BamH*I and *Nhe*I sequences are underlined.
loxP site introducing oligo for *LRRK2*: P40 5'-<u>CTAGTGCTAGC</u>ATAACTTCGTATAGCATACATTATACGAAGTTATG-3' *Spe*I and *Nhe*I sequences are underlined.
LRRK2 PCR Forward Primer for Middle arm amplification: P11 5'-GTCCTACAGGTGGAT<u>TCTAGA</u>CCTACAAGG-3' *Xba*I sequence is underlined.
LRRK2 PCR Reverse Primer for Middle arm amplification: P12 5'-GAC<u>GCGGCCGC</u>GTTTAGTA<u>GCTAG</u>CATGACTATGAAGGAG-3' *Not*I and *Nhe*I sequences are underlined.
LRRK2 PCR Forward Primer for Right arm amplification: P13 5'-GCACTTGAGTCTTAATCTTGGGCAC-3'
LRRK2 PCR Reverse Primer for Right arm amplification: P14 5'-CATTCGAGCAGCTAAGCCTGTAATC-3'
LRRK2 PCR Forward Primer for 5' Southern Probe: P19 5'-CATGGGAGAGAGGGTTTCTCACTTACT-3'
LRRK2 PCR Reverse Primer for 5' Southern Probe: P20 5'-CTTGGACAGCATTGTCAGCCTAGAC-3'
LRRK2 PCR Forward Primer for 3' Southern Probe: P21 5'-GCCAAGCAGTTATTGATGCTGTAGC-3'
LRRK2 PCR Reverse Primer for 3' Southern Probe: P22 5'-CGAGCTGTAAGATGAGCTGGGTACT-3'
LRRK2 PCR Forward Primer for Northern Probe: P35 5'-AGGAAGGCAAGCAGATCGAG-3'
LRRK2 PCR Reverse Primer for Northern Probe: P36 5'-GGCTGAATATCTGTGCATGGC-3'
LRRK2 RT Primer in exons 51: P93 5'-TCGTGTGGAAGATTGAGGTCC-3'
LRRK1 PCR Forward Primer for Genotyping: P25 5'-ATTGGTCTTTGAAGAGACAGCATCTGG-3'
LRRK1 PCR Reverse Primer for Genotyping: P26 5'-TTTCCCTGAGGTGGAGAAGTGACTGG-3'
LRRK1 PCR Reverse Primer for Genotyping: P27 5'-TCACGTCGTCTAAGCCTCCT-3'
LRRK1 PCR Forward Primer for Genotyping: P28 5'-CTTCCTCAGAAGTTAGGTAAACATTG AGTG-3'
LRRK1 PCR Reverse Primer for Genotyping: P29 5'-CTAAGTGACACCGTGTTTCCAAAGTC-3'
LRRK1 PCR Reverse Primer for Genotyping: P30 5'-GGAAAGTTTCACAATTGGAAAAATAAAAATATTTACTGCAGATA-3'
DAT-IRES-Cre PCR Forward Primer for Genotyping: JKM1823 5'-TGGCTGTTGGTGTAAGTGG-3'
DAT-IRES-Cre PCR Reverse Primer for Genotyping: JKM1824 5'-GGACAGGGACATGGTTGACT-3'

> *DAT-IRES-Cre* PCR Reverse Primer for Genotyping: JKM1825
> 5'-CCAAAGACGGCAATATGGT-3'

Mice

All animal use was approved by the IACUC committees of Harvard Medical School and Brigham

and Women's Hospital, and conformed to the USDA Animal Welfare Act, PHS Policy on Humane Care

and Use of Laboratory Animals, the "ILAR Guide for the Care and Use of Laboratory Animals" and other

applicable laws and regulations. Mice were housed in constant humidity- and temperature-controlled

rooms and maintained on a 12 hr light/dark cycle and were given standard rodent chow and water. Mice

of both sexes at multiple ages, from 2 months to 25 months, were used. Floxed *LRRK1* (*fLRRK1*) and the

resulting germline deleted *LRRK1*$^\Delta$ alleles, and floxed *LRRK2* (*fLRRK2*) and the resulting germline

deleted *LRRK2*$^\Delta$ alleles were generated and thoroughly validated at the genomic DNA, mRNA, and

protein levels. *DAT-IRES-Cre* (The Jackson Laboratory, IMSR_JAX:006660), *ACT-FLPe*

(IMSR_JAX:005703), *CMV-Cre* transgenic (IMSR_JAX:006054), and *Rosa26-CAG-LSL-ZsGreen1* mice

(IMSR_JAX:007906) used in the current study were previously characterized and reported (32-35).

LRRK1/LRRK2 cDKO mice (*fLRRK1/fLRRK1; fLRRK2/fLRRK2; DAT-IRES-Cre/+*) and littermate

controls (*fLRRK1/fLRRK1; fLRRK2/fLRRK2*) were obtained from multiple breeding cages of

fLRRK1/fLRRK1; fLRRK2/fLRRK2 and cDKO mice. All mice were maintained on the C57BL6 and 129

hybrid genetic background (F1: IMSR_JAX:101043). All phenotypic analyses were performed in a

genotype blind manner, as previously described (67).

Generation of targeted and floxed *LRRK1* and *LRRK2* alleles

The generation and validation of the *LRRK1* and *LRRK2* targeting vectors, the targeted, floxed,

and deleted *LRRK1* and *LRRK2* alleles by Southern, Northern, RT-PCR, and the sequences of the floxed

LRRK1 and *LRRK2* alleles are included in Figure 1-figure supplement 1-11.

LRRK1

To generate the *LRRK1* targeting vector, we first PCR amplified the *left middle* homologous

region (2,079 bp) containing partial intron 26, exon 27, and partial intron 27 of *LRRK1* from mouse BAC

DNA (clone RP23-213J23, BACPAC Resources Center) using primers, P3 and P4. The PCR fragment

was subcloned into the *pGEM-T* vector (A1360, Promega) to generate pLM1 (for details, see Figure 1-

figure supplement 1). The *right middle* homologous region (3,403) containing *LRRK1* genomic region

from partial intron 27 to partial intron 29 was amplified by PCR using primers, P5 and P6, and then

subcloned into the *pGEM-T* vector (pLM2). The *left middle* homologous region containing *pGEM-T*

plasmid was then digested with *Not*I and *Kpn*I (endogenous *Kpn*I site in the intron 27) and subcloned into

the *Not*I and *Kpn*I sites of the *right middle* homologous region containing *pGEM-T* plasmid (pLM3).

After that, the *middle* homologous region (5.5 kb), from partial intron 26 to partial intron 29, was released

by *Not*I and *Sac*II digestions and was blunted by Klenow, and then subcloned into the *Sma*I site of

PGKneoF2L2DTA vector (#13445, addgene) to generate the *middle* homologous region-

PGKneoF2L2DTA plasmid (pLM4).

The *left* homologous region (2,016 bp) containing partial intron 25, exon 26, and partial intron 27

of *LRRK1* was PCR amplified from mouse BAC DNA (clone RP23-213J23, BACPAC Resources Center)

using primers, P1 and P2. The PCR fragment was digested with *Sac*II (introduced by P1) and *Not*I

(introduced by P2 along with *Hind*III), and was subcloned into the *Sac*II and *Not*I sites of the

PGKneoF2L2DTA vector (pLM5). The *right* homologous region (3,131 bp) containing partial intron 29,

exon 30, and partial intron 30 of *LRRK1* was PCR amplified from mouse BAC DNA (clone RP23-213J23)

using primers, P7 and P8. The PCR fragment was digested with *Sal*I (introduced by P7) and *Hind*III

(introduced by P8), and was subcloned into the *Sal*I and *Hind*III sites of the *PGKneoF2L2DTA* vector

(pLM6).

The *left* homologous region in the *PGKneoF2L2DAT* vector (pLM5) was digested with *Sac*II and

*Not*I, and subcloned into the *Sac*II and *Not*I sites of the *right* homologous region containing

pGKneoF2L2DTA plasmid (pLM6) to generate pLM7. Finally, the *loxP-middle* homologous region-*PGK-*

Neo-loxP fragment was released from pLM4 by *Not*I and *Sal*I digestions, and subcloned into the *Not*I and

*Sal*I sites of pLM7 to generate the final target vector (pLM8), which contains two *loxP* sites (intron 26

and intron 29). Upon Cre-mediated recombination, the endogenous *LRRK1* genomic sequence flanked by

the 5' *loxP* site (1,288 bp upstream of *LRRK1* exon 27) to the 3' *loxP* site (1,023 bp downstream of exon

29) is excised. The *PGK-neo* cassette flanked by two *FRT* sites is under the control of the mouse

phosphoglycerate kinase 1 (*PGK*) promoter and contains the *bovine growth hormone* polyA signal. To

enhance the ratio of ES cells carrying homologous recombination events instead of random insertion of

the targeting vector (68), the negative selection *PGK-DTA* cassette, which encodes diphtheria toxin A

chain, was also included in the *LRRK1* targeting vector.

The *LRRK1* targeting vector was linearized by *Xho*I digestion, and then electroporated into

MKV6.5 embryonic stem (ES) cells (a gift from R. Jaenisch), which were derived from B6/129 F1 mice

(The Jackson Laboratory, IMSR_JAX:101043). G418 was applied to the culture at 150 μg/ml 24 h later,

and after 6 days of G418 selection, the surviving ES clones (480) were picked and screened by Southern

analysis using *Hind*III digestion of genomic DNA followed by hybridization with the 5' external and 3'

external probes to confirm proper recombination events in the 5' and 3' homologous regions, respectively.

25 ES cell clones were confirmed to carry the proper homologous recombination events at the 5'

homologous region, giving rise to the 17.0 kb and the 4.8 kb bands, which represent the wild-type and the

targeted allele, respectively (Figure 1D), and the 3' homologous region, giving rise to the 17.0 kb and the

14.1 kb bands, which represent the wild-type and the targeted allele, respectively. We then expanded the

selected 4 ES cell clones (3A11, 3D8, 3H1, and 3H6) and further verified by Southern using the 5' and 3'

external, and *neo* probes. Two ES cell clones (3D8 and 3H6) were transfected with *pCAGGS-flpE-puro*

(#20733, addgene) to delete the *PGK-neo* cassette by FLP recombinase. Three resulting ES cell clones

(3D8C5, 3D8E5, and 3H6G7) were confirmed by Southern analysis using the 5' and 3' external probes

and the *neo* probe to confirm the floxed *LRRK1* allele by the deletion of the *PGK-neo* cassette.

LRRK2

To generate the *LRRK2* targeting vector, the *left* homologous region (2,579 bp) containing the

LRRK2 promoter region was PCR amplified from mouse BAC DNA (clone RP23-526A2, BACPAC

Resources Center) using primers, P9 and P10, and subcloned into the *pGEM-T* vector to generate

pLRRK2#1 (for details see Figure 1-figure supplement 4). The *BamH*I-*loxP*-*Nhe*I-*Spe*I fragment (56 bp),

which was generated by annealing two complementary oligos, P39 and P40, was introduced to the *BamH*I

and *Spe*I sites of pLRRK2#1 to generate pLRRK2#2. The *middle* homologous region, containing genomic

sequences from the promoter region to partial intron 2, was PCR amplified from mouse BAC DNA (clone

RP23-526A2) using primers, P11 and P12, digested with *Xba*I and *Not*I, and subcloned into the *Xba*I and

*Not*I sites of the *pGEM-T* vector to generate pLRRK2#3, which was digested with *Xba*I and *Not*I, and

subcloned into the *Xba*I and *Not*I sites of pLRRK2#2 to generate pLRRK2#4.

The *right* homologous region (3,503 bp), which contains partial intron 1, exon 2, and partial

intron 2, was PCR amplified from mouse BAC DNA (clone RP23-526A2) using primers, P13 and P14,

and the PCR fragment was subcloned into the *EcoR*V site of *pBluescript II KS* (+) vector (212207,

Agilent) to generate pLRRK2#5. The *right* homologous region in pLRRK2#5 was digested with *BamH*I

followed by Klenow, and then digested with *Cla*I. The released fragment was subcloned into the *EcoR*V

and *Cla*I sites of the *PGKneoF2L2DTA* vector to generate pLRRK2#6. *LFNT-tk/pBS* plasmid (a gift from

S. Tonegawa) was digested with *Sac*II (followed by Klenow) and *Not*I (followed by *Ssp*I digestion to

make it easier to isolate the *Not*I-*Sac*II/KN fragment) to release the *loxP-FRT-PKG-neo-loxP-FRT*

fragment (2,928 bp), and then subcloned into the *Xba*I (followed by Klenow) and *Not*I sites of

pLRRK2#6 to generate pLRRK2#7. Finally, pLRRK2#4 containing the *left-loxP-middle* homologous

regions was digested with *EcoR*V and *Not*I, and subcloned into *Ale*I and *Not*I sites of pLRRK2#7

containing the *loxP-FRT-PKG-neo- loxP-FRT-right* homologous region to generate the final targeting

vector, which contains the *loxP* sites in the promoter region (1,768 bp upstream of the transcription

initiation site) and in *LRRK2* intron 2 (878 bp downstream of exon 2). Upon Cre-mediated recombination,

the floxed endogenous *LRRK2* genomic sequences from the 5' *loxP* site (1,768 bp upstream of *LRRK2*

transcription) to the 3' *loxP* site in intron 2 (878 bp downstream of exon 2) are excised. The negative

selection *PGK-DTA* cassette is also included in the *LRRK2* targeting vector.

The *LRRK2* targeting vector was linearized by *Ahd*I digestion, and then electroporated into

MKV6.5 ES cells. G418 was applied to the culture at 150 ug/ml 24 h later, and after 6 days of G418

selection, the surviving ES clones (480) were picked and screened by Southern analysis using *Nhe*I

digestion of genomic DNA followed by hybridization with the 5' external probe (753 bp, PCR amplified

using P19 and P20). Southern analysis confirmed that two ES cell clones (N24 and N34) carry the proper

recombination event in the 5' homologous arm, giving rise to 11.6 kb and 3.6 kb bands, which represent

the wild-type and targeted alleles, respectively (Figure 1F), followed by genomic PCR confirmation for

the proper recombination in the right arm.

Three (3D8C5, 3A11, 3H6) and one (N24) ES clones for *LRRK1* and *LRRK2*, respectively, were

microinjected into C57BL/6 mouse blastocysts to generate chimera mice, which were bred with B6/129

F1 mice to produce heterozygous floxed *LRRK1* and targeted *LRRK2* mice. Floxed *LRRK1* mice were

confirmed by Southern analysis using the 5' and 3' external probes (Figure 1D). Targeted *LRRK2* mice

were confirmed by Southern (data shown in Figure 1-figure supplement 6; *Nhe*I digestion followed by

hybridization with the 5' external probe and *Sph*I digestion followed by hybridization with the 3' external

probe). Targeted *LRRK2* mice were then bred with *Actin-FLP* deleter mice (IMSR_JAX:005703) (33) to

generate floxed *LRRK2* mice, which were confirmed by Southern analysis using the 5' and 3' external

probes following *Nhe*I digestion (Figure 1F). Heterozygous *fLRRK1*/+ and *fLRRK2*/+ mice were crossed

with each other to obtain homozygous single floxed mice (*fLRRK1/fLRRK1* and *fLRRK2/fLRRK2*) and

double floxed mice (*fLRRK1/fLRRK1*; *fLRRK2/fLRRK2*).

Generation of deleted *LRRK1/LRRK2* alleles and DA neuron-specific cDKO mice

In order to ensure that Cre-mediated deletion of the floxed *LRRK1* and *LRRK2* alleles results in

null alleles, we crossed floxed *LRRK1* and *LRRK2* mice with germline deleter, *CMV-Cre* transgenic mice

(*B6.C-Tg(CMV-cre)1Cgn/J*; IMSR_JAX:006054) (34), to generate *LRRK1* deleted (Δ/Δ) mice (by

removing exons 27-29) and *LRRK2* deleted (Δ/Δ) mice (by removing the promoter region and exons 1-2)

for further molecular characterization. To generate DA neuron-specific *LRRK1/LRRK2* cDKO mice, we

used *DAT-Cre* KI mice (*B6.SJL-Slc6a3tm1.1(cre)Bkmn/J;* IMSR_JAX:006660), which express Cre

recombinase under the control of the endogenous *DAT* promoter (32). We crossed double floxed mice

with *DAT-Cre* mice to generate *LRRK* cDKO mice (*fLRRK1/fLRRK1; fLRRK2/fLRRK2; DAT-Cre/+*).

LRRK cDKO and littermate control mice used in the phenotypic analysis were obtained by crossing

fLRRK1/fLRRK1; fLRRK2/fLRRK2; DAT-Cre/+ with *fLRRK1/fLRRK1; fLRRK2/fLRRK2* mice. We only

used cDKO and control mice that carry all floxed *LRRK1* and *LRRK2* alleles for phenotypic analysis.

Southern analysis

For the identification and validation of the targeted and floxed *LRRK1* alleles, we used the 5'

external, 3' external, and *neo* probes. Genomic DNA from ES cells or mouse tails was digested with

*Hind*III. The 5' external probe (377 bp), which is 839 bp upstream of the 5' homologous region, was PCR

amplified from mouse BAC DNA (clone RP23-213J23) using primers P15 and P16. The 3' external

probe (305 bp), which is 2,736 bp downstream of the 3' homologous region, was PCR amplified from

mouse BAC DNA (clone RP23-213J23) using primers P17 and P18. The *neo* probe (363 bp) was PCR

amplified from pSoriano plasmid using primers P23 and P24. Following *Hind*III digestion, the presence

of the 17.0 kb product using either the 5' or 3' external probe represents the wild-type allele, whereas the

4.8 kb (the 5' external probe) and the 12.2 kb (the 3' external probe) products represent the floxed *LRRK1*

allele. Genomic DNA digested by *Hind*III and hybridized with the *neo* probe further confirmed the wild-

type and floxed alleles (no band) and the targeted *LRRK1* allele (14.1 kb).

For the identification and validation of the targeted and floxed *LRRK2* alleles, we used the 5' and

3' external probes as well as the *neo* probe. Genomic DNA from ES cells or mouse tails was digested

with *Nhe*I or *Sph*I followed by hybridization with the 5' or 3' external probe or the *neo* probe. The 5'

external probe (753 bp), which is 84 bp upstream of the 5' homologous region, was PCR amplified from

mouse BAC DNA (clone RP23-526A2) using primers P19 and P20. The 3' external probe (622 bp),

which is 38 bp downstream of the 3' homologous region, was PCR amplified from mouse BAC DNA

(clone RP23-526A2) using primers P21 and P22. Following *Nhe*I digestion, the presence of the 11.5 kb

product using either the 5' or 3' external probe represents the wild-type allele, whereas the 3.6 kb (the 5'

external probe) and the 5.2 kb (the 3' external probe) products represent the floxed *LRRK2* allele.

PCR genotyping

Genomic PCR was performed to determine the presence of the deleted, the floxed, and/or the

wild-type alleles.   For   *LRRK1*,   the   following   primers   were   used: P25   (5'-

ATTGGTCTTTGAAGAGACAGCATCTGG, forward primer, 392 nt downstream of exon 26), P26 (5'-

TTTCCCTGAGGTGGAGAAGTGACTGG, reverse primer, 567 nt downstream of exon 26), and P27

(5'-TCACGTCGTCTAAGCCTCCT, reverse primer, 1,218 nt downstream of exon 29).   The PCR

products from P25 and P26 are 266 bp and 405 bp, which represent the wild-type and the *fLRRK1* alleles,

respectively, whereas the PCR product from P25 and P27 is 583 bp, which represents the deleted *LRRK1*

allele.

For   *LRRK2*,   the   following   primers   were   used: P28   (5'-

CTTCCTCAGAAGTTAGGTAAACATTGAGTG, forward primer, 2,069 nt upstream of exon 1), P29

(5'-CTAAGTGACACCGTGTTTCCAAAGTC, reverse primer, 1,739 nt upstream of exon 1), and P30

(5'-GGAAAGTTTCACAATTGGAAAAATAAAAATATTTACTGCAGATA, reverse primer at 2,848 nt

downstream of exon 2). The PCR products from P28 and P29 are 305 bp and 367 bp, representing the

wild-type and the *fLRRK2* alleles, respectively, whereas the PCR product from P28 and P30 is 587 bp,

which represents the deleted *LRRK2* allele.

For   *DAT-IRES-Cre*,   the   following   primers   were   used: JKM1823   (5'-

TGGCTGTTGGTGTAAAGTGG, forward primer at exon 16 and 3'-UTR), JKM1824 (5'-

GGACAGGGACATGGTTGACT,   reverse   primer   at   3'-UTR),   and   JKM1825   (5'-

CCAAAAGACGGCAATATGGT, reverse primer at *IRES* sequence). The PCR product from JKM1823

and JKM1824 is 264 bp, which represents the wild-type allele, whereas the PCR product from JKM1823

and JKM1825 is 152 bp, which represents the *DAT-IRES-Cre* KI allele.

Northern analysis

Total RNA was isolated from brains, kidneys, or lungs using TRI reagent (T9424, Millipore-

Sigma) according to the manufacturer's instruction. For the *LRRK1* Northern analysis, polyA+ RNA was

enriched from ~500 μg total RNA using the Poly(A)Purist™ MAG Kit (AM1922, ThermoFisher)

according to the manufacturer's instruction. For the *LRRK2*, ~10 μg of total RNA was used for Northern

analysis. RNA was separated in formaldehyde agarose gel, and transferred into Amersham Hybond-nylon

membrane (RPN303N, GE Healthcare). Probes were synthesized using Prime-It II random labeling kit

(#300385, Agilent) and then used for membrane hybridization at 55°C overnight.

The cDNA probe specific for *LRRK1* exons 2-3 (383 bp) was PCR amplified using primers P31

(5'-CAGGATGAGCGTGTGTCTGCAG) and P32 (5'-CCTTCTCCTGTGAGGATTCGCTCT). The

cDNA probe specific for *LRRK1* exons 27-29 (550 bp) was PCR amplified using primers P33 (5'-

CTGGCCTACCTGCACAAGAA) and P34 (5'-CCTTCCCATCCCAGAACACC).

The cDNA probe specific for *LRRK2* exons 1-5 (437 bp) was PCR amplified using primers P35

(5'-AGGAAGGCAAGCAGATCGAG) and P36 (5'-GGCTGAATATCTGTGCATGGC). The probe

specific for *GAPDH* exons 5-7 (452 bp) was PCR amplified using primers P37 (5'-

ACCACAGTCCATGCCATCAC) and P38 (5'-TCCACCACCCTGTTGCTGTA).

Hybridization was performed using α-^{32}P-dCTP-labeled probes specific to each gene. Specific

signals were detected by autoradiography with Hyperfilm (E3018, Amersham).

RT-PCR

493   Total RNA was isolated from brains, kidneys, or lungs using TRI reagent (T9424, Millipore-

494   Sigma) according to the manufacturer's instruction. Approximately 1 μg of RNA was reverse-transcribed

495   using Superscript III (18080093, Thermo Fisher Scientific) according to the manufacturer's instructions.

496   For RT-PCR analysis of *LRRK1* transcripts in mice carrying the homozygous floxed or deleted alleles, we

used primers P53 and P54 for exons 4-8 (714 bp), P57 and P58 for exons 11-17 (818 bp), P63 and P64 for exons 20-25 (922 bp), or P41 and P42 for exons 25-31 to confirm normal splicing of *LRRK1* mRNA in *fLRRK1/fLRRK1* mice and truncated *LRRK1* transcripts lacking exons 27-29 in Δ/Δ mice (for details see Figure 1-figure supplement 9). For RT-PCR analysis of *LRRK2* transcripts, we used primers P35 and P36 in exons 1-5 to confirm normal splicing of *LRRK2* mRNA (437 bp) in *fLRRK2/fLRRK2* mice and the absence of RT-PCR products in *LRRK2* Δ/Δ mice (Figure 1-figure supplement 11). The identity of the PCR products was confirmed by sequencing.

Western analysis

Fresh tissues were collected and homogenized in an ice-cold stringent RIPA buffer (50 mM Tris-Cl (pH 7.6), 150 mM NaCl, 0.5 mM EDTA, 1% NP40, 0.5% sodium deoxycholate, 0.1% SDS, 1mM PMSF supplement with protease inhibitor cocktail (P8340, Sigma) and phosphatase inhibitor cocktail (P0044, Sigma)), followed by sonication. Homogenates were centrifuged at 14,000xg for 20 min at 4°C to separate supernatants (RIPA buffer-soluble fraction). An equal amount (10-40 μg per lane) of total proteins from each preparation were loaded and separated on NuPAGE gels (Invitrogen), and transferred to nitrocellulose membranes. The membranes were blocked in Intercept (TBS) Blocking Buffer (927-60001, Li-Cor) for 1 hr at room temperature, and incubated at 4°C overnight with specific primary antibodies. Primary antibodies used were rabbit anti-LRRK1 (ANR-101, Alomone Lab, RRID: AB_2756700), rabbit anti-LRRK2 (ab133474, abcam, RRID: AB_2713963), mouse anti-α-Vinculin (05-386, Millipore, RRID: AB_309711). Membranes were then incubated with dye-coupled secondary antibodies, goat anti-mouse IRdye680 (#925-68070, LI-COR, RRID: AB_2651128) or goat anti-rabbit IRdye800 (#925-32211, LI-COR, RRID: AB_2651127). Signals were quantified using the Odyssey Infrared Imaging System (LI-COR).

Histological analysis

Mice were anesthetized with ketamine (100 mg/kg) + xylazine (10 mg/kg) + acepromazine (3 mg/kg), and transcardially perfused with phosphate-buffered saline solution (PBS, pH 7.4) containing

0.25 g/L heparin (H3149, Sigma) and 5 g/L procaine (P9879, Sigma). Brains were post-fixed in 4%

formaldehyde in PBS (pH 7.4) (15710, Electron Microscopy Sciences) at 4˚C overnight and then

processed for paraffin embedding following standard procedures. For frozen sections, post-fixed brains

were immersed in a sucrose series solution (15% and 30% sucrose in PBS) at 4°C overnight for

cryoprotection, and then brains were embedded in an OCT compound (4583, Sakura). Serial coronal

sections (16 μm) of paraffinized brains or frozen brains were obtained using Leica RM2235 or Leica

CM1860, respectively. Coronal sections containing the SNpc or LC were selected for

immunohistochemical analysis.

Histological analyses were performed as described previously (30, 31, 67). Briefly, for DAB-

derived TH-immunohistochemistry, coronal sections were deparaffinized, alcohol-dehydrated, and then

subjected to permeabilization with a solution containing 0.1% Triton X-100 in TBS followed by antigen

retrieval for 5 min in 10 mM sodium citrate buffer, pH 6.0. Endogenous peroxidase activity was quenched

by incubating in 0.3% H_2O_2 in methanol. Sections were then blocked with a solution containing 5%

normal goat serum (S-1000, Vector Laboratories) and 0.1% Triton X-100 in TBS for 1 hr at room

temperature. After blocking, sections were incubated with the primary antibody, rabbit anti-TH (1:750,

ab112, abcam, RRID: AB_297840) overnight at 4°C. Sections were washed three times in 0.1% Triton X-

100 in TBS followed by 1 hr incubation with goat biotinylated anti-rabbit IgG secondary antibody (1:250,

BA-1000, Vector Laboratories, RRID: AB_2313606) at room temperature and 30 min incubation with

Vectastain Elite ABC reagent (PK-6100, Vector Laboratories) and then developed using chromogenic

DAB substrate (SK-4100, Vector Laboratories).

For immunofluorescence staining of paraffin sections, coronal sections were deparaffinized,

alcohol-dehydrated, and then subjected to permeabilization with a solution containing 0.1% Triton X-100

in PBS followed by antigen retrieval for 5 min in 10 mM sodium citrate buffer, pH 6.0 except those for

cleaved-caspase3 immunostaining which were performed antigen retrieval for 10 min. Sections were then

blocked with a solution containing 10% normal goat serum and 0.1% Triton X-100 in PBS for 1 hr at

room temperature. After blocking, sections were incubated with primary antibodies overnight at 4°C. The

primary antibodies used were mouse anti-TH (1:50, sc-25269, Santa Cruz, RRID: AB_628422), rabbit

anti-NeuN (1:400, 12943S, Cell Signaling Technology, RRID: AB_2630395), rabbit anti-cleaved

caspases-3 (1:150, 9661, Cell Signaling Technology, RRID: AB_2341188), or rabbit anti-Iba1 (1:500,

019-19,751, Wako, RRID: AB_839504). Sections were washed three times in 0.1% Triton X-100 in PBS

followed by 1 hr incubation with fluorophore-conjugated secondary antibodies, goat anti-mouse IgG

Alexa Fluor 488 (1:500, A11001, ThermoFisher, RRID: AB_2534069) and goat anti-rabbit IgG Alexa

Fluor 555 (1:500, A32732, ThermoFisher, RRID: AB_2633281) at room temperature. Fluorescence

images were taken (a stack of 2-3 confocal images spaced at 4 μm), projected with maximal intensity

projection mode, and analyzed by FV1000 confocal microscope system (Olympus)

For immunofluorescence staining of cryopreserved sections, coronal brain sections were washed

with PBS to rinse out OCT, and then blocked with a solution containing 5% normal goat serum and 0.1%

Triton X-100 in TBS for 1 hr at room temperature. After blocking with 10% NGS, sections were

incubated with primary antibodies, rabbit anti-GFP (1: 1000, ab290, abcam, RRID: AB_303395), and

mouse anti-TH (1:50, sc-25269, Santa Cruz, RRID: AB_628422), overnight at 4°C. Sections were

incubated with fluorophore-conjugated secondary antibodies, goat anti-rabbit IgG Alexa Fluor 488 (1:500,

A11034, ThermoFisher, RRID: AB_2576217) and goat anti-mouse IgG Alexa Fluor 555 (1:250, A21424,

ThermoFisher, RRID: AB_141780) for 1 hr at room temperature. Fluorescence images were taken and

analyzed by FV1000 confocal microscope system (Olympus).

The number of GFP+ and GFP+/TH+ cells in the SNpc of *DAT-IRES-Cre/+; Rosa26-CAG-LSL-*

ZsGreen1/+ reporter mice were quantified using 3 comparable coronal sections (16 μm in thickness,

spaced 320 μm apart) per brain (n = 3 brains, one hemisphere). The percentage of GFP+ DA neurons in

the SNpc was obtained by dividing the sum of GPF+/TH+ neurons by the sum of total TH+ neurons

quantified.

Quantification of TH+ DA neurons in the SNpc or LC was performed as previously described (30,

31, 67, 69). Briefly, TH+ neurons in the SNpc, which were marked based on morphological features as

previously described (70), were quantified in every 10^{th} serial coronal section (16 μm in thickness)

throughout the SNpc (a total of 6-9 sections, spaced 160 μm apart). Total number of TH+ cells in the

SNpc was calculated as follows: [total number of TH+ DA neurons quantified in all 6-9 sections] x 10

(every 10^{th} section sampled) x 2 (both hemispheres). Similar results were obtained independently by two

investigators using coded samples. Total number of TH+ noradrenergic neurons in the LC (a total of 6-9

sections, spaced 80 μm apart) was calculated as follows: [total number of TH+ noradrenergic neurons

quantified in all 6-9 sections] x 5 (every 5^{th} section sampled) x 2 (both hemispheres).

NeuN+, active Caspase-3+, or Iba1+ cells in the SNpc, which was marked by TH

immunoreactivity, were quantified using serial coronal sections (16 μm in thickness, every 10^{th} section, a

total of 6-9 sections per brain). Compared to chromogenic DAB immunostaining followed by counting

under the stereomicroscope, which captures TH-positive dopaminergic neurons from the single plane

layer of the sections, immunofluorescence immunostaining picks up more cells with higher sensitivity by

maximal intensity projection mode throughout the entire layer of the section under the confocal

microscope. Thus, more TH-positive dopaminergic neurons in the SNpc were captured and quantified

under confocal microscopy compared to DAB staining. The total number of NeuN+, active Caspase-3+,

or Iba1+ cells in the SNpc, which was marked by TH immunoreactivity, was calculated as follows: [total

number of NeuN+, active Caspase-3+, or Iba1+ cells in all sections counted] x 10 (every 10^{th} section

sampled) x 2 (both hemispheres).

Quantitative EM analysis

The collection and quantification of the EM images were performed as described previously (30,

71). Mice were perfused with PBS containing 0.25 g/L heparin and 5 g/L procaine followed by a fixative

solution containing 2.5% paraformaldehyde and 2.5% glutaraldehyde in 0.1 M sodium cacodylate buffer

(pH 7.4) (#1549, Electron Microscopy Science). Brains were dissected and post-fixed overnight in a

fixative solution at 4°C. The dissected tissues were trimmed to 1-2 mm^3 cubes followed by osmication

and uranyl acetate staining, dehydration in graded alcohol, and embedded in TABB 812 Resin (Marivac

Ltd.) at the Harvard Medical School EM facility. 0.5 μm sections were stained with toluidine blue and

viewed under the light stereomicroscope (Nikon Eclips E600) to find the SNpc area for EM viewing.

Then adjacent sections were cut with 80 nm in thickness with the Leica Ultracut S microtome, picked up

on formvar-carbon coated slot Copper grids, stained with 0.2% Lead Citrate, and viewed and imaged

under the JEOL 1200x Electron Microscope. A minimum of 10 micrographs containing the entire cell

body in the SNpc area were analyzed for each brain. The image was analyzed by using the Fiji version of

ImageJ. The number of electron-dense autophagic and lysosomal vacuoles (> 0.5 μm in diameter) in

individual neuronal profiles was quantified. We previously calculated the diameter of electron-dense

autophagic/lysosomal vacuoles by measuring the longest side manually (30, 31). In the current study, we

used Feret's Diameter (72) to accurately measure the longest distance between any of two points in the

electron-dense autophagic/lysosomal vacuoles, resulting in a higher number of electron-dense vacuoles

quantified. The number of electron-dense vacuoles (> 0.5 μm in diameter) in individual neuronal profiles

was quantified. The area of electron-dense vacuoles (> 0.5 μm in diameter) was quantified and averaged

in individual neuronal profiles. Finally, the average number or area of vacuoles was calculated per

genotype. Experiments were done in a genotype blind manner (after scarifying mice, the brain samples

were coded and sent to the Harvard Medical School EM Core, where the images were captured, and the

vacuoles were quantified by another independent experimenter).

Experimental design and statistical analysis

Group sizes were determined based on prior studies. Littermates of control and cDKO mice were

randomly distributed between experimental groups. Data acquisition and quantification were performed in

a genotype blind manner with the exception of molecular analyses (Southern, Northern, RT-PCR, and

Western). All statistical analyses were performed using Prism 9 (GraphPad software) or Excel (Microsoft).

All data are presented as the means ± SEM. The exact sample size, the number of mice or neurons, of

each experiment is indicated in the figure or the legend.

622   Statistical analyses were conducted using a two-tailed Student's *t*-test for the comparison of a

623 given variable in two genotypes or two-way ANOVA followed by Bonferroni's post hoc comparisons for

624 the comparison of more than two conditions. Statistical outliers were identified and excluded using the

625 ROUT method with 1% the maximum desired false discovery rate developed by Prism. All statistical

626 comparisons were performed on data from ≥ 4 biologically independent samples and replicated on

627 different experimental days. Significance is shown as *p < 0.05, **p < 0.01, ***p < 0.001, ****p $<$

628 0.0001, and not significantly different values are often not noted.

629 Data availability

630   All data generated or analyzed during this study are included in the manuscript and supporting

631 file. Source Data files have been provided for Figures 1 to 7.

632 Funding disclosure

633   This work was supported by a grant from the NIH (R37NS071251 to J.S.).

634 Acknowledgments

635   We thank H. Zhao for technical assistance, Y. Yuan for organizing experimental data, and Shen

636 lab members for discussion. We are grateful to M. Ericsson at the Harvard Medical School Electron

637 Microscopy Core for her expert assistance.

638 Author Contributions

639   J.S. conceived and directed the project, J.K. performed Western blotting and quantitative

640 histological and EM analyses of *LRRK* cDKO mice (Figures 1I, 2E, F, 3, 4, 5A, B, 6, 7), G.H. performed

641 Northern blotting and RT-PCR followed by sequencing on mice carrying the floxed and deleted *LRRK1*

642 and *LRRK2* alleles (Figure 1G, H; Figure 1-figure supplement 8-11), preparation of brain sections and

643 histological analysis of TH+ neurons in *LRRK* cDKO mice at all ages (Figure 3A), J.K. and G.H. verified

644 all molecular results and compiled Figure 1-figure supplement 1, 2, 4, 5, L.M. (LRRK1) and Y.T.

645 (LRRK2) generated the targeting vectors and performed ES cell experiments including Southern blotting

(Figure 1C-F; Figure 1-figure supplement 1-6), P.C. performed GFP reporter analysis of *DAT-Cre* mice in

the SNpc (Figure 2A, B) and the LC (Figure 5C), J.K., G.H., P.C. analyzed the data and made the figures,

J.K., G.H., and J.S. wrote the paper.

References

1.   Paisán-Ruíz C, Jain S, Evans EW, Gilks WP, Simón J, van der Brug M, et al. Cloning of the gene
containing mutations that cause PARK8-linked Parkinson's disease. Neuron. 2004;44(4):595-600.

2.   Shen J. Protein kinases linked to the pathogenesis of Parkinson's disease. Neuron.
2004;44(4):575-7.

3.   Zimprich A, Biskup S, Leitner P, Lichtner P, Farrer M, Lincoln S, et al. Mutations in LRRK2
cause autosomal-dominant parkinsonism with pleomorphic pathology. Neuron. 2004;44(4):601-7.

4.   Gilks WP, Abou-Sleiman PM, Gandhi S, Jain S, Singleton A, Lees AJ, et al. A common LRRK2
mutation in idiopathic Parkinson's disease. Lancet. 2005;365(9457):415-6.

5.   Lesage S, Janin S, Lohmann E, Leutenegger AL, Leclere L, Viallet F, et al. LRRK2 exon 41
mutations in sporadic Parkinson disease in Europeans. Arch Neurol. 2007;64(3):425-30.

6.   Kluss JH, Mamais A, Cookson MR. LRRK2 links genetic and sporadic Parkinson's disease.
Biochem Soc Trans. 2019;47:651-61.

7.   Shu L, Zhang Y, Sun Q, Pan H, Tang B. A Comprehensive Analysis of Population Differences in
LRRK2 Variant Distribution in Parkinson's Disease. Front Aging Neurosci. 2019;11:13.

8.   Mata IF, Kachergus JM, Taylor JP, Lincoln S, Aasly J, Lynch T, et al. LRRK2 pathogenic
substitutions in Parkinson's disease. Neurogenetics. 2005;6(4):171-7.

9.   Mata IF, Taylor JP, Kachergus J, Hulihan M, Huerta C, Lahoz C, et al. LRRK2 R1441G in
Spanish patients with Parkinson's disease. Neurosci Lett. 2005;382(3):309-11.

10.   Zabetian CP, Samii A, Mosley AD, Roberts JW, Leis BC, Yearout D, et al. A clinic-based study of
the LRRK2 gene in Parkinson disease yields new mutations. Neurology. 2005;65(5):741-4.

11.   Hatano T, Funayama M, Kubo SI, Mata IF, Oji Y, Mori A, et al. Identification of a Japanese
family with LRRK2 p.R1441G-related Parkinson's disease. Neurobiol Aging. 2014;35(11):2656.e17-.e23.

12.   Takanashi M, Funayama M, Matsuura E, Yoshino H, Li Y, Tsuyama S, et al. Isolated nigral
degeneration without pathological protein aggregation in autopsied brains with LRRK2 p.R1441H
homozygous and heterozygous mutations. Acta Neuropathol Commun. 2018;6(1):105.

13.   Nichols WC, Pankratz N, Hernandez D, Paisan-Ruiz C, Jain S, Halter CA, et al. Genetic
screening for a single common LRRK2 mutation in familial Parkinson's disease. Lancet.
2005;365(9457):410-2.

14.   Di Fonzo A, Rohé CF, Ferreira J, Chien HF, Vacca L, Stocchi F, et al. A frequent LRRK2 gene
mutation associated with autosomal dominant Parkinson's disease. Lancet. 2005;365(9457):412-5.

15.   Hernandez DG, Paisan-Ruiz C, McInerney-Leo A, Jain S, Meyer-Lindenberg A, Evans EW, et al.
Clinical and positron emission tomography of Parkinson's disease caused by LRRK2. Ann Neurol.
2005;57(3):453-6.

16.   Kachergus J, Mata IF, Hulihan M, Taylor JP, Lincoln S, Aasly J, et al. Identification of a novel

LRRK2 mutation linked to autosomal dominant parkinsonism: evidence of a common founder across

European populations. Am J Hum Genet. 2005;76(4):672-80.

17.   Bosgraaf L, Van Haastert PJ. Roc, a Ras/GTPase domain in complex proteins. Biochim Biophys

Acta. 2003;1643(1-3):5-10.

18.   Marin I. The Parkinson disease gene LRRK2: evolutionary and structural insights. Mol Biol Evol.

2006;23(12):2423-33.

19.   Marín I. Ancient origin of the Parkinson disease gene LRRK2. J Mol Evol. 2008;67(1):41-50.

20.   Biskup S, Moore DJ, Rea A, Lorenz-Deperieux B, Coombes CE, Dawson VL, et al. Dynamic and

redundant regulation of LRRK2 and LRRK1 expression. BMC Neurosci. 2007;8:102.

21.   Schulte EC, Ellwanger DC, Dihanich S, Manzoni C, Stangl K, Schormair B, et al. Rare variants

in LRRK1 and Parkinson's disease. Neurogenetics. 2014;15(1):49-57.

22.   Deng J, Lewis PA, Greggio E, Sluch E, Beilina A, Cookson MR. Structure of the ROC domain

from the Parkinson's disease-associated leucine-rich repeat kinase 2 reveals a dimeric GTPase. Proc Natl

Acad Sci U S A. 2008;105(5):1499-504.

23.   Mills RD, Liang LY, Lio DS, Mok YF, Mulhern TD, Cao G, et al. The Roc-COR tandem domain

of leucine-rich repeat kinase 2 forms dimers and exhibits conventional Ras-like GTPase properties. J

Neurochem. 2018;147(3):409-28.

24.   Myasnikov A, Zhu H, Hixson P, Xie B, Yu K, Pitre A, et al. Structural analysis of the full-length

human LRRK2. Cell. 2021;184(13):3519-27 e10.

25.   Tong Y, Yamaguchi H, Giaime E, Boyle S, Kopan R, Kelleher RJ, 3rd, et al. Loss of leucine-rich

repeat kinase 2 causes impairment of protein degradation pathways, accumulation of alpha-synuclein, and

apoptotic cell death in aged mice. Proc Natl Acad Sci U S A. 2010;107(21):9879-84.

26.   Tong Y, Giaime E, Yamaguchi H, Ichimura T, Liu Y, Si H, et al. Loss of leucine-rich repeat kinase

2 causes age-dependent bi-phasic alterations of the autophagy pathway. Mol Neurodegener. 2012;7:2.

27.   Herzig MC, Kolly C, Persohn E, Theil D, Schweizer T, Hafner T, et al. LRRK2 protein levels are

determined by kinase function and are crucial for kidney and lung homeostasis in mice. Hum Mol Genet.

2011;20(21):4209-23.

28.   Tian Y, Lv J, Su Z, Wu T, Li X, Hu X, et al. LRRK2 plays essential roles in maintaining lung

homeostasis and preventing the development of pulmonary fibrosis. Proc Natl Acad Sci U S A.

2021;118(35).

29.   Tong Y, Shen J. Genetic analysis of Parkinson's disease-linked leucine-rich repeat kinase 2.

Biochem Soc Trans. 2012;40(5):1042-6.

30.   Giaime E, Tong Y, Wagner LK, Yuan Y, Huang G, Shen J. Age-Dependent Dopaminergic

Neurodegeneration and Impairment of the Autophagy-Lysosomal Pathway in LRRK Deficient Mice. Neuron. 2017;96(4):796-807.e6.

31.   Huang G, Bloodgood DW, Kang J, Shahapal A, Chen P, Kaganovsky K, et al. Motor Impairments and Dopaminergic Defects Caused by Loss of Leucine-Rich Repeat Kinase Function in Mice. J Neurosci. 2022;42(23):4755-65.

32.   Backman CM, Malik N, Zhang Y, Shan L, Grinberg A, Hoffer BJ, et al. Characterization of a mouse strain expressing Cre recombinase from the 3' untranslated region of the dopamine transporter locus. Genesis. 2006;44(8):383-90.

33.   Rodriguez CI, Buchholz F, Galloway J, Sequerra R, Kasper J, Ayala R, et al. High-efficiency deleter mice show that FLPe is an alternative to Cre-loxP. Nat Genet. 2000;25(2):139-40.

34.   Schwenk F, Baron U, Rajewsky K. A cre-transgenic mouse strain for the ubiquitous deletion of loxP-flanked gene segments including deletion in germ cells. Nucleic Acids Res. 1995;23(24):5080-1.

35.   Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, et al. A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. Nat Neurosci. 2010;13(1):133-40.

36.   Healy DG, Falchi M, O'Sullivan SS, Bonifati V, Durr A, Bressman S, et al. Phenotype, genotype, and worldwide genetic penetrance of LRRK2-associated Parkinson's disease: a case-control study. Lancet Neurol. 2008;7(7):583-90.

37.   Tong Y, Pisani A, Martella G, Karouani M, Yamaguchi H, Pothos EN, et al. R1441C mutation in LRRK2 impairs dopaminergic neurotransmission in mice. Proc Natl Acad Sci U S A. 2009;106(34):14622-7.

38.   Yue M, Hinkle KM, Davies P, Trushina E, Fiesel FC, Christenson TA, et al. Progressive dopaminergic alterations and mitochondrial abnormalities in LRRK2 G2019S knock-in mice. Neurobiol Dis. 2015;78:172-95.

39.   Xing W, Liu J, Cheng S, Vogel P, Mohan S, Brommage R. Targeted disruption of leucine-rich repeat kinase 1 but not leucine-rich repeat kinase 2 in mice causes severe osteopetrosis. J Bone Miner Res. 2013;28(9):1962-74.

40.   Guo L, Girisha KM, Iida A, Hebbar M, Shukla A, Shah H, et al. Identification of a novel LRRK1 mutation in a family with osteosclerotic metaphyseal dysplasia. J Hum Genet. 2017;62(3):437-41.

41.   Sheng Z, Zhang S, Bustos D, Kleinheinz T, Le Pichon CE, Dominguez SL, et al. Ser1292 autophosphorylation is an indicator of LRRK2 kinase activity and contributes to the cellular effects of PD mutations. Science translational medicine. 2012;4(164):164ra1.

42.   Gloeckner CJ, Boldt K, von Zweydorf F, Helm S, Wiesent L, Sarioglu H, et al. Phosphopeptide analysis reveals two discrete clusters of phosphorylation in the N-terminus and the Roc domain of the

Parkinson-disease associated protein kinase LRRK2. J Proteome Res. 2010;9(4):1738-45.

43.   Marchand A, Drouyer M, Sarchione A, Chartier-Harlin MC, Taymans JM. LRRK2 Phosphorylation, More Than an Epiphenomenon. Front Neurosci. 2020;14:527.

44.   Steger M, Tonelli F, Ito G, Davies P, Trost M, Vetter M, et al. Phosphoproteomics reveals that Parkinson's disease kinase LRRK2 regulates a subset of Rab GTPases. eLife. 2016;5:e12813.

45.   Li Y, Liu W, Oo TF, Wang L, Tang Y, Jackson-Lewis V, et al. Mutant LRRK2(R1441G) BAC transgenic mice recapitulate cardinal features of Parkinson's disease. Nat Neurosci. 2009;12(7):826-8.

46.   Lin X, Parisiadou L, Gu XL, Wang L, Shim H, Sun L, et al. Leucine-rich repeat kinase 2 regulates the progression of neuropathology induced by Parkinson's-disease-related mutant alpha-synuclein. Neuron. 2009;64(6):807-27.

47.   Li X, Patel JC, Wang J, Avshalumov MV, Nicholson C, Buxbaum JD, et al. Enhanced striatal dopamine transmission and motor performance with LRRK2 overexpression in mice is eliminated by familial Parkinson's disease mutation G2019S. J Neurosci. 2010;30(5):1788-97.

48.   Melrose HL, Dachsel JC, Behrouz B, Lincoln SJ, Yue M, Hinkle KM, et al. Impaired dopaminergic neurotransmission and microtubule-associated protein tau alterations in human LRRK2 transgenic mice. Neurobiol Dis. 2010;40(3):503-17.

49.   Daher JP, Pletnikova O, Biskup S, Musso A, Gellhaar S, Galter D, et al. Neurodegenerative phenotypes in an A53T α-synuclein transgenic mouse model are independent of LRRK2. Hum Mol Genet. 2012;21(11):2420-31.

50.   Tsika E, Kannan M, Foo CS, Dikeman D, Glauser L, Gellhaar S, et al. Conditional expression of Parkinson's disease-related R1441C LRRK2 in midbrain dopaminergic neurons of mice causes nuclear abnormalities without neurodegeneration. Neurobiol Dis. 2014;71:345-58.

51.   Liu G, Sgobio C, Gu X, Sun L, Lin X, Yu J, et al. Selective expression of Parkinson's disease-related Leucine-rich repeat kinase 2 G2019S missense mutation in midbrain dopaminergic neurons impairs dopamine release and dopaminergic gene expression. Hum Mol Genet. 2015;24(18):5299-312.

52.   Mikhail F, Calingasan N, Parolari L, Subramanian A, Yang L, Flint Beal M. Lack of exacerbation of neurodegeneration in a double transgenic mouse model of mutant LRRK2 and tau. Hum Mol Genet. 2015;24(12):3545-56.

53.   Xiong Y, Neifert S, Karuppagounder SS, Stankowski JN, Lee BD, Grima JC, et al. Overexpression of Parkinson's Disease-Associated Mutation LRRK2 G2019S in Mouse Forebrain Induces Behavioral Deficits and α-Synuclein Pathology. eNeuro. 2017;4(2).

54.   Ramonet D, Daher JP, Lin BM, Stafa K, Kim J, Banerjee R, et al. Dopaminergic neuronal loss, reduced neurite complexity and autophagic abnormalities in transgenic mice expressing G2019S mutant LRRK2. PloS one. 2011;6(4):e18568.

55.   Xiong Y, Neifert S, Karuppagounder SS, Liu Q, Stankowski JN, Lee BD, et al. Robust kinase- 787 and age-dependent dopaminergic and norepinephrine neurodegeneration in LRRK2 G2019S transgenic 788 mice. Proc Natl Acad Sci U S A. 2018;115(7):1635-40.

56.   Zhang P, Fan Y, Ru H, Wang L, Magupalli VG, Taylor SS, et al. Crystal structure of the WD40 790 domain dimer of LRRK2. Proc Natl Acad Sci U S A. 2019;116(5):1579-84.

57.   Greggio E, Zambrano I, Kaganovich A, Beilina A, Taymans JM, Daniels V, et al. The Parkinson 792 disease-associated leucine-rich repeat kinase 2 (LRRK2) is a dimer that undergoes intramolecular 793 autophosphorylation. J Biol Chem. 2008;283(24):16906-14.

58.   Sen S, Webber PJ, West AB. Dependence of leucine-rich repeat kinase 2 (LRRK2) kinase activity 795 on dimerization. J Biol Chem. 2009;284(52):36346-56.

59.   Jorgensen ND, Peng Y, Ho CC, Rideout HJ, Petrey D, Liu P, et al. The WD40 domain is required 797 for LRRK2 neurotoxicity. PloS one. 2009;4(12):e8463.

60.   Dachsel JC, Nishioka K, Vilariño-Güell C, Lincoln SJ, Soto-Ortolaza AI, Kachergus J, et al. 799 Heterodimerization of Lrrk1-Lrrk2: Implications for LRRK2-associated Parkinson disease. Mech Ageing 800 Dev. 2010;131(3):210-4.

61.   Farrer MJ, Stone JT, Lin CH, Dächsel JC, Hulihan MM, Haugarvoll K, et al. Lrrk2 G2385R is an 802 ancestral risk factor for Parkinson's disease in Asia. Parkinsonism Relat Disord. 2007;13(2):89-92.

62.   Kim JM, Lee JY, Kim HJ, Kim JS, Shin ES, Cho JH, et al. The LRRK2 G2385R variant is a risk 804 factor for sporadic Parkinson's disease in the Korean population. Parkinsonism Relat Disord. 805 2010;16(2):85-8.

63.   An XK, Peng R, Li T, Burgunder JM, Wu Y, Chen WJ, et al. LRRK2 Gly2385Arg variant is a risk 807 factor of Parkinson's disease among Han-Chinese from mainland China. Eur J Neurol. 2008;15(3):301-5.

64.   Xie CL, Pan JL, Wang WW, Zhang Y, Zhang SF, Gan J, et al. The association between the 809 LRRK2 G2385R variant and the risk of Parkinson's disease: a meta-analysis based on 23 case-control 810 studies. Neurol Sci. 2014;35(10):1495-504.

65.   Rudenko IN, Kaganovich A, Hauser DN, Beylina A, Chia R, Ding J, et al. The G2385R variant of 812 leucine-rich repeat kinase 2 associated with Parkinson's disease is a partial loss-of-function mutation. The 813 Biochemical journal. 2012;446(1):99-111.

66.   Carrion MDP, Marsicano S, Daniele F, Marte A, Pischedda F, Di Cairano E, et al. The LRRK2 815 G2385R variant is a partial loss-of-function mutation that affects synaptic vesicle trafficking through 816 altered protein interactions. Sci Rep. 2017;7(1):5377.

67.   Kang J, Watanabe H, Shen J. Protocols for assessing neurodegenerative phenotypes in 818 Alzheimer's mouse models. STAR Protocols. 2021;2(3).

68.   Yu H, Kessler J, Shen J. Heterogeneous populations of ES cells in the generation of a floxed

Presenilin-1 allele. Genesis. 2000;26(1):5-8.

69.   Yamaguchi H, Shen J. Histological analysis of neurodegeneration in the mouse brain. Methods

Mol Biol. 2013;1004:91-113.

70.   Nelson EL, Liang CL, Sinton CM, German DC. Midbrain dopaminergic neurons in the mouse:

computer-assisted mapping. J Comp Neurol. 1996;369(3):361-71.

71.   Huang G, Bloodgood DW, Kang J, Shahapal A, Chen P, Kaganovsky K, et al. Motor impairments

and dopamingergic defects caused by loss of LRRK function in mice. J Neurosci. 2022;42(23):4755-65.

72.   Walton WH. Feret's Statistical Diameter as a Measure of Particle Size. Nature.

1948;162(4113):329-30.

Figure 1. Generation and characterization of floxed and deleted *LRRK1* and *LRRK2* alleles

(**A**) Schematic illustrations of human LRRK1 and LRRK2 proteins showing similar functional domains. LRRK1$_{2015}$ protein is derived from exons 2-34 (Ensembl Genome Database: ENSG00000154237). LRRK2$_{2527}$ protein is derived from exons 1-51 (ENSG00000188906). LRR: Leucine rich repeats; Roc: Ras-of-complex; COR: C-terminal of Roc; KIN: Kinase domain.

(**B**) Schematic illustrations of the gene structures of mouse *LRRK1* and *LRRK2*. The boxes in blue are exons that encode the LRRK1 and LRRK2 proteins, and the gray boxes represent the 5' and 3' UTRs. The exons are not drawn in scale. The start codon ATG is in exon 2 of *LRRK1* and exon 1 of *LRRK2*. The exons 27-29 of *LRRK1* and the promoter/exons 1-2 of *LRRK2* are flanked with *loxP* sites (green arrowheads).

(**C**) Targeting strategy for the generation of the targeted, floxed, and deleted *LRRK1* alleles. The red boxes represent the targeted exons 27-29, and the blue boxes represent the untargeted exons. The locations and sizes of the 5' and 3' external probes are shown. The targeting vector contains the 5' and 3' homologous regions (marked by dashed lines) and the middle-targeted region, which includes a *loxP* site (green arrowhead) in intron 26 (1,288 bp upstream of exon 27) and the *PGK-neo* selection cassette flanked by two *FRT* (FLP recognition target) sequences (gray circles) followed by another *loxP* site in intron 29 (1,023 bp downstream of exon 29). A negative selection cassette encoding diphtheria toxin fragment A (DTA) is also included in the targeting vector to reduce ES cells bearing randomly inserted targeting vectors. ES cells carrying the correctly targeted *LRRK1* allele were transfected with *pCAG-FLP* to remove the *PGK-neo* cassette and generate the floxed *LRRK1* allele. Floxed *LRRK1* mice were bred with *CMV-Cre* transgenic mice to generate germline deleted *LRRK1* mice. Detailed strategy for generating targeting vector and DNA sequence of floxed *LRRK1* allele can be found in Figure 1-figure supplement 1 and 2, respectively.

(**D**) Southern analysis of the targeted and floxed *LRRK1* alleles. Genomic DNA from ES cells or mouse tails was digested with *Hind*III and hybridized with the 5' or 3' external probe. For the 5' probe, the resulting 17.0 kb and 4.8 kb bands represent the wild-type (WT) and the targeted (T) or floxed (F) alleles, respectively. For the 3' probe, the resulting 17.0 kb and 12.2 kb bands represent the WT and the floxed (F) alleles, respectively. Detailed Southern strategy can be found in Figure 1-figure supplement 3.

(**E**) Targeting strategy for the generation of the targeted, floxed, and deleted *LRRK2* alleles. The red boxes represent *LRRK2* exons 1 and 2, and the start codon ATG resides in exon 1. The locations and sizes of the 5' and 3' external probes are shown. The targeting vector contains the 5' and 3' homologous regions (marked by dashed lines) and the middle targeted region (from the promoter region to intron 2), which includes a *loxP* site (green arrowhead) upstream (1,768 bp) of the transcription initiation site and the *PGK-neo* selection cassette flanked by two *FRT* sequences (gray circles) and two *loxP* sites (green arrowheads) in intron 2 (878 bp downstream of exon 2). A negative selection cassette encoding DTA is also included in the targeting vector. Mice carrying the correctly targeted *LRRK2* allele were crossed with *Actin-FLP* deleter mice to generate floxed *LRRK2* mice, which were bred with *CMV-Cre* transgenic mice to generate germline deleted *LRRK2* mice. Detailed strategy for generating targeting vector and DNA sequence of floxed *LRRK2* allele can be found in Figure 1-figure supplement 4 and 5, respectively.

(**F**) Southern analysis of the targeted and floxed *LRRK2* alleles. Genomic DNA from ES cells or mouse tails was digested with *Nhe*I and hybridized with the 5' or 3' external probe. For the 5' probe, the resulting 11.5 kb and 3.6 kb bands represent the wild-type (WT) and the targeted (T) or floxed (F) alleles, respectively. For the 3' probe, the resulting 11.5 kb and 5.2 kb bands represent the WT and the floxed (F) alleles, respectively. Detailed Southern strategy can be found in Figure 1-figure supplement 6.

(**G**) Northern analysis of poly(A)+ RNA prepared from the lung of mice carrying homozygous germline deleted (Δ/Δ) *LRRK1* alleles derived from the floxed *LRRK1* alleles using the cDNA probe of exons 2-3 (*left*) and exons 27-29 (*right*). Using the upstream exons 2-3 probe, the *LRRK1* transcripts in wild-type mice are the expected size of ~7.4 kb, whereas the detected *LRRK1* transcripts in *LRRK1* Δ/Δ mice are truncated, consistent with the deletion of exons 27-29 (625 bp), which results in a frameshift and likely degradation of the truncated *LRRK1* mRNA. Using a probe specific for exons 27-29, there is no *LRRK1* transcript in *LRRK1* Δ/Δ mice, as expected. Both blots were hybridized with a *GAPDH* probe as loading controls. Detailed Northern strategy and full-size blots are included in Figure 1-figure supplement 7 and 8, respectively. See also Figure 1-figure supplement 9.

(**H**) Northern analysis of total RNA prepared from the neocortex of mice carrying homozygous germline *LRRK2* deleted (Δ/Δ) alleles using the cDNA probe of exons 1-5 shows the absence of *LRRK2* transcripts. The blot was hybridized with a *GAPDH* probe as a loading control. The full-size blot is included in Figure 1-figure supplement 10. See also Figure 1-figure supplement 11

(**I**) *Left*: Western analysis of wild-type (+/+) and homozygous *LRRK1* Δ/Δ brains show the absence of LRRK1 protein. *Right*: Western analysis of the neocortex of wild-type (+/+) and homozygous *LRRK2* Δ/Δ mice shows the absence of LRRK2 protein. Vinculin was used as a loading control. Full-size blots are included in Figure 1-Source Data 1.

Figure 2. Generation and characterization of DA neuron-specific *LRRK* cDKO mice

(**A**) Immunostaining of GFP and/or TH in the SNpc of *DAT-Cre/+; Rosa26-CAG-LSL-ZsGreen1/+* mice at 2 months of age. Cre recombinase is expressed under the control of the *DAT* endogenous promoter and removes the floxed "stop" cassette, resulting in the expression of EGFP under the control of the ubiquitous *CAG* promoter.

(**B**) Quantification of GFP+/TH+ and TH+ cells shows that 99% of TH+ DA neurons (722 ± 46 TH+ cells per brain) in the SNpc are also GFP+ (713 ± 46 cells per brain), indicating that *DAT-Cre* mediated recombination occurs in essentially all TH+ DA neurons. N = 3 mice, 3 comparable sections per hemisphere, 320 μm apart. Raw quantification data are included in Figure 2-Source Data 1.

(**C**) Similar body weight between *LRRK* cDKO mice and littermate controls at all ages examined ($F_{1,73}$ = 0.0210, p = 0.8853; 2M, 20M, 25M: p > 0.9999; 15M: p = 0.7468, two-way ANOVA with Bonferroni's post hoc multiple comparisons). Raw quantification data are included in Figure 2-Source Data 2.

(**D**) Similar brain weight between *LRRK* cDKO and control mice ($F_{1,61}$ = 0.0440, p = 0.8345; 2M: p = 0.3893; 15M: p > 0.9999; 20M: p > 0.9999; 25M: p = 0.3223, two-way ANOVA with Bonferroni's post hoc multiple comparisons). Raw quantification data are included in Figure 2-Source Data 2.

(**E**) Western analysis of LRRK1 and LRRK2 proteins in the dissected cerebral cortex (CTX) and ventral midbrain (vMB) of *LRRK* cDKO and littermate controls at 2 months of age. Full-size blots are included in Figure 2-Source Data 3.

(**F**) Quantification shows significant decreases of LRRK1 and LRRK2 in the dissected ventral midbrain of *LRRK* cDKO mice (LRRK1, p = 0.0432; LRRK2, p = 0.0162, unpaired two-tailed Student's *t*-test), compared to controls, but not in the dissected cortex of cDKO mice (LRRK1, p = 0.7648; LRRK2, p = 0.2325). Raw quantification data are included in Figure 2-Source Data 4.

The number in the column indicates the number of mice used in the study. Red-filled and open circles represent data obtained from individual male and female mice, respectively. All data are expressed as mean ± SEM. *p < 0.05. Scale bar: 100 μm.

Figure 3. Age-dependent loss of DA neurons in the SNpc of *LRRK* cDKO mice

(**A**) TH immunostaining shows TH+ DA neurons in the SNpc of *LRRK* cDKO and littermate controls at the age of 15, 20, and 24 months. Higher power views of the boxed areas show grossly normal DA neuron morphology in *LRRK* cDKO mice.

(**B**) Quantification of all TH+ DA neurons in the SNpc reveals similar numbers of DA neurons in *LRRK* cDKO mice (10,000 ± 141) and littermate controls (10,077 ± 310, p > 0.9999) at 15 months of age. At 20 months of age, the number of DA neurons in the SNpc of *LRRK* cDKO mice (8,948 ± 273) is significantly reduced, compared to control mice (10,244 ± 220, $F_{1,46}$ = 16.59, p = 0.0002; p = 0.0041, two-way ANOVA with Bonferroni's post hoc multiple comparisons). By 24 months of age, the reduction of DA neurons in the SNpc of *LRRK* cDKO mice (8,188 ± 452) relative to controls (9,675 ± 232, p = 0.0010) is greater, compared to that at 20 months of age. Raw quantification data are included in Figure 3-Source Data 1.

(**C**) Immunohistological analysis of TH and NeuN shows TH+ dopaminergic neurons (green) and NeuN+ neurons (red) in the SNpc of *LRRK* cDKO mice and controls at 24 months of age.

(**D**) Quantification of NeuN+ cells in the SNpc shows that the number of NeuN+ neurons in *LRRK* cDKO mice (17,923 ± 813) is significantly lower than that in control mice (21,907 ± 469, p = 0.0006, Student's *t*-test), indicating loss of neurons in the SNpc of *LRRK* cDKO mice. All TH+ cells are NeuN+. The number of TH+/NeuN+ cells in the SNpc of *LRRK* cDKO mice (10,500 ± 644) is also lower, compared to control mice (14,102 ± 310, p = 0.0001). There is no significant difference in the number of NeuN+/TH- neurons between littermate controls (7,804 ± 249) and cDKO mice (7,423 ± 344, p = 0.3747). Raw quantification data are included in Figure 3-Source Data 2.

The number in the column indicates the number of mice used in the study. Red-filled and open circles represent data obtained from individual male and female mice, respectively. All data are expressed as mean ± SEM. **p < 0.01, ***p < 0.001. Scale bar: 100 μm.

Figure 4. Increases in apoptotic DA neurons in the SNpc of *LRRK* cDKO mice

(**A**) Representative images of TH and active Caspase-3 immunostaining show TH+ DA neurons (green) and active Caspase-3+ apoptotic cells (red) in the SNpc of *LRRK* cDKO and control mice at the age of 24 months.

(**B**) Quantification of active Caspase-3+/TH+ cells shows significant increases of apoptotic DA neurons in the SNpc of *LRRK* cDKO mice (323 ± 38) at 24 months of age, relative to controls (157 ± 8, p = 0.0004, unpaired Student's *t*-test). Raw quantification data are included in Figure 4-Source Data 1.

The number in the column indicates the number of mice used in the study. Red-filled and open circles represent data obtained from individual male and female mice, respectively. All data are expressed as mean ± SEM. ***p < 0.001. Scale bar: 100 μm.

Figure 5. Normal number of TH+ noradrenergic neurons in the LC of *LRRK* cDKO mice

(**A**) Representative images of TH+ noradrenergic neurons in the locus coeruleus (LC) of *LRRK* cDKO mice and littermate controls at 24 months of age.

(**B**) Quantification of TH+ neurons shows similar numbers of TH+ noradrenergic neurons in the LC of *LRRK* cDKO mice (3,350 ± 99) and controls (3,419 ± 86, p = 0.6057, unpaired Student's *t*-test). Raw quantification data are included in Figure 5-Source Data 1.

(**C**) *Top*: Immunostaining of TH and GFP in the LC of *DAT-Cre/+; Rosa26-CAG-LSL-ZsGreen1/+* mice at 2 months of age. There is no GFP+ (green) cell in the LC, indicating that *DAT-Cre* is not expressed in the LC. *Bottom*: Higher power views of the boxed areas.

The number in the column indicates the number of mice used in the study. Red-filled and open circles represent data obtained from individual male and female mice, respectively. All data are expressed as mean ± SEM. Scale bar: 100 μm.

Figure 6. Unchanged number of electron-dense vacuoles in the SNpc of *LRRK* cDKO mice

(**A**, **B**) Representative EM images showing electron-dense vacuoles (arrowheads) in SNpc neurons of cDKO mice and littermate controls at the age of 25 months.

(**C-F**) Higher power views showing various electron-dense vacuoles, autolysosomes (single arrows), autophagosomes (double arrows), and lipid-containing vacuoles (asterisks) in SNpc neurons of littermate control (**C**, **D**) and cDKO (**E**, **F**) mice.

(**G**) *Left*: The average number of electron-dense vacuoles (>0.5 μm in diameter) in the SNpc neuronal profiles per mouse is not significantly different between *LRRK* cDKO mice and littermate controls at the age of 25 months (Control: 6.72 ± 0.43; cDKO: 6.99 ± 0.52, p = 0.6839, unpaired two-tailed Student's *t*-test). Right: The total area of electron-dense vacuoles (>0.5 μm in diameter) in the SNpc neuronal profiles per mouse is similar between *LRRK* cDKO and littermate controls (Control: 4.43 ± 0.44 μm²; cDKO: 4.60 ± 0.49 μm², p = 0.8048). Raw quantification data are included in Figure 6-Source Data 1.

The value in parentheses indicates the number of mice (left) and neuron profiles (right) used in the quantification. Red-filled and open circles represent data obtained from individual male and female mice, respectively. All data are expressed as mean ± SEM. Scale bar: 1 μm.

Figure 7. Elevated microgliosis in the SNpc of *LRRK* cDKO mice

(**A-C**) Representative images of Iba1+ microglia (red, marked by yellow arrowheads) and TH+ dopaminergic neurons (green) in the SNpc of *LRRK* DKO mice and controls at 15 (A), 20 (B), and 24 (C) months of age.

(**D**) Quantification of Iba1+ microglia shows significant increases in the number of Iba1+ microglia in the SNpc of *LRRK* cDKO mice compared to control mice at 15 (control: 1736.7 ± 82.9, cDKO: 2541.3 ± 193.0, p = 0.0017, two-way ANOVA with Bonferroni's post hoc multiple comparisons), 20 (control: 2426.0 ± 67.7, cDKO: 3639.0 ± 127.1, p < 0.0001), and 24 months of age (control: 2640.0 ± 187.1, cDKO: 4088.8 ± 100.3, p < 0.0001). Raw quantification data are included in Figure 7-Source Data 1.

The number in the column indicates the number of mice used in the study. Red-filled and open circles represent data obtained from individual male and female mice, respectively. All data are expressed as mean ± SEM. ****p < 0.0001. Scale bar: 100 μm.

Figure 1-figure supplement 1. Generation of the *LRRK1* targeting vector

1) The <u>left middle</u> arm was amplified by PCR from mouse BAC DNA (Clone #: RP23-213J23, BACPAC Resources Center) using primers P3 and P4. The resulting PCR product (2,079 bp) containing part of *LRRK1* intron 26, exon 27, and part of intron 27 was subcloned into the pGEM-T Vector (Promega, Cat#; A1360) by TA ligation to generate pLRRK1#1 (pLM1).

P3: 5'CCAGTCACTTCTCCACCTCAGGGAAAATGG (30 bp)

P4: 5'CCTTGT*GGTACC*CGGACCTTCTATCACCTTTATCC (35 bp): *Kpn*I (*GGTACC*) is an endogenous restriction site.

2) The <u>right middle</u> arm was amplified from mouse BAC DNA (Clone #: RP23-213J23) using primers P5 and P6. The resulting PCR product (3,403 bp) containing part of *LRRK1* intron 27, exon 28, intron 28, exon 29, and part of intron 29 was subcloned into the pGEM-T Vector by TA ligation to generate pLRRK1#2 (pLM2).

P5: 5'GGTCCG*GGTACC*ACAAGGTGCTGGTTAAGTGCC (33 bp): *Kpn*I (*GGTACC*) is an endogenous restriction site.

P6: 5'AGCAGACCTCTTGCCTTCTACTACTGACTG (30 bp)

3) pLRRK1#1 (pLM1) was digested with *Not*I and *Kpn*I, then subcloned into the *Not*I and *Kpn*I sites of pLRRK1#2 (pLM2) to generate the pLRRK1#3 (pLM3) plasmid which contains the <u>middle</u> homologous region.

4) The middle homologous region was released from pLRRK1#3 (pLM3) using *Not*I and *Sac*II followed by Klenow to blunt the sticky ends, which was then subcloned into the *Sma*I site of pGKneoF2L2DTA (addgene: #13445) to generate pLRRK1#4 (pLM4).

Green arrowhead: *loxP* site, gray circle: *FRT* site

5) The left arm was amplified from mouse BAC DNA (Clone: RP23-213J23) using primers P1 and P2. The resulting PCR product (2,016 bp) containing part of *LRRK1* intron 25, exon 26, and part of intron 26 was digested with *Sac*II and *Not*I, and was then subcloned into the *Sac*II and *Not*I sites of pGKneoF2L2DTA (addgene: #13445) to generate pLRRK1#5 (pLM5).

P1: 5'gacat*CCGCGG*CACCATGTGAGTGGCAGCTGTGGTGAGAAC (41 bp). *Sac*II (*CCGCGG*) is an exogenous site.

P2: 5'gacat*GCGGCCGCAAGCTT*TTTAATAGCCGTTCTTTCTTAGAGAAGGCAG (50 bp). *Not*I (*GCGGCCGC*) and *Hind*III (*AAGCTT*) are exogenous sites.

6) The right arm was amplified from mouse BAC DNA (Clone: RP23-213J23) using primers P7 and P8. The resulting PCR product (3,131 bp) containing part of *LRRK1* intron 29, exon 30, and part of intron 30 was digested with *Sal*I and *Hind*III, and was then subcloned into the *Sal*I and *Hind*III sites of pGKneoF2L2DTA (addgene: #13445) to generate pLRRK1#6 (pLM6).

P7: 5'gacat*GTCGAC*GGATCCGTAGGGAAGACCCACTAGGAGGAAGAAAG (46 bp). *Sal*I (*GTCGAC*) is an exogenous site.

P8: 5'gacat*AAGCTT*TGGTACCTTTCTAAAGGCAGCATTTTGCTTGC (43 bp). *Hind*III (*AAGCTT*) is an exogenous site.

7) pLRRK1#5 (pLM5) was digested with *Sac*II and *Not*I, and was then subcloned into the *Sac*II and *Not*I sites of pLRRK1#6 (pLM6) to generate pLRRK1#7 (pLM7).

8) The middle homologous region was released from pLRRK1#4 by *Not*I and *Sal*I and was then subcloned into the *Not*I and *Sal*I sites of pLRRK1#7 (pLM7) to generate the final targeting vector pLRRK1#8 (pLM8), which was linearized with *Xho*I before electroporation into ES cells.

Restriction sites in black: from vectors

Restriction sites in red: *LRRK1* endogenous sites

Restriction sites in blue: introduced by the primers

Figure 1-figure supplement 2. Genomic DNA sequence of the floxed *LRRK1* allele

Figure 1-figure supplement 3. Southern strategy for the targeted and floxed *LRRK1* alleles

5' external probe: 377 bp (*LRRK1* intron 25)

P15: 5'CAGAATACCCCATGCTGGGGAATTGC (26 bp)

P16: 5'CCGTTTCTAGGATTCTAATTTTTC (24 bp)

*Hind*III digestion   *WT: 17.0 kb*
                               Targeted: 4.8 kb
                               Floxed: 4.8 kb

3' external probe: 305 bp (*LRRK1* exon 32)

P17: 5'AACTCCTTCCTGGTGCTGGCAGGCCTGGCTG (31 bp)

P18: 5'ACAAGTGACGTGACCATGGACGGAGCTGCG (30 bp)

*Hind*III digestion   *WT: 17.0 kb*
                               Targeted: 14.1 kb
                               Floxed: 12.2 kb

Neo **probe (363 bp)**

P23: 5'ATTCGGCTATGACTGGGCAC (20 bp)

P24: 5'GACCACCAAGCGAAACATCG (21 bp)

*Hind*III digestion   *WT and Floxed: no band*
                          Targeted: 14.1 kb

Figure 1-figure supplement 4. Generation of the *LRRK2* targeting vector

1) The left homologous region was amplified from mouse BAC DNA (Clone# RP23-526-A2, BACPAC Resources Center) using primers P9 and P10. The resulting PCR product (2,579 bp) was digested with *EcoR*I, which is an endogenous restriction site upstream of the *LRRK2* promoter, and was then subcloned into the *EcoR*I and *Sma*I sites of pBSK (+) to generate pLRRK2#1 (2,526 bp + 2,961 bp = 5,487 bp).

P9: 5' GAACACACAAGGCTATGGCTATTGTC (26 bp)

P10: 5' GTAGGACTATCATCCACCTGTAGGACTCC (29 bp)

2) Oligoes (P39 and P40) were annealed together to generate a double stranded polylinker (*BamH*I-*loxP*-*Nhe*I-*Spe*I), which was then subcloned into the *BamH*I and *Spe*I sites of pLRRK2#1 to generate pLRRK2#2 (5,487 bp + 50 bp = 5,537 bp).

P39: 5' GATCCATAACTTCGTATAATGTATGCTATACGAAGTTATGCTAGCA (46 bp).
*BamH*I (G'GATCC), *loxP* top strand (ATAACTTCGTATAATGTATGCTATACGAAGTTAT), *Nhe*I (G'CTAGC).

P40: 5' CTAGTGCTAGCATAACTTCGTATAGCATACATTATACGAAGTTATG (46 bp).
*Spe*I (A'CTAGT), *Nhe*I (GCTAGC), *loxP* bottom strand (ATAACTTCGTATAGCATACATTATACGAAGTTAT).

Green arrowhead: *loxP* site

3) The middle arm was amplified from mouse BAC DNA (Clone# RP23-526-A2) using primers P11 (upstream of the *LRRK2* promoter) and P12 (intron 2). The resulting PCR product (2,990 bp) was digested with *Xba*I and *Not*I and was then subcloned into the *Xba*I and *Not*I of pGEM-T Vector (Promega, Cat#: A1360) to generate pLRRK2#3 (2,990 bp + 3,000 bp = 5,990 bp). *Xba*I, *Not*I, and *Nhe*I sites are exogenous sequences introduced into PCR primers.

PCR: P11=5' GTCCTACAGGTGGATTCTAGACCTACAAGG 3' (30 bp) — *Xba*I

P12=5' GACGCGGCCGCGTTTAGTAGCTAGCATGACTATGAAGGAG 3' (40 bp) — *Not*I, *Nhe*I

4) pLRRK2#3 was digested with *Xba*I and *Not*I, and the insert was then subcloned into the *Xba*I and *Not*I sites of pLRRK2#2, which contains the left homologous arm and the *loxP* sequence, to generate pLRRK2#4 (2,958 bp + 5,490 bp = 8,448 bp).

5) The right arm was amplified from mouse BAC DNA (Clone# RP23-526-A2) using primers P13 and P14. The resulting 3,503 bp PCR fragment containing *LRRK2* intron 1, exon 2, and intron 2, was subcloned into the *EcoR*V site of pBSK(+) to generate pLRRK2#5 (3,503 bp + 2,961 bp = 6,464 bp). **Check direction!!!**

PCR: P13=5'GCACTTGAGTCTTAATCTTGGGCAC 3' (25 bp)

P14=5'CATTCGAGCAGCTAAGCCTGTAATC 3' (25 bp)

6) pLRRK2#5 was digested with *BamH*I and then treated with Klenow to blunt the overhang followed by *Cla*I digestion. The digested fragment was subcloned into the *EcoR*V and *Cla*I sites of the pSoriano vector (*PGKneolox2DTA*, addgene 13443) to generate pLRRK2#6 (3,547 bp + 6,307 bp = 9,854 bp).

7) The LFNT-tk/pBS plasmid (a gift from S. Tonegawa) was digested with *Sac*II (followed by Klenow) and *Not*I (followed by *Ssp*I) to release the "*loxP-FRT-PGK-neo-loxP-FRT*" fragment (2,928 bp), which was then subcloned into the *Xba*I (followed by Klenow) and *Not*I sites of pLRRK2#6 to generate pLRRK2#7 (2,928 bp + 7,051 bp = 9,979 bp).

Green arrowhead: *loxP* site, gray circle: *FRT* site

8) pLRRK2#4 was digested with *EcoR*V and *Not*I to release the fragment containing the left arm-*loxP*-middle arm, which was then subcloned into the *Ale*I and *Not*I sites of pLRRK2#7 to obtain pLRRK2#8, the *LRRK2* targeting vector (15,485 bp). The linearized targeting vector (with *Ahd*I) was electroporated into ES cells.

Restriction sites in black: from vectors

Restriction sites in red: *LRRK2* endogenous sites

Restriction sites in blue: introduced by the primers

Figure 1-figure supplement 5. Genomic DNA sequence of the floxed *LRRK2* allele

Figure 1-figure supplement 6. Southern analysis of the targeted and floxed *LRRK2* alleles

5' external probe (753 bp)

P19= 5'CATGGGAGAGAGGGTTTCTCACTTACT (27 bp)

P20= 5'CTTGGACAGCATTGTCAGCCTAGAC (25 bp)

***Nhe*I digestion**　　　　　　　　　*WT: 11.5 kb*
　　　　　　　　　　　　　　　　　Targeted and floxed: 3.6 kb

3' external probe (622 bp)

P21=5' GCCAAGCAGTTATTGATGCTGTAGC (31 bp)

P22= 5' CGAGCTGTAAGATGAGCTGGGTACT (30 bp)

***Sph*I digestion**　　　　　　　　　*WT: 12.3 kb*
　　　　　　　　　　　　　　　　　Targeted: 4.5 kb

***Nhe*I digestion**　　　　　　　　　*WT: 11.5 kb*
　　　　　　　　　　　　　　　　　Floxed: 5.2 kb

Southern blot of targeted *LRRK2* allele

Figure 1-figure supplement 7. Northern strategy for *LRRK1* mRNA

***LRRK1* exons 2-3 probe (377+4=383 bp)**

P31: 5' CAGGATGAGCGTGTGTCTGCAG (CAG exon 1/rest exon2)

P32: 5′ CCTTCTCCTGTGAGGATTCGCTCT (C exon 3/rest exon 4)

WT and floxed alleles: ~7.4 kb

Deleted allele: ~6.8 kb*

* Deletion of *LRRK1* exons 27-29 (625 bp; exon 27: 278 bp, exon 28: 195 bp, exon 29 152 bp) results in a frameshift of downstream exons, likely resulting in degradation of the transcript.

***LRRK1* exons 27-29 probe (550 bp)**

P33: 5' CTGGCCTACCTGCACAAGAA (exon 27)

P34: 5' CCTTCCCATCCCAGAACACC (exon 29)

WT and floxed alleles: 7.4 kb

Deleted allele: no band

***GAPDH* probe (452 bp)**

P37: 5' ACCACAGTCCATGCCATCAC (exon 5)

P38: 5′ TCCACCACCCTGTTGCTGTA (exon 7)

Figure 1-figure supplement 8. Northern analysis of *LRRK1* mRNA in mice carrying *LRRK1* germline deleted alleles

LRRK1 Northern blots (polyA+ RNA from the lung) using exons 2-3 or exons 27-29 probe: The exons 2-3 probe detected the full-length *LRRK1* transcripts (~7.4 kb) in wild-type (+/+) mice and reduced size transcripts in *LRRK1* Δ/Δ mice, likely due to the absence of exons 27-29 (625 bp), which was confirmed by sequencing of RT-PCR products (see Figure1-figure supplement 9). The truncated *LRRK1* mRNA lacking exons 27-29 is present in lower levels, likely due to mRNA degradation by nonsense-mediated decay, as the absence of exons 27-29 (625 bp) results in a frameshift of downstream codons.

The exons 2-3 probe also recognizes the fusion transcript *LRRK1-lacZ* (~4.6 kb) in *LRRK1/2* DKO mice, which contains *LRRK1* exons 1-3 and exon 2 of engrailed, *IRES* and *lacZ* cDNA (Figure 1, Figure S1 in Giaime et al., *Neuron*, 2017). The exons 27-29 probe also recognizes the full-length *LRRK1* transcripts (~7.4 kb) in wild-type (+/+) mice but no truncated *LRRK1* transcripts, due to the germline deletion of the genomic region from intron 26- intron 29, lacking exons 27-29. The exons 27-29 probe also did not detect the *LRRK1-lacZ* fusion transcript, which only contains exons 1-3, in *LRRK1/2* DKO mice. While polyA enrichment reduced levels of 18S and 28S rRNAs, they are still present in the lanes/samples.

Figure 1-figure supplement 9. RT-PCR analysis of the deleted *LRRK1* allele

RT-PCR analysis using a *LRRK1* exon 32-specific primer P47 (5'GGCTCAGGTCATGCTCAGTT) for RT and PCR primer sets in *LRRK1* exons 4-8, 11-17, 20-25, and 25-31 indicates normal splicing in homozygous floxed *LRRK1* (F/F) mice and properly deleted transcripts in *LRRK1* Δ/Δ mice. The identity of all PCR products was confirmed by sequencing.

P53: 5' TTTTGGACACGCCGAAGTAGT (**exon 4**)

P54: 5' AGCCGCTCCAGGTAGTTTTT (**exon 8**)

P57: 5' GGACCTCTCCAGAAACCAGC (**exon 11**)

P58: 5' GCAGGGTTGCTATCCTCTCC (**exon 17**)

P63: 5' GCGGTCAGTGGCAAAGAATG (**exon 20**)

P64: 5' AATGCTGTTCTCACCCTCCG (**exon 25**)

P41: 5' GAATTCTGCTAATGCCCCAGC (**exon 25**)

P42: 5' AGGCTGTAGATATAGATTTTCTGGT (**exon 31**)

The 990 bp product containing exons 25-31 represents the WT allele and confirms the floxed allele. The 365 bp product containing exons 25, 26, 30, and 31 confirms the deletion of exons 27-29.

Figure 1-figure supplement 10. Northern analysis of *LRRK2* mRNA in mice carrying floxed or germline deleted *LRRK2* alleles

LRRK2 exons 1-5 probe (437 bp)

P35= 5' AGGAAGGCAAGCAGATCGAG (exon 1, 20 bp)

P20= 5' GGCTGAATATCTGTGCATGGC (exon 5, 22 bp)

WT and floxed alleles: 8.3 kb
Deletion: no band

GAPDH probe (452 bp)

P37= 5' ACCACAGTCCATGCCATCAC (exon 5, 20 bp)

P38= 5' TCCACCACCCTGTTGCTGTA (exon 7, 20 bp)

LRRK2 Northern blot (total RNA from the brain) using exons 1-5 probe: The probe detected the full-length *LRRK2* transcript (~8.3 kb) in wild-type (+/+) and *fLRRK2/fLRRK2* mice, and no transcript in *LRRK2* Δ/Δ mice, derived from floxed *LRRK2* alleles, and previously generated *LRRK1/2* DKO mice (Giaime et al., *Neuron*, 2017).

Figure 1-figure supplement 11. RT-PCR analysis of the deleted *LRRK2* allele

RT-PCR analysis using an *LRRK2* exon 51-specific primer P93 (5'TCGTGTGGAAGATTGAGGTCC) for RT, and primers in *LRRK2* exons 1 and 5, P35 and P20, for PCR, which shows normal splicing in *fLRRK2/fLRRK2* brains and the absence of PCR products in *LRRK2* Δ/Δ brains. The identity of the PCR product in +/+ and *fLRRK2/fLRRK2* brains was confirmed by sequencing: The 437 bp product contains *LRRK2* exons 1-5.

P35= 5' AGGAAGGCAAGCAGATCGAG (exon 1)

P20= 5' GGCTGAATATCTGTGCATGGC (exon 5)