Genomic analyses of Asiatic Mouflon in Iran provide insights into the

domestication and evolution of sheep

Dong-Feng Wang[1,2,3], Pablo Orozco-terWengel[4], Meng-Hua Li[1*], Feng-Hua Lv[1*]

[1]College of Animal Science and Technology, China Agricultural University, Beijing,

China

[2]CAS Key Laboratory of Animal Ecology and Conservation Biology, Institute of

Zoology, Chinese Academy of Sciences (CAS), Beijing, China

[3]College of Life Sciences, University of Chinese Academy of Sciences (UCAS),

Beijing, China

[4]School of Biosciences, Cardiff University, Cardiff, UK

*lvfenghua@cau.edu.cn, menghua.li@cau.edu.cn

Abstract

The wild ancestor of domestic sheep, Asiatic mouflon (*O. gmelini*), consists of several

subspecies mainly distributed in Iran nowadays. However, the impact of the Asiatic

mouflon subspecies in Iran on sheep domestication remains unclear. Here, we

conducted a comprehensive population genomics analysis of Asiatic mouflons in Iran

with 780 whole-genome sequences, 767 whole mitogenomes, and 239 Y

chromosomes. Whole-genome sequence analyses revealed two subpopulations of the

Asiatic mouflons in Iran: *O. gmelini*_2 limited in Kabudan and *O. gmelini*_1 over a

wide geographic area. Phylogenetic analyses of the Asiatic mouflons in Iran based on

uniparental variants revealed a monophyletic lineage with the mitochondrial

haplogroups C+E, and clustered into a monophyletic with Y-chromosomal lineage

HY2 of sheep. Additionally, introgression tests detected significant signals of genetic

introgression from *O. gmelini*_2 to four sheep populations (e.g., Garut, Garole,

Bangladeshi, and Sumatra) in South and Southeast Asia. In the four sheep

populations, selective tests and introgression signals revealed that the wild

introgression could have contributed to their small body size and local adaptation to

the hot and humid environments in the Indian Peninsula. Our results suggested that

the maternal haplogroups C+E and paternal lineage HY2 could have originated from

the Asiatic mouflon in Iran. Also, our findings provide new insights into sheep

domestication and subsequent introgressions from wild ancestors to domestic

populations.

Key words: Iranian Asiatic mouflon, sheep, whole-genome sequences, uniparental

lineage, domestication, introgression

Introduction

Archeozoological and genetic evidence suggested that Asiatic mouflon (*O. gmelini*),

the wild ancestor of sheep, have underwent a period of multigenerational human

management in southeastern Anatolia by ~10,500 years B.P. [1–4]. Following

domestication, sheep diffused throughout Eurasia with human migrations and

developed into different populations under natural and artificial selections[5–8]. Asiatic

mouflon ranges over a wide geographic area in Armenia, southern Azerbaijan,

Cyprus, northern, southern, and western regions of Iran, and eastern and central

regions of Turkey. It has been classified into five subspecies based on morphological

differences and geographic distributions[99,10], such as Armenian mouflon (*O. gmelini*

gmelini)[11], Anatolian mouflon (*O. gmelini anatolica*)[11], Isfahan mouflon (*O. gmelini*

isphahanica), Laristan mouflon (*O. gmelini laristanica*), and Cyprus mouflon (*O.*

gmelini ophion)[11]. Due to the complex intraspecies taxonomy of Asiatic mouflon, the

genetic contributions of various subspecies to the domestication of sheep remains

unclear[3,12–14].

Previous genetic studies revealed controversial ancestral origins of domestic sheep.

Hiendleder et al. (2002) proposed that sheep were domesticated from two different

subspecies in Turkey and western Iran, corresponding to mtDNA lineages A and B of

domestic sheep, respectively[13]. Through genetic and phylogeographic analyses of

partial mtDNA sequences of modern and ancient samples in wild and domestic sheep,

Demirci et al. (2013) suggested that domestic sheep could have originated from two

distinct ancestral populations of mtDNA lineages A+B+D and C+E of domestic

sheep, respectively[3]. Based on genetic, archeozoological, and geographic evidence of

212 Asiatic mouflon samples covering most of their distribution area, a recent study

suggested that the maternal origin of sheep could be traced to a subspecies of *O.*

gmelini gmelini in eastern Anatolia and central Zagros[4].

Despite the intensive focus on material inheritance, Y-chromosome and whole-

genome analyses have received less attention. Recently, Y-chromosome and whole-

genome analyses of Asiatic mouflon in Iran revealed that they could have diverged

from the direct ancestor of domestic sheep, and showed genetic introgression into

domestic sheep after domestication[8,14,15]. Also, these studies have demonstrated that

genomes of Asiatic mouflon in Iran are crucial for the development of morphological

and adaptive characteristics in domestic sheep[8,15,16]. Nevertheless, the impact and

extent of genetic introgression involving the Asiatic mouflon subspecies in Iran (e.g.,

O. gmelini gmelini, *O. gmelini isphahanica*, and *O. gmelini laristanica*) and their

hybrid populations (*O. gmelini gmelini* × *O. vignei*) on the genomes of domestic

sheep remains unclear.

Here, we comprehensively analyzed whole genome sequence, complete and partial

mtDNA sequences, and Y-chromosome genetic variants in domestic sheep and

Asiatic mouflon, covering a large distribution area in Asia, Europe, Africa, and

America. We aimed to elucidate the contributions of various Asiatic mouflon

subspecies in Iran to the domestication of sheep. Additionally, we investigated the

impact of wild introgression of the Asiatic mouflon subspecies in Iran on the genomes

of domestic sheep populations in South and Southeast Asia, which have adapted to the

hot and humid environments near the Indian Ocean.

Materials and Methods

Datasets

Whole-genome sequences (WGS) of 780 individuals representing 158 domestic sheep

populations and three wild sheep species (32 Asiatic mouflon (*Ovis gmelini*) in Iran, 9

urial (*Ovis vignei*), and 1 bighorn sheep (*Ovis canadensis*)) were obtained from

publicly available data (Supplementary Table S1). Besides these, we retrieved 161

whole mitogenomes from the NCBI (Supplementary Table S2). Also, we included

333 Cytochrome b (*Cytb*) sequences of Asiatic mouflon in the integrated analysis

(Supplementary Table S3).

Variant calling

We implemented quality-based trimming for WGS data with Trimmomatic v0.39 [17]

and mapped high-quality reads to the reference genome

Oar_rambouillet_v1.0_CM022046.1. (https://www.

ncbi.nlm.nih.gov/assembly/GCF_002742125.1/;

https://www.ncbi.nlm.nih.gov/nuccore/CM022046.1/, last accessed May 12, 2020)

using the Burrows–Wheeler aligner (BWA mem) v.0.7.8[18] with the default

parameters. Then, we removed duplicates with the *MarkDuplicates* module in GATK

v 4.1.2.0[19] and called SNPs following the previous procedures[15]. We used the

HaplotypeCaller module in GATK v 4.1.2.0 to generate the GVCF file of each

sample and merged the raw GVCF file of each individual with the *CombineGVCFs*

module. Afterward, we genotyped SNPs using the *GenotypeGVCFs* module and

filtered SNPs using the *VariantFiltering* module of GATK with the parameters

"QUAL < 30.0 || QD < 2.0 || MQ < 40.0 || FS > 60.0 || SOR > 3.0 || MQRankSum < -

12.5 || ReadPosRankSum < -8.0". We obtained a total of 121,252,917 high-quality

autosomal SNPs and 203,938 Y-chromosomal SNPs for the following analyses.

Mitochondrial DNA assembly

We assessed the mitogenome depth of each sample using the SAMtools v.1.9[20] and

kept only the samples with depth > 100 in the below analyses. We assembled the

complete mitogenomes using the following protocols. We first converted BAM files

to the fastq format. Then, we generated the mitogenome consensus sequence from the

fastq files using MIA v1.0 (https://github.com/mpieva/mapping-iterative-assembler).

In addition to the 605 mitogenomes assembled here, we also included 161 whole

mitogenome sequences retrieved from the NCBI GenBank in the analysis, with a final

mitogenome dataset of 767 sequences from 727 domestic sheep, 31 Iranian Asiatic

mouflon, and 9 urial. All sequences were aligned using MAFFT v7.515[21] and

trimmed with trimAl v1.4[22].

Y-chromosomal SNPs

To obtain single-copy Y-chromosomal SNPs, raw SNPs in the 780 samples were

processed as follows. Firstly, we identified ram samples from the whole samples

using PLINK v1.90b6.26[23] with the "--check-sex" option and kept only these ram

samples in the following analyses. We then excluded SNPs in the regions covering the

pseudoautosomal regions (PARs) and the highly repetitive sequences (around 1.06

Mb)[14]. Afterwards, we filtered SNPs with the mean sequencing depth ranging from

$0.5 \times$ to $2 \times$ of the expected depth (half of the genome-wide depth)[14]. Further, we

removed SNPs that were heterozygous in at least one of the male samples. Finally, we

retained 4,096 Y-chromosome SNPs for the haplotype and phylogenetic analyses.

Population structure analysis

To reveal the genetic differentiation within the Asiatic mouflon in Iran, we examined

the population genetic substructure for the total 779 wild and domestic samples using

a set of autosomal SNPs filtered as follows. First, we filtered the total 25,270,188

high-quality SNPs with missing rates < 0.1 and minor allele frequencies > 0.05

(MAF > 0.05) using PLINK v1.90b6.26[23]. Then, we pruned the remaining SNPs with

the options "--indep-pairwise 50 10 0.1" of PLINK v1.90b6.26[23]. Afterwards,

1,566,300 SNPs were retained for the population genetic structure analysis.

We computed the identity by state among samples using PLINK v1.90b6.26[23] with

option "distance square 1-ibs". Next, we generated a neighbor-joining (NJ) tree using

the function "bionj" in the R package *ape* v5.6.2[24]. The tree was rooted using urial (*O.*

vignei) as the outgroup, and its graphical representation was obtained using the R

package *ggtree* v3.6.2[25]. Principal component analyses (PCA) were performed using

the "--pca" option in PLINK v1.90b6.26[23] with the default settings. Furthermore,

model-based clustering was performed using the sNMF v1.2 with clusters K from 2 to

10[26].

The F_{ST} and nucleotide diversity (π) of autosomes were calculated with vcftools

v0.1.17[27] using 2Mbp windows and 1Mbp steps, and P value was calculated with

1000 permutation test. The F_{ST} of mitochondria and Y chromosome were calculated

with R package hierfstat v0.5.11[28] with 1000 permutation test. The haplotype

diversity of mitochondria and Y chromosome were calculated with pegas v1.1[29] with

1000 permutation test.

Phylogenetic analysis

Phylogenetic analyses based on mitogenome and Y-chromosomal SNPs were

performed using the maximum likelihood (ML) algorithm of iqtree2 v2.0.7[30].

Mitogenome sequences were aligned using mafft v7.515[21] with the default setting,

and multiple sequence alignment (MSA) was trimmed to remove ambiguously aligned

regions using trimAl v1.4[22] with parameters "-automated1". To keep high-quality

SNPs, we further removed all the sites with gaps and ambiguous nucleotide bases and

only kept parsimony informative sites using *ClipKIT*[31]. We tested the potential models

of DNA sequence evolution using the option "-m" test in iqtree2 v2.0.7[30] and found

the generalized time reversible model (GTR + G)[32] was the best fitting substitution

model for mitogenome. We constructed a maximum likelihood (ML) tree with the site

model GTR + G with 1000 bootstrap replicates using iqtree2 v2.0.7[30]. The resulting

tree was rooted with bighorn as the outgroup and visualized using the FigTree v1.4.4

(http://tree.bio.ed.ac.uk/software/figtree/). Mitogenome haplotypes were inferred with

the function "haplotype" in the R package *pegas* v1.1[29]. We also constructed the

parsimony mitogenome haplotype network using POPART[33]. We transformed the

genotypes of Y-chromosomal 4,096 SNPs into the fasta format using vcf2phylip.py[34].

Phylogenetic trees were built and visualized following the same pipelines and using

the same programs and substitution model as above.

Demographics and divergence time inference

The population genetic structure analyses above identified two subpopulations (*O.*

*gmelini*_1 and *O. gmelini*_2) in the Asiatic mouflons in Iran. We inferred the dynamic

changes in historical and recent effective population size (N_e) of the two

subpopulations using SMC++ v1.15.2 algorithms[35]. We selected 7 - 10 high-depth

individuals in each of the six geographically differentiated genetic groups of domestic

sheep (Central-and-East Asian, South-and-Southeast Asian, the Middle Eastern,

African, European, and American populations)[15] and the two Asiatic mouflon

subpopulations. We estimated the N_e during the past 1,000 to 100,000 years for each

group using SMC++ v1.15.2[35] under the default setting.

For the SMC++ analysis, the reference genome's low-complexity regions were

masked using snpable (http://lh3lh3.users.sourceforge.net/snpable.shtml, last accessed

October 20, 2022) using in-house scripts. We set the mutation rate and generation

time to be 1.00×10^{-8} per nucleotide site per generation[36] and 3 years[37,38], respectively.

To infer the divergence times between *O. gmelini*_1 and *O. gmelini*_2 as well as

between the Asian mouflon and domestic sheep, we employed the split option of

SMC++ with the same parameter settings as above.

Additionally, we inferred the demographic history of domestic sheep and Asiatic

mouflon based on mitogenomes and Y-chromosome data using Beast v2.7.3[39] and the

Coalescent Bayesian Skyline plot[40]. Firstly, we used a coalescent model to estimate

the mutation rates for mitogenome and Y chromosome chromosomes. The analysis

was performed using the GTR + GAMMA (4) + I site model and the strict clock

model. The most recent common ancestor (MRCA) of Asiatic mouflon and domestic

sheep was set to be 0.653 million years[41]. The chain length was set to be 40,000,000

with a burn-in of 2,000,000. The results were analyzed using Tracer

(http://beast.community/tracer), assuming mutation rates estimates of 2.10×10^{-8} and

7.70×10^{-9} substitutions per site per year for the mitochondrial and Y chromosome

SNPs, respectively.

Subsequently, we utilized the mutation rates estimated above to infer the demographic

history of maternal and paternal lineages in domestic sheep and Asiatic mouflon. We

used the same GTR + GAMMA (4) + I site model as above, with the clock model set

as strict and the mutation rates of 2.10×10^{-8} and 7.70×10^{-9} per site per year for

mitochondria and Y-chromosome, respectively. The prior model remained the

Coalescent Bayesian Skyline, with the chain length of 40,000,000, a burn-in of

2,000,000, and samples drawn every 4,000 steps. The obtained results were further

analyzed using Tracer, and each analysis yielded over 200 effective samples for all

relevant parameters.

Introgression analysis

We conducted introgression analysis to clarify the genetic contributions of the two

Asiatic mouflon subpopulations to South-and-Southeast Asian (SSA) sheep. We

tested introgression using a combination method of D and f_{dM} statistics. Specifically,

we calculated D statistics[42] with a four taxon model ((((P1, P2), P3), P4) using Dsuite

v0.5r50[43]. In this analysis, Menz sheep (MEN) was selected as the reference

population (P1), while SSA populations were the target population (P2), *O. gemlini_1*

or *O. gemlini_2* was the donor (P3), and *O. canadensis* was the outgroup (P4). A *P*

value < 0.05 from a standard block-jackknife procedure was considered as evidence

of gene flow from P3 to P2.

We further implemented sliding window statistics analysis using the Dsuite v0.5r50[41]

to quantify the introgressions. We calculated f_{dM} values using a sliding window of 50

SNPs with a step of 25 SNPs across the genomes with the defined trios (MEN, SSA,

O. gmelini_2), where SSA represents the four South-and-Southeast Asian sheep

populations suggested to be the introgressed populations by the D-statistics and f_4-

ratio tests. We calculated the Z-transformed f_{dm} values, and the windows with $P <$

0.05 and $D > 0$ was defined as the potential introgressed genomic regions. We

evaluated the statistical significance using one-tailed Z-test in R package BSDA

v1.2.1[44].

Selective sweep tests

To assess the potential roles of genomic introgression from Iranian Asiatic mouflons

in Southern and Southeastern Asian (SSA) sheep populations, we conducted a PBS

analysis[45] to identify selective signatures in four introgressed populations (i.e., Garut

(GUR), Garole (GAR), Bangladeshi (BGE), and Sumatra (SUM)). We pooled these

four populations (GUR, BGE, SUM, and GAR) as an introgressed group, and used the

MEN and *O. gmelini_2* as the reference populations. We calculated the PBS values

using PBScan v2020.03.16 with parameters "-win 50 -step 25" described in Hämälä

and Savolainen (2019) [46] and estimated P values by 2000 permutation cycles for

Monte Carlo testing. and detected selective regions with top 1% PBS values.

Subsequently, we identified the overlaps between the selective and the introgressed

regions using BEDtools v2.30.0[47] and annotated functional genes in the overlapping

genomic regions.

Gene annotation and function analyses

We annotated genes in the introgressed regions using the sheep reference assembly

Oar_rambouillet_v1.0 by BEDtools v2.30.0[47]. We also used the sheepQTLdb

(https://www.animalgenome.org/cgi-bin/QTLdb/OA/index)[48] to detect QTLs

associated with the introgressed regions. We examined the LD blocks by

LDBlockShow v1.40[49]. The LD decay was calculated with PopLDdecay v3.42[50]. d_{xy}

and F_{ST} for the introgressed region is calculated with Pixy 1.2.7.beta1[51] with a 20kb

sliding window with 2kb steps. The haplotype heatmap of regions near the

introgression region was plotted with R package ComplexHeatmap v2.14.0[52].

Results

Genetic variants, mitogenome assembly, and Y-chromosome specific SNPs

A total of 121,252,917 SNPs in the 780 wild and domestic sheep samples were

obtained for downstream analyses. We obtained an average depth of 923.33 × (101.92

× - 8,324.72 ×) for the mitogenome assemblies after aligning all mitochondrial reads

from the 606 samples to the reference mitochondrial genome NC_001941.1

(Supplementary Table S2). After quality control for the SNP identification on the Y-

chromosome, we identified 4,096 Y-specific SNPs in a cohort of 239 rams with an

average read depth of 8.26 × (~4.25 × - 16.99 ×) for SNP (Supplementary Table S4).

Population genetic structure of the Asiatic mouflons in Iran

We observed two subpopulations in the 32 Asiatic mouflons in Iran (Figure 1 B/C/D)

with autosome SNPs. The larger subpopulation (*O. gmelini*_1) comprised 25 samples

across eastern and southern Iran, while the smaller subpopulation (*O. gmelini*_2)

consisted of only 7 samples limited in northwestern Iran (Fig. 1A; Supplementary

Table S2). Low kinship values within two subpopulations eliminate inbreeding caused

by sampling (Supplementary Table S5). We also observed significant genetic

divergence and different genetic diversity patterns (Supplementary Table S6-S7). The

result makes sure that the splition is not the artefact of inbreeding.

Phylogenetic analyses of the total 780 wild and domestic sheep samples identified the

same two subpopulations of Asiatic mouflon (*O. gmelini*_1 and *O. gmelini*_2) and six

genetic clusters of domestic sheep, which were congruent with their geographic

distribution (Supplementary Figure 1). Admixture analysis showed inter- and intra-

species genetic divergence similar to the phylogenetic tree (Fig. 1D).

Phylogenetic and haplotype analysis based on uniparental variants

We observed 593 mitochondrial haplotypes from the 767 mitogenomes of wild and

domestic sheep (Supplementary Table S2). The phylogenetic tree inferred from the

mitochondrial haplotypes revealed three monophyletic clades (clades I, II, and III)

(Figure 2B). Clade I comprised of haplogroups A, B, and D of domestic sheep. Clade

II consisted of the domestic sheep's haplogroups C and E and the Asiatic mouflon

samples, while the remaining Asiatic mouflon samples clustered with the urial in

clade III (Figure 2B).

For Y-chromosomal SNPs, the lineage HY2 of domestic sheep was closely related to

Asiatic mouflon in the haplogroups C and E but diverged from the main lineage HY1

(Figure 2C, Supplementary Table S8). However, phylogenetic and network analysis

revealed no shared mitochondrial haplotypes and only one shared Y-haplotypes of

HY2 between domestic sheep and Asiatic mouflon in Iran (Figure 2D, E). We

observed significant genetic divergence (*O. gmelini*_1: $F_{ST} = 0.11$, $P < 0.05$; *O.*

*gmelini*_2: $F_{ST} = 0.21$, $P < 0.05$) between Asiatic mouflons and domestic sheep in

whole genome levels (Supplementary Table S6). The observation implied that the

Asiatic mouflon in Iran might not be the direct ancestor of domestic sheep. Instead,

the relatively close phylogenetic relationships between the Asiatic mouflons in Iran

and domestic sheep suggested by haplogroups C, E, and Y-chromosomal

haplogroupY2 implied their possible gene flows.

Effective population size and divergence time

Inference of demographic history showed the Asiatic mouflons and domestic sheep

shared a similar historical pattern 10,000 years ago (Figure 3A), confirming the

Asiatic mouflon as the wild ancestor of sheep[53]. SMC++ analysis revealed a continual

decline of N_e in domestic sheep and Asiatic mouflon during ~8,000–10,000 years ago.

Nevertheless, the two Asiatic mouflon populations in Iran showed slower declines

than domestic sheep (Figure 3A), which probably reflected the bottleneck effect of

domestication. A sharp decline of N_e was detected ~1,000-5,000 years ago in the *O.*

gmelini_2, demonstrating a pattern different from that for the *O. gmelini_*1 (Figure

3A). We observed similar estimates of divergence time between *O. gmelini_*1 and the

six geographically differentiated genetic groups of domestic sheep during ~9,000-

15,000 years ago (Figure 3B, Supplementary Table 9), which was congruent with the

domestication time of sheep[2]. Additionally, we found the splitting time between the

two Asiatic mouflon populations in Iran around 5,000 years ago (Figure 3B,

Supplementary Table 9), indicating a recent genetic divergence of Asiatic mouflons in

Iran (Figure 1). For the divergence pattern inferred by WGS of the Asian mouflons,

the subpopulation *O. gmelini_*2 clustered with partial individuals of the subpopulation

*O. gmelini_*1 in the mitogenome and Y-chromosomal phylogenetic trees (Figure 1, 3).

The discordant phylogenetic patterns of Asiatic mouflons in Iran inferred by

uniparental and autosomal genetic variants could be explained by the recent

hybridization events[54].

We conducted Bayesian skyline analyses to estimate N_e based on mitogenomes and

Y-chromosomes. Based on the estimated TMRCA (time to the most recent common

ancestor) of Asiatic mouflons and domestic sheep[41], three lineages (lineage o.HCE:

haplogroup of Asiatic mouflons in Iran and partial domestic sheep of haplogroups C

and E, lineage A+B+D: haplogroups A, B, and D, and lineage C+E: haplogroups C

and E) of mitogenome showed different patterns. Lineage A+B+D showed a steep

increase in N_e at approximately ~25 k years ago. For lineage C+E, it also increased at

approximately ~25 k years ago, while lineage o.HCE demonstrated relatively fewer

changes since approximately ~100 k years ago (Figure 3C). Also, we observed

different patterns in the three Y-chromosomal lineages (lineage o.HY2: haplogroups

of Asiatic mouflons in Iran and partial domestic sheep of HY2, lineage HY1, and

lineage HY2). Lineages o.HY2 and HY2 displayed slow decline in N_e at

approximately ~25 k years ago, while lineage HY1 changed sightly since

approximately ~100 k years ago (Figure 3D). Thus, uniparental genetic markers

showed similar demographic patterns between o.HY2 and HY2 as well as between

HCE and C/E, but different patterns between the dominant linages A+B+D and HY1

(Figure 2, 3). The results suggested that the Asiatic mouflons in Iran might not be the

direct ancestor of domestic sheep but be involved in the genetic introgression.

Phylogeography of Asiatic mouflon

We built a *Cyt-b* dataset consisting of 247 Asiatic mouflons and 8 European

mouflons. The samples of Asiatic mouflon covered the current distribution of all the

subspecies (Supplementary Table S3, Figure 4A). To clarify phylogenetic

relationships between different Asiatic mouflon subspecies and domestic sheep, we

performed phylogenetic analysis of *Cyt-b* sequences of Asiatic mouflons and

domestic sheep using argali as the outgroup. In the phylogenetic tree built from *Cyt-b*,

Asiatic mouflon was assigned to 7 major clades supported by high bootstrap values

(94~100) (Figure 4B). The topology confirmed the five mitogenome clades of

domestic sheep. Haplogroups C+E clustered into monophyletic clade, which diverged

from the haplogroups A+B+D (Figure 4B). It was noteworthy that most of the hybrid

samples inferred above were integrated with *O. gmelini* or *O. vignei* samples, and

only around 10% of samples clustered with subspecies such as *O.g.ophion* and

O.g.anatolica (Figure 4B, Supplementary Table S10). Clade 3 comprised haplogroups

C+E of domestic sheep and the Asiatic mouflons in Iran with high frequencies of

shared haplotypes between them (Figure 4C, Supplementary Table S10). Haplogroups

A+B+D of domestic sheep and the Asiatic mouflons from Iran and Turkey fell within

clade 6, but no common haplotypes were detected between domestic and wild sheep

(Figure 4C, Supplementary Table S10). The results were in accordance with the

mitogenome analyses and implied that partial haplotypes C+E in domestic sheep

could have originated from the Asiatic mouflons in Iran by introgression.

Genetic introgression

To investigate the impact of introgression from the Asiatic mouflons in Iran on

domestic sheep, we implemented introgression tests between Iranian Asiatic mouflon

and SSA by D and f_4-ratio statistics[55,56]. We first conducted an admixture analysis

between Asiatic mouflons and the SSA populations. We identified three genetic

clusters with the smallest cross validation error ($K = 3$) (*O. gmelini*_1, *O. gmelini*_2,

and SSA) and detected genetic components of Asiatic mouflons in Iran in some

domestic populations (Figure 5A). Further, we employed the ABBA model ((((P1, P2),

P3), P4) to assess introgression between *O. gmelini*_1 or *O. gmelini*_2 (P3) and SSA

sheep populations (P2). We used Menz sheep (MEN) from Africa as P1, which did

not show genetic introgression from Asiatic mouflon and SSA populations[8,15], and

bighorn sheep as the outgroup P4, which inhabit North America and have no gene

flows with SSA sheep populations. Of the two Asiatic mouflon populations, *O.*

*gmelini*_1 or *O. gmelini*_2 served as P3 and 18 SSA populations as P2. We calculated

the estimates of Patterson's D-statistic and f_4-ratio for all the 18 SSA sheep

populations, and determined introgression based on the following criteria: $D > 0$, *P*-

value < 0.05, and f_4-ratio > 0. Only four sheep populations (Garut, Garole,

Bangladeshi, and Sumatra sheep) showed genetic introgression from *O. gmelini*_2

(Figure 5B, Supplementary Table S11). All the four sheep populations are distributed

in the regions near to the Indian Ocean.

To further examine the genetic introgression in the four sheep populations, we

calculated f_{dm} with a 50-SNP window and 25-SNP step across the sheep genome. We

identified introgressed regions genomic with top 5% f_{dm} and $D > 0$. We obtained a

common introgression region in the four populations spanning 6,384,279 bp (Figure

5C, Supplementary Tables S12 and S13). We then calculated the PBS value of the

four populations to identify regions under selection with MEN sheep and *O.*

*gmelini*_1 as the reference populations. We considered the top 1% of PBS regions as

being under selection (Figure 5D, Supplementary Table S14). We overlapped the

selected regions with the common introgressed regions and detected 19,614 bp region

as the selected introgressed region (Supplementary Table S15). We further overlapped

the introgressed genomic region with sheep QTLs (Supplementary Table S15). We

found that it overlapped with the QTLs for body_weight, carcass_fat, and fat_weight

QTLs, which might be associated with the small body size of Garole, Bangladeshi,

and Garut sheep[57–59]. Additionally, the Nematodirus_FEC QTL could be related to the

phenotypes of Sumatra sheep's susceptibility to nematode parasitism[55]. The results

implied that the adaptive selected introgressed region could have been introgressed

from *O. gmelini*_2, and then have undergone the strong selection to facilitate their

adaptation to the hot and humid environments near the Indian Ocean. We further

overlapped the introgressed genomic region with sheep QTLs (Supplementary Table

S15).

Discussion

Traditionally, Asiatic mouflon had been classified into five subspecies using phenotypic traits, such as coat color, horns, and geographic patterns[11]. However, the intraspecies taxonomy has long lacked genetic evidence. Previous phylogenetic analysis based on partial mtDNA divided Asiatic mouflon into several clades[3,4,58], different from the taxonomy of five subspecies defined by phenotypic traits and geographic patterns[11]. Here, we identified two genetic clusters of the Asiatic mouflons in Iran based on WGS data: *O. gmelini_2* limited in Kaboudan, and *O. gmelini _1* over a wide geographic area in Iran (Figure 1). We detected the divergence time of two clusters around ~5,000 years ago (Figure 3B, Supplementary Table S9), which indicated a recent split of them, and was further supported by the inferred phylogenies based on uniparental genetic markers (Figure 2B, C). All the individuals of *O. gmelini _2* collected in Kabudan inhabited the overlap geographic region with Armenian Mouflon (*O. g.gmelini*), while individuals of *O. gmelini _1* were sampled from the distribution areas of several subspecies such as Armenian Mouflon (*O. g.gmelini*), Isfahan mouflon (*O.g.isphahanica*), and Laristan mouflon (*O.g.laristanica*) (Figure 1A)[10].

These results suggested a significant genetic divergence between Armenian Mouflon and the other subspecies (Supplementary Table S6). Nevertheless, the absence of phenotypic data and other sample information for the Asian mouflons and a

comprehensive sampling throughout Iran limited our in-depth understanding of

within-species genetic differentiation in Asiatic mouflon. Also, Asiatic mouflon is

under the status of "Near Threatened", and their population size has declined in the

past two decades under competition with domestic sheep, poaching, and habitat

deterioration[10,60]. Therefore, the population genetic structure analyses and delimitation

of taxonomic classification based on WGS data are fundamental for conservation

decisions and effective conservation management[61,62].

Genetic variants of uniparental inheritance offer insights into the domestication and

evolutionary history of farm animals, such as pigs[63], sheep[5,14] and goats[64,65]. In

domestic sheep, here we identified five maternal haplogroups (A, B, C, D, and E) in

two monophyletic groups (A+B+D and C+E), and two paternal clades (HY1 and

HY2) (Figure 2B, C). We found that haplotypes of most of the Asiatic mouflons in

Iran clustered with haplogroups C+E based on mitochondrial marker (24/31) and

grouped with haplogroup HY2 based on Y-chromosomal SNPs (12/21)

(Supplementary Table S8). Nevertheless, the Asiatic mouflons in Iran showed higher

divergence from the dominant maternal lineages A and B and paternal lineages HY1

(Figure C, D).

Additionally, *Cyt-b* sequences collected from the whole geographic areas of Asiatic

mouflon also revealed a closer relationship between Asiatic mouflon in Iran and

domestic sheep of maternal haplogroups C+E (Figure 4). The results supported the

hypothesis that mtDNA haplogroups C+E and Y-chromosomal haplogroup HY2

could have originated from Asiatic mouflons in Iran[14]. However, WGS analysis

revealed intraspecies substructure in the Asiatic mouflons in Iran, and genetic

introgression from the *O. gmelini_2* showed a larger impact on domestic sheep than

O. gmelini_1 (Figure 5). Similiar results were observed from Alberto et al., (2018)

that IROO individuals were split into subpopulations and had admixture with some

domestic sheep[38]. Therefore, Iran's two Asiatic mouflon populations showed similar

phylogenetic patterns of uniparental inheritance but different introgression patterns of

WGS, implying multiple introgression events from the Asiatic mouflons to domestic

sheep. In summary, our results implied that Asiatic mouflons in Iran might not be the

ancestor of domestic sheep[4,14], but could have contributed to the gene pool of

domestic sheep through multiple introgressions. Also, the results implied the

complexity of sheep domestication[4].

Recent studies suggested that sheep diffused to South and Southeast Asia via the

Middle East[15]. Our results revealed the genetic introgression from Asiatic mouflon

into some South-and-Southeast Asian sheep populations, supporting previous

findings[8,15]. Further, we detected genetic introgression in the four sheep populations

of Garut, Garole, Bangladeshi, and Sumatra, and were only from *O. gmelini_2* in

Kabudan (Figure 5B). The introgression event is congruent with the demographic

history of *O. gmelini_2* and South-and-Southeast Asian sheep inferred here and

previously: South-and-Southeast Asian sheep descended from Central-and-East Asian

populations around 5.85 to 6.01 ka and then admixed from the Middle Eastern sheep

3.93–4.18 ka[15], and *O.gmelini_2* split from *O.gmelini_1* around ~5,000 years ago

(Figure 3B).

In the four sheep populations, we found that common introgression signals from *O.*

gmelini_2 covered about 4.29% of the total introgressed regions. In contrast to

population-specific introgression, common introgression could be related to a specific

trait or adaptation to similar environments, such as *EPAS1* for the adaptation to

hypoxia[45,66]. We further integrated *PBS* analyses and found four positive selective

regions in the common introgressed regions, which were related to

carcass_fat_percentage, lean_meat_yield_percentage (Figure 5D). This might have

contributed to the small body size phenotype of Garole, Sumatra, and Bangladeshi

sheep[58,67,68]. We also found introgression signals in the *PCDH9* (Figure 5E), which

had been reported to be associated with local adaptation (e.g., hot and arid

environments) in sheep and goats[69,70], and fat deposition[71]. Also, early studies showed

adipogenesis, ECM remodeling, inflammation, and lipid droplet dynamics in mature

adipose tissue played essential roles in response to climate changes in sheep[72]. Further

analysis from Chip-seq and RNA-Seq data will be required to uncover the function

and regulatory mechanisms of *PCDH9* to adapt to hot and humid environments in

sheep.

In summary, we conducted a comprehensive and integrated analysis of Asiatic

mouflon and domestic sheep based on WGS, mitogenome and Y-chromosomal

genetic variants. Our results suggested that the mitochondrial haplogroups C/E and Y-

chromosomal haplogroup HY2 of domestic sheep could have originated from the

Asiatic mouflons in Iran by introgression and further clarified the effects of the

Iranian Asiatic mouflon in the domestication processing. Additionally, our results

revealed the introgression signals of the Asiatic mouflons in Iran in several domestic

sheep populations in South and Southeast Asia and showed that the introgression

could have facilitated their adaptation to hot and humid environments near the Indian

Ocean. Our findings added new insights into the domestication and subsequent

evolutionary history of sheep.

Data availability

All raw sequencing reads in this study were deposited in NCBI's SRA under the

accession number OR160429-OR161034.

Code availability

All scripts used for this work were performed using open-source software tools and

are available from the corresponding authors upon request.

Acknowledgments

This study was financially supported by grants from the National Natural Science

Foundation of China (Nos. U21A20246, 32061133010, and 31972527), the National

Key Research and Development Program-Key Projects (2021YFD1200900 and

2021YFD1300904), the Second Tibetan Plateau Scientific Expedition and Research

Program (STEP) (No. 2019QZKK0501), and Initiation Fund of Sanya Institute of

China Agricultural University (SYND-2022-12). We thank the High-performance

Computing Platform of China Agricultural University for providing computing

resources.

Author contributions

547   F.-H.L. and M.-H.L. conceived and supervised the study. D.-F.W and F.-H.L.

548   conducted the data analysis. D.-F.W, F.-H.L. and P.O wrote the manuscript, and M.-

549   H.L. and P.O revised the paper. All the authors reviewed and approved the final

550   manuscript.

Competing interests

553   The authors declare no competing interests.

References

1.   Zeder, M. A. Domestication and early agriculture in the Mediterranean Basin: Origins, diffusion, and impact. *Proc. Natl. Acad. Sci. U.S.A.* 105, 11597–11604 (2008).

2.   Pedrosa, S. et al. Evidence of three maternal lineages in near eastern sheep supporting multiple domestication events. *Proc. R. Soc. B: Biol. Sci.* 272, 2211–2217 (2005).

3.   Demirci, S. et al. Mitochondrial DNA Diversity of Modern, Ancient and Wild Sheep (*Ovis gmelinii anatolica*) from Turkey: New Insights on the Evolutionary History of Sheep. *PLoS ONE* 8, e81952 (2013).

4.   Her, C. et al. Broad maternal geographic origin of domestic sheep in Anatolia and the Zagros. *Anim. Genet.* 53, 452–459 (2022).

5.   Lv, F.-H. et al. Mitogenomic Meta-Analysis Identifies Two Phases of Migration in the History of Eastern Eurasian Sheep. *Mol. Biol. Evol.* 32, 2515–2533 (2015).

6.   Deng, J. et al. Paternal Origins and Migratory Episodes of Domestic Sheep. *Curr. Biol.* 30, 4085-4095.e6 (2020).

7.   Kalds, P. et al. Genetics of the phenotypic evolution in sheep: a molecular look at diversity-driving genes. *Genet. Sel. Evol.* 54, 61 (2022).

8.   Cheng, H. et al. Long divergent haplotypes introgressed from wild sheep are associated with distinct morphological and adaptive characteristics in domestic

575  sheep. *PLoS Genet.* 19, e1010615 (2023).

576 9. Yeomans, L., Martin, L. & Richter, T. Expansion of the known distribution of

577  Asiatic mouflon (*Ovis orientalis*) in the Late Pleistocene of the Southern Levant.

578  *R. Soc. Open Sci.* 4, 170409 (2017).

579 10. Michel, S. & Ghoddousi, A. *Ovis Gmelini*: The IUCN Red List of Threatened

580  Species 2020. e. T54940218A22147055, https://dx. doi. org/10.2305/IUCN. UK

581  2, (2020).

582 11. Blyth, E. XXVI.— An Amended List of the Species of the Genus *Ovis*. *Ann.*

583  *Mag. Nat.* 7, 195–201 (1841).

584 12. Fernandez H, Taberlet P, Mashkour M, Vigne J-D, Luikart G. The First Steps of

585  Animal Domestication: New Archaeozoological Approaches, eds Vigne JD,

586  Peters J, Helmer D (Oxbow Books, Oxford), pp 50–54 (2005) .

587 13. Hiendleder, S., Kaupe, B., Wassmuth, R. & Janke, A. Molecular analysis of wild

588  and domestic sheep questions current nomenclature and provides evidence for

589  domestication from two different subspecies. *Proc. R. Soc. B: Biol. Sci.* 269,

590  893–904 (2002).

591 14. Li, R. et al. A Hu sheep genome with the first ovine Y chromosome reveal

592  introgression history after sheep domestication. *Sci. China Life Sci.* (2020)

593  doi:10.1007/s11427-020-1807-0.

594 15. Lv, F.-H. et al. Whole-Genome Resequencing of Worldwide Wild and Domestic

595  Sheep Elucidates Genetic Diversity, Introgression, and Agronomically Important

Loci. *Mol. Biol. Evol.* 39, msab353 (2022).

16.   Cao, Y.-H. et al. Historical Introgression from Wild Relatives Enhanced Climatic

Adaptation and Resistance to Pneumonia in Sheep. *Mol. Biol. Evol.* 38, 838–855

(2021).

17.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for

Illumina sequence data. *Bioinformatics* 30, 2114–2120 (2014).

18.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–

Wheeler transform. *Bioinformatics* 25, 1754–1760 (2009).

19.   McKenna, A. et al. The Genome Analysis Toolkit: A MapReduce framework for

analyzing next-generation DNA sequencing data. *Genome Res.* 20, 1297–1303

(2010).

20.   Li, H. et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*

25, 2078–2079 (2009).

21.   Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software

Version 7: Improvements in Performance and Usability. *Mol. Biol. Evol*. 30,

772–780 (2013).

22.   Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for

automated alignment trimming in large-scale phylogenetic analyses.

Bioinformatics 25, 1972–1973 (2009).

23.   Chang, C. C. et al. Second-generation PLINK: rising to the challenge of larger

and richer datasets. *GigaScience* 4, 7 (2015).

24. Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics* 35, 526–528 (2019).

25. Yu, G. Using ggtree to Visualize Data on Tree-Like Structures. *Curr. Protoc.*

Bioinformatics 69, (2020).

26. Frichot, E., Mathieu, F., Trouillon, T., Bouchard, G. & François, O. Fast and

Efficient Estimation of Individual Ancestry Coefficients. *Genetics* 196, 973–983

(2014).

27. Danecek, P. et al. The variant call format and VCFtools. *Bioinformatics* 27,

2156–2158 (2011).

28. Goudet, J. hierfstat, a package for r to compute and test hierarchical F-statistics.

Mol. Ecol. Notes 5, 184–186 (2005).

29. Paradis, E. pegas: an R package for population genetics with an integrated–

modular approach. *Bioinformatics* 26, 419–420 (2010).

30. Minh, B. Q. et al. IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* 37, 1530–1534

(2020).

31. Steenwyk, J. L., Buida, T. J., Li, Y., Shen, X.-X. & Rokas, A. ClipKIT: A

multiple sequence alignment trimming software for accurate phylogenomic

inference. *PLoS Biol.* 18, e3001007 (2020).

32. Gatto, L., Catanzaro, D. & Milinkovitch, M. C. Assessing the Applicability of

the GTR Nucleotide Substitution Model Through Simulations. *Evol. Bioinform.*

Online 2, 145–155 (2007).

33. Leigh, J. W. & Bryant, D. popart : full-feature software for haplotype network

construction. *Methods Ecol. Evol.* 6, 1110–1116 (2015).

34. Ortiz, E. M. vcf2phylip v2.0: convert a VCF matrix into several matrix formats

for phylogenetic analysis. (2019) doi:10.5281/zenodo.2540861.

35. Terhorst, J., Kamm, J. A. & Song, Y. S. Robust and scalable inference of

population history from hundreds of unphased whole-genomes. *Nat. Genet.* 49,

303–309 (2017).

36. Beynon, S. E. et al. Population structure and history of the Welsh sheep breeds

determined by whole genome genotyping. *BMC Genet.* 16, 65 (2015).

37. Zhao, Y.-X. et al. Genomic Reconstruction of the History of Native Sheep

Reveals the Peopling Patterns of Nomads and the Expansion of Early

Pastoralism in East Asia. *Mol. Biol. Evol.* 34, 2380–2395 (2017).

38. Alberto, F. J. et al. Convergent genomic signatures of domestication in sheep and

goats. *Nat. Commun.* 9, 813 (2018).

39. Bouckaert, R. et al. BEAST 2.5: An advanced software platform for Bayesian

evolutionary analysis. *PLoS Comput. Biol.* 15, e1006650 (2019).

40. Drummond, A. J. Bayesian Coalescent Inference of Past Population Dynamics

from Molecular Sequences. *Mol. Biol. Evol.* 22, 1185–1192 (2005).

41. Upadhyay, M. et al. Whole genome sequencing reveals a complex introgression

history and the basis of adaptation to subarctic climate in wild sheep. *Mol. Ecol.*

30, 6701–6717 (2021).

42. Patterson, N. et al. Ancient Admixture in Human History. *Genetics* 192, 1065–

1093 (2012).

43. Malinsky, M., Matschiner, M. & Svardal, H. Dsuite - Fast D-statistics and

related admixture evidence from VCF files. *Mol. Ecol. Resour.* 21, 584–595

(2021).

44. Arnholt, A. T. & Evans, B. BSDA: Basic statistics and data analysis. *R package*

version 1 (2017).

45. Yi, X. et al. Sequencing of 50 Human Exomes Reveals Adaptation to High

Altitude. *Science* 329, 75–78 (2010).

46. Hämälä, T. & Savolainen, O. Genomic Patterns of Local Adaptation under Gene

Flow in *Arabidopsis lyrata*. *Mol. Biol. Evol.* 36, 2557–2571 (2019).

47. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing

genomic features. *Bioinformatics* 26, 841–842 (2010).

48. Hu, Z.-L., Park, C. A. & Reecy, J. M. Bringing the Animal QTLdb and CorrDB

into the future: meeting new challenges and providing updated services. *Nucleic*

Acids Res. 50, D956–D961 (2022).

49. Dong, S.-S. et al. LDBlockShow: a fast and convenient tool for visualizing

linkage disequilibrium and haplotype blocks based on variant call format files.

Brief. Bioinformatics 22, bbaa227 (2021).

50. Zhang, C., Dong, S.-S., Xu, J.-Y., He, W.-M. & Yang, T.-L. PopLDdecay: a fast

and effective tool for linkage disequilibrium decay analysis based on variant call

format files. *Bioinformatics* 35, 1786–1788 (2019).

51. Korunes, K. L. & Samuk, K. pixy : Unbiased estimation of nucleotide diversity

and divergence in the presence of missing data. *Mol. Ecol. Resour.* 21, 1359–

1368 (2021).

52. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and

correlations in multidimensional genomic data. *Bioinformatics* 32, 2847–2849

(2016).

53. Sanna, D. et al. The First Mitogenome of the Cyprus Mouflon (*Ovis gmelini*

ophion): New Insights into the Phylogeny of the Genus *Ovis*. *PLoS ONE* 10,

e0144257 (2015).

54. Teng, H. et al. Population Genomics Reveals Speciation and Introgression

between Brown Norway Rats and Their Sibling Species. *Mol. Biol. Evol.* 34,

2214–2228 (2017).

55. Reich, D., Thangaraj, K., Patterson, N., Price, A. L. & Singh, L. Reconstructing

Indian population history. *Nature* 461, 489–494 (2009).

56. Green, R. E. et al. A Draft Sequence of the Neandertal Genome. *Science* 328,

710–722 (2010).

57. Jakaria, J., Zein, M. S. A., Sulandari, S., Subandriyo, S. & Muladno, M. The use

of microsatellite markers to study genetic diversity in Indonesian sheep. *J.*

Indones. Trop. Anim. Agric. 37, 1–9 (2012).

58. Sharma, R. C., Arora, A. L., Narula, H. K. & Singh, R. N. Characteristics of

Garole sheep in India. *Anim. Genet. Resour. Inf.* 26, (1999).

59. Ramanuj, B., Mandal, P., Pal, U. & Kunal, R. Productivity and Genetic Potential

of Garole Sheep of India-A Review. *Asian J. Anim. Sci.* 4, 170–189 (2010).

60. Bleyhl, B. et al. Assessing niche overlap between domestic and threatened wild

sheep to identify conservation priority areas. *Divers. Distrib.* 25, 129–141

(2019).

61. Yang, L. et al. Evolutionary Conservation Genomics Reveals Recent Speciation

and Local Adaptation in Threatened Takins. *Mol. Biol. Evol.* 39, msac111

(2022).

62. Hu, Y. et al. Genomic evidence for two phylogenetic species and long-term

population bottlenecks in red pandas. *Sci. Adv.* 6, eaax5751 (2020).

63. Ai, H. et al. Human-Mediated Admixture and Selection Shape the Diversity on

the Modern Swine (*Sus scrofa*) Y Chromosomes. *Mol. Biol. Evol.* 38, 5051–5065

(2021).

64. Luikart, G. et al. Multiple maternal origins and weak phylogeographic structure

in domestic goats. *Proc. Natl. Acad. Sci. U.S.A.* 98, 5927–5932 (2001).

65. Naderi, S. et al. The goat domestication process inferred from large-scale

mitochondrial DNA analysis of wild and domestic individuals. *Proc. Natl. Acad.*

Sci. U.S.A. 105, 17659–17664 (2008).

66. Wu, D.-D. et al. Convergent genomic signatures of high-altitude adaptation

among domestic mammals. *Natl. Sci. Rev.* 7, 952–963 (2020).

67. Porter, V. Mason's World Dictionary of Livestock Breeds, Types and Varieties,

6th Edition. (CABI, 2020).

68. Putra, W. P. B., Alwi, Y. & Suryani, H. Morphometrics characterization of Thin-

Tail sheep in lowland and highland areas. *Trop. Anim. Sci. J.* 44, 386–398

(2021).

69. Serranito, B. et al. Local adaptations of Mediterranean sheep and goats through

an integrative approach. *Sci. Rep.* 11, 21363 (2021).

70. Kim, E.-S. et al. Multiple genomic signatures of selection in goats and sheep

indigenous to a hot arid environment. *Heredity* 116, 255–264 (2016).

71. Mastrangelo, S. et al. Genome-wide scan of fat-tail sheep identifies signals of

selection for fat deposition and adaptation. *Anim. Prod. Sci.* 59, 835–848 (2018).

72. Xu, Y.-X. et al. Whole-body adipose tissue multi-omic analyses in sheep reveal

molecular mechanisms underlying local adaptation to extreme environments.

Commun. Biol. 6, 159 (2023).

Fig. 1 Population genetics structure of Asiatic mouflons in Iran (*O. gmelini*). a Geographic distribution of 32 Asiatic mouflons. Purple and blue dots indicate the location of *O. gmelini*_1 and *O. gmelini*_2 subpopulations; **b** Principal components 1 and 2 for the 32 Asiatic mouflon in Iran; **c** Neighbor-joining (NJ) trees of the 32 Asiatic mouflons in Iran based on whole-genome SNPs using the nine urial samples as the outgroup; **d** Population genetic structure of the Asiatic mouflons in Iran and domestic sheep inferred from the program the sNMF v1.2 (*k* = 6) using whole-genome SNPs.

Fig. 2 Phylogeny of Asiatic mouflon, urial, and domestic sheep inferred from mitogenomes and Y-chromosomal genetic variants. a Geographical distribution of the samples in this and previous studies; **b** Phylogeny of Asiatic mouflon, urial, and domestic sheep inferred from 767 mitogenomes; **c** Phylogeny of Asiatic mouflon, urial, and domestic sheep inferred from 239 Y-chromosomal SNPs; **d** Network of mitogenome haplotypes; **e** Network of Y-chromosome haplotypes. Bootstrapping values above 50% were labeled on the phylogenetic tree. Bighorn sheep were used as the outgroup in phylogenetic analyses based on mitogenome and Y-chromosome genetic variants.

Fig. 3 Demographic history of Asiatic mouflon in Iran and domestic sheep. a Effective population sizes (N_e) across six domestic sheep populations and the two subpopulations of Asiatic mouflon in Iran inferred from SMC++; **b** Inference of split times between *O. gmelini*_1 and the other wild and domestic sheep populations. The vertical lines represent the split time between each pair of populations; **c** Bayesian skyline plot (BSP) reconstruction of historical change in effective population size (N_e) of domestic sheep of maternal haplogroups HA+B+D, and HC+E, and the Asiatic mouflons in Iran; **d** Bayesian skyline plot (BSP) reconstruction of historical changes in N_e of paternal lineages HY1 and HY2 of domestic sheep and the Asiatic mouflon in Iran. The middle line represents median estimates of $N_e\tau$, and the shaded area represents 95% highest posterior density (HPD) intervals.

Fig. 4 Geographic distribution of Asiatic mouflon and the phylogenetic relationship among Asiatic mouflon subspecies and domestic sheep. **a** Geographical distribution of Asiatic mouflon and samples information of 32 Asiatic mouflon in Iran and 301domestic sheep samples; **b** Phylogenetic tree inferred based on complete *Cyt-b* sequences using the maximum likelihood algorithms. Bootstrap values above 50% were labeled on the phylogenetic tree. **c** Network of Asiatic mouflon in Iran and Turkey and domestic sheep lineage of C and E built from complete *Cyt-b* sequences.

Fig. 5 Genetic introgression of Asiatic mouflon in SSA domestic populations. a Admixture proportions of Asiatic mouflon in Iran and SSA domestic populations inferred from the program the sNMF v1.2 ($K = 3$) using whole-genome SNPs. **b** Heatmap of D-statistics and f_4-ratio calculated by Dsuite v0.5 applied to data including *O. gmelini*_1, *O. gmelini*_2, and SSA domestic populations; **c,** Venn diagram for the shared introgression regions among four domestic South and Southeast populations; **d** Manhattan plot showing signals of selection in the four introgressed SSA populations (BGE/GAR/GUR/SUM) detected by the PBS with the dashed line marking 99.9% threshold. The introgression region, QTL, and adjacent gene *PCDH9* were labeled with blue, purple, and green bars, respectively. The sheep QTLs overlapped with the top 1% PBS regions are shown, **e** PBS and f_{dm} values around the significant introgressed genomic region on Chr.10.

Fig. 6 Candidate gene *PCDH9* associated with the significant introgression signal on Chr. 10. **a** Linkage disequilibrium (LD) blocks in the significant introgression region and gene *PCDH9* (chr10:41,104,271-43,073,231). The red and yellow bars represent the location of the significant introgression region and gene *PCDH9*; **b** Pattern of LD decay in the introgressed populations. The black line is calculated from the whole 10 chromosomes, the red line is calculated from the beginning of the significant region to the next 800 kb (chr10:41,204,271-42,004,271); **c** D_{xy} and F_{ST} around the significant introgression region on Chr. 10 (chr10:41,104,271-41,313,804), calculated by pixy v1.2.7.beta1; **d** Haplotype pattern of the significant introgression region on Chr. 10 (chr10:41,204,271-41,213,804) (left) and exons of *PCDH9* (right).