Genomic analyses of Asiatic Mouflon in Iran provide

insights into the domestication and evolution of sheep

Dong-Feng Wang[1,2,3], Pablo Orozco-terWengel[4], Meng-Hua Li[1*], Feng-Hua Lv[1*]

[1] College of Animal Science and Technology, China Agricultural University, Beijing,

China

[2] CAS Key Laboratory of Animal Ecology and Conservation Biology, Institute of

Zoology, Chinese Academy of Sciences (CAS), Beijing, China

[3] College of Life Sciences, University of Chinese Academy of Sciences (UCAS),

Beijing, China

[4] School of Biosciences, Cardiff University, Cardiff, UK

*Corresponding author

E-mail addresses:

DFW: atongsa@163.com

POW: Orozco-terWengelPA@cardiff.ac.uk

MHL: menghua.li@cau.edu.cn

FHL: lvfenghua@cau.edu.cn

Abstract

Background

Asiatic mouflon (*O. gmelini*) consists of several subspecies mainly distributed in Armenia, southern Azerbaijan, Cyprus, northern, southern, and western regions of Iran, and eastern and central regions of Turkey nowadays. Genome analyses of Asiatic mouflon in Iran revealed that they could have diverged from the direct ancestor of domestic sheep, and showed genetic introgression into domestic sheep after domestication. However, the impact of the Asiatic mouflon subspecies in Iran on sheep domestication remains unclear.

Results

Here, we conducted a comprehensive population genomics analysis of Asiatic mouflons in Iran with 780 whole-genome sequences, 767 whole mitogenomes, and 239 Y chromosomes. Whole-genome sequence analyses revealed two subpopulations of the Asiatic mouflons in Iran: *O. gmelini_2* limited in Kabudan and *O. gmelini_1* over a wide geographic area. Phylogenetic analyses of the Asiatic mouflons in Iran based on uniparental variants revealed a monophyletic lineage with the mitochondrial haplogroups C+E, and clustered into a monophyletic with Y-chromosomal lineage HY2 of sheep. Additionally, introgression tests detected significant signals of genetic introgression from *O. gmelini_2* to four sheep populations (e.g., Garut, Garole, Bangladeshi, and Sumatra) in South and Southeast Asia. In the four sheep

populations, selective tests and introgression signals revealed that the wild

introgression could have contributed to their small body size and local adaptation to

the hot and humid environments in the Indian Peninsula.

Conclusions

Our results suggested that the maternal haplogroups C+E and paternal lineage HY2

could have originated from the Asiatic mouflon in Iran. Also, our findings provide

new insights into sheep domestication and subsequent introgressions from wild

ancestors to domestic populations.

Background

Archeozoological and genetic evidence suggested that Asiatic mouflon (*O. gmelini*),

the wild ancestor of sheep, have underwent a period of multigenerational human

management in southeastern Anatolia by ~10,500 years B.P. [1–4]. Following

domestication, sheep diffused throughout Eurasia with human migrations and

developed into different populations under natural and artificial selections [5–8].

Asiatic mouflon ranges over a wide geographic area in Armenia, southern Azerbaijan,

Cyprus, northern, southern, and western regions of Iran, and eastern and central

regions of Turkey. It has been classified into five subspecies based on morphological

differences and geographic distributions [9, 10], such as Armenian mouflon (*O.*

gmelini gmelini) [11], Anatolian mouflon (*O. gmelini anatolica*) [11], Isfahan

mouflon (*O. gmelini isphahanica*), Laristan mouflon (*O. gmelini laristanica*), and

Cyprus mouflon (*O. gmelini ophion*) [11]. Due to the complex intraspecies taxonomy

of Asiatic mouflon, the genetic contributions of various subspecies to the

domestication of sheep remains unclear [3, 12–14].

Previous genetic studies revealed controversial ancestral origins of domestic sheep.

Hiendleder et al. (2002) proposed that sheep were domesticated from two different

subspecies in Turkey and western Iran, corresponding to mtDNA lineages A and B of

domestic sheep, respectively [13]. Through genetic and phylogeographic analyses of

partial mtDNA sequences of modern and ancient samples in wild and domestic sheep,

Demirci et al. (2013) suggested that domestic sheep could have originated from two

distinct ancestral populations of mtDNA lineages A+B+D and C+E of domestic

sheep, respectively [3]. Based on genetic, archeozoological, and geographic evidence

of 212 Asiatic mouflon samples covering most of their distribution area, a recent

study suggested that the maternal origin of sheep could be traced to a subspecies of *O.*

gmelini gmelini in eastern Anatolia and central Zagros [4].

Despite the intensive focus on material inheritance, Y-chromosome and whole-

genome analyses have received less attention. Recently, Y-chromosome and whole-

genome analyses of Asiatic mouflon in Iran revealed that they could have diverged

from the direct ancestor of domestic sheep, and showed genetic introgression into

4   4

domestic sheep after domestication [8, 14, 15]. Also, these studies have demonstrated

that genomes of Asiatic mouflon in Iran are crucial for the development of

morphological and adaptive characteristics in domestic sheep [8, 15, 16].

Nevertheless, the impact and extent of genetic introgression involving the Asiatic

mouflon subspecies in Iran (e.g., *O. gmelini gmelini*, *O. gmelini isphahanica*, and *O.*

gmelini laristanica) and their hybrid populations (*O. gmelini gmelini* × *O. vignei*) on

the genomes of domestic sheep remains unclear.

Here, we comprehensively analyzed whole genome sequence, complete and partial

mtDNA sequences, and Y-chromosome genetic variants in domestic sheep and

Asiatic mouflon, covering a large distribution area in Asia, Europe, Africa, and

America. We aimed to elucidate the contributions of various Asiatic mouflon

subspecies in Iran to the domestication of sheep. Additionally, we investigated the

impact of wild introgression of the Asiatic mouflon subspecies in Iran on the genomes

of domestic sheep populations in South and Southeast Asia, which have adapted to the

hot and humid environments near the Indian Ocean.

Methods

Datasets

Whole-genome sequences (WGS) of 780 individuals representing 158 domestic sheep

populations and three wild sheep species (32 Asiatic mouflon (*Ovis gmelini*) in Iran, 9

urial (*Ovis vignei*), and 1 bighorn sheep (*Ovis canadensis*)) were obtained from

publicly available data [See Additional file 1, Table S1]. Besides these, we retrieved

161 whole mitogenomes from the NCBI [See Additional file 1, Table S2]. Also, we

included 333 Cytochrome b (*Cytb*) sequences of Asiatic mouflon in the integrated

analysis [See Additional file 1, Table S3].

Variant calling

We implemented quality-based trimming for WGS data with Trimmomatic v0.39 [17]

and mapped high-quality reads to the reference genome

Oar_rambouillet_v1.0_CM022046.1. (https://www.

ncbi.nlm.nih.gov/assembly/GCF_002742125.1/;

https://www.ncbi.nlm.nih.gov/nuccore/CM022046.1/, last accessed May 12, 2020)

using the Burrows–Wheeler aligner (BWA mem) v.0.7.8 [18] with the default

parameters. Then, we removed duplicates with the *MarkDuplicates* module in GATK

v 4.1.2.0 [19] and called SNPs following the previous procedures [15]. We used the

HaplotypeCaller module in GATK v 4.1.2.0 to generate the GVCF file of each

sample and merged the raw GVCF file of each individual with the *CombineGVCFs*

module. Afterward, we genotyped SNPs using the *GenotypeGVCFs* module and

filtered SNPs using the *VariantFiltering* module of GATK with the parameters

"QUAL < 30.0 || QD < 2.0 || MQ < 40.0 || FS > 60.0 || SOR > 3.0 || MQRankSum < -

12.5 || ReadPosRankSum < -8.0". We obtained a total of 121,252,917 high-quality

autosomal SNPs and 203,938 Y-chromosomal SNPs for the following analyses.

Mitochondrial DNA assembly

We assessed the mitogenome depth of each sample using the SAMtools v.1.9 [20] and

kept only the samples with depth > 100 in the below analyses. We assembled the

complete mitogenomes using the following protocols. We first converted BAM files

to the fastq format. Then, we generated the mitogenome consensus sequence from the

fastq files using MIA v1.0 (https://github.com/mpieva/mapping-iterative-assembler).

In addition to the 605 mitogenomes assembled here, we also included 161 whole

mitogenome sequences retrieved from the NCBI GenBank in the analysis, with a final

mitogenome dataset of 767 sequences from 727 domestic sheep, 31 Iranian Asiatic

mouflon, and 9 urial. All sequences were aligned using MAFFT v7.515 [21] and

trimmed with trimAl v1.4 [22].

Y-chromosomal SNPs

To obtain single-copy Y-chromosomal SNPs, raw SNPs in the 780 samples were

processed as follows. Firstly, we identified ram samples from the whole samples

using PLINK v1.90b6.26 [23] with the "--check-sex" option and kept only these ram

samples in the following analyses. We then excluded SNPs in the regions covering the

pseudoautosomal regions (PARs) and the highly repetitive sequences (around 1.06

Mb) [14]. Afterwards, we filtered SNPs with the mean sequencing depth ranging from

$0.5 \times$ to $2 \times$ of the expected depth (half of the genome-wide depth) [14]. Further, we

removed SNPs that were heterozygous in at least one of the male samples. Finally, we

retained 4,096 Y-chromosome SNPs for the haplotype and phylogenetic analyses.

Population structure analysis

To reveal the genetic differentiation within the Asiatic mouflon in Iran, we examined

the population genetic substructure for the total 779 wild and domestic samples using

a set of autosomal SNPs filtered as follows. First, we filtered the total 25,270,188

high-quality SNPs with missing rates < 0.1 and minor allele frequencies > 0.05 (MAF

> 0.05) using PLINK v1.90b6.26 [23]. Then, we pruned the remaining SNPs with the

options "--indep-pairwise 50 10 0.1" of PLINK v1.90b6.26 [23]. Afterwards,

1,566,300 SNPs were retained for the population genetic structure analysis.

We computed the identity by state among samples using PLINK v1.90b6.26 [23] with

option "distance square 1-ibs". Next, we generated a neighbor-joining (NJ) tree using

the function "bionj" in the R package *ape* v5.6.2 [24]. The tree was rooted using urial

(*O. vignei*) as the outgroup, and its graphical representation was obtained using the R

package *ggtree* v3.6.2 [25]. Principal component analyses (PCA) were performed

using the "--pca" option in PLINK v1.90b6.26 [23] with the default settings.

Furthermore, model-based clustering was performed using the sNMF v1.2 with

clusters K from 2 to 10 [26].

The F_{ST} and nucleotide diversity (π) of autosomes were calculated with vcftools

v0.1.17 [27] using 2Mbp windows and 1Mbp steps, and P value was calculated with

1000 permutation test. The F_{ST} of mitochondria and Y chromosome were calculated

with R package hierfstat v0.5.11 [28] with 1000 permutation test. The haplotype

diversity of mitochondria and Y chromosome were calculated with pegas v1.1 [29]

with 1000 permutation test.

Phylogenetic analysis

Phylogenetic analyses based on mitogenome and Y-chromosomal SNPs were

performed using the maximum likelihood (ML) algorithm of iqtree2 v2.0.7 [30].

Mitogenome sequences were aligned using mafft v7.515 [21] with the default setting,

and multiple sequence alignment (MSA) was trimmed to remove ambiguously aligned

regions using trimAl v1.4 [22] with parameters "-automated1". To keep high-quality

SNPs, we further removed all the sites with gaps and ambiguous nucleotide bases and

only kept parsimony informative sites using *ClipKIT* [31]. We tested the potential

models of DNA sequence evolution using the option "-m" test in iqtree2 v2.0.7 [30]

and found the generalized time reversible model (GTR + G) [32] was the best fitting

substitution model for mitogenome. We constructed a maximum likelihood (ML) tree

with the site model GTR + G with 1000 bootstrap replicates using iqtree2 v2.0.7 [30].

The resulting tree was rooted with bighorn as the outgroup and visualized using the

FigTree v1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/). Mitogenome haplotypes

were inferred with the function "haplotype" in the R package *pegas* v1.1 [29]. We

also constructed the parsimony mitogenome haplotype network using POPART [33].

We transformed the genotypes of Y-chromosomal 4,096 SNPs into the fasta format

using vcf2phylip.py [34]. Phylogenetic trees were built and visualized following the

same pipelines and using the same programs and substitution model as above.

Demographics and divergence time inference

The population genetic structure analyses above identified two subpopulations (*O.*

*gmelini*_1 and *O. gmelini*_2) in the Asiatic mouflons in Iran. We inferred the dynamic

changes in historical and recent effective population size (N_e) of the two

subpopulations using SMC++ v1.15.2 algorithms [35]. We selected 7 - 10 high-depth

individuals in each of the six geographically differentiated genetic groups of domestic

sheep (Central-and-East Asian, South-and-Southeast Asian, the Middle Eastern,

African, European, and American populations) [15] and the two Asiatic mouflon

subpopulations. We estimated the N_e during the past 1,000 to 100,000 years for each

group using SMC++ v1.15.2 [35] under the default setting.

For the SMC++ analysis, the reference genome's low-complexity regions were

masked using snpable (http://lh3lh3.users.sourceforge.net/snpable.shtml, last accessed

October 20, 2022) using in-house scripts. We set the mutation rate and generation

time to be 1.00×10^{-8} per nucleotide site per generation [36] and 3 years [37, 38],

respectively. To infer the divergence times between *O. gmelini*_1 and *O. gmelini*_2 as

well as between the Asian mouflon and domestic sheep, we employed the split option

of SMC++ with the same parameter settings as above.

Additionally, we inferred the demographic history of domestic sheep and Asiatic

mouflon based on mitogenomes and Y-chromosome data using Beast v2.7.3 [39] and

the Coalescent Bayesian Skyline plot [40]. Firstly, we used a coalescent model to

estimate the mutation rates for mitogenome and Y chromosome chromosomes. The

analysis was performed using the GTR + GAMMA (4) + I site model and the strict

clock model. The most recent common ancestor (MRCA) of Asiatic mouflon and

11   11

domestic sheep was set to be 0.653 million years [41]. The chain length was set to be

40,000,000 with a burn-in of 2,000,000. The results were analyzed using Tracer

(http://beast.community/tracer), assuming mutation rates estimates of 2.10×10^{-8} and

7.70×10^{-9} substitutions per site per year for the mitochondrial and Y chromosome

SNPs, respectively.

Subsequently, we utilized the mutation rates estimated above to infer the demographic

history of maternal and paternal lineages in domestic sheep and Asiatic mouflon. We

used the same GTR + GAMMA (4) + I site model as above, with the clock model set

as strict and the mutation rates of 2.10×10^{-8} and 7.70×10^{-9} per site per year for

mitochondria and Y-chromosome, respectively. The prior model remained the

Coalescent Bayesian Skyline, with the chain length of 40,000,000, a burn-in of

2,000,000, and samples drawn every 4,000 steps. The obtained results were further

analyzed using Tracer, and each analysis yielded over 200 effective samples for all

relevant parameters.

Introgression analysis

We conducted introgression analysis to clarify the genetic contributions of the two

Asiatic mouflon subpopulations to South-and-Southeast Asian (SSA) sheep. We

tested introgression using a combination method of D and f_{dM} statistics. Specifically,

we calculated D statistics [42] with a four taxon model (((P1, P2), P3), P4) using

Dsuite v0.5r50 [43]. In this analysis, Menz sheep (MEN) was selected as the reference

population (P1), while SSA populations were the target population (P2), *O. gemlini_1*

or *O. gemlini_2* was the donor (P3), and *O. canadensis* was the outgroup (P4). A *P*

value < 0.05 from a standard block-jackknife procedure was considered as evidence

of gene flow from P3 to P2.

We further implemented sliding window statistics analysis using the Dsuite v0.5r50

[43] to quantify the introgressions. We calculated f_{dM} values using a sliding window of

50 SNPs with a step of 25 SNPs across the genomes with the defined trios (MEN,

SSA, *O. gmelini_2*), where SSA represents the four South-and-Southeast Asian sheep

populations suggested to be the introgressed populations by the *D*-statistics and f_4-

ratio tests. We calculated the Z-transformed f_{dm} values, and the windows with $P < 0.05$

and $D > 0$ was defined as the potential introgressed genomic regions. We evaluated

the statistical significance using one-tailed *Z*-test in R package BSDA v1.2.1 [44].

Selective sweep tests

To assess the potential roles of genomic introgression from Iranian Asiatic

mouflons in Southern and Southeastern Asian (SSA) sheep populations, we conducted

a PBS analysis [45] to identify selective signatures in four introgressed populations

(i.e., Garut (GUR), Garole (GAR), Bangladeshi (BGE), and Sumatra (SUM)). We

pooled these four populations (GUR, BGE, SUM, and GAR) as an introgressed group,

and used the MEN and *O. gmelini_2* as the reference populations. We calculated the

PBS values using PBScan v2020.03.16 with parameters "-win 50 -step 25" described

in Hämälä and Savolainen (2019) [46] and estimated *P* values by 2000 permutation

cycles for Monte Carlo testing. and detected selective regions with top 1% PBS

values. Subsequently, we identified the overlaps between the selective and the

introgressed regions using BEDtools v2.30.0 [47] and annotated functional genes in

the overlapping genomic regions.

Gene annotation and function analyses

We annotated genes in the introgressed regions using the sheep reference assembly

Oar_rambouillet_v1.0 by BEDtools v2.30.0 [47]. We also used the sheepQTLdb

(https://www.animalgenome.org/cgi-bin/QTLdb/OA/index) [48] to detect QTLs

associated with the introgressed regions. We examined the LD blocks by

LDBlockShow v1.40 [49]. The LD decay was calculated with PopLDdecay v3.42

[50]. d_{xy} and F_{ST} for the introgressed region is calculated with Pixy 1.2.7.beta1 [51]

with a 20kb sliding window with 2kb steps. The haplotype heatmap of regions near

the introgression region was plotted with R package ComplexHeatmap v2.14.0 [52].

Results

Genetic variants, mitogenome assembly, and Y-chromosome specific SNPs

A total of 121,252,917 SNPs in the 780 wild and domestic sheep samples were obtained for downstream analyses. We obtained an average depth of 923.33 × (101.92 × - 8,324.72 ×) for the mitogenome assemblies after aligning all mitochondrial reads from the 606 samples to the reference mitochondrial genome NC_001941.1 [See Additional file 1, Table S2]. After quality control for the SNP identification on the Y-chromosome, we identified 4,096 Y-specific SNPs in a cohort of 239 rams with an average read depth of 8.26 × (~4.25 × - 16.99 ×) for SNP [See Additional file 1, Table S4].

Population genetic structure of the Asiatic mouflons in Iran

We observed two subpopulations in the 32 Asiatic mouflons in Iran (Fig. 1) with autosome SNPs. The larger subpopulation (*O. gmelini*_1) comprised 25 samples across eastern and southern Iran, while the smaller subpopulation (*O. gmelini*_2) consisted of only 7 samples limited in northwestern Iran (Fig. 1a) and [See Additional file 1, Table S2]. Low kinship values within two subpopulations eliminate inbreeding caused by sampling [See Additional file 1, Table S5]. We also observed significant

genetic divergence and different genetic diversity patterns [See Additional file 1,

Table S6] and [See Additional file 1, Table S7]. The result makes sure that the

divergence is not the artefact of inbreeding.

Phylogenetic analyses of the total 780 wild and domestic sheep samples identified the

same two subpopulations of Asiatic mouflon (*O. gmelini*_1 and *O. gmelini*_2) and six

genetic clusters of domestic sheep, which were congruent with their geographic

distribution [See Additional file 2, Figure S1] and [See Additional file 2, Figure S2].

Admixture analysis showed inter- and intra-species genetic divergence similar to the

phylogenetic tree (Fig. 1d).

[Figure 1 here]

Phylogenetic and haplotype analysis based on uniparental variants

We observed 593 mitochondrial haplotypes from the 767 mitogenomes of wild and

domestic sheep (Fig. 2a) and [See Additional file 1, Table S2]. The phylogenetic tree

inferred from the mitochondrial haplotypes revealed three monophyletic clades

(clades I, II, and III) (Fig. 2b). Clade I comprised of haplogroups A, B, and D of

domestic sheep. Clade II consisted of the domestic sheep's haplogroups C and E and

the Asiatic mouflon samples, while the remaining Asiatic mouflon samples clustered

with the urial in clade III (Fig. 2b).

For Y-chromosomal SNPs, the lineage HY2 of domestic sheep was closely related to

Asiatic mouflon in the haplogroups C and E but diverged from the main lineage HY1

(Fig. 2c) and [See Additional file 1, Table S8]. However, phylogenetic and network

analysis revealed no shared mitochondrial haplotypes and only one shared Y-

haplotypes of HY2 between domestic sheep and Asiatic mouflon in Iran (Fig. 2d and

e). We observed significant genetic divergence (*O. gmelini*_1: $F_{ST} = 0.11$, $P < 0.05$;

*O. gmelini*_2: $F_{ST} = 0.21$, $P < 0.05$) between Asiatic mouflons and domestic sheep in

whole genome levels [See Additional file 1, Table S6]. The observation implied that

the Asiatic mouflon in Iran might not be the direct ancestor of domestic sheep.

Instead, the relatively close phylogenetic relationships between the Asiatic mouflons

in Iran and domestic sheep suggested by haplogroups C, E, and Y-chromosomal

haplogroupY2 implied their possible gene flows.

[Figure 2 here]

Effective population size and divergence time

Inference of demographic history showed the Asiatic mouflons and domestic

sheep shared a similar historical pattern 10,000 years ago (Fig. 3a), confirming the

Asiatic mouflon as the wild ancestor of sheep [53]. SMC++ analysis revealed a

continual decline of N_e in domestic sheep and Asiatic mouflon during ~8,000–10,000

years ago. Nevertheless, the two Asiatic mouflon populations in Iran showed slower

declines than domestic sheep (Fig. 3a), which probably reflected the bottleneck effect

of domestication. A sharp decline of N_e was detected ~1,000-5,000 years ago in the $O.$

$gmelini\_2$, demonstrating a pattern different from that for the $O.$ $gmelini\_1$ (Fig. 3a).

We observed similar estimates of divergence time between $O.$ $gmelini\_1$ and the six

geographically differentiated genetic groups of domestic sheep during ~9,000-15,000

years ago (Fig. 3b) and [See Additional file 1, Table 9], which was congruent with the

domestication time of sheep [2]. Additionally, we found the splitting time between the

two Asiatic mouflon populations in Iran around 5,000 years ago (Fig. 3b) and [See

Additional file 1, Table 9], indicating a recent genetic divergence of Asiatic mouflons

in Iran (Fig. 1). For the divergence pattern inferred by WGS of the Asian mouflons,

the subpopulation $O.$ $gmelini\_2$ clustered with partial individuals of the subpopulation

$O.$ $gmelini\_1$ in the mitogenome and Y-chromosomal phylogenetic trees (Fig. 1 and

3). The discordant phylogenetic patterns of Asiatic mouflons in Iran inferred by

uniparental and autosomal genetic variants could be explained by the recent

hybridization events [54].

We conducted Bayesian skyline analyses to estimate N_e based on mitogenomes

and Y-chromosomes. Based on the estimated TMRCA (time to the most recent

common ancestor) of Asiatic mouflons and domestic sheep [41], three lineages

(lineage o.HCE: haplogroup of Asiatic mouflons in Iran and partial domestic sheep of

18   18

haplogroups C and E, lineage A+B+D: haplogroups A, B, and D, and lineage C+E:

haplogroups C and E) of mitogenome showed different patterns. Lineage A+B+D

showed a steep increase in N_e at approximately ~25 k years ago. For lineage C+E, it

also increased at approximately ~25 k years ago, while lineage o.HCE demonstrated

relatively fewer changes since approximately ~100 k years ago (Fig. 3c). Also, we

observed different patterns in the three Y-chromosomal lineages (lineage o.HY2:

haplogroups of Asiatic mouflons in Iran and partial domestic sheep of HY2, lineage

HY1, and lineage HY2). Lineages o.HY2 and HY2 displayed slow decline in N_e at

approximately ~25 k years ago, while lineage HY1 changed sightly since

approximately ~100 k years ago (Fig. 3d). Thus, uniparental genetic markers showed

similar demographic patterns between o.HY2 and HY2 as well as between HCE and

C/E, but different patterns between the dominant linages A+B+D and HY1 (Fig. 2 and

3). The results suggested that the Asiatic mouflons in Iran might not be the direct

ancestor of domestic sheep but be involved in the genetic introgression.

[Figure 3 here]

Phylogeography of Asiatic mouflon

We built a *Cyt-b* dataset consisting of 247 Asiatic mouflons and 8 European

mouflons. The samples of Asiatic mouflon covered the current distribution of all the

subspecies (Fig. 4a) and [See Additional file 1, Table S3]. To clarify phylogenetic

relationships between different Asiatic mouflon subspecies and domestic sheep, we

19   19

performed phylogenetic analysis of *Cyt-b* sequences of Asiatic mouflons and

domestic sheep using argali as the outgroup. In the phylogenetic tree built from *Cyt-b*,

Asiatic mouflon was assigned to 7 major clades supported by high bootstrap values

(94~100) (Fig. 4b). The topology confirmed the five mitogenome clades of domestic

sheep. Haplogroups C+E clustered into monophyletic clade, which diverged from the

haplogroups A+B+D (Fig. 4b). It was noteworthy that most of the hybrid samples

inferred above were integrated with *O. gmelini* or *O. vignei* samples, and only around

10% of samples clustered with subspecies such as *O.g.ophion* and *O.g.anatolica* (Fig.

4b) and [See Additional file 1, Table S10]. Clade 3 comprised haplogroups C+E of

domestic sheep and the Asiatic mouflons in Iran with high frequencies of shared

haplotypes between them (Fig. 4c) and [See Additional file 1, Table S10].

Haplogroups A+B+D of domestic sheep and the Asiatic mouflons from Iran and

Turkey fell within clade 6, but no common haplotypes were detected between

domestic and wild sheep (Fig. 4c) and [See Additional file 1, Table S10]. The results

were in accordance with the mitogenome analyses and implied that partial haplotypes

C+E in domestic sheep could have originated from the Asiatic mouflons in Iran by

introgression.

[Figure 4 here]

Genetic introgression

To investigate the impact of introgression from the Asiatic mouflons in Iran on

domestic sheep, we implemented introgression tests between Iranian Asiatic mouflon

and SSA by D and f_4-ratio statistics [55, 56]. We first conducted an admixture

analysis between Asiatic mouflons and the SSA populations. We identified three

genetic clusters with the smallest cross validation error ($K = 3$) (*O. gmelini*_1, *O.*

*gmelini*_2, and SSA) and detected genetic components of Asiatic mouflons in Iran in

some domestic populations (Fig. 5a) and [See Additional file 2, Figure S3]. Further,

we employed the ABBA model (((P1, P2), P3), P4) to assess introgression between *O.*

*gmelini*_1 or *O. gmelini*_2 (P3) and SSA sheep populations (P2). We used Menz

sheep (MEN) from Africa as P1, which did not show genetic introgression from

Asiatic mouflon and SSA populations [8, 15], and bighorn sheep as the outgroup P4,

which inhabit North America and have no gene flows with SSA sheep populations. Of

the two Asiatic mouflon populations, *O. gmelini*_1 or *O. gmelini*_2 served as P3 and

18 SSA populations as P2. We calculated the estimates of Patterson's D-statistic and

f_4-ratio for all the 18 SSA sheep populations, and determined introgression based on

the following criteria: $D > 0$, P-value < 0.05, and f_4-ratio > 0. Only four sheep

populations (Garut, Garole, Bangladeshi, and Sumatra sheep) showed genetic

introgression from *O. gmelini*_2 (Fig. 5b) and [See Additional file 1, Table S11]. All

the four sheep populations are distributed in the regions near to the Indian Ocean.

To further examine the genetic introgression in the four sheep populations, we

calculated f_{dm} with a 50-SNP window and 25-SNP step across the sheep genome. We

identified introgressed regions genomic with top 5% f_{dm} and $D > 0$. We obtained a

common introgression region in the four populations spanning 6,384,279 bp (Fig. 5c),

[See Additional file 1, Table S12] and [See Additional file 1, Table S13]. We then

calculated the PBS value of the four populations to identify regions under selection

with MEN sheep and *O. gmelini*_1 as the reference populations. We considered the

top 1% of PBS regions as being under selection (Fig. 5d) and [See Additional file 1,

Table S14]. We overlapped the selected regions with the common introgressed

regions and detected 19,614 bp region as the selected introgressed region [See

Additional file 1, Table S15]. We further overlapped the introgressed genomic region

with sheep QTLs [See Additional file 1, Table S15]. We found that it overlapped with

the QTLs for body_weight, carcass_fat, and fat_weight QTLs, which might be

associated with the small body size of Garole, Bangladeshi, and Garut sheep [57–59].

Additionally, the Nematodirus_FEC QTL could be related to the phenotypes of

Sumatra sheep's susceptibility to nematode parasitism [55]. The results implied that

the adaptive selected introgressed region could have been introgressed from *O.*

*gmelini*_2, and then have undergone the strong selection to facilitate their adaptation

to the hot and humid environments near the Indian Ocean.

[Figure 5 here]

Discussion

Traditionally, Asiatic mouflon had been classified into five subspecies using

phenotypic traits, such as coat color, horns, and geographic patterns [11]. However,

the intraspecies taxonomy has long lacked genetic evidence. Previous phylogenetic

analysis based on partial mtDNA divided Asiatic mouflon into several clades [3, 4,

58], different from the taxonomy of five subspecies defined by phenotypic traits and

geographic patterns [11]. Here, we identified two genetic clusters of the Asiatic

mouflons in Iran based on WGS data: *O. gmelini*_2 limited in Kaboudan, and *O.*

*gmelini*_1 over a wide geographic area in Iran (Fig. 1). We detected the divergence

time of two clusters around ~5,000 years ago (Fig. 3b) and [See Additional file 1,

Table S9], which indicated a recent split of them, and was further supported by the

inferred phylogenies based on uniparental genetic markers (Fig. 2b and c). All the

individuals of *O. gmelini* _2 collected in Kabudan inhabited the overlap geographic

region with Armenian Mouflon (*O. g.gmelini*), while individuals of *O. gmelini* _1

were sampled from the distribution areas of several subspecies such as Armenian

Mouflon (*O. g.gmelini*), Isfahan mouflon (*O.g.isphahanica*), and Laristan mouflon

(*O.g.laristanica*) (Fig. 1a) [10].

These results suggested a significant genetic divergence between Armenian Mouflon

and the other subspecies [See Additional file 1, Table S6]. Nevertheless, the absence

23   23

of phenotypic data and other sample information for the Asian mouflons and a

comprehensive sampling throughout Iran limited our in-depth understanding of

within-species genetic differentiation in Asiatic mouflon. Also, Asiatic mouflon is

under the status of "Near Threatened", and their population size has declined in the

past two decades under competition with domestic sheep, poaching, and habitat

deterioration [10, 60]. Therefore, the population genetic structure analyses and

delimitation of taxonomic classification based on WGS data are fundamental for

conservation decisions and effective conservation management [61, 62].

Genetic variants of uniparental inheritance offer insights into the domestication

and evolutionary history of farm animals, such as pigs [63], sheep [5, 14] and goats

[64, 65]. In domestic sheep, here we identified five maternal haplogroups (A, B, C, D,

and E) in two monophyletic groups (A+B+D and C+E), and two paternal clades (HY1

and HY2) (Fig. 2b and c). We found that haplotypes of most of the Asiatic mouflons

in Iran clustered with haplogroups C+E based on mitochondrial marker (24/31) and

grouped with haplogroup HY2 based on Y-chromosomal SNPs (12/21) [See

Additional file 1, Table S8]. Nevertheless, the Asiatic mouflons in Iran showed higher

divergence from the dominant maternal lineages A and B and paternal lineages HY1

(Fig. 2c and d).

Additionally, *Cyt-b* sequences collected from the whole geographic areas of

Asiatic mouflon also revealed a closer relationship between Asiatic mouflon in Iran

and domestic sheep of maternal haplogroups C+E (Fig. 4). The results supported the

hypothesis that mtDNA haplogroups C+E and Y-chromosomal haplogroup HY2

could have originated from Asiatic mouflons in Iran [14]. However, WGS analysis

revealed intraspecies substructure in the Asiatic mouflons in Iran, and genetic

introgression from the *O. gmelini*_2 showed a larger impact on domestic sheep than

*O. gmelini*_1 (Fig. 5). Similar results were observed from Alberto et al., (2018) that

IROO individuals were split into subpopulations and had admixture with some

domestic sheep [38]. Therefore, Iran's two Asiatic mouflon populations showed

similar phylogenetic patterns of uniparental inheritance but different introgression

patterns of WGS, implying multiple introgression events from the Asiatic mouflons to

domestic sheep. In summary, our results implied that Asiatic mouflons in Iran might

not be the ancestor of domestic sheep [4, 14], but could have contributed to the gene

pool of domestic sheep through multiple introgressions. Also, the results implied the

complexity of sheep domestication [4, 66].

Recent studies suggested that sheep diffused to South and Southeast Asia via the

Middle East [15]. Our results revealed the genetic introgression from Asiatic mouflon

into some South-and-Southeast Asian sheep populations, supporting previous findings

[8, 15]. Further, we detected genetic introgression in the four sheep populations of

Garut, Garole, Bangladeshi, and Sumatra, and were only from *O. gmelini*_2 in

25   25

Kabudan (Fig. 5b). The introgression event is congruent with the demographic history

of *O. gmelini_2* and South-and-Southeast Asian sheep inferred here and previously:

South-and-Southeast Asian sheep descended from Central-and-East Asian populations

around 5.85 to 6.01 ka and then admixed from the Middle Eastern sheep 3.93–4.18 ka

[15], and *O.gmelini_2* split from *O.gmelini_1* around ~5,000 years ago (Fig. 3b).

In the four sheep populations, we found that common introgression signals from *O.*

gmelini_2 covered about 4.29% of the total introgressed regions. In contrast to

population-specific introgression, common introgression could be related to a specific

trait or adaptation to similar environments, such as *EPAS1* for the adaptation to

hypoxia [45, 67]. We further integrated *PBS* analyses and found four positive

selective regions in the common introgressed regions, which were related to

carcass_fat_percentage, lean_meat_yield_percentage (Fig. 5d). This might have

contributed to the small body size phenotype of Garole, Sumatra, and Bangladeshi

sheep [58, 68, 69]. We also found a candidate gene *PCDH9* downstream ~679 kb of

the AI region (Fig. 5e), which had been reported to be associated with local adaptation

(e.g., hot and arid environments) in sheep and goats [70, 71] and fat deposition [72].

The gene showed a higher linkage with the AI region (Fig. 6a and b). The AI region

was confirmed by d_{XY} and F_{ST} of Menz sheep (MEN) and introgressed breeds (BGE,

GAR, GUR, and SUM) from SSA (Fig. 6c). We observed similar haplotype patterns

among introgressed breeds and *O.gmelini_2* in the AI region contrasting with MEN.

In contrast, no similar haplotype pattern was found in the exon of *PCDH9* (Fig. 6d).

These results implied that the AI region may regulate the function of *PCDH9*. Also,

early studies showed adipogenesis, ECM remodeling, inflammation, and lipid droplet

dynamics in mature adipose tissue played essential roles in response to climate

changes in sheep [73]. Further analysis from Chip-seq and RNA-Seq data will be

required to uncover the function and regulatory mechanisms of *PCDH9* to adapt to

hot and humid environments in sheep.

[Figure 6 here]

Conclusions

In summary, we conducted a comprehensive and integrated analysis of Asiatic

mouflon and domestic sheep based on WGS, mitogenome and Y-chromosomal

genetic variants. Our results suggested that the mitochondrial haplogroups C/E and Y-

chromosomal haplogroup HY2 of domestic sheep could have originated from the

Asiatic mouflons in Iran by introgression and further clarified the effects of the

Iranian Asiatic mouflon in the domestication processing. Additionally, our results

revealed the introgression signals of the Asiatic mouflons in Iran in several domestic

sheep populations in South and Southeast Asia and showed that the introgression

could have facilitated their adaptation to hot and humid environments near the Indian

Ocean. Our findings added new insights into the domestication and subsequent

evolutionary history of sheep.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The whole genome re-sequence data used for the study is publicly available under the

sample accession numbers listed in Additional file 1 Table S1. All mitogenomes

assembled in this study were deposited in NCBI's SRA under the accession number

OR160429-OR161034. All scripts used for this work were performed using open-

source software tools and are available from the corresponding authors upon request.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was financially supported by grants from the National Natural Science

564 Foundation of China (Nos. U21A20246, 32061133010, and 31972527), the National

565 Key Research and Development Program-Key Projects (2021YFD1200900 and

566 2021YFD1300904), the Second Tibetan Plateau Scientific Expedition and Research

567 Program (STEP) (No. 2019QZKK0501), and Initiation Fund of Sanya Institute of

568 China Agricultural University (SYND-2022-12).

Authors' contributions

571 FHL and MHL conceived and supervised the study. DFW and FHL conducted the

572 data analysis. DFW, FHL and POW wrote the manuscript, and MHL and POW

573 revised the paper. All authors read and approved the final manuscript.

Acknowledgements

576 We thank the High-performance Computing Platform of China Agricultural

577 University for providing computing resources.

References

1. Zeder MA. Domestication and early agriculture in the Mediterranean Basin:

Origins, diffusion, and impact. Proc Natl Acad Sci USA. 2008;105:11597–604.

2. Pedrosa S, Uzun M, Arranz J-J, Gutiérrez-Gil B, San Primitivo F, Bayón Y.

Evidence of three maternal lineages in near eastern sheep supporting multiple

domestication events. Proc Biol Sci.. 2005;272:2211–7.

3. Demirci S, Koban Baştanlar E, Dağtaş ND, Pişkin E, Engin A, Özer F, et al.

Mitochondrial DNA diversity of modern, ancient and wild sheep (*Ovis gmelinii*

anatolica) from Turkey: new insights on the evolutionary history of sheep. PLoS

One 2013;8:e81952.

4. Her C, Rezaei H, Hughes S, Naderi S, Duffraisse M, Mashkour M, et al. Broad

maternal geographic origin of domestic sheep in Anatolia and the Zagros. Anim

Genet. 2022;53:452–9.

5. Lv F-H, Peng W-F, Yang J, Zhao Y-X, Li W-R, Liu M-J, et al. Mitogenomic meta-

analysis identifies two phases of migration in the history of eastern eurasian sheep.

Mol Biol Evol. 2015;32:2515–33.

6. Deng J, Xie X-L, Wang D-F, Zhao C, Lv F-H, Li X, et al. Paternal origins and

migratory episodes of domestic sheep. Curr Biol. 2020;30:4085-4095.e6.

7. Kalds P, Zhou S, Gao Y, Cai B, Huang S, Chen Y, et al. Genetics of the phenotypic

evolution in sheep: a molecular look at diversity-driving genes. Genet Sel Evol.

2022;54:61.

8. Cheng H, Zhang Z, Wen J, Lenstra JA, Heller R, Cai Y, et al. Long divergent

haplotypes introgressed from wild sheep are associated with distinct morphological

and adaptive characteristics in domestic sheep. PLoS Genet. 2023;19:e1010615.

9. Yeomans L, Martin L, Richter T. Expansion of the known distribution of Asiatic

mouflon (*Ovis orientalis*) in the late Pleistocene of the southern levant. R Soc open

sci. 2017;4:170409.

10. Michel S, Ghoddousi A. *Ovis gmelini:* The IUCN red list of threatened species

2020. 2020. https://dx.doi.org/10.2305/IUCN.UK.2020-

2.RLTS.T54940218A22147055.en/ Accessed 23 October 2023.

11. Blyth E. XXVI.— an amended list of the species of the genus *Ovis*. Ann Mag Nat

Hist. 1841;7:195–201.

12. Vigne JD, Peters J, Helmer D. The first steps of animal domestication: new

archaeozoological approaches. Oxford: Oxbow Books; 2005.

13. Hiendleder S, Kaupe B, Wassmuth R, Janke A. Molecular analysis of wild and

domestic sheep questions current nomenclature and provides evidence for

domestication from two different subspecies. Proc R Soc Lond B. 2002;269:893–

904.

14. Li R, Yang P, Li M, Fang W, Yue X, Nanaei HA, et al. A hu sheep genome with

the first ovine y chromosome reveal introgression history after sheep

domestication. Sci China Life Sci. 2021; 64:1116-30.

15. Lv F-H, Cao Y-H, Liu G-J, Luo L-Y, Lu R, Liu M-J, et al. Whole-genome

resequencing of worldwide wild and domestic sheep elucidates genetic diversity,

introgression, and agronomically important loci. Mol Biol Evol.

2022;39:msab353.

16. Cao Y-H, Xu S-S, Shen M, Chen Z-H, Gao L, Lv F-H, et al. Historical

introgression from wild relatives enhanced climatic adaptation and resistance to

pneumonia in sheep. Mol Biol Evol. 2021;38:838–55.

17. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for illumina

sequence data. Bioinformatics. 2014;30:2114–20.

18. Li H, Durbin R. Fast and accurate short read alignment with burrows–wheeler

transform. Bioinformatics. 2009;25:1754–60.

19. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al.

The genome analysis toolkit: a mapreduce framework for analyzing next-

generation DNA sequencing data. Genome Res. 2010;20:1297–303.

20. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence

alignment/map format and samtools. Bioinformatics. 2009;25:2078–9.

21. Katoh K, Standley DM. Mafft multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol. 2013;30:772–80.

22. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. Trimal: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics.

2009;25:1972–3.

23. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-

generation plink: rising to the challenge of larger and richer datasets. GigaScience.

2015;4:7.

24. Paradis E, Schliep K. Ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in r. Bioinformatics. 2019;35:526–8.

25. Yu G. Using ggtree to visualize data on tree-like structures. Curr Protoc

Bioinformatics. 2020. 69:e96.

26. Frichot E, Mathieu F, Trouillon T, Bouchard G, François O. Fast and efficient

estimation of individual ancestry coefficients. Genetics. 2014;196:973–83.

27. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The

variant call format and vcftools. Bioinformatics. 2011;27:2156–8.

28. Goudet J. hierfstat, A package for r to compute and test hierarchical f-statistics.

Mol Ecol Notes. 2005;5:184–6.

29. Paradis E. Pegas: an r package for population genetics with an integrated–modular

approach. Bioinformatics. 2010;26:419–20.

30. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von

Haeseler A, et al. Iq-tree 2: new models and efficient methods for phylogenetic

inference in the genomic era. Mol Biol Evol. 2020;37:1530–4.

31. Steenwyk JL, Buida TJ, Li Y, Shen X-X, Rokas A. Clipkit: a multiple sequence

alignment trimming software for accurate phylogenomic inference. PLoS Biol.

2020;18:e3001007.

32. Gatto L, Catanzaro D, Milinkovitch MC. Assessing the applicability of the gtr

nucleotide substitution model through simulations. Evol Bioinform Online.

2007;2:145–55.

33. Leigh JW, Bryant D. Popart: full-feature software for haplotype network

construction. Methods Ecol Evol. 2015;6:1110–6.

34. Ortiz EM. Vcf2phylip v2.0: convert a vcf matrix into several matrix formats for

phylogenetic analysis. 2019. https://zenodo.org/record/2540861 Accessed 15

March 2023.

35. Terhorst J, Kamm JA, Song YS. Robust and scalable inference of population

history from hundreds of unphased whole-genomes. Nat Genet. 2017;49:303–9.

36. Beynon SE, Slavov GT, Farré M, Sunduimijid B, Waddams K, Davies B, et al.

Population structure and history of the welsh sheep breeds determined by whole

genome genotyping. BMC Genet. 2015;16:65.

37. Zhao Y-X, Yang J, Lv F-H, Hu X-J, Xie X-L, Zhang M, et al. Genomic

reconstruction of the history of native sheep reveals the peopling patterns of

nomads and the expansion of early pastoralism in East Asia. Mol Biol Evol.

2017;34:2380–95.

38. Alberto FJ, Boyer F, Orozco-terWengel P, Streeter I, Servin B, de Villemereuil P,

et al. Convergent genomic signatures of domestication in sheep and goats. Nat

Commun. 2018;9:813.

39. Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M,

Gavryushkina A, et al. Beast 2.5: an advanced software platform for bayesian

evolutionary analysis. PLoS Comput Biol. 2019;15:e1006650.

40. Drummond AJ. Bayesian coalescent inference of past population dynamics from

molecular sequences. Mol Biol Evol. 2005;22:1185–92.

41. Upadhyay M, Kunz E, Sandoval-Castellanos E, Hauser A, Krebs S, Graf A, et al.

Whole genome sequencing reveals a complex introgression history and the basis

of adaptation to subarctic climate in wild sheep. Mol Ecol. 2021;30:6701–17.

42. Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, et al. Ancient

admixture in human history. Genetics. 2012;192:1065–93.

43. Malinsky M, Matschiner M, Svardal H. Dsuite - fast d-statistics and related

admixture evidence from vcf files. Mol Ecol Resour. 2021;21:584–95.

44. Arnholt AT, Evans B. Bsda: basic statistics and data analysis. 2017. https://cran.r-

project.org/web/packages/BSDA/index.html. Accessed 28 Sep 2023.

45. Yi X, Liang Y, Huerta-Sanchez E, Jin X, Cuo ZXP, Pool JE, et al. Sequencing of

50 human exomes reveals adaptation to high altitude. Science. 2010;329:75–8.

46. Hämälä T, Savolainen O. Genomic patterns of local adaptation under gene flow in

Arabidopsis lyrata. Mol Biol Evol. 2019;36:2557–71.

47. Quinlan AR, Hall IM. Bedtools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 2010;26:841–2.

48. Hu Z-L, Park CA, Reecy JM. Bringing the animal QTLdb and CorrDB into the

future: meeting new challenges and providing updated services. Nucleic Acids

Res. 2022;50:D956–61.

49. Dong S-S, He W-M, Ji J-J, Zhang C, Guo Y, Yang T-L. Ldblockshow: a fast and

convenient tool for visualizing linkage disequilibrium and haplotype blocks based

on variant call format files. Brief Bioinform. 2021;22:bbaa227.

50. Zhang C, Dong S-S, Xu J-Y, He W-M, Yang T-L. Poplddecay: a fast and

effective tool for linkage disequilibrium decay analysis based on variant call

format files. Bioinformatics. 2019;35:1786–8.

51. Korunes KL, Samuk K. Pixy: unbiased estimation of nucleotide diversity and

divergence in the presence of missing data. Mol Ecol Resour. 2021;21:1359–68.

52. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. Bioinformatics. 2016;32:2847–9.

53. Sanna D, Barbato M, Hadjisterkotis E, Cossu P, Decandia L, Trova S, et al. The

first mitogenome of the Cyprus mouflon (*Ovis gmelini ophion*): new insights into

the phylogeny of the genus *Ovis*. PLoS One. 2015;10:e0144257.

54. Teng H, Zhang Y, Shi C, Mao F, Cai W, Lu L, et al. Population genomics reveals

speciation and introgression between brown Norway rats and their sibling species.

Mol Biol Evol. 2017;34:2214–28.

55. Reich D, Thangaraj K, Patterson N, Price AL, Singh L. Reconstructing indian

population history. Nature. 2009;461:489–94.

56. Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, et al. A draft

sequence of the neandertal genome. Science. 2010;328:710–22.

57. Jakaria J, Zein MSA, Sulandari S, Subandriyo S, Muladno M. The use of

microsatellite markers to study genetic diversity in Indonesian sheep. J Indones

Trop Anim Agric. 2012;37:1–9.

58. Sharma RC, Arora AL, Narula HK, Singh RN. Characteristics of Garole sheep in

India. Anim Genet Resour Inf. 1999;26:57-64

59. Banerjee R, KumarManda P, Kumar Pal U, Ray K. Productivity and genetic

potential of Garole sheep of India-a review. Asian J Anim Sci. 2010;4:170–89.

60. Bleyhl B, Arakelyan M, Askerov E, Bluhm H, Gavashelishvili A, Ghasabian M, et

al. Assessing niche overlap between domestic and threatened wild sheep to

identify conservation priority areas. Divers Distrib. 2019;25:129–41.

61. Yang L, Wei F, Zhan X, Fan H, Zhao P, Huang G, et al. Evolutionary

conservation genomics reveals recent speciation and local adaptation in threatened

takins. Mol Biol Evol. 2022;39:msac111.

62. Hu Y, Thapa A, Fan H, Ma T, Wu Q, Ma S, et al. Genomic evidence for two

phylogenetic species and long-term population bottlenecks in red pandas. Sci Adv.

2020;6:eaax5751.

63. Ai H, Zhang M, Yang B, Goldberg A, Li W, Ma J, et al. Human-mediated

admixture and selection shape the diversity on the modern swine (*Sus scrofa*) y

chromosomes. Mol Biol Evol. 2021;38:5051–65.

64. Luikart G, Gielly L, Excoffier L, Vigne J-D, Bouvet J, Taberlet P. Multiple

maternal origins and weak phylogeographic structure in domestic goats. Proc Natl

Acad Sci USA. 2001;98:5927–32.

65. Naderi S, Rezaei H-R, Pompanon F, Blum MGB, Negrini R, Naghash H-R, et al.

The goat domestication process inferred from large-scale mitochondrial DNA

analysis of wild and domestic individuals. Proc Natl Acad Sci USA.

2008;105:17659–64.

66. Yang L, Zhang X, Hu Y, Zhu P, Li H, Peng Z, et al. Ancient mitochondrial

genome depicts sheep maternal dispersal and migration in eastern Asia. J Gene

Genomics. 2023; https://doi.org/10.1016/j.jgg.2023.06.002 in press.

67. Wu D-D, Yang C-P, Wang M-S, Dong K-Z, Yan D-W, Hao Z-Q, et al.

Convergent genomic signatures of high-altitude adaptation among domestic

mammals. Natl Sci Rev. 2020;7:952–63.

68. Porter V. Mason's world dictionary of livestock breeds, types and varieties, 6th

edition. Wallingford: CABI Publishing; 2020.

69. Putra WPB, Alwi Y, Suryani H. Morphometrics characterization of thin-tail sheep in lowland and highland areas. Trop Anim Sci J. 2021;44:386–98.

70. Serranito B, Cavalazzi M, Vidal P, Taurisson-Mouret D, Ciani E, Bal M, et al. Local adaptations of mediterranean sheep and goats through an integrative approach. Sci Rep 2021;11:21363.

71. Kim E-S, Elbeltagy AR, Aboul-Naga AM, Rischkowsky B, Sayre B, Mwacharo JM, et al. Multiple genomic signatures of selection in goats and sheep indigenous to a hot arid environment. Heredity (Edinb). 2016;116:255–64.

72. Mastrangelo S, Moioli B, Ahbara A, Latairish S, Portolano B, Pilla F, et al. Genome-wide scan of fat-tail sheep identifies signals of selection for fat deposition and adaptation. Anim Prod Sci. 2018;59:835–48.

73. Xu Y-X, Wang B, Jing J-N, Ma R, Luo Y-H, Li X, et al. Whole-body adipose tissue multi-omic analyses in sheep reveal molecular mechanisms underlying local adaptation to extreme environments. Commun Biol. 2023;6:159.

Figures

Figure 1 Population genetics structure of Asiatic mouflons in Iran (*O. gmelini*). a

Geographic distribution of 32 Asiatic mouflons. Purple and blue dots indicate the location of *O.*

*gmelini*_1 and *O. gmelini_2* subpopulations; **b** Principal components 1 and 2 for the 32 Asiatic

mouflon in Iran; **c** Neighbor-joining (NJ) trees of the 32 Asiatic mouflons in Iran based on

whole-genome SNPs using the nine urial samples as the outgroup; **d** Population genetic structure

of the Asiatic mouflons in Iran and domestic sheep inferred from the program the sNMF v1.2 (*k*

= 6) using whole-genome SNPs.

Figure 2 Phylogeny of Asiatic mouflon, urial, and domestic sheep inferred from

mitogenomes and Y-chromosomal genetic variants. a Geographical distribution of the

samples in this and previous studies; **b** Phylogeny of Asiatic mouflon, urial, and domestic sheep

inferred from 767 mitogenomes; **c** Phylogeny of Asiatic mouflon, urial, and domestic sheep

inferred from 239 Y-chromosomal SNPs; **d** Network of mitogenome haplotypes; **e** Network of

Y-chromosome haplotypes. Bootstrapping values above 50% were labeled on the phylogenetic

tree. Bighorn sheep were used as the outgroup in phylogenetic analyses based on mitogenome

and Y-chromosome genetic variants.

Figure 3 Demographic history of Asiatic mouflon in Iran and domestic sheep. a Effective

population sizes (N_e) across six domestic sheep populations and the two subpopulations of

Asiatic mouflon in Iran inferred from SMC++; **b** Inference of split times between *O. gmelini*_1

and the other wild and domestic sheep populations. The vertical lines represent the split time

39   39

between each pair of populations; **c** Bayesian skyline plot (BSP) reconstruction of historical

change in effective population size (N_e) of domestic sheep of maternal haplogroups HA+B+D,

and HC+E, and the Asiatic mouflons in Iran; **d** Bayesian skyline plot (BSP) reconstruction of

historical changes in N_e of paternal lineages HY1 and HY2 of domestic sheep and the Asiatic

mouflon in Iran. The middle line represents median estimates of $N_e\tau$, and the shaded area

represents 95% highest posterior density (HPD) intervals.

Figure 4 Geographic distribution of Asiatic mouflon and the phylogenetic relationship

among Asiatic mouflon subspecies and domestic sheep. **a** Geographical distribution of Asiatic

mouflon and samples information of 32 Asiatic mouflon in Iran and 301domestic sheep samples;

b Phylogenetic tree inferred based on complete *Cyt-b* sequences using the maximum likelihood

algorithms. Bootstrap values above 50% were labeled on the phylogenetic tree. **c** Network of

Asiatic mouflon in Iran and Turkey and domestic sheep lineage of C and E built from complete

Cyt-b sequences.

Figure 5 Genetic introgression of Asiatic mouflon in SSA domestic populations. a

Admixture proportions of Asiatic mouflon in Iran and SSA domestic populations inferred from

the program the sNMF v1.2 ($K = 3$) using whole-genome SNPs. **b** Heatmap of *D*-statistics and f_4-

ratio calculated by Dsuite v0.5 applied to data including *O. gmelini*_1, *O. gmelini*_2, and SSA

domestic populations; **c**, Venn diagram for the shared introgression regions among four domestic

South and Southeast populations; **d** Manhattan plot showing signals of selection in the four

introgressed SSA populations (BGE/GAR/GUR/SUM) detected by the PBS with the dashed line

marking 99.9% threshold. The introgression region, QTL, and adjacent gene *PCDH9* were

labeled with blue, purple, and green bars, respectively. The sheep QTLs overlapped with the top

1% PBS regions are shown, **e** PBS and f_{dm} values around the significant introgressed genomic

region on Chr.10.

Figure 6 Candidate gene *PCDH9* associated with the significant introgression signal on

Chr. 10. **a** Linkage disequilibrium (LD) blocks in the significant introgression region and gene

PCDH9 (chr10:41,104,271-43,073,231). The red and yellow bars represent the location of the

significant introgression region and gene *PCDH9*; **b** Pattern of LD decay in the introgressed

populations. The black line is calculated from the whole 10 chromosomes, the red line is

calculated from the beginning of the significant region to the next 800 kb (chr10:41,204,271-

42,004,271); **c** D_{xy} and F_{ST} around the significant introgression region on Chr. 10

(chr10:41,104,271-41,313,804), calculated by pixy v1.2.7.beta1; **d** Haplotype pattern of the

significant introgression region on Chr. 10 (chr10:41,204,271-41,213,804) (left) and exons of

PCDH9 (right).

Additional files

833

834 **Additional file 1 Table S1**

835 Format: excel

836 Title: 780 samples information for autochromosome analysis.

837 **Additional file 1 Table S2**

838 Format: excel

839 Title: 770 samples information for mitochondria chromosome analysis.

840 **Additional file 1 Table S3**

841 Format: excel

842 Title: 337 samples information of *Cyt-b* sequence in this study.

843 **Additional file 1 Table S4**

844 Format: excel

845 Title: 239 samples for Y chromosome analysis in this study.

846 **Additional file 1 Table S5**

Format: excel

Title: Kinship of 32 *O. gmelini* samples.

Additional file 1 Table S6

Format: excel

Title: The FST of autosome, mitochondria and Y chromosome of *O. gmelini*_1, *O. gmelini*_2

and *O. aries.*

Additional file 1 Table S7

Format: excel

Title: The nucleotide diversity / haplotype diversity of autosome, mitochondria and Y

chromosome of *O. gmelini*_1, *O. gmelini*_2 and *O. aries.*

Additional file 1 Table S8

Format: excel

Title: Haplotype information of mitogenomes and Y chromosomes.

Additional file 1 Table S9

Format: excel

Title: Split times estimated by SMC++.

43　　　　　　　　　　　　　　　　　　43

Additional file 1 Table S10

Format: excel

Title: Haplotype information of *Cyt-b* sequences.

Additional file 1 Table S11

Format: excel

Title: *D*-statistics from Iranian mouflon to SSA populations.

Additional file 1 Table S12

Format: excel

Title: Details for chromosome-level introgression across four breeds.

Additional file 1 Table S13

Format: excel

Title: Common introgressed blocks from *O.gmelini_2* into four SSA populations.

Additional file 1 Table S14

Format: excel

Title: Population branch statistic (PBS) of putative selective regions in the SSA population.

Additional file 1 Table S15

Format: excel

Title: Overlaps regions among selected regions identified by PBS, introgression regions and

QTLs regions.

Additional file 2 Figure S1

Format: word

Title: 780 autosome NJ-tree showing the same two cluster of *O. gmelini* as Fig.1b and domestic

sheep samples clustered based on their geographic distribution.

Additional file 2 Figure S2

Format: word

Title: Admixture of 770 samples for *K*=2-9.

Additional file 2 Figure S3

Format: word

Title: Admixture of SSA 18 breeds with two *O. gmelini* populations for *K*= 2-9.

45   45

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6